INDEX FILICUM

CARL CHRISTENSEN

INDEX FILICUM

SIVE

ENUMERATIO OMNIUM GENERUM
SPECIERUMQUE

FILICUM ET HYDROPTERIDUM

AB ANNO 1753 AD FINEM ANNI 1905
DESCRIPTORUM

ADJECTIS

SYNONYMIS PRINCIPALIBUS, AREA GEOGRAPHICA
ETC.

HAFNIAE 1906
APUD H. HAGERUP

TYPIS TRIERS BOGTRYKKERI (G. L. LIND & NUMA FRÆNKEL)

PREFACE

At the point of completing the edition of »Index Filicum« it only remains to me to return my sincere thanks for the unanimous recognition, with which the professionals have received my work and particularly for the valuable assistance that has been rendered me from different sides.

The book, as a whole, is my own work; only in a few instances have I found it necessary to seek advice and help. In the first place I must here mention the name of the famous pteridologist Dr. H. CHRIST, Basle, without whose assistance I should hardly have succeeded in getting the book published. From the very first Dr. CHRIST has embraced my work with the greatest interest and during the time of editing he has very often helped me in different ways. For this I thank Dr. CHRIST most heartily. Next I must mention Mr. A. GEPP of British Museum. Mr. GEPP has with the utmost readiness excerped several works to which I had no access, and thereby he has contributed to my work being well nigh complete. I also take the opportunity to beg Mr. GEPP accept my best thanks. Further I must thank Prof. Dr. G. HIERONYMUS, Berlin, who has unravelled certain difficult groups of forms of *Asplenium* for me.

In conclusion it is my pleasant duty to thank the DIRECTORS OF THE CARLSBERG FUND, Copenhagen, for the pecuniary support they have given me as an aid towards the publication, without which it would have been impossible for me to bring out the book.

I herewith surrender my work to the pteridologists and others studying the science of botany, and I only beg that they will forgive the no doubt numerous mistakes and errors, which it has been impossible to avoid. As it is my intention (time and health permitting), at some future time to issue a Supplement, I shall be most grateful for all communications pointing out omissions and errors.

Copenhagen, Sept. 15. 1906.

Carl Christensen.

In explanation of the principles I have adopted in preparing the Index only a few words are needed, as everybody using the book will soon be able to understand the plan of it. I shall here confine myself to following: — As a basis of the whole work I have chosen the system followed in ENGLER und PRANTL: »Die natürlichen Pflanzenfamilien«; still I have found it correct to depart from this system in some few cases, which are all emphasized in the first part of the Index, partly from systematic and partly from nomenclaturic reasons. Although I am far from agreeing with DIELS in his view of genera, still I have considered it right upon the whole to follow his system, because Engler und Prantl certainly is or will be the basis of the arrangement of species in many museums and private collections. Only in a very few cases (for instance by adopting *Lepicystis* as a separate genus), I have not thought it fit to follow DIELS. In other cases the deviations are owing to later investigations, to which I everywhere have paid the greatest attention.

Besides stating where the original description is to be found, I have under each species quoted the page, where the species is described or mentioned in HOOKER and BAKER: »Synopsis Filicum«, 2. Ed. 1874 (abbreviated to HB), in H. CHRIST: »Die Farnkräuter der Erde«. 1897. (Chr.), and in »Die Natürlichen Pflanzenfamilien« Teil 1, Abt. 4. 1899—1902. (NPfl). These quotations will, I think, be of considerable value, as you generally in works easy of access quickly may find the desired information about a species without being obliged to resort to the original diagnosis, which often is so difficult to get at. Besides, in genera monographically dealt with in recent times, the monograph in question is quoted as much abridged.

With regard to the limitation of species I have, of course, in most cases had to follow others. My principle has been: In genera, treated monographically, I have chiefly followed the monograph concerned; otherwise I have relied on the identifications of authors having had the original specimens for examination, or others, who to a great extent have been enabled to study the species in the field (for inst. JENMAN as far as the West-Indian, and BEDDOME as far as the Indian species are concerned). Still I have no doubt that a great number of names, reduced to synonyms, are representing good species, and especially many forms by HOOKER and BAKER reduced to mere synonyms, according to their comprehensive ideas of species, probably will prove to be good species. The numerous species described as new these last years have, of course, been adopted without criticism.

MS. names I have not included. An exceedingly great number of these are given as synonyms in different pteridographical works. They have no claim of priority as published names, and the authors are doing the nomenclature a bad service by reprinting such names without descriptions. A continuation of this practise ought to be carefully avoided in the future. If I were to include these MS. names, which is done in several Indices, my book would have been larger by several sheets. Names of garden forms are rather fully included only from the time of the last 10 years, of older forms only such as are still being cultivated.

I must point out that not all the synonyms of a species are given under the species; I have preferably only referred to those representing forms, which are often considered as being valid species and without exception forms that are being dealt with as species in one of the three mentioned principal works. In addition I have everywhere of each species as well as of each of its synonyms named as far as it was possible all the genera, to which each form has been referred, by whom and when, in chronological order. This should give

one an idea of how the systematical classification of the species has been understood in the course of times; in a certain sense it may be said to give us its systematical history and render it possible in a moment to make any conceivable combination of names with the correct author-citation.

The nomenclaturic principles, underlying the designation of genera and species, will plainly appear from the work itself. I have a clear understanding that some people wont approve of the names chosen, but it is a comfort to me that the case would have been just the same had I followed any other principles. I shall only mention here that no name published before 1st edition of Linné's Species plantarum 1753 has been included, neither any name published since 31. Dec. 1905.

Still a remark about the great desirability that authors when naming new species should be careful in ascertaining, as for as it is possible, that the contemplated name of species has not already been used (even if now it has been reduced to a synonym) not only in the genus in question, but also in the subgenera of the latter. For instance: No *Polypodium* should be given a name that has been used under *Campyloneurum, Goniophlebium, Pleopeltis,* etc. In exceedingly many instances you never know, if this older name may not be adopted by a later author, the form in question represented by the name proving to be a good species and accordingly will be referred to *Polypodium*. Hence the necessity of having a new name for the new species. This may be easily avoided by consulting 1st part of Index and see which genera (adopted or not adopted) should be looked over in the list of species (or synonyms). Another instance: No new species belonging to *Polypodium § Phegopteris* Hk. & Bak. ought to be given a name which formerly has been used either under *Polypodium, Dryopteris, Nephrodium, Lastrea, Aspidium, Phegopteris, Leptogramma, Goniopteris,* etc. This has been my principal nomenclaturic point of view, and it has been the cause of a great part of the versions of names done. If this rule is conscientiously followed in the future, I firmly believe that the confusion of the pteridological nomenclature will be reduced considerably.

In the first part I have attempted to arrange all the generic synonyms under each adopted genus in such a manner as will enable you to find out under which generic name a given group of allied species has been placed. The order of the group-names under each subgenus is following the arrangement of the species given in »Die Natürl. Pflanzenfam.« and is of no systematic value but a guide to the student in the mystery of pteridological nomenclature. — In the enumeration of the species of the larger genera each species is prefixed a letter signifying the subgenus — adopted in the first part — to which the species is meant to belong. In the case of *Trichomanes* these subgenera are named by different names in the first and second part, the names in the first part in my opinion being the best.

In the »Catalogue of Literature« I have confined myself to enumerating only the works which are quoted in the Index. Several others should be included if a complete bibliography of works dealing with ferns, systematically or geographically, was to be compiled, but the place did not permit this.

Explicatio abbreviationum essentialium.

HB. = Hooker and Baker: Synopsis Filicum. Second Edition. London 1874.
Chr. = H. Christ: Die Farnkräuter der Erde. Jena 1897.
NPfl. = Engler und Prantl: Die natürlichen Pflanzenfamilien. I Teil, Abteilung 4. Leipzig 1902.

Numerus cit. post »Mett. n.« = numerus speciei in Mettenii: Farngattungen:
I. Polypodium (cit. sub Cyclophoro et Polypodio).
IV. Phegopteris und Aspidium (cit. sub Aspidio, Dryopteride, Polysticho).
V. Cheilanthes (cit. sub Adiantopside, Cheilanthide, Notholaena).
VI. Asplenium (cit. sub Asplenio, Athyrio, Diplazio).

Signum - inter nomina terrarum = usque ad.
Signo ; synonyma absoluta separantur.
Signo — varietates vel formae speciei separantur.
Signum * ante nomen subspeciem indicat.

Series scriptorum periodicorum citatorum numero Romano (II) indicatur.
Volumen libri citati numero crassiusculo (**2**) indicatur.
Pagina libri citati numero tenui (2) indicatur.
t. = tabula.
f. = figura.

Nomina specierum litteris **crassiusculis** impressa.
» specierum valde dubiarum et hortulanorum litteris *cursivis* impressa.
Synonyma litteris rectis impressa.

ENUMERATIO GENERUM SYSTEMATICA

CUM

SYNONYMIS OMNIBUS CHRONOLOGICE ET SYSTEMATICE ORDINATIS

FILICALES.

1. FILICALES LEPTOSPORANGIATAE.

A. EUFILICINEAE.

I. **HYMENOPHYLLACEAE.**

I. VERAE.

1. TRICHOMANES Linné 1753 emend. Smith 1793. — Species: 228.

Typ. ?

1764. Pyxidaria Gleditsch.
1790. Achomanes Neck. (§ 5).
1824. Feea Bory. (§ 6).
— Hymenostachys Bory. (§ 6 a).
1827. Didymoglossum Desv. (§ 2 et 8 b).
1843. Abrodictyum Pr. (§ 8 b).
— Achomanes (Neck. 1790) Pr. (Trichomanes §). (§ 5).
— Cardiomanes Pr. (§ 4).
— Cephalomanes Pr. (§ 7 b).
— Chilodium pt. Pr. (Didymoglossum §). (§ 8 h).
— Crepidium Pr. (Didymoglossum §). (§ 8 a).
— Hemiphlebium Pr. (§ 2 b).
— Lecanium Pr. (§ 2 c).
— Meringium Pr. (?).
— Microgonium Pr. (§ 2 a).
— Neurophyllum Pr. (non Torr. et Gray 1838: gen. Umbelliferar.). (§ 5 f).
— Pachychaetum Pr. (Trichomanes §). (§ 8 h).
1843. Ragatelus Pr. (§ 5 b).
1848. Homoeotes Pr. (§ 6 b).
— Macroglena Pr. (Trichomanes §). (§ 8 i).
1849. Crepidomanes Pr. (§ 8 a).
— Leucomanes Pr. (§ 8 c).
— Odontomanes Pr. (§ 5 g).
— Pleuromanes Pr. (§ 8 c).
— Pseudachomanes Pr. (Trichomanes §). (§ 2 c).
— Taschneria Pr. (§ 8 b).
1851. Neuromanes Trev.; v.d.B. 1859. (§ 5 f).
1857. Bergera Schaffn. msc. (§ 2).
1861. Adiantopsis v.d.B. (non Fée 1850-52). (§ 8 j).
— Craspedoneuron v.d.B. (§ 8 c).
— Davalliopsis v.d.B. (§ 8 f).
— Gonocormus v.d.B. (§ 3).
— Lacostea v.d.B. (§ 7 a).
— Maschalosorus v.d.B. (§ 6 c).
— Phlebiophyllum v.d.B. (§ 8 d).
— Ptilophyllum v.d.B. (§ 5).

1872. Mülleria Schaffn. msc. (§ 5 c).
1875. Acarpacrium Prantl (Ptilophyllum §). (§ 5 e).
— Lacosteopsis Prantl (Trichomanes §). (§ 8 g).
— Leptomanes Prantl (Trichomanes §). (§ 8 e).
1875. Microtrichomanes Mett.; Prantl (Gonocormus §). (§ 3 a).
— Selenodesmium Prantl (Trichomanes §). (§ 8 h).
— Trigonophyllum Prantl (Ptilophyllum §). (§ 5 d).
1897. Holophlebium Christ (Trichomanes §). (§ 2, 3, 5, 6, 7, 8).

§ 1. **Serpyllopsis** Christ 1897 (caespitosum *).
Serpyllopsis v.d.B. 1861.

§ 2. **Didymoglossum** (hymenoides).
Didymoglossum pt. Desv. 1827 = Didymoglossum § Eudidymoglossum pt. Pr. 1843 = Didymoglossum v.d.B. 1859 = Hemiphlebium Prantl 1875 = Trichomanes § Hemiphlebium Christ 1897.
Bergera Schaffn. 1857 msc.
a. Microgonium Pr. 1843 (cuspidatum).
b. Hemiphlebium Pr. 1843 (pusillum).
c. Lecanium Pr. 1843 (membranaceum).

§ 3. **Gonocormus** (proliferum).
Gonocormus v.d.B. 1861.
a. Microtrichomanes Mett. Prantl 1875 (digitatum).

§ 4. **Cardiomanes** (reniforme).
Cardiomanes Pr. 1843.

§ 5. **Achomanes** Pr. 1849 (crispum).
? Achomanes Neck. 1790 = Ptilophyllum v.d.B. 1861 emend. Prantl 1875 part.
a. Euachomanes (crispum) = Ptilophyllum § Achomanes Prantl 1875.
b. Ragatelus Pr. 1843 (crinitum).
c. Pseudachomanes Pr. 1849 (polypodioides) = Mülleria Schaffn. 1872.
d. Trigonophyllum Prantl 1875 (arbuscula).
e. Acarpacrium Prantl 1875 (alatum).
f. Neuromanes Trev. 1851 (pinnatum) = Neurophyllum Pr. 1843 = Ptilophyllum § Feea pt. Prantl 1875.
g. Odontomanes Pr. 1849 (Hostmannianum).

§ 6. **Feea** (botryoides).
Feea Bory 1824 = Ptilophyllum § Feea pt. Prantl 1875.
a. Hymenostachys Bory 1824 (diversifrons).
b. Homoeotes Pr. 1848 (heterophyllum) = Ptilophyllum § Homoeotes Prantl 1875.
c. Maschalosorus v.d.B. 1861 (osmundoides v. Mougeotii).

§ 7. **Cephalomanes** (javanica).
Cephalomanes Pr. 1843 emend. = Lacostea Prantl 1875.
a. Lacostea v.d.B. 1861 (pedicellatum).
b. Cephalomanes Pr. 1843 (javanica).

---

*) nomen parenthesi inclusum est nomen speciei typicae subgeneris vel sectionis ut in Indice specierum nominatae.

§ 8. **Eutrichomanes** (?).

Trichomanes Prantl 1875.

a. Crepidium Pr. 1843 (humile) = Trichomanes § Crepidomanes Prantl 1875 — Crepidomanes Pr. 1849 (intramarginale).

b. Taschneria Pr. 1849 (bipunctatum) = Didymoglossum pt. Desv. 1827 = Didymoglossum § Eudidymoglossum pt. Pr. 1843 = Trichomanes § Didymoglossum Prantl 1875. Christ 1897.

c. Pleuromanes Pr. 1849 (pallidum) = Craspedoneuron v.d.B. 1861 = Trichomanes § Craspedoneuron Prantl 1875 — Leucomanes Pr. 1849.

d. Phlebiophyllum v.d.B. 1861 (venosum).

e. Leptomanes Prantl 1875 (capillaceum).

f. Abrodictyum Pr. 1843 (Cumingii) = Habrodictyon v.d.B. 1861.

(- Eutrichomanes Prantl 1875).

g. Lacosteopsis Prantl 1875 (rupestre).

h. Pachychaetum Pr. 1843 (rigidum) = Trichomanes § Selenodesmium Prantl 1875.

i. Macroglena Pr. 1848 (meifolium).

j Davalliopsis v.d.B. 1861 (elegans) = Adiantopsis v.d.B. 1861.

2. HYMENOPHYLLUM Smith 1793. — Species: 231.

Typ. Trichomanes tunbrigense L.

1753. Trichomanes sp. L.
1793. Hymenophyllum Sm.
1800. Ptychomanes Hedw.
1843. Chilodium pt. Pr. (Didymoglossum §). (§ 2 d).
— Craspedophyllum Pr. (Hymenophyllum §). (§ 1 a).
— Cycloglossum Pr. (Hymenophyllum §). (§ 1 d).
— Didymoglossum pt. Pr.
— Hymenoglossum Pr. (§ 1 a).
— Leptocionium Pr. (§ 2).
— Myrmecostylum Pr. (§ 2 b).
1843. Ptychophyllum Pr. (§ 2 c).
— Sphaerocionium Pr. (§ 1 c).
— Sphaerodium Pr. (Hymenophyllum §). (§ 2 a).
1849. Amphipterum Pr. (§ 1 e).
— Dermatophlebium Pr. (§ 1 i).
— Mecodium Pr. (§ 1 h).
(1856. Euphorophyllum v.d.B. (Hymenophyllum §)).
(— Charithophyllum v.d.B. (Hymenophyllum §)).
1861. Diploöphyllum v.d.B. (§ 1 f).
— Pachyloma v.d.B. (§ 1 b).

§ 1. **Euhymenophyllum** (?).

a. Hymenoglossum Pr. 1843 (cruentum).

b. Craspedophyllum Pr. 1843 (marginatum) = Pachyloma v.d.B. 1861.

c. Sphaerocionium Pr. 1843 (?).

d. Cycloglossum Pr. 1843 (Ex.: decurrens).

e. Amphipterum Pr. 1849 (fuscum).

f. Diploöphyllum v.d.B. 1861 (dilatatum).

g. Euphorophyllum v.d.B. 1856.

h. Mecodium Pr. 1849 (sanguinolentum).

i. Dermatophlebium Pr. 1849 (sericeum).

§ 2. **Leptocionium** (fucoides: dicranotrichum).

Leptocionium Pr. 1843, emend. v.d.B. 1861.

a. Sphaerodium Pr. 1843 (tunbridgense).
b. Myrmecostylum Pr. 1843 tortuosum).
c. Ptychophyllum Pr. 1843 (dichotomum).
d Chilodium Pr. 1843 (denticulatum).

II. DUBIAE.

3. LOXSOMA R.Brown 1836. (Loxoma). — Species: 1.
Typ. L. Cunninghami R.Br.

4. LOXSOMOPSIS Christ 1904. — Species: 2.
Typ. L. costaricensis Christ.

II. CYATHEACEAE.

I. DICKSONIEAE.

1. BALANTIUM Kaulfuss 1824 emend. — Species 3.
Typ. Dicksonia culcita L'Hérit.

1836. Culcita Pr.
Dicksonia § Eudicksonia pt. Hk. Bak. 1866.
Dicksonia sp. Christ 1897.

2. DICKSONIA L'Héritier 1788 reduct. - Species: 17.
Typ. D. arborescens L'Hérit.

Balantium pt. Klf. 1824. Pr. 1836.
Dicksonia § Eudicksonia pt. Hk. Bak. 1866.

Obs. Genus 1 Balantium Diels 1899 (non Presl 1836) certe Dicksonia rectius appellandum est, et genus 2. Dicksonia Diels = Balantium Klf. emend. Presl 1836 Balantium.

3. CIBOTIUM Kaulfuss 1820. 1824. — Species: 9.
Typ. C. Chamissoi Klf.

1824. Pinonia Gaud. 1824.
1827. Dicksonia sp. Desv. Hk. et Bak. 1866.
1844. Hiatea Menzies.
Dicksonia § Cibotium Hk. et Bak. 1866.

II. THYRSOPTERIDEAE.

4. THYRSOPTERIS Kunze 1834. — Species: 1.
Typ. T. elegans Kze.

? Chonta Molina (non vidi).
1836. Panicularia Colla.

III. CYATHEEAE.

5. CYATHEA Smith 1793 reduct. — Species: 182.

Typ. Polypodium arboreum L.

1753. Polypodium sp. L.
1793. Cyathea Sm. pt.
1801. Sphaeropteris Bernh. (medullaris).
1836. Disphenia Pr. (arborea).
— Notocarpia Pr. (Cyathea §) (sp. soris in medio dorso venae venulaeve).
1838. Schizocaena J. Sm. (moluccana).
1850—52. Hemithelia § Disphenia Fée (Pr. 1836).
1858. Cyatheopsis Karst. (Cyathea §) (sp. pinnis pinnulisque articulatis).
1873. Eatoniopteris Bomm. (sp. american. indusio semigloboso ad maturitatem sori irregulariter soluto).

6. HEMITELIA R.Brown 1810. — Species: 59.

Typ. Cyathea multiflora Sm.

1753. Polypodium sp. L.
1793. Cyathea sp. Sm.
1810. Hemitelia R.Br.
1836. Cnemidaria Pr. (§ 1).
1842. Amphicosmia Gardn. (§ 3).
— Hymenostegia pt. J. Sm. (Alsophila §) (§ 3).
1844. Eleutheria Kze. (Hemitelia §). (§ 1 d).
1848. Actinophlebia Pr. (§ 1 c).
— Hemistegia Pr. (§ 1 b).
— Microstegnus Pr. (§ 1 a).
1848. Notophoria Pr. (Hemitelia §). (§ 1 d).
1850-52. Hemithelia § Euhemithelia Fée. (§ 2).
— Hemistegia Fée (Pr. 1848 emend). (§ 1).
(1854. Cormophyllum Newm).
1857. Amphicosmia Moore (Gardn. 1842 emend). (§ 2).
— Chlamydia Moore (Amphicosmia §) (§ 2 a).
1898. Hemiteliella Diels (Hemitelia §) (§ 3).

§ 1. **Cnemidaria** (horrida).
Cnemidaria Pr. 1836 = Hemistegia Fée 1850-52 = Hemitelia § Euhemitelia Hk. 1865.
a. Microstegnus Pr. 1848 (grandifolia [Willd.]).
b. Hemistegia Pr. 1848 (grandifolia Hk.).
c. Actinophlebia Pr. 1848 (horrida).
d. Eleutheria Kze. 1844 (speciosa) = § Notophoria Pr. 1848.

§ 2. **Euhemitelia** Fée 1850-52 (multiflora) (non § Euhemitelia Hk. 1865).
Amphicosmia Moore 1857 = Hemitelia § Amphicosmia Hk. 1865. **Diels** 1899.
a. Chlamydia Moore 1857 (Walkeræ).

§ 3. **Amphicosmia** (capensis).
Hemitelia Pr. 1836 = Amphicosmia Gardn. 1842 = Alsophila § Hymenostegia pt. J. Sm. 1842 = Cyathea sp. Christ 1877 = Hemitelia § Hemiteliella **Diels** 1899.

7. **ALSOPHILA** R. Brown 1810. — Species: 185.

Typ. Polypodium extensum Forst.

1753. Polypodium sp. L.
1801. Cyathea sp. Sw.
1810. Alsophila R.Br.
1822. Trichipteris Pr. (§ 1 b).
1824. Chnoophora Klf. (§ 1 e).
1828. Gymnosphaera Bl. (§ 3 a).
1834. Chnoophora Mart. (non Klf. 1824 Alsophila §). (§ 1 b).
— Dicranophlebia Mart. (Alsophila §). (§ 1 d).
— Haplophlebia Mart. (Alsophila §). (§ 1 c).
— Amphidesmium Schott (§ 1 a).
1834. Trichopteris Schott (Trichipteris Pr. 1822 . (§ 1 b).
1836. Metaxya Pr. (§ 1 a).
1842. Hymenostegia pt. J. Sm. (Alsophila §).
— Trichostegia J. Sm. (Alsophila §).
1848. Dichorexia Pr. (§ 3 b).
— Lophosoria Pr. (§ 1 f).
1849. Trichosorus Liebm. (§ 1 f).
1858. Alsophilopsis Karst. (Alsophila §). (§ 1 g).
1873. Fourniera Bomm. (§ 3 c).

§ 1. **Species americanae.**

a. Amphidesmium Schott 1834 (blechnoides) = Metaxya Pr. 1836.
b. Trichopteris Pr. Schott 1834 = Trichipteris Pr. 1822 (elegans) = Alsophila § Chnoophora Mart. 1834.
c. Haplophlebia Mart. 1834 (sp. venis simplicibus).
d. Dicranophlebia Mart. 1834 (sp. venis furcatis).
e. Chnoophora Klf. 1824 (villosa).
f. Lophosoria Pr. 1848 (quadripinnata) = Trichosorus Liebm. 1849.
g. Alsophilopsis Karst. (sp. pinnis pinnulisque articulatis).
- Hymenostegia pt. J. Sm. 1842 (sp. soro indusio minuto instructo; Ex. aspera).
- Trichostegia J. Sm. 1842 (sp. receptaculo pilis articulatis instructo).

§ 2. **Species africanae.**

§ 3. **Species asiaticæ et oceanicae.**

a. Gymnosphaera Bl. 1828 (glabra).
b. Dichorexia Pr. 1848 (latebrosa).
c. Fourniera Bomm. 1873. (sp. soro indusio e squamis laciniatis patulis constituto instructo: novae caledoniae) = ? Alsophila § Hymenostegia pt. J. Sm. 1842.

III. POLYPODIACEAE.

I. WOODSIEAE.

1. WOODSIINAE.

1. **DIACALPE** Blume 1828. — Species: 1.

Typ. D. aspidioides Bl.

2. PERANEMA Don 1825. — Species: 1.

Typ. P. cyatheoides Don.

1828. Sphaeropteris Wall. (non Bernh. 1801). Hk. 1844. Hk. et Bak. 1866.
1830. Podeilema R.Br.
1845. Nematopera Kze. (corr.).

3. WOODSIA R. Brown 1810. 1815. — Species: 25.

Typ. Acrostichum ilvense L.

1753. Acrostichum sp. L.
1789. Polypodium sp. Vill.
1806. Aspidium sp. Sw. 1806.
1810. Woodsia R.Br.
1829. Physematium Klf. (§ 1).
1831. Hymenocystis C. A. Mey. (§ 1).
1836. Phegopteris pt. Pr. (Polypodium § 2 b). (§ 2).
1844. Perrinia Hk. (Woodsia §) (§ 1).
1875. Diacalpe sp. Trev. (§ 1).

§ 1. **Physematium** (mollis).
Physematium Klf. 1829 = Diacalpe sp. Trev. 1875.
— Hymenocystis C. A. Mey. 1831 (fragilis).
— § Perrinia Hk. 1844 (obtusa).

§ 2. **Euwoodsia** (ilvensis).

4. HYPODERRIS R. Brown 1830. — Species: 3.

Typ. H. Brownii J. Sm.

1856. Woodsia sp. Mett.

5. CYSTOPTERIS Bernhardi 1806. — Species: 13.

Typ. Polypodium fragile L.

1753. Polypodium sp. L.
1757. ? Filix Ludwig.
1793. Cyathea sp. Sm.
1801. Aspidium sp. Sw.
1804. Athyrium sp. Spr.
1806. Cystopteris Bernh.
1821. Cyclopteris Gray.
1828. Cystea Sm.
1900. Filix Underw.

6. ACROPHORUS Presl. 1836. — Species: 1.

Typ. Aspidium nodosum Bl.

Davallia sp. Bak. 1867. (Vide Syn. in Ind. spec.).

2. ONOCLEINEAE.

7. MATTEUCCIA Todaro 1866. — Species: 3.

Typ. Osmunda Struthiopteris L.

1753. Osmunda sp. L.
1795. Onoclea sp. Hoffm. Hk. Bak. 1866.
1809. Struthiopteris Willd. (non Weis 1770 nec Bernh. 1801).
1866. Matteuccia Tod.
1891. Pterinodes sp. O.Ktze.

8. **ONOCLEA** Linné 1753. — Species: 1.

Typ. O. sensibilis L.

1753. Onoclea L. 1751.
1763. Angiopteris Adans. (Mitch. 1748).
1802. Calypterium Bernh.
1803. Riedlea Mirb.
1836. Ragiopteris Pr.

II. ASPIDIEAE.

1. ASPIDIINAE.

9. **DRYOPTERIS** Adanson 1763, emend. — Species: 741.

Typ. Polypodium filix mas L.

1753. Polypodium sp. L.
1763. Dryopteris Adans.
1790. Gleichenia Necker.
— Psidopodium Necker.
1791. Meniscium Schreber (§ 7).
1800. Polystichum sp. Roth.
1801. Aspidium sp. Sw.
— Nephrodium sp. Rich. **Diels** 1899.
— Tectaria sp. Cav.
1824. Lastrea Bory (§ 1 e et 2).
1828. Stegnogramma Bl. (§ 6).
— Arthrobotrys Wall. (§ 1 f).
1833. Cyclosorus Link (§ 4).
— Hypodematium Kze. (§ 1 k).
1834. Dryopteris Schott (§ 1).
— Nephrodium Schott (§ 4).
— Thelypteris Schott (non Adans. 1763). (§ 1 e).
1836. Grammitis § Eugrammitis Pr. (non Sw. 1801). (§ 3).
— Goniopteris Pr. (§ 5).
— Monogonia Pr. (§ 5).
— Lastrea Pr. §§ Dryopteris, Thelypteris, Arthrobotrys (§ 1).
— Phegopteris Pr. (Polypodium §). (§ 2).
1838. Pleurogramma R.Br. (non Bl. 1828). (§ 3).
1841. Dichasium A.Br. (§ 1 f).
1842. Leptogramma J.Sm. (§ 3).
1843. Abacopteris Fée (§ 4 a).
— Amauropelta Kze. (§ 1 c).
1847. Arsenopteris Webb et Berth. (§ 1 et 4).
1848. Ampelopteris Kze. (§ 2).
— Glaphyropteris Pr. (§ 2).
1849. Anisocampium Pr. (§ 4 b).
— Haplodictyum Pr. (§ 4 a).
— Lastreastrum Pr. (Lastrea §) (§ 1 f).
— Pronephrium Pr. (§ 4 a).
1850-52. Aspidium Fée (§ 1).
— Camptodium Fée (§ 1 a).
— Dichasium (A.Br. 1841) Fée (§ 1 f).
— Gymnogramme Fée (non Desv. 1811). (§ 3).
— Oochlamys Fée (§ 1 d).
— Phegopteris (Pr. 1836) Fée. Mett. 1856. (§ 2).
1851. Oligocampia Trev. (§ 4 b).
— Gymnothalamium Zenker (§ 1 i).
— Gymnocarpium Newm. (§ 2).
Hemestheum Newm. (§ 1 e).
— Lophodium Newm. (§ 1 g).
1854. Syneuron J. Sm. (§ 6).
1855. Pycnopteris Moore (§ 1 b).
1857. Eleutherophlebia Gris. (Aspidium §). (§ 1).
— Anopodium J. Sm. (Phegopteris §). (§ 2).
— Desmopodium J. Sm. (§ 2).
— Catapodium J. Sm. (Phegopteris §). (§ 2).
— Oreopteris J.Sm. (Lastrea §) (§ 1d).
— Pachyderis J. Sm. (§ 1).
— Diclisodon Moore (§ 1 h).
— Monophlebia Moore (Lastrea §) (§ 1 d).
— Tectaria Moore (Nephrodium §) (§ 4).
1866. Microdictyon Fée (Goniopteris §) (§ 5).

1873. Parapolystichum Klys. (Polystichum §). (§ 1).

1875. Megalastrum J. Sm. (Lastrea §) (§ 1 i).

1875. Polystichopsis pt. J. Sm. (Lastrea §). (§ 1 j).

§ 1. **Eudryopteris** (filix mas).

Dryopteris Adans. 1763. Schott 1834 = Lastrea Pr. 1836 (Bory 1824 pt.) = Aspidium Fée 1850-52 (Sw. 1801 pt.) = Aspidium § Eleutherophlebia Gris. 1857 = Nephrodium § Lastrea Hk. 1862. Bak. 1867. **Diels** 1899 pt. = Aspidium § Lastrea Christ 1897 = Aspidium § Arsenopteris pt. Webb et Berth. 1847. — Polystichum § Parapolystichum Keys. 1873 (effusa, decomposita).

a. Camptodium Fée 1850-52 (pedata) = Pachyderis J. Sm. 1857.
b. Pycnopteris Moore 1855 (Sieboldii).
c. Amauropelta Kze. 1843 (limbata).
d. Monophlebia Moore 1857 (sp. ven. simplic.) = Lastrea § Dryopteris Pr. 1836 = Lastrea § Oreopteris J. Sm. 1857. — Oochlamys Fée 1850-52 (opposita).
e. Hemestheum Newm. 1851 (thelypteris) = Lastrea pt. Bory 1834 = Thelypteris Schott 1834.
f. Filix mas (filix mas) — Arthrobotrys Wall. 1828. Dichasium (A.Br. 1841) Fée 1850-52. — Lastrea § Lastreastrum Pr. 1849.
g. Lophodium Newm. 1851 (spinulosa) = Lastrea § Arthrobotrys Pr. 1836.
h. Diclisodon Moore 1857 (deparioides).
i. Megalastrum J. Sm. 1875 (villosa) — ? Gymnothalamium Zenker 1851 (scabrosa).
j. Polystichopsis pt. J. Sm. 1875 (pubescens).
k. Hypodematium Kze. 1833 (crenata).

§ 2. **Phegopteris** (phegopteris).

Lastrea pt. Bory 1824 = Polypodium § Phegopteris Pr. 1836. Hk. 1862 = Phegopteris Fée 1850-52 = Gymnocarpium Newm. 1851 = Polypodium § Euphegopteris Bak. 1867 = Phegopteris § Euphegopteris Christ 1897. Desmopodium J. Sm. 1857. Phegopteris § Anopodium et Catapodium J. Sm. 1875 — Glaphyropteris Pr. 1848 (decussata) — Ampelopteris Kze. 1848 (prolifera?).

§ 3. **Leptogramma** (totta)

Grammitis § Eugrammitis Pr. 1836 = Pleurogramma R.Br. 1838 = Leptogramma J. Sm. 1841 = Gymnogramme Fée 1850-52 = Grammitis § Leptogramma Moore 1857 = Phegopteris sp. Mett 1858. Christ 1897 = Gymnogramme § Eugymnogramme pt. Hk. 1864 = Gymnogramme § Leptogramme Bak. 1868.

§ 4. **Cyclosorus** (gongylodes).

Cyclosorus Link 1833 = Nephrodium Schott 1834 (non Rich. 1801) = Nephrodium § Eunephrodium Hk. 1862. Bak. 1867. **Diels** 1899 = Aspidium § Nephrodium Christ 1897. Aspidium § Arsenopteris pt. Webb et Berth. 1847.

a. Abacopteris Fée 1843 (lineata) = Pronephrium Pr. 1849. Haplodictyum Pr. 1849.
b. Anisocampium Pr. 1849 (otaria) = Oligocampia Trev. 1851.

§ 5. **Goniopteris** (vivipara).

Goniopteris Pr. 1836 = Polypodium § Goniopteris Hk. 1864. Bak. 1867 = Phegopteris § Goniopteris Christ 1897 = Nephrodium § Goniopteris **Diels** 1899. Goniopteris § Microdictyon Fée 1866 (Ghiesbreghtii) — Monogonia Pr. 1836.

§ 6. **Stegnogramma** (stegnogramme).

Stegnogramma Bl. 1828 = Gymnogramme § Stegnogramme Hk. 1864. Bak. 1868 = Syneuron J. Sm. 1854.

§ 7. **Meniscium** (reticulata).

Meniscium Schreb. 1791. Bak. 1868 = Phegopteris § Meniscium Christ 1897 = Nephrodium § Meniscium **Diels** 1899.

10. LUERSSENIA Kuhn 1882. — Species: 1.

Typ. Aspidium Kehdingianum Luerss.

Aspidium § Luerssenia Christ 1897.

11. FADYENIA Hooker 1840. — Species: 1.

Typ. Asplenium proliferum Sw.

Aspidium § Fadyenia Christ 1897.

12. MESOCHLAENA R. Brown 1838. — Species: 1.

Typ. Aspidium polycarpum Bl.

1839. Sphaerostephanos J. Sm.
1850-52. Stegnogramme pt. Fée.
1867. Didymochlaena sp. Bak.
1897. Aspidium § Mesochlaena Christ 1897.

13. DIDYMOCHLAENA Desvaux 1811. — Species: 1.

Typ. Adiantum lunulatum Houtt.

1824. Tegularia Reinw.
1827. Hippodium Gaud. msc.
— Monochlaena Gaud.
1850-52. Hysterocarpus Langsd. msc. ap. Fée.

14. CYCLOPELTIS J. Smith 1846. — Species: 2.

Typ. Polypodium semicordatum Sw.

1801. Aspidium sp. Sw. *Bak.* 1867. *Christ* 1897.
1836. Lastrea § 3 Arthrobotrys B. pt. Pr. (non Arthrobotrys Wall. 1828).
1846. Cyclopeltis J. Sm.
1850-52. Hemicardion Fée.
1857. Polystichum § Cyclopeltis Moore.

15. ASPIDIUM Swartz 1801 reduct. — Species: 138.

Typ. Polypodium trifoliatum L.

1753. Polypodium sp. L.
1799. *Tectaria* Cav. (nom. opt.). (§ 1).
1801. Aspidium sp. Sw.
1834. Aspidium Schott (§ 1).
1836. Bathmium Pr. (Aspidium §). (§ 1).
1836. Dictyopteris Pr. (non Lamour 1809). (§ 3).
— Digrammaria Pr. (§ 2 g).
— Pleocnemia Pr. (§ 4).
— Sagenia Pr. (§ 2).

1841. Bathmium Link (non Pr. 1836) (§ 1).
1849. Heterogonium Pr. (§ 2 g).
— Microbrochis Pr. (§ 2 a).
— Polydictyum Pr. (§ 2 b).
— Proferea Pr. (§ 4).
1850-52. Bathmium Fée (Link 1841 pt.) (§ 1).
— Cardiochlaena Fée (§ 2 e).
— Dryomenis Fée (§ 2 c).
— Phlebiogonium Fée (§ 2 d).
— Podopeltis Fée (§ 1).
1852. Deparia § Trichiocarpa Hk. (§ 2 f).
— Cionidium Moore. (§ 2 f).
1855. Dictyocline Moore (§ 2 h).
1856. Trichiocarpa (Hk. 1852) J. Sm. (§ 2 f).
1857. Dictyophlebia Gris. (Aspidium §). (§ 1).
— Patanema J. Sm. (§ 2 f).
1858. Phegopteris sp. Mett. (§ 3).
1862. Nephrodium § Pleocnemia Hk. (§ 4).
1864. Hemionitis sp. Hk. (§ 2 h).
1866. Grammatosorus Regel. (§ 2 i).
1867. Melanopteris J. Sm. (§ 1).
— Nephrodium § Pleocnemia et Sagenia Bak. (§ 2 et 4).
— Polypodium § Dictyopteris Bak. (§ 3).
1868. Gymnogramme § Digrammaria Bak. (§ 2 g).
1891. Depariopsis Bak. (Polypodium §) (§ 3).
1903. Arcypteris Underw. (§ 3).

§ 1. **Tectaria** (trifoliatum).

Tectaria Cav. 1799 = Aspidium Schott 1834 = Bathmium Link 1841. Fée 1850-52 = Aspidium § Dictyophlebia Gris. 1857 = Aspidium § Euaspidium Hk. 1862 pt. Bak. 1867, Diels 1899 pt. — Aspidium § Bathmium Pr. 1836 (singaporianum) = Podopeltis Fée 1850-52 — Melanopteris J. Sm. 1867 (plantaginea).

§ 2. **Sagenia** (?).

Sagenia Pr. 1836 = Aspidium § Euaspidium pt. Hk. 1862. **Diels** 1899 = Nephrodium § Sagenia Bak. 1867 = Aspidium § Sagenia pt. Christ 1897.

a. Microbrochis Pr. 1849 (cicutarium * apiifolium).
b. Polydictyum Pr. 1849 (menyanthidis).
c. Dryomenis Fée 1850-52 (siifolium).
d. Phlebiogonium Fée 1850 52 (brachiatum).
e. Cardiochlaena Fée 1850-52 (martinicense).
f. Cionidium Moore 1852 (Moorei) = Deparia § Trichiocarpa Hk. 1852 = Trichiocarpa J. Sm. 1856 = Patanema J. Sm. 1857 = Deparia sp. Bak. 1866.
g. Digrammaria Pr. 1836 (ambigua) = Heterogonium Pr. 1849 = Gymnogramme § Digrammaria Bak. 1868.
h. Dictyocline Moore 1855 (Griffithii) = Hemionitis sp. Hk. 1864. Bak. 1868.
i. Grammatosorus Regel.

§ 3. **Arcypteris** (irregulare).

Dictyopteris Pr. 1836 pt = Polypodium § Dictyopteris Hk. 1864. Bak. 1867 = Phegopteris § Dictyopteris Christ 1897 = Arcypteris Und. 1903. — Polypodium § Depariopsis Bak. 1891 (depariopsis).

§ 4. **Pleocnemia** (Leuzeanum).

Pleocnemia Pr. 1836 = Nephrodium § Pleocnemia Hk. 1862. Bak. 1867. Aspidium § Pleocnemia Christ 1897. **Diels** 1899 — Proferea Pr. 1849 (excellens).

16. NEOCHEIROPTERIS Christ 1905. — Species: 1.

Typ. Polypodium palmatopedatum Bak.

1898. Cheiropteris Christ 1898. **Diels** 1899 (non Kurr, gen. palaeophyt.).

16a. ADENODERRIS J. Smith 1875, Maxon 1905. — Species: 2.

Typ. Aspidium glandulosum Hk. et Grev. Vide Ind. spec. pag. 660.

Polystichum sp. **Diels** 1899.

17. POLYSTICHUM Roth 1799, 1800 emend. Schott 1834. — Species: 112.

Typ. Polypodium aculeatum L.

1753. Polypodium sp. L.
1800. Polystichum sp. Roth.
1801. Aspidium sp. Sw.
— Tectaria sp. Cav.
1803. Hypopeltis Michx.; Bory 1833 (nom. opt.?). (§ 1).
1819. Rumohra Raddi (§ 1 b).
1828. ? Arachniodes Bl. (§ 1 b).
1833. Tectaria Link (§ 1 b).
1834. Polystichum Schott (§ 1).
1836. Cyrtomium Pr. (§ 3).
(1841). Hemigonum J. Sm. 1875. (§ 3).
1850-52. Peltochlaena Fée (§ 3 a).
1858. Phegopteris sp. Mett.
1862. Aspidium § Polystichum Hk.
1863. Cyrtomiphlebium Hk. (Polypodium §). (§ 3 b).
1865. Porpaea Pr. msc. (§ 1 a).
1875. Polystichopsis pt. J. Sm. (Lastrea §). (§ 1 c).
1884. Ptilopteris Hance (§ 2).

§ 1. **Eupolystichum** (aculeatum).

Hypopeltis Michx. 1803 = Polystichum Schott 1834 (Roth 1800 pt.) = Aspidium § Polystichum Hk. 1862. Bak. 1867. Christ 1897.

a. Porpaea Pr. 1865 (vestitum).

b. Rumohra Raddi 1819 (adiantiforme) = Tectaria Link 1833. ? Arachniodes Bl. 1828.

c. Lastreopsis (hispida) = Lastrea § Polystichopsis pt. J. Sm. 1875.

§ 2. **Ptilopteris** (Maximowiczii).

Ptilopteris Hance 1884 = Phegopteris § Ptilopteris Christ 1897.

§ 3. **Cyrtomium** (falcatum).

Cyrtomium Pr. 1836 = Hemigonum J. Sm. (1841) = Aspidium § Cyrtomium pt. Hk. 1862. Bak. 1867. Christ 1897 = Polystichum § Cyrtomium pt. **Diels** 1899.

a. ? Peltochlaena Fée 1850-52 (? abbreviatum).

b. Cyrtomiphlebium (dubium) = Polypodium § Cyrtomiphlebium Hk. 1863. Bak. 1867.

18. PHANEROPHLEBIA Presl 1836. Und. 1902. — Species: 10.

Typ. Polypodium juglandifolium H.B. Willd.

Amblia Pr. 1836. = Aspidium § Cyrtomium pt. Hk. 1862. Bak. 1867. Christ 1897 = Polystichum § Cyrtomium pt. **Diels** 1899.

19. PLECOSORUS Fée 1850-52. — Species: 3.

Typ. Cheilanthes speciosissimus A.Br.

Cryptostigma A.Br. msc. 1856.
Cheilanthes sp. Bak. 1867.

20. CYCLODIUM Presl 1836. — Species: 2.

Typ. Aspidium meniscioides Willd.

Aspidium § Cyclodium Hk. 1862. Bak. 1867 (pt.). Christ 1897.

21. POLYBOTRYA Humb. et Bonpl. Willdenow 1810. — Species: 30.

Typ. P. osmundacea H.B. Willd.

1753. Osmunda sp. L.
1806. Acrostichum sp. Sw.
1810. Polybotrya H.B. Willd.
1819. Olfersia Raddi (§ 5).
1824. Gymnogramma sp. Klf. (§ 1).
1834. Egenolfia Schott (§ 1).
1845. Soromanes Fée (§ 6).
— Ectoneura Fée (Polybotrya §) (§ 1).
1846. Lacaussadea Gaud. (§ 1).
1846. Botryothallus Kl. (§ 6).
1849. Dorcapteris Pr. (§ 5).
— Psomiocarpa Pr. (§ 2).
1850-52. Granulina Bory. Fée (Egenolfia §). (§ 1)
1851. Amphisoria Trevis. (§ 2).
1854. Lomagramma Brack. (non J. Sm. 1842). (§ 4).
1869. Teratophyllum sp. Mett. (§ 4).
1875. Arthrobotrya J. Sm. (§ 4).

§ 1. **Egenolfia** (appendiculata).

Egenolfia Schott 1834 = Polybotrya § Ectonevra Fée 1845 = Lacaussadea Gaud. 1846 = Acrostichum § Egenolfia Hk. 1864. Bak. 1868. Egenolfia § Granulina Bory; Féc 1850-52.

§ 2. **Psomiocarpa** (apiifolia).

Psomiocarpa Pr. 1849 = Amphisoria Trev. 1851.

§ 3. **Eupolybotrya** (osmundacea).

Acrostichum § Polybotrya Hk. 1864. Bak. 1868.

§ 4. **Arthrobotrya** (articulata).

Lomagramma Brack. 1854 = Teratophyllum Mett. 1869 pt. = Arthrobotrya J. Sm. 1875 = Polybotrya § Teratophyllum Christ 1897. **Diels** 1899.

§ 5. **Olfersia** (cervina).

Olfersia Raddi 1819 = Dorcapteris Pr. 1849 = Acrostichum § Olfersia Bak. 1868.

§ 6. **Soromanes** (serratifolia).

Soromanes Fée 1845 = Botryothallus Kl. 1846 = Acrostichum § Soromanes Bak. 1868.

22. STENOSEMIA Presl 1836. — Species: 2.

Typ. Acrostichum auritum Sw.

Polybotrya sp. Bl. 1828 = Acrostichum § Stenosemia Hk. 1864. Bak. 1868 = Gymnopteris § Stenosemia Christ 1897.

23. LEPTOCHILUS Kaulfuss 1824. — Species: 64.

Typ. Acrostichnm axillare Cav.

1775. Polypodium sp. Aublet.
1781. Acrostichum sp. L.fil. *Bak.* 1868.
1783. Ophioglossum sp. Houtt.
1789. Osmunda sp. Jacq.
1806. Gymnopteris sp. Bernh. (non 1799).
1824. Leptochilus Klf. (§ 1).
(1827. Poikilopteris Eschweiler (§ 3).
1834. Bolbitis Schott (§ 3).
1836. Anapausia Pr. (Gymnopteris §) (§ 2).

1836. Campium Pr. (§ 3 a).
— Gymnopteris Pr. (non Bernh. 1799) **Diels 1899.**
— Poecilopteris Pr. (§ 3).
1840. Jenkinsia Hk. (§ 3 b).
1841. Cyrtogonium J. Sm. (§ 3).
— Lomagramma J. Sm. (§ 4).
1845. Cheilolepton Fée (§ 4).
— Gymnopteris Fée (§ 2).
— Heteronevron Fée (§ 3).
1845. Leptochilus Fée (§ 1).
1849. Anapausia Pr. (1836 §). (§ 2).
— Dendroglossa Pr. (§ 1).
— Euryostichum Pr. (Anapausia §) (§ 2).
1855. Chorizopteris Moore (§ 4).
1856. Chrysodium sp. Mett. (non Fée 1845).
1857. Neurocallis § Cheilolepton et Chorizopteris (§ 4).

§ 1. **Euleptochilus** (axillaris).

Leptochilus Klf. 1824 = Gymnopteris § Gymnopteris B. Pr. 1836 = Acrostichum § Gymnopteris A. B. pt. et § Chrysodium A. pt. Bak. 1868.

— Dendroglossa Pr. 1849 (zeylanicus).

§ 2. **Anapausia** (alienus).

Gymnopteris § Anapausia Pr. 1836 = Gymnopteris Fée 1845 = Anapausia Pr. 1849 = Acrostichum § Gymnopteris C. pt. Bak. 1868 — Anapausia § Euryostichum Pr. 1849 (nicotianifolius).

§ 3. **Bolbitis** (serratifolius).

(Poikilopteris Eschw. 1827) = Bolbitis Schott 1834 = Poecilopteris Pr. 1836 = Cyrtogonium J. Sm. 1841 = Heteronevron Fée 1845 = Acrostichum § Gymnopteris pt. et § Chrysodium pt. Bak. 1868.

a. Campium Pr. 1836 (virens).

b. Jenkinsia Hk. 1840 (scalpturatus f. monstr.).

§ 4. **Lomagramma** (lomarioides) (an gen. propr.?).

Lomagramma J. Sm. 1841 = Cheilolepton Fée 1845 = Neurocallis § Cheilolepton Moore 1857 — Chorizopteris Moore 1855 = Neurocallis § Chorizopteris Moore 1857 (f. monstr.).

II. 2. DIPTERIDINAE.

24. DIPTERIS Reinwardt 1824. — Species: 6.

Typ. D. conjugata Reinw.

Polypodium § Phymatodes sect. Dipteris Bak. 1868.

III. OLEANDREAE.

25. OLEANDRA Cavanilles 1799. — Species: 10.

Typ. O. neriiformis Cav.

1801. Aspidium sp. Sw.
1824. Ophiopteris Reinw.
1825. Neuronia Don.
1857. Haplophlebia Gris. (Aspidium §)

IV. DAVALLIEAE.

26. ARTHROPTERIS J. Smith 1854. — Species: 4.

Typ. Polypodium tenellum Forst.

Polypodium et Nephrolepis sp. Bak. 1867. Christ 1897. (vide Ind. spec.).

27. NEPHROLEPIS Schott 1834. — Species: 17.

Typ. Aspidium biserratum Sw.

1753. Polypodium sp. L.
1801. Aspidium sp. Sw.
1825. Nephrodium Pr. (non Rich. 1801).
1833. Hypopeltis sp. Bory (H. legitimae) (non Michx. 1803).
1834. Nephrolepis Schott.
1836. Leptopleuria Pr. (§ 4).
1842. Isoloma sp. J. Sm. (§ 1).
1850-52. Lepidonevron Fée.
1857. Cardiostegia Moore (Nephrolepis §). (§ 2).

§ 1. **Lindsayoideae** Christ 1897 (acutifolia).
Isoloma sp. J. Sm. 1842. Lindsaya sp. Bak. 1867.

§ 2. **Aspidioideae** Christ 1897 (biserrata).
Lepidonevron pt. Fée 1850-52 = Nephrolepis § Cardiostegia Moore 1857.

§ 3. **Davallioideae** Christ 1897 (acuminata).

§ 4. **Dicksonioideae** Christ 1897 (abrupta).
Leptopleuria Pr. 1836. Dicksonia sp. Bak. 1866.

28. HUMATA Cavanilles 1802. — Species: 14.

Typ. Adiantum repens L.fil.

1793. Davallia sp. Sm. Bak. 1867. Christ 1897.
1802. Humata Cav.
1808. Trichomanes sp. Poir.
1836. Pachypleuria Pr. (Davallia §).
1849. Pachypleuria Pr. (1836 §).
1850-52. Pteronevron Fée.

Humata Cav. 1802 = Davallia § Pachypleuria Pr. 1836 = Davallia § Humata Hk. 1846. Bak. 1867. Christ 1897 = Pachypleuria Pr. 1849 = Humata § Euhumata **Diels** 1899.

— Pteronevron Fée 1850-52 (Gaimardiana).

29. SACCOLOMA Kaulfuss 1820. 1824, emend. Mett. 1861. — Species: 8.

Typ. S. elegans Klf.

1804. Davallia sp. Spr.
1820. Saccoloma Klf. (§ 1).
1827. Neuropteris Desv. (§ 1).
1836. Microlepia sp. Pr. Fée 1850-52 (§ 2).
— Cibotium sp. Pr.
1842. Cystodium J. Sm. (§ 3).
1844. Dicksonia sp. Hk.
1849. Patania sp. Pr. (§ 2).
1873. Parasaccoloma Keys. (Saccoloma §). (§ 2).

§ 1. **Eusaccoloma** Keys. (Pol. Cyath. Hb. Bung. 4. 1873 (elegans).
Saccoloma Klf. 1824 = Neuropteris Desv. 1827 = Davallia sp. Bak. 1867.

§ 2. **Parasaccoloma** Keys. 1873 (inaequale).
Microlepia sp. Pr. 1836 = Davallia sp. Bak. 1867 et Dicksonia sp. Bak. 1866.

§ 3. **Cystodium** (sorbifolium).
Cystodium J. Sm. 1842 = Dicksonia sp. Bak. 1866.

30. DIELLIA Brackenridge 1854. — Species: 7.

Typ. D. falcata Brack. (?).

1857. Schizoloma sp. Moore.
1867. Lindsaea § Diellia Hk. Bak. Christ 1897.

31. LEPTOLEPIA Mettenius; Kuhn 1882. — Species: 2.

Typ. Davallia novae-zelandiae Col.

1846. Davallia § Leucostegia sp. Hk. Bak. 1867.
1854. Acrophorus sp. Moore.
1856. Microlepia sp. J. Sm. Christ 1897.

32. DAVALLIA Smith 1793 reduct. — Species: 67.

Typ. Trichomanes canariense L.

1753. Trichomanes sp. L.
1793. Davallia Sm.
1801. Wibelia Bernh. (§ 3).
1827. Humata sp. Desv. **Diels** 1899.
1836. Colposoria Pr. (Davallia §). (§ 3).
— Leucostegia Pr. (§ 4).
— Prosaptia Pr. (§ 1).
— Stenolobus Pr. (§ 3 et 4).
1842. Polypodium § Ctenopteris sp. J. Sm. (§ 1).
1849. Parestia Pr. (§ 3).
— ? Gymnogrammitis Griff. (? § 4).
— ? Stenochlamys Griff.
1850-52. Scyphularia Fée (§ 2).
1854. Acrophorus sp. Moore (§ 4).

§ 1. **Prosaptia** (contigua).
Prosaptia Pr. 1836 = Polypodium § Ctenopteris pt. J. Sm. 1842.

§ 2. **Scyphularia** (pentaphylla).
Stenolobus sp. Pr. 1836 = Scyphularia Fée 1850-52.

§ 3. **Eudavallia** Hk. 1846 (canariensis).
Davallia § Compositae **Diels** 1899 = Humata sp. Desv. 1827 — Davallia § Colposoria pt. Pr. 1836 — Wibelia Bernh. 1801 (denticulata) = Parestia Pr. 1849 — Stenolobus Pr. 1836 (solida).

§ 4. **Leucostegia** (immersa).
Leucostegia Pr. 1836 = Acrophorus sp. Moore 1854 = Humata sp. Mett. 1856 = Humata § Leucostegia **Diels** 1899 — ? Gymnogrammitis Griff. 1849.

33. MICROLEPIA Presl 1836 emend. — Species: 29.

Typ. Polypodium speluncae L.

1753. Polypodium sp. L.
1784. Trichomanes sp. Thbg.
1801. Dicksonia sp. Sw.
— Davallia sp Sw. R.Br. 1810.
1809(?) ? Scyphofilix Thouars. (nom. opt.?)
1827. Humata sp. Desv.
1836. Microlepia pt. Pr.
1837. Selenidium Kze. (Dicksonia ? §).
1850-52. Saccoloma sp. Fée.
1875. Scypholepia J. Sm.
1897. Wilfordia Christ (Davallia §).
1905. Davallodes Copeland (Microlepia §).

a. Davallodes Copeland 1905 (hirsuta).
b. Scypholepia J. Sm. 1875 (Hookeriana) = Saccoloma sp. Fée 1850-52.
c. Wilfordia Christ 1897 (Wilfordii).

34. ODONTOSORIA (Presl 1836) Fée 1850-52 emend. — Species: 19.

Typ. Trichomanes chinense L.

1753. Adiantum sp. L.
— Trichomanes sp. L.
1793. Davallia sp. Sm.
1836. Odontosoria Pr. (Davallia §).
1850-52. Odontosoria Fée (Pr. 1836 §).
— Stenoloma Fée.
1854. Acrophorus sp. Moore.
1856. Microlepia sp. Mett.
— Lindsaya sp. Mett. 1861.
— Odontoloma sp. Mett.
1873. Davalliastrum pt. Fourn. (Lindsaya §).
1882. Schizoloma sp. Kuhn.
— Lindsayopsis Kuhn.

§ 1. **Euodontosoria** (chinensis).

Davallia § Odontosoria Pr. 1836 = Stenoloma sp. Fée 1850-52 = Odontosoria J. Sm. 1854 (non Fée 1850-52) = Microlepia sp. Mett. 1856 = Lindsaya sp. Mett. 1861 = Davallia § Stenoloma pt. Bak. 1867 = Schizoloma sp. Kuhn 1882 = Lindsaya § Stenoloma pt. Christ 1897.

§ 2. **Stenoloma** (aculeata).

Stenoloma Fée 1850-52 (pt.) = Odontosoria Fée 1850-52 = Acrophorus sp. Moore 1854 = Lindsaya sp. Mett. 1861 = Davallia § Stenoloma pt. Bak. 1867 = Lindsayopsis Kuhn 1882 = Schizoloma sp. Kuhn 1882 = Lindsaya § Lindsayopsis Christ 1897.

35. TAPEINIDIUM (Presl 1849) C. Chr. 1906. — Species: 4.

Typ. Davallia pinnata Cav.

1802. Davallia sp. Cav. Bak. 1867.
1836. Saccoloma sp. Pr.
1842. Microlepia sp. J. Sm.
1849. Tapeinidium Pr. (Microlepia §).
1850-52. Wibelia Fée (non Bernh. 1801). Kuhn 1882. **Diels** 1899.
1869. Lindsaya sp. Mett.
1897. Davallia § Wibelia Christ.

36. DENNSTAEDTIA Bernhardi 1801. — Species: 57.

Typ. Trichomanes flaccidum Forst.

1786. Trichomanes sp. Forst.
1788. Polypodium sp. Sw.
1801. Dicksonia sp. Sw. *Bak.* 1866.
— Dennstaedtia Bernh. Moore 1857.
1827. Sitobolium Desv.
1836. Patania Pr.
1841. Adectum Link.
1842. Sitolobium J. Sm.
1844. Deparia sp. Hk. *Bak.* 1866.
1854. Litolobium Newm.
1866. Dicksonia § Patania *Bak.*

a. Dennstaedtia Bernh. 1801 (flaccida).
b. Sitobolium Desv. 1827 (punctilobula) = Adectum Link 1841 = Sitolobium J. Sm. 1842 (1841) = Litolobium Newm. 1854.
c. Patania Pr. 1836 (obtusifolia).

37. MONACHOSORUM Kunze 1848. — Species: 2.

Typ. Aspidium subdigitatum Bl.

Polypodium sp. Bak. 1867. (vide Ind. spec.).

38. SCHIZOLOMA Gaudichaud 1824. — Species: 15.

Typ. S. cordatum Gaud.

1775. Adiantum sp. Aublet.
1797. Lindsaea sp. Dry.
1824. Schizoloma Gaud.
1828. Pericoptis Wall. (nomen).
1841. Isoloma J. Sm. (§ 1).
1851-52. Schizolepton Fée (§ 2 a).
1873. Pellaeastrum pt. Fourn. (Lindsaya §).
1875. Guerinia J. Sm. (§ 2 b).

§ 1. **Isoloma** (divergens).

Isoloma J. Sm. 1841 = Lindsaya § Isoloma Sect * et ** Bak. 1867. Christ 1897.

§ 2. **Euschizoloma** (cordatum).

Schizoloma Gaud. 1824 = Lindsaya § Schizoloma Bak. 1867 (pt.) Christ 1897.

a. Schizolepton Fée 1851-52 (cordatum).
b. Guerinia J. Sm. 1875 (Guerinianum).

39. DICTYOXIPHIUM Hooker 1840. — Species: 1.

Typ. D. panamense Hk.

40. LINDSAYA Dryander 1793. 1797 reduct. — Species: 90.

Typ. L. trapeziformis Dry.

1763. Adiantum sp. L.
1793. Lindsaea sp. Dry.
1827. Hymenotomia Gaud. (§ 1).
1836. Saccoloma sp. Pr. (§ 3).
1837. Davallia sp. Kze. Hk. 1846 (§ 3).
1841. Synaphlebium J. Sm. (§ 4).
1850-52. Lindsaynium Fée (§ 1 b).
— Stenoloma sp. Fée (§ 2).
1854. Acrophorus sp. Moore (§ 3).
1873. Paralindsaya Keys. (Lindsaya §) (§ 1 a).
— Davalliastrum pt. Fourn. (Lindsaya §). (§ 2).
1875. Odontosoria sp. J. Sm. (§ 2).
1882. Schizoloma sp. Kuhn. (§ 2).

§ 1. **Eulindsaya** (lancea).

a. Paralindsaya Keys. 1873 (linearis).
b. Lindsaynium Fée 1850-52 (rigida).

§ 2. **Hymenotomia** (microphylla).

Hymenotomia Gaud. 1827 = Stenoloma sp. Fée 1850-52 = Lindsaya § Isoloma sect.*** pt. Bak. 1867 = Lindsaya § Davalliastrum (pt.) Fourn. 1873 = Odontosoria sp. J. Sm. 1875 — Schizoloma sp. Kuhn 1882 = Lindsaya § Stenoloma pt. Christ 1897.

§ 3. **Odontoloma** (repens).

Saccoloma sp. Pr. 1836 = Odontoloma J. Sm. 1841 = Acrophorus sp. Moore 1854 = Davallia § Odontoloma Bak. 1867.

§ 4. **Synaphlebium** (decomposita).

Synaphlebium J. Sm. 1841. Schizoloma sp. Moore 1857.

V. ASPLENIEAE.

V. 1. ASPLENIINAE.

41. ATHYRIUM Roth 1799. 1800. — Species: 86.

Typ. Polypodium filix femina L.

1753. Polypodium sp. L.
1799.1800. Athyrium Roth.
1801. Aspidium sp. Sw.
— Asplenium sp. Sw. Bernh. 1806.
1802. Tectaria sp. Cav.
1803. Nephrodium sp. Michx.
1810. Allantodia R.Br. (f).
1831. Deparia Hk. et Grev. (a).
1849. Brachysorus Pr. (e).
1850-52. Hypochlamys Fée (b).
1851. Solenopteris Zenker msc. (d).
— Pseudathyrium Newm. (c).
1856. Phegopteris sp. Mett.
1880. Pseudallantodia Clarke. (Asplenium §).

Asplenium § Athyrium Bak. 1867.

a. Deparia Hk. et Grev. 1831 (proliferum).
b. Hypochlamys Fée 1850-52 (costale).
c. Pseudathyrium Newm. 1851 (alpestre) = Phegopteris sp. Mett. 1856 = Polypodium sp. Bak. 1867.
d. Solenopteris Zenker 1859 (solenopteris).
e. Brachysorus Pr. 1849 (silvaticum).
f. Allantodia R.Br. 1810 (umbrosum) = Asplenium § Pseudallantodia Clarke 1880.

42. DIPLAZIUM Swartz 1801. — Species: 208.

Typ. Asplenium plantaginifolium L.

1759. Asplenium sp. L.
1801. Diplazium Sw.
1804. Callipteris Bory (§ 1).
1836. Anisogonium Pr. (§ 2).
— Hemidictyum Pr. (§ 3).
— Oxygonium pt. Pr. (§ 2 b).
1840. Digrammaria Hk. (non Pr. 1836) (§ 2 c).
1841. Callipteris J. Sm. (§ 2).
1842. Callipteris J. Sm. (§ 2 c).
1843. Pteriglyphis Fée 1850-52. (§ 2 a).
1847. Lotzea Kl. (§ 1 b).
1849. Microstegia Pr. (§ 2 c).
— Ochlogramma Pr. (§ 2 a).
1850-52. Callipteridastrum Fée (Diplazium §). (§ 1 a).
1857. Didymochlamys Moore (Diplazium §). (§ 1 c).
1866. Triblemma J. Sm. (Asplenium §) (§ 1 a).
1870. Athyrium §§§ Callipteris, Hemidictyum pt., Diplazium Milde.
1875. Micropodium sp. J. Sm. (§ 1 a).
1905. Allantodia sp. Racib.

§ 1. **Eudiplazium** (plantaginifolium).

Asplenium § Diplazium Bak. 1867 = Athyrium § Diplazium Milde 1870.

a. Callipteridastrum Fée 1850-52. Asplenium § Triblemma J. Sm. 1866 (lanceum) = Micropodium sp. J. Sm. 1875.
b. Lotzea Kl. 1847 (Klotzschii).
c. Didymochlamys Moore 1857 (hians).

§ 2. **Anisogonium** (fraxinifolium).

Anisogonium Pr. 1836 = Callipteris J. Sm. 1841 (Bory 1804 pt.) = Asplenium § Anisogonium Bak. 1867 = Diplazium § Calli-

pteris Christ 1897. **Diels** 1899. = Athyrium § Callipteris Milde 1870.

a. Pteriglyphis Fée 1843 = Ochlogramma Pr. 1849 (Cumingii).
b. Oxygonium Pr. 1836 (pt.) (cordifolium).
c. Microstegia Pr. 1849 (esculentum) = Digrammaria Hk. 1840 = Callipteris J. Sm. 1842.

§ 3. **Hemidictyum** (marginatum).

Hemidictyum Pr. 1836. Christ 1897 = Asplenium § Hemidictyum pt. Bak. 1867. = Athyrium § Hemidictyum Milde 1870 (pt.) = Allantodia sp. Racib. 1905.

43. DIPLAZIOPSIS C. Chr. 1906. — Species: 1.

Typ. Asplenium javanicum Bl.

Allantodia Wall. 1830. Bak. 1867. Christ 1897. **Diels** 1899 (non R.Br. 1810).

44. TRIPHLEBIA Baker 1886. — Species: 3.

Typ. Scolopendrium pinnatum J. Sm.

1825. Scolopendrium sp. Pr. Bak. 1867. Christ 1897.
1866. Micropodium sp. Mett.
1886. Triphlebia Bak.
1891. Phyllitis sp. O.Ktze.
Scolopendrium § Triphlebia Christ 1897.

45. DIPLORA Baker 1873. — Species: 2.

Typ. D. integrifolia Bak.

Scolopendrium § Diplora Christ 1897.

46. PHYLLITIS Ludwig 1757 emend. — Species: 9.

Typ. Asplenium Scolopendrium L.

1753. Asplenium sp. L.
1757. Phyllitis Ludwig (§ 1).
1763. Scolopendrium Adans. Sm. 1793.
1819. Glossopteris Rafin. (§ 1).
1836. Antigramma Pr. (§ 2).
— Hemidictyum sp. Pr. (§ 2).
1841. Camptosorus sp. Link. (§ 2).
1857. Schaffneria Fée (§ 3).

§ 1. **Euphyllitis** (scolopendrium).

Phyllitis Ludwig 1757 = Scolopendrium Adans. 1763 = Glossopteris Rafin. 1819 = Scolopendrium § Euscolopendrium Hk. 1862. Bak. 1867 (pt.) Christ 1897. **Diels** 1899.

§ 2. **Antigramma** (brasiliensis).

Antigramma Pr. 1836 = Camptosorus sp. Link 1841 = Scolopendrium § Antigramma Hk. 1862. Bak. 1867. Christ 1897. **Diels** 1899.

§ 3. **Schaffneria** (nigripes).

Schaffneria Fée 1857 = Scolopendrium § Schaffneria Hk. 1862. Bak. 1867. Christ 1897. **Diels** 1899 = Antigramma § Schaffneria J. Sm. 1875.

47. CAMPTOSORUS Link 1833. — Species: 2.

Typ. Asplenium rhizophyllum L.

1753. Asplenium sp. L.
1862. Scolopendrium §Camptosorus Hk. (Endl. 1840). Bak. 1867. Christ 1897. **Diels** 1899.
1875. Antigramma § Camptosorus J. Sm.
1891. Phyllitis sp. O.Ktze.

48. ASPLENIUM Linné 1753 reduct. — Species: 429.

Typ. ?

1753. Asplenium L.
1763. Ceterach pt. Adans.
1786. Caenopteris Berg. (§ 5).
— Trichomanes sp. Forst. (§ 6).
1789. Darea Juss. (§ 5).
1790. Onopteris Necker (§ 4 f).
1794. Phyllitis Moench (non Ludwig 1757). (§ 4).
1801. Davallia sp. Sw. (§ 6).
1803. Belvisia Mirb. (pt.?) (§ 3).
1833. Acropteris Link (§ 3).
1836. Thamnopteris Pr. (Asplenium §) (§ 1).
1841. Neottopteris J. Sm. (§ 1).
1844. Amesium Newm. (§ 3, 4).
1847. Homaloneuron Kl. (Asplenium §) (§ 4: sp. venis flabellatis).
1849. Tarachia Pr. (§ 4 d).
— Thamnopteris Pr. (1836 §). (§ 1).
1850-52. Acropteridastrum Fée (Asplenium §). (§ 4 d).
- Dareastrum Fée (Asplenium §) (§ 4 g).
— Hemionitidastrum Fée (Asplenium §). (§ 2).
— Neottopteridastrum Fée (Asplenium §). (§ 1 a).
1853. Loxoscaphe Moore (§ 6).
1854. Asplenidictyum J. Sm. 1875. (§ 4 h).
1856. Sphenopteris Mett. (Asplenium §) (§ 4 d).
1857. Hemidictyum § Asplenidictyum Moore (§ 4 h).
— Acropteris Moore (Asplenium §) (§ 4 d).
1866. Phyllitis J. Sm. (Asplenium §) (§ 1 a).
— Micropodium pt. Mett. (§ 1 b).
1873. Lobium Keys. (Asplenium §)(§4c).
— Microsplenium Keys. (Asplenium §). (§ 4 a).
— Parasplenium Keys. (Asplenium §) (§ 4 b).
1875. Asplenites J. Sm. (Asplenium §) (§ 4).
— Cheilosorum J. Sm. (Asplenium §) (§ 4 e).
— Hemionites J. Sm. (Asplenium §) (§ 2).
1877. Eremopodium Trev. (§ 1 b).
1901. Arcasplenium Moore (§ 4 i).

§ 1. **Thamnopteris** Pr. 1836 (nidus).
Neottopteris J. Sm. 1841 = Thamnopteris Pr. 1849 — Asplenium § Neottopteris **Diels** 1899.
a. Neottopteridastrum Fée 1850-52 (serratum) = Asplenium § Phyllitis J. Sm. 1866.
b. Micropodium pt. Mett. 1866 (vittaeforme) = Eremopodium Trev. 1877.

§ 2. **Hemionitidastrum** Fée 1850-52 (hemionitis).
Asplenium § Hemionites J. Sm. 1875.

§ 3. **Acropteris** (septentrionale).
Belvisia pt. Mirb. 1803 = Acropteris Link 1833 = Amesium sp. Newm. 1844.

§ 4. **Euasplenium** (?).

Asplenium § Euasplenium pt. Bak. 1867 = Asplenium § Asplenites J. Sm. 1875. Phyllitis Moench 1794.

a. Micrasplenium Keys. 1873 (trichomanes).
b. Parasplenium Keys. 1873 (monanthes).
c. Lobium Keys. 1873 (pumilum)
d. Tarachia pt. Pr. 1849 (praemorsum) = Asplenium § Acropteridastrum Fée 1850-52 = Asplenium § Sphenopteris Mett. 1856 = Asplenium § Acropteris pt. Moore 1857.
e. Cheilosorum J. Sm. 1875 (cheilosorum).
f. Onopteris Neck. 1790 (adiantum nigrum) = Tarachia sp. Pr. 1849.
g. Dareastrum Fée 1850-52 (cristatum).
h. Asplenidictyum J. Sm. 1854 (Purdicanum) = Hemidictyum § Asplenidictyum Moore 1857 = Asplenium § Hemidictyum pt. Bak. 1867.
i. Arcasplenium Moore 1901 (Sanderi).

§ 5. **Caenopteris** (achilleifolium).

Caenopteris Berg. 1786 = Darea Juss. 1789 = Asplenium § Darea Bak. 1867. Christ 1897. **Diels** 1899.

§ 6. **Loxoscaphe** (theciferum).

Loxoscaphe Moore 1853 = Davallia § Loxoscaphe Bak. 1867.

49. CETERACH Adanson 1763 emend. Kuhn 1879. — Species: 4.

Typ. Asplenium Ceterach L.

1753. Asplenium sp. L.
1763. Ceterach Adans. pt.
1798. Scolopendrium sp. Sym.
1805. Ceterach DC. (§ 2).
1806. Grammitis sp. Sw.
— Gymnopteris sp. Bernh.
1827. Gymnogramma sp. Spr.
1844. Notolepeum Newm (§ 2).
1857. Lepichroa Moore (Grammitis §). (§ 3).
1860. Hemidictyum pt. Hk. (Asplenium §). (§ 2).
1875. Ceterachopsis J. Sm. (Asplenium §). (§ 1).
1876. Hemidictyum sp. Bedd. (§ 2).
1879. Ceterach Kuhn.

§ 1. **Ceterachopsis** J. Sm. 1875 (Dalhousiae) = Asplenium sp. Bak. 1867. Christ 1897.

§ 2. **Euceterach** (officinarum).

Ceterach DC. 1805. Christ 1897 = Notolepeum Newm. 1844 = Asplenium § Hemidictyum pt. Bak. 1867.

§ 3. **Lepichroa** (cordatum).

Grammitis § Lepichroa Moore 1857 = Gymnogramme sp. Bak 1868. Christ 1897.

50. PLEUROSORUS Fée 1850-52. — Species: 3.

Typ. Gymnogramme papaverifolia Kze.

1827. Gymnogramma sp. Desv. Bak. 1867. Christ 1897.
1856. Ceterach sp. Mett.

V. 2. BLECHNINAE.

51. BLECHNUM Linné 1753 emend. Mett. 1856. — Species: 138.

Typ. B. occidentale (et orientale) L.

1753. Blechnum L. (§ 1).
— Osmunda sp. L. (§ 3).
1770. Struthiopteris Weis (§ 3 c).
1795. Onoclea sp. Hoffm. Sw. 1801. 1806. (§ 3).
1802. Asplenium sp. Bernh. (§ 3).
1809. Lomaria Willd. (§ 3).
1810. Stegania R.Br. (§ 3).
1841. Salpichlaena J. Sm. (§ 2).
1842. Stenochlaena sp. J. Sm. (§ 3). (filiforme).
1847. Salpiglaena Kl. (§ 2).
1849. Blechnopsis Pr. (§ 1 c).
— Diafnia Pr. (Blechnopsis §). (§ 1 d).
— Distaxia Pr. (§ 1 b).
— Lomaridium Pr. (§ 3 b).
— Mesothema Pr. (§ 1 a).
1849. Orthogramma Pr. (§ 1 e).
— Parablechnum Pr. (§ 1 e).
— Salpinchlaena Pr (§ 2).
— Spicanta Pr. (§ 3 c).
1850-52. Paralomaria Fée (Lomaria §). (§ 1 a et 3 e).
— Salpichlaena Fée (§ 2 et 1 a).
1851. Blechnopteris Trev. (§ 1 a et 3 d. e).
1860 (?) Blechnidium Moore (§ 1 a).
1869. Salpinchlaena Trev. (§ 2 et 1 c).
— Struthiopteris Trev. (§ 3 c. et 1 a).
1875. Lomariocycas J. Sm. (Lomaria §). (§ 1 e).
— Loxochlena J. Sm. (Lomaria §). (§ 1 d).
— Stegania J. Sm. (Lomaria §). (§ 1 a).
1891. Spicanta O.Ktze. 1891 (§ 1. 2. 3).

§ 1. **Eublechnum** (occidentale).
Blechnum L. 1753 = Blechnum § Eublechnum Bak. 1867 = Spicanta pt. O.Ktze. 1891.
a. Mesothema Pr. 1836 (pt.) (australe) = Lomaria § Paralomaria Fée 1850-52 (pt.) = Blechnopteris Trev. 1851 (pt.) = Struthiopteris Trev. 1869 (pt.).
b. Distaxia Pr. 1849 (fraxinea).
c. Blechnopsis Pr. 1849 (occidentale) = Salpichlaena pt. Fée 1850-52. Trev. 1869.
d. Diafnia Pr. 1849 (serrulatum).
e. Blechnidium Moore 1860 (melanopus).

§ 2. **Salpichlaena** (volubile).
Salpichlaena J. Sm. 1841 = Salpiglaena Kl. 1847 = Salpinchlaena Pr. 1849 = Blechnum § Salpichlaena Bak. 1867.

§ 3. **Lomaria** (spicant).
Osmunda sp. L. 1753 = (Struthiopteris Weis 1770) = Onoclea sp. Hoffm. Sw. 1801 = Lomaria Willd. 1809 = Stegania R.Br. 1810 = Lomaria § Eulomaria Bak. 1867 = Spicanta pt. O.Ktze. 1891.
a. Stegania R.Br. 1810 et emend. J. Sm. 1875 (Patersoni).
b. Lomaridium Pr. 1849 (Plumieri).
c. Struthiopteris Weis 1770. Trev. 1869 pt. (spicant) = Spicanta Pr. 1849.
d. Loxochlena J. Sm. 1875 (punctulatum) = Mesothema pt. Pr. 1849 = Blechnopteris pt. Trev. 1851.
e. Parablechnum Pr. 1849 (pt.) (capense) = Lomaria § Paralomaria (pt.) Fée 1850-52 = Blechnopteris pt. Trev. 1851 = Lomaria § Lomariocycas J. Sm. 1875. — Orthogramma Pr. 1849 (Gilliesii).

52. SADLERIA Kaulfuss 1824. — Species: 4.

Typ. S. cyatheoides Klf.

1827. Blechnum sp. Gaud.
1856. Woodwardia sp. Mett.

53. BRAINEA J. Smith 1856. — Species: 1.

Typ. Bowringia insignis Hk.

1853. Bowringia Hk. (non Benth. 1852 gen. Papilionac.).

54. STENOCHLAENA J. Smith 1841. 1842. — Species: 11.

Typ. Acrostichum sorbifolium L.

1753. Acrostichum sp. L. Bak. 1867.
1801. Onoclea sp. Sw.
1810. Lomaria sp. Willd.
1836. Olfersia sp. Pr.
? Leptophyllum Bl. (Olfersia §). (§ 1).
1841. Stenochlaena J. Sm. (§ 1).
1845. Lomariopsis Fée. Christ 1897. (§ 1).
1849. Cafraria Pr. (Stenochlaena §). (§ 2).
1850-52. Lomariobotrys Fée (§ 2).
1856. Polybotrya sp. Mett.
1866. Teratophyllum (pt.) Mett. (§ 1).
1871. Chrysodium sp. Luerss.
1875. Lomariopsis J. Sm. (§ 1).
— Stenochlaena J. Sm. (§ 2).

Acrostichum § Stenochlaena Bak. 1867.

§ 1. **Eustenochlaena** (sorbifolia).
Stenochlaena J. Sm. 1841 = Lomariopsis Fée 1845. J. Sm. 1875. — Leptophyllum Bl. (gracilis). Teratophyllum Mett. 1866 (f. tenuisectae S. sorbifoliaé et S. palustris).

§ 2. **Cafraria** Pr. 1849 (tenuifolia).
Lomariobotrys Fée 1850-52 = Stenochlaena J. Sm. 1875.

55. WOODWARDIA Smith 1793. — Species: 7.

Typ. Blechnum radicans L.

1753. Acrostichum sp. L.
1771. Blechnum sp. L.
1793. Woodwardia Sm.
1836. Doodia sp. Pr. (§ 1).
1849. Anchistea Pr. (§ 1).
— Lorinseria Pr. (§ 3).
1850-52. Woodwardia § Euwoodwardia Fée (§ 2).

§ 1. **Anchistea** (virginica).
Anchistea Pr. 1849.

§ 2. **Euwoodwardia** Fée 1850-52 (radicans).

§ 3. **Lorinseria** (areolata).
Lorinseria Pr. 1849.

56. DOODIA R. Brown 1810 (rectius Doodya). — Species: 6.

Typ. ?

1850-52. Woodwardia § Doodya Fée.

VI. PTERIDEAE.

VI. 1. GYMNOGRAMMINAE.

57. PTEROZONIUM Fée 1850-52. — Species: 2.

Typ. Gymnogramme reniformis Mart.

Gymnogramme § Eugymnogramme pt. Bak. 1868 = Gymnogramme § Pterozonium Christ 1897. Trichiogramme sp. Kuhn 1882.

58. SYNGRAMMA J. Smith 1845. — Species: 16.

Typ. S. alismifolia J. Sm.

1836. Oxygonium sp. Pr. J. Sm. 1842.
1845. Syngramma J. Sm.
1850-52. Callogramme Fée (§ 2).
1856. Gymnogramme sp. Mett.; Hk. 1864 (§ 1. 2. 3).
1857. Dictyogramma sp. Moore; Trev. 1877 (non Fée 1850-52). (§ 4).
1864. Hemionitis sp. Hk. (§ 4).
1873. Austrogramme (pt.) Fourn. (§ 1).
1875. Asplenitis J. Sm. (Gymnogramma §). (§ 1).
1877. Toxopteris Trev (§ 2).
1882. Trichiogramme Kuhn.
1897. Neurogramme § Trichiogramme Christ.

Syn. Trichiogramme Kuhn 1882.

§ 1. **Austrogramme** (marginata).
Austrogramme Fourn. 1873 (pt.) = Gymnogramme § Asplenitis J. Sm. 1875 = Gymnogramme § Eugymnogramme sp. Bak. 1868.

§ 2. **Toxopteris** (borneensis).
Toxopteris Trev. 1877 = Gymnogramme § Syngramme pt. Bak. 1868.

§ 3. **Callogramme** (alismifolia).
Callogramme Fée 1850-52 = Gymnogramme § Syngramme p Bak. 1868.

§ 4. (Hookeri).
Dictyogramme sp. Moore 1857 = Hemionitis sp. Bak. 1868. Christ 1897.

59. ANOGRAMMA Link 1841. — Species: 9.

Typ. Polypodium leptophyllum L.

1806. Grammitis sp. Sw.
1811. Gymnogramma sp. Desv.
1841. Anogramma Link.
1854. Dicranodium Newm.
1882. Gymnogramme Kuhn.

Gymnogramme § Eugymnogramme Bak. 1868 (pt.). Christ 1897.

60. GYMNOGRAMMA Desvaux 1811 reduct. — Species: 50.

Typ. Grammitis flexuosa Humb. et Bonpl.

1809. Grammitis sp. H. et B.
1810. Hemionitis sp. Willd. (§ 4).
1811. Gymnogramma Desv. (pt.).
1844. Jamesonia sp Kze.
1850-52. Anogramme sp. Fée.
1850-52. Ceropteris sp. Fée (§ 4).
— Eriosorus Fée (§ 1 b).
1866. Ampelogramma J. Sm. (Gymnogramma §). (§ 1 a).

1875. Pleurosorus pt. J. Sm. (Gymnogramma §).
1882. Psilogramme § Eupsilogramme Kuhn (§ 1 a).
1889. ?Cerosora Bak. (Gymnogramme §) (§ 5).
1899. Cerogramme Diels (Gymnogramme §). (§ 4).
1904. Isgnogramme Hieron. (Gymnogramme §). (§ 2).
— Definitifolium Hieron. (Gymnogramme §). (§ 3).

§ 1. **Eugymnogramme** Diels (flexuosa).

Gymnogramme § Eugymnogramme pt. Bak. 1868 = Psilogramme § Eupsilogramme Kuhn 1882 = Gymnogramme § Psilogramme Christ 1897.

a. Ampelogramma J. Sm. 1866 (flexuosa).
b. Eriosorus Fée 1850-52 (aureo-nitens).

§ 2. **Isgnogramme** Hieron. 1904 (Lehmannii).
§ 3. **Definitifolium** Hieron. 1904 (longipetiolata).
§ 4. **Cerogramme** Diels 1899 (sulphurea).

Hemionitis sp. Willd. 1810 = Ceropteris sp. Fée 1850-52 = Gymnogramme § Ceropteris pt. Bak. 1868. Christ 1897.

§ 5. ?**Cerosora** Bak. 1889 (chrysosora).

61. JAMESONIA Hook. et Grev. 1830. — Species: 14.

Typ. Pteris imbricata Cav.

1836. Allosorus sp. Pr.
1847. Gymnogramme § Jamesonia Kl.
1882. Psilogramme § Jamesonia Kuhn.

62. CONIOGRAMME Fée 1850-52. — Species: 2.

Typ. Gymnogramme javanica Bl.

1811. Gymnogramma sp. Desv. Bak. 1868.
(1849. Dyctiogramme Pr.
— Notogramme Pr.).
1850-52. Coniogramme Fée (fraxinea).
1850-52. Dictyogramme Fée (non Pr. 1849) (japonica).
1851. Neurosorus Trev. = Coniogramme.
1897. Neurogramme sp. Christ 1897.

63. HEMIONITIS Linné 1753 (reduct.). — Species: 8.

Typ. H. palmata L.

1833. Gymnogramma sp. Link.
1841. Schizogramma Link.
1875. Cardinitis J. Sm. (Hemionitis §).
1877. Dictyogramme sp. Trev.

Hemionitis § Euhemionitis pt. Bak. 1868.

a. Schizogramma Link 1833 (pedata).
b. Cardinitis J. Sm. 1875 (arifolia).

64. GYMNOPTERIS Bernhardi 1799. — Species: 12.

Typ. Pteris ruffa L.

1753. Pteris sp. L.
1786. Asplenium sp. Lam.
1799. Gymnopteris Bernh. (non Pr. 1836) (§ 1).
1801. Hemionitis sp. Sw. (§ 1).
1811. Gymnogramme sp. Desv.
1841. Neurogramma Link (§ 1). (Diels 1899).
1847. Stenogramme Kl. (Gymnogramme §). (§ 2).

1857. Syngramma sp. Moore (§ 3).
1866. Grammitis § Paraceterach F. Muell. (§ 3).
1872. Stegnogramme Fourn (Kl. 1847 § corr.) (non Bl. 1828). (§ 2).
1875. Sericonitis J. Sm. (Hemionitis §) (§ 3).
1876. Bommeria Fourn. 1880 (§ 2).
1877. Dictyogramme sp. Trev. (§ 2).

§ 1. **Neurogramma** (rufa).
Gymnopteris Bernh. 1799 = Neurogramma Link 1841 = Gymnogramme § Eugymnogramme pt. Bak. 1868 = Neurogramme § Euneurogramme (pt.) Christ 1897. **Diels** 1899.

§ 2. **Bommeria** (pedata).
Gymnogramme § Stenogramme Kl. 1847 = Stegnogramme Fourn. 1872 = Bommeria Fourn. 1876 = Gymnogramme § Eugymnogramme pt. et § Selliguea pt. Bak. 1868 = Gymnogramme § Bommeria Christ 1897. Dictyogramme sp. Trev. 1877.

§ 3. **Paraceterach** (Muelleri).
Syngramma sp. Moore 1857 = Grammitis § Paraceterach Muell. 1866 = Hemionitis § Sericonitis J. Sm. 1875 = Gymnogramme § Eugymnogramme pt. Bak. 1867 = Gymnogramme § Vestitae Christ 1897.

65. CEROPTERIS Link 1841. — Species: 7.

Typ. Acrostichum calomelanos L.

1753. Acrostichum sp. L.
1811. Gymnogramma sp. Desv.
1836. *Calomelanos* Pr. (Gymnogramme §) (nom. opt.!).
1841. Ceropteris Link.
1850-52. Argyria Fée (Ceropteris §).
— Chrysodia Fée (Ceropteris §).
1899. Neurogramme sp. **Diels.**

Gymnogramme § Ceropteris pt. Bak. 1868. Christ 1897.

66. TRISMERIA Fée 1850-52. — Species: 2.

Typ. Acrostichum trifoliatum L.

Gymnogramme § Ceropteris pt. Bak. 1868. Christ 1897.

VI. 2. CHEILANTHINAE.

67. PELLAEA Link 1841. — Species: 68.

Typ. Pteris atropurpurea L.

1753. Pteris sp. L.
1806. Cheilanthes sp. Sw.
1824. Cassebeera sp. Klf. **Diels** 1899 (§ 4).
— Gymnogramma sp. Klf. Mett. 1859 (§ 5).
1836. Allosorus Pr. p. p. max. (non Bernh. 1806).
1841. Pellaea Link (§ 2).
— Platyloma J. Sm. (§ 1).
— Argyrochosma J. Sm. (Notholaena §). (§ 5).
1841 Notholaena sp. J. Sm. (§ 5).
1850-52. Cincinalis Fée (non Gled. 1754 nec Desv. 1811). (§ 5).
1857. Synochlamys Fée (§ 2 a).
1858. Crypteris Nutt. msc. (§ 2 b).
1867. Holcochlaena Bak. (Pellaea §) (§ 6 a).
1875. Ormopteris J. Sm. (§ 4).
— Pellaeopsis J. Sm. (§ 6 a).
1876. Mildella Trev. (§ 3).
1879. Choristosoria Mett.; Kuhn (§ 8).

1879. Pteridella Mett: Kuhn (§ 6 et 7).
1882. Pteridellastrum Prantl (Pellaea §) (§ 7).
1891. Allosorus O.Ktze.
1891. Bakeriopteris sp. O.Ktze. (§ 4).
1896. Hymenoloma Dav. (Pellaea §). (§ 2 c).

§ 1. **Platyloma** (paradoxa).
Platyloma J. Sm. 1841.

§ 2. **Eupellaea** (atropurpurea).
Pellaea Link 1841 = Pellaea § Allosorus pt. Bak. 1868 pt. Christ 1897.
a. Synochlamys Fée 1857 (ambigua).
b. Crypteris Nutt. 1858 (andromedifolia).
c. Hymenoloma Dav. 1896 (membranacea).

§ 3. **Mildella** (intramarginalis).
Mildella Trev. 1876 = Pellaea § Pteridomorpha pt. Christ 1897.

§ 4. **Ormopteris** (gleichenioides).
Cassebeera sp. Klf.? 1824. Bak. 1867. Christ 1897. **Diels** 1899 = Bakeriopteris sp. O.Ktze. 1891.

§ 5. **Argyrochosma** (nivea).
Gymnogramme sp. Klf. 1824 = Notholaena § Argyrochosma J. Sm. 1841 = Cincinalis Fée 1850-52 = Notholaena § Cincinalis Bak. 1868 (pt.). Christ 1897 (pt.) = Pellaea § Cincinalis Prantl 1882. **Diels** 1899.

§ 6. **Pteridella** Prantl 1882 (Doniana).
Pellaea § Platyloma pt. Bak. 1867. Christ 1897 = Pteridella Mett. 1879.
a. Holcochlaena Bak. 1867 (angulosa) = Pellaeopsis J. Sm. 1875.

§ 7. **Pteridellastrum** Prantl 1882 (viridis).
Pellaea § Platyloma pt. Bak. 1867. Christ 1897 = Pteridella pt. Mett. 1879 = Pellaea § Pteridomorpha pt. Christ 1897.

§ 8. **Choristosoria** (pteroides).
Choristosoria Mett. 1879. Cheilanthes sp. Bak. 1867. Christ 1897.

68. DORYOPTERIS J. Smith 1841, 1842 emend. Diels 1899. — Species: 39.

Typ. Pteris pedata L.

1753. Pteris sp. L.
1824. Cassebeera pt. Klf. 1824 (typ.) **Diels** 1899 (non Dennst. 1818: gen. Melastomat.) (§ 1 a).
1836. Litobrochia sp. Pr. (§ 2).
1841. Doryopteris J. Sm. (§ 2).
1847. Lytoneuron Kl. (Doryopteris §). (§ 1 b).
1850-52. Pellaea sp. Fée (§ 1).
1857. ? Cheiloplecton Fée (§ 1 c).
1869. Heteropteris Fée (non Fée 1843). (§ 2).
— Doryopteridastrum Fée (Pellaea §). (§ 1).
1891. Allosorus sp. O.Ktze. (§ 1).
— Bakeriopteris sp. O.Ktze. (§ 1 a).

§ 1. **Doryopteridastrum** (concolor).
Pellaea § Doryopteridastrum Fée 1869 = Pellaea § Cheiloplecton Bak. 1867 pt. Christ 1897 = Allosorus sp. O.Ktze.
a. Bakeriopteris O.Ktze. 1891 (triphylla) = Cassebeera Klf. 1824. Bak. 1867. Christ 1897. **Diels** 1899.
b. Lytoneuron Kl. 1847 (lomariacea) = Pteris § Heterophlebium pt. Bak. 1867.
?c. Cheiloplecton Fée 1857 (ambigua).

§ 2. **Eudoryopteris** (pedata).

Doryopteris J. Sm. 1841 = Litobrochia § Doryopteris Moore 1857 = Pteris § Doryopteris Bak. 1867. Christ 1897 = Pellaea § Doryopteris Prantl 1882.

— Heteropteris Fée 1869 (lonchophora).

69. ADIANTOPSIS Fée 1850-52. — Species: 15.

Typ. Adiantum radiatum L.

1753. Adiantum sp. L.
1806. Cheilanthes sp. Sw.
1846. Actinopteris J. Sm. (Cheilanthes §) (non Actiniopteris Link 1841).
1852. Hypolepis sp. Hk.
1857. Adiantastrum Fée (Adiantopsis §).
— Cheilanthastrum Fée (Adiantopsis §).

Cheilanthes § Adiantopsis p. p. max. Bak. 1867. Christ 1897.

a. Euadiantopsis (Adiantastrum) Fée 1857 (radiata).
b. Cheilanthastrum Fée 1857 (capensis).

70. ASPLENIOPSIS Mettenius; Kuhn 1882. — Species: 1.

Typ. Gymnogramme decipiens Mett.

Gymnogramme sp. Bak. 1868. Gymnogramme § Aspleniopsis Christ 1897.

71. NOTHOLAENA R Brown 1810. — Species: 47.

Typ. Acrostichum Marantae L.

1753. Acrostichum sp. L.
— Pteris sp. L.
1810. Notholaena R.Br.
1811. Cincinalis Desv. (non Gled. 1764). (§ 1).
1836. Gymnogramme sp. Pr. Mett. 1856. 1859.
1841. Eriochosma J. Sm. (Notholaena §). (§ 1 a).
1841. Lepichosma J. Sm. pt. (Notholaena §). (§ 1 b).
1857. Alloesthes Moore (Nothochlaena §). (§ 1).
1859. Cheilanthes sp. Mett.
1866. Cosentinia Todaro (§ 1 c).
1872. Aleuritopteris sp. Fourn. (§ 2).
1875. Chrysochosma J. Sm. (Nothochlaena §). (§ 2).

1. Cincinalis Desv. 1811 = Nothochlaena § Eunothochlaena Bak. 1867. (Christ 1897) = Nothochlaena § Alloesthes Moore 1857.
   a. Eriochosma J. Sm. 1841 (rufa).
   b. Lepichosma J. Sm. 1841 pt. (Marantae).
   c. Cosentinia Todaro 1866 (vellea).
2. Chrysochosma J. Sm. 1875 (sulphurea) = Nothochlaena § Cincinalis pt. Bak. 1867. Christ 1897.

72. CHEILANTHES Swartz 1806. — Species: 101.

Typ. ?

1768. Pteris sp. Gmel.
1781. Adiantum sp. L.fil.
1806. Cheilanthes Sw.
1825. Gymnia Ham. msc. (§ 2).
1833. Othonoloma Link (§ 1 b).
1836. Allosorus sp. Pr. (§ 1 b).
1836 Physapteris Pr. (Cheilanthes §). (§ 3).
1841. Cassebeera sp. J. Sm. (§ 2).
— Notholaena § Lepichosma pt. J. Sm. (§ 3).
1850-52. Aleuritopteris Fée (§ 2).

1850-52. Myriopteris Fée (§ 1).
— Cheilanthastrum Fée (Myriopteris §). (§ 3 b).
1867. Pellaea sp. Bak. (§ 1 b).
— Oeosporangium Vis. (§ 3 a).
1873. Parallosorus Keys. (Cheilanthes §). (§ 1 b).
1875. Pteridopsis J. Sm. (non Link 1841) (Cheilanthes §). (§ 1).
1877. Cheilosoria Trev. (§ 1 a).

§ 1. **Eucheilanthes** Bak. 1867 (pteridioides).
Cheilanthes § Pteridopsis J. Sm. 1875.
a. Cheilosoria Trev. 1877 (tenuifolia).
b. Othonoloma Link 1833 (marginata) = Allosorus sp. Pr. 1836 = Pellaea § Allosorus pt. Bak. 1867 = Cheilanthes § Parallosorus Keys. 1873.
§ 2. **Aleuritopteris** (farinosa).
Gymnia Ham. 1825 msc. = Cassebeera sp. J. Sm. 1841 = Aleuritopteris Fée 1850-52.
§ 3. **Physapteris** Pr. 1836 (lendigera).
Notholaena § Lepichosma pt. J. Sm. 1841 = Myriopteris Fée 1850-52.
a. Oeosporangium Vis. 1867 (persica).
b. Cheilanthastrum Fée 1850-52 (scariosa).

73. HYPOLEPIS Bernhardi 1806. — Species: 29.

Typ. Lonchitis tenuifolia Forst.

1753. Lonchitis sp. L.
1806. Cheilanthes sp. Sw. (§ 2).
— Hypolepis Bernh.
1810. Adiantum sp. Poir.
1852. Aspidotis Nutt. msc. (§ 2).
1873. Phegopteris sp. Keys.

§ 1. **Euhypolepis** Bak. 1867 (tenuifolia).
§ 2. **Aspidotis** Hk. 1852 (californica).
Aspidotis Nutt. 1852. Cheilanthes sp. Bak. 1867. Christ 1897.

74. LLAVEA Lagasca 1816. — Species: 1.

Typ. L. cordifolia Lag.

1839. Ceratodactylis J. Sm.
1850-52. Botryogramme Fée.
1882. Cryptogramme § Llavea Prantl.

75. ONYCHIUM Kaulfuss 1820. 1824. — Species: 6.

Typ. O. auratum Klf.

1825. Leptostegia Don.
1827. Phorolobus pt. Desv.
1882. Cryptogramme § Onychium Prantl. **Diels** 1899.

76. CRYPTOGRAMMA R.Brown 1823. — Species: 4.

Typ. C. acrostichoides R.Br.

1806. *Allosorus* pt. Bernh. (an nom. opt.?).
1823. Cryptogramma R.Br.
1827. Phorolobus pt. Desv.
1845. Homopteris Rupr. (Allosorus §) (§ 1 b).
1858. Pellaea sp. Hk. (§ 1 b).
1882. Cryptogramme § Eucryptogramme Prantl. **Diels** 1899.

a. Cryptogramma R.Br. (acrostichoides).
b. Homopteris Rupr. 1845 (Stelleri) = Pellaea sp. Bak. 1867. Christ 1897.

76 a. NEURORSORIA Mett.; Kuhn 1869. — Species: 1.

Typ. Acrostichum pteroides R.Br.

77. PLAGIOGYRIA (Kze. 1850) Mettenius 1858. — Species: 11.

Typ. Lomaria euphlebia Kze.

1848. Lomaria sp. Kze.
1849. *Polygramma* Presl (an nomen veter.?).
1850-52. Lomaria § Plagiogyria Kze. Bak. 1867.
1858. Plagiogyria Mett.

VI. 3. ADIANTINAE.

78. ADIANTUM Linné 1753. — Species: 184.

Typ. ?

1836. Adiantellum Pr. (Adiantum §).
1841. Hewardia J. Sm. (§ 1 b).
1850-52. Apotomia Fée (Adiantum §).
— Synechia Fée (Adiantum §).
1857. Mesopleuria Moore (Adiantum §).
— Isotes Moore (Adiantum §). (§ 1 a).
1881. Adiantellum Kuhn (Adiantum §) (non Pr. 1836). (§ 2).

Adiantum § Adiantum Pr. 1836 = § Synechia Fée 1850-52 = § Mesopleuria Moore 1857 (species soris continuis).

§ Adiantellum Pr. 1836 = § Apotomia Fée 1850-52 (species soris globosis distinctis).

§ 1. **Euadiantum** Kuhn 1881 (?).
a. Isotes Moore 1857 (Leprieurii).
b. Hewardia J. Sm. 1841 (adiantoides) = Adiantum § Hewardia Bak. 1867.

§ 2. **Adiantellum** Kuhn 1881 (?).

VI. 4. PTERIDINAE.

79. ACTINIOPTERIS Link 1841. — Species: 1.

Typ. Acrostichum australe L.fil.

Asplenium § Actiniopteris J. Sm. 1875.

80. AMPHIBLESTRA Presl 1836. — Species: 1.

Typ. Pteris latifolia H.B.Willd.

Pteris § Amphiblestra Bak. 1867. Christ 1897.

81. ANOPTERIS (Prantl) Diels 1899. — Species: 1.

Typ. Pteris heterophylla L.

Pteris sp. Bak. 1867. Christ 1897. Cryptogramme § Anopteris Prantl 1882.

82. OCHROPTERIS J. Smith 1841. — Species: 1.

Typ. Adiantum pallens Sw.

Cryptogramme § Ochropteris Prantl 1882.

83. PTERIS Linné 1753 reduct. - Species: 157.

Typ. P. longifolia L.

1753. Pteris L.
1763. ? Thelypteris Adans
1819. Lemapteris Rafin.
— Peripteris Rafin.
1836. Campteria Pr. (§ 1 d).
— Litobrochia pt Pr. (§ 3).
1841. Pteridopsis Link (Pteris §). (§ 1 b).
1847. Macropteris Webb et Berth. (Pteris §). (§ 1).
1849. Pycnodoria Pr. (§ 1 a).
1850-52. Heterophlebium Fée (§ 2).
1865. Heteropteris Pr. msc. (non Fée 1843). (§ 1 c).
1873. Parapteris Keys. (Pteris §). (§ 1).
1874. Schizopteris Hill. (§ 1 e).
1888. Schizostege Hill. (§ 1 e).

§ 1. **Eupteris** (longifolia).

Pteris § Macropteris Webb et Berth. 1847 = Pteris § Parapteris et Eupteris Keys. 1873.

a. Pycnodoria Pr. 1849 (opaca).
b. Pteridopsis Link 1841 (longifolia).
c. Heteropteris Pr. 1865 (lineata).
d. Campteria Pr. 1836 (biaurita) = Litobrochia § Campteria Fée 1850-52 = Pteris § Campteria Bak. 1867.
e. Schizostege Hill. 1888 (Lydgatei) = Schizopteris Hill. 1874 msc. = Cheilanthes sp. Bak. 1874.

§ 2. **Heterophlebium** (grandifolia).

Heterophlebium Fée 1850-52.

§ 3. **Litobrochia** (denticulata).

Litobrochia p. p. max. Pr. 1836.

84. HISTIOPTERIS (Agardh 1839) J. Smith 1875. — Species: 2.

Typ. Pteris incisa Thbg.

Pteris § Histiopteris Agardh 1839. Christ 1897.
Pteris § Litobrochia pt. Bak. 1867.

85. LONCHITIS Linné 1753 emend. Kuhn 1879. — Species: 8.

Typ. L. hirsuta L. vel L. aurita L.

1810. Pteris sp. Willd. Mett. 1856. Christ 1897.
1850-52. Lonchitidium Fée (Pteris §).
1851. Anisosorus Trev.
1882. Antiosorus Roemer; Kuhn.

§ 1. **Eulonchitis** (aurita).

Lonchitis auctt. Bak. 1867.

§ 2. **Anisosorus** (hirsuta).

Anisosorus Trev. 1851 = Pteris § Lonchitideum Fée 1850-52 = Antiosorus Roem.; Kuhn 1882 = Pteris sp. Bak. 1867.

86. PTERIDIUM Gleditsch 1760. — Species: 1.

Typ Pteris aquilina L.

1753. Pteris sp. L.
1760. Pteridium Gled. Scopoli.
1764. Cincinalis Gled. Trev. 1874.
1836. Aquilini(a) Pr. (Allosorus §).
1839. Ornithopteris pt. Ag. (Pteris §) (non Bernh. 1806).

1845. Eupteris Newm.
1847. Nymphopteris Webb et Berth. (Pteris §).
1875. Ornithopteris pt. J Sm. (Ag. 1839 §).

Pteris § Paesia pt. Bak. 1867 = Pteris § Pteridium Christ 1897.

87. PAESIA St. Hilaire 1833 emend. Kuhn 1822 — Species: 7.

Typ. P. viscosa St. Hil.

1824. Pteris sp. Lab.
1833. Paesia St. Hil.
1836. Allosorus § Aquilini pt. Pr.
1839. Ornithopteris pt. Ag. (Pteris §).
1875. Ornithopteris pt. J. Sm. (non Bernh. 1806).

Pteris § Paesia Bak. 1867 (pt.). Christ 1897.

VII. VITTARIEAE.

VII. 1. VITTARIINAE.

88. MONOGRAMMA Schkuhr 1809. Desvaux 1811. — Species: 15.

Typ. Pteris graminea Poir.

1801. Grammitis sp. Sw.
1809. Monogramma Schkuhr.
1820. Cochlidium Klf. 1824. Schott 1834 (§ 1 a).
1824. Taenitis sp. Klf. (§ 2). Mett. 1856 (§ 1 a. 2).
1827. Micropteris sp. Desv.
1828. Pleurogramme Bl. (Antrophyum §). (§ 2).
1836. Pleurogramme Pr. (Bl. 1828 §). (§ 2).
1843. Vaginularia Fée (§ 1 b).
1854. Diclidopteris Brack. (§ 1 c).

§ 1. **Eumonogramma** (graminea).

Monogramma Schkuhr 1809. Christ 1897.

a. Cochlidium Klf. 1824 (graminoides) = Pleurogramme sp. Fée 1851-62. J. Sm. 1875 = Taenitis sp. Mett. 1856.

b. Vaginularia Fée 1843 (trichoidea).

c. Diclidopteris Brack. 1854 (paradoxa) = Pleurogramme sp. Fée 1851-52.

§ 2. **Pleurogramme** (seminuda).

Taenitis sp. Klf. 1824. Mett. 1856 = Antrophyum § Pleurogramme Bl. 1828 = Pleurogramme Pr. 1836. Christ 1897.

89. VITTARIA Smith 1793. — Species: 46.

Typ. Pteris lineata L.

1753. Pteris sp. L.
1790 ? Oetosis Necker (t. Greene).
1793. Vittaria Sm.
1810. Taenitis sp. R.Br. (§ 3).
1827. Pteropsis sp. Desv. (vide Drymoglossum).
1836. Haplopteris Pr. (§ 2).
1842. Taeniopteris Hook. (§ 2 a).
1842. Taeniopsis J. Sm. (§ 2 a).
1851-52. Pteropsis Fée (§ 3).
1854. Aristaria — Parenchymaria — Runcinaria (Vittaria §§) K. Müll. (sectiones structura squamarum signatae).
1900. Oetosis Greene (non O.Ktze. 1891).

§ 1. **Euvittaria** (elongata).

§ 2. **Haplopteris** (lineata) — (nomen Euvittaria hujus sectionis proprius est).
Haplopteris Pr. 1836 = Taeniopsis J. Sm. 1842 = Vittaria § Taeniopsis Fée 1851-52. Bak. 1868. Christ 1897. **Diels** 1899.
a. Taeniopteris Hk. 1842 (scolopendrina) = Haplopteris Pr. 1836.

§ 3. **Pteropsis** (angustifolia) — (nomen malum).
Pteropsis Fée 1851-52. Christ 1897 (Desv. 1827 p. p. min.) = = Taenitis sp. Bak. 1868.

VII. 2. ANTROPHYINAE.

90. HECISTOPTERIS J. Smith 1842. — Species: 1.

Typ. Gymnogramma pumila Spr.

Gymnogramme sp. Bak. 1868.

91. ANTROPHYUM Kaulfuss 1824. — Species: 27.

Typ. (Hemionitis plantaginea Cav.).

1753. Hemionitis sp. L.
1822. Solenopteris Wall. pt. (nomen).
1824. Antrophyum Klf.
1827. Polytaenium Desv. (a).
1857. Scoliosorus Moore (b).
1877. Dictyogramme sp. Trev. (c).

a. Polytaenium Desv. 1827 (lineatum).
b. Scoliosorus Moore 1857 (ensiforme).
c. § B. Diels = Dictyogramme sp. Trev. 1877.

92. ANETIUM (Kunze) Splitgerber 1840. — Species: 1.

Typ. Acrostichum citrifolium L.

Acrostichum § Anetium Kze. hb.
Hemionitis § Anetium Bak. 1868.

VIII. POLYPODIEAE.

VIII. 1. TAENITIDINAE.

93. DRYMOGLOSSUM Presl 1836. — Species: 9.

Typ. Acrostichum heterophyllum L.

1759. Pteris sp. L.
1790. ?Oetosis Neck. (t. O.Ktze.).
1822. Solenopteris Wall. pt. (nomen).
1824. Nothochlaena sp. Klf.
1827. *Pteropsis* Desv. (pro parte max. certe nom. opt.).
1836. Drymoglossum Pr.
1849. Lemmaphyllum Pr. (a).
1856. Taenitis sp. Mett.
1857. Schizolepton sp. Moore (non Fée 1851-52) (b).
1860. Asteroglossum J. Sm. msc. (a).
1891. Oetosis sp. O.Ktze.

a. Lemmaphyllum Pr. 1849 (carnosum) = Asteroglossum J. Sm. 1860.
(b. Schizolepton Moore 1857 pt. (rigidum)).

94 (?). DRYMOTAENIUM Makino 1901. — Species: 1.

Typ. Taenitis Miyoshianum Mak.

95. ESCHATOGRAMME Trevisan 1851. — Species: 1.

Typ. Pteris furcata L.

1810. Taenitis sp. Willd. Bak. 1868.
1827. Pteropsis sp. Desv.
1851-52. Cuspidaria Fée. Christ 1897 (non DC. 1838 : Gen. Bignoniac).
1851. Eschatogramme Trev.
1854. Dicranoglossum J. Sm. **Diels** 1899.

96. TAENITIS Willdenow; Sprengel 1804. — Species: 2.

Typ. Pteris blechnoides Willd.

Taenitis auctt. Bak. 1868 valde reduct.

97. PLATYTAENIA Kuhn 1882. — Species: 1.

Typ. Acrostichum Requinianum Gaud.

98. PALTONIUM Presl 1849. — Species: 2.

Typ. Pteris lanceolata L.

1810. Taenitis sp. R.Br. Bak. 1868.
1828. Chilogramme pt. Bl. (Antrophyum §).
1843. Heteropteris Fée. **Diels** 1899 (non HBK. gen. Malpighiac).
1849. Paltonium Pr.
1851-52. Neurodium Fée. Christ 1897.
1900. Cheilogramma Und.

99. HYMENOLEPIS Kaulfuss 1824. — Species: 4.

Typ. Acrostichum spicatum L.fil.
(Hymenolepis ophioglossoides Klf.).

1803. Belvisia pt. Mirb.
1824. Hymenolepis Klf. (non Cass. 1817: Gen. Compositar).
1836. Gymnopteris § Gymnopteris Pr.
1849. Macroplethus Pr. (platyrhynchos).
1850. Hyalolepis Kze. (non DC. 1837: Gen. Compositar).
1856. Taenitis sp. Mett.
1903. Belvisia Und.

Acrostichum § Hymenolepis Bak. 1868.

99 a. CHRISTOPTERIS Copeland 1905. — Species: 1.

Typ. Polypodium Sagitta Christ.

VIII. 2. POLYPODIINAE.

100. POLYPODIUM Linné 1763 reduct. — Species: 784.

Typ. ?

1753. Polypodium L.
1801. Grammitis pt. Sw. (§ 1 a).
1810. Pleopeltis Humb. et Bonpl. (§ 6 o).
1824. Selliguea Bory (§ 7).
— Marginaria Bory (§ 3).
— Adenophorus Gaud. (non Desv. 1808 gen. algar.) (§ 1 l).
1824. Xiphopteris Klf. (§ 1 e).
1827. Amphoradenium Desv. (§ 1 l).
— Micropteris Desv. (§ 1 e).
1828. Diagramma Bl. (Grammitis §). (§ 7).
— Paragramma Bl. (Grammitis §). (§ 6 h).

1828. Atactosia pt. Bl. (Polypodium §). (§ 6 c).
— Loxogramme Bl. (Antrophyum §). (§ 8).
— Ctenopteris Bl. Kze. 1846.
— Goniophlebium Bl (Polypodium §). (§ 2).
1833. Microsorium Link (§ 6 b).
1834. Aglaomorpha Schott (§ 6 q).
— Anaxetum Schott (§ 6 e).
1836. Calymmodon Pr. (§ 1 f).
— Campyloneurum Pr. (§ 5).
— Chilopteris Pr. (Grammitis §). (§ 1 a).
— Dictyopteris pt. Pr. (§ 6 g).
— Goniophlebium Pr. (Bl. 1829 §). (§ 2).
— Loxogramme Pr. (Bl. 1828 §). (§ 8).
— Marginaria Pr. emend. (Bory 1824). (§ 3 et 2).
— Microgramma Pr. (§ 6 j).
— Phymatodes Pr. (§ 6 (6 m)).
— Pleuridium Pr. (Phymatodes §). (§ 6 e).
— Pleurogonium Pr. (Marginaria §). (§ 3 a).
— Psygmium Pr. (§ 6).
— Synammia Pr. (§ 2 a).
1838. Cyrtophlebium R.Br. (Polypodium §). (§ 5).
— Phlebodium R.Br. (Polypodium §). (§ 4).
1840. Lopholepis J. Sm. (Goniophlebium §). (§ 3 a).
— Schellolepis J. Sm. (Goniophlebium §). (§ 2 b).
1841. Dicranopteris »Bl.« Endl. (Polypodium §). (§ 1).
— Chrysopteris Link (§ 4 et 6 m).
— Craspedaria Link (§ 3 a).
— Diblemma J. Sm. (§ 6 a).
— Phyllitidis J. Sm. (Drynaria §).
— Cyrtophlebium J. Sm. (R.Br. 1838 §). (§ 5).
— Drynaria § Phymatodes J. Sm. (§ 6).
— Phlebodium J. Sm. (R.Br. 1838 §). (§ 4).
— Lepicystis J. Sm. (Goniophlebium §). (§ 3).
1842. Lopholepis J. Sm. (1840 §). (§ 3 a).
— Lepicystis J. Sm. (1841 §). (§ 3).
1843. Cryptosorus Fée (§ 1 g).
1846. Anapeltis J. Sm. (Phlebodium §). (§ 6 i).
— Dictymia J. Sm. (§ 6 g).
— Lepisorus J. Sm. (Drynaria §). (§ 6 f).
1846. Symplecium Kze. (Polypodium §). (§ 6 p).
1847. Mecosorus Kl. (§ 1 a et 6).
1849. Colysis Pr. (§ 6 d et 7).
— Dictyogramma Pr. (Colysis §). (§ 7).
— Crypsinus Pr. (§ 2 d).
— Microterus Pr. (§ 6 l).
1850-52. Drynaria Fée p. p. max. (non Bory 1825). (§ 6).
— Pleuridium Fée (Pr. 1836 §). (§ 6 e).
— Plectopteris Fée (§ 1 f).
1851. Trichocalymma Zenker msc. (§ 1 c).
— Trichothemeleum Kze. (Grammitis §). (§ 1 c).
1854. Glyphotaenium J. Sm. (Cteno pteris §). (§ 1 h).
1857. Coelopteris A.Br. msc. (§ 1 g).
— Thylacopteris Kze. msc. J. Sm. 1875 (§ 1 i).
— Anapeltis J. Sm. (1846 §). (§ 6 i).
1857. Allothecium Moore (Pleopeltis §). (§ 6 n).
— Arthromeris Moore (Pleopeltis §). (§ 6 r).
— Cephalosorium Moore (Campyloneurum (§ 5 a).
— Holcosorus Moore (§ 1 d).
— Marginariopsis Moore (Phlebodium §). (§ 2 c).
— Paragramma Moore (Bl. 1828 §). (§ 1 h).
— Phlebodiopsis Moore (Pleopeltis §). (§ 1 k et i).
— Pleopeltis Moore emend. (H.B. Willd. 1810). (§ 6).
— Prosechium Moore (Polypodium §). (§ 1 j).
— Themelium Moore (Polypodium §). (§ 1 k).
1857. Catenularia Zippel msc. (§ 1 a).

1857. Leptostegia Zippel msc. (non Don 1825). (§ 1 g).
— Dictyophlebia Gris. (Polypodium §); (§ 2-6; sp. venis anastom.).
— Eleutherophlebia Gris. (Polypodium §). (§ 1).
1861. Cystidium J. Sm. msc. (§ 1 g).
1864. Gymnogramme § Selliguea pt. Hk. (§ 7).
1866. Schellolepis J. Sm. (1840 §). 1875. (§ 2 b).
1870. Prosaptia sp. Mett. (§ 1 g).
1872. Pteropsis Fourn. (non Desv. 1827). (§ 6 o).
1873. Austrogramme Fourn. pt. (§ 1 a).
— Grammitastrum Fourn. (Polypodium §). (§ 1 a).
— Tomophyllum Fourn. (Polypodium §). (§ 1 k).
1875. Arthromeris J. Sm. (Moore 1857 §). (§ 6 r).
— Colysis J. Sm. (non Pr. 1849). (§ 6 (6 d)).
1875. Ctenopteris J. Sm. emend. (Bl. 1829). (§ 1).
— Glyphotaenium J. Sm. (1854 §). (§ 1 h).
— Lomaphlebia J. Sm. (§ 1 b).
— Micropteris J. Sm. (non Desv. 1827). (§ 1).
— Phymatopsis J. Sm. (§ 6 s et k).
1879. Clathropeltis Kuhn (Polypodium §). (§ 6 f).
1897. Myrmecophila Christ (Polypodium §). (§ 9).
1899. Lepicystis Diels emend. (J. Sm. 1842). (§ 1. 3. 6 o).
Goniolepicystis Diels (Lepicystis §). (§ 3).
— Phlebiolepicystis Diels (Lepicystis §). (§ 6 o).
— Aspidopodium Diels (Polypodium §). (§ 9).
1905. Drynariopsis Copeland (Polypodium §). (§ 6 t).

§ 1. **Eupolypodium** (vulgare).

Polypodium § Eleutherophlebia Gris. 1857.
Ctenopteris Bl. 1828 (sp. venis simplicibus).
§ Dicranopteris Bl. Endl. 1841 (sp. venis furcatis).
Ctenopteris J. Sm. 1875 (sp. foliis fasciculatis ad caudicem non articulatis).
— Micropteris J. Sm. 1875 (sp. foliis remotis ad caudicem non articulatis).
— Lepicystis § Eulepicystis **Diels** 1899 (sp. foliis squamis peltatiformibus instructis).

a. Grammitis Sw. 1801 pt. (marginellum: sp. foliis integris soris elongatis) = Grammitis § Chilopteris Pr. 1836 = Catenularia Zippel 1857 = Polypodium § Grammitastrum Fourn. 1873 = Mecosorus pt. Kl. 1847.
b. Lomaphlebia J. Sm. 1875 (gramineum).
c. Trichothemeleum Kze. 1851 (mediale v. parasiticum) = Trichocalymma Zenker msc. 1851.
d. Holcosorus Moore 1857 (bisulcatum).
e. Xiphopteris Klf. 1824 (serrulatum) = Micropteris Desv. 1827.
f. Calymmodon Pr. 1836 (cucullatum) = Plectopteris Fée 1850-52.
g. Cryptosorus Fée 1843 (obliquatum) = Ctenopteris (Bl. 1828 pt.) Kze. 1846 = Coelopteris A.Br. msc. 1857 = Leptostegia Zippel msc. 1857 = Cystidium J. Sm. msc. 1861 = Prosaptia sp. Mett. 1870.
h. Glyphotaenium J. Sm. 1854 (goniopteroides).
i. Thylacopteris Kze. msc. 1857. J. Sm. 1875 (papillosum).

j. Prosechium Moore 1857 (suspensum, pendulum).
k. Themelium Moore 1857 (tenuisectum) = Tomophyllum Fourn. 1873 (millefolium).
l. Amphoradenium Desv. 1827 (hymenophylloides etc.) = Adenophorus Gaud. 1824.

§ 2. **Goniophlebium** Bl. 1828 (Ex. brasiliense).

Goniophlebium Pr. 1836. Marginaria § Marginaria p. p. max. Pr. 1836.
a. Synammia Pr. 1836 (synammia).
b. Schellolepis J. Sm. 1840 (subauriculatum).
c. Marginariopsis Moore 1857 (Lowei).
d. ? Crypsinus Pr. 1849 (pyrolifolium).

§ 3. **Marginaria** (polypodioides).

Marginaria Bory 1824. 1826 (p. p. max.) = Lepicystis J. Sm. 1842 = Polypodium § Goniophlebium pt. Bak. 1867 = Polypodium § Lepicystis Christ 1897 = Lepicystis § Goniolepicystis Diels 1899.
a. Pleurogonium Pr. 1836 (vacciniifolium) = Lopholepis J. Sm. 1840 = Craspedaria Link 1841 = Polypodium § Craspedaria Christ 1897 = Goniophlebium § Crypsinus Moore 1857.

§ 4. **Phlebodium** R.Br. 1838 (aureum).

Chrysopteris Link 1841 p. p. max. Fée 1850-52 = Phlebodium ** J. Sm. 1841.

§ 5. **Campyloneurum** (phyllitidis).

Campyloneurum Pr. 1836 = Campylonevron Fée 1850-52 = Polypodium § Cyrtophlebium R.Br. 1838 = Cyrtophlebium J. Sm. 1841.
a. Cephalosorium Moore 1857 (decurrens).

§ 6. **Pleopeltis** (?).

Pleopeltis Humb. et Bonpl. 1810 emend. Moore 1857 = Phymatodes Pr. 1836 = Drynaria § Phymatodes J. Sm. 1841 = Drynaria p. p. max. Fée 1850-52 = Polypodium § Phymatodes Bak. 1867 (pt.). Cylysis J. Sm. 1875.
a. Diblemma J. Sm. 1841 (tenuilore).
b. Microsorium Link 1833 (punctatum).
c. Atactosia pt. Bl. 1828 (musifolium).
d. Colysis Pr. 1849 (pt.) (hemionitideum). J. Sm. 1875 pt.
e. Anaxetum Schott 1834 (crassifolium) = Phymatodes § Pleuridium Pr. 1836 = Pleuridium Fée 1850-52.
f. Lepisorus J. Sm. 1846 (excavatum et lineare) = Clathropeltis Kuhn 1879 (lineare).
g. Dictymia J. Sm. 1846 (Brownii) = Dictyopteris pt. Pr. 1836.
h. Paragramma Bl. 1828 (revolutum). Moore 1857.
i. Anapeltis J. Sm. 1846 (lycopodioides) = Craspedaria pt. Fée 1850-52 = Pleopeltis § Phlebodiopsis pt. Moore 1857 = Polypodium § Craspedaria pt. Christ 1897.
j. Microgramma Pr. 1836 (persicariifolium).
k. Phlebodiopsis pt. Moore 1857 (accedens) = Phymatopsis pt. J. Sm. 1875.
l. Microterus Pr. 1849 (neglectum).
m. Phymatodes Pr. 1836 reduct. (phymatodes) = Chrysopteris pt. Link 1841.
n. Allothecium Moore 1857 (pteropus).

o. Pleopeltis H.B.Willd. 1810 (angustum) = Lepicystis § Phlebiolepicystis **Diels** 1899. Pteropsis Fourn. 1872.
p. Symplecium Kze. 1846 (insigne).
q. Aglaomorpha Schott 1834 (Meyenianum) = Psygmium Pr. 1836.
r. Arthromeris Moore 1857 (Wallichianum). J. Sm. 1875.
s. Phymatopsis J. Sm. 1875 pt. (palmatum).
t. Drynariopsis Copeland 1905 (heracleum).

§ 7. **Selliguea** (Feei).
Selliguea Bory 1824 = Grammitis § Diagramma Bl. 1828 = Colysis sp. Pr. 1849 = Gymnogramme § Selliguea pt. Hk. 1864. Bak. 1868 = Polypodium § Selliguea Christ 1897 = Polypodium § Pleopeltis pt. **Diels** 1899. Colysis § Dictyogramma Pr. 1849 (ellipticum).

§ 8. **Loxogramme** Christ 1897 (Blumeanum).
Antrophyum § Loxogramme Bl. 1828 = Loxogramme Pr. 1836 = Gymnogramme § Selliguea pt. Bak. 1868 = Polypodium § Pleopeltis pt. **Diels** 1899.

§ 9. **Myrmecophila** Christ 1897 (sinuosum).
Polypodium § Phymatodes pt. Bak. 1868 = Polypodium § Aspidopodium **Diels** 1899.

101. ENTEROSORA Baker 1886. — Species: 1.

Typ. E. Campbellii Bak.

102. CYCLOPHORUS Desvaux 1811. — Species: 73.

Typ. Polypodium adnascens Sw.

1784. Acrostichum sp. Thbg.
1786. Polypodium sp. Forst.
1803. ? Pyrrosia Mirbel.
— *Candollea* Mirbel p. p. max. (nom. opt.!).
1811. Cyclophorus Desv.
1824. Niphobolus Klf. **Diels** 1899.
1836. Scytopteris Pr. (Niphobolus §). (§ 2).
1848. Digramma Kze. (Antrophyum §). (§ 1 a).
1849. Apalophlebia Pr. (§ 1 a).
— Cyclophorus Pr. (Desv. 1811).
— Galeoglossa Pr. (§ 1 d).
— Gyrosorium Pr. (§ 1 f).
— Polycampium Pr. (§ 1 c).
— Scytopteris Pr. (1836 §). (§ 2).
— Sphaerostichum Pr. (§ 1 g).
1857. Niphopsis J. Sm. (§ 2 a).
1875. Niphidium J. Sm. (§ 1 b).

§ 1. **Eucyclophorus** (adnascens).
Polypodium § Niphobolus Bak. 1868. Christ 1897 = Niphobolus § Euniphobolus **Diels** 1899.
a. Digramma Kze. 1848 (flocciger).
b. Niphidium J. Sm. 1875 (americanus).
c. Polycampium Pr. 1849 (lingua).
d. Galeoglossa Pr. 1849 (nummularifolius).
e. Apalophlebia Pr. 1849 (stigmosus) = ? Pyrrosia Mirb. 1803.
f. Gyrosorium Pr. 1849 (africanus).
g. Sphaerostichum Pr. 1849 (sphaerostichus).

§ 2. **Scytopteris** (confluens).

Niphobolus § Scytopteris Pr. 1836 = Scytopteris Pr. 1849 = Niphopsis J. Sm 1857 = Polypodium § Phymatodes sp. Bak. 1868 = Polypodium § Pleopeltis sect. Niphopsis Christ 1897 = Niphobolus § Niphopsis **Diels** 1899.

a. Niphopsis J. Sm. 1857 (angustatus).

103. LECANOPTERIS Reinwardt 1825. — Species: 7.

Typ. L. carnosa Bl.

1824. Onychium Reinw. (non Klf. 1820).
1897. Polypodium § Lecanopteris Christ.

104. DRYOSTACHYUM J. Smith 1841. — Species: 3.

Typ. D. splendens J. Sm.

Polypodium sp. Bak. 1868. Acrostichum § Photinopteris pt. Bak. 1868. Polypodium § Dryostachyum pt. Christ 1897.

105. PHOTINOPTERIS J. Smith 1841. — Species: 1.

Typ. Lomaria speciosa Bl.

Acrostichum § Photinopteris pt. Bak. 1868. Polypodium § Dryostachyum pt. Christ 1897.

106. DRYNARIA (Bory 1825) J. Smith 1841 emend. Moore 1857. — Species: 14.

Typ. Polypodium quercifolium L.

1753. Polypodium sp. L.
1825 Drynaria Bory (Polypodium §).
1836. Phymatodes § Drynaria Sect. B. Pr.
1842. Drynaria § Drynaria vera J. Sm.
1857. Drynaria Moore.
1875. Poronema J. Sm. (Drynaria §).

Polypodium § Drynaria Bak. 1868. Christ 1897.

§ 1. **Eudrynaria** (quercifolia).
§ 2. **Poronema** J. Sm. 1875 (rigidula).

IX. ACROSTICHEAE.

IX. 1. ACROSTICHINAE.

107. ELAPHOGLOSSUM Schott 1834 emend. Christ 1899. — Species: 280.

Typ. Acrostichum conforme Sw.

1753. Acrostichum sp. L.
1834. Elaphoglossum Schott.
— Rhipidopteris Schott.
1836. Aconiopteris Pr. (§ 1 a. α).
— Olfersia p. p. max. Pr. (non Raddi 1819).
1841. Peltapteris Link (§ 1 a. β).
1845. Acrostichum Fée.
— Hymenodium Fée (§ 2 b. α).
1846. Dictyoglossum J. Sm. (§ 2 b. α).
1849. Microstaphyla Pr. (§ 2 a. α).
1849. Nebroglossa Pr. (Aconiopteris §) (§ 1 a. α).
— Anetium sp. Pr. (§ 2 b. α).
1857. Oligolepidum, — Polylepidum } Moore (§§).
1899. Condyloneura, — Craspedoglossa, — Gymnoglossa, — Lepidoglossa, — Pachyglossa, — Platyglossa, — Stenoneura } Christ (§§).

Acrostichum §§ Elaphoglossum, Rhipidopteris, Aconiopteris Bak. 1868.

§ 1. **Stenoneura** Christ 1899 (sp. venis apicibus non incrassatis).

a. Craspedoglossa Christ 1899 (sp. foliis nudis vel subnudis) = § Oligolepidum pt. Moore 1857.

α. Platyglossa Christ 1899 (sp. foliis majoribus acuminatis vel subacuminatis: Ex. latifolium).

— Aconiopteris pt. Fée 1845 (longifolium) = Nebroglossa Pr. 1849.

β. Pachyglossa Christ 1899 (sp. foliis minoribus acutis vel obtusis: Ex. conforme).

— Aconiopteris Pr. 1836 (nervosum).

— Rhipidopteris Schott 1834 (flabellatum). **Diels** 1899 = Peltapteris Link 1841 = Acrostichum § Rhipidopteris Bak. 1868.

b. Lepidoglossa Christ 1899 (sp. foliis ciliatis squamis pseudopeltatiformibus ± vestitis) = § Polylepidum pt. Moore 1857.

§ 2. **Condyloneura** Christ 1899 (sp. venis apicibus incrassatis).

a. Gymnoglossa Christ 1899 (sp. foliis nudis vel subnudis) = § Oligolepidum pt. Moore 1857.

α. Microstaphyla Pr. 1849. **Diels** 1899 (furcatum).

b. Polytrichia Sod. 1893. Christ 1899 (sp. foliis squamis setiformibus instructis) = § Polylepidum pt. Moore 1857.

α. Hymenodium pt. Fée 1845 (crinitum) = Dictyoglossum J. Sm. 1846 = Anetium sp. Pr. 1849 = Acrostichum § Chrysodium pt. Bak. 1868 = Elaphoglossum § Hymenodium **Diels** 1899.

108. TRACHYPTERIS André; Christ 1899. — Species: 1.

Typ. Hemionitis pinnata Hk.fil.

Acrostichum § Chrysodium pt. Bak. 1868 = Elaphoglossum § Heteroglossum **Diels** 1899.

109. ACROSTICHUM Linné 1753 reduct. — Species: 4.

Typ. A. aureum L.

1845. Chrysodium Fée. Christ 1897.
— Neurocallis Fée (praestantissimum).
Acrostichum § Chrysodium pt. Bak. 1868.

IX. 2. PLATYCERIINAE.

110. CHEIROPLEURIA Presl 1849. — Species: 1.

Typ. Polypodium bicuspe Bl.

Acrostichum § Chrysodium pt. Bak. 1868.

111. PLATYCERIUM Desvaux 1827. — Species: 14.

Typ. Acrostichum alcicorne Willem.

1826. Alcicornium Gaud. Und. 1905.
1836. Neuroplatyceros Endl. (Acrostichum §).
1845. NeuroplatycerosFée (Endl.1836 §).
1850-52. Scutigera Fée (Neuroplatyceros §).

§ 1. **Euplatycerium** (bifurcatum).

§ 2. **Scutigera** Fée 1850-52 (coronarium).

IV. PARKERIACEAE.

1. CERATOPTERIS Brongniart 1821. — Species: 1.

Typ. Acrostichum thalictroides L.

1823. Teleozoma R.Br.
— Cryptogenis Rich.
1824. Ellobocarpus Klf.
1825. Parkeria Hk.
1827. Furcaria Desv.

V. MATONIACEAE.

1. MATONIA R.Brown 1829. — Species: 2.

Typ. M. pectinata R.Br.

1828. Prionopteris Wall. (nomen).

VI. GLEICHENIACEAE.

1. STROMATOPTERIS Mettenius 1861. — Species: 1.

Typ. S. moniliformis Mett.

Gleichenia sp. Hk. 1865 = Gleichenia § Stromatopteris Christ 1897.

2. GLEICHENIA Smith 1793. — Species: 79.

Typ. Onoclea polypodioides L.

1753. Polypodium sp. L. (§ 2).
1771. Onoclea sp. L. (§ 2).
1793. Gleichenia Sm. (non Necker 1790) (§ 2).
1804. Mertensia Willd. (non Roth 1797: Gen. Boraginac.). (§ 1).
1806. *Dicranopteris* Bernh. (an nom. opt.?) (§ 1).
1810. Platyzoma R.Br. Hk. 1865 (§ 3).
1836. Sticherus Pr. (§ 1 c).
1836. Calymella Pr. (§ 2 a).
1848. Gleicheniastrum Pr. (§ 2 b).
1849. Hicriopteris Pr. (§ 1 b).
1856. Mesosorus Hassk. (§ 1).
1900. Acropterygium } Diels (§§) (§ 1 b-e).
— Diplopterygium
— Heteropterygium
— Holopterygium
1905. Protogleichenia Copeland (Dicranopteris §). (§ 1 a).

§ 1. **Dicranopteris** (linearis).

Mertensia Willd. 1804 = Dicranopteris Bernh. 1806 = Mesosorus Hassk. 1856 = Gleichenia § Mertensia Hk. 1865. Christ 1897. **Diels** 1900.

a. Protogleichenia Copeland 1905 (dolosa).
b. Diplopterygium Diels 1900 (glauca). — Hicriopteris Pr. 1849 (glauca var).
c. Holopterygium Diels 1900 (furcata).
— Sticherus Pr. 1836 (laevigata).
d. Acropterygium Diels 1900 (pectinata).
e. Heteropterygium Diels 1900 (linearis).

§ 2. **Eugleichenia** (polypodioides).
Gleichenia Sm. 1793.
a. Calymella Pr. 1836 (alpina).
b. Gleicheniastrum Pr. 1849 (circinnata).

§ 3. **Platyzoma** (platyzoma).
Platyzoma R.Br. 1810. Hk. 1865.

VII. SCHIZAEACEAE.

1. SCHIZAEA Smith 1793. — Species: 25.

Typ. ? Acrostichum pectinatum L.

1753. Acrostichum sp. L.
1792. Lophidium Rich.
1793 (vel 1791). Schizaea Sm.
1801. Ripidium Bernh.
1828. Actinostachys Wall.

§ 1. **Actinostachys** (digitata).
Actinostachys Wall. 1828.

§ 2. **Euschizaea** (pectinata).

§ 3. **Lophidium** (elegans).
a. Ripidium Bernh. 1801 (dichotoma).
b. Lophidium Rich. 1792 (elegans).

2. LYGODIUM Swartz 1801. — Species: 26.

Typ. Ophioglossum flexuosum L.

1753. Ophioglossum sp. L.
1801. Lygodium Sw.
— Ugena Cav.
— Ramondia Mirb.
— Odontopteris Bernh.
— Gisopteris Bernh. (§ 1).
1802. Hydroglossum Willd.
1803. Cteisium Michx. (§ 1).
1809. Vallifilix Thouars.
1842. Lygodictyon J. Sm. (§ 2 a).
1845. Hydroglossum Pr. (Willd. 1802 emend.) (§ 2 a).
— Arthrolygodes Pr. (Lygodium §) (§ 1 a).

§ 1. **Gisopteris** (palmatum).
Gisopteris Bernh. 1801 = Cteisium Michx. 1803 = Lygodium § Palmata Prantl 1881. **Diels** 1900.
a. Arthrolygodes Pr. 1845 (articulatum).

§ 2. **Volubilia** Prantl 1881 (volubile).
a. Lygodictyon J. Sm. 1842 (reticulatum) = Hydroglossum Pr. 1845.

§ 3. **Flexuosa** Prantl 1881 (flexuosum).

3. MOHRIA Swartz 1806. — Species: 3.

Typ. Polypodium caffrorum L.

4. ANEIMIA Swartz 1806. — Species: 64.

Typ. ? Osmunda phyllitidis L.

1753. Osmunda sp. L.
1806. Anemia Sw.
— Ornithopteris Bernh. (§ 3).
1824. Aneimia Klf.
1842. Anemidictyon J. Sm. (§ 2 a).
— Coptophyllum Gardn. (§ 2 a).

1842. Trochopteris Gardn. (§ 1).
1843. Mohria sp. J. Sm. (§ 2 c).
1845. Spathepteris Pr. (?).
1854. Anemirhiza J. Sm. (§ 3).
1869. Aneimiaebotrys Fée (§ 2 b).
1881. Hemianeimia Prantl (Aneimia §). (§ 2).

§ 1. **Trochopteris** (elegans).
Trochopteris Gardn. 1842. Bak. 1868.

§ 2. **Hemianeimia** Prantl 1881 (?).
a. Anemidictyon J. Sm. 1842 (phyllitidis: sp. venis anastom.).
b. Aneimiaebotrys Fée 1869 (aspera).
c. Coptophyllum Gardn. 1842 (buniifolia) = Mohria sp. J. Sm. 1843.

§ 3. **Ornithopteris** (adiantifolia).
Ornithopteris Bernh. 1806 = Anemirhiza J. Sm. 1854 = Aneimia § Aneimiorrhiza **Diels** 1900.

VIII. OSMUNDACEAE.

1. TODEA Willdenow 1802. — Species: 1.

Typ. Acrostichum barbarum L.

Todea § Eutodea Bak. 1868.

2. LEPTOPTERIS Presl 1845. — Species: 7.

Typ. Todea hymenophylloides Rich.

Todea § Leptopteris Bak. 1868. Osmunda sp. Armstr. 1881.

3. OSMUNDA Linné 1753 reduct. — Species: 9.

Typ. O. regalis L.

1801. Struthiopteris Bernh.
1802. Aphyllocalpa Cav. (§ 3).
1836. Plenasium Pr. (§ 1).
1845. Osmundastrum Pr. (Osmunda §). (§ 2).
1847. Osmundastrum Pr. (1845 §). (§ 2).

§ 1. **Plenasium** (banksiifolia).
Plenasium Pr. 1836.

§ 2. **Osmundastrum** Pr. 1845 (cinnamomea).
Osmundastrum Pr. 1847 pt.

§ 3. **Euosmunda** (regalis).
Aphyllocalpa Cav. 1802.

B. HYDROPTERIDINEAE.

IX. SALVINIACEAE.

1. AZOLLA Lamarck 1783. — Species: 5.

Typ. A. filiculoides Lam.

1808. Carpanthus Rafin.
1834. Rhizosperma Meyen.

§ 1. **Euazolla** (filiculoides).
§ 2. **Rhizosperma** (pinnata).
Rhizosperma Meyen 1834.

2. SALVINIA (Micheli) Adanson 1763. — Species: 13

Typ. Marsilea natans L.

X. MARSILEACEAE.

1. MARSILEA Linné 1753. — Species: 56.

Typ. M. quadrifolia L.

1763. Lemma Adans.
1775. Zalusianskya Necker.

§ 1. (polycarpa).
§ 2. **Eumarsilea** (quadrifolia).

2. REGNELLIDIUM Lindman 1904. — Species: 1.

Typ. R. diphyllum Lindm.

3. PILULARIA Linné 1753. — Species: 6.

Typ. P. globulifera L.

1891. Calamistrum O.Ktze. (L. 1737).

---

2. MARATTIALES.

XI. MARATTIACEAE.

1. ANGIOPTERIS Hoffmann 1796. — Species: 62.

Typ. Polypodium evectum Forst.

1796. Angiopteris Hoffm. 1796 (non Adans. 1763).
1803. *Clementea* Cav. (nom. opt.).
1845. Psilodochea Pr. (§ 2 a).
— Pseudangiopteris Pr. (Angiopteris §). (§ 2).

§ 1. **Euangiopteris** (evecta).
§ 2. **Pseudangiopteris** Pr. 1845 (crassipes).
a. Psilodochea Pr. 1845 (salicifolia).

2. ARCHANGIOPTERIS Christ et Giesenhagen 1899. — Species: 1.

Typ. A. Henryi Christ et Gies.

3. MARATTIA Swartz 1788. Species: 28.

Typ. M. alata Sw.

(1786. Celanthera Thouin ?).
1788. Marattia Sw.
1789. Myriotheca Juss.
1842. Eupodium J. Sm. (Kaulfussii).
1845. Discostegia Pr. (alata).
— Gymnotheca Pr. (cicutifolia).
— Stibasia Pr. (Douglasii).

4. CHRISTENSENIA Maxon 1905. — Species: 1.

Typ. Aspidium aesculifolium Bl.

1828. Kaulfussia Bl. Bak. 1868. Christ 1897. **Diels** 1900 (non Dennst. 1818: Gen. Polygalac. nec Nees 1820: gen. Compositarum).
1836. Macrostoma Griff. (non Hedw. 1806).

5. DANAEA Smith 1793. — Species: 26.

Typ. Acrostichum nodosum L.

1845. Arthrodanaea Pr. (Leprieurii).
— Danaeopsis Pr. (paleacea).
1845. Heterodanaea Pr. (stenophylla).
— Holodanaea Pr. (alata).

---

3. OPHIOGLOSSALES.

XII. OPHIOGLOSSACEAE.

1. OPHIOGLOSSUM Linné 1753 reduct. — Species: 43.

Typ. O. vulgatum L.

1828. Ophioderma Bl. (Ophioglossum §). (§ 2).
1836. Ophioderma Endl. (Bl. 1828 §). (§ 2).
1845. Cheiroglossa Pr. (§ 3).
— Rhizoglossum Pr. (§ 1 a).
1847. Cassiopteris Karst. (§ 3).
1883. Paraneura Prantl (Ophioglossum §). (§ 1 b).
— Ptiloneura Prantl (Ophioglossum §). (§ 1 c).

§ 1. **Euophioglossum** (vulgatum).
a. Rhizoglossum Pr. 1845 (Bergianum).
b. Paraneura Prantl 1883.
c. Ptiloneura Prantl 1883.

§ 2. **Ophioderma** Bl. 1828 (pendulum).
Ophioderma Endl. 1836.

§ 3. **Cheiroglossa** (palmatum).
Cheiroglossa Pr. 1845 = Cassiopteris Karst. 1847.

2. BOTRYCHIUM Swartz 1801. — Species: 34.

Typ. Osmunda lunaria L.

1753. Osmunda sp. L.
1801. Botrychium Sw.
— Botrypus Rich.
1867. Osmundopteris Milde (Botrychium §).
1883. Phyllotrichium Prantl (Botrychium §).
1905. Sceptridium Lyon.

§ 1. **Eubotrychium** (lunaria).
§ 2. **Phyllotrichium** Prantl 1883 (ternatum).
— Sceptridium Lyon 1905.
a. Osmundopteris Milde 1867 (virginianum).

3. HELMINTHOSTACHYS Kaulfuss 1822. 1824. — Species: 1.

Typ. Osmunda zeylanica L.

1825. Botryopteris Pr.
1827. Ophiala Desv.

Summa generum specierumque.

Fam.	I.	Hymenophyllaceae.	Verae Gen.	2	Sp. 459
	—		Dubiae	2	3
—	II.	Cyatheaceae		7	456
—	III.	Polypodiaceae		114	4527
—	IV.	Parkeriaceae		1	1
—	V.	Matoniaceae		1	2
—	VI.	Gleicheniaceae		2	80
—	VII.	Schizaeaceae		4	118
—	VIII.	Osmundaceae		3	17
—	IX.	Salviniaceae		2	18
—	X.	Marsileaceae		3	63
—	XI.	Marattiaceae		5	118
—	XII.	Ophioglossaceae		3	78
			Genera:	**149**	Sp. **5940**

Summa synonymorum generum	670	
— — specierum		15787
— specierum dubiarum et hortulanorum		953
	819	22680

Summa nominum omnium in Indice enumeratorum:

23.499.

ENUMERATIO
GENERUM SPECIERUMQUE
ALPHABETICA

ENUMERATIO GENERUM SPECIERUMQUE ALPHABETICA

ABACOPTERIS Fée, Congr. sci. de France Xme sess. **1**. 178. 1843; Gen. Fil. 309. 1850-52 = **Dryopteris.**

elegans Fée, 1 mém. 14 t. 1 f. 23. 1844 (nomen); Gen. 310. 1850-52 — Cochinchina. — Nephrodium Moore 1857.

glandulosa Fée, Gen. 310. 1850-52.

Philippinarum Fée, Congr. sci. de France X sess. **1**. 178. 1843; Gen. 310 = [? D. glandulosa.

simplicifolia Fée, Gen. 310. 1850-52 = D. canescens.

truncata Fée, Gen. 310. 1850-52 = D. lineata.

ABRODICTYUM Presl, Hymenoph. 20. 1843 = **Trichomanes.**

Cumingii Pr. Hym. 20 t. 7. 1843.

Acarpacrium Prantl, Hymenoph. 48. 1875 *(Ptilophyllum §)* = Trichomanes.

ACHOMANES Necker, Element. bot. **3**. 313. 1790 = **Trichomanes.**

Trichomanes § Achomanes Pr. Hym. 15. 1843.

ACONIOPTERIS Presl, Tent. Pterid. 236. 1836 = **Elaphoglossum.**

glabrescens Pr. Epim. 166. 1849 = E. tomentosum.

gorgonea J. Sm. Hist. Fil. 128. 1875.

gorgonea Bedd. Handb. 427. 1883 = Asplenium sp. ster. (nidus?)

longifolia Fée, Acrost. 80 t. 41. 1845.

nervosa J. Sm. Ferns br. and for. 107. 1866.

obtusa Fée, Acrost. 80 t. 40 f. 2. 1845 = E. gorgoneum.

Richardi Bory; Fée, Acrost. 80 t. 16 f. 4. 1845.

subdiaphana Pr. Tent. 236. 1836 = E. nervosum.

ACROPHORUS Presl, Tent. Pterid. 93. 1836 — (Polypodiaceae Gen. **6**.)

adiantoides Moore, Ind. XCI. 1857 = Lindsaya.

affinis Moore, Proc. Linn. Soc. **2**. 286. 1854 = Davallia hymenophylloides.

assamicus Bedd. Ferns br. Ind. t. 94. 1866 = Davallia.

bifidus Moore, Ind. XCI. 1857 = Odontosoria.

chaerophyllus Moore, Proc. Linn. Soc. **2**. 286. 1854 = Davallia pulchra.

cuneifolius Moore, Ind. XCI. 1857 = Lindsaya adiantoides.

falcinellus Moore, Ind. 2. 1857 = Davallia.
fallax Regel, Gartenfl. **6**. 343. 1857 = Dryopteris amplissima.
Goudotianus Moore, Ind. 2. 1857 = Lindsaya.
hemipterus Moore, Ind 295. 1861 = Lindsaya repens var.
hispidus Moore, Proc. Linn. Soc. **2**. 286. 1854 = Leptolepia novae zelandiae.
Hookeri Moore, Ind. 2. 1857 (nomen) = Davallia Clarkei.
Hookeri Bedd. Ferns br. Ind. t. 95. 1866 = Polypodium dareiforme.
hymenophylloides Moore, Ind. 2. 1857 = Lindsaya.
hymenophyllus Parish; Bedd. Ferns br. Ind. t. 96. 1866 = Davallia.
immersus Moore, Proc. Linn. Soc. **2**. 286. 1854 = Davallia.
Imrayanus Moore, Ind. 295. 1861 = Saccoloma.
jamaicensis Moore, Proc. Linn. Soc. **2**. 286. 1854 =? Microlepia speluncae.
?javensis Moore, Ind. 2. 1857 = Dryopteris punctata.
membranulosus Moore, Proc. Linn. Soc. **2**. 286. 1854 = Davallia.
multidentatus Bedd. Ferns br Ind. Correct. 1870 = Polypodium dareiforme.
nodosus Pr. Tent. 94 t. 3 f 2. 1836 = A stipellatus.
Parkeri Moore, Ind. XCI. 1857 = Lindsaya.
parvulus Moore, Proc. Linn. Soc. **2**. 286. 1854 = Davallia.
pseudocystopteris Moore, Ind. 3. 1857 = Davallia.
pulcher Moore, Ind. 3. 1857 = Davallia.
repens Moore, Ind. XCI. 1857 = Lindsaya.
stipellatus (Wall.) Moore, Gard. Chr. **1854**. 135 — India bor.-Malesia. Davallia Wall. List n. 260. 1828 (nomen); Aspidium nodosum Bl. Enum. 171. 1828 (non Willd. 1810); Acrophorus Pr. 1836. Chr. 285. NPfl. 164; Davallia? Hk. 1846. HB. 92; Cystopteris Mett. 1864; Leucostegia Bedd. 1876.
tenuifolius Moore, Ind. XCI. 1857 = Lindsaya.
Thomsoni Moore, Ind. 4. 1857 (nomen) = Davallia multidentata.

Acropteridastrum Fée, Gen. Fil. 190. 1850-52 *(Asplenium §)* = Asplenium.
ACROPTERIS Link, Hort. Berol. **2**. 55. 1833.
australis Link, Hort. Berol. **2**. 56. 1833 } = Actiniopteris australis.
radiata Link, Hort. Berol. **2**. 56. 1833 }
Seelosii Heufl. Verh. zool. bot. Ges. **6**. 345. 1856 = Asplenium.
septentrionalis Link, Hort. Berol. **2**. 56. 1833 = Asplenium.

Acropterygium Diels, Nat. Pfl. **1**⁴. 353. 1900 *(Gleichenia §)* = Gleichenia.

ACROSTICHUM Linné (Gen. Plant. 785. 1737); Spec. plant. **2**. 1067. 1753 — (Polypodiaceae Gen. **109**).
accedens Mett; Kuhn, Linn. **36**. 49. 1869 = Elaphoglossum.
achroalepis Bak. JoB. **1880**. 371 = Elaphoglossum.
acrocarpon Bak. Syn. ed. I. 408. 1868 = Elaphoglossum Mathewsii.
acrocarpum Mart. Ic. Cr. Bras. 85 t. 23. 1834 = Elaphoglossum.
actinotrichum Mart. Ic. Cr. Bras. 86. 1834 = Elaphoglossum.
aculeatum L. Syst. Nat. ed. X. **2**. 1320. 1759 = Odontosoria fumarioides.
aculeatum Desv. Berl. Mag. **5**. 310. 1811 = Ceropteris calomelanos var.
aculeatum Racib. Pterid. Buit. 53. 1898 = Stenochlaena.
acuminans Fée, Cr. vasc. Br. **1**. 12 t. 1 f. 2. 1869 = Elaphoglossum plumosum.
acuminatum Willd. sp. **5**. 116. 1810 = Leptochilus nicotianifolius.
acuminatum Juss.; Poir. Enc. Suppl. **1**. 120. 1810 = Elaphoglossum petiolosum.
acuminatum Hk. sp. **5**. 245. 1864 = Polybotrya scandens.
acuminatum Karst.; Christ, Mon. Elaph. 78. 1899 = Elaphoglossum cuspidatum.
acutum Fée; Kuhn, Linn. **36**. 44. 1869 = Elaphoglossum spathulatum.

acutum Fée, Cr. vasc. Br. **2**. 9 t. 83 f. 3. 1872-73 = Elaphoglossum villosum.
adenolepis Kze. Linn. **9**. 27. 1834 = Elaphoglossum.
aemulum Klf. Enum. 63. 1824 = Elaphoglossum.
aemulum Bl. Enum. 101. 1828 = Elaphoglossum laurifolium.
affine Mart. et Gal. Mém. Ac. Brux. **15**. 24 t. 3 f. 1. 1842 = Elaphoglossum.
Afzelii Carr. Cat. Welw. pl. **2**^{2}. 277. 1901 = Leptochilus acrostichoides.
alatum Roxb. Calc. Journ. **4**. 480. 1844 = Stenochlaena sorbifolia var. 3.
alatum Fée, Acrost. 35. t 5 f. 2. 1845 = Elaphoglossum pteropus.
albescens Sod. Rec. 75. 1883 = Elaphoglossum.
albidulum Cav.; Sw. Syn 16, 205 t. 1 f. 2. 1806 = Pellaea nivea.
album Vell. Fl. flum. **11** t. 91. 1827; Arch. Mus. Nac. Rio Jan. **5**. 453. 1881 = Ceropteris calomelanos.
alcicorne Willem. Usteri Ann. **4**^{12}. 61. 1796 = Platycerium.
alcicorne Wall. List n. 19. 1828 = Platycerium Wallichii.
alienum Sw. Schrad. Journ. **1800**^{2}. 13. 1801 = Leptochilus.
alismaefolium Fée, Acrost. 28 t. 3. 1845 = Elaphoglossum latifolium var.
alpestre Gardn.; Field. et Gardn. Sert. pl. t. 25. 1844 = Elaphoglossum.
alpinum Bolton, Fil. Brit. 76 t. **42**. 1790 = Woodsia.
ambiguum Mett.; Christ, Mon. Elaph. 60. 1899 = Elaphoglossum flaccidum.
amplissimum Fée, Cr. vasc. Br. **1**. 5 t. 6. 1869 = Elaphoglossum.
amygdalifolium Mett.; Kuhn, Linn. **36**. 41. 1869 = Elaphoglossum.
andicola Fée, Acrost. 28 t. 2 f. 1. 1845 = Elaphoglossum latifolium var.
angulatum Bl. Enum. 101. 1828 = Elaphoglossum.
angustatum Schrad. Gött. gel. Anz. **1818**. 915 = Elaphoglossum conforme var.
angustissimum Fée, 10 mém. 6 t. 29 f. 3. 1865 = Elaphoglossum petiolatum var.
angustum Fée, Cr. vasc. Br. **2**. 12 t. 86 f. 3. 1872-73 = Elaphoglossum.
antrophyoides Bak. Journ. Linn. Soc. **22**. 231. 1886 = Leptochilus.
aphlebium Kze.; Kl. Linn. **20**. 419. 1847 = Elaphoglossum conforme.
apiifolium Hk. sp. **5**. 248. 1864 = Polybotrya.
apodum Klf. Enum. 59. 1824 = Elaphoglossum.
appendiculatum Willd. sp. **5**. 114. 1810 = Polybotrya.
araneosum Eat. Proc. Amer. Acad. **22**. 461. 1887 = Elaphoglossum.
areolatum L. sp. **2**. 1069. 1753 = Woodwardia.
argyrophyllum Sod. Rec. 85. 1883; Cr. vasc. quit. 477. 1893 = Elapho-[glossum Lindigii.
articulatum Hk. sp. **5**. 247. 1864 = Polybotrya.
aspidioides Bak. Syn. 414. 1868 = Polybotrya.
aspidiolepis Bak. JoB. **1880**. 371 = Elaphoglossum.
aspleniifolium Bory, Bél. Voy. Bot. **2**. 21 t. 3. 1833 = Polybotrya appendiculata.
assurgens Bak. Syn. ed. II. 408. 1874 = Elaphoglossum Mathewsii.
asterolepis Bak. JoB. **1880**. 371 = Elaphoglossum.
attenuatum Fée, Gen. 43. 1850-52 (nomen); 6 mém. 1 t. 1 f. 1. 1853 = Elaphoglossum hybridum var. melanopus.
attenuatum Kze.; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860; Kuhn, Linn. **36**. 56. 1869 = Elaphoglossum firmum.
Aubertii Desv. Berl. Mag. **5**. 309. 1811 = Elaphoglossum.
aureo-nitens Hk. Ic. pl. t. 933. 1854 = Trachypteris pinnata.
aureum L. sp. **2**. 1069. 1753 HB. 423. NPfl. 336 — Trop.-Subtrop.
Chrysodium Mett. 1856. Chr. 47; C. vulgare Fée 1845 — Acrostichum inaequale Willd. 1810; Chrysodium Fée 1845 — Acrostichum speciosum Willd. 1810; Chrysodium Fée 1845 — Acrostichum daneaefolium Langsd.

et Fisch. 1810; Chrysodium Fée 1845 — Acrostichum fraxinifolium R. Br. 1810; Chrysodium Fée 1845 — Acrostichum obliquum Bl. 1828 — Chrysodium Urvillei, scalpturatum, hirsutum Fée 1845 — C. lomarioides Jenm. 1885; Acrostichum Jenm. 1898 (non Bory 1833).

aureum Cav. Anal. Cienc. **4**. 104. 1801 = Ceterach.

auricomum Kze. Linn. **9**. 28. 1834 = Elaphoglossum.

auriculatum Lam. Enc. **1**. 36. 1783 = Leptochilus.

auritum Sw. Schrad. Journ. **1800**^{2}. 12. 1801 = Stenosemia.

auritum Poir. Enc. Suppl. **1**. 122. 1810 = Leptochilus auriculatus.

australe L.fil. Suppl. 444. 1781 = Actiniopteris.

axillare Cav. Anal. Hist. Nat. **1**. 101. 1799 = Leptochilus.

backhousianum Bak. Ann. Bot. **5**. 491. 1891 = Elaphoglossum.

Bakeri Sod. Rec. 77. 1883 = Elaphoglossum.

Banksianum Fée, Acrost. 64. 1845 = Elaphoglossum.

barbarum L. sp. **2**. 1072. 1753 = Todea.

barbatum Karst.; Mett Ann. sc. nat. V. **2**. 200. 1864 (nomen); Karst. Fl. Col. **2**. 155 t. 181. 1869 = Elaphoglossum scolopendrifolium.

Barteri Bak. Syn. 519. 1874 = Elaphoglossum.

Beaurepairii Fée, Cr. vasc. Br. **2**. 10 t. 79 f. 2. 1872-73 = Elaphoglossum [Aubertii.

beccarianum Bak. Malesia **3**. 27. 1886 = Elaphoglossum.

Bellermannianum Kl. Linn. **20**. 426 1847 = Elaphoglossum.

Bernoullii Kuhn; Christ, Bull. Soc. bot. Belg. **35**. 244. 1896 = Leptochilus.

bicolor Cav. Anal. Hist. Nat. **1**. 103. 1799; Descr. 238. 1802 — Ins. Mariannae — (prob. Cyclophorus sp.).

bicuspe Hk. sp. **5**. 271. 1864 = Cheiropleuria

bifidum Poir. Enc. Suppl. **1**. 126. 1810 = Schizaea.

biforme Sw. Schrad. Journ. **1800**^{2}. 11. 1801 = Platycerium coronarium.

bifurcatum Cav. Anal. Hist. Nat. **1**. 105. 1799; Descr. 240. 1802 = Platycerium alcicorne.

bifurcatum Sw. Schrad. Journ. **1800**^{2}. 13. 1801 = Elaphoglossum furcatum.

Blanchetii Mett.; Kuhn, Linn. **36**. 55. 1869 = Elaphoglossum.

blepharodes Fée, Acrost. 48 t. 24 f. 3. 1845 = Elaphoglossum hybridum var.

Blumeanum Fée, Acrost. 62. 1845 = Elaphoglossum.

Blumeanum Hk. sp. **5**. 268. 1864 = Leptochilus lomarioides.

Boivini Mett.; Kuhn, Fil. Afr. 43. 1868 = Elaphoglossum Aubertii.

Boivini Welw.; Carr. Cat. Welw. pl. **2**^{2}. 276. 1901 = Leptochilus.

bonariense Willd. sp. **5**. 114. 1810 = Notholaena.

boragineum Sod. Rec. 80. 1883 = Elaphoglossum.

borbonicum Bak. Ann. Bot. **5**. 491. 1891 = Elaphoglossum.

Borneense Burck, Ann. Jard. Buit. **4**. 99. 1884 = Elaphoglossum.

Boryanum Fée, Acrost. 40 t. 1. 1845 = Elaphoglossum.

botryoides Bak. JoB. **1881**. 206 = Polybotrya.

Bourgaeanum Fourn. Mex. pl. **1**. 64 t. 1 f. 1. 1872 = Elaphoglossum tambillense.

brachynevron Fée, Acrost. 49 t. 22 f. 1. 1845 = Elaphoglossum.

brachystachyum Racib. Pterid. Buit. 51. 1898 = Hymenolepis.

brasiliense Spr.; Kuhn, Linn. **36**. 54. 1869 = Elaphoglossum simplex.

? Breutelianum Kze. Farnkr. **2**. 3 t. 102. 1848 = { Elaphoglossum petiolatum.
? Polypodium laevigatum.

brevipes Kze. Ind. Fil. Hort. Lips. 1845; Fée, Acrost. 29. 1845 = Elaphoglossum.

Brightiae F. Muell. Fragm. **7**. 119. 1870 = Stenochlaena sorbifolia var. 13.

brunneum Willd. sp. **5**. 113. 1810 = Leptochilus alienus.
Burchellii Bak. Syn. 401. 1868 = Elaphoglossum.
buxifolium Kze. Farnkr. **1**. 171 t. 72. 1845 = Stenochlaena pollicina.
Caenopteris Kze.; Hk. sp. **5**. 256. 1864 = Polybotrya serratifolia.
caespitosum Sod. Rec. 74. 1883 = Elaphoglossum.
Calaguala Kl. Linn. **20**. 421. 1847 = Elaphoglossum huacsaro.
callaefolium Bl. Enum. 100. 1828 = Elaphoglossum.
callaefolium Link, Hort. Berol. **2**. 46. 1833 = Elaphoglossum brevipes.
callipteris Ehrh. Beitr. **3**. 77. 1788 = Dryopteris cristata.
callolepis Fée, 8 mém. 69. 1857 = Elaphoglossum.
calomelanos L. sp. **2**. 1072. 1753 = Ceropteris.
Calomelas Sw. Obs. bot. 396. 1791 = Ceropteris calomelanos.
calophyllum Kze. Linn. **9**. 27. 1834 = Elaphoglossum.
canaliculatum Hk. sp. **5**. 247. 1864 = Polybotrya.
canariense Willd. sp. **5**. 121. 1810 = Notholaena Marantae.
cardiophyllum Hk. Ic. pl. t. 715. 1845 = Elaphoglossum.
cartilagineum Mett.; Kuhn, Linn. **36**. 58. 1869 = Elaphoglossum.
castaneum Bak. JoB. **1877**. 166 = Elaphoglossum.
catanense Cosentini, Atti Acad. Gioen. **2**. 207, 218 c. tab. 1827 = Notholaena vellea.
caudatum Cav. Descr. 242. 1802 = ? Leptochilus alienus.
caudatum Hk. Ic. pl. t. 215. 1840 = Elaphoglossum petiolosum.
caudatum Hk. sp. **5**. 244. 1864 = Polybotrya.
caulolepia Karst. Fl. Col. **1**. 121 t. 60. 1860 = Elaphoglossum.
cayennense Pr. Epim. 181. 1849 = A. aureum.
celebicum Bak. Kew Bull. **1901**. 145 = Leptochilus.
cervinum Sw. Syn. 14, 200. 1806 = Polybotrya.
chartaceum Bak.; Jenm. JoB. **1882**. 325 = Elaphoglossum.
chrysoconium Desv. Prod. 212. 1827 — Peru. —. Gymnogramma Moore 1857 — (Ceropteris sp. dub.).
chrysolepis Fée, Cr. vasc. Br. **1**. 10 t. 2 f. 1. 1869 = Elaphoglossum plumosum.
chrysolepis Sod. Cr. vasc. quit. 485. 1893 = Polybotrya altescandens.
chrysophyllum Sw. Schrad. Journ. **1800**[2]. 14. 1801 = Ceropteris calomelanos var.
cicutarium Hk. sp. **5**. 257. 1864 = Stenosemia aurita var.
ciliare Thouars, Fl. Trist. d'Ac. 32. 1804 = Elaphoglossum spathulatum.
ciliare Carm. Tr. Linn. Soc. **12**. 510. 1818 = Elaphoglossum hybridum.
ciliatum Desv. Berl. Mag. **5**. 310. 1811 = Elaphoglossum hybridum.
ciliatum Pr. Rel. Haenk. **1**. 15. 1825 = Elaphoglossum.
cinnamomeum Bak. Syn. 409. 1868 = Elaphoglossum.
citrifolium L. sp. **2**. 1067. 1753 = Anetium.
cladorrhizans Spr. Nova Acta **10**. 225. 1821 = Leptochilus alienus.
clarenceanum Bak. Kew Bull. **1896**. 42 = Elaphoglossum.
cochleariaefolium Fée, Gen. 42. 1850-52; 6 mém. 2 t. 1 f. 3. = Elaphoglossum cardiophyllum.
cochleatum Bory; Fée, Acrost. 63 t. 16 f. 1. 1845 = Elaphoglossum pilosum.
commutatum Mett.; Kuhn, Ann. Lugd. Bat. **4**. 292. 1869 = Elaphoglossum laurifolium.
compactum Mett.; Kuhn, Linn. **36**. 59. 1869 = Elaphoglossum conforme.
conforme Sw. Syn. 10, 192 t. 1 f. 1. 1806 = Elaphoglossum.
conforme Bl. Fl. Javae Fil. 23 t. 5. 1828 = Elaphoglossum laurifolium.

conopodium van Houtte, Cat. 1868; Christ, Mon. Elaph. 49 = Elaphoglossum conforme.
consobrinum Kze. Flora **1839**[1]. Beibl. 44 = Elaphoglossum latifolium.
contaminans Wall. List n. 22. 1828 = Leptochilus virens.
contaminoides Lindm. Hedwigia **43**. 311. 1904 = Leptochilus.
contractum Wall. List n. 2166. 1829 — India austr.
cordatum Thbg. Prod. Fl. Cap 171. 1800 = Ceterach.
Corderoanum Sod. Cr. vasc. quit. 470. 1893 = Elaphoglossum.
coriaceum Wall. List n. 14. 1828 = ? Elaphoglossum Sieberi.
corpulentum Kze.; Kuhn, Linn. **36**. 61. 1869 = Elaphoglossum pachyphyllum.
costatum Wall. List n. 26. 1828; Hk. sp. **5**. 262. 1864 = Leptochilus scalpturatus.
costulatum Cesati, Rend. Acad. Napoli **16**. 27, 30. 1877 = Polypodium.
craspedariaefolium Fée, Cr. vasc. Br. **1**. 261. 1869 = Elaphoglossum squamipes.
crassifolium Gaud. Freyc. Voy. Bot. 303. 1827 = Elaphoglossum reticulatum.
crassifolium Wall.; Pr. Tent. 241. 1836 = A. aureum.
crassinerve Kze. Ind. Fil. Hort. Lips. 1845; Fée, Acrost. 29. 1845 = Elaphoglossum latifolium var.
crassipes Hier. Engl. Jahrb. **22**. 409. 1896 = Elaphoglossum.
crinitum L. sp. **2**. 1068. 1753 = Elaphoglossum.
crinitum Mart. et Gal. Mém. Ac. Brux. **15**. 25. 1842 = Elaphoglossum hybridum var. blepharodes.
crispatulum Wall. List n. 24. 1828 = Leptochilus virens var.
crispatulum Fée, Gen. 42. 1850-52; 6 mém. 2 t. 1 f. 2 = Elaphoglossum ovatum.
crispum Vill. Hist. pl. Dauph. **3**. 838. 1789 = Cryptogramma.
cruciatum L. sp. **2**. 1072. 1753 (Plum. t. 48 B) — Hispaniola — Gymnogramma Moore 1857.
Cruegerianum Kze.; Kuhn, Engl. Jahrb. **24**. 139 1897 = Elaphoglossum.
Cuacsaro Bert. Opusc. sci. Bol. **1**. 241 t. 8. 1817 = Elaphoglossum huacsaro.
cubense Mett.; Kuhn, Linn. **36**. 46. 1869 = Elaphoglossum.
cultratum Bak. Syn. 524. 1874 = Leptochilus lomarioides.
Cumingii Fée, Acrost. 34. 1845 = Elaphoglossum decurrens var.
Curtisii Bak. Ann. Bot. **5**. 489. 1891 = Elaphoglossum.
Curupirae Lindm. Hedwigia **43**. 311. 1904 = Leptochilus.
curvans Kze. Linn. **9**. 30. 1834 = Elaphoglossum.
cuspidatum Willd. sp. **5**. 106. 1810 = Elaphoglossum.
cylindricum Hk. sp. **5**. 246. 1864 = Polybotrya osmundacea var.
daneaefolium Langsd. et Fisch. Ic. Fil. 5 t. 1. 1810 = A. aureum
Deckenii Kuhn, Fil. Deck. 16. 1867 = Elaphoglossum
decoratum Kze. Linn. **9**. 25. 1834 = Elaphoglossum.
decrescens Bak. Syn. 412. 1868 = Stenochlaena.
decurrens Desv. Berl. Mag. **5** 310. 1811 = Elaphoglossum.
decurrens Wall. List n. 18. 1828; Thwait. En. pl. Zeyl. 381. 1864 = Leptochilus.
decurrens Benth. Fl. Hongk. 443. 1861 = Leptochilus Harlandii.
deltoideum Sod. Cr. vasc. quit. 466. 1893 = Elaphoglossum.
dendricolum Bak. Syn. 521. 1874 = Elaphoglossum.
denticulatum Colmeiro, Enum. Cr. Esp. y Port. **1**. 8. 1867 — Notholaena vellea var.
deorsum Karst. Fl. Columb. **1**. 118 t. 59. 1860 = Elaphoglossum lepidotum.
Deppei Kze.; Christ, Mon. Elaph. 118. 1899 = Elaphoglossum gratum.
dichotomum L. sp. **2**. 1068. 1753 = Schizaea.

dichotomum Forsk. Fl. Aeg. Arab. 184. 1775 = Actiniopteris australis.
dicksonioides Desv. Prod. 212. 1827 = ? Polybotrya osmundacea.
didynamum Fée, Acrost. 57 t. 16 f. 2. 1845 = Elaphoglossum conforme.
digitatum L. sp. **2**. 1068. 1753 = Schizaea.
dimorphum Hk. et Grev. Ic. Fil. t. 145. 1829 = Elaphoglossum.
discolor Kuhn, Linn. **36**. 53. 1869 = Elaphoglossum.
dissimile Kze. Linn. **9**. 28. 1834 = Elaphoglossum.
diversifolium Bl. Enum. 103. 1828 = Leptochilus.
Dombeyanum Fée, Acrost. 59 t. 17 f. 2. 1845 = Elaphoglossum muscosum.
drynarioides Hk. sp. **5**. 282. 1864 = Dryostachyum.
dubium Poir. Enc. Suppl. **1**. 120. 1810 = ? Cyclophorus spissus.
durum Kze. Linn. **22**. 575. 1849 = Elaphoglossum simplex var. martinicense.
Eatoni Bak. Ann. Bot. **5**. 493. 1891 } = Elaphoglossum
Eatonianum E. Britton, Bull. Torr. Cl. **15**. 252. 1888 } Eatonianum.
ebeneum L. sp. **2**. 1071. 1753 = Ceropteris calomelanos.
Eggersii Bak. JoB. **1888**. 34 = Elaphoglossum.
elegans Vahl, Symb. **2**. 104 t. 50. 1791 = Schizaea.
elegans Fée, Cr. vasc. Br. **2**. 11 t. 85 f. 1. 1872-73 = Elaphoglossum plumosum.
ellipticum Fée, Acrost. 30 t. 4. 1845 = Elaphoglossum.
elongatum Kze. Linn. **9**. 31. 1834 = Elaphoglossum.
emarginatum Buch.; Roxb. Calc. Journ. **4**. 480 t. 27. 1844 = A. aureum.
Engelii Karst. Fl. Col. **1**. 118 t. 59. 1860 = Elaphoglossum.
erinaceum Fée, Acrost. 41. 1845 = Elaphoglossum hybridum.
erythrodes Kze. Flora **1839**[1]. Beibl. 46 = Stenochlaena marginata.
erythrolepis Fée, Acrost. 60. 1845 = Elaphoglossum.
erythrotrichum Salom. Nom. 5. 1883 = Elaphoglossum.
eximium Mett. Ann. sc. nat. V. **2**. 199. 1864 = Elaphoglossum.
exsculptum Bak. JoB. **1888**. 326 = Leptochilus.
falcatum Fée, Acrost. 44 t. 21 f. 1. 1844 = Elaphoglossum obductum.
fasciculatum (Fourn.) C. Chr. Ind. 9. 1905 — N. Caledonia.
Chrysodium Fourn. Ann. sc. nat. V. **18**. 272. 1873 — (an A. aureum?).
Feei Bory: Fée, Acrost. 48 t. 18 f. 2. 1845 = Elaphoglossum.
Féejanum Ettingsh. Denkschr. Ak. Wien **22**. 54. t. 2 f. 13. 1864 = Leptochilus lanceolatus.
Feejeense Hk. sp. **5**. 199. 1864 = Elaphoglossum.
Fendleri Bak. JoB. **1887**. 100 = Leptochilus.
Fendleri Mett. (Christ. Mon. Elaph. 44. 1899) = Elaphoglossum conforme var.
ferrugineum L. sp. ed. II. **2**. 1525. 1763 = Polypodium polypodioides.
ferrugineum Linden, Cat. 1856; Bak. Syn. 522. 1874 = Elaphoglossum.
filare Forsk. Fl. Aeg. Arab. 184. 1775 = ? Asplenium praemorsum.
fimbriatum Cav. Anal. Hist. Nat. **1**. 102. 1799 — Peru. — (Elaphoglossum sp.; forte = E. scolopendrifolium).
fimbriatum Sod. Rec. 84. 1883 = Elaphoglossum ecuadorense.
Finlaysonianum Wall. List n. 2162. 1829 = Leptochilus auriculatus.
firmum Mett.; Kuhn, Linn. **36**. 55. 1869 = Elaphoglossum.
fistulosum Poir. Enc. Suppl. **1**. 125. 1810 = Schizaea.
flabellatum H. B. Willd. sp. **5**. 110. 1810 = Elaphoglossum.
flabellifolium Link, Fil. sp. 165. 1841 — Hort.
flaccidum Fée, Acrost. 35 t. 7 f. 2. 1845 = Elaphoglossum.
flagelliferum Wall.; Hk. et Grev. Ic. Fil. t. 23. 1827 = Leptochilus heteroclitus.
flavens Sw. Syn. 16, 204. 1806 = Pellaea.

floridum Poir. Enc. Suppl. **1**. 122. 1810 = Stenosemia aurita.
fluviatile Hk. sp. **5**. 274. 1864 = Leptochilus.
foeniculaceum Hk. et Grev. Ic. Fil. t. 119. 1829 = Elaphoglossum.
Fonki Phil. Linn. **29**. 104. 1857 = Elaphoglossum Mathewsii.
formosum Pr. Del. Prag. **1**. 160. 1822 = A. aureum.
Forsteri Montr.; Salom. Nom. 6. 1883 (nomen) — N. Caledonia.
fractiseriale Bak. Syn. 414. 1868 = Polybotrya.
Fraseri Mett.; Kuhn, Linn. **36**. 43. 1869 = Elaphoglossum.
fraxinifolium R. Br. Prod. Fl. N. Holl. 145. 1810 = A. aureum.
fraxinifolium Pr. Del. Prag. **1**. 161. 1822 = Leptochilus serratifolius.
frigidum Linden, Cat. 1856 = Elaphoglossum perelegans.
fuciforme Wall. List n. 20. 1828 = Platycerium coronarium.
fulvum Mart. et Gal. Mém. Ac. Brux. **15**. 24 t. 3 f. 2. 1842 = Elaphoglossum [vestitum.
Funckii Fée, Acrost. 36 t. 6 f. 1. 1845 = Elaphoglossum.
furcatum L. Syst. Nat. ed. X. **2**. 1321 1759 = Gleichenia.
furfuraceum Mett.; Kuhn, Linn. **36**. 51. 1869 = Elaphoglossum.
furfuraceum Bak. JoB. **1877**. 166 = Elaphoglossum vulcanicum.
Gaboonense Hk. sp. **5**. 270. 1864 = Leptochilus.
Galvinii Glaziou; Bak. Ann. Bot. **5**. 492. 1891 = Elaphoglossum angustum.
Gardneri Hk. sp. **5**. 233. 1864 } = Elaphoglossum
Gardnerianum Kze.; Fée, Acrost. 55 t. 15 f. 3. 1845 } Gardnerianum.
Gayanum Fée, Acrost. 37 t. 19 f. 2. 1845 = Elaphoglossum.
gillianum Bak. JoB. **1882**. 310 = Trachypteris pinnata.
glabellum Kl. Linn. **20**. 421. 1847 = Elaphoglossum.
glabratum Mett. Ann. sc. nat. IV. **15**. 55. 1861 = Elaphoglossum.
glabrescens Kuhn, Fil. Afr. 45. 1868 = Elaphoglossum tomentosum.
glandulosum Carm; Hk. et Gr. Ic. Fil. t. 3. 1827 = Elaphoglossum conforme.
glaucum Cav. Anal. Hist. Nat. **1**. 107. 1799 = Cheilanthes.
glaucum Fée, 8 mém 67. 1857; 10 mém. 4 t. 28 f. 1 = Elaphoglossum conforme.
Glaziovii Fée, Cr. vasc. Br. **1**. 6 t. 1 f. 1. 1869 = Elaphoglossum scolopendrifolium.
glossoides André ap. Christ, Mon. Elaph. 97. 1999 = Elaphoglossum.
glutinosum Spruce ap. Christ, Mon. Elaph. 120. 1899 = Elaphoglossum.
gorgoneum Klf. Enum. 63. 1824 = Elaphoglossum.
gorgoneum Bl. Enum. 101 1828 = Elaphoglossum conforme.
gorgoneum Clarke, Tr. Linn. Soc. II. Bot. **1**. 578. 1880 = Asplenium sp. ster. (nidus?)
gracile Fée, Cr. vasc. Br. **2**. 8 t. 83 f. 2. 1872-73 = Elaphoglossum lineare.
gracile Racib. Pterid. Buitenz. 53. 1898 = Stenochlaena.
gracilipes Fee; Kuhn, Linn. **36**. 47. 1869 = Elaphoglossum.
gramineum Jenm. JoB. **1879**. 263 = Elaphoglossum.
graminoides Sw. Prod. 128. 1788 = Monogramma.
grandifolium Mett.; Kuhn, Linn. **36**. 62. 1869 = Elaphoglossum.
gratum Fée, 8 mém. 69. 1857 = Elaphoglossum.
Guatemalense Kl. Allg. Gartenzeit. **1855**. 66 = Elaphoglossum.
guianense Bak. Fl. bras. **1**². 591. 1870 = Leptochilus.
guineense Carr. Cat. Welw. pl. **2**². 276. 1901 = Lomariopsis.
Hackelianum Sod. Cr. vasc. quit. 491. 1893 — Ecuador. — Polybotrya? sp.
Hamiltonianum Wall. List n. 29. 1828 = Polybotrya appendiculata.

Harlandii Hk, sp. **5**. 274. 1864 = Leptochilus.
Hartii Bak. JoB. **1888**. 371 = Polybotrya.
Hartwegii Fée, Acrost. 53 t. 9 f. 2. 1845 = Elaphoglossum Mathewsii.
hastatum Thbg. Fl. Jap. 331 t. 34. 1784 = Cyclophorus.
hastatum Liebm. Vid. Selsk. Skr. V. **1**. 172 (seors. 20). 1849 = Leptochilus alienus.
Hayesii Mett.; Kuhn, Linn. **36**. 43. 1869 = Elaphoglossum spathulatum var.
Haynaldii Sod. Rec. 80. 1883; Cr. vasc. quit. 452. 1893 = Elaphoglossum
heliconiaefolium Sod. Rec. 80. 1883 = Elaphoglossum. [villosum var.
helleri Und. ap. Heller, Minn. Bot. St. **1**. 771 t. 42. 1897 = Elaphoglossum aemulum.
Herminieri Bory et Fée; Fée, Acrost. 43 t. 11. 1845 = Elaphoglossum.
heteroclitum Pr. Rel. Haenk. **1**. 15 t. 2 f. 2. 1885 = Leptochilus.
heterolepis Fée, Acrost. 56 t. 15 f. 1. 1845 = Elaphoglossum tomentosum.
heterolepis Bak. Syn. 521. 1874 = ? Elaphoglossum squamipes.
heteromorphum Kl. Linn. **20**. 424. 1847 = Elaphoglossum.
heterophyllum L. sp. **2**. 1067. 1753 = Drymoglossum.
heterophyllum Raddi, Pl. Bras. **1**. 5 t. 17. 1825 = Blechnum pteropus.
Heudelotii Hk. sp. **5**. 264. 1864 = Leptochilus.
hirsutum Mett.; Christ, Mon. Elaph. 70. 1899 = Elaphoglossum strictum var.
hirtipes Fée, Cr. vasc. Br. **1**. 238 t. 76 f. 1. 1869 = Elaphoglossum Aubertii.
hirtipes Sod. Cr. vasc. quit. 442. 1893 = Elaphoglossum confusum.
hirtum Sw. Schrad. Journ. **1800^{2}**. 10. 1801 = Elaphoglossum.
Hoffmanni Mett.; Kuhn, Linn. **36**. 56. 1869 = Elaphoglossum.
Hollrungii Bak. Ann. Bot. **5**. 495. 1891 = Leptochilus.
Hookerianum Thwait. En. pl. Zeyl. 380. 1864 = Leptochilus subcrenatus.
horridulum Klf. Enum. 58. 1824 = Elaphoglossum spathulatum var.
Huacsaro Ruiz, Disertac. sobre la raiz de la Ratanhia **42**. 1799; Bertol. Amoen. Ital. 53 t. 2. 1819 = Elaphoglossum.
Humblotii Bak. JoB. **1884**. 144 = Leptochilus.
humile Mett. Ann. sc. nat. V. **2**. 202. 1864 = Elaphoglossum revolutum.
hybridum Bory, Voy. **3**. 95. 1804 = Elaphoglossum.
hybridum Raddi, Opusc. sci. Bol. **3**. 283. 1819 = Elaphoglossum scolopendrifolium.
hymenodiastrum Fée, Cr. vasc. Br. **1**. 3 t. 5. 1869 = Elaphoglossum latifolium.
hyperboreum Liljebl. Vet. Ak. Nya Hdl. **1793**. 201 t. 8 = Woodsia alpina.
hystrix Kze. Linn. **9**. 26. 1834 = Elaphoglossum.
ilvense L. sp. **2**. 1071. 1753 = Woodsia.
impressum Fée, Acrost. 33 t. 5 f. 3. 1845 = Elaphoglossum latifolium.
inaequale Willd. sp. **5**. 117. 1810 = A. aureum.
inaequalifolium Jenm. JoB. **1886**. 273 = Elaphoglossum.
incisum Hk. sp. **5**. 245. 1864 = Polybotrya osmundacea.
insigne Fée, Cr. vasc. Br. **2**. 6 t. 82 f. 1. 1872-73 = Elaphoglossum scolo-
insigne Bak. JoB. **1887**. 167 = Polybotrya andina. [pendrifolium.
intermedium Fée, 8 mém. 69. 1857 = Elaphoglossum elongatum.
inversum Cord. Bull. Soc. sci. et arts Réunion **1890-91**. 84. 1891 = Elaphoglossum.
irregulare Liebm. Vid. Selsk. Skr. V. **1**. 173 (seors. 21). 1849 = Leptochilus
isophyllum Sod. Cr. vasc. quit. 441. 1893 = Elaphoglossum. [alienus.
Jamesoni Hk. et Grev. Ic. Fil. t. 86. 1829 = Elaphoglossum.
japurense Mart. Ic. Cr. Bras. 86 t. 24. 1834 = Stenochlaena marginata.
javense Willd. sp. **5**. 126. 1810 = Notholaena hirsuta.
juglandifolium Klf. Enum. 65. 1824 = A. aureum.

juglandifolium Bak. JoB. **1881**. 207 = Polybotrya Kalbreyeri.
Junghuhnianum Kze Bot. Zeit. **1848**. 101 = Elaphoglossum callifolium.
Karstenianum Kze. Linn. **23**. 214, 298. 1850 = Elaphoglossum petiolatum var.
Klotzschii Moritz; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860 = Elaphoglossum Aubertii.
laciniatum Gilib. Exerc. Phyt. **2**. 555. 1792 = Asplenium septentrionale.
Lagolo Blanco, Fl. Filip. 826. 1837 — Ins. Philipp.
laminarioides Bory; Fée, Acrost. 57 t 12. 1845 = Elaphoglossum.
lanceolatum L. sp. **2**. 1067. 1753 = { Cyclophorus acrostichoides (f. Sw.) / Leptochilus Linnaeanus (f. Fée).
lanceolatum Roxb. ap. Beats. St. Helena 296. 1816 = Elaphoglossum nervosum.
lanceolatum Hk. sp **5**. 276. 1864 = Leptochilus.
lancifolium Desv. Berl. Mag. **5**. 310. 1811 = Elaphoglossum petiolatum.
Langsdorffii Hk. et Grev. Ic. Fil. t. 164. 1829 = Elaphoglossum muscosum.
lanuginosum Desf. Fl. Atlant. **2**. 400 t. 256. 1800 = Notholaena vellea.
lanuginosum Willd. Schr. Akad. Erf. **1802**. 31 t. 3 f. 4 = Notholaena scariosa.
Lastii Bak. Ann. Bot. **5**. 491. 1891 = Elaphoglossum.
latifolium Sw. Prod. 128 1788 = Elaphoglossum.
Laurentii Christ, Bull. Soc. bot. Belg. **36**[2]. 94. 1897 = Leptochilus.
laurifolium Thouars, Fl. Trist. d'Ac. 31. 1804 = Elaphoglossum.
laurifolium Hk. sp. **5**. 250. 1864 = Stenochlaena.
Lechlerianum Mett. Fil. Lechl. **1**. 3. 1856 = Elaphoglossum.
Lechlerianum Hk. sp. **5**. 246. 1864 = Polybotrya.
Lepervanchii Bory; Fée, Acrost. 37 t. 9 f. 1. 1845 = Elaphoglossum conforme.
lepidoglossum Bak. Ann. Bot. **5**. 492. 1891 = Elaphoglossum.
lepidopteris Langsd. et Fisch. Ic. Fil. 5 t. 2. 1810 = Polypodium.
lepidotum Willd. sp. **5**. 102 1810 = Elaphoglossum plicatum.
lepidotum? Raddi, Opusc. sci. Bol. **3**. 283. 1819 = Elaphoglossum strictum.
leprosum Mett.; Kuhn, Linn. **36**. 48. 1869 = Elaphoglossum.
leptophlebium Bak. Tr. Linn. Soc. II. Bot. **2**. 295. 1886 = Elaphoglossum.
leptophyllum Lam. et DC. Fl. franç. **2**. 565. 1805 = Anogramma.
leptophyllum Fée, Acrost. 45 t. 17 f. 1. 1845 = Elaphoglossum.
Lessoni Mett.; Kuhn, Linn. **36** 60. 1869 = Elaphoglossum.
Liaisianum Glaziou; Fée, Cr. vasc. Br. **2**. 13 t. 86 f. 2. 1872-73 = Elaphoglossum strictum.
Lindbergii Mett.; Kuhn, Linn. **36**. 46. 1869 = Elaphoglossum hybridum.
Lindenii Bory; Fée, Acrost. 48 t. 18 f. 3. 1845 = Elaphoglossum.
Lindigii Karst. Fl. Col. **1**. 5 t. 3. 1858 = Elaphoglossum.
Lindigii Bak. Syn. 423. 1868 = Leptochilus.
lineare Spr. Schrad. Journ. **1799**[2]. 267 = Lomaria woodwardioides.
lineare Hassk. Cat. Pl. Bogor. 3. 1844 = ? Photinopteris speciosa.
lineare Fée, Acrost. 47 t. 15 f. 2. 1845 = Elaphoglossum.
linearifolium Pr. Del. Prag. **1**. 159. 1822 = Polybotrya cervina var.
lineatum Cav. Anal. Hist. Nat. **1**. 106. 1799 = Blechnum spicant.
lineatum Kuhn; Christ, Mon. Elaph. 146. 1899 = Elaphoglossum simplex var.
Lingua Thbg. Fl. Jap. 330 t. 33. 1784 = Cyclophorus.
Lingua Raddi, Opusc. sci. Bol. **3**. 283. 1819 = Elaphoglossum.
linguaeforme Cav. Anal. Hist. Nat. **1**. 103. 1799 = Elaphoglossum.
Linnaeanum Hk. 2 Cent. t. 26. 1860 = Leptochilus.
Listeri Bak. Journ. Linn. Soc. **25**. 361. 1889 = Leptochilus.

Lloense Hk. Ic. pl. t. 657. 1844 = Elaphoglossum.
lomarioides Bory, Bél. Voy. Bot. **2**. 21 t. 2. 1833 = Stenochlaena pollicina var.
lomarioides Christ, Verh. Nat. Ges. Basel **11**. 252. 1895 = Leptochilus.
lomarioides Jenm. Bull. Dep. Jam. **5**. 154. 1898 = A. aureum var.
lonchophorum Kze. Farnkr. **1**. 5 t. 2. 1840 = Leptochilus cuspidatus.
lonchophyllum Fée, 8 mém. 68. 1857 = Elaphoglossum flaccidum var.
longifolium Burm. Fl. Ind. 228. 1768 = Cyclophorus.
longifolium Jacq. Coll. **2**. 105. 1788 = Elaphoglossum.
Lorentzii Hier. Engl. Jahrb. **22**. 408. 1896 = Elaphoglossum.
Lowei Fée, Acrost. 4, 9. 1845 = Elaphoglossum hirtum.
ludens Wall. List n. 2685. 1829 = Polybotrya appendiculata.
luridum Fée, Acrost. 35 t. 19 f. 1. 1845 = Elaphoglossum Schomburgkii.
luteum Desv. Prod. 211. 1827 = Notholaena.
Macahense Fée, Cr. vasc. Br. **2**. 2 t. 79 f. 1. 1872-73 — Elaphoglossum Aubertii var.
macrophyllum Mett.; Kuhn, Linn. **36**. 61. 1869 = Elaphoglossum.
macropodium Fée, Acrost. 30 t. 6 f. 2. 1845 = Elaphoglossum Sieberi.
macrorhizum Bak. Ann. Bot. **5**. 490. 1891 = Elaphoglossum.
madagascariense Kuhn; Bak. Ann. Bot. **5**. 490. 1891 = Elaphoglossum.
magnum Bak. Gard. Chr. n. s. **20**. 135. 1883 = Elaphoglossum.
Mandoni Mett.; Kuhn, Linn. **36**. 47. 1869 = Elaphoglossum.
Mannianum Mett.; Kuhn, Fil. Afr. 46. 1868 = Elaphoglossum cinnamomeum.
Marantae L. sp. **2**. 1071. 1753 = Notholaena.
Marantae Pallas, Reis. d. russ. Reich. **3**. 293. 1776 — Woodsia ilvensis.
marginale Bak. Syn. 519. 1874 = Elaphoglossum.
marginatum L. sp. ed. II. **2**. 1526. 1763 = Pteris grandifolia.
marginatum Schkuhr, Kr. Gew. **1**. 185 t. 3 b. 1809 = A. aureum.
marginatum Wall. List n. 17. 1828; Fée, Acrost. 31. 1845 = Elaphoglossum conforme.
martinicense Desv.; Berl. Mag. **5**. 309. 1811 = Elaphoglossum simplex var.
martinicense Fée, Acrost. 45 t. 16 f. 3. 1845 = Elaphoglossum glabellum.
mascarenense Spr. Syst. **4**. 37. 1827 = Leptochilus auriculatus.
Mathewsii Fée, Acrost. 54 t. 2 f. 2. 1845 = Elaphoglossum.
maximum Bak. Ann. Bot. **5**. 490. 1891 = Elaphoglossum.
melanolepis Fée, Gen. 43. 1850-52 = Elaphoglossum.
melanopus Kze. Linn. **23**. 214, 298. 1850 = Elaphoglossum hybridum var.
melanostictum Bl. Fl. Javae Fil. 26 t. 7. 1828 = Elaphoglossum.
meridense Kl. Linn. **20**. 427. 1847 = Elaphoglossum.
Mettenii Kuhn, Linn. **36**. 48. 1869 = Elaphoglossum.
mexicanum Fourn. Mex. pl. **1**. 63. 1872 = Elaphoglossum hybridum.
Meyerianum Hk. Gard. Ferns t. 16. 1862 = Stenochlaena tenuifolia.
micans Mett.; Kuhn, Linn. **36**. 50. 1869 = Elaphoglossum hirtum var.
micradenium Fée, Acrost. 43 t. 8 f. 1. 1845 = Elaphoglossum.
microlepis Kze. Linn. **23**. 214, 299. 1850 = Elaphoglossum.
microlepis Sod. Cr. vasc. quit. 437. 1893 = Elaphoglossum subnudum.
microphyllum Bertol. Propag. agric. **6**. 343. 1856; Misc. bot. **18**. 19 t. 14. 1857. (Rend. Sess. Acc. Sci. Bol. **1856-57**. 53. 1857) = Notholaena persica.
microphyllum Mett.; Kuhn, Fil. Afr. 46. 1868 = ? Elaphoglossum hybridum.
Miersii Bak. Ann. Bot. **5**. 490. 1891 = Elaphoglossum.

miniatum Christ, Verh. Nat. Ges. Basel **11**. 254 t. 3 f. 25—27. 1895 = Elapho-
minus Mett. Fil. Lips. 20. 1856 = Leptochilus. [glossum.
minutum Pohl; Fée, Acrost. 39 t. 10 f. 3. 1845 = Elaphoglossum.
modestum Bak. Journ. Linn. Soc. **22**. 231. 1886 = Leptochilus.
mollissimum Fée, Cr. vasc. Br. **1**. 7 t. 2 f. 3. 1869 = Elaphoglossum villosum var.
Moorei E. Britton, Mem. Torr. Cl. **4**. 271. 1895 = Elaphoglossum.
Moritzianum Kl. Linn. **20**. 423. 1847 = Elaphoglossum villosum var.
Mülleri Fourn. Mex. pl. **1**. 68 t. 1 f. 2. 1872 = Elaphoglossum.
muscosum Sw. Prod. 128. 1788 = Elaphoglossum.
nanum Liebm. Vid. Selsk. Skr. V. **1**. 171 (seors. 19). 1849 = Polypodium (Craspedaria) sp. stat. juv.!
nanum Hk. sp. **5**. 249. 1864 = Blechnum lanceolatum.
neglectum Bailey, Tr. Linn. Soc. N. S. Wales **5**. 32. 1881; F. Muell. Fragm. **11**. 131. 1881 = Leptochilus.
nemorale Lam. Fl. franç. **1**. 11. 1778 = Blechnum spicant.
neriifolium Wall. List n. 16. 1828 = Elaphoglossum petiolatum var.
nervosum Bory, Dup. Voy. Bot. **1**. 252. 1828 = Elaphoglossum.
nicotianaefolium Sw. Syn. 13, 199. 1806 = Leptochilus.
nigrescens Hk. sp. **5**. 214. 1864 = Elaphoglossum.
nitidum Liebm. Vid. Selsk. Skr. V. **1**. 168 (seors. 16). 1849 = Elaphoglossum
niveum Desv. Prod. 212. 1827 = Pellaea. [hybridum var.
nivosum Kze. Bot. Zeit. **1845**. 281 = Elaphoglossum tectum.
nodosum L. sp. **2**. 1070. 1753 = Danaea.
Norrisii Hk. sp. **5**. 215. 1864 = Elaphoglossum melanostictum.
Norrisii Cesati, Atti Acad. Napoli **7**^{8}. 31. 1876 = Elaphoglossum Beccarianum.
notatum Fée, Acrost. 38 t. 10 f. 1. 1845 = Elaphoglossum.
nudum Kze.; Kl. Linn. **20**. 422. 1847 = Elaphoglossum Gayanum.
nummularifolium Sw. Syn. 191, 419 t. **2** f. 1. 1806 = Cyclophorus.
nutans Hk. sp. **5**. 244. 1864 = Polybotrya.
obcrenatum Fée, 11 mém. 4 t. 1 f. 2. 1866 — Guadeloupe — (Elaphoglossum?
obductum Klf.; Spr. Syst. **4**. 34. 1827 = Elaphoglossum. [sp. dubia).
obliquatum Fée, Cr. vasc. Br. **1**. 261. 1869 = Elaphoglossum Sellowianum.
obliquum Bl. Enum. 101. 1828 = A. aureum.
oblongum Desv. Berl. Mag. **5**. 308. 1811 = Elaphoglossum conforme.
obovatum Bl. Enum. 102. 1828 = Cyclophorus nummularifolius var.
obscurum Fourn. Mex. pl. **1**. 68. 1872 = Elaphoglossum.
obtusatum Carm. Tr. Linn. Soc. **12**. 510. 1818 = Elaphoglossum spathulatum var.
obtusifolium Willd. sp. **5**. 107. 1810 — India orient. — (Leptochilus? sp. dub.)
obtusifolium Spr. Syst. **4**. 34. 1827; Bl. Enum. 102. 1828 = Elaphoglossum
oligarchicum Bak. Syn. 418. 1868 = Leptochilus. [decurrens.
oligodictyon Bak. Journ. Linn. Soc. **24**. 261. 1887 = Leptochilus.
omphalodes Fée, Cr. vasc. Br. **2**. 7 t. 81 f. 2. 1872-73 = Elaphoglossum villosum vel E. Lindenii.
ophioglossoides Goldm. Nova Acta **19** Suppl. I. 451. 1843 = Elaphoglossum decurrens.
Orbignyanum Fée, Acrost. 56 t. 13 f. 2. 1845 = Elaphoglossum laminarioides.
ornatum Mett.; Kuhn, Linn. **36**. 44. 1869 = Elaphoglossum.
osmundaceum Hk. sp. **5**. 246. 1864 = Polybotrya.
ovalifolium Boj. Hort. Maur. 414. 1837 (nomen) = Elaphoglossum conforme.
ovalifolium Fée, Cr. vasc. Br. **1**. 3 t. 4 f. 1. 1869 = Elaphoglossum.

ovatum Hk. et Grev. Ic. Fil. t. 146. 1829 = Elaphoglossum.
ovatum Liebm. Vid. Selsk. Skr. V. **1**. 164 (seors. 12). 1849 = Elaphoglossum squamipes.
ovatum Fée, Cr. vasc. Br. **2**. 3 t. 80 f. 2. 1872-73 = Elaphoglossum conforme.
pachydermum Fée, Acrost. 47. 1845 = Elaphoglossum.
pachyphyllum Kze. Linn. **9**. 26. 1834 = Elaphoglossum.
paleaceum Hk. et Grev. Ic. Fil. sub t. 235. 1831 = Elaphoglossum hirtum.
pallidum Bak. JoB. **1879**. 263 = Elaphoglossum.
pallidum Beyrich; Christ, Mon. Elaph. 60. 1899 = Elaphoglossum flaccidum.
palustre Hk. sp. **5**. 214. 1864 = Elaphoglossum.
palustre Clarke, Tr. Linn. Soc. II. Bot. **1**. 577. 1880 = Stenochlaena.
pandurifolium Hk. sp. **5**. 271. 1864 = Leptochilus.
papillosum Bak. JoB **1877**. 167 = Elaphoglossum.
papyraceum Fée, Cr. vasc. Br. **1**. 4 t. 3 f. 1. 1869 = Elaphoglossum flaccidum.
Patini Bak. Syn. 519. 1874 = Elaphoglossum.
pectinatum L. sp. **2**. 1068. 1753 = Schizaea.
pellucido-marginatum Christ, Verh. Nat. Ges. Basel **11**. 255. 1895 = Elaphoglossum gorgoneum.
peltatum Sw. Schrad. Journ. **1800²**. 11. 1801 = Elaphoglossum.
pennula Poir. Enc. Suppl. **1**. 125. 1810 = Schizaea.
perelegans Fée, Acrost. 55 t. 23. 1845 = Elaphoglossum.
pervium Lindm. Ark. för Bot. **1**. 254 t. 8 f. 7. 1903 = Leptochilus serratifolius.
petiolatum Sw. Prod. 128. 1788 = Elaphoglossum.
petiolosum Desv. Berl. Mag. **5**. 309. 1811 = Elaphoglossum.
phanerodictyon Bak. Boll. Soc. Brot. **4**. 156 t. 2 f. C. 1886 = Leptochilus.
phlebodes Kze. Lin. **9**. 33. 1834 = Stenochlaena marginata.
Photinopteris »Bak.« Salom. Nom. 10. 1883 — Ins. Admiralitatis.
pilosella Spr. Syst. **4**. 34. 1827 } = Elaphoglossum spathulatum var.
piloselloides Pr. Rel. Haenk. **1**. 14 t. 2 f. 1. 1825 } = Elaphoglossum spathulatum var.
pilosiusculum Wikstr. Vet. Ak. Hdl. **1825**. 439. 1826 = ? Dryopteris africana.
pilosum H.B. Willd. **5**. 103. 1810 = Elaphoglossum.
pilosum L. v. Buch, Beschr. Canar. Ins. 189. 1825 (nomen) = Dryopteris africana.
Pittieri Christ, Bull. Soc. bot. Belg. **35**. 243. 1896 = Stenochlaena.
platyneuron L. sp. **2**. 1069. 1753 = Asplenium.
platynevron Fée, Acrost. 43 t. 4 f. 1. 1845 = Elaphoglossum apodum.
platyrhynchos Hk. sp. **5**. 280. 1864 = Hymenolepis.
plicatum Cav. Anal. Hist. Nat. **1**. 103. 1799 = Elaphoglossum.
plumbicaule Bak. Syn. 413. 1868 = Polybotrya.
Plumieri Desv. Prod. 209. 1827 = Elaphoglossum petiolatum.
Plumieri Fée, Acrost. 50. 1845 = Elaphoglossum villosum var.
plumosum Fée, Acrost. 54 t. 20 f. 1. 1845 = Elaphoglossum.
podotrichum Desv. Berl. Mag. **5** 309. 1811 = Elaphoglossum undulatum.
Poeppigianum Fée, Gen. 43. 1850-52 = Elaphoglossum villosum.
polybotryoides Bak. JoB. **1881**. 207 = Polybotrya p.
polyphyllum Hk. sp. **5**. 243. 1864 = Stenochlaena decrescens.
polyphyllum Hk. sp. **5**. 269. 1864 = Polybotrya.
polypodioides L. sp. **2**. 1068. 1753 = Polypodium p.
polypodioides Thouars, Fl. Trist. d'Ac. t. 2. 1804 = Blechnum penna marina.
polystichum Mett.; Kuhn, Linn. **36**. 57. 1869 = Elaphoglossum Hoffmanni.

polytrichoides Thours, Fl. Trist. d'Ac. 32 t. 2 (polypodioides). 1804 = Blechnum penna marina.
Poolii Bak. Ann. Bot. **5**. 493. 1891 = Elaphoglossum.
Portoricense Spr. Nova Acta **10**. 226. 1821 = Leptochilus alienus.
potentillaefolium Bomm.; Christ, Mon. Elaph. 59. 1899; (Bull. Soc. bot. Belg. **35**. 242. 1896) = Elaphoglossum peltatum.
praecelsum André ap. Christ, Mon. Elaph. 121. 1899 = Elaphoglossum.
praelongum Fée, Cr. vasc. Br. **1**. 9 t. 3 f. 2. 1869 = Elaphoglossum Preslianum.
?**praestantissimum** Bory ap. Fée, Acrost. 89 1845 (syn.). HB. 423. NPfl. 336 — India occ. — Neurocallis Fée, Acrost. 89 t. 52. 1845; Chrysodium Mett. 1856; Lomagramme Gris. 1864.
Preslianum Fée, Acrost. 46. (t. 24 f. 1?). 1845 = Elaphoglossum.
Preslianum Hk. sp. **5**. 265. 1864 = Leptochilus.
Prestoni Bak. Gard. Chr. **1872**. 1555 = Elaphoglossum scolopendrifolium var.
Prieuriana Kl. Linn. **20**. 429. 1847 = Stenochlaena marginata.
Pringlei Dav. Bot. Gaz. **21**. 253. 1894 = Elaphoglossum.
procurrens Mett.; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860 = Elaphoglossum.
producens Fée, Cr. vasc. Br. **2**. 4 t. 80 f. 1. 1872 = Elaphoglossum lingua.
proliferum Bl. Enum 104. 1828 = Leptochilus cuspidatus.
proliferum Hk. Ic. pl. t. 681-682. 1844 = Leptochilus subcrenatus.
propinquum Mett.; Kuhn, Linn. **36**. 45. 1869 = Elaphoglossum.
proximum Bomm. Bull. Soc. bot. Belg. **35**. 241. 1896 = Elaphoglossum lloense.
pterioides Bernh. Schrift. Akad. Erfurt **1802**. 18. f. 11. = Notholaena trichomanoides.
pterioides R. Br. Prod. Fl. N. Holl. 145. 1810 = Neurosoria.
pubens Bak. Syn. 414. 1868 = Polybotrya.
pulchellum Bak. Fl bras. 1^2. 600. 1870 = Elaphoglossum Lindenii.
pulchrum L. sp. **2**. 1072. 1753 = Asplenium adiantum nigrum.
pumilum Mart. et Gal. Mém. Ac. Brux. **15**. 23 t. 2 f. 2. 1842 = Elaphoglossum spathulatum.
punctatum L sp. ed. II. **2**. 1524. 1763 = Polypodium.
punctatum L.fil. Suppl. 444. 1781 } = Leptochilus auriculatus.
punctulatum Sw. Schrad. Journ. $\mathbf{1800^2}$. 12. 1801 } = Leptochilus auriculatus.
punctulatum Pr. Rel. Haenk. **1**. 16. 1825 = Leptochilus Preslianus.
pusillum Mett.; Kuhn, Linn. **36**. 43. 1869 = Elaphoglossum.
pygmaeum Mett.; Kuhn, Linn. **36**. 42. 1869 = Elaphoglossum.
quercifolium Retz. Obs. **6**. 39. 1791 = Leptochilus zeylanicus.
quitense Bak. Ann. Bot. **5**. 493. 1891 = Elaphoglossum.
Quoyanum Gaud. Freyc. Voy. Bot. 306 t. 3. 1827 = Leptochilus cuspidatus var.
rabdolepis Fée, Gen. 42 t. 1 f. 1. 1850-52 = Elaphoglossum ovatum.
Raddianum Hk. et Gr. Ic. Fil. t. 4. 1827 = Elaphoglossum spathulatum.
Raddianum Kze.; Hk. sp. **5**. 264. 1864 = Leptochilus guianensis.
Raddii Desv. Prod. 209. 1827 = Elaphoglossum spathulatum.
radiatum König; Poir. Enc. Suppl. **1**. 128. 1810 = Actiniopteris australis.
ramentaceum Roxb. Calc. Journ. **4**. 479. 1844 = Hemionitis arifolia.
ramosissimum Fée, Acrost. 53 t. 22 f. 3. 1845 = Elaphoglossum ovatum.
rampans Bak. Syn. 518. 1874 = Elaphoglossum.
Rawsoni Bak. Ann. Bot. **5**. 496. 1891 = Leptochilus.
recognitum Kze.: Fourn. Mex. pl. **1**. 67. 1872 = Elaphoglossum villosum.
Reichenbachii Moritz; Kuhn, Linn. **36**. 42. 1869 = Elaphoglossum.
repandum Bl. Enum. 104. 1828 = Leptochilus cuspidatus.

reptans Cav. Anal. Hist. Nat. **1**. 104. 1799 = Elaphoglossum.
Requinianum Gaud. Freyc. Voy. Bot. 304 t. **4**. 1827 = Platytaenia.
reticulatum Klf. Enum. 64. 1824 = Elaphoglossum.
revolutum Liebm. Vid. Selsk. Skr. V. **1**. 163 (seors. 11). 1849 = Elaphoglossum.
revolvens Kze.; Mett. Ann. sc. nat. V. **2**. 203. 1864 (nomen); Kuhn, Linn. **36**. 57. 1869 = Elaphoglossum.
Richardi Hk. sp. **5**. 256. 1864 = Elaphoglossum.
rigens Pr. Epim. 180. 1849 = A. aureum.
rigidum Wall. List n. 27. 1828 = Photinopteris speciosa.
rigidum Fée, Cr. vasc. Br. **2**. 7 t. 83 f. 1. 1872-73 = Elaphoglossum simplex.
Rimbachii Sod. Cr. vasc. quit. 478. 1893 = Elaphoglossum.
rivulare Wall. List n. 2165. 1829 ≐ Leptochilus decurrens.
rivulare Hk. sp. **5**. 263. 1864 = Leptochilus.
Roezlii Schaffn.; Fée, 8 mém. 69. 1857 = Elaphoglossum.
rubescens Kuhn ap. Christ, Mon. Elaph. 93. 1899 (syn.) = Elaphoglossum.
rubicundum Pohl; Ettingsh. Farnkr. 4 t. 3 f. 2. 1865 = Elaphoglossum latifolium.
rubiginosum Fée, Acrost. 47 t. 5 f. 1, t. 13 f. 1. 1845 = Elaphoglossum tectum.
rufescens Liebm. Vid. Selsk. Skr. V. **1**. 171 (seors. 18). 1849 = Elaphoglossum.
rufidulum Willd.; Kuhn, Fil. Afr. 47. 1868 = Elaphoglossum.
rufum L. Syst. Nat. ed. X. **2**. 1320. 1759 = Gymnopteris.
rufum Spr. Syst. **4**. 31. 1827 = Blechnum discolor.
rupestre Karst. Fl. Col. **1**. 118 t. 59. 1860 = Elaphoglossum.
Ruta muraria Lam. Illustr. t. 865 f. 1. —. = Asplenium.
salicifolium Willd.; Klf. Enum. 58. 1824 = Elaphoglossum petiolatum.
salicinum Hk. sp. **5**. 265. 1864 = Leptochilus.
Salvini Bak.; Hemsl. Biol. Centr. Amer. **3**. 688. 1885 = Elaphoglossum.
samoense Bak. Syn. 407. 1868 = Elaphoglossum.
sanctum L. Syst. Nat. ed. X. **2**. 1320. 1759 = Dryopteris.
Sartorii Liebm. Vid. Selsk. Skr. V. **1**. 166 (seors. 14). 1849 = Elaphoglossum [dissimile.
savaiense Bak. Ann. Bot. **5**. 494. 1891 = Elaphoglossum.
scalpellum Mart. Ic. Cr. Bras. 86. 1834 = Elaphoglossum latifolium.
scalpturatum Kze. Bot. Zeit. **1848**. 103 = Leptochilus.
scalpturatum Pr. Epim. 181. 1849 = A. aureum.
scalpturatum Lindm. Ark. för Bot. **1**. 253 t. 8 f. 9. 1903 = Leptochilus curupirae.
scandens Raddi, Opusc. sci. Bol. **3**. 284. 1819; Pl. bras. **1**. 6 t. 18. 1825 = Leptochilus guianensis.
scandens Bory; Fée, Acrost. 33. 1845 = Elaphoglossum lingua var.
scandens Hk. sp. **5**. 249. 1864 = Stenochlaena palustris.
scapellum Fée, Acrost. 32 t. 10 f. 2. 1845 (err.) = A. scalpellum Mart.
scapellum Kze.; Fée, 9 mém. 1. 1857 = Elaphoglossum.
scariosum Sw. Syn. 16. 1806 = Notholaena.
Schaffneri Fée; Fourn. Mex. pl. **1**. 64. 1872 = ? Elaphoglossum flaccidum.
Schiedei Kze. Anal. 10. 1837 = Elaphoglossum dissimile.
schizolepis Bak. Journ. Linn. Soc. **15**. 421. 1876 = Elaphoglossum.
schizolepis Bak. JoB. **1881**. 368 = Elaphoglossum Curtisii.
Schlimense Fée, 8 mém. 68. 1857 = Elaphoglossum.
Schmitzii Mett.; Kuhn, Linn. **36**. 51. 1869 = Elaphoglossum dissimile.
Schomburgkii Fée, Acrost. 32 t. 8 f. 2. 1845 = Elaphoglossum.
scolopendrifolium Raddi, Pl. Bras. **1**. 4 t. 16. 1825 = Elaphoglossum.

seetacoonense Roxb. Calc. Journ. **4**. 480. 1844 — Chittagong. (Plagiogyria vel Stenochlaena sp.).
sejunctum Kuhn, Fil. Afr. 47. 1868 (nomen) — Sierra Leone. — (Elaphoglossum sp.).
Sellowianum Kl.; Kuhn, Linn. **36**. 52. 1869 = Elaphoglossum.
semicordatum Bak. Syn. 422. 1868 = Leptochilus virens.
semipinnatifidum Hk. sp. **5**. 273. 1864 = Leptochilus alienus var.
semipinnatum Roxb. Calc. Journ. **4**. 480. 1844 = ? Leptochilus latifolius.
septentrionale L. sp. **2**. 1068. 1753 = Asplenium
serratifolium Mertens; Klf. Enum. 66. 1824 = Leptochilus.
serratifolium Lindm. Ark. för Bot. **1**. 252 t. 8 f. 8. 1903 = Leptochilus contaminoides.
serratum Lam. Illustr. t. 865 f. 2; Poir. Enc. Suppl. **1**. 121. 1810 = Polypodium minimum (f. Moore); = Schizaea serrata (f. Fée).
serratum Bak. Syn. 524. 1874 = Leptochilus.
serrulatum Sw. Prod. 128. 1788 = Polypodium.
serrulatum Willd. sp. **5**. 113. 1810 (Plum. t. 81) — Hispaniola. Polybotrya ? Plumieri Moore 1857.
sessile Fée, Gen. 43. 1850-52 (nomen) = Elaphoglossum.
setigerum Sod. Cr. vasc. quit. 453. 1893 = Elaphoglossum Aubertii.
setosum Wall. List n. 30. 1828 = Polybotrya appendiculata.
setosum Liebm. Vid. Selsk. Skr. V. **1**. 169 (seors. 17). 1849 = Elaphoglossum [villosum.
Sherringii Bak. Ann. Bot. **6**. 100. 1892 = Elaphoglossum.
Sieberi Hk. et Grev. Ic. Fil. t. 237. 1831 = Elaphoglossum.
siliquoides Jenm. JoB. **1881**. 53 = Elaphoglossum.
siliquosum L. sp. **2**. 1070. 1753 = Ceratopteris thalictroides.
simplex Sw. Prod. 128. 1788 = Elaphoglossum.
simplex Blanco, Fl. Filip. 826. 1837 — Ins. Philipp.
sinuatum Lagasca; Sw. Syn. 14. 1806 = Notholaena.
Smithii Bak. Ann. Bot. **5**. 169. 1891 = Elaphoglossum.
Smithii Racib. Bull. int. Ac. Cracovie **1902**. 59 = Stenochlaena Raciborskii.
Sodiroi Bak. JoB. **1877**. 167 = Elaphoglossum.
sorbifolium L. sp. **2**. 1069. 1753 = Stenochlaena.
spathulatum Bory, Voy. **1**. 363 t. 20 f. 1. 1804 = Elaphoglossum.
spathulinum Raddi, Pl. Bras. **1**. 3 t. 15 f. 2. 1825 = Elaphoglossum spathu- [latum var.
speciosum Willd. sp. **5**. 117. 1810 = A. aureum.
spectabile Zoll. Syst. Verz. 33. 1854 = A. aureum.
spectabile Racib. Pterid. Buit. 54. 1898 = Stenochlaena sorbifolia var. 8.
sphenophyllum Kze. Anal. 11 t. 7. 1837 = Elaphoglossum flabellatum.
Spicant Willd. Fl. Berol. Prod. 289. 1787 = Blechnum.
spicatum L.fil. Suppl. 444. 1781 = Hymenolepis.
spinulosum Noronha, ex Thouars, Mél. bot. et voy. —. 1811 — Madagascar.
spissum Fée, Cr. vasc. Br. **1**. 8 t. 2 f. 2. 1869 = Elaphoglossum conforme.
splendens Bory; Willd. sp. **5**. 104. 1810; Poir. Enc. Suppl. **1**. 120. 1810 = Elaphoglossum hirtum var.
sporadolepis Kze.; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860 (nomen); Kuhn, Linn. **36**. 59. 1869 = Elaphoglossum.
Spruceanum Fée: Christ. Mon. Elaph. 32. 1899 = Elaphoglossum latifolium.
Sprucei Bak. Syn. 411. 1868 = Elaphoglossum.
squamatum Sw. Syn. 11. 1806 = A. squamosum.
squamipes Hk. Ic. Pl. t. 197. 1837 = Elaphoglossum.

squamosum Cav. Anal. Hist. Nat. **1**. 104. 1799; Descr. 239. 1802 — Ins. Mariannae. Peru. — A. squamatum Sw. 1806; Olfersia Pr. 1836; Elaphoglossum Moore 1857.
squamosum Sw. Schrad. Journ. **1800**². 11. 1801 = Elaphoglossum hirtum.
squarrosum Kl. Linn. **20**. 424. 1847 = Elaphoglossum.
staphyleum **Link, Hort. Berol. 2. 47. 1833 — Hort.**
stelligerum Wall. List n. 2167. 1829; Bak. Syn. 521. 1874 = Elaphoglossum petiolatum.
Stemaria Pal. Beauv. Fl. d'Oware **1**. 2 t. 2. 1804 = Platycerium.
stenophyllum Sod. Cr. vasc. quit. 468. 1893 = Elaphoglossum.
Stenopteris Kl. Linn. **20**. 420. 1847 = Elaphoglossum brachyneuron
stenosemioides Bak. Journ. Linn. Soc. **22**. 230. 1886 = Polybotrya.
stigmatolepis Fée, Acrost. 62 t. 24 f. 2. 1845 = Elaphoglossum.
stipitatum Bory; Fée, Acrost. 38 t. 4 f. 3. 1845 = Elaphoglossum.
stramineum Mett. Ann. sc. nat. V. **2**. 200. 1864 = Elaphoglossum.
stramineum Fée, Cr. vasc. Br. **1**. 238. 1869 = Elaphoglossum Aubertii.
strictum Raddi, Pl. Bras. **1**. 3 t. 15 f 3. 1825 = Elaphoglossum.
subcordatum Cav. Anal. Cienc. **4**. 97. 1801 = Notholaena Marantae.
subcrenatum Hk. et Grev. Ic. Fil. t. 110. 1829 = Leptochilus.
subdiaphanum Hk. et Grev. Ic. Fil. t. 205. 1831 = Elaphoglossum nervosum.
suberectum Bak. JoB. **1881**. 207 = Polybotrya.
subrepandum Hk. sp. **5**. 275. 1864 = Leptochilus Zollingeri.
subsessile Bak. JoB. **1884**. 143 = Elaphoglossum.
subsessile Bak. Kew Bull. **1897**. 300 = Elaphoglossum malgassicum.
succisaefolium Thouars, Fl. Trist. d'Ac. 31. 1804 (?) (an succisum?) = Elaphoglossum.
sulphureum Sw. Prod. 129. 1788 = Gymnogramma.
taccaefolium Hk. sp. **5**. 278. 1864 = Leptochilus latifolius var.
tahitense Carr. in Seem. Fl. Vit. 373. 1873 = Elaphoglossum.
tambillense Hk. Ic. pl. t. 636. 1844 = Elaphoglossum.
Tanajense Kuhn, Linn. **36**. 50. 1869 = Elaphoglossum auricomum.
tartareum Cav. Descr. 242. 1802 = Ceropteris.
Tatei Bak. Syn. 518. 1874 = Elaphoglossum amygdalifolium.
Taylori Bail. Trans. Accl. Soc. Queensl. 1884; Syn. Queensl. Fl. Suppl. I. 65. 1886 = Leptochilus.
tectum H.B. Willd. sp. **5**. 102. 1810 = Elaphoglossum.
tenellum Desv. Prod. 210. 1827 = Elaphoglossum spathulatum. (an E. lineare?).
tenerum Fée; Kuhn. Linn. **36**. 52. 1869 = Elaphoglossum.
tenerum Bak. JoB. **1878**. 302 = Elaphoglossum Balansae.
tenue Retz. Obs. **6**. 39. 1791 = Cheilanthes tenuifolia.
tenuiculum Fée, 10 mém. 6 t. 29 f. 2. 1865 = Elaphoglossum.
tenuifolium Liebm. Vid. Selsk. Skr. V. **1**. 163 (seors. 11). 1849 = Elaphoglossum.
tenuifolium Bak. Syn. 412. 1868 = Stenochlaena.
tereticaulon Desv. Berl. Mag. **5**. 310. 1811 = ? Pellaea nivea var.
terminans Wall. List n. 2168. 1829 = Leptochilus virens.
Teysmannianum Bak. Malesia **3**. 56. 1886 = Stenosemia.
thalictroides L. sp. 2. 1070. 1753 = Ceratopteris.
Thelypteris L. sp. 2. 1071. 1753 = Dryopteris.
Thomsoni Bak. Journ. Linn. Soc. **15**. 111. 1876 = Dryostachyum.
Tolimense Kuhn ap. Christ. Mon. Elaph. 124. 1899 (syn.) = Elaphoglossum.
tomentosum Bory; Willd. sp. **5**. 101. 1810 = Elaphoglossum.
Tonduzii Christ, Mon. Elaph. 45. 1899 = Elaphoglossum.

Tosaense Yatabe, Bot. Mag. Tokyo **5**. 149 t. 24. 1891 = Elaphoglossum.
Tovarense Moritz; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860; Kuhn, Linn. **36**. 60. 1869 = Elaphoglossum latifolium.
tricholepis Bak. JoB. **1891**. 5 = Elaphoglossum.
tricuspe Hk. sp. **5**. 272 t. 304. 1864 = Leptochilus.
trifoliatum L. sp. **2**. 1070. 1753 = Trismeria.
trifoliatum Houtt. Pfl. Syst. **13**[1]. 90 t. 95 f. 3. 1786 = Pteris ensiformis.
trifrons Commers.; Mirb. Dict. sci. nat. **1**. 244. 1816 = Stenochlaena pollicina.
trinerve Hassk. Cat. Pl. Bogor. 3. 1844 = Cheiropleuria bicuspis.
tripartitum Hk. et Grev. Ic. Fil. t. 118. 1829 = Elaphoglossum flabellatum.
tripinatum Blanco, Fl. Filip. 826. 1837 — Ins. Philipp.
triquetrum Wall. List n. 23. 1828 = Plagiogyria euphlebia.
triste Vell. Fl. flum. **11**. t. 90. 1827; Arch. Mus. Nac. Rio Jan. **5**. 453. 1881 = ? Leptochilus serratifolius.
trivittatum Sod. Rec. 82. 1883 = Elaphoglossum.
truncicola Karst. Fl. Col. **1**. 121 t. 60. 1860 = Elaphoglossum.
umbrosum Liebm. Vid. Selsk. Skr. V. **1**. 174 (seors. 22). 1849 = Leptochilus.
undulatum Willd. sp. **5**. 105. 1810 = Elaphoglossum.
unitum Bory; Fée, Acrost. 44. 1845 = Elaphoglossum affine.
Urvillei Pr. Epim. 181. 1849 = A. aureum.
vagans Mett.; Kuhn, Linn. **36**. 58. 1869 = Elaphoglossum.
variabile Hk. sp. **5**. 277. 1864 = Leptochilus decurrens.
varians Mett. Ann. sc. nat. IV. **15**. 56. 1861 = Leptochilus.
velleum Ait. Hort. Kew. **3**. 457. 1789 = Notholaena.
venustum Liebm. Vid. Selsk. Skr. V. **1**. 168 (seors. 16). 1849 = Elaphoglossum Lindenii.
venustum Fée, 8 mém. 68. 1857 = Elaphoglossum araneosum.
versatile Sod. Rec. 76. 1883 = Elaphoglossum scolopendrifolium.
Vespertilio Mett. Fil. Lips. 20. 1856 = Cheiropleuria bicuspis.
vestitum Ham.; Wall. List n. 2164. 1829 (nomen) = Elaphoglossum tomentosum.
vestitum Schl. et Cham. Linn. **5**. 605. 1830 = Elaphoglossum.
vestitum Lowe; Hk. et Grev. Ic. Fil. t. 235 (paleaceum). 1831 = Elaphoglossum hirtum.
Vieillardii Mett. Ann. sc. nat. IV. **15**. 55. 1861 = Elaphoglossum.
villosum Sw. Prod. 128. 1788 = Elaphoglossum.
virens Wall. (List n. 1033. 1828. dub.); Hk. et Grev. Ic. Fil. t. 221. 1831 = Leptochilus.
virens Bak. Syn. 420. 1868 = Leptochilus virens, subcrenatus, scalpturatus, acrostichoides.
viride Fourn. Mex. pl. **1**. 66. 1872 = Elaphoglossum.
viridifolium Jenm. JoB. **1886**. 273 = Elaphoglossum.
viscidum Fée, Cr. vasc. Br. **2**. 12 t. 86 f. 1. 1872-73 = Elaphoglossum.
viscosum Sw. Syn. 10, 193. 1806 = Elaphoglossum petiolatum.
viviparum L.fil. Suppl. 444. 1781 = Asplenium.
viviparum Cav. Descr. 240. 1802 = ? Onychium siliculosum.
viviparum Ham.; Spr. Syst. **4**. 36. 1827 = Polybotrya appendiculata var.
Volkensii Hier. in Engl. Pflw. Ostafr. C. 81. 1895 = Elaphoglossum.
Wageneri Kze. Linn. **25**. 746. 1853 = Elaphoglossum plumosum.
Wallii Bak. JoB. **1872**. 146 = Leptochilus.
Wawrae Luerss. Flora **1875**. 420 = Elaphoglossum.
Webbii Bory; Fée, Acrost. 51 t. 24. 1845 = Elaphoglossum apodum.

Welwitschii Bak. Syn. 521. 1874 = Elaphoglossum.
Wightianum Wall. List n. 2163. 1829 = Polybotrya appendiculata.
Wightianum Pr. Tent. 241. 1836 = A. aureum.
Wilkesianum Hk. sp. **5**. 247. 1864 = Polybotrya.
Wrightii Mett.; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860 = Elaphoglossum.
xanthoneuron Kze.; Eat. Mem. Amer. Acad. n. s. **8**. 194. 1860 = Elaphoglossnm.
Yapurense vide japurense.
Yatesii Sod. Anal. Univ. Quito **4**. 174. 1890 = Elaphoglossum.
Yoshinagae Yatabe, Bot. Mag. Tokyo **5**. 109 t. 23. 1891 = Elaphoglossum.
yunnanense Bak. Kew Bull. **1898**. 233 = Elaphoglossum petiolatum var.
Zollingeri Kze. Bot. Zeit. **1846**. 419 = Leptochilus.

ACTINIOPTERIS Link, Fil. spec. 73, 79. 1841 — (Polypodiaceae Gen. **79**).
australis (L.fil.) Link, Fil. sp. 80. 1841 — Africa. Asia trop.
Acrostichum L.fil. Suppl. 444. 1781; Asplenium Sw. 1801; Belvisia Mirb. 1802; Pteris Thouars —.; Hk. et Grev. 1827; Acropteris Link 1833; Blechnum flabellatum Pr. 1836 — Acrostichum dichotomum Forsk. 1775 (non L. 1753); Pteris Kuhn 1867; Actiniopteris Kuhn 1871; Acrostichum radiatum König msc.; Poir. 1810; Asplenium Sw. 1801; Acropteris Link 1833; Blechnum Pr. 1836; Pteris Boj. 1837; Actiniopteris Link 1841. HB. 246. Chr. 175. NPfl. 288.
dichotoma Kuhn, Bot. Zeit. **1871**. 504 = A australis.
radiata Link, Fil. sp. 80. 1841 = A. australis.

ACTINOPHLEBIA Presl, Abh. böhm. Ges. V. **5**. 355. 1848 = **Hemitelia.**
horrida Pr. Abh. böhm. Ges. V. **5**. 355, 356. 1848.
obtusa Pr. Abh. böhm. Ges. V. **5**. 356. 1848 = H. subincisa.

Actinopteris J. Smith, Bot. Mag. **72** Comp. 20. 1846. *(Cheilanthes §)* = Adiantopsis.

ACTINOSTACHYS Wallich, List n. 1. 1828; Hook. Gen. t. 111. 1842 = **Schizaea.**
digitata Wall. List n. 1. 1828.
Germani Fée, 11 mém. 123 t. 29 f. 3. 1866.
pennula Hk. Gen. t. 111 A. 1842.
subtrijuga Pr. Suppl. 73. 1845 = S. penicillata.
trilateralis J. Sm. Lond. JoB. **1**. 202. 1842 = S. pennula.

ADECTUM Link, Fil. spec. 41, 42. 1841 = **Dennstaedtia.**
pilosiusculum Link, Fil. sp. 42. 1841 = D. punctilobula.

ADENODERRIS J. Smith, Hist. Fil 222. 1875 = **Polystichum.**
glandulosa J. Sm. Hist. Fil. 223. 1875 = P viscidulum.

ADENOPHORUS Gaudichaud, Ann. sc. nat. **3**. 508. 1824 = **Polypodium.**
bipinnatus Gaud. Ann. sc. nat. **3**. 508. 1824 = P. tamariscinum.
hymenophylloides Hk. et Grev. Ic. Fil. t. 176. 1830.
minutus Gaud. Ann. sc. nat. **3**. 508. 1824 = P. hymenophylloides.
? pinnatifidus Gaud. Freyc. Voy. Bot. 365. 1827 = P. adenophorus.
Tamarisci Hk. et Grev. Ic. Fil. t. 175. 1830 = P. tamariscinum.
tripinnatifidus Gaud. Ann. sc. nat. **3**. 508. 1824 = P. tamariscinum.

Adiantastrum Fée, 8 mém. 115. 1857 *(Adiantopsis §)* = Adiantopsis.

Adiantellum Presl, Tent. Pterid. 157. 1836 *(Adiantum §)* = Adiantum.

ADIANTOPSIS Fée. Gen. Fil. 145. 1850-52 — (Polypodiaceae Gen. **69**).
alata Prantl, Gartenfl. **32**. 99 t. 1115. 1883. NPfl. 272 — Brasilia. Guiana.
californica Moore, Ind. XXXVII. 1857 = Hypolepis.
capensis (Thbg.) Fée, Gen. 145. 1850-52 — Africa austr.

Adiantum Thbg. Prod. Fl. Cap. 173. 1800; Cheilanthes Sw. 1806. Mett. n. 75. HB. 132; Hypolepis Hk. 1852.

chlorophylla (Sw.) Fée, Gen. 145. 1850-52. NPfl. 272 — Amer. trop.
Cheilanthes Sw. Vet. Ak. Handl. **1817**. 76. Mett. n. 60. HB. 133. Chr. 143; C. brasiliensis Raddi 1819. C. spectabilis Klf. 1824; Hypolepis Link 1841; Adiantopsis Fée 1850-52.

dichotoma (Cav.) Moore, Ind. 17. 1857.; NPfl. 272 — Brasilia. Uruguay. Ecuador.
Pteris Cav. msc.; Cheilanthes Sw. Syn. 129, 335 t. 3 f. 7. 1806. Mett. n. 40. HB. 133. Chr. 143; Adiantum Poir. 1810; Pellaea Prantl 1882.

elongata Kl. Allg. Gartenzeit. **1855**. 392 = A. pedata.

Fordii (Bak.) C. Chr. Ind. 22. 1905 — Canton.
Cheilanthes Bak. JoB. 1879. 304.

incisa Moore, Ind. 243. 1861 = Hypolepis.

Lindigii (Mett.) Prantl, Gartenfl. **32**. 101. 1883 — Columbia.
Cheilanthes Mett. Ann. sc. nat. V. **2**. 218. 1864. HB. 475.

madagascariensis (Bak.) C. Chr. Ind. 22. 1905 — Madagascar.
Cheilanthes Bak. Journ. Linn. Soc. **16**. 198. 1877.

microphylla (Kl.) Prantl; Salom. Nom. 15. 1883 (nomen) — Brasilia.
Hypolepis Kl. msc.

monticola (Gardn.) Moore, Ind. XXXVII. 1857. NPfl. 272 — Brasilia.
Cheilanthes Gardn. in Hk. Ic. pl. t. 487. 1842. Mett. n. 57. HB. 131. Chr. 143; Hypolepis Hk. 1852; H. Gardneri Hk. 1852 — Cheilanthes trifurcata Bak. 1901!

obtusissima Fée, Cr. vasc. Br. **1**. 52 t. 13 f. 1. 1869 = A. regularis.

paupercula (Kze.) Fée, Gen. 145. 1850-52 — Cuba. Jamaica.
Adiantum Kze. Farnkr. **2**. 65 t. 127. 1850; Hypolepis Hk. 1852: Cheilanthes Mett. 1856. n. 55. HB. 132.

pedata (Hk.) Moore, Ind. 18. 1857. NPfl. 272 — Jamaica. Peru.
Hypolepis Hk. sp. **2**. 73 t. 92 A. 1852; Cheilanthes A.Br. 1857. Mett. n. 59. HB. 132 — Adiantopsis elongata Kl. 1855 (Peru).

propinqua (Mett.) Prantl, Gartenfl. **32**. 101. 1883 — Columbia.
Cheilanthes Mett. Ann. sc. nat. V. **2**. 218. 1864. HB. 475.

pteroides Moore, Ind. XXXVII. 1857 = Pellaea.

radiata (L.) Fée, Gen. 145. 1850-52. NPfl. 272 — Amer. trop.
Adiantum L. sp. **2**. 1094. 1753; Cheilanthes R.Br.; J. Sm. 1841. Mett. n. 58. HB. 132. Chr. 142; Hypolepis Hk. 1852.

Reesii (Jenm.) C. Chr. Ind. 22. 1905 — Jamaica.
Cheilanthes Jenm. JoB. **1886**. 267.

regularis (Kze.) Moore, Ind. 252. 1861. NPfl. 272 — Brasilia.
Adiantum Kze. 1850 (nomen); Cheilanthes Mett. Cheil. 41 n. 56. 1859. HB. 131; Adiantopsis obtusissima Fée 1869.

Schimperi Moore, Ind. XXXVII. 1857 = Hypolepis.

spectabilis Fée, Gen. 145. 1850-52 = A. chlorophylla.

ternata Prantl, Gartenfl. **32**. 101. 1883 — Orinoco.

ADIANTUM Linné (Gen. Pl. 782. 1737); Spec. plant. **2**. 1094. 1753 — (Polypodiaceae Gen. **78**).

abscissum Schrad. Gött. gel. Anz. **1824**. 872 — Brasilia.

achilleaefolium Lam. Enc. **1**. 43. 1783 = Asplenium.

acrocarpon Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 110 c. fig. — China.

aculeatum L. sp. **2**. 1096. 1753 = Odontosoria.

acuminatum Desv. Berl. Mag. 5. 327. 1811 = A. villosum.
adiantoides (J. Sm.) C. Chr. Ind. 23. 1905 — Guiana. Brasilia.
Hewardia J. Sm. JoB. 3. 432 t. 16-17. 1841; Adiantum Hewardia Kze. 1842. HB. 127. NPfl. 284 — Hewardia diphylla Fée 1869.
aemulum Moore, Gard. Chr. n. s. 8. 584 f. 114. 1877 — Brasilia.
aethiopicum L. Syst. Nat. ed. X. 2. 1329. 1759. HB. 123 (pt.). Chr. 139 (pt.) NPfl. 286 — ?Hispania. Afr. trop. et austr. Australia. N. Zealand. — A. assimile Sw. 1801 — A. trigonum Lab. 1806
affine Willd. sp. 5. 448. 1810 — N. Zealand.
affine Mart. et Gal. Mém. Ac. Brux. 15. 70. 1842 = A. concinnum.
affine Hk. sp. 2. 32. 1851 = A. diaphanum.
affine Moore, Ind. 19. 1857. HB. 117 et auctt. = A. Cunninghami.
africanum R. Br. in Tuckey, Narrat. of an Exp. to Zaire (App. V.) 463 nota 1818 = A. capillus veneris.
Alarconianum Gaud. Voy. Bonite Bot. t. 99. 1846 — Mexico-Ecuador. Galapagos. A. incisum Pr. Rel. Haenk. 1. 61 t. 10 f. 3. 1825. HB. 122. Chr. 138. NPfl. 283 (non Forsk. 1775).
altissimum Boj. Hort. Maur. 405. 1837 = Hypolepis sparsisora.
amabile Liebm. Vid. Selsk. Skr. V. 1. 265 (seors. 113). 1849 = A. andicola.
amabile Moore, Gard. Chr. 1868. 1090 = A. Moorei.
Amelianum Glaziou; Bak. JoB. 1882. 309 — Brasilia.
amoenum Wall. List n. 78. 1828; Hk. et Grev. Ic. Fil. t. 103. 1829 = A. [flabellulatum.
amplum Pr. Rel. Haenk. 1. 63. 1825. NPfl. 284 — Mexico.
A. glaucophyllum Hk. 1851. HB. 124 (pt.).
andicola Liebm. Vid. Selsk. Skr. V. 1. 266 (seors 114). 1849 — Mexico.
A. mexicanum Pr. 1836 (nomen) — A. amabile Liebm. 1849. A. glaucophyllum pt. Hk. 1854. HB. 124 (pt.).
aneitense Carr. in Seem. Fl. Vit. 346. 1873. HB. 472 — Aneiteum.
angustatum Klf. Enum. 202. 1824 — Brasilia.
apiculatum Schrad. Gött. gel. Anz. 1824. 872 = A. pulverulentum.
arborescens Poir. Enc. Suppl. 1. 144. 1810 = Hypolepis tenuifolia.
arcuatum Sw. Syn. 122. 1806 = A. lunulatum.
argutum Splitgb. Tijdschr. Nat. Gesch. 7. 427. 1840 = A. latifolium.
argutum Pr.; Ettingsh. Farnkr. 80 t. 41 f. 19. 1865 = A. venustum.
asarifolium Willd. sp. 5. 427. 1810 = A. reniforme*.
asperum Desv. Berl. Mag. 5. 327. 1811 = A. lucidum.
asperum Fée, Gen. 115. 1850-52 = A. striatum.
assimile Sw. Schrad. Journ. 1800[2]. 83. 1801 = A. aethiopicum.
assimile Link, Hort. Berol. 2. 17. 1833 = A. tenerum.
Aubertii Desv. Prod. 310. 1827 = A. Poiretii.
auriculatum Thbg. Prod. Fl. Cap. 173. 1800 = Pellaea.
Balansae Bak. JoB. 1890. 263 — Tonkin. China austr.
Balfourii Bak. in Hk. Ic. pl. t. 1630. 1886. NPfl. 283 — Socotra.
Bausei Moore, Gard. Chr. n. s. 12. 456 f. 69-70, 73. 1879 — Hort.
bellum Moore, Gard. Chr. n. s. 11. 172 f. 24. 1879 — Bermuda. Guiana.
Bennettii Carr. in Seem. Fl. Vit. 346. 1873. HB. 473 — Ins Hawaii.
Bessoniae Jenm. W. Ind. and Guiana Ferns 96. 1899 — Trinidad.
A. Bessonianum hrt. O'Brien: Gard. Chr. III. 20. 75. 1896 (nomen).
betulinum Klf. Enum. 207. 1824 = A. subcordatum.
Birkenheadii Moore, Gard. Chr. n. s. 25. 648. 1886 — Hort.

Bonii Christ, Journ. de Bot. 8. 150. 1894 — Tonkin. Siam.
Bonplandii Desv. Prod. 309. 1827 = A. serrato-dentatum.
borbonicum Jacq. Coll. **3**. 286 t. 21 f. 1. 1789 = Asplenium achilleifolium.
boreale Pr. Tent. 158. 1836 = A. pedatum.
brasiliense Raddi, Pl. Bras. **1**. 57 t. 76. 1825. Chr. 141 — Brasilia.
A. curvatum β HB. 474. A. pubescens Raddi 1819 (non Schkuhr 1809).
brasiliense Link, Hort. Berol. **2**. 13. 1833 = A. latifolium.
Braunii Mett.; Kuhn, Linn. **36**. 75. 1869 — Mexico. Venezuela.
A. convolutum Fourn. 1872. HB. 474.
Burnii Moore, Gard. Chr. III. **1**. 447. 1887 — Hort.
caffrorum L.fil. Suppl. 447. 1781 = Mohria.
caffrorum Sw. Schrad. Journ. **1800**². 85. 1801 = Cheilanthes hirta.
calcareum Gardn. in Hk. Ic. pl. t. 467. 1842. HB. 115. NPfl. 283 — Brasilia.
cantoniense Hance, Ann. sc. nat. IV. **15**. 229. 1861 = A. capillus junonis.
capense Thbg. Prod. Fl. Cap. 173. 1800 = Adiantopsis.
capillus Sw. Schrad. Journ. **1800**². 83. 1801 = A. capillus veneris.
Capillus Gorgonis Webb in Hk. Niger Fl. 192. 1849 = A. caudatum.
capillus junonis Rupr. Distr. Cr. vasc. Ross. 49. 1845; Milde, Bot. Zeit. **1867**. 148. HB. 114. NPfl. 283 — China.
A. cantoniense Hance 1861. HB. ed. I. 114.
capillus veneris L. sp. **2**. 1096. 1753. HB. 123. Chr. 138. NPfl. 284 — Europa occ. et austr. Africa. Asia temp.-Himalaya. Ceylon. Queensland. Polynesia. U. S. A. merid. et occ.-Columbia-Amazonas. ? Ind. occ.
A. capillus Sw. 1801 — A. emarginatum Bory; Willd. 1810 — A. Wattii Bak. 1881 — A. Levingei Bak. 1891. NPfl. 284.
cardiochlaena Kze. Linn. **17**. 569. 1843 = A. polyphyllum.
cassioides Desv. Prod. 309. 1827 = A. serrato-dentatum.
caudatum L. Mant. 308. 1771. HB. 115. Chr. 134. NPfl. 283 — Africa et Asia tropica. Ins. N. Hebridae.
— A. hirsutum Bory 1804 — A. ciliatum Bl. 1828.
cayennense Willd.; Kl. Linn. **18**. 552. 1844 — Guiana. Amazonas.
celebicum Christ, Verh. Nat. Ges. Basel **11**. 428. 1896; Ann. Jard. Buit. 15. 105. 1897 — Celebes.
Charlottae ht.; Möller's Gärtner Zeit. **1890**. 377; Gartenfl. **53**. 216. 1904 — Hort.
chilense Klf. Enum. 207. 1824. (HB. 123) — Peru-Chile. Juan Fernandez.
chinense Burm. Fl. Ind. 236. 1768 = Odontosoria.
chusanum L. sp. **2**. 1095. 1753 = Odontosoria chinensis.
cicutaefolium Lam. Enc. **1**. 44. 1783 = Cheilanthes tenuifolia.
cicutaefolium Noronha, Prod. Phyt. ex Thouars, Mél. bot. et voy. —. 1811 — [Madagascar.
ciliatum Bl. Enum. 215. 1828 = A. caudatum.
Claesianum L. Lind. et Rod. L'Ill. Hort. **1894**. 137 t. 9 — Hort. (e Brasilia).
Claussenii Fée, Gen. 115. 1850-52 = A. pulverulentum.
clavatum L. sp. **2**. 1096. 1753 = Odontosoria.
clavatum Forst. Prod. 83. 1786 = Odontosoria angustifolia.
Collisii Moore, Gard. Chr. n. s. **25**. 681. 1886 — Hort.
concinnum H.B. Willd. sp. **5**. 451. 1810. HB. 123. Chr. 139. NPfl. 286 — Amer trop. — A. tenerum Schkuhr 1809 (non Sw. 1788).
concinnum β subscandens Bak. Syn. ed. I. 123. 1867 = A. subvolubile.
colpodes Moore, Gard. Chr. **1865**. 530. HB. 124. NPfl. 286 — Ecuador. Peru. Argentina.

confine Fée, 10 mém. 14 t. 32 f. 1. 1865 — Ins. Comorae. Madagascar. Réunion.
conicum Vell. Fl. flum. **11** t. 97. 1827; Arch. Mus. Rio Jan. **5**. 455. 1881 = A. subcordatum.
convolutum Fourn. Mex. pl. **1**. 127 t. 6. 1872 = A. Braunii.
Cooperi Bak. JoB. **1887**. 25 — Costa Rica.
coriandrifolium Lam. Fl. franç. **1**. 29. 1778 = A. capillus veneris.
crenatum Willd. sp. **5**. 446. 1810 (Plum. t. 48) — Hispaniola.
A. quadriternatum Desv. 1811. (an A. cristatum?).
crenatum Poir. Enc. Suppl. **1**. 137. 1810 = A. Poiretii.
crenatum Bak. Syn. 120. 1867 = A. Wilesianum.
cristatum L. Syst. Nat. ed. X. **2**. 1328. 1759. HB. 118. Chr. 136. NPfl. 284 — Ind. occ. Guiana. Venezuela. Brasilia. — — A. tomentellum Fée 1869.
Cubense Hk. sp. **2**. 8 t. 73 A. 1851 = A. melanoleucum.
cultratum Willd. Phyt. 14 t. 10 f. 2. 1794 = Lindsaya.
cultratum Pr. Tent. 157. 1836 (nomen) — Brasilia. — (an seq.?).
cultratum J. Sm.; Hk. sp. **2**. 34. 1851 = A. trapezoides.
cultrifolium Noronha, Prod. Phyt. ex Thouars, Mél. bot. et voy. —. 1811 — [Madagascar.
cuneatum Forst. Prod. 84. 1786 = Lindsaya.
cuneatum Langsd. et Fisch. Ic. Fil. 23 t. 26. 1810. HB. 124. Chr. 138. NPfl. 286 — Brasilia-Argentina-Peru.
cuneifolium Stokes, Bot. Mat. Med. **4**. 612. 1812 = A. capillus veneris.
Cunninghami Hk. sp. **2**. 52 t. 86 A. 1851. NPfl. 286 — N. Zealand c. ins. adj. Australia. — A. affine auctt. HB. 117 (non Willd. 1810 t. Kuhn).
curvatum Klf. Enum. 202. 1824. HB. 126 (excl. var. 474). NPfl. 284 — Brasilia.
A. tetragonum Bak. Syn. ed. I. 126. 1867 (non Schrad. 1824).
cycloides Zenker, Pl. Ind. 11 t. 11. 1835 = A. Poiretii.
cyclosorum Moore, Gard. Chr. III. **1**. 547. 1887 — Ecuador.
Daddsii Moore, Gard. Chr. III. **1**. 42. 1887 — Hort.
Davidi Franch. Nouv. Arch. Mus. II. **10**. 112. 1887. NPfl. 284 — Tibet orient.
decipiens Desv. Prod. 307. 1827 = A. rhizophorum.
decorum Moore, Gard. Chr. **1869**. 582 = A. tinctum (Wagneri).
decurrens Jacq. Coll. **2**. 103 t. 2 f. 1, 2. 1788 = Hymenophyllum.
deflectens Mart. Ic. Cr. Bras. 94. 1834. NPfl. 283 — Venezuela. Guiana. Brasilia. — A. lunatum pt. Keys. 1875 — A. dolabriforme Hk. 1837 — A. flagellum Fée 1850-52 — A. subaristatum Fée 1869 — A. tremulum Kze. msc.; A. lunulatum β HB. 114.
delicatulum Mart. Ic. Cr. Bras. 93 t. 56 f. 2. 1834. NPfl. 283 — Brasilia.
A. filiforme Gardn. 1843. A. lunulatum pt. HB. 114. A. lunatum pt. Keys. 1875.
deltoideum Sw. Prod. 134. 1788. HB. 121. Chr. 138. NPfl. 283 — Ind. occ.
Allosorus domingensis Pr. 1836 — Adiantum jamaicense Fée 1866.
denticulatum Burm. Fl. Ind. 236. 1768 = Davallia.
denticulatum Sw. Prod. 135. 1788 = A. latifolium.
diaphanum Bl. Enum. 215. 1828. HB. 117. Chr. 135. NPfl. 286 — Asia trop. Polynesia. Australia. N. Zealand.
A. setulosum J. Sm. 1846. A. affine Hk. 1851 (non Willd. 1810) — A. heteromorphum Col.; Field 1890.
dichotomum Poir. Enc. Suppl. **1**. 144. 1810 = Adiantopsis.
digitatum Pr. Tent. 159. 1836 (nomen); Hk. sp. **2**. 38. 1851; Mett. Fil. Lechl. **2**. 12. 1859. HB. 125; Chr. 140. NPfl. 284 — Brasilia. Ecuador-Bolivia.
A. speciosum Hk. 1851.

dilatatum Schrad. Gött. gel. Anz. **1824**. 872 = A. subcordatum.

dioganum Glaziou: Bak. JoB. **1882**. 310 — Brasilia.

dissimile Schrad. Gött. gel. Anz. **1824**. 872 — Brasilia. (an A. delicatulum?).

dissimulatum Jenm. Gard. Chr. III. **16**. 656. 1894 — Jamaica.

dolabriforme Hk. Ic. pl. t. 191. 1837 = A. deflectens.

dolosum Kze. Linn. **21**. 219. 1848. HB. 127. Chr. 137. NPfl. 283 — Amer. trop. Hewardia Fée 1850-52. Lindsaya macrophylla Kze. 1837 (Anal. 37 pt. non t. 25); Hk. Ic. pl. t. 193. 1837 (non Klf. 1824) — Hewardia serrata Fée 1850-52. — *Adiantum Wilsoni Hk. 1851; Hewardia Fée 1850-52.

duale Jenm. Gard. Chr. III. **13**. 10. 1893 (err.) = Asplenium.

Edgeworthii Hk. sp. **2**. 14 t. 81 B. 1851. HB. 472 — China. India.

elatum Desv. Berl. Mag. **5**. 327. 1811 = A. tetraphyllum.

elegans Moore, Gard. Chr. n. s. **25**. 200. 1886 — Hort.

elegantissimum ht.; Gard. Chr. III. **12**. 249. 1892 — Hort.

emarginatum Bory; Willd. sp. **5**. 449. 1810 = A. capillus veneris var.

emarginatum Poir. Enc. Suppl. **1**. 139. 1810 = Schizoloma reniforme.

emarginatum Hk. sp. **2**. 39 t. 75 A. 1851 = A. Jordani.

eminens Pr. Tent. 158. 1836 = A. trapeziforme.

ensifolium Poir. Enc. Suppl. **1**. 139. 1810 = Schizoloma.

erectum Kze. Bot. Zeit. **1848**. 211 = A. diaphanum.

erythrochlamys Diels, Engl. Jahrb. **29**. 201. 1900 — China.

excisum Kze. Linn. **9**. 82. 1834; Anal. 33 t. 21. HB. 123. NPfl. 286 — Chile. Panama. Mexico.

extensum Fée, Gen. 114. 1850-52 (nomen); 8 mém. 72. 1857 = ? A. amplum.

Faberi Bak. JoB. **1888**. 225. NPfl. 284 — China.

falcatum Sw. Schrad. Journ. **1800**^{2}. 82. 1801 = A. villosum.

falcatum Vell. Fl. flum. **11** t. 96. 1827; Arch. Mus. Rio Jan. **5**. 455. 1881 = [A. tetraphyllum.

falcatum Blanco, Fl. Filip. 833. 1837 — Ins. Philipp.

falcinellum Desv. Berl. Mag. **5**. 326. 1811 — »Amer. trop.«

Farleyense Moore, Journ. Roy. Hort. Soc. n. s. **1**. 82 —; Gard. Chr. **1866**. 730 — Hort. (ex India occ.).

fasciculatum ht. May; Gard. Chr. III. **22**. 9. 1897 — Hort.

Faulkneri ht. Rochford; Gard. Chr. III. **24**. 241. 1898 — Hort.

Feei Moore; Fée, 7 mém. 29 t. 24 f. 1. 1857. HB. 125 (pt.). NPfl. 284 — Mexico.

Fergusoni Moore, Gard. Chr. n. s. **22**. 360. 1884 — Hort.

festum Moore, Gard. Chr. III. **1**. 110. 1887 — Hort.

filicaule Kze. Bot. Zeit. **1848**. 210 = A. diaphanum.

filiforme Gardn. in Hk. Ic. pl. t. 503. 1843 = A. delicatulum.

flabellifolium Lodd. Cat. 1849 = ? A. flabellulatum.

flabellulatum L. sp. **2**. 1095. 1753. HB. 126. Chr. 141. NPfl. 284 — Japonia. China-Asia trop. — A. amoenum Wall. 1828; Hk. et Grev. 1829.

flabellulatum Wall. List n. 2177. 1829 = A. hispidulum.

flabellum »Fée«; Christ, Farnkr. 134. 1897 (err.) = A. flagellum.

flagelliferum Wall. List n. 76. 1828 = A. caudatum.

flagellum Fée, Gen. 117. 1850-52; 6 mém. 4 t. 2 f. 1 = A. deflectens.

Flemmingii ht. van Houtte = A. concinnum var. hort.

flexuosum Hk. 2 Cent. t. 61. 1861 — Amer. centr. — A. Feei pt. HB. 125.

fontanum Salisb. Prod. 404. 1796 = A. capillus veneris.

formosissimum Kl. Linn. **18**. 556. 1844 = A. trapeziforme.

formosum R. Br. Prod. Fl. N. Holl. 155. 1810. HB. 119. Chr. 137. NPfl. 286 — Australia. N. Zealand.

formosum A. Cunn. in Hk. Comp. Bot. Mag. **2**. 366. 1836 = A. Cunninghami.

fovearum Raddi, Pl. Bras. **1**. 58 t. 77. 1825 = A. latifolium.

fragile Sw. Prod. 135. 1788. HB. 124. Chr. 139. NPfl. 286 — India occ. — A. parvifolium Fée 1857.

fragile Boj. Hort. Maur. 405. 1837 = Dennstaedtia anthriscifolia.

fragrans L.fil. Suppl. 447. 1781 = Cheilanthes pteridioides.

fragrantissimum Henders.; Moore, Gard. Chr. III. **1**. 447. 1887 — Hort.

fructuosum Kze.: Spr. Syst. **4**. 113. 1827; Kze. Linn. **9**. 81. 1834 = A. tetraphyllum var.

fruticosum Noronha, Prod. Phyt. ex Thouars, Mél. bot. et voy. —. 1811 — [Madagascar.

fuliginosum Fée, Gen. 116. 1850-52 = A. hirtum.

fulvum Raoul, Choix de Plantes 9. 1846. HB. 120. NPfl. 286 — Polynesia. Australia. N. Zealand.

fumarioides Willd. sp. **5**. 452. 1810 — Réunion.

furcatum L.fil. Suppl. 447. 1881 = Elaphaglossum.

fuscum Retz. Obs. **2**. 28 t. 5. 1781 = A. flabellulatum.

Galeottianum Hk. sp. **2**. 10 t. 80 B. 1851. HB. 116. NPfl. 286 — Mexico.

Ghiesbreghtii ht. Backh. (Gartenfl. **13**. 246. 1864) = A. tenerum.

gibbosum Roemer; Kuhn, Jahrb. bot. Gart. Berlin **1**. 344. 1881 — Brasilia.

glanduliferum Link, Hort. Berol. **2**. 18. 1833 = A. chilense.

glareosum Lindm. Ark. för Bot. **1**. 202 t. 9 f. 5. 1903 — Brasilia.

glaucescens Kl. Linn. **18**. 552. 1844. HB. 118 — Guiana-Amazonas. Ecuador.

glaucinum Kze.: Ettingsh. Farnkr. 82 t. 48 f. 7. 1865; Kuhn. Linn. **36**. 74. 1869 — Brasilia.

glaucophyllum Hk. sp. **2**. 40. 1851 (pt.). HB. 124 pt. = A. andicola.

glaucophyllum Hk. Ic. pl. t. 961. 1854. HB. 124 pt. = A. amplum.

Glaziovii Bak. JoB. **1882**. 309 — Brasilia.

globatum Poir. Enc. Suppl. **1**. 144. 1810 = Cheilanthes multifida.

gomphophyllum Bak. Ann. Bot. **8**. 124. 1894 — Malacca.

gracile Fée, Gen. 116. 1850-52; 7 mém. 27 t. 11 f. 1 — Brasilia.

gracillimum Moore, Gard. Chr. n. s. **1**. 14. 1874 — Hort.

grande Fée, Gen. 116. 1850-52 = A. polyphyllum.

gratum Fée, Gen. 119. 1850-52; 7 mém. 29 t. 12 f. 3 = A. Poiretii.

Gravesii Hance, JoB. **1875**. 197. Chr. 140. NPfl. 284 — China austr. Tonkin. A. Mariesii Bak. 1880.

Groenewegenianum Regel, Gartenfl. **16**. 16. 1867 — Hort. (India).

grossum Mett. Ann. sc. nat. V. **2**. 215. 1864. HB. 472. Chr. 135 — Columbia.

guianense Aublet, Hist. pl. Guiane **2**. 963. t. 365. 1775 = Lindsaya.

Haenkeanum Pr. Rel. Haenk. **1**. 62. 1825 = ? A. intermedium.

hastatum L.fil. Suppl. 447. 1781 = Pellaea viridis.

Hemsleyanum ht. May; Gard. Chr. III. **23**. 332 f. 124 1898 — Hort.

Hendersoni Linden, L'Ill. Hort. **20** t. 127. 1873 = A. tetraphyllum v. fruticosum.

Henslovianum Hk. fil. Tr. Linn. Soc. **20**. 169. 1847. HB. 118 (pt.) — Venezuela. Galapagos.

heteroclitum Christ, Bull. Boiss. II. **4**. 1094. 1904 — Costa Rica.

heteromorphum Col.: Field. Ferns N. Zeal. 80. 1890 = A. diaphanum var.

heterophyllum Poir. Enc. Suppl. **1**. 139. 1810 = Schizoloma.

heterophyllum Col. Tr. N. Zeal. Inst. **20**. 218. 1888 = A. Cunninghami.

Hewardia Kze. Farnkr. **1**. 104 t. 49. 1842 = A. adiantoides.
hexagonum L. sp. **2**. 1097. 1753 = Anopteris.
hians Moore, Gard. Chr. III. **1**. 41. 1887; Bak. Ann. Bot. **5**. 207. 1891 — [Fiji. N. Caledonia.
hippocrepis A.Br.; Salom. Nom. 19. 1883 (nomen) — ?
hirsutum Bory, Voy. **1**. 198. 1804 = A. caudatum.
hirtum Poir. Enc. Suppl. **1**. 142. 1810 = Cheilanthes.
hirtum Splitgb. Tijdschr. Nat. Gesch. **7**. 428. 1840. HB. 119 (pt.) — Guiana. Brasilia.
hirtum Kl. Linn. **18**. 563. 1844; Hk. sp. **2**. 20 t. 82 A. 1851. HB. 119 (pt.) = A. terminatum.
hispidulum Sw. Schrad. Journ. **1800**^{2}. 82. 1801. HB. 126. Chr. 140. NPfl. 286 — India austr.-Malesia-Polynesia. Australia. N. Zealand. Afr. austr. cum. ins. adj. ? Jamaica. ? Peru.
A. pedatum Forst. 1786 (non L. 1753). A. pubescens Schkuhr 1809 — A. rigidum Fourn. 1873 (non Link 1841).
Hornei Bak. JoB. **1879**. 294 — Fiji.
Hosei Bak. JoB. **1888**. 324 — Borneo.
humile Kze. Linn. **9**. 80. 1834 = ? A. intermedium.
incertum Lindm. Ark. för Bot. **1**. 204 t. 9 f. 4. 1903 — Brasilia.
incisum Forsk. Fl. Aeg. Arab. 187. 1775 = A. caudatum.
incisum Pr. Rel. Haenk. **1**. 61 t. 10 f. 3. 1825 = A. Alarconianum.
integrifolium Poir. Enc. Suppl. **1**. 139. 1810 = Lindsaya falcata.
intermedium Sw. Vet. Ak. Handl. **1817**. 76 — Brasilia.
intermedium auctt. HB. 116 = A. latifolium.
Irvinianum Linden, Cat. 1854 — Hort.
Jacobinae Fée. Gen. 115. 1850-52 = A. serrato-dentatum.
jamaicense Fée, 11 mém. 19 t. 33 f. 3. 1866 = A. deltoideum.
Jordani K. Müll. Bot. Zeit. **1864**. 26 t. 1 f. 1 — U. S. A. pacif.
A. emarginatum Hk. sp. **2**. 39 t. 75 A. 1851. (Chr. 139). (non Bory 1810).
Kalbreyeri C. Chr. Ind. 28. 1905 — Columbia.
A. pilosum Bak. Ann. Bot. **5**. 207. 1891 (non Fée 1850-52).
Kaulfussii Kze. Linn. **21**. 221. 1848 = A. petiolatum.
Kendalli Jenm. JoB. **1886**. 39 — Jamaica.
Klotschianum Pr. Tent. 158. 1836 = A. subcordatum.
Klotzschianum Hk. sp. **2**. 21 t. 82 C. 1851 = A. tomentosum.
Kohautianum Pr. Tent. 158. 1836 = A. tetraphyllum.
Kunzeanum Pr. Tent. 157. 1836 = A. pulverulentum.
Kunzeanum Kl. Linn. **18**. 555. 1844 = A. melanoleucum.
Kunzei Miq. Diar. Inst. Reg. Bat. **1843**. 5 t. 1 = A. serrato-dentatum.
laetum Pr. Tent. 158. 1836 (nomen) = A. melanoleucum.
laetum Mett.; Kuhn, Linn. **36**. 76. 1869 — Peru. — A. Henslovianum pt. [HB. 118.
Lambertianum ht.: Wien. Ill. Gartenzeit. **1890**. 345 — Hort.
lancea L. sp. ed. II. **2**. 1557. 1763 = Lindsaya.
lancea Bak. Fl. bras. **1**^{2}. 373. 1870 = A. tetraphyllum.
lancea Keys. Mém. Ac. St. Pétersb. **22**^{2}. 6, 28. 1875 = A. macrophyllum.
lanceolatum Poir. Enc. Suppl. **1**. 134. 1810 = Schizoloma ensifolium.
lanceolatum Fée, Gen. 115. 1850-52 = A. villosum.
Lathomi ht. Wills: Moore, Gard. Chr. n. s. **16**. 685. 1881 — Hort.
latifolium Lam. Enc. **1**. 43. 1783 — Amer. trop.
A. denticulatum Sw. Prod. 135. 1788 (non Burm. 1768). Chr. 134. NPfl.

283\. A. fovearum Raddi 1825. A. intermedium auctt. HB. 116. ? Chr. 135. (an adhuc A. petiolatum Desv. 1811 cum syn.?).
laxum Kze. Linn. **9**. 79. 1834 = A. tetraphyllum.
Legrandii ht. Veitch; Gard. Chr. III. **15**. 663. 1894 — Hort.
lendigerum Poir. Enc. Suppl. **1**. 142. 1810 = Cheilanthes.
Leprieurii Hk. sp. **2**. 31 t. 82 B. 1851. HB. 127 — Guiana.
Hewardia Fée 1850-52.
Levingei Bak. Ann. Bot. **5**. 207. 1891 = A. capillus veneris var.
Lindeni Moore, Gard. Chr. **1866**. 778. HB. 473 — Amazonas.
lindsaea Cav. Descr. 271. 1802 — Ecuador.
lineare Poir. Enc. Suppl. **1**. 139. 1810 = Lindsaya.
lineatum L. Linden, L'Ill. Hort. **1895**. 185, 312 t. 44 — Hort.
littorale Jenm. W. Ind. and Guiana Ferns 96. 1899 — Jamaica.
lobatum Poir.; Steud. Nomenkl. **2**. 275. 1824 = Lindsaya decomposita.
lobatum Pr. Rel. Haenk. **1**. 62 t. 10 f. 4. 1825 = ? A. chilense. (an seq.?).
lobatum Kze.; Kuhn, Jahrb. bot. Gart. Berlin **1**. 348. 1881 (an Pr.?) — Mexico.
Lobbianum Hk. sp. **2**. 51 t. 86 C. 1851 = A. pulchellum.
lobulatum Kze. Bot. Zeit. **1846**. 445 (nomen); Ettingsh. Farnkr. 82 f. 32 t. 47 f. 7. 1865 = A. hispidulum.
Lorentzii Hier. Engl. Jahrb. **22**. 393. 1896 — Argentina.
lucidum (Cav.?) Sw. Syn. 121. 1806. HB. 121. Chr. 137. NPfl. 283 — Amer. trop. — ? Pteris Cav. Descr. 266. 1802. P. aspera Poir. 1804; Adiantum Desv. 1811 — ? A. varium H.B. Willd. 1810.
Lüddemannianum ht; Veitch, Cat. **1878**. 68 c. fig. — Hort.
lunatum Cav. Descr. 272. 1802 = A. lunulatum.
lunatum Keys. Mém. Ac. St. Pétersb. **22**^{2}. 2, 25. 1875 = A. lunulatum, deflectens, delicatulum.
lunulatum Burm. Fl. Ind. 235. 1768. HB. 114 (pt.). Chr. 134 (pt.). NPfl. 283 — Africa, Asia, Australia, Polynesia trop. Amer. centr.
Pteris Retz. 1781. ? Adiantum philippense L. 1753. A lunatum Cav. 1802. A. arcuatum Sw. 1806. — conf. A. deflectens et delicatulum.
lunulatum Houtt. Nat. Hist. **14**. 209. 1783; Pfl. Syst. **13**^{1}. 252 t. 100 f. 1. 1786 = Didymochlaena truncatula.
lutescens Moug.; Fée, Gen. 119. 1850-52 = A. concinnum.
lyratum Blanco, Fl. Filip. 832. 1837 — Ins. Philipp.
macrocarpum Pr. Tent. 158. 1836 = A tetraphyllum.
macrocladum Kl. Linn. **18**. 554. 1844 — Peru. — A. polyphyllum pt. HB. 118.
macrodus Kze. Flora. **1839**^{1}. Beibl. 42 = A. obliquum.
macrophyllum Sw. Prod. 135. 1788. HB. 121. Chr. 137. NPfl. 283 — Amer. trop. — A. lancea Keys. 1875 (non L. 1753).
macropterum Miq.; Kuhn. Linn. **36**. 72. 1869 — Guiana. — (an A. olivaceum?).
macropus Schrad. Gött. gel. Anz. **1824**. 872 = A. obliquum.
maderense Lowe ? = A. capillus veneris.
Mairisii Moore, Gard. Chr. n. s. **24**. 294. 1885 — Hort.
malaliense Jenm. Gard. Chr. III. **20**. 182. 1896 — Guiana.
manicatum ht.; Garden **88**. 94. 1900 — Hort.
marginatum Schrad. Gött. gel. Anz. **1818**. 918 = Adiantopsis capensis.
Mariesii Bak. Gard. Chr. n. s. **14**. 494. 1880 = A. Gravesii.
Mathewsianum Hk. sp. **2**. 35 t. 84 A. 1851 — Peru. — A. polyphyllum pt. HB. 118.

melanoleucum Willd. sp. **5**. 443. 1810. Chr. 136. NPfl. 284 — Ind. occ.
A. Kunzeanum Kl. 1844 — A. cubense Hk. 1851. HB. 119.
Mettenii Kuhn, Fil. Afr. 65. 1868 (nomen). HB. 472; Kuhn, Jahrb. bot. Gart. Berlin **1**. 338. 1881 — Afr. occ. trop. India. Ins. Philipp.
mexicanum Pr. Tent. 158. 1836 = A. andicola.
Meyerianum Zoll. Nat. Gen. Arch. **2**. 204. 1845; Hassk Flora **1847**[1]. 318 = [A. hispidulum.
microcarpum Pr. Tent. 158. 1836 (nomen) — ?
microphyllum Sw. Prod. 135. 1788 = Cheilanthes.
microphyllum Poir. Enc. Suppl. **1**. 140. 1810 = Lindsaya.
microphyllum Hornem. Ht. reg. bot. Hafn. Suppl. 110. 1819 = ? Cheilanthes.
microphyllum Klf. Enum. 204. 1824 = A. pyramidale.
microphyllum Roxb. Calc. Journ. **4**. 513. 1844 = A venustum.
micropteris Poir. Enc. Suppl. **1**. 141. 1810 = Cheilanthes.
microsorum C. Chr. Ind. 30. 1905 — Peru. — A. Veitchianum Moore, Gard. Chr. **1868**. 1090. HB. 473 (non A. Veitchii Hance 1861).
modestum Und. Bull. Torr. Cl. **28**. 46. 1901 — New Mexico.
monochlamys Eat. Proc. Amer. Acad. **4**. 110. 1858. HB. 125. Chr. 140. NPfl. 284 — Japonia. China. — — A. Veitchii Hance 1861.
monosoratum Willd. sp. **5**. 445. 1810 = A. pulverulentum.
monosorum Bak. Syn. 472. 1874. NPfl. 286 — N. Caledonia.
monotis Nees, Linn. **19**. 684. 1847 = A. villosum.
Moorei Bak. Gard. Chr. **1873**. 811. HB. 474 — Peru.
A. amabile Moore, Gard. Chr. **1868**. 1090 (non Liebm. 1849).
Moritzianum Link, Fil. sp. 71. 1841 = A. capillus veneris.
Muelleri (F. Muell. Fragm. **7**. 121. 1870) = A. diaphanum.
multifidum Sw. Schrad. Journ. **1800**[2]. 85. 1801 = Cheilanthes.
multiforme A.Br.; Fourn. Mex. pl. **1**. 128. 1872 = A. amplum.
mundulum Moore, Gard. Chr. n. s. **12**. 84. 1879 — Hort.
myriophyllum Pr. Tent. 158. 1836 (nomen) = A. macrocladum.
myriosorum Bak. Kew Bull. **1898**. 230 = A. pedatum.
nebulosum ht. Pynaert; Gard. Chr. III. **13**. 415. 1893 — Hort.
neoguineense Moore, Gard. Chr. n. s. **7**. 12. 1877; Bak. Ann. Bot. **5**. 208. 1891 — N. Guinea.
nervosum Sw. Syn. 123. 1806 = A hispidulum. (an seq.?).
nervosum Kuhn, Jahrb. bot. Gart. Berlin **1**. 342. 1881 (an Sw.?) = A. serrato-dentatum.
nigrescens Fée, Gen. 117. 1850-52; 7 mém. 28 t. 11 f. 2. HB. 117 — Ind. [occ. — (an A. striatum?).
nobile Fisch.; ht. = A. tenerum.
novae caledoniae Keys. Mém. Ac. St. Pétersb. **22**[2]. 4, 27 c. tab. 1875 — N. Caledonia.
Oatesii Bak. in Oates, Matabeleland, App. V. c. fig. 1881 — Africa austr.
oblique-truncatum Fée, 11 mém. 18 t. 7 f. 2. 1866 = A. villosum.
obliquum Willd. sp. **5**. 429. 1810. HB. 115 — Amer. trop.
obliquum Klf. Enum. 200. 1824; Hk. et Grev. Ic. Fil. t. 190. 1830 = A. petio- [latum.
oblongatum Schrad. Gött. gel. Anz. **1824**. 872 = A. petiolatum.
obtusum Desv. Berl. Mag. **5**. 327. 1811 = A. serrato-dentatum.
odoratum Poir. Enc. Suppl. **1**. 142. 1810 = Cheilanthes pteridioides.
odoratum Wallis — Hort.
odorum DC. Fl. franç. **5**. 237. 1815 = Cheilanthes pteridioides.
olivaceum Bak. Syn. 127. 1867. NPfl. 284 — Guiana. — (conf. A. macropterum).

orbiculatum Lam. Enc. **1**. 41. 1783 = Lindsaya.

Orbignyanum Mett.; Kuhn, Linn. **36**. 78. 1869 — Bolivia. Peru.

ornithopodum Pr. Tent. 158. 1836 (nomen): Kuhn, Linn. **36**. 74. 1869 — [Brasilia.

ovalescens Fée, 8 mém. 72. 1857 — S. Domingo.

ovatifolium Noronha, Prod. Phyt. ex Thouars, Mél. bot. et voy. —. 1811 — [Madagascar.

Oweni Moore, Gard. Chr. III. **1**. 110. 1887 — Hort.

oyapokense Jenm. W. Ind. and Guiana Ferns 85. 1899 — Guiana.

pachysorum Rchb.; Kze. Linn. **21**. 224. 1848 = A. tetraphyllum.

Pacotti ht.; Gard. Chr. n. s. **25**. 14. 1886 — Hort.

pallens Sw. Schrad. Journ. **1800**^{2}. 84. 1801 = Ochropteris.

palmatum Schum. Vid. Selsk. Afh. **4**. 234 (seors. 460). 1827 = Doryopteris concolor.

palmatum Moore, Gard. Chr. n. s. **7**. 40 f. 5. 1877 — Hort. (Peru. Ecuador).

papyraceum Desv. Prod. 307. 1827 — Réunion. — (Asplenium sp. stat. ster.).

Paradiseae Bak. Gard. Chr. III. **6**. 558. 1889 — Africa austr.

paradoxum R.Br. Prod. Fl. N. Holl. 155. 1810 = Pellaea.

parallelogrammum Vell. Fl. flum. **11**. t. 99. 1827; Arch. Mus. Nac. Rio Jan. **5**. 456. 1881 = A. polyphyllum.

Parishii Hk. Fil. Exot. t. 51. 1858. HB. 114. Chr. 133. NPfl. 282 — Birma.

parvifolium Fée, 7 mém. 27 t. 23 f. 2. 1857 = A. fragile.

parvilobum Sw. Schrad. Journ. **1800**^{2}. 85. 1801 = Cheilanthes hirta var.

parvulum Hk. fil. Tr. Linn. Soc. **20**. 168. 1847 — Galapagos.

patens Willd. sp. **5**. 439. 1810. HB. 126. Chr. 141. NPfl. 286 — Mexico -Ecuador. Galapagos.

pauperculum Kze. Farnkr. **2**. 65 t. 127. 1850 = Adiantopsis.

Pearcei Phil. Anal. Univ. Chile **1861**. 68; Linn. **33**. 304. 1864 — Chile.

pectinatum Kze., Ettingsh. Farnkr. 85 t. 45 f. 14-16. 1865. HB. 120. Chr. 136. NPfl. 284 — Brasilia. Peru.

pedatum L. sp. **2**. 1095. 1753. HB. 125. Chr. 140. NPfl. 284 — Alaska. Amer. bor. temp. Japonia. China. Himalaya. — A. boreale Pr. 1836 (Unalaschka) — A. myriosorum Bak. 1898 (China austr.).

pedatum Forst. Prod. 83. 1786 = A. hispidulum.

pedatum Raddi, Opusc. sci. Bol. **3**. 295. 1819 = A. brasiliense.

pellucidum Mart. et Gal. Mém. Ac. Brux. **15**. 72 t. 19. 1842 = A. Poiretii.

pensile Kze.; Fée, Gen. 114. 1850-52 = A. sinuosum.

pentadactylon Langsd. et Fisch. Ic. Fil. 22 t. 25. 1810 = A. trapeziforme var.

persimile Pr.; Ettingsh. Farnkr. 85 t. 46 f. 4-5. 1865 = A. deflectens.

peruvianum Kl. Linn. **18**. 555. 1844. HB. 116. Chr. 134. NPfl. 284 — Peru.

petiolatum Desv. Berl. Mag. **5**. 326. 1811. — Amer. centr. et austr. trop. A. obliquum Klf. 1824 (non Willd. 1810); A. Kaulfussii Kze. 1848. HB. 115 — (an A. latifolium?).

philippense L. sp. **2**. 1094. 1753 = ? A. lunulatum.

phyllitidis J. Sm. Lond. JoB. **1**. 197. 1842. HB. 121. NPfl. 283 — Amer. centr. -Peru. Guadeloupe. — A. Poeppigianum Pr. 1836 (nomen).

pilosum Fée, Gen. 118. 1850-52 = A. chilense.

pilosum Bak. Ann. Bot. **5**. 207. 1891 = A. Kalbreyeri.

platyphyllum Sw. Vet. Akad. Hdl. **1817**. 74 t. 3 f. 6 — Amer. centr. Brasilia. Bolivia. — A. Seemanni Hk. 1851. HB. 121. Chr. 137.

platyphyllum Kze. Anal. 31 t. 20. 1837 = A. petiolatum.

platyphyllum Bak. Fl. bras. **1**^{2}. 363. 1870 = A. obliquum et petiolatum.

plicatum Klf. Enum. 201. 1824 = A. hispidulum.

Poeppigianum Pr. Tent. 157. 1836 = A. phyllitidis.
Poiretii Wikstr. Vet. Akad. Hdl. **1825**. 443. 1826 — Africa trop. et subtrop. Penins. Indiae orient. Mexico-Amer. austr. trop.
A. crenatum Poir. Enc. Suppl. **1**. 137. 1810 (non Willd. 1810). NPfl. 286; A. Aubertii Desv. Prod. 1827. A. thalictroides Willd.; (Schlecht. 1832.) Kze. 1836 — A. pellucidum Mart. et Gal. 1842. A. gratum Fée 1850-52. A. aethiopicum pt. auctt. HB. 123. — (conf. Kuhn, Abh. Nat. Ges. Halle **11**. 29.)
politum H.B. Willd. sp. **5**. 442. 1810 — Cumana.
politum J. Sm. Lond. JoB. **1**. 198. 1842 = A. polyphyllum.
polymorphum Poir. Enc. Suppl. **1**. 142. 1810 = Doryopteris pilosa.
polymorphum Col. Tr. N. Zeal. Inst. **20**. 215. 1888 = A. diaphanum.
polyphyllum Willd. sp. **5**. 454. 1810. HB. 118. Chr. 135. NPfl. 284 — Venezuela-Peru. — A. cardiochlaena Kze. 1843.
Prattii Bak. Journ. Linn. Soc. **29**. 321. 1892 — China.
princeps Moore, Gard. Chr. n. s. **4**. 197 f. 43-44. 1875 — Columbia.
prionophyllum HBK. Nov. Gen. et Sp. **1**. 20. 1815 = A. tetraphyllum.
proliferum Roxb. Calc. Journ. **4**. 512. 1844 = A. caudatum.
propinquum Fée, Gen. 114. 1850-52 = Guiana.
proximum Gaud. Freyc. Voy. Bot. 403. 1827 = A. serrato-dentatum.
pseudo-capillus Fée, Gen. 118. 1850-52: 7 mém. 29 t. 1 = A. capillus veneris.
pseudotinctum Hier. Engl. Jahrb. **22**. 394. 1896. NPfl. 286 — Paraguay.
pteroides L. Mant. 130. 1767 = Pellaea.
pubescens Schkuhr, Kr. Gew. **1**. 108 t. 116. 1809 = A. hispidulum.
pubescens Poir. Enc. Suppl. **1**. 141. 1810 = Cheilanthes microphylla.
pubescens Raddi, Opusc. sci. Bol. **3**. 295. 1819 = A. brasiliense.
? pubescens Pr. Rel. Haenk. **1**. 63. 1825 = A. chilense.
pulchellum Bl. Enum. 216. 1828. HB. 120. NPfl. 286 — Java.
A. Lobbianum Hk. 1851.
pullum Col. Tr. N. Zeal. Inst. **25**. 319. 1893 — N. Zealand.
pulverulentum L. sp. **2**. 1096. 1753. HB. 122. Chr. 138. NPfl. 283 — Ind. occ. -Brasilia. — A. monosoratum Willd. 1810 — A. serrulatum L. 1753.
pumilum Sw. Prod. 134. 1788. HB. 472. NPfl. 283 — Jamaica.
pusillum All. (Poll. Fl. Veron. **3**. 293. 1824) = Cheilanthes pteridioides.
pyramidale (L.) Willd. sp. **5**. 442. 1810 — Ind. occ.
Polypodium L. sp. **2**. 1093. 1753. Adiantum microphyllum Klf. Enum. 204. 1824. HB. 122 (t. Jenm.). A. striatum Schkuhr 1809 (non Sw. 1788).
quadriternatum Desv. Berl. Mag. **5**. 327. 1811 = A. crenatum Willd.
Raddianum Pr. Tent. 158. 1836 = A. cuneatum.
radiatum L. sp. **2**. 1094. 1753 = Adiantopsis.
radicans Fée, Gen. 118 t. 29 f. 2. 1850-52 = A. caudatum.
rectangulare Lindm. Ark. för Bot. **1**. 204 t. 9 f. 3. 1903 — Brasilia.
refractum Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 224 — China.
reginae Veitch, Cat. 1888; Wien. Ill. Gartenzeit. **1888**. 107 — Hort.
regulare Kze. Farnkr. **2**. 66. 1850 = Adiantopsis.
reniforme L. sp. **2**. 1094. 1753. HB. 114. Chr. 133. NPfl. 283 — Ins. Atlantidis.
— * A. asarifolium Willd. 1810. NPfl. 283 — Ins. Mascarenae.
— var. crenatum Bak. JoB. **1891**. 4. NPfl. 283 — Madagascar.
reniforme Sessé et Mociño, Fl. Mex. 261. 1896 — Mexico.
repandum Tausch in Sieb. Fil. exs. n. 176 = A. capillus veneris.
repens L.fil. Suppl. 446. 1781 = Humata.

rhizophorum Sw. Syn. 320, 422. 1806. HB. 115 (A. caudatum β). Chr. 134. NPfl. 283 — Afr. austr. cum ins. adjac. Timor.
rhizophyllum Pr. Tent. 157. 1836 = A. rhizophytum.
rhizophytum Schrad. Gött. gel. Anz. **1824**. 872; Mart. Ic. Cr. Bras. 92 t. 62. (HB. 472) — Brasilia.
rhodophyllum Moore, Gard. Chr. n. s. **21**. 372. 1884 — Hort.
rhomboideum Schkuhr, Kr. Gew. **1**. 114 t. 122. 1809 = A. trapeziforme.
rhomboideum HBK. Nov. Gen. et Sp. **1**. 20. 1815 = A. serrato-dentatum.
rigidulum Mett.; Kuhn, Linn. **36**. 76. 1869. HB. 474 — Ind. occ.
rigidum Pr. Tent. 158. 1836 = A. serrato-dentatum.
rigidum Link, Fil. sp. 69. 1841 = A. tetraphyllum.
rigidum Fourn. Ann. sc. nat. V. **18**. 329. 1873 = A. hispidulum.
Roborowskii Maxim. Mél. biol. **11**. 867. 1883 — China occ.
rotundatum Desv. Prod. 310. 1827 = A. chilense.
rotundatum Kze. Linn. **10**. 528. 1836 = A. rhizophorum.
rubellum Moore, Gard. Chr. **1868**. 866. HB. 474 — Bolivia.
rufopunctatum Mett.; Kuhn, Jahrb. bot. Gart. Berlin **1**. 350. 1881 — Bolivia.
Ruizianum Kl. Linn. **18**. 551. 1844; Bak. Ann. Bot. **5**. 206. 1891 — Peru.
sagittatum Aublet, Hist. pl. Guiane **2**. 964 t. 366 1775 = Schizoloma.
sanctae catherinae ht.; J. Sm. Ferns br. and for. ed. II. 310. 1877 — Hort. [(e Brasilia).
scabrum Klf. Enum. 207. 1824. NPfl. 286 — Chile.
scabrum Wall. List n. 79. 1828 = A. hispidulum.
scandens Lour. Fl. Cochinch. **2**. 837. 1790 = Lygodium japonicum.
scandicinum Willd.; Sw. Syn. 326. 1806 (in obs.) = Cheilanthes mysurensis.
Schaffneri Fourn. Bull. Soc. Fr. **27**. 328. 1880 — Mexico.
schizophyllum Moore, Wiener Illustr. Gartenzeit. **1887**. 377 — Hort.
Schmidtchenii Hier. Engl. Jahrb. **34**. 487. 1904 — Columbia.
Schweinfurthii Kuhn, Sitzungsber. Ges. nat. Fr. Berlin **1869**. 40; Bot. Zeit. **1870**. 89. (HB. 472). NPfl. 283 — Afr. centr. trop.
scutum ht. = A. tenerum.
Seemanni Hk. sp. **2**. 5 t. 81 A. 1851 = A. platyphyllum.
Sellowianum Pr. Tent. 159. 1836 (nomen) — Brasilia.
Senae Bak. JoB. **1885**. 217 — Brasilia.
sericeum Eat. Bot. Zeit. **1869**. 361. HB. 473. NPfl. 283 — Cuba.
serrato-dentatum Willd. sp. **5**. 445. 1810. Chr. 136 — Amer. trop.
A. obtusum Desv. 1811. HB. 119. A. rhomboideum HBK. 1815.
A. nervosum Kuhn 1881 (an Sw. 1806?).
serrulatum L. sp. **2**. 1095. 1753 = A. pulverulentum.
serrulatum (Sw. Prod. 135. 1788?) Keys. Mém. Ac. St. Pétersb. **22**[2]. 9. 37. 1875 = A. tetraphyllum.
sessilifolium Hk. sp. **2**. 44 t. 85 B. 1851 = A. Henslovianum.
setulosum J. Sm. Bot. Mag. **72** Comp. 22. 1846 = A. diaphanum.
Shepherdi Hk. sp. **2**. 9 t. 73 B. 1851. HB. 116. NPfl. 286 — Mexico.
sinuosum Gardn. Hk. Ic. pl. t. 504. 1843. HB. 117 — Brasilia.
soboliferum Wall. List n. 74. 1828; Hk. sp. **2**. 13 t. 74 A. 1851. HB. 115 — [India.
sordidum Lindm. Ark. för Bot. **1**. 205 t. 9 f. 1. 1903 — Brasilia.
speciosum Hk. sp. **2**. 45 t. 85 C. 1851 = A. digitatum.
Steerei Harr. Journ. Linn. Soc. **16**. 34. 1877 — Peru.
stellatum Warsz.; Salom. Nom. 22. 1883 — Guatemala.
stenochlamys Bak. Ann. Bot. **5**. 209. 1891 — Borneo. Pulo Gaya.

striatum Sw. Prod. 135. 1788 — Ind. occ. (conf. A. nigrescens).
striatum Schkuhr, Kr. Gew. **1**. 109 t. 118 f. a-g. 1809 = A. pyramidale.
strictum Sw. Prod. 135. 1788 = Lindsaya.
suaveolens Poir. Enc. Suppl. **1**. 142. 1810 = Cheilanthes pteridioides.
subaristatum Fée, Cr. vasc. Br. **1**. 33 t. 8 f. 2. 1869 = A. deflectens.
subcordatum Sw. Vet. Ak. Hdl. **1817**. 75. HB. 116. Chr. 135 — Guiana. Brasilia
A. truncatum Raddi 1819. A. betulinum Klf. 1824.
subemarginatum Christ, Bull. Boiss. II. **3**. 511. 1903 — China.
?sublobatum Schum. Vid. Selsk. Afh. **4**. 235 (seors. 461). 1827 = Arthropteris obliterata.
subramosum Fée. Cr. vasc. Br. **1**. 36 t. 9 f. 1. 1869 = A. curvatum.
subsulphureum Remy; Gay, Fl. Chil. **6**. 486. 1853 = A. sulphureum.
subtrapezoideum Christ, Bull. Boiss. II. **4**. 1094. 1904 — Costa Rica.
subvolubile Mett.; Kuhn, Linn. **36**. 77. 1869. HB. 473 — Ecuador.
A. concinnum β subscandens Bak. 1867. HB. ed. I. 123 (sine descript.)
sulphureum Klf. Enum. 207. 1824 — Chile.
tenellum Jacq. Coll. **3**. 287 t. 21 f. 3. 1789 = Hymenophyllum.
tenellum Moore, Veitch Cat. 1855 = A. hispidulum.
tenerum Sw. Prod. 135. 1788. HB. 124. Chr. 139. NPfl. 286 — Bermuda. Ind. occ. Mexico-Juan Fernandez.
tenerum Schkuhr, Kr. Gew. **1**. 112 t. 121. 1809 = A. concinnum.
tenerum L. v. Buch, Abh. Akad. Berlin **1816-17**. 360. 1819. Roxb. Calc. Journ. **4**. 513. 1844 = A. capillus veneris.
tenuifolium Lam. Enc. **1**. 44. 1783 = Odontosoria chinensis.
tenuifolium Sw. Schrad. Journn. **1800**². 85. 1801 = Cheilanthes.
tenuifolium Blanco, Fl. Filip. 832. 1837 — Ins. Philipp.
tenuissimum Taubert, Engl. Jahrb. **21**. 421. 1896 — Brasilia centr.
terminatum Kze.; Miq. »Diar. Inst. Reg. Bat. **1843**. 3«; Kze. Linn. **21**. 222. 1848. NPfl. 284 — Guiana. Brasilia.
A. hirtum Kl. Linn. **18**. 536. 1844. Hk. 1851. HB. 119 (pt.) (non Splitgb. 1840).
ternatum H.B. Willd. sp. **5**. 436. 1810 = A. tetraphyllum.
tetragonum Schrad. Gött. gel. Anz. **1824**. 872; Mart. Ic. Cr. Bras. 93 t. 63. HB. 475. NPfl. 284 — Brasilia.
tetragonum Bak. Syn. ed. I. 126. 1867 = A. curvatum.
tetraphyllum H.B. Willd. sp. **5**. 441. 1810. HB. 120. Chr. 136. NPfl. 283 — Amer. trop. Afr. occ. trop.
Synaphlebium Fée 1850-52. Adiantum prionophyllum HBK. 1815; A. lancea Bak. 1870 (non L. 1753) — A. ternatum H. B. Willd. 1810 — A. fructuosum Kze.; Spr. 1827 — A. Vogelii Mett.; Keys. 1875 (Afr.).
thalictroides Willd.; (Schlecht. Adumbr. 53 t. 33 ined.) Kze. Linn. **10**. 530. 1836 = A. Poiretii.
tinctum Moore, Gard. Chr. **1862**. 932. HB. 474 — Costa Rica-Peru.
— A. Wagneri Mett. 1869. HB. 473. A. decorum Moore 1869.
tomentellum Fée, Cr. vasc. Br. **1**. 37 t. 9 f. 2. 1869 = A. cristatum var.
tomentosum Kl. Linn. **18**. 553. 1844. NPfl. 284 — Guiana. Amazonas.
A. Klotzschianum Hk. 1851.
trapeziforme L. sp. **2**. 1097. 1753. HB. 118. (excl. γ). Chr. 134 — Amer. trop.
A. rhomboideum Schkuhr 1809 — A. pentadactylon Langsd. et Fisch. 1810.
trapeziforme Huds. Fl. angl. 391. 1762 = Asplenium marinum.
trapeziforme Forst. Prod. 84. 1786 = A. affine.

trapezoides Fée, Gen. 117. 1850-52 — Brasilia. Mexico. Jamaica.
A. cultratum J. Sm.; Hk. 1851 (non Willd. 1794); A. trapeziforme γ HB. 118.
tremulum Kze. (Hk. Bak. Syn. 114. 1867) = A. deflectens var.
triangulare Poir. Enc. Suppl. **1**. 140. 1810 = Lindsaya orbiculata var. tenera.
triangulatum Klf. Enum. 204. 1824 (an Gris. Fl. br. W. Ind. 664. 1864?) = ? A. intermedium.
tricholepis Fée, 8 mém. 72. 1857 — Mexico. New Mexico. Texas.
trichomanoides Poir. Enc. Suppl. **1**. 140. 1810 = Lindsaya cuneata.
trifidum Willd.; Fée, Gen. 114. 1850-52; Bolle, Bonpl. **3**. 121. 1855 = A. capillus [veneris.
trifoliatum L. sp. **2**. 1095. 1753 = Davallia.
trifoliatum Sessé et Mociño, Fl. Mex. 261. 1896 (2 spec.) — Mexico.
trigonum Labill. Nov. Holl. pl. spec. **2**. 99 t. 248 f. 2. 1806 = A. aethiopicum.
trilobum L. sp. **2**. 1095. 1753 = Davallia.
triphyllum Lam. Enc. **1**. 41. 1783 = Doryopteris.
triquetrum Pr.; Ettingsh. Farnkr. 81 t. 42 f. 9, 10. 1865 — »Amer. trop.«
truncatum Burm. Fl. Ind. 235 t. 66 f. 1. 1768 = Acacia decipiens.
truncatum Raddi, Opusc. sci. Bol. **3**. 295. 1819; Pl. Bras. **1**. 59 t. 78 f. 1. 1825 = A. subcordatum.
tuberosum Col. Tr. N. Zeal. Inst. **20**. 217. 1888 = A. diaphanum.
umbrosum Willd. sp. **5**. 447. 1810 = A. pulverulentum.
urophyllum Hk. sp. **2**. 24 t. 84 B. 1851. (HB. 118). NPfl. 284 — Ecuador.
varians Poir. Enc. Suppl. **1**. 143. 1810 = Cheilanthes tenuifolia.
varium H.B. Willd. sp. **5**. 435. 1810 = ? A. lucidum.
Veitchianum Moore, Gard. Chr. **1868**. 1090 = A. microsorum.
Veitchii Hance, Ann. sc. nat. IV. **15**. 229. 1861 = A. melanochlamys var.
velutinum Moore, Gard. Chr. **1866**. 777. HB. 472 — Columbia.
venustum Don, Prod. Fl. Nepal. 17. 1825. HB. 125. Chr. 139. NPfl. 284 — Afghanistan-India bor.
versaillense ht.; Garden and Forest **1**. 376. 1888 — Hort.
vestitum Spr. Anleit. **3**. 122. 1804 = Cheilanthes lanosa.
vestitum Wall. List n. 75. 1828 = A. caudatum.
Victoriae Moore, Gard. Chr. n. s. **17**. 428. 1882 — Hort.
villosissimum Mett.; Kuhn, Linn. **36**. 73. 1869. HB. 473 — Panama.
villosum L. Syst. Nat. ed. X. **2**. 1328. 1759. HB. 122. Chr. 138. NPfl. 283 — Ind. occ. Amer. austr. trop. — — A. falcatum Sw. 1801.
viride Vahl, Symb. **3**. 104. 1794 = Pellaea.
viridescens Col. Tr. N. Zeal. Inst. **27**. 400. 1895 — N. Zealand.
Visianii Schloss et Vukot, Fl. Croat. 1319. 1869 = A. capillus veneris.
Vogelii Mett.; Kuhn, Fil. Afr. 66. 1868 (syn.); Keys. Mém. Ac. St. Pétersb. **22**[3]. 8, 31. 1875 = A. tetraphyllum var.
Wagneri Mett.; Kuhn. Linn. **36**. 77. 1869 = A. tinctum.
Waltoni Moore, Gard. Chr. III. **1**. 111. 1887 — Hort.
Wattii Bak. Journ. Linn. Soc. **18**. 381 t. 14 f. A. 1881 = A. capillus veneris var.
Weigandi Moore, Gard. Chr. n. s. **20**. 748. 1883 — Hort.
Wercklearnum Christ, Bull. Boiss. II. **4**. 1093. 1904 — Costa Rica.
Wilesianum Hk. sp. **2**. 50 t. 83 C. 1851 — Mexico. ? Ind. occ.
A. crenatum Bak. 1867. HB. 120 (vix Willd. 1810).
Williamsii Moore, Gard. Chr. n. s. **10**. 45 f. 4. 1878; Bak. Ann. Bot. **5**. 207. [1891 — Peru.
Wilsoni Hk. sp. **2**. 6 t. 72 A. 1851 = A. dolosum*.
Zollingeri Mett.; Kuhn, Ann. Lugd. Bat. **4**. 280. 1869 — Java. Angola.

AGLAOMORPHA Schott, Gen. Fil. t. 20. 1834 = **Polypodium.**
Meyeniana Schott, Gen. t. 20. 1834.
AITHEROBOTRYS Wallich, Tidsskr. for Naturv. **1**. 259. 1822; Flora **1823**[1]. 67 (Aithenobotrys) (nomen) — ?
ALCICORNIUM Gaudichaud, Freyc. Voy. Bot. 48. 1826-27 = **Platycerium.**
vulgare Gaud. Freyc. Voy. Bot. 307. 1827 = P. alcicorne.
ALEURITOPTERIS Fée, Gen. Fil. 153. 1850-52.
affinis Fourn. Mex. pl. 1. 121. 1872 = Notholaena.
argentea Fée, Gen. 154. 1850-52 = Cheilanthes.
argyrophylla Fée, Gen. 154. 1850-52 = Cheilanthes farinosa.
candida Fée, Gen. 154. 1850-52 = Notholaena sulphurea*.
cretacea Fourn. Mex. pl. 1. 121. 1872 = Notholaena.
dealbata Fée, Gen. 154 t. 12 B f. 2. 1850-52 = Cheilanthes farinosa var.
farinosa Fée, Gen. 154 t. 12 B f. 1. 1850-52 = Cheilanthes.
Indica Fée, Gen. 154. 1850-52 = Cheilanthes farinosa.
lutea Fourn. Mex. pl. **1**. 121. 1872 = Cheilanthes aurantiaca.
Mexicana Fée, Gen. 154. 1850-52 = Cheilanthes farinosa.
Preslii Fourn. Mex. pl. **1**. 121. 1872 = Notholaena sulphurea.
pulveracea Fée, 8 mém. 117. 1857 = Notholaena sulphurea.
Schaffneri Fourn. Bull. Soc. Fr. **27**. 328. 1880 = Notholaena.
sulfurea Fée, Gen. 154. 1850-52 = Notholaena sulphurea.
ALLANTODIA R. Brown, Prod. Fl. N. Holl. 149. 1810 = **Athyrium.**
aemula Desv. Prod. 265. 1827 = Dryopteris aemula.
aspidioides Kze. Bot. Zeit. **1848**. 191 = A. scandicinum.
aspidioides de Vriese; Kze. Linn. **24**. 269 in obs. 1851 = Diplazium polypodioides.
asplenioides Kze. Linn. **9**. 72. 1834 = Diplazium aspl.
australis R. Br. Prod. Fl. N. Holl. 149. 1810 = A. umbrosum*.
axillaris Klf. Enum. 178. 1824 = A. filix femina*.
cordifolia Desv. Prod. 265. 1827 = Llavea c.
costalis Desv. Prod. 265. 1827.
(decurtata Kze. Linn. 24. 268. 1851.)
deflexa Kze. Bot. Zeit. **1848**. 191 = Diplazium Petersenii.
Fieldingiana Kze. Linn. 24. 268. 1851.
Hohenackeriana Kze. Farnkr. 2. 63 t. 126. 1850.
nitidula Kze. Bot. Zeit. 1848. 191.
oligantha Desv. Prod. 265. 1827 = Dryopteris filix mas *oligodonta.
paludosa Zippel; Kze. Bot. Zeit. **1848**. 191 = A. nigripes.
pinnata Blanco, Fl. Filip. ed. II. 571. 1845 — Ins. Philipp.
rupestris Klf.; Zoll. Syst. Verz. 42. 1854 (nomen) — Java.
? scabra Kze. Bot. Zeit. **1848**. 192 = Alsophila tristis.
scandicina Klf. Enum. 179. 1824.
Solenopteris Kze. Linn. 24. 266. 1851.
strigosa Bevis; Loud. Hort. Brit. Suppl. ed. nov. 485. 1850 — Madeira.
sylvatica Bl. Enum. 173. 1828 = A. silvaticum.
tenera R. Br. Prod. Fl. N. Holl. 149. 1810 = A. umbrosum.
umbrosa Klf. Enum. 179. 1810.
ALLANTODIA Wallich, Pl. asiat. rar. **1**. 44 t. 22. 1830 = **Diplaziopsis.**
Brunoniana Wall. List p. 63. 1829; Pl. as. rar. **1**. 44. t. 52. 1830 = D. javanica.
javanica Trevis. Nu. Giorn. bot. It. **7**. 159 (nota). 1875.
Alloesthes Moore, Ind. Fil. LXX. 1857 *(Nothochlaena §)* = Notholaena.

ALLOSORUS Bernhardi, Schrad. Neu. Journ. 1². 5, 36. 1806.
acclivis Kze. Farnkr. **2**. **6**. 1848 = Paesia.
acrostichoides Spr. Syst. **4**. 66. 1827 = Cryptogramma.
acutifolius Pr. Tent. 154. 1836 = Pteridium aquilinum.
adiantoides Pr. Tent. 153. 1836 = Pellaea viridis.
alabamensis O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Cheilanthes.
ambiguus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
andromedaefolius Klf.; Kze. Linn. **9**. 56. 1834 = Pellaea.
angulosus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
angustifolius Pr. Tent. 152. 1836 = Cheilanthes.
aquilinus Pr. Tent. 153. 1836 = Pteridium.
arachnoideus Pr. Tent. 153. 1836 = Pteridium aquilinum var.
argenteus Pr. Tent. 153. 1836 = Cheilanthes.
argyrophyllus Pr. Tent. 153. 1836 = Cheilanthes farinosa.
asper O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea scabra.
atropurpureus Kze.; Pr. Tent. 153. 1836 = Pellaea.
aurantiacus Pr. Tent. 153. 1836 = Cheilanthes.
auratus Pr. Tent. 152. 1836 = Onychium siliculosus.
auriculatus Pr. Tent. 153. 1836 = Pellaea.
Barklyae O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
bellus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea brachyptera var.
Boivini O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
brachypterus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
Breweri O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
Bridgesii Luerss. Flora **1876**. 290 = Pellaea.
Brunonianus J. Sm. JoB. **4**. 49. 1841 = Cryptogramma.
caespitosus Pr. Tent. 152. 1836 = Cheilanthes Belangeri.
caespitosus Kze. Linn. **23**. 218. 1850 = Cheilanthes marginata.
Calomelanos Pr. Tent. 153. 1836 = Pellaea hastata.
cambodiensis O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
capensis Pr. Tent. 152. 1836 = Onychium japonicum.
capensis Pappe et Raws. Syn. Fil. Afr. austr. 32. 1858 = Pteridium aquilinum*.
cartilagineus Pr. Tent. 153. 1836 = Doryopteris rigida.
caudatus Pr. Tent. 154. 1836 = Pteridium aquilinum*.
chaerophyllus Mart. et Gal. Mém. Ac. Brux. **15**. 47 t. 11. 1842 = Cheilanthes marginata.
ciliatus Pr. Rel. Haenk. **1**. 59. 1825 = Cheilanthes marginata.
concolor O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Doryopteris.
consobrinus Pappe et Raws. Syn. Fil. Afr. austr. 31. 1858 = Pellaea quadripinnata.
cordatus Pr. Tent. 153. 1836 = Pellaea.
cordifolius Keys. Pol. Cyath. Hb. Bung. 30. 1873 = Llavea.
cordifolius Pappe et Raws. Syn. Fil. Afr. austr. 31. 1858 = Pteridium aquilinum *capense.
crispatulus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Doryopteris.
crispus Bernh. (Schrad. Neu. Journ. 1². 36. 1806); Spr. Syst. **4**. 65. 1827 = Cryptogramma.
cuneatus Pr. Tent. 153. 1836 = Cheilanthes angustifolia var.
cuspidatus Hochst. in Jaub. et Spach, Ill. pl. orient. **3**. 3 t. 201. 1847 = Onychium melanolepis.
dealbatus Pr. Tent. 153. 1836 = Cheilanthes farinosa var.

decompositus Mart. et Gal. Mém. Ac. Brux. **15.** 48 t. 10 f. 2. 1842 = Cheilanthes angustifolia.
decompositus Kze; Mett. Fil. Lechl. **1.** 11. 1856 = Cheilanthes glauca.
deltodeus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Doryopteris deltoidea.
densus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
depilatus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
domingensis Pr. Tent. 153. 1836 = Adiantum deltoideum.
Donianus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
durus Pr. Tent. 153. 1836 = Pellaea.
esculentus Pr. Tent. 154. 1836 = Pteridium aquilinum*.
falcatus Kze. Linn. **23.** 219. 1850 = Pellaea.
farinosus Pr. Tent. 153. 1836 = Cheilanthes.
farinosus Kze. Farnkr. **2.** 5 t. 103 1848 = Ceropteris tartarea var.
flavescens O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
flexuosus Klf.; Kze. Linn. **13.** 136. 1839 = Pellaea.
formosus Liebm. Vid. Selsk. Skr. V. **1.** 220 (seors. 68). 1849 = Pellaea pul-
Fournieri O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Doryopteris. [chella.
foveolatus Rupr. Distr. Cr. vasc. Ross. 46. 1845 = Cryptogramma acrostichoides.
fumariaefolius O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Cryptogramma crispa
glaucus J. Sm. Hist. Fil. 167. 1875 = Cheilanthes. [var.
gracilis Pr. Tent. 153. 1836 = Cryptogramma Stelleri.
Greggii O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
hastatus Pr. Tent. 153. 1836 = Pellaea viridis.
? heterophyllus Bernh. (Schrad. Neu. Journ. **1^2.** 36. 1806) = Anopteris hexagona.
heterophyllus Pr. Tent. 152. 1836 = Doryopteris pilosa.
hirsutus Pr. Rel. Haenk. **1.** 59 t. 10 f. 1. 1825 = Cheilanthes glauca.
hottentottus Pr. Tent. 154. 1836 = Pteridium aquilinum *capense.
imbricatus Pr. Tent. 153. 1836 = Jamesonia.
integricuspis O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
intramarginalis Pr. Tent. 153. 1836 = Pellaea.
? involutus Pr. Tent. 153. 1836 = Pellaea.
Karwinskii Kze. Linn. **13.** 138. 1839 = Llavea cordifolia.
Kitchingii O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
lancifolius O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
lanuginosus Pr. Tent. 154. 1836 = Pteridium aquilinum *capense.
leucomelas O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
lomariaceus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Doryopteris.
lorigerus Pr. Tent. 154. 1836 = Pteridium aquilinum.
marginatus J. Sm. Bot. Voy. Herald 341. 1854 = Cheilanthes.
melanolepis Dcne. Arch. Mus. **2.** 189. 1841 = Onychium.
microphyllus Liebm. Vid. Selsk. Skr. V. **1.** 219 (seors. 67). 1849 = Cheilanthes.
microphyllus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Doryopteris.
minutus Turcz.; Trautv. Pl. Imag. Fl. Ross. 9 t. 3. 1844 = Cryptogramma
mucronatus Eat. Am. Journ. Sci. II. **22.** 138. 1856 = Pellaea. [Stelleri.
myrtifolius O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea myrtillifolia.
namaquensis O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
nitidulus Pr. Tent. 152. 1806 = Pellaea.
nitidus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.
ochraceus Hk. in Benth. Pl. Hartw. 55. 1840 = Cheilanthes aurantiaca.
ornithopus O. Ktze. Rev. Gen. Pl. **2.** 806. 1891 = Pellaea.

pallidus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
paradoxus Kze. Linn. **23**. 219. 1850 = Pellaea.
Pearcei O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
pectiniformis O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea Goudotii.
pilosus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Doryopteris.
Pringlei O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
psittacinus Pr. Tent. 153. 1836 = Pteridium aquilinum var.
pulchellus Pr. Tent. 152. 1836 = Cheilanthes.
pulchellus Mart. et Gal. Mém. Ac. Brux. **15**. 47 t. 10 f. 1. 1842 = Pellaea.
pulveraceus Pr. Tent. 153. 1836 = Notholaena sulphurea.
pusillus Bernh. Schrad. Neu. Journ. 1^2. 36. 1806 = Cheilanthes pteridioides.
quadripinnatus Pr. Tent. 154. 1836 = Pellaea.
recurvatus Pr. Tent. 154. 1836 = Pteridium aquilinum.
Riedelii O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
rigidus Kze. Linn. **9**. 55. 1834 = Cheilanthes Poeppigiana.
rigidus Pr. Tent. 153. 1836 = Doryopteris.
robustus Kze. Linn. **10**. 502. 1836 = Doryopteris.
rotundifolius Kze. Linn. **23**. 219. 1850 = Pellaea.
?rugulosus Guill. Ann. sc. nat. II. **6**. 314. 1836 = Paesia rugosula.
sagittatus Pr. Tent. 153. 1836 = Pellaea cordata var.
scaberulus Pr. Tent. 154. 1836 = Paesia.
Seemanni O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
sitchensis Rupr. Distr. Cr. vasc. Ross. 47. 1845 = Cryptogramma.
Skinneri O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Doryopteris.
Stelleri Rupr. Distr. Cr. vasc. Ross. 47. 1845 = Cryptogramma.
subverticillatus Pr. Tent. 153. 1836 = Pellaea ternifolia.
sulphureus Pr. Tent. 153. 1836 = Notholaena.
Tamburii O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Cheilanthes argentea.
tauricus Pr. Tent. 154. 1836 = Pteridium aquilinum.
ternifolius Kze.; Kl. Linn. **20**. 339. 1847 = Pellaea.
tripinnatus O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
villosus Pr. Tent. 154. 1836 = Pteridium aquilinum.
viridis O. Ktze. Rev. Gen. Pl. **2**. 806. 1891 = Pellaea.
Zippelii Miq. Ann. Lugd. Bat. **4**. 98. 1868-69 = Pellaea.

ALLOSURUS Endlicher, Genera plant. 59. 1836 et auctt. = Allosorus.

Allothecium Moore, Ind. Fil. LXXXVIII. 1857 *(Pleopeltis §)* = Polypodium.

ALSOPHILA R. Brown, Prod. Fl. N. Holl. 158. 1810 — (Cyatheaceae Gen. **7**).
aculeata J. Sm. Lond. JoB. **1**. 667. 1842 = A. microdonta.
aculeata Kl. Linn. **18**. 540. 1844: **20**. 442. 1847 = A. armata.
aculeata Hk. sp. **1**. 49. 1844 = A. aspera.
acuminata J. Sm. Lond. JoB. **1**. 667. 1842 = A. Miersii.
acuta Pr. Abh. böhm. Ges. V. **5**. 343. 1848 = A. glauca.
adspersa Klf.; Kze. Bot. Zeit. **1844**. 314 — Brasilia.
aethiopica Welw.; Hk. Syn. 44. 1846. NPfl. 135 (althiopica) — Afr. occ. trop.
affinis Fée, Gen. 346. 1850-52 = A. quadripinnata. [Ins. Comorae.
alata Kze.; Fée, Gen. 346. 1850-52 (nomen) = A. phalerata.
alata Fourn. Ann. sc. nat. V. **18**. 349. 1873. NPfl. 138 — N. Caledonia.
alata Sod. Rec. 19. 1883 = A. Sodiroi.
albosetacea Bedd. Ferns. br. Ind. Suppl. 2. 1876. NPfl. 136 — Ins. Nicobar.
alternans Hk. Syn. 41. 1866 = Cyathea.

althiopica Diels, Nat. Pfl. 1⁴. 135. 1899 (err. typ.) = A. aethiopica.
alutacea Kze. Bot. Zeit. **1844**. 327 = A. phalerata var.
amazonica Linden, Cat. 1871 — (Brasilia) Hort.
Andersonii Scott; Bedd. Ferns br. Ind. t. 310. 1869. HB. 459. NPfl. 136 — [Sikkim.
aperta Fée, Cr. vasc. Br. **1**. 158 t. 54 f. 2. 1869 — Brasilia.
aquilina Christ, Engl. Jahrb. **24**. 83. 1897 — Cuba. Brasilia.
arbuscula Pr. Tent. 62. 1836; Bak. Fl. bras. 1². 322. 1870. (HB. 33). NPfl. 134 — [Brasilia.
armata Mart. Ic. Cr. Bras. 72 t. 28, 48. 1834 = A. microdonta.
armata (Sw.) Pr. Tent. 62. 1836. HB. 25. Chr. 325. NPfl. 134 — Amer. trop.
Polypodium Sw. Prod. 134. 1788; Alsophila Swartziana Mart. 1834. Cyathea aculeata Willd.; Klf. 1824; Disphenia Pr. 1836; Alsophila Kl. 1844; Hemitelia Fée 1850-52. Alsophila hirta Klf. 1824.
? armigera Kze. Linn. **9**. 98. 1834 = A. procera var.
articulata J. Sm.; Moore et Houlst. Gard. Mag. Bot. **3**. 331 f. 81. 1852 = A. aspera.
aspera (L.) (R. Br. Prod. Fl. N. Holl. 158. 1810) Spr. Syst. **4**. 124. 1827. HB. 33. Chr. 325. NPfl. 134 — Ind. occ. Costa Rica.
Polypodium L. sp. **2**. 1093. 1753; Cyathea Sw. 1801; Disphenia Pr. 1836; Hemitelia Fée 1850-52 — Gymnosphaera aculeata J. Sm. 1842; Alsophila Hk. 1844; A. echinata Moore 1857 — A. Eatoni Jenm. 1887.
aterrima Hk. Syn. 38. 1866. NPfl. 134 — Peru.
atrovirens (Langsd. et Fisch.) Pr. Tent. 61. 1836. HB. 34. NPfl. 134 — Brasilia austr. — Polypodium Langsd. et Fisch. Ic. Fil. 12 t. 14. 1810. Alsophila Hookeriana Kl.; Hk. 1844.
atrovirens ht. Sander, Rev. Hort. **1893**. 200 — Hort. — (an praec.?)
aurea Fée, 8 mém. 109. 1857 = A. myosuroides.
australis R.Br. Prod. Fl. N. Holl. 158. 1810. HB. 40. Chr. 328. NPfl. 138 — [Australia. Tasmania.
axillaris Moore, Ind. 48. 1857 = A. leucolepis.
Bakeri Zeiller, Bull. Soc. Fr. **32**. 72. 1885 = A. Kingi.
Bakeri Sod. Cr. vasc. quit. 532. 1893 = A. quitensis.
Baroni Bak. Journ. Linn. Soc. **21**. 455. 1885. NPfl. 135 — Madagascar.
baroumba L. Linden, La Sem. Hort. **4**. 435. 1900 — Congo.
Batesii Bak. Kew Bull. **1897**. 299 — Kamerun.
batjanensis Christ; Diels, Nat. Pfl. 1⁴. 138. 1899; Christ in Warb. Monsunia **1**. 90. 1900 — Ins. Batjan.
Beyrichiana J. Sm. Hist. Fil. 249. 1875 = Hemitelia setosa.
bicrenata (Liebm.) Fourn. Mex. pl. **1**. 134. 1872. NPfl. 134 — Mexico.
Cyathea Liebm. Vid. Selsk. Skr. V. **1**. 289 (seors. 137). 1849.
bipinnatifida Bak. Syn. 456. 1874. NPfl. 132 — Guiana.
Blancheti Trevis. Atti Ist. Veneto II. **2**. 165. 1851 } = A. phalerata.
Blanchetiana Pr. Epim. 28. 1849 }
blechnoides (Rich.) Hk. sp. **1**. 35. 1844. HB. 32. Chr. 324. NPfl. 132 — Amer. trop.
Polypodium Rich. 1791 (nomen); Sw. 1806 (nomen); Amphidesmium Kl. 1847; Polypodium rostratum H.B. Willd. sp. **5**. 193. 1810 (non Burm. 1768 nec Cav. 1802); Aspidium HBK. 1815; Alsophila Mart. 1834; Metaxya Pr. 1836; Amphidesmium J. Sm. 1842 — Polypodium Parkeri Hk. et Grev. 1831; Amphidesmium Schott 1834; Metaxya J. Sm. 1842.
Boivini Mett.; Ettingsh. Farnkr. 220 t. 155 f. 2-3. 1865; **Kuhn**, Fil. Afr. 161. 1868. HB. 460. Chr. 328. NPfl. 135 — Comorae. Madagascar.
Bongardiana Mett.; Bak. Syn. 41. 1874 — Ins. Bonin.

Boninsimensis Christ in Warb. Monsunia **1**. 90. 1900 = Hemitelia.
brevis J. Sm. Lond. JoB. **1**. 667. 1842 (nomen) — Brasilia.
Brunoniana Wall. List n. 7073. 1829 = Cyathea.
Brunoniana Bedd. Ferns br. Ind. t. 66. 1866 = A. glauca.
bullata Bak. Journ. Linn. Soc. **15**. 412. 1876. NPfl. 135 — Madagascar.
Burbidgei Bak. JoB. **1879**. 38. NPfl. 136 — Borneo.
camerunensis Diels, Nat. Pfl. 1^4. 135. 1899 — Kamerun.
capensis J. Sm. Lond. JoB. **1**. 666. 1842 = Hemitelia.
caracasana Kl. Linn. **18**. 541. 1844. HB. 457. NPfl. 134 — Amer. austr. septentr.
castanea Bak. JoB. **1891**. 3. NPfl. 136 — Madagascar.
caudata J. Sm. JoB. **3**. 419. 1841; Hk. sp. **1**. 52 t. 20B. 1844. HB. 42. NPfl. 137 — Luzon. — Hemitelia Mett. 1859.
? **celebica** (Bl.) Mett. Ann. Lugd. Bat. **1**. 53. 1863. HB. 41 — Celebes. Ternate. Cyathea? Bl. Enum. 245. 1828.
Ceropteris Fée, Cr. vasc. Br. **1**. 167 t. 57 f. 1. 1869 = A. arbuscula.
chimborazensis Hk. Syn. 37. 1866. NPfl. 134 — Ecuador.
chnoodes Christ, Bull. Boiss. II. **4**. 958. 1904 — Costa Rica.
Colensoi Hk.fil. Fl. N. Zeal. **2**. 8 t. 73. 1854. HB. 40. NPfl. 138 — N. Zealand.
commutata Mett. Ann. Lugd. Bat. **1**. 53. 1863. HB. 53. NPfl. 136 — Malacca. Gymnosphaera squamulata J. Sm. 1841 et in Hk. Gen. t. 100. 1842 (non Bl. 1828); Alsophila Hk. 1844 (pt.).
comosa Wall. List p. 64 (n. 319). 1829; Hk. sp. **1**. 53 t. 20 A. 1844 = A. squamulata.
comosa Scott, Tr. Linn. Soc. **30**. 36 t. 16 B. 1874 = A. ornata.
compta Mart. Ic. Cr. Bras. **1**. 66 t. 41. 1834. HB. 33. NPfl. 134 — Mexico -Ecuador-Brasilia. — Cyathea Mart. Denkschr. bot. Ges. Regensb. **2**. 146 t. 2 f. 1, 2. 1822.
concinna Bak. Syn. 459. 1874 — Louisiade Archip.
conjugata Spruce; Hk. Syn. 37. 1866. NPfl. 134 — Ecuador.
contaminans Wall. List pag. 64. 1829 = A. glauca.
contracta Fée, Cr. vasc. Br. **1**. 167 t. 59 f. 2. 1869 — Brasilia.
Cooperi F. Muell. Fragm. **5**. 117. 1866. HB. 459. NPfl. 138 — Australia.
corcovadensis (Raddi) C. Chr. Ind. 41. 1905 — Brasilia. Polypodium Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 26. t. 40. 1825; P. Taenitis Roth 1821; Alsophila Kze. 1834. HB. 32. Chr. 324. NPfl. 133: Trichipteris excelsa Pr. 1822; Alsophila Mart. 1834 (non infra); Chnoophora Mart. 1834.
corcovadensis Fée, Cr. vasc. Br. **1**. 163 t. 56 f. 2. 1869 = A. Glaziovii.
cordata Kl. Bot. Zeit. **1846**. 104; Linn. **20**. 441. 1847 — Columbia.
costalis Christ. Bull. Boiss. II. **4**. 951. 1904 — Costa Rica.
crassa Karst. Fl. Col. **2**. 187 t. 199. 1869 = A. phalerata var.
crenata Kze. Bot. Zeit. **1844**. 312 = A. elegans var.
crenulata Mett.; Hk. Syn. 44. 1865 = Hemitelia.
crinita Hk. Ic. pl. t. 671. 1844. HB. 42. Chr. 327. NPfl. 136 — India austr. Ceylon. Java.
dealbata Pr. Abh. böhm. Ges. V. **5**. 343. 1848 = A. glauca.
debilis de Vriese; Jungh. Java ed. germ. 310, 476. 1852 — Java. (= ? A. latebrosa).
decipiens Scott; Bedd. Ferns br. Ind. t. 311. 1869 = Cyathea.
decipiens Fée. Cr. vasc. Br. **2**. 81 t. 103 f. 1. 1872-73 — Brasilia.
Deckeriana Kl.; Kze. Linn. **23**. 220. 408. 1850 = A. quadripinnata.

decomposita Karst. Fl. Col. **2**. 185 t. 198. 1869 — Columbia.
A. oligocarpa Fée 1850-52 (nomen). HB. 39. 1874. NPfl. 135.
decurrens Hk. sp. **1**. 51. 1844. HB. 39. NPfl. 138 — Polynesia.
decussata Christ, Prim. Fl. Costar. **3**. 41. 1901 — Costa Rica.
denticulata Bak. JoB. **1885**. 102 — Formosa.
denticulata ht. Lind. et ht. Veitch — Hort.
dichromatolepis Fée, Cr. vasc. Br. **1**. 164 t. 57 f. 2. 1869 — Brasilia.
dimorpha Christ, Ann. Jard. Buit. II. **4**. 41. 1904 — Celebes.
dissitifolia Bak. JoB. **1886**. 182. NPfl. 138 — Fiji.
Dombei Desv. Prod. 320. 1827 (rectius Dombeyi) — Peru.
dorsalis (Fée) Christ, Bull. Boiss. II. **2**. 648. 1902 — Brasilia. — Lophosoria Fée, Cr. vasc. Br. **1**. 173 t. 51. f. 3. 1869. (t. Bak. = Hemitelia setosa).
dubia Bedd. JoB. **1888**. 1. t. 279 A. NPfl. 136 — Malacca. Borneo.
Eatoni Jenm. JoB. **1887**. 98 = A. aspera var. aculeata.
echinata Moore, Syn. CV. 1857 = A. aspera var. aculeata.
elegans Mart. Ic. Cr. Bras. 63 t. 38. 1834. HB. 32. Chr. 324. NPfl. 133 — Brasilia — Trichopteris Pr. 1836 — Alsophila crenata Kze. 1844; Trichopteris [Fée 1869.
elegantissima Linden; (Salom. Nom. 28. 1883) — Hort.
Elliottii Bak. Ann. Bot. **6**. 96. 1892 — Grenada. (Ind. occ.).
elongata Hk. sp. **1**. 43. 1844. HB. 34. NPfl. 134 — Amer. centr.-Columbia. Rio Janeiro.
A. Tumacensis J. Sm. 1842 (nomen) — A. Tijucensis Fée 1869.
eriocarpa Fée, Cr. vasc. Br. **1**. 162 t. 66 f. 1. 1869 = A. paleolata.
erubescens Kze. Bot. Zeit. **1844**. 344 = Dryopteris crinita.
excelsa R. Br. Prod. Fl. N. Holl. 158. 1810 (nomen); Endl. Prod. Fl. Norf. 16. 1833. HB. 40. NPfl. 138 — Ins. Norfolk. Queensland.
excelsa Mart. Ic. Cr. Bras. 63 t. 27, 37. 1834 = A. corcovadensis.
extensa (Forst.) (R. Br. Prod. Fl. N. Holl. 158. 1810) Spr. Syst. **4**. 124. 1827 — Polynesia. N. Guinea. Celebes.
Polypodium Forst. Prod. 83. 1786 (n. 453); Cyathea Sw. 1801; Hemitelia Pr. 1848 — Polypodium lunulatum Forst. Prod. 83. 1786 (n. 456); Alsophila (R. Br. 1810) Spr. 1827. HB. 41. NPfl. 137; Phegopteris Fée 1850-52 — Alsophila vitiensis Carr. 1873.
extensa Desv. Prod. 319. 1827 = Cyathea medullaris.
extensa Bl. Enum. 246. 1828 = Hemitelia crenulata.
falcata Mett.; Kuhn, Linn. **36**. 155. 1869. HB. 458. NPfl. 134 — Panama (Ins. Gorgona).
farinosa Karst.: Mett. Ann. sc. nat. V. **2**. 262. 1864 (nomen); Karst. Fl. Col. **2**. 163 bis t. 186. 1869 — Columbia.
Fauriei Christ, Bull. Boiss. II. **1**. 1019. 1901 — Liu Kiu.
Feeana C. Chr. Ind. 42. 1905 — Brasilia.
A. Glaziovii Bak. Fl. bras. 1^2. 592. 1870. HB. 456. NPfl. 133 (non Fée 1869); Trichopteris elegans Fée 1872-73 (non Pr. 1836).
ferox Pr. Tent. 62. 1836 = A. microdonta.
? *Finlaysonianum* Wall.; Moore, Ind. 50. 1857 (nomen) — India.
Polypodium Wall. List n. 2221. 1829 (nomen).
Fischeriana Regel, Ind. sem. ht. Petr. 1855; Linn. **28**. 366. 1856 = Dryopteris grandis *macroptera.
flexuosa Fée, Cr. vasc. Br. **1**. 159. 1869 — Brasilia.
floribunda Hk.; Bak. Syn. 458. 1874. NPfl. 134 — Peru.

formosana Bak. Ann. Bot. **5**. 190. 1891. NPfl. 136 — Formosa.
fragilis Zoll. et Moritz, Nat. Gen. Arch. **1**. 400. 1844 = Dryopteris lineata.
fragrans »J. Sm.«; Salom. Nom. 28. 1883 — Ins. Philipp.
frigida Karst. Fl. Col. **1**. 61. t. 30. 1859. HB. 38. NPfl. 135 — Columbia.
fuliginosa Christ, Bull. Boiss. **6**. 138. 1898 — Luzon.
fulva Mart. et Gal. Mém. Ac. Brux. **15**. 78 t. 23. 1842 = Cyathea schanschin.
fumata Kl. (Hk. sp. **1**. 42. 1842. syn.); Sal. Nom. 28. 1883 = A. phalerata.
furcata Christ, Bull. Bois. II. **4**. 957. 1904 — Costa Rica.
Fusagasuga Trevis. Atti Ist. Veneto II. **2**. 165. 1851 = A. armata.
Gardneri Hk. sp. **1**. 40. 1844 = A. paleolata.
gazellae Kuhn, Forschr. Gazelle **4** Farne 13. 1889 — N. Hannover.
gibbosa Kl. Linn. **18**. 542. 1844. HB. 457. NPfl. 134 — Guiana. Peru. Puerto Rico.
gigantea Wall. List pag. 64 (n. 321). 1829 = A. glabra.
glabra (Bl.) Hk. sp. **1**. 51. 1844. HB. 43. Chr. 326. NPfl. 136 — Asia trop.
Gymnosphaera Bl. Enum. 242. 1828 — Polypodium giganteum Wall. 1828 (non Desv. 1827); Alsophila Wall. 1829; Gymnosphaera Hk. 1842; Dichorexia Pr. 1849 — Alsophila Metteniana Hance 1868.
glauca (Bl.) J. Sm. JoB. **3**. 419. 1841 — Asia trop..
Chnoophora Bl. Enum. 243. 1828. Polypodium contaminans Wall. 1828; Alsophila Wall. 1829. HB. 41. Chr. 327. NPfl. 136.
glauca Goldm. Nova Acta **19** Suppl. I. 465. 1843. (an praec.? — Manila.
glauca Urban, Symb. Ant. **4**. 12. 1903 = A. quadripinnata.
glauca Hort.; Mett. Fil. Lips. 109. 1856 — Hort.
glaucescens Wall. List n. 7074. 1829; Hk. sp. **1**. 55. 1844 = A. glauca.
Glaziovii Fée, Cr. vasc. Br. **1**. 160 t. 55. f. 2. 1869 — Brasilia.
— A. corcovadensis Fée 1869 (non supra).
Glaziovii Bak. Fl. bras. **1**². 592. 1870 = A. Feeana.
glumacea Fée, Cr. vasc. Br. **1**. 170 t. 61 f. 2. 1869 = A. leucolepis.
Godmani Hk. Syn. 36. 1866. NPfl. 134 — Guatemala.
goyazensis Christ in Schwacke, Pl. Nov. Mineiras **2**. 33. 1900: Bull. Boiss. II. **2**. 646. 1902 — Brasilia centr.
gracilis Und. et Maxon, Bull. Torr. Cl. **29**. 577. 1902 — Cuba.
Grevilleana Wall. List n. 7075. 1829 = Microlepia speluncae.
guianensis Regel, Ann. sc. nat. IV. **12**. 372. 1859: Gartenfl. **6**. 119 t. 288. 1860 — Hort. (Guiana).
guimaraensis Fée et Glaz.: Fée, Cr. vasc. Br. **2**. 81 t. 103 f. 2. 1872-73 — Brasilia.
Haenkei Pr. Rel. Haenk. **1**. 68. 1825 = A. extensa.
Helferiana Pr. Abh. böhm. Ges. V. **5**. 341. 1848 = A. glabra.
Henryi Bak. Kew Bull. **1898**. 229 — Yunnan.
hirsuta Kze. Linn. **9**. 98. 1834 = A. armata.
hirsutissima Fée, 11 mém. 95. 1886 — Guadeloupe.
hirta Klf. Enum. 249. 1824 = A. armata.
hispida Bak. JoB. **1881**. 202. NPfl. 135 — Columbia.
Holstii Hier. in Engl. Pflw. Ostafr. C. 87. 1895. NPfl. 135 — Afr. orient.
Hookeriana Kl.: Hk. sp. **1**. 39. 1844 = A. atrovirens.
Hornei Bak. JoB. **1879**. 293 — Fiji.
Hostmanni J. Sm. Bot. Mag. **72**. Comp. 37. 1846 = Hemitelia multiflora.
Humboldtii Kl.; Kze. Linn. **23**. 220. 1850 = A. villosa.
humilis J. Sm. Lond. JoB. **1**. 667. 1842 = A. villosa.
hypolampra Kze.: Kuhn, Linn. **36**. 158. 1869 — Merida. ? Columbia.

impressa Fée, Cr. vasc. Br. **1**. 166 t. 58 f. 1. 1869 — Brasilia.
infesta Kze. Linn. **9**. 98. 1834 = A. phalerata.
intermedia Mett. Ann. sc. nat. IV. **15**. 83. 1861. (HB. 41) — N. Caledonia. [N. Guinea.
Junghuhniana Kze. Bot. Zeit. **1848**. 284 = Hemitelia.
Kalbreyeri C. Chr. Ind. 44. 1905 — Columbia.
A. podophylla Bak. JoB. **1881**. 202 (non Hk. 1857).
khasyana Moore; Kuhn. Linn. **36**. 154. 1869. NPfl. 136 — Khasia. (prob. = A. latebrosa var.).
Kingi Clarke: Bedd. Hdb. 475. 1883. NPfl. 136 — Perak. — A. Bakeri Zeiller [1885.
laeta Kze. Bot. Zeit. **1846**. 476 = A. squamulata.
laevis J. Sm. Lond. JoB. **1**. 666. 1842 = Hemitelia multiflora.
lanuginosa Pr. Epim. 29. 1849 = A. tomentosa.
lasiosora Mett.: Kuhn, Linn. **36**. 157. 1869 — Peru. — A. infesta β HB. 457.
latebrosa Wall.; Pr. Tent. 62. 1836 (nomen); Hk. sp. **1**. 37. 1844. HB. 43. Chr. 327. NPfl. 136 — India bor.-Malesia.
Polypodium Wall. 1828 (nomen); Dichorexia Pr. 1848; Hemitelia Mett. 1856.
latebrosa var. hemitelioides Scott, Tr. Linn. Soc. **30**. 34. 1874 = Cyathea [Brunoniana.
latevagans Bak. JoB. **1881**. 203. NPfl. 133 — Columbia.
Lechleri Mett. Fil. Lechl. **2**. 28. 1859 — Peru.
Leichhardtiana F. Muell. Fragm. **5**. 53. 1865. HB. 40. NPfl. 138 — Australia.
A. Macarthurii Hk. 1866. A Moorei J. Sm. 1866.
lepidophora Kze. Bot. Zeit. **1844**. 345. (corr. nom.) = A. lepifera.
lepidotricha (Fourn.) Diels, Nat. Pfl. 1^4. 138. 1899 — N. Caledonia.
Fourniera Fourn. Ann. sc. nat. V. **18**. 349. 1873.
lepifera J. Sm. JoB. **3**. 419. 1841; Hk. sp. **1**. 54. 1844 = A. tomentosa.
?Leprieuriana Kze. Linn. **21**. 235. 1848 = Hemitelia multiflora.
leptocladia Fée, Cr. vasc. Br. **1**. 161 t. 55 f. 1. 1869 — Brasilia.
Leschenaultiana (Wall.) Moore, Ind. 52. 1857 — India austr.
Polypodium Wall. List n. 323. 1828 (nomen).
leucolepis Mart. Ic. Cr. Bras. 70 t. 46. 1834. HB. 34. NPfl. 134 — Brasilia. Costa Rica.
Polypodium axillare Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 27 t. 41. 1825 (non Ait. 1789); Phegopteris Fée 1850-52; Alsophila Moore 1857 – A. glumacea, Ludoviciana, nigrescens, pectinata Fée 1869.
Loddigesii Kze. Linn. **20**. 7. 1847. HB. 458. NPfl. 138 — Queensland. N. S. Wales.
Loubetiana L. Linden, La Sem. Hort. **4**. 435, 453 f. 152. 1900 — Congo.
Ludoviciana Fée, Cr. vasc. Br. **1**. 169 t. 60 f. 2. 1869 = A. leucolepis.
lunulata R. Br. (Prod. Fl. N. Holl. 158. 1810) Spr. Syst. **4**. 124. 1827 = A. extensa.
lunulata Bl. Enum. 246. 1828 = Hemitelia Junghuhniana.
lurida (Bl.) Hk. sp. **1**. 55. 1844. HB. 43 — Java. Celebes.
Chnoophora Bl. Enum. 244. 1828.
Macarthurii Hk. Syn. ed. I. 40. 1866 = A. Leichhardtiana.
Macgillivrayi Bak. Syn. 458. 1874 — Louisiade Archip.
macrosora Bak. Tr. Linn. Soc. II. Bot. **2**. 288. 1887. NPfl. 134 — Guiana.
Hemitelia Jenm. 1898.
manilensis Pr. Tent. 62. 1836 = Hemitelia.
marginalis Kl. Linn. **18**. 542. 1844. HB. 32. NPfl. 133 — Guiana.
Hemitelia Jenm. 1898; Trichopteris J. Sm. msc.
marianna Gaud. Freyc. Voy. Bot. 365. 1827 = A. extensa.
Marshalliana L. Linden, Gard. Chr. III. **15**. 663. 1894 — Hort. (Brasilia?)
martinicensis Spr. Neue Entd. **3**. 7. 1822 = Dryopteris subincisa*.

melanopus Hassk. JoB. **7**. 325. 1855; Obs. Bot. Fil. **1**. 42. 1856 — Java.
melanopus Hk. Syn. 37. 1866 = A. nigripes.
Mertensiana Kze. Bot. Zeit. **1848**. 586 — Peel Ins. Ins. Bonin-Sima.
Hemitelia Pr. 1849.
Mertensii Trevis. Atti Ist. Veneto II. **2**. 165. 1851 = A. glauca.
Metteniana Hance, JoB. **1868**. 175 = A. glabra.
mexicana Mart. Ic. Cr. Bras. 70 t. 45. 1834. HB. 38. NPfl. 134 — Mexico.
microdonta Desv. Prod. 319. 1827 — Amer. trop.
Polypodium Desv. Berl. Mag. **5**. 319. 1811. P. aculeatum Raddi 1819 (non L. 1753); Alsophila J. Sm. 1842. Cyathea ferox Pr. 1822; Alsophila Pr. 1836. HB. 34. Chr. 325. NPfl. 134. A. armata Mart. 1834.
microphylla Kl. Linn. **18**. 541. 1844. HB. 36. NPfl. 134 — Venezuela. — A. squamata Kl. 1844.
microptera »Bak.« (an hort?); Salom. Nom. 29. 1883 — Venezuela.
Miersii Hk. sp. **1**. 38. 1844. HB. 32 — Brasilia. Venezuela.
A. acuminata J. Sm. 1842 (nomen).
millefolium Desv. Prod. 320. 1827 = A. quadripinnata.
mindanensis Christ; Diels, Nat. Pfl. **1**[4]. 136. 1899; Christ in Warb. Monsunia **1**. 90. 1900 — Mindanao.
Miquelii Kze. Linn. **23**. 221. 1850 — Hort. (Java. Surinam?).
modesta Bak. JoB. **1880**. 210. NPfl. 136 — Sumatra.
mollissima Kze. Bot. Zeit. **1844**. 328 = A. villosa.
mollissima Moore, Ind. 54. 1857 = A. armata.
monosticha Christ, Bull. Soc. bot. Belg. **33**[2]. 94. 1894 — Guiana.
monticola Mart. Ic. Cr. Bras. 75. 1834 = A. quadripinnata var.
Moorei J. Sm. Ferns br. and for. 245. 1866 = A. Leichhardtiana.
Moorei J. Sm. Hist. Fil. 249. 1875 = Hemitelia.
mucronata Christ, Bull. Soc. bot. Belg. **35**. 178. 1896 — Costa Rica.
multiflora Pr. Tent. 61. 1836 = Hemitelia.
munita Klf.; Pr. Tent. 62. 1836 = A. paleolata.
muricata Desv. Prod. 319. 1827 = Cyathea.
myosuroides Liebm. Vid. Selsk. Skr. V. **1**. 286 (seors. 134). 1849. HB. 38. NPfl. 134 — Mexico. — A. aurea Fée 1857.
Naumanni Kuhn, Forschr. Gazelle **4** Farne 13. 1889 — N. Pommern.
nigra Mart. Ic. Cr. Bras. 71 t. 30 f. 5 t. 47. 1834. HB. 35. NPfl. 134 — Brasilia.
nigra Jenm. W. Ind. and Guiana Ferns 38. 1898 — Trinidad.
nigrescens Fée, Cr. vasc. Br. **1**. 170 t. 54 f. 1. 1869 = A. leucolepis.
nigripes C. Chr. Ind. 45. 1905 — Ecuador.
A. melanopus Hk. Syn. 37. 1866. NPfl. 134 (non Hassk. 1855).
nitens J. Sm. Lond. JoB. **1**. 667. 1842 (nomen); Gris. Fl. br. W. Ind. 705. 1864 — Ind. occ. Brasilia.
A. nitida Kze. Ettingsh. 1865. A. aspera pt. HB. 33.
nitida Kze.; Ettingsh. Farnkr. 222 t. 154 f. 4, 8 t. 155 f. 1, 7. 1865 = A. nitens.
novae caledoniae Mett. Ann. sc. nat. IV. **15**. 82. 1861. HB. 39. Chr. 329. NPfl. 138 — N. Caledonia. — Fourniera Bomm.; Fourn. 1873.
oblonga Kl. Linn. **18**. 540. 1844. HB. 458. NPfl. 134 — Guiana.
obscura Scort.; Bedd. JoB. **1887**. 321 t. 278 f. 2. NPfl. 136 — Perak.
obtusa Kl. Allg. Gartenzeit. **1852**. 41; Bot. Zeit. **1854**. 439 — Venezuela. Columbia.
obtusiloba Hk. Syn. 45. 1866. NPfl. 135 — Afr. occ. trop.
Hemitelia Mett.; Salomon 1883.

Oldhami Bedd. Ferns br. Ind. t. 343. 1870 = A. ornata.
oligocarpa Fée, Gen. 346. 1850-52 = A. decomposita.
oligosora Miq.; Kze. Linn. **23**. 221. 1850; Miq. Verh. Nederl. Inst. III. **4**. 55. 1851 = A. squamulata.
ornata Scott; Bedd. Ferns br. Ind. t. 342. 1870. HB. 460. NPfl. 136 — Sikkim.
A. Oldhami Bedd. 1870 — A. Scottiana Bak. 1872. HB. 460 — A. comosa Scott 1874 — A. sikkimensis Clarke et Bak. 1888.
paleolata Mart. Ic. Cr. Bras. 68 t. 43. 1834. HB. 33. NPfl. 134 — Brasilia. Columbia.
Polypodium alsophilum Link 1833 (an nomen opt.?). Alsophila Gardneri Hk. 1844 — ? Cyathea Sellowiana Pr. 1836; Hemitelia Pr. 1848 — Alsophila eriocarpa, scrobiculata, Unguis cati Fée 1869.
pallescens Sod. Rec. 20. 1883; Cr. vasc. quit. 535. 1893. NPfl. 134 — Ecuador.
Parkeri J. Sm. Hist. Fil. 249. 1875 = Hemitelia multiflora*.
parvula Jenm. JoB. **1879**. 258 = Hemitelia.
pauciflora (Kze.) Pr. Abh. böhm. Ges. V. **5**. 343. 1848; Kuhn, Linn. 36. 156. 1869 — Venezuela. — Cyathea Kze. 1846 (nomen). Alsophila paleolata pt. HB. 33.
paucifolia Bak. Syn. 456. 1874 — Ecuador.
pectinata Fée, Cr. vasc. Br. **1**. 168 t. 60 f. 1. 1869 = A. leucolepis.
Perriniana Spr. Nova Acta **10**. 232. 1821 = Woodsia obtusa var.
peruviana Kl. Linn. **20**. 441. 1847 = A. phalerata.
petiolulata Karst.; Mett. Ann. sc. nat. V. **2**. 263. 1864 (nomen) = Cyathea divergens.
phalerata Mart. Ic. Cr. Bras. 67 t. 30 f. 1. t. 42. 1834. (HB. 457). NPfl. 134 — Amer. trop.
Cyathea Mart. Denkschr. Regensb. bot. Ges. **2**. 146 t. 2 f. 3. 1822 — Alsophila infesta Kze. 1834. HB. 34, 457 (excl. β) — A. alutacea Kze. 1844. A. crassa Karst. 1869. — *Obs.* t. Kuhn est A. infesta Kze = Cyathea multiflora Sm.; conf. Hemitelia multiflora. Vide Engl. Jahrb. **24**. 83.
phegopteroides Hk. Syn. 32. 1865. NPfl. 132 — Peru.
philippinense ht. Veitch; Gard. Chr. n. s. **4**. 179. 1875 — Hort. (Ins. Philipp.)
pilosa Mart. et Gal. Mém. Ac. Brux. **15**. 78 t. 22. 1842 = Dryopteris rudis.
pilosissima Bak. Syn. 457. 1874 — Peru.
pinnula Christ, Prim. Fl. Costar. **3**. 43. 1901 — Costa Rica.
plagiopteris Mart. Ic. Cr. Bras. 73 t. 50. 1834. HB. 35. NPfl. 134 — Brasilia. [Ecuador.
platyphylla Pr. Epim. 29. 1849 — Guiana.
podophylla Hk. JoB. **9**. 334. 1857. HB. 43. Chr. 355. NPfl. 136 — Liu Kiu. China austr.-Siam. — Cyathea Bonii Christ 1890.
podophylla Bak. JoB. **1881**. 202 = A. Kalbreyeri.
Poeppigii Hk. sp. **1**. 43. 1844. HB. 35. NPfl. 134 — Peru. ? Brasilia.
A. villosa Kze. 1834 (non Desv. 1827).
Pohlii Pr. Tent. 62. 1836 = A. armata.
polycampta Kze. Bot. Zeit. **1846**. 475 = A. glabra.
polyphlebia Bak. Journ. Linn. Soc. **15**. 104. 1876 — Ins. Aru.
polystichoides Christ. Bull. Soc. bot. Belg. **35**. 177. 1896 — Costa Rica.
Poolii C. Chr. Ind. 46. 1905 — Madagascar.
A. vestita Bak. Journ. Linn. Soc. **15**. 412. 1876. NPfl. 136 (non Pr. 1849).
praecincta Kze. Flora 1839[1] Beibl. 53. HB. 458. Chr. 326. NPfl. 134 — Brasilia.
procera (Willd.) Desv. Prod. 319. 1827. HB. 33 (pt.). Chr. 324. NPfl. 134 — Brasilia. Peru.
Polypodium Willd. sp. **5**. 206. 1810 — Alsophila armigera Kze. 1834 (Peru).

pruinata Klf.; Kze. Linn. **9**. 99. 1834 = A. quadripinnata.

pterorachis Bak. Syn. 456. 1874. NPfl. 133 — Peru.

pubescens Bak. Syn. ed. I. 449. 1868; ed. II. 32. 1874. NPfl. 133 — Columbia-Peru.

pungens (Willd.) Klf.; Pr. Tent. 61. 1836. HB. 457. NPfl. 134 — Guiana.
Polypodium Willd. sp. **5**. 206. 1810.

pustulosa Christ, Bull. Boiss. II. **1**. 1019. 1901 — Liu Kiu.

pycnocarpa Kze. Linn. **9**. 97. 1834. HB. 36 — Peru.

quadripinnata (Gmel.) C. Chr. Ind. 47. 1905 — Amer. trop. ad Patagoniam, Chiloe, Juan Fernandez.
Polypodium Gmel. Syst. Nat. 2^2. 1314. 1791; P. glaucum Sw. Prod. 134. 1788 (non Thbg. 1784); Lophosoria Kuhn 1897; Alsophila Urban 1903 (non J. Sm. 1841); Polypodium pruinatum Sw. 1801; Alsophila Klf.; Kze. 1834. HB. 38. Chr. 326. NPfl. 135; Lophosoria Pr. 1848; Polypodium cinereum Cav. 1802; P. griseum Schkuhr 1806 — Lophosoria sp. plur. Pr., Fée, Kl., Fourn. — Trichosorus sp. Liebm. 1849 — Alsophila monticola Mart. 1834; Cyathea Pr. 1848.

quitensis C. Chr. Ind. 47. 1905 — Ecuador.
A. Bakeri Sod. Cr. vasc. quit. 532. 1893 (non Zeiller 1885).

radens Klf. Enum. 248. 1824; Mett. Fil. Lips. 109. 1856. HB. 35. NPfl. 134 — Brasilia.
Obs. t. Bak. Fl. bras. 1^2. 320 A. radens Klf. (non Mett.) est A. atrovirens (L. et F.); nomen plantae Mettenii ergo mutandum est.

ramispina Hk. Syn. 42. 1866; Bak. Journ. Linn. Soc. **22**. 292. Chr. 327 — Borneo.

Rebeccae F. Muell. Fragm. **5**. 53. 1865. HB. 40. NPfl. 136 — Queensland.

rheosora Bak. JoB. **1890**. 292. NPfl. 136 — Tonkin.

rheosora Christ, Bull. Boiss. **6**. 972. 1898 = A. Henryi.

Ridleyi Bak. Ann. Bot. **5**. 122. 1891 — Singapore.

rigidula Mart. Ic. Cr. Bras. 74 t. 51. 1834 = A. villosa.

Robertsiana F. Muell. Fragm. **5**. 54. 1865. HB. 459. NPfl. 138 — Queensland.

robusta de Vriese; Jungh. Java ed. germ. **1**. 310, 476. 1852 — Java.

rostrata Mart. Ic. Cr. Bras. 64 t. 39. 1834 = A. blechnoides.

rufa Fée, Cr. vasc. Br. **1**. 166 t. 59 f. 1. 1869 = A. armata.

sagittifolia Hk. Syn. ed. I. 37. 1866. ed. II. 456. 1874. Chr. 325. NPfl. 133 — Trinidad. — Hemitelia Jenm. 1898.

Salvinii Hk. Syn. 36. 1866. NPfl. 135 — Guatemala.

samoensis Brack. Expl. Exp. **16**. 287 t. 40 f. 1. HB. 39. NPfl. 138 — Samoa. Louisiade Archip. — Hemitelia Mett. 1859.

sangirensis Christ; Diels, Nat. Pfl. 1^4. 138. 1899; Christ in Warb. Monsunia **1**. 90. 1900 — Celebes.

scaberula Christ in Schum. et Laut. Fl. deut. Südsee 110. 1901 — N. Guinea.

Schaffneriana Fée, 8 mém. 109. 1827 — Mexico. — Lophosoria Fourn. 1872.

Schiedeana Pr. Tent. 62. 1836 (nomen). HB. 33. NPfl. 134 — Mexico.
Polypodium sp. indet. Schlecht. et Cham. Linn. **5**. 609. 1830.

Schlimii Mett. Ann. sc. nat. V. **2**. 263. 1864 (nomen); Kuhn, Linn. **36**. 157. 1859. HB. 458 — Costa Rica-Columbia.

Scottiana Bak. Gard. Chr. **1872**. 699. fig. 169 C = A. ornata.

scrobiculata Fée, Cr. vasc. Br. **1**. 157 t. 53. f. 1. 1869 = A. paleolata.

senilis Kl. Bot. Zeit. **1846**. 101; Linn. **20**. 442. 1847. NPfl. 134 — Columbia. Venezuela. — A. paleolata pt. HB. 33.

serrata J. Sm. Lond. JoB. **1**. 666. 1842 = A. aspera.

sessilifolia Jenm. JoB. **1882**. 325 — Jamaica. — Hemitelia Jenm. 1898.

setosa Klf. Enum. 249. 1824 = Hemitelia.
Shepherdii Bull, Cat. 1871 — Hort.
sikkimensis Clarke et Bak. Journ. Linn. Soc. **24**. 409. 1888 = A. ornata.
simulans Bak. JoB. **1891**. 3. NPfl. 136 — Madagascar.
Smithiana Pr. Abh. böhm. Ges. V. **5**. 342. 1848 } A. glauca.
Smithii Trevis. Atti Ist. Veneto II. **2**. 165. 1851 }
Sodiroi Bak. Ann. Bot. **5**. 149. 1891; Sod. Cr. vasc. quit. 534. 1893. NPfl. 134 — Ecuador. — A. alata Sod. Rec. 19. 1883 (non Fourn. 1873).
Sollyana Griff. Notul. **2**. 624. 1849; Ic. pl. As. rar. t. 130 f. 3-4 = A. glauca.
speciosa (Meyen) Pr. Tent. 62. 1836 — Brasilia.
Polypodium Meyen, Reise um die Erde **1**. 108. 1834.
Sprengeliana Mart. Ic. Cr. Bras. 75. 1834 = A. aspera.
Sprucei Hk. Syn. 37. 1866. NPfl. 134 — Ecuador.
squamata Kl. Linn. **18**. 541. 1844 = A. microphylla.
squamulata (Bl.) Hk. sp. **1**. 51. 1844 (pt.); Mett. Ann. Lugd. Bat. **1**. 52. 1863 — Malesia. — Gymnosphaera Bl. Enum. 243. 1828. Polypodium comosum Wall. 1828; Alsophila Wall. 1829. HB. 41. NPfl. 136; Athyrium Pr. 1836 (ex err.); Cystopteris Pr. 1836.
squamulata Hk. sp. **1**. 51. 1844 (pt.); Syn. ed. I. 43. 1866 = A. commutata.
stipularis Christ, Bull. Boiss. II. **4**. 958. 1904 — Costa Rica.
strigosa J. Sm. Lond. JoB. **1**. 666. 1842 = Hemitelia multiflora *Parkeri.
subaculeata Splitgb. Tijdschr. Nat. Gesch. **7**. 430. 1840 = Hemitelia multi- [flora.
subaspera Christ, Prim. Fl. Costar. **3**. 43. 1901 — Costa Rica.
subglandulosa Hance, Ann. sc. nat. V. **5**. 253. 1866. HB. 44 — Formosa.
Surinamensis J. Sm. Ferns br. and for. 244. 1866 = Hemitelia multiflora*.
Swartziana Mart. Ic. Cr. Bras. 73. t. 49. 1834 = A. armata.
Taenitis Kze. Linn. **9**. 90. 1834 = A. corcovadensis.
tahitensis Brack. Expl. Exp. **16**. 288. t. 40. 1854 = Hemitelia.
Telfairiana Wall.; Hk. sp. **1**. 56. 1844 = Cyathea borbonica.
tenera J. Sm. Lond. JoB. **1**. 666. 1842 = Cyathea.
tenerifrons Christ, Bull. Boiss. II. **4**. 959. 1904 — Costa Rica.
tenuisecta Bl.; Moore, Ind. 58. 1857 = Cystopteris.
temulata J. Sm. Lond. JoB. **1**. 666. 1842 (err. typ.) = A. lunulata.
Tijucensis Fée, Cr. vasc. Br. **1**. 173 t. 73 f. 1. 1869 = A. elongata.
tomentosa (Bl.) Hk. sp. **1**. 55. 1844. HB. 42. NPfl. 138 — Java. Ins. Camarinae. Formosa. — Chnoophora Bl. Enum. 244. 1828; Cyathea Zoll. et Moritz 1844 — Alsophila lepifera J. Sm. 1841.
tomentosa Pr. Tent. 62. 1836 = A. villosa.
trichodesma Scort.; Bedd. JoB. **1887**. 321. NPfl. 136 — Perak.
trichophlebia Bak. Ann. Bot. **5**. 189. 1891. NPfl. 134 — Paraguay.
?**tristis** Bl.; Moore, Ind. 58. 1857. HB. 44. Chr. 328. NPfl. 138 — Java. Aspidium Bl. Enum. 169. 1828; Lastrea Moore 1858; Cystopteris Mett. 1864. Davallia Racib. 1898. Allantodia? scabra Kze. 1848; Athyrium? Pr. 1849.
truncata Brack. Expl. Exp. **16**. 289 t. 41. 1854. HB. 39. Chr. 329. NPfl. 138 — Fiji. Samoa. Celebes. Batjan. — Hemitelia Christ 1897.
Tumacensis J. Sm. Lond. JoB. **1**. 667. 1842 = A. elongata.
umbrosa Wall. List pag. 64 (n. 336). 1829 = A. glabra.
Unguis Cati Fée, Cr. vasc. Br. **1**. 165. t. 58 f. 2. 1869 = A. paleolata.
Van Gaerdtii van Houtte; (Salom. Nom. 31. 1883) = ? *A. Van Geertii* van Geert; (J. Sm. Ferns br. and for. ed. II. 321. 1877) — Hort.

Veitchii Bak. Syn. ed. II. 41. 1874 — Aneiteum.
venulosa Wall. List pag. 63. 1829 = A. glabra.
vernicosa Mett.; Kuhn, Linn. **36**. 155. HB. 37. NPfl. 134 — Venezuela.
vestita J. Sm. Lond. JoB. **1**. 667. 1842 (nomen) = A. armata.
vestita Pr. Epim. 27. 1849 — Guiana.
vestita Bak. Journ. Linn. Soc. **15**. 412. 1876 = A. Poolii.
vexans Ces. Atti Ac. Napoli 7^8. 4. 1876 = A. glabra.
villosa (H.B. Willd.) Desv. Prod. 319. 1827. HB. 35. Chr. 325. NPfl. 134 — Amer. austr. trop. — Cyathea H.B. Willd. sp. **5**. 495. 1810: Chnoophora Humboldtii Klf. 1824; Alsophila Kl.; Kze. 1850. — A. rigidula Mart. 1834.
villosa Kze. Linn. **9**. 99. 1834 = A. Poeppigii.
vitiensis Carr. in Seem. Fl. Vit. 334. 1873 = A. extensa var.
Walkerae J. Sm. Hist. Fil. 249. 1875 = Hemitelia.
Wallacei Mett.; Kuhn, Linn. **36**. 153. 1869. HB. 459. NPfl. 136 — Borneo.
Wallichiana Pr. Tent. 62. 1836 = A. glauca.
Warburgii Christ, Ann. Jard. Buit. **15**. 80 t. 13 f. 3. 1897 = Dryopteris subarborea var.
Weigeltii Roem.; Pr. Tent. 61. 1836 = Hemitelia multiflora *Parkeri.
Wendlandi Mett.; Kuhn, Linn. **36**. 158. 1869. HB. 457 — Costa Rica.
Woolsiana F. Muell. Fragm. 8. 179. 1874. NPfl. 138 — Queensland.
xantholepis Christ; Diels, Nat. Pfl. 1^4. 138. 1899 — Ins. Batjan.
Zenkeri Hier.; Diels, Nat. Pfl. 1^4. 135. 1899 — Kamerun.

Alsophilopsis Karst. Fl. Columb. **1**. 4. 1858 *(Alsophila §)* = Alsophila.

AMAUROPELTA Kunze, Farnkr. **1**. 86, 109. 1843 = **Dryopteris.**
Breutelii Kze. Farnkr. **1**. 109 t. 51. 1843 = D. limbata.

AMBLIA Pr. Tent. Pter. 184. 1836 (Amblya Fée, Gen. 283. 1850-52) = **Phanerophlebia.**
juglandifolia Pr. Tent. 185 t. 7 f. 2. 1836.
latifolia Fée, 8 mém. 101. 1857 = P. juglandifolia.
serrata Fée, 10 mém. 35 t. 40 f. 2. 1865 — St. Helena (patria prob. erronea). Aspidium Kuhn 1868 (non Sw. 1806).

AMESIUM Newman, Hist. brit. Ferns ed. II. 10. 1844 = **Asplenium.**
germanicum Newm. Hist. br. Ferns ed. II. 10. 1844.
Ruta muraria Newm. Hist. br. Ferns ed. II. 10. 1844.
septentrionale Newm. Hist. br. Ferns ed. II. 10. 1844.

Ampelogramma J. Smith, Ferns brit. and for. 284. 1866 *(Gymnogramma §)* = Gymnogramma.

?AMPELOPTERIS Kunze, Bot. Zeit. **1848**. 114 = **Dryopteris.**
elegans Kze. Bot. Zeit. **1848**. 114 }
firma Kze. Linn. **24**. 251. 1851 } = D. prolifera.

Ampelopteris Klotzsch, Linnaea **20**. 430. 1847 *(Taenitis §)* = Eschatogramme.

AMPHIBLESTRA Presl, Tent. Pter. 150. 1836 — (Polypodiaceae Gen. **80**).
latifolia (H.B. Willd.) Pr. Tent. 151 t. 6 f. 1. 1836. NPfl. 288 — Venezuela. Pteris H.B. Willd. sp. **5**. 370. 1810. HB. 173. Chr. 174.
?*longifolia* Pr. Tent. 151. 1836 (nomen) — Chile.
simplex Fourn. Bot. Zeit. **1873**. 8 = Dictyoxiphium panamense.

AMPHICOSMIA Gardner, Lond. Journ. Bot. **1**. 441. 1842 = **Hemitelia.**
?alternans Moore, Ind. CIV. 1857 = Cyathea a.
australis Moore, Ind. 59. 1857.
Beyrichiana Moore, Ind. CIV. 1857 = H. setosa.
Brunoniana Bedd. Hdb. 10. 1883 = Cyathea B.

capensis Kl. Allg. Gartenzeit. **1856**. 107.
Cumingii Moore, Ind. 60. 1857 = H. denticulata.
decipiens Bedd. Ferns br. Ind. Suppl. 1. 1876 = Cyathea d.
Hostmanni Moore, Ind. CIV. 1857 = H. multiflora*.
javanica Moore, Ind. 60 1857 = H. Junghuhniana.
Kegelii Moore, Ind. 60. 1857 = H. multiflora.
laevis Moore, Ind. CIV. 1857 = H. multiflora.
lingulata Moore, Ind. 60. 1857 = H. multiflora.
macrocarpa Moore, Ind. 60. 1857.
manilensis Moore, Ind. 61. 1857.
multiflora Gardn. Lond. JoB. **1**. 441. 1842.
nigricans Moore, Ind. 61. 1857.
Parkeri Moore, Ind. 61. 1857 = H. multiflora*.
riparia Gardn. Lond. JoB. **1**. 441 t. 12. 1842 = H. capensis.
strigosa Moore, Ind. 61. 1857 = H. multiflora *Parkeri.
tahitensis Moore, Ind. 61. 1857.
urolepis Moore, Ind. 61. 1857 = H. multiflora *Parkeri.
Walkerae Moore, Ind. CIV. 1857.

AMPHIDESMIUM Schott, Gen. Fil. ad t. 5. 1834 = **Alsophila.**
blechnoides Kl. Linn. **20**. 372. 1847 }
Parkeri Schott, Gen. Fil. ad t. 5. 1834 } = A. blechnoides.
rostratum J. Sm. Lond. JoB. **1**. 201. 1842 }

AMPHIPTERUM Presl, Epim. Bot. 258. 1849.
fuscum Pr. Epim. 258. 1849 = Hymenophyllum.

AMPHISORIA Trevisan, Atti dell' Ist. Veneto II. **2**. 168. 1851 = **Polybotrya.**

AMPHORADENIUM Desv. Prod. 335. 1827 = **Polypodium.**
australe Desv. Prod. 336. 1827 }
gaudichaudii Desv. Prod. 336. 1827 } = P. tamariscinum.
minutum Desv. Prod. 336. 1827 = P. hymenophylloides.

ANAPAUSIA Presl, Epim. Bot. 185. 1849 = **Leptochilus.**
Gymnopteris § Anapausia Pr. Tent. 244. 1836.
acuminata Pr. Epim. 188. 1849 = L. nicotianifolius.
aliena Pr. Epim. 187. 1849.
bicuspis Moore, Ind. XXI. 1857 = Cheiropleuria.
decurrens Pr. Epim. 186. 1849.
dentata Pr. Epim. 188. 1849 = L. alienus.
Heydelotii Pr. Epim. 187. 1849 = L. Heudelotiii.
nicotianaefolium Pr. Epim. 189. 1849.
portoricensis Pr. Epim. 188. 1849 = L. alienus.
semipinnatifida Pr. Epim. 187. 1849 = L. alienus var.
vespertilionis Moore, Ind. XXI. 1857 = Cheiropleuria bicuspis.
Zollingeri Pr. Epim. 187. 1849.

ANAPELTIS J. Smith, Cat. cult. Ferns 5. 1857 = **Polypodium.**
Phlebodium § Anapeltis J. Sm. Bot. Mag. **72**. Comp. 12. 1846.
geminata J. Sm. Ferns br. and for. 86. 1866.
lycopodioides J. Sm. Cat. cult. ferns 6. 1857.
nitida J. Sm. Cat. cult. ferns 6. 1857 = P. lycopodioides.
owariensis J. Sm. Ferns br. and for. 86. 1866 = P. lycopodioides.
serpens J. Sm. Cat. cult. ferns 5. 1857 = P. runcinatum.
squamulosa J. Sm. Cat. cult. ferns 6. 1857.

stigmatica J. Sm. Ferns br. and for. 86. 1866.
vacciniifolia J. Sm. Cat. cult. ferns 5. 1857.
venosa J. Sm. Cat. cult. ferns 6. 1857 = P. stigmaticum.
ANAXETUM Schott. Gen. Fil. t. 1. 1834 = **Polypodium.**
crassifolium Schott, Gen t. 1. 1834.
ANCHISTEA Presl. Epim. Bot. 71. 1849 = **Woodwardia.**
virginica Pr. Epim. 71. 1849.
ANEIMIA Swartz, Syn. Fil. 6, 155. 1806 — (Schizaeaceae Gen. 4).
(Anemia Sw.; Aneimia Klf. Enum. 51. 1824).
abscissa Schrad. Gött. gel. Anz. **1824**. 864 = A. hirta.
adiantifolia (L.) Sw. Syn. 157. 1806. HB. 434. Prantl, Schiz. 126. Chr. 349. NPfl. 371 — Amer. trop. Florida. — Osmunda L. sp. **2**. 1065 1753; Ornithopteris Bernh. 1806 — Osmunda asplenifolia Sav. 1797; Anemia Sw. 1806; Anemirhiza J. Sm. (1854) — Anemia carvifolia Pr. 1825.
affinis Bak. Syn. 525. 1874. Prantl, Schiz. 128 — Mexico.
ahenobarba Christ in Schwacke, Pl. Nov. Mineiras **2**. 37. 1900; Bull. Boiss. II. **2**. 696. 1902 — Brasilia centr.
anthriscifolia Schrad. Gött. gel. Anz. **1824**. 865. Prantl, Schiz. 97; Lindm. Ark. för Bot. **1**. 258 t. 12 f 1. — Mexico-Paraguay. — A. tomentosa pt. HB. 433.
aspera (Fée) Bak. Journ. Linn. Soc. **14**. 27. 1873. Prantl, Schiz. 99 — Brasilia. Costa Rica. — Aneimiaebotrys Fée, Cr. vasc. Br. **1**. 267 t. 78 f. 2. 1869.
asplenifolia Sw. Syn. 157. 1806 = A. adiantifolia.
aurita Sw. Syn. 157. 1806. HB. 435. Prantl, Schiz. 122. NPfl. 371 — Jamaica. Puerto Rico. — Osmunda Sw. Prod. 127. 1788; Mohria J. Sm. 1843.
bipinnata Sw. Syn. 157. 1806 = Osmunda cinnamomea.
bipinnata Moore, Ind. CXVI. 1857 = A. cicutaria.
Breuteliana Pr. Suppl. 90. 1845 = A. hirta.
buniifolia (Gardn.) Moore, Ind. CXVI. 1857 — Brasilia.
Coptophyllum Gardn. Lond. JoB. **1**. 133. 1842; Hk. Ic. pl. t. 477. 1842; Mohria J. Sm. 1843. Aneimia dichotoma Gardn.; Pr. Suppl. 80. 1845. HB. 435. Prantl, Schiz. 100. Chr. 349. NPfl. 368 — A. tenuifolia Pr. 1848, 1849.
carvifolia Pr. Rel. Haenk. **1**. 74. 1825 = A. adiantifolia.
caudata Klf. Enum. 52. 1824 = A. radicans.
cheilanthoides Klf. Enum. 53. 1824 = A. tomentosa.
cicutaria Kze.: Spr. Syst. **4**. 31. 1827; Kze. Linn. **9**. 22. 1834; Anal. 9 t. 5 f. 2. Prantl, Schiz. 124. NPfl. 371 — Bahama. Cuba. Yucatan.
Coptophyllum Kl. 1844; Ornithopteris Und. 1902; Anemia bipinnata Moore 1857. HB. 435 (non Sw. 1806). Mohria intermedia J. Sm. 1843.
ciliata Pr. Del. Prag. **1**. 158. 1822 = A. repens.
ciliata × Phyllitidis Prantl, Schiz. 121. 1881 = A. laciniata.
coccinea Loud. Hort. brit. ed. nov. 488. 1850 — Hort.
collina Raddi, Opusc. sci. Bol. **3**. 282. 1819; Pl. Bras. **1**. 71 t. 12. 1825. HB. 432. Prantl. Schiz. 114. Chr. 348. NPfl. 371 — Brasilia. — A. vellea Schrad. 1824.
collina × phyllitidis Prantl, Schiz. 120. 1881 — Hort.
cordifolia Pr. Rel. Haenk. **1**. 73 t. 11 f. 3. 1825 = A. phyllitidis.
coriacea Gris. Cat. Pl. Cub. 272. 1866. (HB. 433). Prantl, Schiz. 123. NPfl. 371. [— Cuba.
cornea Prantl, Schiz. 104. 1881 — Mexico.
cuneata Kze.: Spr. Syst. **4**. 32. 1827: Kze. Anal. 8 t. 8 f. 1. 1837. HB. 434. Prantl, Schiz. 125. NPfl. 371 — Cuba.
deltoidea Sw. Syn. 156. 1806 = A. tomentosa.

densa Link. Hort. Berol. **2**. 142. 1833 = A. phyllitidis.
dentata Gardn.; Field. et Gardn. Sert. pl. ad. t. 70. 1844 = A. filiformis.
dichotoma Gardn.; Pr. Suppl. 80. 1845 = A. buniifolia.
dimorphostachys Bak. Engl. Jahrb. **17**. 522. 1893 — Brasilia.
dissecta Pr. Rel. Haenk. **1**. 74 t. 11. f. 4. 1825 = A. tenella.
distans Fée, 9 mém. 41 (33). 1857; Prantl. Schiz. 127 — Mexico.
diversifolia Schrad. Gött. gel. Anz. **1824**. 864 = Brasilia.
— A. Vespertilio Schrad. 1824. A. Schraderiana Mart. Ic. Br. Bras. 113 t. 58. 1834. HB. 434. Prantl, Schiz. 118.
Dregeana Kze. Linn. **10**. 493. 1836. HB. 431. Prantl, Schiz. 116. Chr. 348. NPfl. 371 — Afr. austr.
elegans (Gardn.) Pr. Suppl. 81. 1845. Prantl, Schiz. 90. Chr. 351. NPfl. 368 — Brasilia. (? Cuba). — Trochopteris Gardn. Lond. JoB. **1**. 74 t. 4. 1842. [HB. 436.
eximia Taubert, Engl. Jahrb. **21**. 422. 1896 — Brasilia.
A. Schwackeana Christ, Farnkr. 351. c. fig. 1897. NPfl. 368.
ferruginea HBK. Nov. Gen. et Sp. **1**. 32. 1815 = A. fulva.
filiculifolia (L.) Sw. Syn. 158. 1806 — Ind. occ. — Osmunda L. sp. **2**. 1065. 1753 (Plum. t. 161) — (an Polybotrya osmundacea?).
filiformis (Sav.) Sw. Syn. 156. 1806. HB. 431 (pt.). Prantl, Schiz. 106. NPfl. 370 — Amer. austr. trop. Jamaica.
Osmunda Sav. in Lam. Enc. **4**. 652. 1797 — Anemia dentata Gardn. 1844 — A. Schomburgkiana Pr. 1845 — A. gracilis Schrad. 1824.
flexuosa (Sav.) Sw. Syn. 156. 1806. Prantl, Schiz. 94 — Amer. trop.
Osmunda Sav. in Lam. Enc. **4**. 652. 1797. Anemia Raddiana Link 1833. A. tomentosa pt. HB. 433. — A. oblonga Sturm 1859.
— *A. villosa H.B. Willd. sp. **5**. 92. 1810; Osmunda Poir 1816.
fraxinifolia Raddi, Opusc. sci. Bol. **3**. 282. 1819; Pl. Bras. **1**. 69 t. 8 bis. 1825 = A. phyllitidis.
fulva (Cav.) Sw. Syn. 157. 1806. Prantl, Schiz. 98. NPfl. 368 — Amer. austr. trop.
Osmunda Cav. Icon. **6**. 70 t. 593 f. 2. 1801. Anemia ferruginea HBK. 1815. — A. rubrostipes Pohl; Sturm 1859. A. tomentosa β HB. 433.
Gardneri Hk. Ic. pl. t. 190. 1837. HB. 431. Prantl, Schiz. 91. NPfl. 368 — Brasilia. — A. Gardneriana Chr. 346. 1897 (non Pr. 1845) — A. Glaziovii [Fée 1869.
Gardneriana Pr. Suppl. 82. 1845 = A. glareosa.
Gardneriana Christ, Farnkr. 346. 1897 = A. Gardneri.
glareosa Gardn. in Field. et Gardn. Sert. pl. t. 70. 1844. HB. 432. Prantl, Schiz. 91. Chr. 347. NPfl. 368 — Brasilia. Columbia. — A. Gardneriana Pr. 1845.
Glaziovii Fée, Cr. vasc. Br. **1**. 207 t. 74 f. 2. 1869 = A. Gardneri.
glomerata Gardn.; Moore, Ind. 66. 1857 = A. rutifolia.
gracilis Schrad. Gött. gel. Anz. **1824**. 865 = A. filiformis.
Haenkei Pr. Rel. Haenk. **1**. 74. 1825 = A. phyllitidis.
helveola Fée, 9 mém. 41 (32). 1857; Prantl, Schiz. 127 — Mexico.
heterodoxa Christ, Ann. Cons. Jard. bot. Genève **3**. 45. 1899 — Brasilia.
hirsuta (L.) Sw. Syn. 156. 1806. HB. 433 (pt.). Chr. 349 (pt.). NPfl. 370 (pt.). Lindm. Ark. för. Bot. **1**. 259 t. 12 f. 3. — Amer. trop. (excl. Brasilia?).
Osmunda L. sp. **2**. 1064. 1753; Ornithopteris Bernh. 1806.
hirsuta Prantl, Schiz. 108. 1881 (pt.?) (et auctt.?) = A. repens et tenella.
hirsuta × phyllitidis Prantl, Schiz. 121. 1881 — Columbia.
hirta (L.) Sw. Syn. 155. 1806. Prantl, Schiz. 115. Chr. 348. NPfl. 371 — Ind.

occ. Costa Rica. Brasilia. — Osmunda L. sp. **2**. 1064. 1753; Aneimidictyum Pr. 1845. Aneimia Breuteliana Pr. 1845. HB. 432.

hirta Raddi, Opusc. sci. Bol. **3**. 281. 1819 = A. phyllitidis.

hispida Kze. Linn. **9**. 20. 1834. (HB. 433) — Peru.

humilis (Cav.) Sw. Syn. 156. 1806. Prantl, Schiz. 103. NPfl. 370 — Panama. Columbia. Brasilia. — Osmunda Cav. Icon. **6**. 69 t. 592 f. 3. 1801; Anemia oblongifolia β HB. 431 — A. Seemanni Hk. 1848.

humilis Hk. Exot. Flora **1**. t. 28. 1823 = A. filiformis.

imbricata Sturm. Fl. bras. **1**². 205. 1859. Prantl, Schiz. 93 — Brasilia.

incisa Schrad. Gött. gel. Anz. **1824**. 865. (HB. 433). Prantl, Schiz. 115 — Brasilia Aneimidictyum Pr. 1845.

Karwinskyanum (Pr.) Prantl, Schiz. 99. 1881 — Mexico.
A. villosa ζ Karwinskyana Pr. Suppl. 83. 1845.

Kunzeana Kl. Linn. **18**. 526 (nota). 1844 (nomen) — ?

laciniata Link. Fil. sp. 25. 1841. (HB. 435) — Venezuela. Mexico.
Aneimidictyum Pr. 1845. Aneimia media Link 1841; A. ciliata × Phyllitidis Prantl, Schiz. 121. 1881.

lanceolata Lodd. Bot. Cab. **15** index (t. 1416). 1828 = A. phyllitidis.

Langsdorffiana Pr. Suppl. 89. 1845 = A. phyllitidis var.

lanuginosa Bongard; Sturm, Fl. bras. **1**². 210. 1859. Prantl, Schiz. 92 — Brasilia.

laxa Lindm. Ark. för Bot. **1**. 261 t. 13. 1903 — Brasilia.

longifolia Raddi, Pl. bras. **1**. 69. t. 8. 1825 = A. phyllitidis.

longistipes (Liebm.) C. Chr. Ind. 53. 1905 — Mexico. Venezuela. Trinidad.
A. pilosa β longistipes Liebm. Vid. Selsk. Skr. V. **1**. 301 (seors. 149). 1849
A. pastinacaria Prantl, Schiz. 110. 1881.

mandioccana Raddi, Opusc. sci. Bol. **3**. 282. 1819; Pl. bras. **1**. 70 t. 9 f. 1. 1825. HB. 432. Prantl, Schiz. 113. Chr. 348 — Brasilia. Jamaica.

mandioccana Hk. Gen. t. 90. 1842 = A. Breuteliana.

media Link, Fil. sp. 25. 1841 = A. laciniata.

mexicana Kl. Linn. **18**. 526. 1844. HB. 433. Prantl, Schiz. 123. Chr. 348 NPfl. 371 — Texas-Mexico-Guatemala.
Ornithopteris Und. 1900. — Aneimia speciosa Pr. 1845.

microstachys Fée, Cr. vasc. Br. **1**. 206 t. 74 f. 3. 1869 = { A. radicans (f. Prantl) / A. filiformis (f. Bak).

millefolia Gardn.: Pr. Suppl. 80. 1845. HB. 436. Prantl, Schiz. 101. Chr. 350. NPfl. 368 — Brasilia. Columbia. — Coptophyllum Gardn. Lond. JoB. **1**. 133. 1842 et in Hk. Ic. pl. t. 478. 1842: Mohria J. Sm. 1843.

nana Bak. Engl. Jahrb. **17**. 522. 1893 — Brasilia.

nervosa Pohl; Sturm, Fl. bras. **1**². 193. 1859. Prantl, Schiz. 117. NPfl. 371 — [Brasilia

obliqua Schrad. Gött. gel. Anz. **1824**. 864 = A. phyllitidis var.

oblonga Sturm. Fl. bras. **1**². 206. 1859 = A. flexuosa var.

oblongifolia (Cav.) Sw. Syn. 156. 1806. HB. 431 (excl. β). Prantl, Schiz. 105. Chr. 346. NPfl. 369 — Amer. trop.
Osmunda Cav. Icon. **6**. 69 t. 592 f. 2. 1801 — Aneimia pumila Kl. 1844.

obtusa Desv. Berl. Mag. **5**. 308. 1811; Prantl, Schiz. 127 — Bonaria.

opaca Fée, 9 mém. 41 (33) 1857 = A. repens.

ouropretana Christ in Schwacke, Pl. Nov. Mineiras **2**. 36. 1900: Bull. Boiss. II. **2**. 693. 1902 — Brasilia.

pallida Gardn. in Field. et Gardn. Sert. pl. ad t. 70. 1844 (HB. 433). Prantl Schiz. 110 — Brasilia.

palmarum Lindm. Ark. för Bot. **1**. 261 t. 14. 1903 — Brasilia.

pastinacaria Prantl, Schiz. 110. 1881 = A. longistipes.

phyllitidis (L.) Sw. Syn. 155. 1806. HB. 435. Prantl, Schiz. 119 (pt.). Chr. 348. NPfl. 371 – Amer. trop. — Osmunda L. sp. **2**. 1064. 1753; Anemidictyon J. Sm. 1842 — Anemia fraxinifolia Raddi 1819; Anemidictyon J. Sm. 1843 — Anemia sorbifolia Schrad. 1824. — A. Haenkei Pr. 1825; Aneimidictyum Pr. 1845 — Anemia longifolia Raddi 1825 — A. densa Link 1833; Anemidictyon J. Sm. 1843 — Anemia obliqua Schrad. 1824; Aneimidictyon Fée 1866 — Aneimia Langsdorffiana Pr. 1845. HB. 433 (Langsd. et Fisch. t. 28).

pilosa Mart. et Gal. Mém. Ac. Brux. **15**. 19 t. 2 f. 1. 1842. Prantl, Schiz. 104. NPfl. 370 — Mexico-Columbia.

pilosa Pr. Suppl. 86. 1845 = A. Presliana.

pilosa *β* longistipes Liebm. Vid. Selsk. Skr. V. **1**. 301 (seors. 149). 1849 = A. longistipes.

Pohliana Sturm, Fl. bras. 1^2. 195. 1859. (HB. 433). Prantl, Schiz. 116. NPfl. 371 — Brasilia.

Presliana Prantl, Schiz. 104. 1881 — Brasilia. Guatemala. Columbia. A. pilosa Pr. Suppl. 86. 1845. (HB. 431) (non Mart. et Gal. 1842).

pulchra Pohl; Prantl, Schiz. 109. 1881 — Brasilia. — A. filiformis pt. auctt. HB. 431.

pumila Kl. Linn. **18**. 526. 1844 = A. oblongifolia.

pyrenaea Taubert, Engl. Jahrb. **21**. 422. 1896 — Brasilia.

Raddiana Link, Hort. Berol. **2**. 144. 1833 = A. flexuosa.

radicans Raddi, Opusc. sci. Bol. **3**. 282. 1819; Pl. bras. **1**. 70 t. 10. 1825. Prantl, Schiz. 112. Chr. 347. NPfl. 370 — Brasilia. — A. caudata Klf. 1824. HB. 432.

radicans *β* Raddi, Pl. bras. **1**. 70 t. 11. 1825 = A. rotundifolia.

repens Raddi, Opusc. sci. Bol. **3**. 282. 1819; Pl. bras. **1**. 71 t. 9 f 2 b (*β* major; non *α*) 1825 — Columbia. Venezuela. Brasilia. A. ciliata Pr. 1822. Prantl, Schiz. 107 (excl. *β*). NPfl. 370. A. hirsuta pt. auctt. HB. 433. Prantl, Schiz. 115. Chr. 349. NPfl. 370.

repens *α* minor Raddi, Pl. bras. **1**. 71 t. 9 f. 2 a 1825 = A. filiformis.

rotundifolia Schrad. Gött. gel. Anz. **1824**. 865. HB. 432. Prantl. Schiz. 112. Chr. 347. NPfl. 371 — Brasilia. — A. radicans *β* Raddi 1825.

rubrostipes Pohl; Sturm, Fl. bras. 1^2. 202. 1859 = A. fulva.

rutifolia Mart. Ic. Cr. Bras. 112 t. 55 f. 1. 1834. HB. 434. Prantl, Schiz. 101 — Brasilia.

scandens Spr. Nova Acta **10**. 233. 1821 = Stenochlaena sorbifolia.

Schimperiana Pr. Suppl. 84. 1845. (HB. 434). Prantl, Schiz. 96. Chr. 349. NPfl. 368 — Afr. trop. India austr. — A. Wightiana Gardn. 1847 (India).

Schomburgkiana Pr. Suppl. 86. 1845 = A. filiformis.

Schraderiana Mart. Ic. Cr. Bras. 113 t. 58. 1834 = A. diversifolia.

Schwackeana Christ, Farnkr. 351. 1897 = A. eximia.

Seemanni Hk. Lond. JoB. **7**. 564 t. 16. 1848 = A. humilis.

sorbifolia Schrad. Gött. gel. Anz. **1824**. 864 = A. phyllitidis.

speciosa Pr. Suppl. 89. 1845 = A. mexicana.

tenella (Cav.) Sw. Syn. 156. 1806; Lindm. Ark. för Bot. **1**. 260 t. 12 f. 4. 1903 — Brasilia. — Osmunda Cav. Icon. **6**. 69 t. 592 f. 1. 1801. Anemia dissecta Pr. 1825. A. hirsuta *β* HB. 433; Prantl, Schiz. 127.

tenera Pohl; Sturm, Fl. bras. 1^2. 208. 1859. Lindm. Ark. för Bot **1**. 260 t. 12 f. 2 — Brasilia. — A. ciliata *β* tenera Prantl, Schiz. 108.

tenuifolia Pr. Abh. böhm. Ges. V. **5.** 327. 1848; Epim. 10 t. 4. 1849 = A. buniifolia var.

tomentosa (Sav.) Sw. Syn. 157. 1806. HB. 433 (pt). Prantl, Schiz. 95. Chr. 349. NPfl. 368 — Amer. austr. trop.

Osmunda Sav. in Lam. Enc. **4.** 652. 1797. O. deltoidea Cav. 1801; Anemia Sw. 1806 — A. cheilanthoides Klf. 1824.

trichorhiza Gardn. in Hk. Ic. pl. t. 876. 1852. HB. 434. Prantl, Schiz. 100. Chr. 350. NPfl. 368 — Brasilia.

triniaefolia Gardn.; Fée, Cr. vasc. Br. **1.** 206. 1869 (err.) = A. tenuifolia.

Tweedieana Hk. Ic. pl. t. 906. 1854. HB. 434 — Uruguay; Brasilia austr.

Anemidictyon Moore 1857. Aneimia Phyllitidis pt. Prantl, Schiz. 119.

Ulei Christ in Schwacke, Pl. Nov. Mineiras **2.** 36. 1900; Bull. Boiss. II. **2.** 694. 1902 — Brasilia.

vellea Schrad. Gött. gel. Anz. **1824.** 865 = A. collina.

verticillata (L.) Sw. Syn. 158. 1806 — Ind. occ. — Osmunda L. sp. **2.** 1065. 1763 (Plum. t 160); Spathepteris Pr. 1845. (planta dubia, certe non Aneimia).

vespertilio Schrad. Gött. gel. Anz. **1824.** 865 = A. diversifolia.

villosa H.B. Willd. sp. **5.** 92. 1810 = A. flexuosa*.

Warmingii Prantl, Schiz. 113. 1881 — Brasilia.

Wightiana Gardn. Calc. Journ. **7.** 10 t. 1. 1847 = A. Schimperiana.

Wrightii Bak. Syn. 435. 1868. Prantl, Schiz. 125 — Cuba. Yucatan.

Ornithopteris Millsp. 1903.

ANEIMIAEBOTRYS Fée, Crypt. vasc. Brésil **1.** 267. 1869 = **Aneimia.**

aspera Fée, Cr. vasc. Br. **1.** 267 t. 78 f. 2. 1869.

ANEIMIDICTYON J. Smith in Hook. Gen. Fil. t. 103. 1842 = **Aneimia.**

(Anemidictyon J. Sm.; Aneimidictyum Pr. Suppl. 91. 1845).

densum J. Sm. Lond. JoB. **2.** 387. 1843 = A. phyllitidis.

fraxinifolium J. Sm. Lond. JoB. **2.** 387. 1843 = A. phyllitidis.

Haenkei Pr. Suppl. 94. 1845 = A. phyllitidis.

hirtum Pr. Suppl. 92. 1845 = { A. hirta. / A. phyllitidis. }

incisum Pr. Suppl. 95. 1845.

laciniatum Pr. Suppl. 94. 1845.

obliquum Fée, 11 mém. 124. 1866 = A. phyllitidis var.

Phyllitidis J. Sm. in Hk. Gen. Fil. t. 103. 1842.

Tweedieanum Moore, Ind. CXVI. 1857.

ANEMIA Sw. = **Aneimia.**

ANEMIRHIZA J. Smith, Bot. Voy. Herald 243. 1854 = **Aneimia.**

adiantifolia J. Sm. (Bot. Herald 243. 1854).

ANETIUM (Kunze) Splitgerber, Tijdschr. Nat. Gesch. **7.** 395. 1840 — (Polypodiaceae Gen. **92**). — Acrostichum § Anetium Kunze hb.

citrifolium (L.) Splitgb. Tijdschr. Nat. Gesch. **7.** 395. 1840. Chr. 56. NPfl. 302 — Amer. trop. — Acrostichum L. sp. **2.** 1067. 1753; Antrophyum Fée 1851-52; Hemionitis Hk. 1864. HB. 399 — H. parasitica L. 1759 — Antrophyum pendulum Lepr.; Fée 1851-52.

crinitum Pr. Epim. 176. 1849 = Elaphoglossum.

pachyphyllum Pr. Epim. 176. 1849 = Elaphoglossum.

reticulatum Pr. Epim. 176. 1849 = Elaphoglossum.

Sprucei Moore, Ind. 72. 1857 = A. citrifolium.

ANGIOPTERIS (Mitchell 1748); Adanson, Fam. des pl. **2.** 21. 1763 = **Onoclea.**

ANGIOPTERIS Hoffmann, Comm. Soc. Reg. Gotting. **12**. 29. (1793?) 1796 — (Marattiaceae Gen. **1**).

Obs. Numerosissimae species Vrieseanae certe non distinctae sunt, sed quia monographia recentior hujus generis non exstat, DE VRIESE sequor. BAKER (HB. 440) omnes formas sub una specie (A. evecta Hoffm.) conjunxit.

acrocarpa de Vriese, Mon. 20. 1853 — Ins. Societatis.

affinis de Vriese, Ned. Kr. Arch. **3**. 192. 1852 — Java.

alata Nadeaud, Journ. de Bot. **11**. 120. 1897 — Tahiti.

amboinensis de Vriese, Ned. Kr. Arch. **3**. 195. 1852; Mon. 32 — Amboina.

angkolana de Vriese vide ankolana.

angustata Miquel, Ind. Sem. ht. Amst. 1849; Verh. Nederl. Inst. III. **4**. 50. t. 6 f. B. 1851; de Vriese, Mon. 26 — Java.

angustifolia Pr. Suppl. 21. 1845; de Vriese, Mon. 18. NPfl. 438 — Luzon.

ankolana de Vriese, Ned. Kr. Arch. **3**. 190. 1852 (angkolana): Mon. 19 t. 3 f. 9, t. 4 f. 9. 1853. NPfl. 438 — Java. Sumatra.

aphanosorus de Vriese, Ned. Kr. Arch. **3**. 194. 1852; Mon. 19 — Sumatra.

approximata de Vriese, Mon. 25. 1853 — Sumatra.

Arnottiana Miquel, Verh. Nederl. Inst. III. **4**. 53 t. 7 f. A. 1851: de Vriese, Mon. 28 — India.

assamica de Vriese, Mon. 33. 1853 — Assam.

attenuata Brack. Expl. Exp. **16**. 310. 1854 — Ins. Philipp.

aurata de Vriese, Mon. 22. 1853 — ? N. Zealand.

badioneura de Vriese, Ned. Kr. Arch. **3**. 187. 1852 = A. microsporangia.

Beecheyana de Vriese, Mon. 22. 1853 — Ins. Carolinae.

Brongniartiana de Vriese, Mon. 30 t. 3 f. 5, t. 4 f. 5. 1853. NPfl. 439 — [Tahiti.

campsophlebia de Vriese, Mon. 31. 1853 — India.

cartilagidens Christ, Bull. Boiss. **6**. 207 t. 4 f. 8. 1898 — Luzon.

caudata de Vriese, Mon. 20. 1853. NPfl. 438 — Ins. Philipp. Bismarck's Archip.

cochinchinensis de Vriese, Mon. 23 t. 3 f. 22, t. 4 f. 22. 1853 — Cochinchina.

commutata Pr. Suppl. 25. 1845; de Vriese, Mon. 33 — Tahiti.

crassifolia de Vriese, Ned. Kr. Arch. **3**. 193. 1852; Mon. 17. NPfl. 438 — Java.

crassipes Wall. List n. 187. 1828; Pr. Suppl. 23. 1845; de Vriese Mon. 27. NPfl. 438 — Ind. bor. Birma. China austr.

cupreata de Vriese, Mon. 21. 1853 — Ins. Societatis.

cuspidata de Vriese, Ned. Kr. Arch. **3**. 188. 1852; Mon. 18 — Java.

distans Pr. Suppl. 23. 1845; de Vriese, Mon. 31 — India.

Dregeana de Vriese, Mon. 17 t. 3 f. 8. t. 4 f. 8. 1853 = A. javanica.

D'Urvilleana de Vriese, Mon. 17 t. 3 f. 11, t. 4 f. 11. 1853. NPfl. 438 — Ins. Philipp.-Tahiti.

erecta Desv. Prod. 206. 1827 et auctt. (err.) = A. evecta.

evecta (Forst.) Hoffm. Comm. Soc. Reg. Gott. **12**. 29 t. 5. 1796; de Vriese, Mon. 16. HB. 440 (pt.). Chr. 358 (pt.). NPfl. 438 — Tahiti (Africa, Asia trop.).
Polypodium Forst. Prod. 81. 1786; Danaea Spr. 1799. ? Clementea palmiformis Cav. 1803 (f. Presl = A. angustifolia).

Gaudichaudiana de Vriese, Mon. 30 t. 3 f. 13; t. 4 f. 13. 1853 — Ind. bor.

Griffithsiana de Vriese, Mon. 29. 1853 (rectius Griffithiana) — Mergui.

Hartingeana de Vriese, Ned. Kr. Arch. **3**. 192. 1852; Mon. 25 — Java.

Helferiana Pr. Suppl. 22. 1845; de Vriese, Mon. 22 — Birma.

Hookeriana de Vriese, Mon. 29. 1853 — India.

Hügeliana Pr. Suppl. 25. 1845; de Vriese, Mon. 75 — India.

hypoleuca de Vriese, Ned. Kr. Arch. 2^2. 286. 1851; Mon. 21 — Java.
indica Desv. Berl. Mag. **5**. 307. 1811 — India.
javanica Pr. Suppl. 20. 1845 — Java.
A. Dregeana de Vriese, Mon. 17. 1853. NPfl. 438.
laciniata de Vriese, Mon. 30. 1853 - India (Sharapour).
Lasegueana de Vriese, Mon. 25. 1853 — »Huachine«. (? Ins. Societatis).
latifolia Pr. Suppl. 24. 1845: de Vriese, Mon. 27 — India (Pundoa).
Leschenaultiana de Vriese, Mon. 31 t. 3 f. 14. t. 4 f. 14. 1853 — Ceylon.
longifolia Grev. et Hk. Bot. Misc. **3**. 227. 1833; de Vriese Mon. 19. NPfl. 438 — Ins. Pitcairn. Ins. Societatis.
macrocephala Pr. Suppl. 26. 1845: de Vriese, Mon. 32 — India (Punjab).
macrophylla de Vriese, Mon. 34. 1853 — Hort.
madagascariensis de Vriese, Mon. 23. 1853 — Madagascar.
magnifica Miquel, Verh. Nederl. Inst. III. **4**. 54 t. 7 f. B. 1851; de Vriese, Mon. 32 — Ceylon.
marginata de Vriese, Mon. 29. 1853. NPfl. 439 — Ceylon.
microsporangia de Vriese, Ned. Kr. Arch. **3**. 190. 1852; Mon. 18 — Java, Sumatra. — A. badioneura de Vriese 1852.
Miqueliana de Vriese, Ned. Kr. Arch. **3**. 194. 1852; Mon. 26 — Java.
muricata Pr.; de Vriese, Ned. Kr. Arch. **3**. 195. 1852: Mon. 30 — Borneo.
pallescens de Vriese, Ned. Kr. Arch. **3**. 191. 1852; Mon. 25 — Sumatra. Java.
plagiocarpa de Vriese, Mon. 34 t. 3 f. 15. t. 4 f. 15. 1853 — Ceylon.
polysporangia de Vriese, Mon. 23. 1853. NPfl. 438 — Ceylon.
Presliana de Vriese, Ned. Kr. Arch. 2^2. 284. 1851; Mon. 20 — Java.
pruinosa Kze. Bot. Zeit. **1846**. 417; de Vriese, Mon. 26. NPfl. 438 — Java.
punctata de Vriese, Mon. 31 t. 3 f. 3, t. 4 f. 3. 1853 — Ceylon.
repandula de Vriese, Mon. 30 t. 3 f. 4, t. 4 f. 4. 1853 — India (Sharapour),
salicifolia (Pr.) de Vriese, Mon. 34. 1853. NPfl. 432 — India (Lahore).
Psilodochea Pr. Suppl. 28. 1845.
similis Pr.; de Vriese, Ned. Kr. Arch. **3**. 189. 1852: Mon. 17 — Java.
Smithii Racib. Bull. int. Acad. Cracovie **1902**. 54 — Cult. in hort. Bogor.
subopposititifolia de Vriese, Mon. 23. 1853 — Ceylon. Peel Ins. Bonin-Sima.
sylhetensis de Vriese, Mon. 27. 1853. NPfl. 438 — India (Sylhet).
Teysmanniana de Vriese, Ned. Kr. Arch. 2^1. 228. 1850; Mon. 24 t. 1, 2. NPfl. 438 — Java.
uncinata de Vriese, Mon. 29 t. 3 f. 6. 1853 — Amboyna. N. Caledonia.
Wallichiana Pr. Suppl. 22. 1845; de Vriese, Mon. 27 — India (Nepal).
Wightiana de Vriese, Mon. 28. 1853 — India (Peninsula).
Willinkii Miquel; de Vriese, Mon. 21. 1853 — Java.

ANISOCAMPIUM Presl, Epim. Bot. 58. 1849 = **Dryopteris.**
Cumingianum Pr. Epim. 59. 1849 = D. otaria.

ANISOGONIUM Presl, Tent. Pterid. 115. 1836 = **Diplazium.**
alismaefolium Christ, Verh. Nat. Ges. Basel **11**. 233. 1895 = D. Cumingii.
attenuatum Pr. Tent. 116. 1836 = D. proliferum.
cordifolium Bedd. Ferns br. Ind. t. 331. 1870.
decussatum Pr. Tent. 116 t. 4 f. 13. 1836 = D. proliferum.
elegans Pr. Epim. 93. 1849 = D. fraxinifolium.
esculentum Pr. Tent. 116. 1836.
fraxinifolium Pr. Tent. 116 t. 4 f. 18. 1836.
grossum Pr. Epim. 93. 1849 = D. fraxinifolium.

heterophlebium Bedd. Ferns br. Ind. t. 329. 1869.
integrifolium Pr. Tent. 116. 1836 = D. cordifolium.
lineolatum Bedd. Ferns br. Ind. t. 330. 1869 = D. fraxinifolium
ovatum Fée, 1 mém. 14 t. 1 f. 24. 1844 = D. cordifolium.
pinnatifidum Pr. Tent. 116. 1836.
serampurense Pr. Tent. 116. 1836 = D. esculentum.
serrulatum Pr. Tent. 116. 1836 = D. manilense.
Smithianum Bedd. Ferns br. Ind. Suppl. 13. 1876.
sylvaticum Pr. Tent. 116. 1836 = D. silvaticum.
Zollingeri Pr. Epim. 92. 1849 = D. cordifolium.

ANISOSORUS Trevisan, Atti dell' Ist. Veneto II. 2. 166. 1851 = **Lonchitis.**

ANOGRAMMA Link, Fil. spec. 137. 1841 — (Polypodiaceae Gen. 59).

ascensionis (Hk.) Diels, Nat. Pfl. 1[4]. 258. 1899 — Ascension. — Grammitis Hk. Ic. pl. t. 967. 1854; Gymnogramme Hk. 1864. HB. 383. Chr. 66.

Biardii Fée, Cr. vasc. Br. **1**. 241 t. 77 f. 1. 1869 = Gymnogramma.

bifida Fée, 10 mém. 25. 1865 = ? Gymnogramma flexuosa.

chaerophylla (Desv.) Link, Fil. sp. 138. 1841. NPfl. 258 — Amer trop.
Gymnogramma Desv. Berl. Mag. **5**. 305. 1811. HB. 383. Chr. 65; Hemionitis Poir. 1813 — Anogramme villosa Fée 1850-52; Gymnogramma Moore 1858 (non Link 1833).

conspersa Fée, Gen. 184. 1850-52 = Gymnogramma argentea.

? davallioides Fée, Gen. 184. 1850-52 = Monachosorum subdigitatum.

Eggersii Christ in C. Chr. Ind. 58. 1905 — Cuba.
Gymnogramme Christ. Bull. Soc. bot. Belg. **33**[2]. 92. 1894.

Fauriei Christ in C. Chr. Ind. 58. 1905 — Japonia.
Gymnogramme Christ, Bull. Boiss. II. **2**. 561. 1902

flabellata Fée, Gen. 184. 1850-52 = Gymnogramma.

? hispidula Fée, Gen. 184. 1850-52 = Gymnogramma.

laserpitiifolia Fée, Gen. 184. 1820-52 = Gymnogramma hirta.

leptophylla (L.) Link, Fil. sp. 137. 1841. NPfl. 258 — Europa atlant. Reg. mediterr. Ins. Atlant. Abyssinia. Afr. austr. Madagascar. India. Australia. N. Zealand. Mexico-Argentina.
Polypodium L. sp. **2**. 1092. 1753; Asplenium Sw. 1791; Osmunda Sav. 1797; Acrostichum Lam. et DC. 1805; Grammitis Sw. 1806; Gymnogramma Desv. 1811. HB. 383. Chr. 65; Hemionitis Lag. 1816; Dicranodium Newm. 1854.

Lorentzii (Hier.) Diels, Nat. Pfl. 1[4]. 258. 1899 — Uruguay. Brasilia austr.
Gymnogramme Hier. Engl. Jahrb. **22**. 397. 1896 — G. Regnelliana Lindm. 1903 (non Christ 1900) = var. megaspora Lindm. Hedwigia **43**. 309 c. fig. 1904.

Makinoi (Maxim.) Christ in C. Chr. Ind. 58. 1905 — Japonia.
Gymnogramme Maxim.; Mak. Bot. Mag. Tokyo **8**. 481 c. tab. 1894.

microphylla (Hk.) Diels, Nat. Pfl. 1[4]. 259. 1899 — India bor. China.
Gymnogramme Hk. Ic. pl. t. 916. 1854. HB. 383. Chr. 66; Grammitis Bedd. 1866.

Ottonis Fée, Gen. 184. 1850-52 = Gymnogramma.

? paradoxa Fée, Gen. 184. 1850-52 = { Elaphoglossum dimorphum. / Elaphoglossum furcatum. }

petroselinifolia Fée, Gen. 184. 1850-52 = Gymnogramma hirta.

refracta Fée, Gen. 184 t. 19 A f. 2. 1850-52 = Gymnogramma flexuosa.

rosea Fée, Gen. 184. 1850-52 = Gymnogramma argentea.
Ruiziana Fée, Gen. 184. 1850-52 = Gymnogramma flexuosa.
schizophylla (Bak.) Diels, Nat. Pfl. 1^4. 258. 1899 = Cuba. Jamaica.
Gymnogramme Bak. JoB. **1877**. 266. Chr. 65; Psilogramme Und. 1902.
Schomburgkiana Fée, Gen. 184. 1850-52 = Gymnogramma.
villosa Fée, Gen. 184. 1850-52 = A. chaerophylla.

Anopodium J. Smith, Cat. cult. ferns 16. 1857 (*Phegopteris §*) = Dryopteris.

ANOPTERIS (Prantl) Diels, Nat. Pflanzenfam. 1^4. 288. 1899 — (Polypodiaceae Gen. **81**).
Cryptogramme § Anopteris Prantl, Engl. Jahrb. **3**. 414. 1882.
heterophylla Diels, Nat. Pfl. 1^4. 289. 1899 = A. hexagona.
hexagona (L.) C. Chr. Ind. 59. 1905 — Ind. occ. Brasilia.
Adiantum L. sp. **2**. 1097. 1753; Pteris heterophylla L. Syst. Nat. ed. X. **2**. 1322. 1759. HB. 160. Chr. 170; Allosorus Bernh. 1806; Cryoptogramme Prantl 1882; Onychium Kuhn 1897; Anopteris Diels 1899. NPfl. 289.

ANTIGRAMMA Presl, Tent. Pterid. 120. 1836 = **Phyllitis.**
brasiliensis Moore, Ind. LIII. 1857.
Douglasii Hk. Gen. ad t. 55A, 57A. 1840 = P. plantaginea.
Lageana Fée et Glaz.; Fée, Cr. vasc. Br. **1**. 264. 1869 = P. plantaginea.
lancifolia Pr. Tent. 120. 1836 = ? P. plantaginea.
nigripes J. Sm. Hist. Fil. 331. 1875.
oblongata Pr. Tent. 120. 1836 = P. brasiliensis.
plantaginea Pr. Tent. 120. 1836.
populifolia Pr. Tent. 120. 1836.
repanda Pr. Tent. 120 t. 4. f. 9-10. 1836 = P. brasiliensis.
rhizophylla J. Sm. JoB. **4**. 176. 1841 = Camptosorus r.
sibirica J. Sm. Hist. Fil. 331. 1875 = Camptosorus s.
subsessilis Fée, Gen. 210. 1850-52 = P. brasiliensis var.

ANTIOSORUS Roemer; Kuhn, Chaetopterides 347. 1882 = **Lonchitis.**
hirsutns Kuhn, Chaetopt. 347. 1882.
occidentalis Kuhn, Chaetopt. 347. 1882.

ANTROPHYUM Kaulfuss, Enum. Fil. 197. 1824 — (Polypodiaceae Gen. **91**).
alatum Brack. Expl. Exp. **16**. 64. 1854 = A. callifolium vel A. semicostatum.
anetioides Christ, Bull. Boiss II. **5**. 12. 1905 — Costa Rica.
angustatum Brack. Expl. Exp. **16**. 63. 1854 = A. plantagineum var.
avenium Bl. Fl. Javae Fil. 86 t. 37 f 2. 1828 = Polypodium Blumeanum var.
Boryanum (Willd.) Spr. Syst. **4**. 67. 1877. HB. 394 = Madagascar. Mascarenae.
Hemionitis Willd. sp. **5**. 128. 1810; Dictyogramme Trevis. 1877; Hemionitis reticulata Bory 1804 (non Forst. 1786).
Boryanum Bl. Fil. Javae Fil. 75 t. 31. 1828 = A. latifolium.
brasilianum (Desv.) C. Chr. Ind. 59. 1905 — Amer trop.
Hemionitis Desv. Prod. 216. 1827; Antrophyum subsessile Kze. Anal. 29 t. 19 f. 1. 1837. HB. 394. NPfl. 301. A. spathulatum Fée 1851-52.
Brookei Hk. 2 Cent. t. 79. 1861 = A. subfalcatum.
callifolium Bl. Enum. 111. 1828; Fl. Javae Fil. 83 t. 35 — Malesia-Polynesia.
Hemionitis Pr. 1836 — Antrophyum falcatum Bl. 1838; Hemionitis de Vriese 1846 — Antrophyum marginale Bl. 1828; Pteropsis Pr. 1836; Taenitis Moore 1857; Pteropsis Blumei Fée 1851-52; Taenitis Hk. 1864 — Antrophyum nanum Fée 1851-52. (HB. 393).
carnosum Liebm. Vid. Selsk. Skr. V. **1**. 161 (seors. 9). 1849 = A. ensiforme.

cayennense (Desv.) Spr. Syst. **4**. 67. 1827. HB. 394 — Puerto Rico. Amer. austr. trop. — Hemionitis (cajenensis) Desv. Berl. Mag. **5**. 311. 1811.

citrifolium Fée, 4 mém. 51. 1851-52 = Anetium.

coriaceum (Don) Wall. List n. 43. 1828. HB. 393. NPfl. 301 — India bor.-Malesia. — Hemionitis Don, Prod. Fl. Nep. 13. 1825. Antrophyum plicatum Fée 1851-52. Solenopteris lanceolata Wall. hb. (1823).

coriaceum Bl. Fl. Javae Fil. 85 t. 37 f. 1. 1828 = Polypodium Blumeanum var.

Cumingii Fée, 4 mém. 42 t. 4 f. 7. 1851-52; Trevis. Atti. Ist. Veneto II. **2**. 167. 1851 = A. Grevillei.

Desvauxii Moore, Ind. 80. 1858 — Ins. St. Thomas (Ind. occ.). — Hemionitis gigantea Desv. Prod. 216. 1827. (si sp. bona, hoc. nom. anteponendum est).

discoideum Kze. Bot. Zeit. **1848**. 702 = A. brasilianum.

discoideum Liebm. Vid. Selsk. Skr. V. **1**. 177 (seors. 25). 1849 = A. ensiforme.

D'Urvillaei Bory, Dup. Voy. Bot. **1**. 254. 1828 = A. semicostatum.

elongatum (Bl.) Fée, 4 mém. 43. 1851-52 — Java.
A. parvulum β elongatum Bl. Fl. Javae Fil. 78. 1828.

ensiforme Hk. in Benth. Fl. Hartw. 73. 1841. HB. 393. NPfl. 301 — Mexico. Guatemala. — Scoliosorus Moore 1857; Dictyogramme Trevis. 1877. Antrophyum falcatum Mart. et Gal. 1842 (non Bl. 1828); A. Galeottii Fée 1851-52.

falcatum Bl. Enum. 109. 1828 = A. callifolium.

falcatum Mart. et Gal. Mém. Ac. Brux. **15**. 49 t. 12. 1842 = A. ensiforme.

Feei Schaffn.; Fée, 7 mém. 42 t. 22 f. 1. 1857 — Mexico. Ecuador.

Galeottii Fée, 4 mém. 51 t. 5 f. 4. 1851-52 = A. ensiforme.

giganteum Bory, Bél. Voy. Bot. **2**. 36. 1833. HB. 394. Chr. 59. NPfl. 301 — Ins. Mascarenae. — Dictyogramme Trevis. 1877. (nom. malum; conf. A. Desvauxii).

Grevillei Balf.; Grev. Ann. Mag. Nat. Hist. II. **2**. 10 t. 1. 1848 — Ins. Philipp. Polynesia. — Polytaenium Moore 1857. Antrophyum Cumingii Fée 1851-52. HB. 393.

Hookerianum Kze. Bot. Zeit. **1848**. 198; Fée, 4 mém. 46. 1851-52 = A. immersum.

immersum (Bory) Mett. Ann. Lugd. Bat. **4**. 171. 1868-69. HB. 393 — Ins. Mascarenae. Malesia. — Hemionitis Bory; Willd. sp. **5**. 127. 1810. Antrophyum pumilum Klf. 1824.

involutum Bl. Fl. Javae Fil. 87. 1828 = Polypodium scolopendrinum.

japonicum Mak. Phan. Pterid. Jap. Ic. **1**. t. 26. 1899 — Japonia.

Johannis Trevis. Atti Ist. Veneto II. **2**. 167. 1851 = A. callifolium (nanum).

lanceolatum (L.) Klf. Enum. 198. 1824. HB. 394. Chr. 58. NPfl. 301 — Amer. trop. Hemionitis L. sp. **2**. 1077. 1753; Dictyogramme Trevis. 1877.

lanceolatum Bl. Fl. Javae Fil. 84 t. 36. 1828 = Polypodium loxogramme.

latifolium Bl. Fl. Javae Fil. 75. 1828. HB. 394. NPfl. 301 — China austr. Ind. bor.-Malesia. — Hemionitis Keys. 1873; Dictyogramme Trevis. 1877; Hemionitis Boryana Bl. Enum. 111. 1828 (non Willd 1810); Antrophyum Bl. Fl. Javae Fil. 75 t. 31. 1828; Hemionitis Blumeana Pr. 1836 — Antrophyum obovatum Bak. 1898.

latipes Kze. Bot. Zeit. **1848**. 209 = A. semicostatum.

Lessonii Bory, Dup. Voy. Bot. **1**. 255 t. 28 f. 2. 1828 = A. plantagineum var.

lineatum (Sw.) Klf. Enum. 199. 1824. HB. 392. Chr. 57. NPfl. 301 — Amer. trop. Hemionitis Sw. Prod. 129. 1788; Loxogramme Pr. 1836; Polytaenium J. Sm. 1841; Vittaria lanceolata Sw. 1799; Polytaenium Desv. 1827.

Mannianum Hk. 2 Cent. t. 73. 1861. HB. 394. Chr. 59. NPfl. 301 — Afr. trop. Dictyogramme Trevis. 1877. Antrophyum Meniscium Mett. 1868.

marginale Bl. Enum. 111. 1828 = A. callifolium.
Meniscium Mett.; Kuhn, Fil. Afr. 56. 1868 = A. Mannianum.
minimum Bak. Ann. Bot. **5**. 488. 1891 — Costa Rica.
nanum Fée, 4 mém. 44. 1851-52 = A. callifolium.
niphoboloides Kze. Bot. Zeit. **1848**. 209 = Cyclophorus flocciger.
obovatum Bak. Kew Bull. **1898**. 233 = A. latifolium var.
Obtusatum Bory, Dup. Voy. Bot. **1**. 256 t. 29 f. 2. 1828 = A. obtusum.
obtusum (Bory); Spr. Syst. **4**. 67. 1827. (HB. 394) — Madagascar. Ins. Mascarenae.
Hemionitis Bory; Willd. sp. **5**. 127. 1810; Dictyogramme Trevis. 1877.
obtusum Bl. Enum. 110. 1828 = A. immersum.
parvulum Bl. Enum. 110. 1828; Fl. Javae Fil. 78 t. 34 f. 3 (HB. 393) — Java. — Hemionitis Pr. 1836.
pendulum Leprieur; Fée, 4 mém. 51. 1851-52 = Anetium citrifolium.
petiolatum Bak.; Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 202 (nomen) — China.
plantagineum (Cav.) Klf. Enum. 197. 1824. HB. 393. Chr. 57. NPfl. 301 — India bor. Ceylon. Malesia. Polynesia.
Hemionitis Cav. Descr. 260. 1802 — Antrophyum Lessonii Bory 1828; Hemionitis Pr. 1836. Antrophyum angustatum Brack. 1854.
plicatum Fée, 4 mém. 44 t. 5 f. 1. 1851-52 = A. coriaceum.
pumilum Klf. Enum. 197. 1824 = A. immersum.
reticulatum (Forst.) Klf. Enum. 198. 1824. HB. 393. Chr. 57. NPfl. 301 — India bor.-Malesia-Polynesia. Austr. trop. Madagascar.
Hemionitis Forst. Prod. 79. 1786.
semicostatum Bl. Enum. 110. 1828; Fl. Javae Fil. 77 t. 33. HB. 393. Chr. 58. NPfl. 301 — Ceylon. Perak. Malacca. Malesia-Polynesia.
Hemionitis Pr. 1836. Antrophyum D'Urvillaei Bory 1828 — A. latipes Kze. 1848; Dictyogramme Trevis. 1877 — (an Hemionitis rigida Cav. 1802?).
sessilifolium (Cav.) Spr. **4**. 67. 1827 — Ins. Philipp.
Hemionitis Cav. Descr. 261. 1802.
spathulatum Fée, 4 mém. 46 t. 4 f. 6. 1851-52 = A. brasilianum.
Sprucei »Moore«; Salom. Nom. 40. 1883 = Anetium citrifolium.
stenophyllum Bak. Kew Bull. **1898**. 233 — Yunnan.
strictum Mett. Ann. Lugd. Bat. **4**. 171 t. 7 f. 1-3. 1868-69 — N. Guinea.
subfalcatum Brack. Expl. Exp. **16**. 65. 1854. HB. 392 — Borneo. Celebes. Polynesia. — A. Brookei Hk. 1861.
subsessile Kze. Anal. 29 t. 19 f. 1. 1837 = A. brasilianum.
vittarioides Bak. JoB. **1890**. 267. NPfl. 301 — Tonkin.
Werckleanum Christ, Bull. Boiss. II. **5**. 11. 1905 — Costa Rica.
zosteraefolium Fée, 4 mém. 52. 1851-52 — India. — Hemionitis falcata Willd. sp. **5**. 126. 1810. (si sp. bona, hoc nom. anteponendum est).

APALOPHLEBIA Presl, Epim. Bot. 137. 1849 = **Cyclophorus.**
costata Pr. Epim. 138. 1849 = C. Beddomeanus.
?flocculosa Pr. Epim. 139. 1849.
splendens Pr. Epim. 138. 1849.
venosa Pr. Epim. 139. 1849 = C. stigmosus.

APHYLLOCALPA Cavanilles, Anal. Cienc. **5**. 164. 1802; Descr. 556. 1803 = [**Osmunda.**
regalis Cav. Anal. Cienc. **5**. 164. 1802; Descr. 556. 1803

Apotomia Fée. Gen. Fil. 112. 1850-52 *(Adiantum §)* = Adiantum.

AQUILINA Presl msc. (Ettingsh. Farnkr. 91. 1865) = **Pteridium.**

ARACHNIODES Blume. Enum. Pl. Javae 241. 1828 = **Polystichum.**
aspidioides Bl. Enum. 242. 1828 = P. aristatum.
ARCASPLENIUM Moore msc. (Baker. Kew Bull. **1901**. 145) = **Asplenium.**
grande Moore msc. = A. Sanderi.
ARCHANGIOPTERIS Christ et Giesenhagen, Flora **1899**. 72. 77 — (Marattiaceae Gen. **2**).
Henryi Christ et Gies. Flora **1899**. 78. NPfl. 439 — Yunnan.
ARCYPTERIS Underwood, Bull. Torr. Cl. **30**. 678. 1903 = **Aspidium.**
difformis Und. Bull. Torr. Cl. **30**. 678. 1903 = A. irregulare.
Argyria Fée, Gen. Fil. 183. 1850-52 (*Ceropteris §*) = Ceropteris.
Argyrochosma J. Smith, Journ. Bot. **4**. 50. 1841 (*Notholaena §*) = Pellaea.
Aristaria K. Müller, Bot. Zeit. **1854**. 545 (*Vittaria §*) = Vittaria.
Arsenopteris Webb et Berthelot, Hist. Nat. Canar. **3**². pars 3. 437. 1847 (*Aspidium §*) = Dryopteris.
ARTHROBOTRYA J. Smith, Hist. Fil. 141. 1875 = **Polybotrya.**
articulata J. Sm. Hist. Fil. 142. 1875.
ARTHROBOTRYS Wallich, List n. 395. 1828 = **Dryopteris.**
Avara Wall. List n. 1034. 1828 } = D. cochleata.
macrocarpa Wall. List n. 395. 1828 } = D. cochleata.
Arthrodanaea Presl, Suppl. Tent. Pter. 37. 1845 (*Danaea §*) = Danaea.
Arthrolygodes Presl, Suppl. Tent. Pter. 101. 1845 (*Lygodium §*) = Lygodium.
ARTHROMERIS (Moore) J. Smith, Hist. Fil. 110. 1875 = **Polypodium.**
Pleopeltis § Arthromeris Moore, Ind. Fil. LXXVIII. 1857.
juglandifolia J. Sm. Hist. Fil. 111. 1875 = P. Wallichianum.
venusta J. Sm. Hist. Fil. 111. 1875 = P. himalayense.
ARTHROPTERIS J. Smith in Hook. fil. Fl. N. Zeal. **2**. 43. 1854 — (Polypodiaceae Gen. **26**).
albopunctata J. Sm. Cat. cult. ferns 62. 1857 — Dryopteris orientalis.
altescandens (Colla) J. Sm. Hist. Fil. 225. 1875. NPfl. 205 — Juan Fernandez.
Polypodium Colla, Mem. Ac. Torino **39**. 48. 1836. Mett. Pol. n. 90; Nephrolepis Bak. 1867. HB. 301. (Chr. 288; vix Luerss. Fil. Graeff. 203); Polypodium procurrens Kze. 1837.
Beckleri Mett. Novara Exp. Bot. **1**. 213. 1870 = A. obliterata.
filipes Moore, Gard. Chr. **1855**. 368 = A. tenella stat. juv.
heteromorpha J. Sm. Hist. Fil. 225. 1875 = Polypodium.
obliterata (R. Br.) J. Sm. Cat. cult. ferns 62. 1827 — Asia, Australia, Polynesia, Africa trop.
Nephrodium R. Br. Prod. Fl. N. Holl. 148. 1810; Aspidium Spr. 1827; Nephrolepis J. Sm. 1841. Aspidium Palisoti Desv. 1811; A. ramosum Pal. Beauv. post 1811; Nephrolepis Moore 1858. HB. 301. Chr. 288; Arthropteris Mett. 1870. NPfl. 205 — Nephrolepis trichomanoides J. Sm. 1841; Lepidonevron Fée 1850-52 — Polypodium Beckleri Hk. 1862; Arthropteris Mett. 1870.
pectinata Kuhn in Schum. Fl. Kais. Wilh. Land 10. 1889 = Dryopteris orientalis.
ramosa Mett. Novara Exp. Bot. **1**. 213. 1876 = A. obliterata.
subbiaurita J. Sm. Hist. Fil. 225. 1875 = Dryopteris orientalis.
tenella (Forst.) J. Sm. in Hk. fil. Fl. N. Zeal. **2**. 43. t. 82. 1854. NPfl. 205 — Austr. N. Zealand. N. Caledonia.
Polypodium Forst. Prod. 81. 1786. Mett. Pol. n. 89. HB. 337. Chr. 84.

ASPIDIUM Swartz, Schrad. Journ. **1800**[2]. 29. 1801 — (Polypodiaceae Gen. **15**).

Subgenera: Ar. = **Arcypteris** (Dictyopteris.)
Pl. = **Pleocnemia.**
S. = **Sagenia.**
T. = **Tectaria** (Euaspidium).

abbreviatum Poir. Enc. Suppl. **4**. 516. 1816 = Dryopteris filix mas.
abbreviatum Schrad. Gött. gel. Anz. **1824**. 869 = Polystichum.
abortivum Bl. Enum. 154. 1828 = Dryopteris.
abruptum Bl. Enum. 154. 1828 = Dryopteris truncata.
abruptum Kze. Linn. **9**. 93. 1834 = Dryopteris Kunzeana.
abruptum Mart. et Gal. Mém. Ac. Brux. **15**. 65. 1842 = Dryopteris patens.
abyssinicum Döll, Gefässkr. Grossh. Badens 23. 1855 = Atthyrium Schimperi.
acanthophyllum Franch. Bull. Soc. Fr. **32**. 28. 1885 = Polystichum.
achalense Hier. Engl. Jahrb. **22**. 371. 1896 = Dryopteris.
acrocarpon Fée, 11 mém. 84. 1866 = Dryopteris protensa.
acrostichoides Sw. Syn. 44. 1806 = Polystichum.
aculeatum Sw. Schrad. Journ. **1800**[2]. 37. 1801 = Polystichum.
aculeatum × lonchitis Haláczy, Consp. Fl. Graec. **3**. 475. 1904 = Polystichum.
acuminatum Willd. sp. **5**. 221. 1810 = Nephrolepis biserrata.
acuminatum Lowe, Ferns **6**. t. 11. 1857 = Dryopteris decomposita.
(S) **acutilobum** Hier. Engl. Jahrb. **34**. 450. 1904 — Columbia.
acutum Schkuhr. Kr. Gew. **1**. 32. t. 31. 1806 = Nephrolepis biserrata.
acutum Fée, Cr. vasc. Br. **2**. 76. 1872-73 = Dryopteris.
Adenopteris Mett. Pheg. u. Asp. 107. n. 255. 1858 = Dryopteris pennigera.
adiantoides Bl. Enum. 145. 1828 = Lindsaya.
adnatum Bl. Enum. 162. 1828 = Dryopteris filix mas var.
adscendens Hk. sp. **4**. 32 t. 224. 1862 vide ascendens.
adultum Wikstr. Vet. Ak. Handl. **1825**. 437. 1826 = Dryopteris parasitica.
aduncum Wall. List n. 384. 1828 = Dryopteris hirtipes.
aemulum Sw. Schrad. Journ. **1800**[2]. 42. 1801 = Dryopteris.
aesculifolium Bl. Enum. 143. 1828 = Kaulfussia.
affine Bl. Enum. 148. 1828 = Dryopteris lineata.
affine Wall. List n. 370. 1828 = Polystichum speciosum.
affine Lowe. JoB. **1**. 26. 1834 (nomen); Tr. Cambr. Phil. Soc. 525. 1838 = Dryopteris filix mas.
affine Fisch. et Mey. Bull. Soc. nat. Moscou **1838**. 240 = Dryopteris filix mas.
affine Reichb.: Vis. Fl. Dalm. **1**. 39. 1842 = Dryopteris rigida *australis.
agatolepis Fée, 8 mém. 106. 1857 = Dryopteris patula.
alatellum Christ, Bull. Boiss. II. **1**. 454. 1901 = Dryopteris.
alatum Wall. List n. 378. 1828 = A. vastum.
alatum Guill.; Brack. Expl. Exp. **16**. 179. 1854 = A. decurrens.
albescens Mett.; Salom. Nom. 42. 1883 = Nephrodium.
albicaule Fée, 8 mém. 102. 1857 = Dryopteris patens.
albolineatum Boivin; Salom. Nom. 42. 1883 = Dryopteris mauritiana.
albopunctatum Bory; Willd. sp. **5**. 242. 1810 = Dryopteris orientalis.
alcicorne Christ. Bull. Soc. bot. It. **1901**. 295 = Polystichum.
Alfarii Christ, Bull. Boiss. II. **5**. 259. 1905 = Dryopteris.
alienum Mett.: Kuhn. Linn. **36**. 120. 1869 = A. variolosum.
alpestre Hoppe. Neu. Taschenb. **1805**. 216 = Athyrium.

alpinum Sw. Schrad. Journ. **1800**[2]. 42. 1801 = Cystopteris fragilis *regia.
alsophilaceum Kze. Mett. Pheg. u. Asp. 90. n. 218. 1858 } = Dryopteris.
alsophileum Kze. Flora **1837**[1]. 326 (nomen) } tenuifolia.
amabile Bl. Enum. 165. 1828 = Polystichum.
amaurolepis Fée, Cr. vasc. Br. **1**. 137 t. 44 f. 2. 1869 = Dryopteris submarginalis.
ambiguum (Pr.) Diels, Nat. Pfl. **1**[4]. 188. 1899 — Ins. Philipp.
Digrammaria Pr. Tent. 117 t. 4 f. 12. 1836; Gymnogramme Hk. 1864. HB. 378; G. philippinensis Fée 1850-52; Phegopteris Mett. n. 55. var. 1. 1858. Heterogonium aspidioides Pr. 1849.
amblyotus Kze. Bot. Zeit. **1848**. 283 = Polystichum aculeatum var. 32.
amboinense Willd. sp. **5**. 228. 1810 = Dryopteris parasitica*.
americanum Torrey, Cat. Pl. N. York 80. 1818 = Dryopteris Goldiana.
americanum Dav. Amer. Nat. **12**. 714. 1878 = Dryopteris spinulosa *intermedia.
ameristonevron Fée, 8 mém. 104. 1857 = Dryopteris.
ammifolium Desv. Berl. Mag. **5**. 321. 1811 = Polystichum aculeatum var. 1.
ammifolium Mett.; Salom. Nom. 42. 1883 = Dryopteris cubensis.
amplificatum Fée; Fourn. Mex. pl. **1**. 94. 1872 = Dryopteris.
amplissimum Mett. Pheg. u. Asp. 68 n. 160. 1858 = Dryopteris.
amplum Mett. Pheg. u. Asp. 74 n. 176. 1858 = Dryopteris.
amplum Kuhn, Fil. Afr. 125. 1868 = Dryopteris inaequalis.
anateinophlebium Kuhn, v. Deck. Reis. **3**[3]. Bot. 65. 1879 = Dryopteris.
(Ar) **andaiense** (Bak.) C. Chr. Ind. 64. 1905 — N. Guinea.
Polypodium Bak. Malesia **3**. 45. 1886.
(S) **andinum** (Bak.) C. Chr. Ind. 64. 1905 — Peru.
Nephrodium Bak. Ann. Bot. **5**. 331. 1891.
aneitense Fourn. Ann. sc. nat. V. **18**. 297. 1873 = Dryopteris Lenormandi.
(Ar) **angelicifolium** (Schum.) C. Chr. Ind. 64. 1905 — Guinea.
Polypodium Schum. Vid. Selsk. Afh. **4**. 228 (seors. 454). 1827; Phegopteris Mett.; Kuhn 1868.
angulare Kit.; Willd. sp. **5**. 257. 1810 = Polystichum aculeatum et Braunii.
(T) **angulatum** (Willd.) J. Sm.; Mett. Ann. Lugd. Bat. **1**. 239. 1864. HB. 258. Chr. 228. NPfl. 185 — Malesia. N. Guinea.
Polypodium Willd. sp. **5**. 185. 1810; Dryopteris O. Ktze. 1891. Aspidium sanctum Bl. 1828 (non Mett. 1858).
angulatum Hk. sp. **4**. 44. 1862 = A. platanifolium.
angustifolium Mett.; Salom. Nom. 43. 1883 = Dryopteris hispidula.
angustifrons Miq. Ann. Lugd. Bat. **3**. 178. 1867 = ? Dryopteris gracilescens.
angustum Willd. sp. **5**. 277. 1810 = Athyrium filix femina.
anomalum Hk. sp. **4**. 27. 1862 = Polystichum aculeatum var. 37.
anomophyllum Zenker, Pl. Ind. **1**. t. 1. 1835 = Polystichum falcatum var.
anopteron Kze.; Kuhn, Linn. **36**. 113. 1869 = Dryopteris.
antarcticum Fourn. Compt. rend. **81**. 1140. 1875 = Dryopteris filix mas var.
(S) **antioquianum** (Bak.) C. Chr. Ind. 64. 1905 — Columbia.
Nephrodium Bak. JoB. **1881**. 205.
apertum Fée, 8 mém. 106. 1857 = Dryopteris.
apicale Benth. Fl. austr. **7**. 758. 1878 = Dryopteris.
apiciflorum Wall. List n. 345. 1828 = Dryopteris.
apifolium Schkuhr, Kr. Gew. **1**. 198 t. 56 b. 1809 = A. cicutarium*.
apiifolium Mett.; Kuhn, Bot. Zeit. **1869**. 165 = Polystichum.
appendiculatum Bl. Enum. 165. 1828 = Polystichum aristatum.

appendiculatum Wall. List n. 349. 1828 = Dryopteris cana et prolixa.
appendiculatum Christ, Bull. Soc. Fr. **52** Mém. I. 36. 1905 = Dryopteris squamaestipes var.
aquapimense Schum. Vid. Selsk. Afh. **4**. 230 (seors. 456). 1827 — Guinea.
aquilinoides Bolle, Zeitschr. Ges. f. Erdk. **1**. 221. 1866 = Dryopteris filix mas *oligodonta.
aquilinoides Mett.; Kuhn, Fil. Afr. 126. 1868 = Dryopteris.
arborescens Luerss. Fil. Graeff. 176. 1871 = Dryopteris ludens.
arbuscula Willd. sp. **5**. 233. 1810 = Dryopteris.
Araguata (Moritz) Reichardt, Denkschr. Ak. Wien **17**². 38 t. 2 f. 28-31. 1859 — Caracas. — Polypodium Araguato Moritz 1854 (nomen).
arcuatum Desv. Berl. Mag. **5**. 321. 1811 = Dryopteris serra.
Arechavaletae Hier. Engl. Jahrb. **22**. 370. 1896 = Dryopteris.
argentinum Hier. Engl. Jahrb. **22**. 367. 1896 = Dryopteris.
argutum Klf. Enum. 242. 1824 = Dryopteris rigida*.
aridum Don, Prod. Fl. Nepal. 4. 1825 = Dryopteris.
aristatum Sw. Schrad. Journ. **1800**². 37. 1801 = Polystichum.
articulatum Sw. Schrad. Journ. **1800**². 30. 1801 (pt.); Willd. sp. **5**. 212. 1810 = Oleandra.
articulatum Schkuhr, Kr. Gew. **1**. 28 t. 27. 1806 = Oleandra nodosa.
articulatum Sieb. Syn. Fil. exsicc. — = Nephrolepis biserrata.
articulatum Lowe, Ferns **6**. t. 49. 1857 = Dryopteris megaphylla.
(Pl) **artinexum** (Clarke) C. Chr. Ind. 65. 1905 — Sikkim.
Nephrodium Clarke 1880; Pleocnemia Clarkei Bedd. Ferns br. Ind. Suppl. 15 t. 368. 1876 (non Nephrodium Bak. 1874).
ascendens Hew. Mag. Nat. Hist. II. **2**. 463. 1838 = Polystichum apiifolium.
Ascencionis Kuhn, Fil. Afr. 126. 1868 = Dryopteris.
asperulum Fée, 11 mém. 80 t. 23 f. 1. 1866 = Dryopteris decumbens.
asperum Gray, Nat. Arr. br. Pl. **2** —. 1821 = Polystichum lonchitis.
asperum Mett. Ann. Lugd. Bat. **1**. 235. 1864 = Dryopteris urophylla.
asplenioides Sw. Syn. 60. 1806 = Athyrium filix femina.
asplenioides Gris. Fl. br. W. Ind. 693. 1864 = Dryopteris.
assamicum Kuhn, Linn. **36**. 108. 1869 = Polystichum aristatum var.
asterolepis Trevis. Hedwigia **13**. 145. 1874 = Polystichum vestitum.
athamanticum Kze. Linn. **18**. 133. 1844 = Dryopteris.
athyrioides Mart. et Gal. Mém. Ac. Brux. **15**. 67 t. 18. 1842 = Dryopteris.
Atkinsoni Clarke, Tr. Linn. Soc. II. Bot. **1**. 506. 1880 = Polystichum.
atomarium Mühl.: Willd. sp. **5**. 279. 1810 = Cystopteris bulbifera.
atratum Wall. List n. 380. 1828 = Dryopteris hirtipes.
atrorubens Mett.; Kuhn, Linn. **36**. 112. 1869 = Dryopteris.
attenuatum Sw. Schrad. Journ. **1800**². 34. 1801 = Dryopteris dissecta.
attenuatum Kze.; Mett. Pheg. u. Asp. 96. n. 233. 1858 = Dryopteris stenobasis.
augescens Link, Fil. sp. 103. 1841 = Dryopteris serra.
aureo-vestitum Gris. Fl. br. W. Ind. 692. 1864 = Dryopteris vellea.
auriculatum Sw. Schrad. Journ. **1800**². 31. 1801 = Polystichum.
auriculatum Schkuhr, Kr. Gew. **1**. 31 t. 30. 1806 = Polystichum acrostichoides.
auriculatum Wall. List n. 2233. 1829 = Nephrolepis cordifolia.
auritum Bak. Ann. Bot. **5**. 313. 1891 = Phanerophlebia.
auritum Christ. Bull. Boiss. II. **4**. 616. 1904 = Dryopteris.
axillare Sw. Schrad. Journ. **1800**². 42. 1801 = Athyrium filix femina*.

Bakerianum Atkins.; Bak. Ic. pl. t. 1656. 1886 = Polystichum.
(S) **Balansae** C. Chr. Ind. 66. 1905 — Tonkin. — Nephrodium stenopteron Bak. JoB. **1890**. 265 (non Aspidium stenopteris Kze. 1849).
Balbisii Kuhn, JoB. **1877**. 231 = Dryopteris.
bantamense Bl. Enum. 141. 1828 = Oleandra neriiformis.
(Ar) **Barberi** (Hk.) C. Chr. Ind. 66. 1905 — Malesia; Ins. Philipp.
Polypodium Hk. sp. **5**. 100. 1864. HB. 317; Dictyopteris Bedd. 1869; Phegopteris Mett.; Kuhn 1869.
barbigerum Christ, Farnkr. 260. 1897 = Dryopteris.
(S) **Barclayi** (Carr.) C. Chr. Ind. 66. 1905 — Fiji. — Lastrea Carr. in Seem. Fl. Vit. 359. 1873; Nephrodium Bak. 1874. HB. 504. Dryopteris O. Ktze. 1891.
Baromez Willd. sp. **5**. 268. 1810 = Cibotium barometz.
(S) **Barteri** J. Sm. Ferns br. and for. 286. 1866; Mett; Kuhn, Fil. Afr. 126. 1868. NPfl. 185 — Africa occ. trop.
Nephrodium Bak. 1867. HB. 299; Dryopteris O. Ktze. 1891.
basiattenuatum Jenm. Gard. Chr. III. **17**. 132. 1895 = Dryopteris.
basilare Fée, Cr. vasc. Br. **1**. 135 t. 43 f. 2. 1869 — Bahia. (Dryopteris sp. dubia).
basipinnatum Bak. JoB. **1889**. 176 = Polystichum.
(Ar) **Beccarianum** (Ces.) Diels, Nat. Pfl. **1**[4]. 186. 1899 — N. Guinea.
Polypodium Cesati, Rend. Ac. sci. Napoli **16**. 27, 30. 1877.
Beccarianum Bak. Malesia **3**. 43. 1886 = A. Cesatianum.
Beddomei Prantl, Verh. zool. bot. Ges. **31**. 119. 1882 = Dryopteris.
Benoitianum Gaud. Freyc. Voy. Bot. (331) t. 11. 1827 = Polystichum.
Bergianum Mett. Pheg. u. Asp. 79. n. 188. 1858 = Dryopteris.
Berterianum Colla, Mem. Ac. Torino **39**. 42 t. 70. 1836 = Polystichum.
Berteroanum Fée, 11 mém. 72 t. 22 f. 1. 1866 = Dryopteris opposita.
biaristatum Bl. Enum. 164. 1828 = Polystichum aculeatum var. 30.
Bicknellii Christ in Burnat, Matér. p. fl. Alp. Marit. —. 1900 = Polystichum aculeatum × lobatum.
bidentatum Spr. Syst. **4**. 99. 1827 = Nephrolepis.
bifidum Carm. Tr. Linn. Soc. **12**. 511. 1818 = Dryopteris tomentosa.
bifidum Pr. Del. Prag. **1**. 173. 1822 = A. martinicense.
biforme Fée, Cr. vasc. Br. **1**. 139. 1869 = Dryopteris amplissima.
Biolleyi Christ, Prim. Fl. Costaric. **3**. 31. 1901 = Dryopteris.
biserratum Sw. Schrad. Journ. **1800**[2]. 32. 1801 = Nephrolepis.
Blumei Kze.; Mett. Fil. Lips. 94 t. 22 f. 5. 1856 = Dryopteris canescens.
Blumei Thwait. En. pl. Zeyl. 392. 1864 = Dryopteris rhodolepis.
Bodinieri Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 248 c. fig. = Dryopteris.
Boivini Kuhn, v. Deck. Reis. **3**[3]. Bot. 65. 1879 = Dryopteris.
Boottii Tuckerm. Hovey's Mag. **9**. 145. 1843 = ? Dryopteris cristata × spinulosa.
Boryanum Willd. sp. **5**. 285. 1810 = Dryopteris.
Boryanum Boj. Hort. Maur. 391. 1837; Kuhn, Fil. Afr. 127 (pt.). 1868 = Dryopteris filix mas *elongata.
bosniacum Formánek, Oest. Bot. Zeit. **38**. 243. 1888 = Polystichum aculeatum.
Bourgaei Fourn. Mex. Pl. **1**. 98. 1872 = Dryopteris.
Boutonianum Hk. Ic. pl. t. 931. 1854 = Dryopteris orientalis.
Boydiae Eat. Bull. Torr. Cl. **6**. 361. 1879 = Dryopteris cyatheoides var.
(S) **brachiatum** Zoll. et Moritz; Nat. Geneesk. Arch. **1**. 399. 1844 — Java.
A. Zollingerianum Kze. 1846; Nephrodium Bak. 1867. HB. 298 (pt.); Dryopteris O. Ktze. 1891. Phlebiogonium impressum Fée 1850-52 Aspidium Hk. 1862.

brachyneuron Fée, Cr. vasc. Br. **1**. 133. 1869 — Bahia. — (Dryopteris sp. dub.).
brachyodon Gris. Fl. br. W. Ind. 693. 1864 = Dryopteris.
brachypterum Kze. Linn. **24**. 288. 1851 = Polystichum lobatum var. 4.
Brackenridgei Mett. Ann. sc. nat. IV. **15**. 75. 1861 = Dryopteris.
brasilianum Pr. Del. Prag. **1**. 176. 1822 = Hypolepis.
Braunianum Karst. Fl. Col. **1**. 63 t. 31. 1859 = Dryopteris.
Braunii Spenn. Fl. Frib. **1**. 9 t. 2. 1825 = Polystichum.
Breutelianum Mett. Fil. Lips. 85 t. 18 f. 12. 1856 } = Dryopteris limbata.
Breutelii Mett. Pheg. u. Asp. 85. n. 204. 1858 } = Dryopteris limbata.
Bridgesii Sturm, En. pl. vasc. cr. chil. —. 1858 = Polystichum aculeatum var. 11.
Brongniartianum Sturm, En. pl. vasc. cr. chil. —. 1858 = Polystichum aculeatum var. 12.
(Ar) **Brongniartii** (Bory) Diels, Nat. Pfl. **1**[4]. 186. 1899 — Malesia-Ins. Philipp. Polypodium Bory, Dup. Voy. Bot. **1**. 263 t. 34. 1828. HB. 318; Phegopteris Mett. n. 66. 1858. Polypodium pteroides Pr. 1825 (non Retz. 1791); Dictyopteris Pr. 1836.
Brunonianum Wall. List n. 344. 1828 = Dryopteris.
Buchholzii Kuhn, v. Deck. Reisen **3**[3]. Bot. 47. 1879 = Dryopteris.
bulbiferum Sw. Schrad. Journ. **1800**[2]. 41. 1801 = Cystopteris.
(Pl) **Cadieri** Christ, Journ. de Bot. **19**. 62. 1905 — Annam.
caducum HBK. Nov. Gen. et Sp. **1**. 12. 1815 = Cyclopeltis semicordata.
caducum Wall. List n. 381. 1828 = Polystichum Hookerianum.
caespitosum Wall. List n. 367. 1828 = Polystichum obliquum.
calcaratum Bl. Enum. 159. 1828 = Dryopteris.
(T) **calcareum** (J. Sm.) Pr. Epim. 63. 1849. Mett. n. 283. HB. 259 — Ins. Philipp. Sagenia J. Sm. 1841 (nomen); Dryopteris O. Ktze. 1891.
calcareum Baumg. En. stirp. Transs. **4**. 29. 1846 = Dryopteris Robertiana.
calcareum? Ces. Rend. Ac. sci. Napoli **16**. 26. 1877 = A. Cesatianum.
calcigenum Fée, 10 mém. 38. 1865 = Dryopteris orientalis.
californicum Eat. Pr. Amer. Acad. **6**. 555. 1865 = Polystichum.
Callipteris Wilms, Verh. Rhein. Ges. **9**. 577. 1852 = Dryopteris spinulosa.
callopsis Franch. et Sav. En. pl. Jap. **2**. 240, 634. 1879 = Dryopteris.
callosum Bl. Enum. 152. 1828 = Dryopteris.
calyptratum Desv. Prod. 251. 1827 = Polystichum adiantiforme.
Cameroonianum Mett.; Kuhn, Fil. Afr. 127. 1868 = A. varians.
campylopterum Kze. Am. Journ. Sci. II. **6**. 83, 84. 1848 = Dryopteris spinulosa *dilatata.
canariense A. Br. Flora **1841**. 708 = Dryopteris filix mas *oligodonta.
canescens Wall. List n. 354. 1828 = Dryopteris parasitica.
canescens Kze. Bot. Zeit. **1848**. 556 = Dryopteris unita.
canescens Christ, Ann. Jard. Buit. **15**. 130. 1897 = Dryopteris.
canum Wall. List n. 387. 1828 = Dryopteris calcarata et didymosora.
capense Sw. Schrad. Journ. **1800**[2]. 42. 1801 = Hemitelia.
capense Willd. sp. **5**. 267. 1810 = Polystichum adiantiforme.
capense Desv. Prod. 250. 1827 = Polystichum pungens.
capillipes Bak. JoB. **1888**. 228 = Polystichum.
Capitainei Fée et L'Herm.; Fée, 11 mém. 80 t. 22 f. 3. 1866 = Dryopteris trichophora.
caripense Mett. Fil. Lips. 90 t. 18 f. 9. 1856 = Dryopteris submarginalis.
caruifolium Bak. JoB. **1888**. 228 = Polystichum omeiense.
carvifolium Kze. Linn. **24**. 292. 1851 = Polystichum.

caryotideum Wall. List n. 376. 1828 = Polystichum falcatum var.
catacolobum Kze.; Ettingsh. Farnkr. 182 t. 126 f. 9, 10. 1865 = Dryopteris [anoptera.
catocarpum Kze. Linn. **9**. 95. 1834 = Dryopteris.
catophoron Kze. Bot. Zeit. **1848**. 262 = Dryopteris sparsa.
catopteron Kze. Linn. **10**. 550. 1836 = Dryopteris lanuginosa.
caucasicum A.Br. Flora **1841**. 707 = Dryopteris filix mas.
caudatum Sw. Syn. 55. 1806 = Athyrium umbrosum.
caudatum Sw. Vet. Ak. Hdl. **1817**. 62 = Polystichum aculeatum var. 25.
caudatum Jenm. JoB. **1879**. 260 = Polystichum viviparum.
caudiculatum Sieb.; Kze. Linn. **24**. 280. 1851 = Dryopteris truncata.
caudiculatum Mett.; Salom. Nom. 46. 1883 = Dryopteris immersa.
Cavalerii Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 116 = Dryopteris.
(T) **Cesatianum** C. Chr. Ind. 68. 1905 — N. Guinea. — A. Beccarianum Bak. Malesia **3**. 43. 1886 (non supra); A. calcareum? Ces. 1877.
chaerophyllaefolium Zippel; Mett. Ann. Lugd. Bat. **1**. 228. 1864 = Dryopteris.
chaerophylloides Moritz (Mett. Pheg. u. Asp. 65. 1858. syn.) = Dryopteris patula var.
Championi Benth. Fl. Hongk. 456. 1861 = Dryopteris filix mas.
chattagrammicum Diels, Nat. Pfl. **1**[4]. 186. 1899 = A. membranifolium.
cheilanthoides Kze. Linn. **22**. 578. 1849 = Dryopteris.
cheiloplotium Fée, 8 mém. 103. 1857 = Dryopteris.
(S) **chimborazense** C. Chr. Ind. 68. 1905 — Chimborazo.
Nephrodium elatum Bak. Syn. 298. 1867 (non pag. 502 nec Desv. 1827); Aspidium J. Sm. 1877 (non Mett.; Kuhn 1868).
chlorophyllum »Hk.«; Salom. Nom. 46. 1883 (nomen) — Cuba.
chontalense Fourn. Bull. Soc. Fr. **19**. 254. 1872 = Dryopteris.
Christianae Jenm. Bull. Dep. Jam. **2**. 285. 1895 — Jamaica.
(sp. dub.; an Polypodium miserum Hew. 1838?).
chrysocarpon Fée, 8 mém. 103. 1857 = Dryopteris filix mas.
Chrysocoma Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 253 = Dryopteris.
chrysolepis Fée, 8 mém. 107. 1857 = Dryopteris crenata.
chrysolobum Klf.; Link, Hort. Berol. **2**. 117. 1833 = Dryopteris.
chrysotrichum Christ, Engl. Jahrb. **23**. 352. 1896 = Dryopteris.
(S) **cicutarium** (L.) Sw. Schrad. Journ. **1800**[2]. 36. 1801. Mett. n. 278. Chr. 230. NPfl. 186 — Trop. — Polypodium L. Syst. Nat. ed. X. **2**. 1326. 1759; Sagenia Moore 1857; Nephrodium Bak. 1867. HB. 299; Dryopteris O. Ktze. 1891 —? Polydictyum heterophyllum Pr. 1849.
— *Aspidium apifolium Schkuhr 1809; ? Nephrodium Hk. et Arn. 1832; Sagenia J. Sm. 1841; Microbrochis Pr. 1849.
— Aspidium tenuifolium Mett. 1869. — A. Gaudichaudii Mett. 1869. 1870.
— *Aspidium latifolium Pr. 1825. Mett. n. 279 (non J. Sm. 1841); Sagenia Pr. 1836. (Amer. trop.).
— *Aspidium coadunatum Wall. 1828. Mett. n. 280 (non Klf. 1824); Sagenia J. Sm. 1841 — S. macrodonta Fée 1850-52 (Asia. Afr.).
cicutarium Willd. sp. **5**. 215. 1810; Kze. Flora **1839**[1]. Beibl. 33 = Dryopteris protensa.
ciliatum Wall. List n. 351. 1828 = Dryopteris calcarata var.
ciliatum Mett.; Salom. Nom. 46. 1883 = Dryopteris excelsa.
cinnamomeum Sw. Syn. 55. 1806 = Polystichum denticulatum.
cirrhosum Schum. Vid. Selsk. Afh. **4**. 231 (seors. 457). 1827 = Dryopteris.

clypeolarium Desv. Prod. 249. 1827 = Polystichum adiantiforme.
clypeolatum Mett.; Salom Nom. 47. 1883 = Nephrodium clypeolutatum.
coadunatum Klf. Enum. 239. 1824 — Brasilia.
Nephrodium Moore 1858. (Dryopteris sp. dub.).
coadunatum Wall. List n. 377. 1828; Hk. et Grev. Ic. Fil. t. 202. 1831 = A. cicutarium*.
coarctatum Kze. Bot. Zeit. **1845**. 287 = Dryopteris.
Cochinchinae Mett.; Salom. Nom. 47. 1883 = Cyclopeltis Presliana.
cochleatum Spr. Syst. **4**. 106. 1827 = Dryopteris.
cognatum Fée, Gen. 292. 1850-52 (nomen); Mett. Pheg. u. Asp. 61. n. 145. 1858 = Dryopteris.
commutatum Franch. et Sav. En. pl. Jap. 240. 1876 = Dryopteris Maximowiczii.
Comorense Kuhn, Fil. Afr. 128. 1868 = Dryopteris Spekei.
concinnum Link, Hort. Berol. **2**. 120. 1833 = A. Linkii.
concinnum Mett. Fil. Lips. 89. 1856 = Dryopteris.
concinnum Thwait. En. pl. Zeyl. 392. 1864 = Dryopteris deparioides var.
confertum Klf. Enum. 232. 1824 = Cyclodium meniscioides.
confluens Fée, Gen. 293. 1850-52 — Amer. austr.
Lastrea Moore 1858. (Dryopteris sp. dub.).
confluens Mett.; Kuhn, Linn. **36**. 125. 1869 = A. Mülleri.
congener Bl. Enum. 165. 1828 = Polystichum aculeatum var. 32.
coniifolium Pr. Del. Prag. **1**. 175. 1822 = Adiantopsis chlorophylla.
coniifolium Wall. List n. 341. 1828 = Polystichum carvifolium.
conioneuron Mett. Pheg. u. Asp. 102. n. 243. 1858 = Dryopteris extensa.
conjugatum Bl. Enum. 169. 1828 = A. Leuzeanum.
connexum Mett.; Salom. Nom. 47. 1883 = Dryopteris Martini.
consanguineum Kze.; Mett. Pheg. u. Asp. 101 n. 241 b. 1858 = D. gongylodes.
consanguineum Fée, 11 mém. 76 t. 20 f. 3. 1866 = Dryopteris opposita.
consobrinum Bory: Fée, Gen. 292. 1850-52 (nomen); 11 mém. 85. 1866 — Guadeloupe. — Lastrea Moore 1858. (Dryopteris sp. dub.).
conspersoides Fée, 8 mém. 105. 1857 = Dryopteris lugubris.
conspersum Kze. Flora **1839**[1]. Beibl. 32 = Dryopteris patens.
conterminum Willd. sp. **5**. 249. 1810 = Dryopteris opposita.
continuum Desv. Berl. Mag. **5**. 320. 1811 = Dryopteris gongylodes.
contractum Link, Fil. sp. 104. 1841 — Hort. — Nephrodium? Moore 1858.
contractum Sod. Cr. vasc. quit. 220. 1893 = A. quitense.
controversum Hance, Ann. sc. nat. IV. **18**. 235. 1862 = Polystichum amabile.
Corazonense Christ, Farnkr. 254. 1897 = Dryopteris.
cordatum Steud. Nomencl. **2**. 61. 1824 = Athyrium filix femina.
cordifolium Sw. Schrad. Journ. **1800**[2]. 32. 1801 = Nephrolepis.
cordifolium Pr. Epim. 63. 1849 = A. brachiatum.
coriaceum Sw. Schrad. Journ. **1800**[2]. 40. 1801 = Polystichum adiantiforme.
coriaceum Lowe, Ferns **6** t. 36. 1857 = Polystichum Berterianum.
coriandrifolium Sw. Schrad. Journ. **1800**[2]. 36. 1801 = A. cicutarium vel A. hippocrepis.
cormosum Kl.; Salom. Nomenkl. 47. 1883 (nomen) — Ceylon.
cornu cervi Don, Prod. Fl. Nepal. 5. 1825 = Polystichum aristatum var.
coronopus Desv. Prod. 246. 1827 = A. cicutarium.
costale Bl. Enum. 170. 1828 = Athyrium nigripes.
costale Mett.; Kuhn, Linn. **36**. 111. 1869 = Dryopteris densisora.

(S) **costaricanum** C. Chr. Ind. 70. 1905. — Costa Rica. — Nephrodium athyrioides Bak. JoB. **1884**. 363 (non Aspidium Mart. et Gal. 1843).
costulare Kuhn, v. Deck. Reisen **3**[3]. Bot. 65. 1879 = Dryopteris.
Crameri Christ, Farnkr. d. Schw. 153. 1900 = Dryopteris oyamensis.
craspedosorum Maxim. Bull. Ac. St. Pétersb. **15**. 231. 1871 = Polystichum.
crassifolium Bl. Enum. 158. 1828 = Dryopteris.
crenatum Sommerf. Vet. Ak. Hdl. **1834**. 104. 1835 = Athyrium.
crenatum Mett. Fil. Lips. 89. 1856 = Polystichum aculeatum var. 4.
crenatum Kuhn, Fil. Afr. 129. 1868 = Dryopteris.
crenulans Fée, Cr. vasc. Br. **1**. 139 t. 47 f. 1. 1869 = Dryopteris villosa.
crinitum Wall. List n. 357. 1828 = Dryopteris.
crinitum Mart. et Gal. Mém. Ac. Brux. **15**. 66 t. 17 f. 2. 1842 = Dryopteris filix mas.
crinulosum Desv. Prod. 249. 1827 = Polystichum aculeatum var. 1.
cristatum Sw. Schrad. Journ. **1800**[2]. 37. 1801 = Dryopteris.
cristatum × marginale Dav. Bot. Gaz. **19**. 494. 1894 = Dryopteris.
cristatum × spinulosum Milde, Nova Acta **26**. 533. 1858 = Dryopteris.
cruciatum Willd. sp. **5**. 278. 1810 = Dryopteris.
crystallinum Mett. (Fil. Lips. 90. 1856) = ? Dryopteris hirta var.
Ctenitis Link. Hort. Berol. **2**. 122. 1833 = Dryopteris.
Cubense Kuhn, Linn. **36**. 108. 1869 = Dryopteris.
cucullatum Bl. Enum. 151. 1828 = Dryopteris unita.
cucullatum β rigidum Bl. Enum. 152. 1828 = Dryopteris Haenkeana.
cultratum Pr. Del. Prag. **1**. 174. 1822 = Didymochlaena truncatula.
Cumingianum Kze. Farnkr. **1**. 17 t. 9 f. 2. 1840 = Dryopteris.
Cumingianum Sturm, En. pl. vasc. cr. chil. —. 1858 = Polystichum adiantiforme.
cuneatum Schkuhr, Kr. Gew. **1**. 198 t. 56 b. 1809 = Asplenium fissum.
Cunninghami Kze. Linn. **23**. 225. 1850 = Dryopteris pennigera.
Cunninghamianum Colenso, Tasm. Journ. **2**. —. 1845 = Polystichum adiantiforme.
Currori Mett.; Kuhn, Fil. Afr. 130. 1868 = Dryopteris.
curvifolium Kze. Bot. Zeit. **1848**. 283 = Polystichum carvifolium.
cuspidatum Desv. Prod. 244. 1827. (Plum. t. 153) — Hispaniola.
cuspidatum Mett. Pheg. u. Asp. 92. n. 223. 1858 = Dryopteris khasiana.
cyatheoides Klf. Enum. 234. 1824 = Dryopteris.
cycadinum Franch. et Sav. En. pl. Jap. **2**. 236, 630. 1879 = Dryopteris.
cyclochlamys Fée, Gen. 296. 1850-52 — Réunion.
Lastrea Moore 1858. — f. Cordemoy fortasse Dryopteris nudata.
cystolepidotum Miq. Ann. Lugd. Bat. **3**. 177. 1867 = Dryopteris.
Cystostegia Hk. sp. **4**. 26 t. 227. 1862 = Polystichum.
(Ar) **Dahlii** (Hier.) Diels in Schum. et Laut. Fl. deut. Südsee 116. 1901 — Neu Lauenburg. — Phegopteris Hier. Notizbl. bot. Gart. Berl. **2**. 84. 1898.
dasychlamys Fée, 10 mém. 38 t. 41 f. 4. 1865 — Réunion. (Dryopteris sp. dub.).
davallioides Sw. Schrad. Journ. **1800**[2]. 33. 1801 = Nephrolepis acuminata.
davallioides Kuhn, Verh. zool. bot. Ges. **19**. 578. 1869 = Dryopteris.
Dayi Bedd. JoB. **1888**. 4 = Dryopteris.
decipiens Luerss. Engl. Jahrb. **4**. 360. 1883 = Dryopteris.
decompositum Spr. Syst. **4**. 109. 1827 = Dryopteris.
decrescens Kze.; Mett. Pheg. u. Asp. 84. n. 202. 1858 = Dryopteris.

(S) **decurrens** Pr. Rel. Haenk. **1**. 28. 1825. Chr. 229. NPfl. 186. — China austr. et Ind. bor.-Malesia-Polynesia.
Sagenia Houlst.; Moore 1857; Nephrodium Bak. 1867. HB. 299; Dryopteris O. Ktze. 1891. Aspidium decurrens Kze. 1846. Mett. n. 284; Sagenia Moore 1858 — Cardiochlaena alata Fée 1850-52.
decurrens Lowe, Ferns **6** t. 5. 1857 = Dryopteris decursive-pinnata.
decursive-pinnatum Kze. Bot. Zeit. **1848**. 555 = Dryopteris.
decurtatum Kze. Linn. **23**. 276, 299. 1850 = Dryopteris abortiva.
decussatum Sieb. Syn. Fil. exs. n. 189 —? — Lastrea Pr. 1836 (nomen).
deltodon Bak. Gard. Chr. n. s. **14**. 494. 1810 = Polystichum.
deltoideum Sw. Schrad. Journ. **1800²**. 34. 1801 = Dryopteris.
densum Wall. List n. 390. 1828 = Dryopteris sparsa.
dentatum Sw. Syn. 59. 1806 = Cystopteris fragilis.
denticulatum Sw. Schrad. Journ. **1800²**. 40. 1801 = Polystichum.
deparioides Hk. Fil. exot. t. **3**. 1858 = Dryopteris.
deparioides Diels, Nat. Pfl. **1⁴**. 186. 1899 = A. dapariopsis.
depariopsis C. Chr. Ind. 71. 1905 — Fiji.
Polypodium (Depariopsis) deparioides Bak. JoB. **1879**. 297; Aspidium Diels 1899. NPfl. 186 (non Hk. 1858).
depastum Schkuhr, Kr. Gew. **1**. 50 t. 51. 1806 = Dryopteris filix mas.
Desvauxii Mett.; Kuhn, Fil. Afr. 131. 1868 = Dryopteris obtusiloba.
deversum Kze. Linn. **23**. 226, 299. 1850 = Dryopteris patens et parasitica.
(T?) **devexum** Kze. Bot. Zeit. **1848**. 259 — China austr. Formosa. Ceylon. Malesia. Ins. Philipp. — Dryopteris O. Ktze. 1891. Polypodium membranaceum Hk. 1862 (non Don 1825); Aspidium Hk. 1864. HB. 259. NPfl. 188; Sagenia Bedd. 1870; Pleocnemia Bedd. 1876. Aspidium giganteum var. minor Hk. sp. **4**. 50. 1862; Sagenia Bedd. 1863.
diaphanum Zoll. et Moritz, Nat. Geneesk. Arch. **2**. 203. 1845 = Polystichum.
diastematocarpon Fée, 8 mém. 107. 1857 = Dryopteris patula.
Dickinsii Franch. et Sav. En. pl. Jap. **2**. 236, 629. 1879 = Dryopteris.
dicksoniaefolium A. Rich. Sert. Astrol. XXXIX. 1834 = Microlepia speluncae.
dicksonioides Fée, Cr. vasc. Br. **1**. 143 t. 49 f. 1. 1869 = Dryopteris protensa.
diffractum Christ, Bull. Boiss. **7**. 17. 1899 = Dryopteris.
difforme Bl. Enum. 160. 1828 = A. irregulare.
dilaceratum Kze. Linn. **23**. 226, 300. 1850 = A. cicutarium.
dilatatum Sm. Fl. brit. **3**. 1125. 1804 = Dryopteris spinulosa*.
dimidiatum Mett.; Kuhn, Fil. Afr. 131. 1868 = Dryopteris.
dimorphum Kze. Bot. Zeit. **1848**. 261 = Dryopteris heterocarpa var.
diplazioides Moritz; Mett. Pheg. u. Asp. 83. n. 200. 1858 = Dryopteris Moritziana.
discolor Langsd. et Fisch. Ic. Fil. 16 t. 18. 1810 = Polystichum adiantiforme.
discretum Don, Prod. Fl. Nep. 4. 1825 = Polystichum lobatum var. 3.
dissectum Fée, 8 mém. 107. 1857 = Polystichum denticulatum.
dissectum Mett. Ann. Lugd. Bat. **1**. 232. 1864 = Dryopteris invisa.
dissectum Christ, Engl. Jahrb. **23**. 353. 1896 = Dryopteris.
(Pl) **dissidens** Mett. Pheg. u. Asp. 116. n. 275 b. 1858 — Puerto Rico.
Nephrodium Hk. 1862. HB. 295; Pleocnemia Moore 1858; Dryopteris O. Ktze. 1891.
distans Viv. App. ad Fl. Cors. Prod. 8. 1825 = Woodsia ilvensis (t. Milde; t. Trevisan Polystichum sp.).

distans Kuhn, Fil. Afr. 209. 1868 = Dryopteris.
distans Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 253 = Dryopteris brunnea.
diversilobum Mett. Pheg. u. Asp. 100. n. 239. 1858 = Dryopteris canescens vel D. parasitica.
divisum Wall. List n. 393. 1828 = Dryopteris Boryana.
(S) **domingense** (Bak.) C. Chr. Ind. 72. 1905 — San Domingo.
Nephrodium Bak. Ann. Bot. **5**. 329. 1891.
Donianum Spr. Syst. **4**. Suppl. 320. 1827 = Dryopteris filix mas*.
(Ar) **draconopterum** Eat. Mem. Amer. Acad. n. s. **8**. 211. 1860. NPfl. 186 — Columbia-Ecuador. — Polypodium Hk. 1863. HB. 319. Phegopteris Mett. 1864. Dryomenis Purdiei J. Sm. 1854 (nomen); Pleopeltis Moore 1862.
Dregei Fée, Gen. 292. 1850-52 = Dryopteris thelypteris var. squamigera.
drepanopteron Mett. Fil. Lips. 93 t. 19 f. 1-4. 1856 = Athyrium.
drepanum Sw. Schrad. Journ. **1800²**. 38. 1801 = Polystichum.
drepanum Schkuhr, Kr. Gew. **1**. t. 47. 1806 = Dryopteris spinulosa *dilatata.
Dryopteris Baumg. Enum. stirp. Transs. **4**. 29. 1846 = Dryopteris d.
(S) **dubium** (Clarke et Bak.) Bedd. Handb. Suppl. 47. 1892 — Assam.
Nephrodium cicutarium var.? dubia Clarke et Bak. Journ. Linn. Soc. **24**. 417. 1888.
Dubrueillianum Gaud. Freyc. Voy. Bot. (333) t. 9. 1827 = Dryopteris cyatheoides.
dumetorum Sm. Engl. Fl. **4**. 281. 1828 = Dryopteris spinulosa *dilatata.
duriusculum Sturm, Enum. pl. vasc. cr. chil. —. 1858 = Polystichum.
Durvillaei Bory, Dup. Voy. Bot. **1**. 266. 1828 = A. melanocaulon.
Duthiei Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 532 t. 6. 1899 = Polystichum.
ebenum J. Sm. Cat. cult. ferns 52. 1857 = A. pica.
eburneum Wall. List n. 389. 1828 = Dryopteris cana vel Athyrium drepa-[nopterum.
echinatum Mett. Ann. Lugd. Bat. **1**. 230. 1864 = Dryopteris.
Ecklonii Kze. Linn. **10**. 546. 1836 = Dryopteris gongylodes.
edentulum Kze. Bot. Zeit. **1846**. 474 = Dryopteris.
edule Spr. Syst. **4**. 98. 1827 = Nephrolepis cordifolia.
effusum Gris. Fl. br. W. Ind. 690. 1864 = Dryopteris.
elatior Fée, Cr. vasc. Br. **1**. 132 t. 42 f. 3. 1869 = Dryopteris opposita.
elatum Boj. Hort. Maur. 390. 1837 (nomen) = Dryopteris mauritiana var.
elatum Mett.; Kuhn, Fil. Afr. 130. 1868 = Dryopteris.
elatum J. Sm. Ferns br. and for. ed. II. 304. 1877 = A. chimborazense.
elegans (Moore et Houlst.) Mett. Pheg. u. Asp. 71. n. 168. 1858 (nomen) — Ceylon.
Lastrea Moore et Houlst. (1852); J. Sm. Cat. cult. ferns 57. 1857. — (Dryopteris sp. aff. D. recedens).
elongatum Sw. Schrad. Journ. **1800²**. 38. 1801 = Dryopteris filix mas*.
elongatum Willd. sp. **5**. 269. 1810 = Dryopteris filix mas *oligodonta.
emarginatum Willd. sp. **5**. 235. 1810 (Plum. t. 62 A) — Hispaniola.
Polystichum? Moore 1858 — (an Dryopteris subobliquata?).
eminens Wikstr. Vet. Ak. Hdl. **1825**. 436. 1826. = Nephrolepis exaltata.
eminens Bedd. JoB. **1888**. 4 = Dryopteris megaphylla.
(S) **Endresi** (Bak.) C. Chr. Ind. 72. 1905 — Costa Rica.
Nephrodium Bak. Ann. Bot. **5**. 329. 1891.
ensifolium Schkuhr, Kr. Gew. **1**. 32 t. 32. 1806 = Nephrolepis biserrata.
equestre Kze. Linn. **18**. 347. 1844 = Dryopteris excelsa.
eriocarpum Wall. List n. 342. 1828 = Dryopteris crenata.
eriocaulon Fée, Cr. vasc. Br. **1**. 136 t. 44 f. 1. 1869 = Dryopteris.

eriosorus Fée, Cr. vasc. Br. **2**. 73 t. 101. 1872-73 = Dryopteris.
erosum Schkuhr, Kr. Gew. **1**. 46 t. 45. 1806 = Dryopteris filix mas.
erubescens Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 251 = Dryopteris.
erythrosorum Eat. in Perry, Narr. Exp. to China **2**. 330. 1856 = Dryopteris.
erythrostemma Christ, Bull. Boiss. II. **4**. 961. 1904 = Dryopteris.
(S) **eurylobum** Christ, Bull. Soc. bot. Belg. **35**. 206. 1896. (Chr. 229). NPfl. 186 — Costa Rica. Ecuador.
eusorum Thwait. En. pl. Zeyl. 391. 1864 = Dryopteris truncata.
(S) **Everettii** (Bak.) C. Chr. Ind. 73. 1905 — Ins. Natuna (Malesia).
Nephrodium Bak. Kew Bull. **1896**. 41.
exaltatum Sw. Schrad. Journ. **1800**^{2}. 32. 1801 = Nephrolepis.
(Pl) **excellens** Bl. Enum. 160. 1828 — Java.
Proferea Pr. 1849; Nephrodium Hk. 1862. HB. 295; Dryopteris O. Ktze. 1891.
excultum Mett. Pheg. u. Asp. 69. n. 162. 1858 ≐ Dryopteris effusa*.
exiguum Kze.; Mett. Pheg. u. Asp. 76. n. 180. 1858 = Dryopteris philippina.
exile Hance, JoB. **1883**. 268 = Polystichum aristatum.
expansum Willd. sp. **5**. 284. 1810 = Athyrium.
expansum Desv. Berl. Mag. **5**. 320. 1811 = A. martinicense.
expansum Dietr. Fl. Jenensis —. 1826 = Dryopteris filix mas.
expansum Mart.; Fée, Gen. 292. 1850-52 — Brasilia. — Lastrea Moore 1858.
expansum Fée, 10 mém. 42. 1865 = Dryopteris Trianae.
exsudans Fourn. Mex. pl. **1**. 93. 1872 = Dryopteris oligocarpa.
extensum Bl. Enum. 156. 1828 = Dryopteris.
extensum Fée, Gen. 294. 1850-52 = Dryopteris Trianae.
Fadyenii Mett. Fil. Lips. 95 t. 23 f. 13-14. 1856 = Fadyenia.
falcatum Sw. Schrad. Journ. **1800**^{2}. 31. 1801 = Polystichum.
falciculatum Raddi, Opusc. sci. Bol. **3**. 289. 1819 = Dryopteris.
falcilobum Hk.; Benth. Fl. Hongk. 455. 1861 = Dryopteris calcarata var.
falcinellum Sw. Syn. 46, 243. 1806 = Polystichum.
fallax Fisch.; Kze. Linn. **23**. 226. 1850; Fisch. Ind. Sem. ht. Petrop. 1855; Linn. **28**. 367 = Dryopteris amplissima.
Fargesii Christ, Bull. Soc. Fr. **52** Mém. I. 42. 1905 = Dryopteris.
fasciculatum Fourn. Ann. sc. nat. V. **18**. 295. 1873 = Dryopteris.
Fauriei Christ, Bull. Boiss. **6**. 193. 1898 = Athyrium.
Fendleri Eat. Mem. Amer. Acad. n. s. **8**. 210. 1860 = Dryopteris.
(Ar) **fernandense** (Bak.) Diels, Nat. Pfl. **1**^{4}. 186. 1899 — Fernando Po.
Polypodium Bak. Ann. Bot. **5**. 462. 1891.
ferox Bl. Enum. 153. 1828 = Dryopteris.
ferrugineum Sw. Syn. 56. 1806 = Polystichum adiantiforme.
ferrugineum Fée, Gen. 293. 1850-52 — India orient.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
ferrugineum Diels, Nat. Pfl. **1**^{4}. 186. 1899 = A. Zippelianum.
festinum Hance, JoB. **1883**. 269 = Polystichum Standishii.
fibrillosum Kuhn, v. Deck. Reisen **3**^{3}. Bot 65. 1879 = Dryopteris Poolii.
Filix femina Sw. Schrad. Journ. **1800**^{2}. 41. 1801 = Athyrium.
Filix mas Sw. Schrad. Journ. **1800**^{2}. 44. 1801 = Dryopteris.
Filix mas Pursh, Fl. Am. sept. **2**. 662. 1814 = Dryopteris Goldiana.
Filix mas var. Chrysocoma Christ, Bull. Boiss. **6**. 966. 1898 = Dryopteris chrysocoma.
Filix mas × spinulosum A. Br. in Döll. Fl. Bad. 30. 1857 = Dryopteris.

Filix-Panna Lucan. — = Dryopteris athamantica.
fimbriatum Willd. sp. **5**. 213. 1810 = A. Plumieri vel A. trifoliatum.
fimbriatum Wall. List n. 339. 1828 = Athyrium.
Finlaysonianum Wall. List n. 2235. 1829 —? — Lastrea falcifera Moore 1858.
Fischeri Mett. Pheg. u. Asp. 80. n. 192. 1858 = Dryopteris.
fissum Kze. Bot. Zeit. **1848**. 258 = A. menyanthidis vel A. pachyphyllum.
flaccidum Bl. Enum. 161. 1828 = Dryopteris.
flaccidum Fourn. Bull. Soc. Fr. **27**. 328. 1880 = Dryopteris Fournieri.
flagelliferum Roxb.; Wall. List n. 2234. 1829 = Nephrolepis exaltata.
flexile Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 252 = Dryopteris.
flexum Kze. Anal. 44. 1837 = Polystichum Berterianum.
flexuosum Fée, Cr. vasc. Br. **1**. 138 t. 46 f. 2. 1869 = Dryopteris.
floccigerum Bl. Enum. 147. 1828 = Nephrolepis.
floridanum Eat. in Chapm. Fl. South. U. St. 595. 1860 = Dryopteris.
foeniculaceum Hk. sp. **4**. 36 t. 237. 1862 = Polystichum.
Foenisecii Lowe, JoB. **1**. 26. 1834 (nomen) = Dryopteris aemula.
? foliolosum Wall. List n. 359. 1828 = Acrophorus stipellatus.
fontanum Sw. Schrad. Journ. **1800**². 40. 1801 = Asplenium.
Forbesii Hance, JoB. **1875**. 198 = Dryopteris subtripinnata.
formosanum Christ, Bull. Boiss. II. **4**. 615. 1904 = Dryopteris.
formosum Fée, Gen. 296. 1850-52 = Polystichum denticulatum.
Forsteri Kze. Bot. Zeit. **1846**. 462 = A. latifolium.
fragile Sw. Schrad. Journ. **1800**². 40. 1801 = Cystopteris.
fragrans Sw. Schrad. Journ. **1800**². 35. 1801 = Dryopteris.
fragrans Gray, Nat. Arr. br. Pl. **2**. 9. 1821 = Dryopteris rigida.
Francoanum Fourn. Bull. Soc. Fr. **19**. 255. 1872 = Dryopteris Skinneri.
Fraseri Mett.; Kuhn, Linn. **36**. 109. 1869 = Dryopteris.
fraternum Mett.; Kuhn, Fil. Afr. 132. 1868 = Dryopteris Vogelii.
fraxinellum Christ, Bull. Boiss. **7**. 15. 1899 = Polystichum.
fraxinifolium Schrad. Gött. gel. Anz. **1824**. 1868 = A. martinicense.
Freyreissii Wikstr. Vet. Ak. Hdl. **1825**. 443. 1826 = Polystichum aculeatum var. 25.
frondosum Wikstr. Vet. Ak. Hdl. **1825**. 437. 1826 = Polystichum denticulatum.
frondosum Lowe, Tr. Cambr. Phil. Soc. **4**. 6. 1830 = Polystichum Webbianum.
frondulosum Fée, 10 mém. 40. 1865 — Réunion. (Dryopteris sp. dub.).
fructuosum Christ, Bull. Soc. Fr. **52** Mém. I. 38. 1905 = Dryopteris.
fumarioides Pr. Rel. Haenk. **1**. 39. 1825 = Cystopteris fragilis.
Funckii Mett. Ann. sc. nat. V. **2**. 246. 1864 = Dryopteris.
funestum Kze. Linn. **9**. 96. 1834 = Dryopteris protensa*.
furcatum Kl. Linn. **20**. 371. 1847 = Dryopteris.
fuscatum Willd. sp. **5**. 256. 1810 = Polystichum aculeatum vel P. mohrioides.
fuscipes Wall. List n. 361. 1828 = A. membranifolium et Dryopteris sagenioides.
fuscipes Bedd. Ferns br. Ind. Suppl. 15 t. 366. 1876 = A. membranifolium.
(S) **gaboonense** Kuhn, Fil. Afr. 133. 1868 (nomen) — Africa occ. trop.
Polypodium tenerifrons pt. Hk. sp. **5**. 104. 1864.
Galanderi Hier. Engl. Jahrb. **22**. 369. 1896 = Dryopteris.
Gardneri Christ, Farnkr. 229. 1897 = A. organense.
Gardnerianum Mett. Pheg. u. Asp. 71 n. 170. 1858 = ? Dryopteris dissecta.
Gaudichaudii Mett.; Kuhn, Linn. 36. 123. 1869; Mett. Novara Exp. Bot. **1**. 219. 1870 = A. cicutarium*.

Gaudichaudii Trevis. Hedwigia **13**. 148. 1874 = Cyclopeltis semicordata.
Geisbreghti Christ, Bull. Boiss. II. **4**. 962. 1904 (err.) = Dryopteris Ghiesbreghtii.
gelidum Kze.; Kl. Linn. **20**. 365. 1847 = Polystichum aculeatum var. 27.
Germani L'Herm.; Fée, 11 mém. 82 t. 23 f. 3. 1866 — Guadeloupe.
(Dryopteris sp.; non D. Germaniana infra).
Germinyi Linden, L'Ill. Hort. **27**. 163 t. 402. 1880 = Dryopteris obliquata var.
geropogon Fée, 10 mém. app. 1865 = Dryopteris.
Ghiesbreghtii Fourn. Mex. pl. **1**. 94. 1872 = Dryopteris oligocarpa.
gibbosum Willd. sp. **5**. 222. 1810 = Nephrolepis biserrata.
(Pl) **giganteum** Bl. Enum. 159. 1828 — India austr. Ceylon. Perak. Java.
Pleocnemia Pr. 1849; Sagenia Moore 1858; Nephrodium Bak. 1874. HB. 503; Dryopteris O. Ktze. 1891; Pleocnema Trimeni Bedd. 1883; Aspidium Diels 1899. NPfl. 188.
glabellum Lowe, Ferns **6** t. 32. 1857 = Dryopteris.
glaberrimum A. Rich. Sert. Astrol. XVIII. 1834 — N. Guinea.
Nephrodium Moore 1858 — (Dryopteris sp. dub.).
glabratum Mett.; Kuhn, Fil. Afr. 133. 1868 = Dryopteris.
glabrum Mett. Pheg. u. Asp. 59. n. 140. 1858 = Dryopteris.
glanduliferum Wall. List n. 347. 1828 = Dryopteris ochthodes.
glanduliferum Karst.; Kl. Linn. **20**. 369. 1847 = Dryopteris Balbisii.
glanduliferum Kuhn, Verh. zool. bot. Ges. **19**. 577. 1869 = Dryopteris Haenkeana (f. Fourn.) vel D. tannensis (f. Bak.).
glanduliferum Mett. Ann. Lugd. Bat. **1**. 232. 1864 = Dryopteris subspinosa.
glanduligerum Kze. Anal. 44. 1837 = Dryopteris gracilescens*.
glandulosum Bl. Enum. 144. 1828 = Dryopteris.
glandulosum Hk. et Grev. Ic. Fil. t. 140. 1829 = Polystichum viscidulum.
Glaziovii Christ, Bull. Boiss. II. **2**. 633. 1902 = Dryopteris.
gleichenioides Christ, Bull. Boiss. II. **4**. 960. 1904 = Dryopteris.
globuliferum Mann, Pr. Amer. Acad. **7**. 217. 1868 = Dryopteris.
(S) **Godeffroyi** (Luerss.) Christ, Farnkr. (230) 370. 1897. NPfl. 186 — Fiji.
Dennstaedtia Luerss. Fil. Graeff. 222 t. 19. 1871; Deparia Bak. 1891.
Goeringianum Kze. Bot. Zeit. **1848**. 557 = Athyrium.
goggilodus Schkuhr, Kr. Gew. **1**. 193 t. 33 c. 1809 = Dryopteris gongylodes.
Goldianum Hk.; Goldie, Edinb. Phil. Journ. **6**. 333. 1822 = Dryopteris.
Goldieanum Christ, Bull. Boiss. II. **2**. 828. 1902 = Dryopteris monticola.
gongylodes auctt. Mett. Pheg. u. Asp. 101. n. 241. 1858 = Dryopteris.
gracilescens Bl. Enum. 155. 1828 = Dryopteris.
gracilipes Fée, Cr. vasc. Br. **1**. 146 t. 49 f. 2. 1869 = Polystichum denticulatum.
Grammatosorus Salom. Nomenkl. 54. 1883 = Grammatosorus Blumeanus.
grammitoides Christ, Bull. Boiss. **6**. 193. 1898 = Dryopteris.
grande J. Sm JoB. **3**. 410. 1841 (nomen); Mett. Pheg. u. Asp. 121. n. 285. 1858 = A. grandifolium.
grande Fée, Gen. 295. 1850-52 = Dryopteris filix mas *elongata.
(S) **grandifolium** Pr. Epim. 64. 1849 — Ins. Philipp.
A. grande J. Sm. 1841 (nomen); Mett. n. 285. 1858; Sagenia Moore 1857; Nephrodium Bak. 1891 — Cardiochlaena laevis et sinuosa Fée 1850-52.
grandifolium Mett. Pheg. u. Asp. 134. n. 294. 1858 = A. polymorphum.
Grayanum Regel, Ind. sem. ht. Petr. 1855; Linn. **28**. 367. 1856 = Dryopteris spinulosa *intermedia.

Griffithii (Moore) Diels, Nat. Pfl. 1^4. 186. 1899 — Japonia. China-Ind. bor.
Dictyocline Moore, Ind. LIX. 1857; Hemionitis Hk. f. et Thoms. hb. Hk. 1864. HB. 399. Chr. 62; Dictyogramme Trevis. 1877; Gymnogramme Hance 1887 — Hemionitis Wilfordii Hk. 1859; Dictyocline J. Sm. 1875; Dictyogramme Trevis. 1877.

Griffithii Bedd. Ferns br. Ind. Suppl. 15. 1876 = A. multicaudatum.

Grunowii Bolle, Bonpl. **3**. 123. 1855 = Dryopteris.

guadalupense Christ, Bull. Soc. bot. Belg. **35**. 210. 1896 = Dryopteris.

Gueintzianum Mett. Pheg. u. Asp. 83. n. 201. 1858 = Dryopteris.

guianense Kl. Linn. **20**. 364. 1847 = Polystichum abbreviatum*.

guineense Schum. Vid. Selsk. Afh. **4**. 229. (seors. 455). 1827 = Nephrolepis biserrata.

Haenkeanum Fourn. Ann. sc. nat. V. **18**. 298. 1873 = Dryopteris.

(T) **Haenkei** Pr. Rel. Haenk. **1**. 30. 1825. NPfl. 186 — Ins. Mariannae. Timor.

Haleakalense Mann, Pr. Amer. Acad. **7**. 216. 1868 = Polystichum lobatum.

Halleri Willd. sp. **5**. **274**. 1810 = Asplenium fontanum.

Hamiltonii Spr. Syst. **4**. 108. 1827 }
Hamiltonianum Wall. List n. 2232. 1829 } = Polystichum aristatum var.

Hancockii Bak. Ann. Bot. **5**. 314. 1891 = Polystichum.

Harrisonii Christ, Bull. Soc. bot. Belg. **35**. 209. 1896 = Dryopteris.

Hartwegii Kl. Linn. **20**. 366. 1847 = Polystichum aculeatum var 4.

Harveyi Mett.; Kuhn, Linn. **36**. 115. 1869 = Dryopteris.

hastatum Kuhn (Engl. Jahrb. **24**. 124. 1897. syn.) = Dryopteris.

hastifolium Gris. Fl. br. W. Ind. 694. 1864 = Dryopteris sagittata.

hastulatum Tenore, Atti Ist. Incorr. Nap. **5**. 149 t. 4 f. 7 A. b. 1830 = Polystichum aculeatum var. 42.

Hawaiiense Hill. Fl. Haw. 575. 1888 = Dryopteris filix mas*.

(Ar) **Haynaldii** (Sod.) C. Chr. Ind. 76. 1905 — Ecuador.
Polypodium Sod. Rec. 61. 1883; Cr. vasc. quit. 305. 1893.

(S) **hederifolium** (Bak.) Diels, Nat. Pfl. 1^4. 185. 1899 — Ins. Salomonis.
Nephrodium Bak. Journ. Linn. Soc. **19**. 295. 1882; Ic. pl. t. 1665.

helveolum Fée, Cr. vasc. Br. **1**. 132 t. 42 f. 2. 1869 = Dryopteris opposita.

hemicardion Trevis. Hedwigia **13**. 148. 1874 = Cyclopeltis semicordata.

(S) **heptaphyllum** (Bak.) C. Chr. Ind. 76. 1905 — Samoa.
Nephrodium Bak. JoB. **1879**. 296.

heracleifolium Willd. sp. **5**. 217. 1810 = A. trifoliatum.

heracleifolium Bl. Enum. 145. 1828 = ? A. brachiatum.

Herusianum Fourn. Ann. sc. nat. V. **18**. 295. 1873 (nomen) = ? Dryopteris [truncata.

heterocarpon Bl. Enum. 155. 1828 = Dryopteris.

heterocarpum Bedd. Ferns br. Ind. Suppl. 15. 1876 = A. heterosorum.

heterodon Schrad. Gött. gel. Anz. **1824**. 869 = Cyclodium meniscioides.

heterodon Bl. Enum. 157. 1828 = Mesochlaena polycarpa

heterophyllum Hk. Ic. pl. t. 920. 1854 = Dryopteris canescens*.

heteropteron Mett.; Kuhn, Fil. Afr. 134. 1868 = Dryopteris.

(S) **heterosorum** (Bak.) Bedd. Handb. Suppl. 46. 1892 — Birma. Assam.
Nephrodium Bak. 1874. HB. 504; Dryopteris O. Ktze. 1891; Sagenia heterocarpa Bedd. Ferns br. Ind. t. 47. 1866; Aspidium Bedd. 1876. NPfl. 185 (non Bl. 1828).

Hillebrandii Hill. Fl. Haw. 568. 1888 = Polystichum lobatum var. 6.

Hilsenbergii Sieb. Syn. fil. exsic. n. 49. = Dryopteris parasitica.

(S) **hippocrepis** (Jacq.) Sw. Syn. 51. 1806. Chr. 230. NPfl. 186 — Ind. occ.
Polypodium Jacq. Collect. 3. 186. 1879; Nephrodium Pr. 1825?; Desv. 1827; Sagenia Pr. 1836; Polydictyum Pr. 1849. Aspidium vel Nephrodium cicutarium part. auctt.

hirsutulum Sw. Schrad. Journ. **1800**². 32. 1801 = Nephrolepis.

hirsutulum Ham.; Wall. List n. 2240. 1829 = ? Dryopteris parasitica.

hirsutulum Wall. List n. 7083. 1829 = Dryopteris calcarata.

hirsutum Kze.; Mett. Pheg. u. Asp. 107. n. 256. 1858 = Dryopteris adenophora.

hirtipes Bl. Enum. 148. 1828 = Dryopteris.

hirtum Sw. Schrad. Journ. **1800**². 39. 1801 = Dryopteris.

hispidulum Dcne. Nouv. Ann. Mus. **3**. 346. 1834 = Dryopteris.

hispidum Sw. Schrad. Journ. **1800**². 39. 1801 = Polystichum.

Hoffmannseggii Poir. Enc. Suppl. **4**. 509. 1816 = Nephrolepis biserrata.

Hookeri Wall. List p. 64. 1829; Hk. Ic. pl. t. 922. 1854 = Dryopteris arbuscula.

Hookeri Sweet, Hort. brit. (ed. II?) 579. 1830 = Fadyenia Fadyenii.

Hookeri Kl. Linn. **20**. 364. 1847 = Cyclodium meniscioides.

Hookeri Bak. Syn. 257. 1867 = Dryopteris orbicularis.

Hornei Kuhn, v. Deck. Reisen **3**³. Bot. 69. 1879 = Dryopteris.

(S) **Hosei** (Bak.) C. Chr. Ind. 77. 1905 — Borneo.
Nephrodium Bak. 1891; N. stenophyllum Bak. Journ. Linn. Soc. **22**. 227 t. 11. 1886 (non Sod. 1883 nec Bak. 1884); Aspidium Diels 1899. NPfl. 185.

Hudsonianum Mann, Pr. Amer. Acad. **7**. 217. 1868 = Dryopteris truncata.

humile Willd. sp. **5**. 220. 1810 = Pteris longifolia.

hymenodes Mett.; Kuhn, Linn. **36**. 123. 1869 = A. latifolium.

hymenophylloides Bl. Enum. 172. 1828 = Davallia.

ilicifolium Don, Prod. Fl. Nepal. 3. 1825 = Polystichum.

illyricum Borbás, Oest. Bot. Zeit. **41**. 354. 1891 = ? Polystichum lobatum × lonchitis.

imbecile Kze.; Ettingsh. Farnkr. 176 t. 107 f. 10, t. 109 f. 1. 1865 = Dryopteris patula.

imbricatum Klf.; Spr. Syst. **4**. 97. 1827 = Nephrolepis cordifolia.

imbricatum Fourn. Mex. pl. **1**. 96. 1872 = Dryopteris.

immersum Bl. Enum. 156. 1828 = Dryopteris.

immersum Hk. sp. **4**. 58. 1862 = A. brachiatum.

impressum Kze. Linn. **23**. 222. 1850 = Dryopteris immersa.

impressum Fée, Gen. 296. 1850-52 — Ceylon.
Lastrea Moore 1858 — (Dryopteris sp. dub.).

? Imrayanum Fée, 11 mém. 79. 1866 = Dryopteris rotundata*.

inaequale Schlecht Adumbr. 23 t. 12. 1825 = Dryopteris.

inaequilaterale Luerss. Fil. Graeff. 181. 1871 = Dryopteris Brackenridgei.

incisum Sw. Syn. 47. 1806 = Dryopteris serra var.

incisum Gris. Fl. br. W. Ind. 694. 1864 = Dryopteris.

indecorum Fourn. Mex. pl. **1**. 97. 1872 = Dryopteris.

inquinans Fée, 8 mém. 106. 1857 = Dryopteris indecora.

inquinans Kuhn, Fil. Afr. 135. 1868 = Nephrodium.

insigne Mett. Ann. sc. nat. V. **2**. 247. 1864 = Dryopteris.

intermedium Willd. sp. **5**. 262. 1810 = Dryopteris spinulosa*.

intermedium Sadl. Descr. pl. epiph. Hung. 16. 1820 = Polystichum lobatum.

intermedium Link, Enum. alt. **2**. 459. 1822 = Athyrium filix femina.

intermedium Bl. Enum. 161. 1828 = Dryopteris rhodolepis.
intermedium J. Sm. Bot. Voy. Herald 428. 1857 = A. cicutarium.
invisum Sw. Schrad. Journ. **1800**[2]. 34. 1801 = Dryopteris patens*.
invisum Raddi, Opusc. sci. Bol. **3**. 289. 1819 = Dryopteris patens.
invisum Christ, Engl. Jahrb. **23**. 350. 1896 = Dryopteris.
irregulare Brack. Expl. Exp. **16**. 180. 1854 = A. latifolium.
(Ar) **irregulare** (Pr.) C. Chr. Ind. 78. 1905. — Malesia-Ins. Philipp. Tenasserim. Polypodium Pr. Rel. Haenk. **1**. 25 t. 4 f. 3. 1825; Dictyopteris Pr. 1836; Phegopteris Mett. 1864. Aspidium difforme Bl. 1828. NPfl. 186; Polypodium Bl. 1828. HB. 318; Phegopteris Mett. 1856. n. 67. Chr. 267; Dictyopteris Moore 1858; Arcypteris Und. 1903.
irriguum Sm. Engl. Bot. **31** t. 2199. 1810 = Athyrium filix femina.
(S) **irriguum** J. Sm. JoB. **3**. 410. 1841 (nomen); Pr. Epim. 62. 1849. NPfl. 185 — Ins. Philipp. — Nephrodium Bak. 1867. HB. 297; Dryopteris O. Ktze. 1891; Microsorium trifidum Fée 1850-52.
isabellinum Fée, Cr. vasc. Br. **1**. 137 t. 45 f. **2** 1869 = Dryopteris.
jaculosum Christ, Bull. Boiss. II. **4**. 615. 1904 = Dryopteris.
(S) **Jardini** Mett.; Kuhn, Linn. **36**. 121. 1869 — Ins. Marquesas.
javanicum Mett. Pheg. u. Asp. 103. n. 245. 1858 = Mesochlaena polycarpa.
javense Willd. sp. **5**. 284. 1810 = Dryopteris punctata.
Jenmani Christ, Engl. Jahrb. **24**. 116. 1897 = Dryopteris.
jucundum Fée, 10 mém. 41 t. 42 f. 1. 1865 = Polystichum denticulatum.
juglandifolium Kze.; Kl. Linn. **20**. 363. 1847 = Phanerophlebia.
juglandifolium Christ, Engl. Jahrb. **23**. 354. 1896 = Nephrodium.
juglandifolium v. macrosorum Bak. JoB. **1887**. 25 = Phanerophlebia macrosora.
(S) **kanakorum** (Fourn.) C. Chr. Ind. 78. 1905 — N. Caledonia. Bathmium Fourn. Ann. sc. nat. V. **18**. 301. 1873; Nephrodium Bak. 1891.
Karsteni A. Br. Ind. sem. ht. Berol. 1856 — Caracas. (vix Lastrea similis J. Sm.).
Karwinskyanum Mett. Pheg. u. Asp. 59. n. 141. 1858 = Dryopteris.
Kaulfussii Link, Hort. Berol. **2**. 117. 1833 = Dryopteris oligocarpa*.
(S) **Keckii** Luerss. Bot. Centralbl. **11**. 76. 1882 — Sumatra. Nephrodium Bak. 1891.
Kehdingiana Luerss.; Christ, Farnkr. 235. 1897 = Luerssenia.
kiboschense Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
Kilemense Kuhn, v. Deck. Reisen **3**[3]. Bot. 46. 1879 } = Dryopteris kilemensis.
Kilmense Kuhn, Fil. Deck. 24. 1867 (err. typ.) }
Kingii Hance, Ann. sc. nat. IV. **18**. 237. 1862 = Cyclopeltis Presliana var.
Klotzschii Hk. Ic. pl. t. 923. 1854 = Polystichum denticulatum *laetum.
Koordersii Christ, Ann. Jard. Buit. **15**. 128. t. 15 f. 7. 1897 = Dryopteris.
Krameri Christ, Bull. Soc. Fr. **52** Mém. I. 37. 1905 = Dryopteris oyamensis.
(Ar) **Kuhnii** C. Chr. Ind. 78. 1905 — Malacca. Phegopteris polycarpa Mett.; Kuhn, Linn. **36**. 124. 1869 (non Aspidium Bl. 1828); Polypodium Bak. 1874. HB. 506; Dictyopteris Bedd. 1876.
(T) **Künstleri** Bedd. Handb. Suppl. 44. 1892 — Perak.
Kunzei Fée, 10 mém. 37 t. 41 f. A, B. 1865 = Dryopteris oochlamys.
Labordei Christ, Bull. Soc. Fr. **52** Mém. I. 40. 1905 = Dryopteris.
(S) **labrusca** (Hk.) Christ, Ann. Jard. Buit. **15**. 137. 1897. NPfl. 185 — Java, Celebes, Borneo. — Polypodium Hk. sp. **5**. 73 t. 285 B. 1863. HB. 361; Colysis J. Sm. 1875; Nephrodium Racib. 1902.
lacerum Sw. Schrad. Journ. **1800**[2]. 39. 1801 = Dryopteris.

Lachenense Hk. sp. **4**. 8 t. 212. 1862 = Polystichum.
laetevirens Lowe; Kze. Linn. **23**. 227. 1850 (nomen) = Polystichum Webbianum.
laetum Sw. Vet. Ak. Hdl. **1817**. 63 t. 4 f. 3 = Polystichum denticulatum*.
laetum Moritz, Bot. Zeit. **1854**. 855 = Dryopteris effusa *exculta.
laeve Mett. Pheg. u. Asp. 104. n. 249. 1858 = Dryopteris.
lancastriense Spr. Anleit. **3**. 134. 1804 = Dryopteris cristata.
lanceolatum Bak. Gard. Chr. n. s. **14**. 494. 1880 = Polystichum.
lanceum Kze. Bot. Zeit. **1846**. 473 = Athyrium macrocarpum.
lanigerum Kuhn, Fil. Afr. 135. 1868 = Dryopteris.
lanosum Sw. Syn. 58. 1806 = Cheilanthes.
lanuginosum Willd.; Klf. Enum. 244. 1824 = Dryopteris.
laserpitiifolium Mett. Ann. Lugd. Bat. **1**. 227 t. 6. f. 3. 1864 = Polystichum Standishii.
lasiesthes Kze. Linn. **23**. 227, 300. 1850 = Dryopteris mollicula.
Lastii Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
lastreoides Mett.; Salom. Nom. 57. 1883 = Pronephrium.
latebrosum Kze. Bot. Zeit. **1848**. 261 = Dryopteris.
latifolium Pr. Rel. Haenk. **1**. 30. 1825 = A. cicutarium*.

(S) **latifolium** (Forst.) J. Sm. JoB. **3**. 410. 1841. NPfl. 185 — Celebes. Polynesia.
Polypodium Forst. Prod. 83. 1786; Phymatodes Pr. 1836; Drynaria Fée 1850-52; Pleopeltis Moore 1862; Nephrodium Bak. 1867. HB. ed. I. 297, ed. II. 298; Sagenia Carr. 1873; Dryopteris O. Ktze. 1891. Aspidium Forsteri Kze. 1846. Mett. n. 295 — A. irregulare Brack. 1854; Nephrodium Bak. 1867 — Aspidium hymenodes Mett.; Kuhn 1869 — ? Bathmium Seemanni Fourn. 1873.

latifrons Mann, Pr. Amer. Acad. **7**. 217. 1868 = Dryopteris.
latipinna Hance, Journ. Linn. Soc. **13**. 141. 1873 = Dryopteris parasitica var.
latissimum Fée, Cr. vasc. Br. **1**. 142 t. 48 f. 2. 1869 = Dryopteris amplissima.
latum Kze. Mett. Pheg. u. Asp. 95. n. 231. 1858 = Dryopteris crassifolia.

(S) **Lawrenceanum** (Moore) Diels, Nat. Pfl. 1^4. 185. 1899 — Madagascar.
Sagenia Moore, Gard. Chr. n. s. **15**. 9. 1881; Nephrodium Bak. 1884.

laxum Franch. et Sav. En. pl. Jap. **2**. 237, 631. 1879 = Dryopteris.
Lechleri Mett. (Hk. sp. **4**. 21. 1862) = Polystichum aculeatum var. 4.
lentum Don, Prod. Fl. Nepal. **4**. 1825 = Polystichum.
lepidocaulon Hk. sp. **4**. 12 t. 217. 1862 = Polystichum.
lepidotrichum Desv. Berl. Mag. **5**. 321. 1811 = Dryopteris hirta.
Lepinei Kuhn, Linn. **36**. 118. 1869 = Dryopteris.

(Ar) **Leprieurii** Mett.; Diels, Nat. Pfl. 1^4. 187. 1899 — Panama-Guiana.

leptorachis Kze. Linn. **18**. 346. 1844 = Dryopteris patula.
Lessonii Bory, Dup. Voy. Bot. **1**. 265. 1828 — Tahiti.
leuconeuron Kuhn, Fil. Afr. 136. 1868 = Nephrodium.
leucophlebium Christ, Bull. Boiss. II. **4**. 961. 1904 = Dryopteris.
leucopis Fée, Gen. 292. 1850-52 (err.: »leucolepis«) = Dryopteris setigera.
leucosticton Kze. Linn. **23**. 227; 301. 1850 = Dryopteris orientalis.
leucostipes Christ, Bull. Boiss. II. **4**. 615. 1904 = Dryopteris.

(Pl) **Leuzeanum** (Gaud.) Kze. Bot. Zeit. **1846**. 474. Mett. n. 276. Chr. 231. NPfl. 187 — China austr. Ind. bor.-Malesia-Polynesia.
Polypodium Gaud. Freyc. Voy. Bot. 361 t. 6. 1827; Pleocnemia Pr. 1836; Nephrodium Hk. 1862. HB. 295; Dryopteris O. Ktze. 1891.

Levyi Fourn. Bull. Soc. Fr. **19**. 255. 1872 = Dryopteris.

L'Herminieri Kze.; Mett. Pheg. u. Asp. 85. n. 205. 1858 = Dryopteris.
libanoticum Rosenstock, Mém. l'Herb. Boiss. n. 9. 1900 = Dryopteris.
(Ar) **lifuense** (Fourn.) C. Chr. Ind. 80. 1905 — N. Caledonia.
Dictyopteris Fourn. Ann. sc. nat. V. 18. 291. 1873.
ligulatum Kze.; Mett. Pheg. u. Asp. 88. n. 213. 1858 = Dryopteris.
ligusticifolium Desv. Prod. 250. 1827 = Polystichum denticulatum.
limbatum Sw. Schrad. Journ. **1800**². 35. 1801 = Dryopteris.
Lindeni Kuhn, Linn. **36**. 116. 1869. = Dryopteris.
Lindeni Fourn. Mex. pl. **1**. 97. 1872 = Dryopteris Moreletii.
lineatum Bl. Enum. 144. 1828 = Dryopteris.
Linkii A. Br. Ind. sem. ht. Berol. 1856 — Amer. merid.
A. concinnum Link, Hort. Berol. **2**. 120. 1833 — (Dryopteris sp. dubia).
(S) **Lizarzaburui** (Sod.) C. Chr. Ind. 80. 1905 — Ecuador. — Nephrodium Sod. Rec. 55. 1883; Cr. vasc. quit. 272. 1893 (-oi, ex err.; corr. p. 640).
lobatum Sw. Schrad. Journ. **1800**². 37. 1801 = Polystichum.
lobatum × aculeatum Christ, Farnkr. d. Schw. 124. 1900 = Polystichum aculeatum × lobatum.
lobatum × Braunii Luerss. in Rabenh. Krypt. Fl. ed. II. **3**. 357 f. 143. 1884-89 = Polystichum Braunii × lobatum.
lobatum × Lonchitis Murbeck, Lunds Univ. Årsskr. **27**⁵. 16. 1892 = [Polystichum.
(S) **Lobbii** Hk. sp. **4**. 59 t. 232. 1862 — Borneo.
Nephrodium Bak. 1867. HB. 297; Dryopteris O. Ktze. 1891.
lobulatum Bl. Enum. 142. 1828 = A. decurrens vel A. singaporianum.
lobulatum Christ, Bull. Boiss. II. **4**. 614. 1904 = Dryopteris taiwanensis.
Loherianum Christ, Bull. Boiss. **6**. 191. 1898 = Dryopteris.
lomatopelta Kze.; Mett. Pheg. u. Aspid. 72 n. 172. 158 = Dryopteris.
lomatopus Kze. Bot. Zeit. **1848**. 238 = Oleandra neriiformis.
lonchitiforme Halacsy, Consp. Fl. Graec. **3**. 475. 1904 = Polystichum aculeatum × lonchitis.
Lonchitis Sw. Schrad. Journ. **1800**². 30. 1801 = Polystichum.
Lonchitis × lobatum Aschers. Syn. Mitt. Eur. Fl. **1**. 42. 1896 = Polystichum lobatum × lonchitis.
lonchitoides Christ, Bull. Boiss. **7**. 16. 1899 = Polystichum.
lonchodes Eat. Mem. Amer. Acad. n. s. **8**. 210. 1860 = Dryopteris.
longicuspe Kuhn, v. Deck. Reisen **3**³. Bot. 65. 1879 = Dryopteris.
longifolium Desv. Berl. Mag. **5** 319. 1811 = A. martinicense.
longipes Bl. Enum. 155. 1828 = Dryopteris.
Lorentzii Hier. Engl. Jahrb. **22**. 368. 1896 = Dryopteris.
lorifrons Kze. Farnkr. **1**. 36. 1840 (in nota) = Oleandra neriiformis.
lucens Boj.; Kze. Linn. **23**. 228. 1850 = ? Dryopteris gongylodes.
luctuosum Kze. Linn. **10**. 548. 1836 = Polystichum lobatum var. 1.
ludens Christ, Engl. Jahrb. **23**. 352. 1896 = Dryopteris.
Ludovicianum Kze. Amer. Journ. Sci. II. **6**. 93. 1848 = Dryopteris filix mas*.
Luerssenii Dörfler, Oest. Bot. Zeit. **40**. 227. 1890 = Polystichum Braunii [× lobatum.
Luerssenii Christ, Bull. Boiss. **6**. 193. 1898 = Dryopteris.
lugubre Mett. Pheg. u. Asp. 94. n. 230. 1858 = Dryopteris.
Lunanense Christ, Bull. Boiss. **6**. 966. 1898 = Dryopteris.
lutescens Willd. sp. **5**. 272. 1810 = Polypodium pulverulentum.
Macleaii Bak. in Hk. Ic. pl. t. 1654. 1886 = Polystichum.
macrocarpon Bl. Enum. 162. 1828 = Athyrium.

macrochlamys Fée, Gen. 295. 1850-52 = Dryopteris sparsa.
(Ar) **macrodon** (Reinw.) Keys. Pol. Cyath. Hb. Bung. 40. 1873. NPfl. 186 — Malesia-Ins. Philipp.-Fiji. — Polypodium Reinw. hb. HB. 318. 1867; Dictyopteris Pr. 1836 (nomen); Phegopteris Mett. Pheg. u. Asp. 31 n. 68. 1858. Chr. 266. Dictyopteris Cumingiana Pr. 1849 (nomen); Polypodium Hk. 1864.
macrolepidum Desv. Prod. 248. 1827 = Polystichum aculeatum ? var. 9.
macrophyllum Rudolphi 1805; Sw. Syn. 43, 239. 1806 = A. martinicense.
macrophyllum Bl. Enum. 144. 1828 = A. pachyphyllum.
macropus Mett. Fil. Lechl. **2**. 20. 1859 = Dryopteris Swartzii var.
(S) **macrosorum** (Bak.) Diels, in Schum. et Laut. Fl. deut. Südsee 115. 1901 — Ins. Salomonis. — Nephrodium Bak. Journ. Linn. Soc. **19**. 295. 1882.
macrosorum Christ, Bull. Boiss. II. **4**. 963. 1904 = Phanerophlebia.
macrostegium Fée, Cr. vasc. Br. **2**. 76. 1872-73 = Dryopteris.
macrourum Klf. Flora **1823**[1]. 365 = Dryopteris patens*.
macrum Fée, Cr. vasc. Br. **1**. 142 t. 48 f. 1. 1869 = Dryopteris amplissima var.
Madagascariense Fée, Gen. 296. 1850-52 — Madagascar.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
Malaccense Fée, Gen. 292. 1850-62 = ? Dryopteris extensa.
malgassicum Kuhn, Fil. Afr. 136. 1868 = Dryopteris madagascariensis.
(S) **mamillosum** (Moore) C. Chr. Ind. 81. 1905 — Ins. Moluccae.
Sagenia Moore, Gard. Chr. n. s. **26**. 38. 1886; L'Ill. Hort. IV. **6**. t. 598; Nephrodium Bak. 1891.
manilense Mett.; Salom. Nom. 59. 1883 = Lastrea.
Manmeiense Christ, Bull. Boiss. **6**. 965. 1898 = Polystichum nepalense.
maranguense Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
marginale Sw. Schrad. Journ. **1800**[2]. 35. 1801 = Dryopteris.
marginatum Schkuhr, Kr. Gew. **1**. 195 t. 45 b. 1809 = Dryopteris marginalis.
marginatum Wall. List n. 366. 1828 = Polystichum nepalense.
marginatum Wall. List n. 391. 1828 = Dryopteris filix mas* et ? D. punctata.
marginiferum Mett.; Salom. Nom. 59. 1883 = Hypopeltis.
(S) **martinicense** Spr. Anleit. **3**. 133. 1804 — Amer. trop.
Dryopteris O. Ktze. 1891; Aspidium macrophyllum Rudolphi 1805; Sw. 1806. Mett. n. 289. Chr. 231. NPfl. 186; Bathmium Link 1841; Cardiochlaena Fée 1850-52; Sagenia Moore 1857; Nephrodium Bak. 1867. HB. 300 — Aspidium repandum Vahl 1807 — A. bifidum Pr. 1822 — Cardiochlaena pilosa Fée 1850-52. Cardiochlaena sp. plur. Fée. (an Polypodium expansum Poir. 1804?).
mascarenense Klf.; Spr. Syst. **4**. 101. 1827 = Dryopteris truncata.
mascarenense Kuhn, v. Deck. Reis. **3**[3]. Bot. 65. 1879 = Dryopteris mascarensis.
Mascarenhense Fée, Gen. 296. 1850-52 = Dryopteris crinita.
Matsumurae Mak. Bot. Mag. Tokyo —. 1892 = Dryopteris.
mauritianum Desv. Berl. Mag. **5**. 320. 1811 = Nephrolepis biserrata.
mauritianum Kuhn, Fil. Afr. 136. 1868 = Dryopteris.
Maximowiczianum Miq. Ann. Lugd. Bat. **3**. 178. 1867 = Polystichum aristatum.
maximum Bojer; Bak. Syn. 502. 1874 = Dryopteris mauritiana var.
medium Carm. Tr. Linn. Soc. **12**. 511. 1818 = Athyrium.
(Pl) **megalocarpum** (Hk.) C. Chr. Ind. 81. 1905 — Java. Perak.
Polypodium Hk. sp. **5**. 102. 1864. HB. 318; Dictyopteris J. Sm. 1875; Phegopteris Salomon 1883; Pleocnemia Bedd. 1892 — Nephrodium oligodictyon Bak. 1891.

megalodus Mett. Fil. Lechl. **2**. 21. 1859 = Dryopteris.

megaphyllum Mett. Ann. Lugd. Bat. **1**. 233. 1864 = Dryopteris.

megaphyllum Christ, Bull. Boiss. **6**. 194. 1898 = Dryopteris subarborea.

(S) **melanocaulon** Bl. Enum. 161. 1828 — China austr. Tonkin. Malesia. Ins. Philipp. N. Guinea. — Sagenia Moore 1858; Nephrodium Bak. 1867. HB. 296; Dryopteris O. Ktze. 1891 — ? Aspidium Durvillaei Bory 1828 — Cardiochlaena subpinnatifida Fée 1850-52.

melanocaulon F. Muell. Fragm. **5**. 133. 1865 = A. Muelleri.

melanochlamys Fée, Gen. 294. 1850-52 = Polystichum.

(T ?) *melanopodon* Desv. Berl. Mag. **5**. 320. 1811 — Terrae Magellanicae.

melanopus Spr. Syst. **4**. 101. 1827 = A. melanopodon.

(S) **melanorachis** (Bak.) C. Chr. Ind. 82. 1905 — Borneo.
Nephrodium Bak. JoB. **1888**. 325.

melanorhizum Desv. Prod. 251. 1827 = A. cicutarium.

melanorhizum Christ, Bull. Soc. bot. It. **1901**. 295 = Dryopteris subthe-

melanostictum Kze. Linn. **13**. 148. 1839 = Polystichum. [lypteris.

membranaceum Fée, Gen. 292. 1850-52 = A. membranifolium vel Dryo-

membranaceum Hk. sp. **5**. 105. 1864 = A. devexum. [pteris dissecta.

(Pl) **membranifolium** (Pr.) Kze. Bot. Zeit. **1848**. 261. Mett. n. 270. NPfl. 188 — China austr. India bor. Birma-Malacca. ? Malesia. Ins. Philipp.
Nephrodium Pr. Rel. Haenk. **1**. 36. t. 5 f. 3. 1825; Lastrea Pr. 1836; Pleocnemia Bedd. 1883; Sagenia Christ 1902; Aspidium fuscipes pt. Wall. 1828; Bedd. 1876; Nephrodium Clarke 1880. N. dissectum pt. HB. 282 — Polypodium chattagrammicum Clarke 1880; Dictyopteris Bedd. 1883; Aspidium Diels 1899. NPfl. 186.

menisciicarpon Bl. Enum. 142. 1828 = Dryopteris lineata.

menisciicarpon Mett. Pheg. u. Asp. 121. n. 287. 1858 = A. siifolium.

menisciinerve Gaud. Freyc. Voy. Bot. 244. 1827 — Ins. Moluccae.
Nephrodium Moore 1858 — (Dryopteris sp. dub.).

meniscioides Willd. sp. **5**. 218. 1810 = Cyclodium.

(S) **menyanthidis** Pr. Rel. Haenk. **1**. 28. 1825. Mett. n. 293 — N. Guinea. Ins. Philipp. Ins. Salomonis.
Polydictyum Pr. 1849; Cardiochlaena Fée 1850-52; Sagenia Moore 1858; Nephrodium (Meyanthidis) Bak. 1867. HB. 297; Dryopteris O. Ktze. 1891.

mexicanum Kze. Linn. **13**. 147. 1839 = Dryopteris patula.

micranthum Bl. Enum. 141. 1828 = Oleandra neriiformis.

microcarpon Bl. Enum. 142. 1828 = Polypodium punctatum.

microcarpon Fée, 8 mém. 105. 1857 = Dryopteris microchlaena.

microcarpum Kuhn, Fil. Afr. 136. 1868 = Polypodium.

microchlaena Fée, 8 mém. 102. 1857 = Dryopteris.

microchlamys Christ, Bull. Boiss. **7**. 830. 1899 = Polystichum Bakerianum.

microphyllum Bl. Enum. 163. 1828 = Polystichum aculeatum var. 32.

microsorum Klf. in Sieb. Syn. Fil. exs. n. 101 = Dryopteris decomposita.

microsorum Pr. Epim. 61. 1849 = A. melanocaulon vel A. latifolium.

microsorum Kuhn, Fil. Afr. 136. 1868 = Dryopteris unita.

microthecium Fée, 10 mém. 37 t. 41 f. 2 A. 1865 = ? Dryopteris syrmatica.

Mildeanum Göppert, Denkschr. 50 jähr. Best. vat. Ges. Schles. 193. 1853 = Dryopteris filix mas.

Milnei Luerss. Fil. Graeff. 184. 1871 = Dryopteris.

Miquelianum Maxim.; Franch. et Sav. En. pl. Jap. **2**. 240, 634. 1879 = Dryopteris.

mohrioides Bory, Mém. Soc. Linn. Paris **4**. 597 —. Dup. Voy. Bot. **1**. 267 t. 35 f. 1. 1828 = Polystichum.
mohrioides auctt. quoad pl. amer. bor. = Polystichum Lemmoni.
molle Sw. Schrad. Journ. **1800**[2]. 34. 1801 = Dryopteris parasitica (vel D. patens).
molliculum Mett. Fil. Lechl. **1**. 19. 1856 (an Fourn. Mex. pl. **1**. 94. 1872?) = Dryopteris.
mollissimum Christ, Bull. Boiss. **6**. 968. 1898 = Dryopteris.
molliusculum Kuhn, Bot. Zeit. **1868**. 41 = Dryopteris.
moluccense Bl. Enum. 168. 1828 = Polystichum aculeatum var. 33.
mongolicum Franch. »Pl. David. exsicc. n. 2273« = Dryopteris.
monostichum Kze.; Mett. Pheg. u. Asp. 94. n. 228. 1858 = Dryopteris tristis var.
monotis Christ, Bull. Soc. bot. It. **1901**. 294 = Polystichum.
montanum Sw. Schrad. Journ. **1800**[2]. 42. 1801. = Cystopteris.
montanum Aschers. Fl. Brand. **3**. 133. 1859 = Dryopteris oreopteris.
montevidense Hier. Engl. Jahrb. **22**. 364. 1896 = Polystichum aculeatum var. 9.
monticola Christ, Bull. Soc. Fr. **52** Mém. I. 39. 1905 = Dryopteris.
Moorei (Hk.) Diels, Nat. Pfl. **1**[4]. 186. 1899 — N. Caledonia.
Deparia Hk. JoB. **4**. 55 t. 3. 1852. HB. 55. (Chr. 230); Cionidium Moore 1852; Trichiocarpa J. Sm. 1856.
Moritzianum Kl. Linn. **20**. 367. 1847 = Polystichum aculeatum var. 4.
Moritzii Kze. Bot. Zeit. **1848**. 238 = Oleandra neriiformis.
Mulmeinense Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 247 = Dryopteris moulmeinensis.
moupinense Franch. Nouv. Arch. Mus. II. **10**. 115. 1887 = Polystichum.
mucronatum Sw. Schrad. Journ. **1800**[2]. 30. 1801 = Polystichum echinatum.
mucronatum Don, Prod. Fl. Nepal. 5. 1825 = Polystichum aristatum var.
mucronatum Beyrich; Mett. Pheg. u. Asp. 90 n. 219 b. 1858 = Dryopteris falciculata.
mucronifolium Bl. Enum. 164. 1828 = Polystichum aculeatum var. 32.
(mucronulatum Opiz; Steud. Nomencl. **2**. 63. 1824 —?).
(S) **Mülleri** C. Chr. Ind. 83. 1905 — Queensland.
A. confluens Mett.; Kuhn, Linn. **36**. 125. 1869 (non Fée 1850-52); Nephrodium Müll. msc. Bak. 1874. HB. 504; Dryopteris O. Ktze. 1891; Aspidium melanocaulon F. Müll. 1865 (non Bl. 1828); Sagenia Bail. 1874.
Münchii Christ, Bull. Boiss. II. **1**. 1120. 1901 = Polystichum.
(S) **multicaudatum** Wall.; Bedd. Hdb. 222. 1883 — Ind. bor. (Khasia). Birma. Perak.
Nephrodium Clarke, Tr. Linn. Soc. II. Bot. **1**. 540 t. 77. 1880 — N. Griffithii Bak. Syn. 300. 1867; Sagenia Bedd. 1870; Aspidium Bedd. 1876 (non supra); Dryopteris O. Ktze. 1891.
multidentatum Wall. List n. 346. 1828 = Davallia.
multifidum Mett. Fil. Lechl. **1**. 20 t. 3 f. 14-17. 1856 = Polystichum.
multijugum Wall. List n. 348. 1828 = Dryopteris extensa.
multilineatum Wall. List n. 353. 1828 = Dryopteris truncata et D. megaphylla.
multilineatum Mett. Pheg. u. Asp. 108. n. 258. 1858 = Dryopteris Haenkeana.
multilineatum Christ, Farnkr. 249. 1897 = Dryopteris moulmeinensis.
multisorum Desv. Prod. 246. 1827 = A. trifoliatum.
munitum Klf. Enum. 236. 1824 = Polystichum.
munitum Sadl. Fil. ver. Hung. 34. 1830 = Polystichum aculeatum.
muricatum Willd. sp. **5**. 264. 1810 = Polystichum aculeatum var. 4.
(T) **Murrayi** Bak. Ann. Bot. **5**. 315. 1891 — Ins. St. Lucia.

musaefolium Bl. Enum. 141. 1828 = Oleandra.
muscosum Sw. Adnot. 68. 1829 = Dryopteris pubescens var.
muticum Franch. et Sav. En. pl. Jap. **2**. 240, 635. 1879 = Dryopteris.
(S) **myriosorum** Christ, Bull. Boiss. II. **5**. 256. 1905 — Costa Rica.
Napoleonis Fée, Gen. 292. 1850-52 = Dryopteris.
Natalense Fée, 8 mém. 102. 1857 = Dryopteris prolixa.
(S) **nebulosum** (Bak.) C. Chr. Ind. 84. 1905 — Sumatra.
Nephrodium Bak. JoB. **1880**. 213.
(Ar) **neglectum** (Mett.) C. Chr. Ind. 84. 1905 — Assam.
Phegopteris Mett.; Kuhn, Linn. **36**. 124. 1869.
nemophilum Kze. Linn. **9**. 95. 1834 = Dryopteris catocarpa.
nemorale Gray, Nat. Arr. br. Pl. **2**. —. 1821 = Dryopteris filix mas.
nemorosum Willd. sp. **5**. 255. 1810 = Dryopteris.
nepalense Spr. Syst. **4**. 97. 1827 = Polystichum.
nephrodioides Kl. Linn. **20**. 370. 1847 — Columbia. — Lastrea Moore 1858. — (Dryopteris sp. dub.; f. Salomon = Cyclodium meniscioides).
nephrodioides Hk. sp. **4**. 42 t. 235. 1862 = Dryopteris orbicularis.
nephrodioides Fée, Cr. vasc. Br. **1**. 138 t. 46 f. 1. 1869 = Dryopteris patens.
neriiforme Sw. Syn. 42, 237. 1806 = Oleandra.
nervatum Fée, Cr. vasc. Br. **1**. 136. 1869 = Dryopteris submarginalis.
nevadense Boiss. Elench. pl. nov. 93. 1838; (Milde, Bot. Zeit. **1868**. 360) = Dryopteris rigida var.
nevadense Eat. Ferns N. Am. **1**. 73 t. 10. 1878 = Dryopteris oregana.
(T) **nicaraguense** (Fourn.) Bak. Syn. 493. 1874. NPfl. 184 — Nicaragua.
Bathmium Fourn. Bull. Soc. Fr. **19**. 256. 1872; Dryopteris O. Ktze. 1891.
(Ar) **nicotianifolium** (Bak.) Diels, Nat. Pfl. **1**[4]. 186. 1899 — Ecuador.
Polypodium Bak. Syn. ed. I. 455. 1868; JoB. **1877**. 165.
(S) **nigrescens** Mett.; Kuhn, Fil. Afr. 137. 1868. Chr. 231. NPfl. 186 — Afr. occ. trop.
Nephrodium Bak. 1874. HB. 504; Dryopteris O. Ktze. 1891.
nigricaule Fée, 8 mém. 104. 1857 = ? Dryopteris sagenioides.
nigripes Bl. Enum. 162. 1828 = Athyrium.
nigripes Kl. Allgm. Gartenzeit. **1856**. 189 = Hort. (an A. cicutarium?).
Nigritianum Mett.; Kuhn, Fil. Afr. 138. 1868 = Dryopteris tomentella.
nigropunctatum Spr. Neue Entd. **3**. 7. 1822 = Nephrolepis biserrata.
nipponicum Franch. et Sav. Enum. pl. Jap. **2**. 242, 636. 1879 = Dryopteris.
nitidulum Wall. List n. 392. 1828; Kuhn, Linn. **36**. 117. 1869 = Dryopteris sparsa var.
nitidulum (Pr.) Kze.; Mett. Pheg. u. Asp. 105. n. 249 c. 1858 — Ins. Philipp.
Nephrodium Pr. Epim. 46. 1849. — (Dryopteris sp. dub.).
nitidulum Kze.; Ettingsh. Farnkr. 188 t. 133 f. 3. 1865 = Dryopteris anoptera.
nitidum Bory; Willd. sp. **5**. 251. 1810 = Dryopteris crinita var.
nobile Schlecht. et Cham. Linn. **5**. 610. 1830 = Phanerophlebia.
nodosum Willd. sp. **5**. 211. 1810 = Oleandra.
nodosum Bl. Enum. 171. 1828 = Acrophorus stipellatus.
(Pl) **novae caledoniae** C. Chr. Ind. 84. 1905 — N. Caledonia.
Pleocnemia Vieillardii Fourn. Ann. sc. nat. V. **18**. 299. 1873 (non Aspidium Mett. 1861); Nephrodium Fournieri Bak. Ann. Bot. **5**. 328 (non 317). 1891.
Novae-Zeelandiae (Pr.) Ettingsh. Denkschr. Ak. Wien **23**. 103. 1864 — N. Zealand.
Goniopteris Pr. hb. — (an Dryopteris pennigera?).

noveboracense Sw. Schrad. Journ. **1800²**. 38. 1801 = Dryopteris.
(S) **nudum** (Bak.) Diels in Schum. et Laut. Fl. deut. Südsee 116. 1901 — Borneo; Bismarck's Archip. — Nephrodium Bak. JoB. **1879**. 41.
nymphale Schkuhr, Kr. Gew. **1**. 36 t. 34. 1806 = Dryopteris parasitica.
nymphale Bl. Enum. 157. 1828 = Dryopteris procurrens.
obliquatum Mett. Ann. sc. nat. IV. **15**. 75. 1861 = Dryopteris.
obliquum Don, Prod. Fl. Nepal. 3. 1825 = Polystichum.
obliteratum Spr. Syst. **4**. 99. 1827 = Arthropteris.
obscurum Bl. Enum. 150. 1828 = Dryopteris arida.
obscurum Fisch. et Mey. Ind. sem. ht. Petr. 1833; Ann. sc. nat. II. **5**. 128. 1836 et in Linn. **10**. Litteraturb. 83. 1836; Regel, Ind. sem. ht. Petr. 1855 = Dryopteris Fischeri.
obtusatum Sw. Schrad. Journ. **1800²**. 33. 1801 = Dryopteris pteroides.
obtusatum Schkuhr, Kr. Gew. **1**. t. 33 c. 1806; Bl. Enum. 151. 1828 = Dryopteris gongylodes.
obtusifolium Willd. sp. **5**. 231. 1810 = Nephrolepis radicans.
obtusilobum Fée, 8 mém. 105. 1857 = Dryopteris huatuscensis.
obtusilobum Prantl, Verh. zool. bot. Ges. **31**. 119. 1882 = Dryopteris peranemiformis var.
obtusissimum Mett.; Kuhn, Linn. **36**. 119. 1869 = Dryopteris sparsa var.
obtusum Sw. Syn. 420. 1806 = Woodsia.
obtusum Kze.; Mett. Pheg. u. Asp. 51. n. 120. 1858 = Polystichum.
ocellatum Wall. List n. 360. 1828 = Polystichum lentum.
ochthodes Kze. Linn. **24**. 282. 1851 = Dryopteris.
oculatum Hk. sp. **4**. 24 t. 228. 1862 = Polystichum Richardi var.
odoratum Bory; Willd. sp. **5**. 286. 1810 = Dryopteris crenata.
odoratum Mett. Pheg. u. Asp. 115. n. 273. 1858 = Dryopteris lanuginosa.
odoriferum Gray, Nat. Arr. br. Pl. **2**. 6. 1821 = Dryopteris oreopteris.
Okuboanum Mak. Bot. Mag. Tokyo —. 1892 = Athyrium.
oliganthum Desv. Berl. Mag. **5**. 321. 1811 = Dryopteris filix mas *oligodonta.
oligocarpum HBK. Nov. Gen. et Sp. **1**. 13. 1815 = Dryopteris.
oligodonton Desv. Berl. Mag. **5**. 322. 1811 = Dryopteris filix mas*.
oligophlebium Christ, Bull. Boiss. II. **4**. 616. 1904 = Dryopteris.
opacum Benth. Fl. Hongk. 456. 1861 = Polystichum varium.
Opizii Wierzb. Flora **1840**. 379 = Dryopteris filix mas.
oppositum Klf.; Spr. Syst. **4**. 108. 1827 = Dryopteris mascarenarum.
oppositum Sw. Adnot. 67. 1829 = Dryopteris.
opulentum Klf. Enum. 238. 1824 = Dryopteris pteroides.
orbiculatum Desv. Berl. Mag. **5**. 321. 1811 = Polystichum aculeatum var. 13.
ordinatum Kze. Linn. **18**. 347. 1844 = Polystichum aculeatum var. 4.
Oreopteris Sw. Schrad. Journ. **1800²**. 35. 1801 = Dryopteris.
(S) **organense** C. Chr. Ind. 85. 1905 — Brasilia.
Nephrodium Gardneri Bak. Ann. Bot. **5**. 330. 1891: Aspidium Christ 1897. 229. NPfl. 186 (non A. Gardnerianum Mett. 1858).
orientale Desv. Prod. 250. 1827 = Polystichum adiantiforme.
Orizabae Fée, 8 mém. 104. 1857 = Dryopteris.
ornatum Christ, Ann. Jard. Buit. **15**. 120. 1897 = Dryopteris.
Oshimense Christ, Bull. Boiss. II. **1**. 1018. 1901 = Dryopteris.
Otaria Kze.; Mett. Pheg. u. Asp. 34. n. 73. 1858 = Dryopteris.
otarioides Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 247 = Dryopteris.

otophorum Franch. Nouv. Arch. Mus. II. **10**. 116. 1887 = Polystichum.
oxyodon Franch. Nouv. Arch. Mus. II. **7**. 163. 1883 = Dryopteris filix mas*.
pachychlamys Fée, 11 mém. 77 t. 21 f. **2**. 1866 = Dryopteris opposita.
(S) **pachyphyllum** Kze. Bot. Zeit. **1848**. 259. Mett. n. 286. Chr. 229. NPfl. 185 — Malesia-Polynesia. — Sagenia Moore 1857; Nephrodium Bak. 1867. HB. 299; Dryopteris O. Ktze. 1891.
pachyrachis Kze.; Mett. Pheg. u. Asp. 83. n. 199. 1858 = Dryopteris.
paleaceum Sw. Syn. 52. 1806 = Dryopteris filix mas*.
paleaceum Don, Prod. Fl. Nepal. 4. 1825 = Dryopteris filix mas *patentissima.
Palisoti Desv. Berl. Mag. **5**. 320. 1811 = Arthropteris obliterata.
pallens Mett.; Salom. Nom. 63. 1883 = Lastrea.
pallidivenium Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris striata.
pallidum Bl. Enum. 169. 1828 — Ins. Moluccae.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
pallidum Link, Fil. sp. 107. 1841 = Dryopteris rigida *australis.
pallidum Fourn. Mex. pl. **1**. 96. 1872 = Dryopteris chlorophylla.
(T) **palmatum** Mett. Ann. Lugd. Bat. **1**. 238. 1864 — Borneo. Sumatra.
palmipes Kze. Linn. **24**. 287. 1851 = Polystichum carvifolium.
paludosum Raddi, Pl. bras. **1**. 29 t. 44. 1825 = Nephrolepis biserrata.
paludosum Bl. Enum. 168. 1828 = Dryopteris brunnea.
paludosum Mett. Pheg. u. Asp. 101. n. 239 b. 1858 = Dryopteris gongylodes.
palustre Gray, Nat. Arr. br. Pl. **2**. 9. 1821 = Dryopteris thelypteris.
palustre Fée, Cr. vasc. Br. **2**. 76. 1872-73 = Dryopteris.
pandiforme Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 255 = Dryopteris.
paradoxum Fée, Gen. 293. 1850-52 — Ceylon.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
paraënse Willd. sp. **5**. 228. 1810 = Nephrolepis biserrata.
parallelogrammum Kze. Linn. **13**. 146. 1839 = Dryopteris filix mas var.
parallelum Desv. Prod. 245. 1827 — »Amer. trop.«
parallelum Kuhn, v. Deck. Reisen **3**³. Bot. 65. 1879 = Dryopteris.
parasiticum Sw. Schrad. Journ. **1800**². 35. 1801 = Dryopteris.
parathelypteris Christ, Bull. Soc. Fr. **52** Mém. I. 36. 1905 = Dryopteris.
patens Sw. Schrad. Journ. **1800**². 34. 1801 = Dryopteris.
patens Bl. Enum. 157. 1828 et auctt. plur. = Dryopteris parasitica.
patens β Sprengelii Kze. Linn. **10**. 547. 1836 = Dryopteris Bergiana.
patentissimum Wall. List n. 340. 1828 = Dryopteris filix mas*.
patentissimum Franch. Nouv. Arch. Mus. II. **10**. 118. 1887 = Dryopteris filix [mas *khasiana.
patulum Sw. Vet. Ak. Hdl. **1817**. 64 = Dryopteris.
paucicuspe Sturm, En. pl. cr. vasc. chil. —. 1858 = Polystichum aculeatum.
pauciflorum Klf.; Spr. Syst. **4**. 101. 1827 = Dryopteris crinita.
paucijugum Kl. Linn. **20**. 368. 1847 — Columbia.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
pauper Fée, 8 mém. 103. 1857 = Dryopteris patens.
paupertinum Röm.; Kze. Linn. **18**. 345. 1844 = Dryopteris patula.
pectinatum Willd. sp. **5**. 223. 1810 = Nephrolepis.
pectinatum Christ, Farnkr. 250. 1897 = Dryopteris orientalis.
pedatum Desv. Prod. 244. 1827 = Dryopteris.
pedicularis Kuhn, Fil. Afr. 138. 1868 = Polypodium.
pellitum Willd. sp. **5**. 242. 1810 = ? Dryopteris parasitica.
pellucidum Franch. Nouv. Arch. Mus. II. **10**. 119. 1887 = Dryopteris.

pendulum Raddi, Opusc. sci. Bol. **3**. 289. 1819; Pl. Bras. **1**. 30 t. 45. 1825 = Nephrolepis cordifolia*.
pendulum Splitgb. Tijds. Nat. Gesch. **7**. 412. 1840 = Oleandra neriiformis.
pennigerum Sw. Schrad. Journ. **1800²**. 34. 1801 = Dryopteris.
pennigerum Bl. Enum. 153. 1828 = Dryopteris megaphylla.
pentagonum Kuhn, Fil. Afr. 139. 1868 = Dryopteris inaequalis.
pentangularum Col. Tasm. Journ. **2**. —. 1845 = Dryopteris decomposita.
pentaphyllum Willd. sp. **5**. 216. 1810 — Martinique. (an A. trifoliatum?).
perakense Bedd. JoB. **1888**. 4 = Dryopteris.
perelegans Col. Tr. N. Zeal. Inst. **29**. 416. 1897 = Polystichum.
phaeochlamys Fée, Cr. vasc. Br. **1**. 140 t. 47 f. 2. 1869 = Dryopteris amplissima.
phaeostigma Cesati, Rend. Ac. Napoli **16**. 26, 29. 1877 = Dryopteris.
Phegopteris Baumg. Enum. stirp. Transs. **4**. 28. 1846 = Dryopteris.
philippinense Christ, Bull. Boiss. **6**. 191. 1898 = Dryopteris basilaris.
Philippinum Fée, Gen. 292. 1850-52 = Dryopteris ligulata.
phyllarthron Kze. Bot. Zeit. **1848**. 237 = Oleandra neriiformis.
Phyllitidis Roxb. Calc. Journ. **4**. 466. 1844 = A. singaporianum.
physematioides Kuhn et Christ, Engl. Jahrb. **24**. 115. 1897 = Dryopteris.
(S) **pica** (L.fil.) Desv. Berl. Mag. **5**. 319. 1811. Mett. n. 288. Chr. 227. NPfl. 185 — Ins. Mascarenae, Madagascar. ? Polynesia.
Polypodium L.fil. Suppl. 446. 1781; Sagenia Moore 1858; Nephrodium Bak. 1867. HB. 298; Dryopteris O. Ktze. 1891. Bathmium ebeneum Fée, 1850-52; Aspidium J. Sm. 1857. A. trifoliatum auctt. quoad pl. Mascar; Nephrodium Bory 1833.
pilosiusculum Mett. Ann. Lugd. Bat. **1**. 233. 1864 = Dryopteris appendiculata.
pilosulum Wall. List n. 337. 1828 = Dryopteris crenata et Microlepia speluncae.
pilosulum Kl. et Karst.; Kze. Linn. **23**. 229. 1850 = ? Dryopteris mollicula.
pilosum Langsd. et Fisch. Ic. Fil. 14 t. 16. 1810 = Nephrolepis hirsutula.
pilosum Schur, »Siebenb. Ver. **2**. 168. 1851«; Oest. Bot. Zeit. **8**. 195. 1858 = Polystichum Braunii.
pinnatifido-serratum Mett.; Kuhn, Fil. Afr. 139. 1868 = Dryopteris Welwitschii.
pistillare Sw. Schrad. Journ. **1800²**. 30. 1801 = Oleandra neriiformis.
(A) **plantagineum** (Jacq.) Gris. Abh. Ges. Wiss. Gött. **7**. 286. 1857. Mett. n. 296. HB. 258. Chr. 226. NPfl. 183 — Ind. occ.-Brasilia.
Polypodium Jacq. Coll. **2**. 104 t. 3 f. 1. 1788; Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Podopeltis Fée 1850-52; Dryomenis J. Sm. 1854; Pleopeltis Moore 1862; Bathmium Fourn. 1872; Dryopteris O. Ktze. 1891. Polypodium latifolium Vahl 1807 (non Forst. 1786). Bathmium macrocarpum Fée 1850-52 — B. sinuatum Fée 1850-52; Aspidium Moore 1858 (non Lab. 1824); Podopeltis J. Sm. 1875.
Plantianum Kuhn, Fil. Afr. 139. 1868 = Nephrodium.
Plaschnickianum Kze. Linn. **23**. 229, 302. 1850 = Polystichum.
(T) **platanifolium** Mett. Ann. Lugd. Bat. **1**. 239. 1864. HB. 258. NPfl. 185 — Malesia.
Dryopteris O. Ktze. 1891. Aspidium angulatum Hk. 1862 (non J. Sm.).
platynotus Kze. Linn. **23**. 229. 1850 = A. decurrens.
platyphyllum Willd. sp. **5**. 255. 1810 = Polystichum aculeatum var. 23.
platyphyllum Pr. Epim. 65. 1849 = A. repandum.
platypus Kze. Bot. Zeit. **1848**. 262 = Polystichum aristatum var.
platyrachis Fée, Cr. vasc. Br. **2**. 71 t. 100 f. 2. 1872-73 = Dryopteris.

(S) **pleiotomum** (Bak.) Kuhn, v. Deck. Reisen **3**[3]. Bot. 66. 1879 — Ins. Sechellae. Madagascar.
Nephrodium Bak. Syn. ed. II. 297. 1874; Dryopteris O. Ktze. 1891.
plicatum Poepp.; Kze. Linn. **9**. 94. 1834 = Polystichum mohrioides.
Plukenetii Steud. Nomencl. **2**. 64. 1824 = Polystichum lobatum.
(T) **Plumierii** Pr. Rel. Haenk. **1**. 29. 1825. Mett. n. 291. Chr. 227. NPfl. 185 — Ind. occ. — A. trifoliatum part. HB. 258 — an A. fimbriatum Willd. 1810?
plumiferum Kuhn, Forschr. Gazelle. **4**. Farne 9. 1889 = Dryopteris arida.
podophyllum Hk. JoB. **5**. 236 t. 1. 1853 = Dryopteris.
podophyllum Lowe, Ferns **6** t. 34-35. 1857 = Dryopteris Sieboldii.
(T) **Poeppigii** Pr. Epim. 62. 1849. Mett. n. 292 — Nicaragua. Peru.
Bathmium Fourn. 1872.
Pohlianum Pr. Del. Prag. **1**. 173. 1822 = Dryopteris gongylodes.
politum Desv. Prod. 250. 1827 = Polystichum adiantiforme.
polyblepharon Röm.; Kze. Bot. Zeit. **1848**. 572 = Polystichum lobatum var. 3.
polycarpon Bl. Enum. 156. 1828 = Mesochlaena.
polylepis Franch. et Sav. Enum. pl. Jap. **2**. 236, 631. 1879 = Dryopteris.
(S) **polymorphum** Wall. List n. 382. 1828; Hk. sp. **4**. 54. 1862. Chr. 229. NPfl. 185 — India bor. Birma-Malesia. — Nephrodium Bak. 1867. HB. 297; Dryopteris O. Ktze. 1891. Aspidium grandifolium Mett. Pheg. u. Asp. 124 n. 294. 1858 (non Pr. 1849) — Nephrodium Wightii Clarke 1880.
polyphyllum Klf. Flora **1823**[1]. 362 = Dryopteris opposita.
polyphyllum Mett. Fil. Lechl. **1**. 20. 1856 = Polystichum aculeatum var. 24.
polypodiforme Mak. Bot. Mag. Tokyo —. 1892 = Dryopteris.
polystichoides Mett.; Salom. Nom. 67. 1883 = Lastrea.
Pontederae Willd. sp. **5**. 273. 1810 = Athyrium filix femina.
porphyrophlebium Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 117 = Dryopteris.
Powellii Christ, Engl. Jahrb. **23**. 354. 1896 = Dryopteris.
praetervisum Kuhn, Linn. **36**. 111. 1869 = Dryopteris.
Prenticei Christ, Engl. Jahrb. **23**. 352. 1896 = Dryopteris.
Prescottianum Wall. List n. 363. 1828 = Polystichum.
Prescottianum var. Bakeriana Clarke, Tr. Linn. Soc. II. Bot. **1**. 510 t. 66. 1880 = Polystichum Bakerianum.
Preslianum Mett. Pheg. u. Asp. 36 n. 81. 1858 = Cyclopeltis.
Preslianum Sturm, En. pl. cr. vasc. chil. —. 1858 = Dryopteris concinna.
prionophyllum Wall. List n. 355. 1828 = Dryopteris extensa et immersa.
procerum Spr. Syst. **4**. 102. 1827 = Nephrodium.
procerum Boj.; Kuhn, Fil. Afr. 139. 1868 = Dryopteris mauritiana.
procurrens Mett. Ann. Lugd. Bat. **1**. 231. 1864 = Dryopteris.
producens Fée, 10 mém. 38. 1865 = Dryopteris diplazioides.
productum Klf. Enum. 237. 1824 = Dryopteris.
proliferum R. Br. Prod. Fl. N. Holl. 147. 1810 = Polystichum aculeatum var. 29.
proliferum Hk. et Grev. Ic. Fil. t. 96. 1829 = Fadyenia Fadyenii.
proliferum Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
prolificum Maxim.; Franch. et Sav. En. pl. Jap. **2**. 239, 632. 1879 = Dryopteris erythrosora f. monstr.
prolixum Willd. sp. **5**. 251. 1810 = Dryopteris.
prolongum Fée, Gen. 292. 1850-52 (nomen) — Ins. Philipp.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
prominulum Christ, Bull. Boiss. **4**. 659. 1896 = Dryopteris.

propinquoides Mett.; Salom. Nom. 66. 1883 = Hypopeltis.
propinquum Sw. Adnot. 67. 1829 = Dryopteris gongylodes*.
propinquum Fée, Gen. 292. 1850-52 = Dryopteris Preslii.
protensum Afz.; Sw. Schrad. Journ. **1800**^{2}. 36. 1801 = Dryopteris.
(S) **psammiosorum** C. Chr. Ind. 89. 1905 — Trinidad.
Nephrodium Sherringiae Jenm. **1887**. 99 (non 1879).
pseudo-Filix-mas Fée, 8 mém. 103. 1857 = Dryopteris filix mas.
pseudo-lonchitis Dumort —? = Polystichum lobatum.
pseudomontanum Hier. Engl. Jahrb. **22**. 373. 1896 = Dryopteris.
pseudovarium Christ, Bull. Soc. Fr. **52** Mém. I. 42. 1905 = Dryopteris.
(S) **psilopodum** C. Chr. Ind. 89. 1905 — Borneo. — Nephrodium subdigitatum Bak. Journ. Linn. Soc. **24**. 259. 1887 (non Aspidium Bl. 1828).
Ptarmica Kze.; Mett. Pheg. u. Asp. 80. n. 191. 1858 = Dryopteris.
pterifolium Mett.; Kuhn, Linn. **36**. 110. 1869 = Dryopteris.
pteroides Sw. Schrad. Journ. **1800**^{2}. 33. 1801 = Dryopteris.
pteroides var. a Mett. Ann. Lugd. Bat. **1**. 231. 1864 = Dryopteris extensa.
pteropodum Diels, Nat. Pfl. **1**^{4}. 183. 1899 = Polypodium Scortechinii.
pteropus Kze. Bot. Zeit. **1846**. 462 = A. decurrens.
(Ar) **puberulum** Desv. Prod. 245. 1827 (nomen) — Ins. Comorae; Madagascar; Ins. Mascarenae. — Sagenia Moore 1857; Polypodium triphyllum Desv. Berl. Mag. **5**. 315. 1811 (non Jacq. 1788); Phegopteris Kuhn 1868. Aspidium vel Nephrodium subtriphyllum auctt. quoad pl. afr.
puberulum Gaud. Freyc. Voy. Bot. 342. 1827 = A. martinicense.
puberulum Fée, 10 mém. 40. 1865 = Dryopteris Feei.
puberum Wall. List n. 338. 1828 = { Microlepia speluncae. / Dryopteris calcarata var. falciloba. }
pubescens Sw. Schrad. Journ. **1800**^{2}. 39. 1801 = Dryopteris.
pubescens Lowe, Ferns **6** t. 25. 1857 = Dryopteris decomposita.
pubescens Fée, Cr. vasc. Br. **2**. 73. 1872-73 = Dryopteris oligocarpa.
pubirachis Christ, Engl. Jahrb. **23**. 354. 1896 = Dryopteris.
pulchellum Bl. Enum. 167. 1828 = Polystichum aristatum.
pulchellum Kuhn, Fil. Afr. 140. 1868 = Polypodium.
pulcherrimum Col. Tasm. Journ. **2**. —. 1845 = Polystichum vestitum.
pulchrum Bory; Willd. sp. **5**. 253. 1810 = Dryopteris strigosa.
pulverulentum Desv. Prod. 251. 1827 = Polypodium.
pumilum Mart. et Gal. Mém. Ac. Brux. **15**. 64 t. 17 f. 1. 1842 = Phanerophlebia.
pumilum Lowe, Ferns **6** t. 15. 1857 = Dryopteris filix mas.
puncticaule Bl. Enum. 159. 1828 = Athyrium macrocarpum.
punctilobulum Sw. Syn. 60. 1806 } = Dennstaedtia punctilobula.
punctilobum Willd. sp. **5**. 279. 1810 }
punctulatum Sw. Syn. 46. 1806 = Nephrolepis biserrata*.
pungens Klf. Enum. 242. 1824 = Polystichum.
pungens Wall. List n. 368. 1828 = Polystichum ilicifolium.
(S) **Purdiaei** Jenm. Gard. Chr. III. **22**. 282. 1897 — Trinidad.
purpurascens Bl. Enum. 169. 1828 = Dryopteris sparsa*.
pusillum Mett. Ann. sc. nat. V. **2**. 245. 1864 = Dryopteris.
pycnolepis Kze.; Kl. Linn. **20**. 365. 1847 = Polystichum aculeatum var. 27.
quadrangulare Fée, Cr. vasc. Br. **1**. 145 t. 50 f. 2. 1869 = Dryopteris pseudotetragona.
quinquangulare Kze. Linn. **23**. 230. 302. 1850 = ? Dryopteris decomposita.

(S) **quinquefidum** (Bak.) Diels, Nat. Pfl. 1[4]. 186. 1899 — Tonkin.
Nephrodium Bak. JoB. **1890**. 265.
(T) **quitense** C. Chr. Ind. 90. 1905 — Ecuador.
Aspidium contractum Sod. Cr. vasc. quit. 220. 1893 (non Link. 1841).
Raddianum Mett. Pheg. u. Asp. 91. n. 221. 1858 = Dryopteris deflexa.
radicans Sieb. Syn. Fil. exsicc. n. 104 = Polystichum aculeatum var. 29.
radicans Fée, Gen. 294. 1850-52 — Caracas.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
ramosum Pal. Beauv. Fl. d'Oware **2**. 54 t. 91. post 1811 = Arthropteris obliterata.
ramosum Fée, Cr. vasc. Br. **1**. 141 t. 47 f. 3. 1869 = Dryopteris amplissima.
rampans Christ, Bull. Boiss. **6**. 968. 1898 = Dryopteris.
recedens Lowe, Ferns **7** t. 1. 1859 = Dryopteris.
recurvum Bree, Phytol. **1**. 773. 1843 = Dryopteris aemula.
reductum Bak. JoB. **1888**. 105 = Polystichum Hancockii.
refractum A. Br. Ind. sem. ht. Berol. 1856 = Dryopteris.
regium Sw. Schrad. Journ. **1800**[2]. 41. 1801 = Cystopteris fragilis*.
Reinwardtianum Kze. Bot. Zeit. **1848**. 261 = Dryopteris calcarata.
remotum A. Br. Verjüngung 330. 1850 = Dryopteris filix mas × spinulosa.
remotum Christ in Schwacke, Pl. Nov. Min. **2**. 29. 1900 = Polystichum.
(T) **repandum** Willd. sp. **5**. 216. 1810. Mett. n. 298. HB. 258 — Ins. Philipp. Perak.
Bathmium Fée 1850-52; Sagenia Moore 1858; Dryopteris O. Ktze. 1891. —? Tectaria crenata Cav. 1802; Aspidium tectaria Desv. 1827.
repandum Bl. Enum 144. 1828 = Dryopteris urophylla.
repandum Sod. Rec. 42. 1883 = A. subrepandum.
reptans Mett. Pheg. u. Asp. 99. n. 237. 1858 { var. 1 = Dryopteris cordata. var. 2 = Dryopteris sagittata. var. 3 et 4 = Dryopteris reptans.
resiniferum Klf. Enum. 237. 1824 = Dryopteris gongylodes.
resinofoetidum Christ, Bull. Soc. bot. Belg. **35**. 213. 1896 = Dryopteris.
retroflexum Willd. sp. **5**. 238. 1810 = ? Dryopteris refracta.
retusum Mett. Novara Exp. Bot. **1**. 218. 1870 = Dryopteris.
rhaeticum Willd. sp. **5**. 280. 1810 = { Cystopteris fragilis. Athyrium filix femina.
(Ar) **rheosorum** (Bak.) C. Chr. Ind. 90. 1905 — Costa Rica.
Polypodium Bak. JoB. **1884**. 363 (non 1891).
rheticum Sw. Schrad. Journ. **1800**[2]. 41. 1801 = { Athyrium alpestre et A. filix femina.
rhizophyllum Sw. Schrad. Journ. **1800**[2]. 31. 1801 = Polystichum.
rhomboideum Wall. List n. 364. 1828 = Polystichum amabile.
Richardi Hk. sp. **4**. 23. t. 222. 1862 = Polystichum.
Riedleanum Gaud. Freyc. Voy. Bot. 326 (syn.). 1827 = Polystichum.
rigidissimum C. Chr. Bot. Tidsskr. **25**. 79. 1902 = Cyclodium.
rigidulum Mett.; Kuhn, Linn. **36**. 109. 1869 = Dryopteris scalpturoides.
rigidum Sw. Schrad. Journ. **1800**[2]. 37. 1801 = Dryopteris.
rigidum var. remotum A. Br.; Döll, Rhein. Fl. 16. 1843 = Dryopteris filix mas × spinulosa.
riparium Bory, Willd. sp. **5**. 250. 1810 = Dryopteris tomentosa.
riparium Moritz, Bot. Zeit. **1854**. 855 = Dryopteris oligocarpa.
(S) **rivale** Mett.: Kuhn, Linn. **36**. 120. 1869 — Panama-Ecuador.

Rivoirei Fée, 11 mém. 76 t. 21 f. 3. 1866 = Dryopteris oligocarpa.
rivularioides Fée, Cr. vasc. Br. **1**. 148 t. 50 f. 1. 1869 = Dryopteris opposita.
rivulorum Link, Hort. Berol. **2**. 119. 1833 = Dryopteris opposita.
Robertianum Luerss. in Aschers. et Graebner Syn. **1**. 22. 1896 = Dryopteris.
robustum Kze.; Mett. Fil. Lechl. **1**. 20. 1856 = Polystichum aculeatum var. 26.
Rochaleanum Christ, Bull. Boiss. II. **2**. 558. 1902 = Polystichum aculeatum [var. 21.
roseum Fourn. Mex. pl. **1**. 97. 1872 = Dryopteris patula.
rostratum HBK. Nov. Gen. et Sp. **1**. 12. 1815 = Alsophila blechnoides.
rostratum Wall. List n. 383. 1828 = A. polymorphum.
rotundatum Willd. sp. **5**. 247. 1810 = Dryopteris.
rotundatum Vahl; Hornem. Animadvers. Havn. 8. 1819 (nomen) — Sierra Leone.
rubiginosum Mann, Proc. Amer. Acad. **7**. 217. 1868 = Dryopteris.
rufescens Bl. Enum. 168. 1828 = Dryopteris.
rufescens Kze. Flora **1839**[1]. Beibl. 34 = Nephrolepis hirsutula.
rufidulum Sw. Syn. 58. 1806 = Woodsia ilvensis.
rufo-barbatum Wall. List n. 369. 1828 = Polystichum lobatum var. 4
rufostramineum Christ, Bull. Soc. Fr. **52** Mém. I. 36. 1905 = Dryopteris.
rufum Mett. Novara Exp. Bot. **1**. 218. 1870 = Dryopteris.
rutaceum Willd. sp. **5**. 266. 1810 = Asplenium.
Sabaei Franch. et Sav. Enum. pl. Jap. **2**. 259, 632. 1879 = Dryopteris.
sagenioides Mett. Pheg. u. Asp. 113. n. 269. 1858 = Dryopteris.
sagittaefolium Bl. Enum. 153. 1828 = Dryopteris.
salaccense Bl. Enum. 140. 1828 = Oleandra neriiformis.
Sancti-Gabrielis Fée, Cr. vasc. Br. **2**. 75. 1872-73 = Dryopteris.
sanctoides Fée, Gen. 292. 1850-52 = Dryopteris opposita.
sanctum Bl. Enum. 143. 1828 = A. angulatum.
sanctum Mett. Pheg. u. Asp. 76. n. 181. 1858 = Dryopteris.
savaiiense Christ, Engl. Jahrb. **23**. 350. 1896 = Dryopteris.
(Ar) **saxicola** Bl. Enum. 160. 1828 — Java.
Polypodium petrophyum Bl. Fl. Jav. Fil. 163 t. 71. 1829. (HB. 319); Dictyopteris Moore 1861; Phegopteris Mett. 1864.
scabriusculum Mett.; Salom. Nom. 69. 1883 = Dryopteris patens.
scabriusculum Dav. Bot. Gaz. **21**. 255. 1896 = Dryopteris Davenportii.
scabrosum Kze. Linn. **24**. 286. 1851 = Dryopteris.
Scallanii Christ, Bull. Soc. Bot. It. **1901**. 296 = Dryopteris.
scandens Raddi, Pl. Bras. **1**. 34 t. 49. 1825 = Polybotrya.
scandicinum Willd. sp. **5**. 285. 1810 = Athyrium.
scariosum Roxb. Calc. Journ. **4**. 466. 1844 = Polystichum speciosum.
Schaffneri Kuhn, Linn. **36**. 114. 1869 = Dryopteris.
Schimperianum Hochst.; A.Br. Flora **1841**. 708 = Dryopteris.
Schizolobium Kuhn, Fil. Afr. 141. 1868 = Polystichum aculeatum var. 2.
Schkuhrii Bl. Enum. 147. 1828 = Nephrolepis hirsutula.
Schomburgkii Kl. Linn. **20**. 369. 1847 — Guiana.
Lastrea Moore 1857. — (Dryopteris sp. dub.).
Schottianum Kze. (Bak. Syn. 262. 1867) = Dryopteris chrysoloba var.
Schweinitzii Beck (Gray, Man. ed. I. 633. 1848) = Polystichum acrostichoides.
sclerophyllum Kze.; Spr. Syst. **4**. 99. 1827; Kze. Linn. **9**. 92. 1834 = Dryopteris asplenioides.
scolopendrioides Mett. Pheg. u. Asp. 97 n. 235. 1858, var. 1. incisa = Dryopteris incisa; var. 2. subpinnata = Dryopteris scolopendrioides; var. 3. pinnata = Dryopteris asplenioides.

securidiforme Mett.; Kuhn, Fil. Afr. 141. 1868 = Dryopteris.
Seemanni Hk. sp. **4**. 34. t. 230. 1862 = Polystichum.
(S) **semibipinnatum** Wall. List n. 388. 1828; Hk. sp. **4**. 59 t. 231. 1862 — Penang. Borneo. — Polypodium Wall. 1829; Sagenia Moore 1858; Nephrodium Bak. 1867. HB. 297; Dryopteris O. Ktze. 1891.
semicordatum Sw. Schrad. Journ. **1800**[2]. 31. 1801 = Cyclopeltis.
semihastatum Kze. Linn. **9**. 91. 1834 = Dryopteris.
senanense Franch. et Sav. Enum. pl. Jap. **2**. 241, 636. 1879 = Dryopteris.
sericeum Fée, Cr. vasc. Br. **1**. 144 t. 42 f. 1. 1869 = Dryopteris opposita.
serra Sw. Schrad. Journ. **1800**[2]. 33. 1801 = Dryopteris.
serra Schkuhr, Kr. Gew. **1**. 35 t. 33 b. 1806 = Dryopteris gongylodes.
serratum Sw. Syn. 47, 246. 1806 = Dryopteris gongylodes.
serratum Kuhn, Fil. Afr. 142. 1868 = Amblia.
serrulatum Mett. Pheg. u. Asp. 106. n. 252. 1858 = Dryopteris.
sesquipedale Willd. sp. **5**. 230. 1810 = Nephrolepis rivularis.
setigerum Sw. Adnot. 66. 1829 — Hispaniola.
Lastrea Moore 1858 — (Dryopteris sp. aff. D. hirta).
setigerum Kuhn, Verh. zool. bot. Ges. **19**. 578. 1869 = Dryopteris.
setosum Sw. Schrad. Journ. **1800**[2]. 39. 1801 = Polystichum varium.
setosum Wall. List n. 371. 1828 = Polystichum lobatum var. 3.
setosum Kl. Linn. **20**. 371. 1847 = Dryopteris pseudotetragona.
Sewellii Kuhn, v. Deck. Reisen **3**[3]. Bot. 65. 1879 = Dryopteris.
Shepherdi Kze. Linn. **23**. 230. 1850 = Dryopteris decomposita.
shikokianum Mak. Bot. Mag. Tokyo —. 1892 = Dryopteris.
siambonense Hier. Engl. Jahrb. **22**. 372. 1896 = Dryopteris.
sibiricum Turcz.; Besser, Flora **1834**[1]. Beibl. 28 = Athyrium crenatum.
Sieberi Steud. Nomencl. **2**. 64. 1824 = Dryopteris limbata.
Sieberianum Mett.; Salom. Nom. 70. 1883 = Dryopteris crinita.
Sieboldii van Houtte; Mett. Fil. Lips. 87 t. 20 f. 1-4. 1856 = Dryopteris.
(S) **siifolium** (Willd.) Mett. Ann. Lugd. Bat. **1**. 237. 1864. NPfl. 185 — Malesia. Ins. Philipp. N. Guinea. — Polypodium Willd. sp. **5**. 196. 1810; Nephrodium Bak. 1867. HB. 299; Dryopteris O. Ktze. 1891. Drynaria menisciicarpa J. Sm. 1841; Aspidium Mett. n. 287. 1858 (non Bl. 1828); Dryomenis Moore 1857; Polypodium Hk. 1863 (pt.). Dryomenis Phymathodes Fée 1850-52 — Aspidium Teijsmannianum Hk. 1862.
siifolium Bl. Enum. 143. 1828 = A. pachyphyllum.
sikkimense Bak. Syn. 256. 1867 = Dryopteris.
(S) **Simonsii** (Bak.) Bedd. Ferns br. Ind. Suppl. 15 t. 367. 1876 — India bor. Nephrodium Bak. Syn. 504. 1874; Dryopteris O. Ktze. 1891.
simplex Hance, Journ. Linn. Soc. **13**. 140. 1873 = Dryopteris.
simplicifolium Hk. Ic. pl. t. 919. 1854 = Dryopteris canescens*.
simplicissimum Christ, Bull. Boiss. II. **4**. 959. 1904 = Dryopteris.
simulatum Dav. Bot. Gaz. **19**. 495. 1894 = Dryopteris.
(T) **singaporianum** Wall.; Hk. et Grev. Ic. Fil. t. 26. 1827. Mett. n. 299. Chr. 227. NPfl. 183 — Perak. Malacca. Penang. Malesia. China centr. Podopeltis Fée 1850-52; Nephrodium Bak. 1867. HB. 296; Dryopteris O. Ktze. 1891 — ? Aspidium lobulatum Bl. 1828.
Sintenisii Kuhn (Bak. Ann. Bot. **5**. 327. 1891); Christ, Engl. Jahrb. **24**. 119. 1897 = Dryopteris.

(S) **sinuatum** Labill. Sert. Austr. Cal. 1 t. 1. 1824 — N. Caledonia.
Sagenia Moore 1858; Bathmium Fourn. 1873; B. Billardierii Fée 1850-52.
sinuatum Gaud. Freyc. Voy. Bot. 343. 1827 = A. cicutarium var. Gaudichaudii.
sinuatum Moore, Ind. 104. 1858 = A. plantagineum.
sinuatum Mett. Ann. sc. nat. IV. **15**. 76. 1861 = A. novae caledoniae.
Skinneri Hk. Ic. pl. t. 924. 1854 = Dryopteris.
(S) **Sodiroi** (Bak.) Hier. Engl. Jahrb. **34**. 451. 1904 — Ecuador. Columbia.
Nephrodium Bak. JoB. **1877**. 164.
solutum Wall. List n. 350. 1828 = Dryopteris parasitica.
sophoroides Sw. Schrad. Journ. **1800**^{2}. 33. 1801 = Dryopteris.
sorbifolium Willd. sp. **5**. 223. 1810 = ? Polybotrya polyphylla.
(Ar) **sparsiflorum** (Hk.) Diels, Nat. Pfl. **1**^{4}. 185. 1899 — Africa occ. trop.
Polypodium Hk. sp. **5**. 92. 1863. HB. 319.
sparsum Spr. Syst. **4**. 106. 1827 = Dryopteris.
speciosum Don, Prod. Fl. Nepal. 5. 1825 = Polystichum.
speciosum Mett.; Ettingsh. Farnkr. 180. t. 106 f. 12, t. 109 f. 10, t. 110 f. 8. 1865; Kuhn, Fil. Afr. 142. 1868 = Dryopteris Vogelii.
spectabile Bl. Enum. 158. 1828 = Dryopteris syrmatica.
spectabile Wall. List n. 372. 1828 = Dryopteris Hendersoni.
Spekei Kuhn, Fil. Afr. 209. 1868 = Dryopteris.
speluncae Willd. sp. **5**. 269. 1810 = Microlepia.
spinescens (Moore et Houlst.) Lowe, Ferns **7** t. 6. 1859 — India.
Lastrea Moore et Houlst. (1852?) — (Dryopteris sp. dub.).
spinulosum Sw. Schrad. Journ. **1800**^{2}. 38. 1801 = Dryopteris.
spinulosum × cristatum Lasch, A.Br. et Milde, Jahresber. schles. Ges. **1855**. 94 = Dryopteris cristata × spinulosa.
splendens Willd. sp. **5**. 220. 1810 = Nephrolepis biserrata.
splendens Ham.; Wall. List n. 2241. 1829 —?
splendens Christ, Farnkr. 259. 1897 = Dryopteris.
sporadosorum Kze. Bot. Zeit. **1848**. 556 = Polystichum carvifolium.
Sprengelii Klf. Flora **1823**^{1}. 365 = Dryopteris Balbisii.
squamatum Willd. sp. **5**. 256. 1810 = Didymochlaena truncatula.
squamatum Kze. Linn. **23**. 230. 1850 = Polystichum.
squamigerum Fée, 8 mém. 104. 1857 = Dryopteris thelypteris*.
squamigerum Mann, Proc. Am. Acad. **7**. 217. 1868 = Dryopteris.
squamisetum Kuhn, Fil. Deck. 24. 1867 = Dryopteris.
squamosum Poir. Enc. Suppl. **4**. 521. 1816 = Didymochlaena truncatula.
squarrosum Don, Prod. Fl. Nepal. 4. 1825 = Polystichum lobatum var. 4.
squarrosum Wall. List n. 356. 1828 = Athyrium macrocarpum.
stenolepis Christ, Bull. Boiss. **7**. 22. 1899 = Dryopteris.
stenophyllum Diels, Nat. Pfl. **1**^{4}. 185. 1899 = A. Hosei.
stenopteris Kze. Farnkr. **2**. 48 t. 120. 1849 = Dryopteris incisa.
stenopteron Christ, Journ. de Bot. **19**. 61. 1905 = A. Balansae.
stimulans Kze.; Mett. Pheg. u. Asp. 43. n. 97. 1858 = Polystichum ilicifolium.
stipellatum Bl. Enum. 152. 1828 = Dryopteris.
stipitatum Mett. Pheg. u. Asp. 60. n. 143. 1858 = Dryopteris.
stipulaceum Mett. Pheg. u. Asp. 82. n. 198. 1858 = Dryopteris heteroptera.
stipulare Willd. sp. **5**. 239. 1810 = Dryopteris patens*.
stramineum Klf.; Spr. Syst. **4**. 105. 1827 = Polystichum aculeatum var. 1.
striatum Schum. Vid. Selsk. Afh. **4**. 230 (seors. 456). 1827 = Dryopteris.

strigilosum Dav. Bot. Gaz. **21**. 256. 1896 = Dryopteris.
strigosum Willd. sp. **5**. 249. 1810 = Dryopteris.
strigosum Mett. Pheg. u. Asp. 89 n. 217. 1858 = Dryopteris crinita.
strigosum Fée, 11 mém. 78 t. 22 f. 2. 1866 = Dryopteris opposita.
strigosum Christ, Bull. Soc. bot. Belg. **35**. 210. 1896 = Dryopteris Alfarii.
(S) **subconfluens** Bedd. Ferns br. Ind. Suppl. 14 t. 364. 1876. NPfl. 186 — India bor. — Nephrodium Clarke 1880.
subcrenulatum Kuhn, v. Deck. Reis. 3^3. Bot. 65. 1879 = Dryopteris.
(Ar) **subdecurrens** (Luerss.) C. Chr. Ind. 94. 1905 — Singapore. Phegopteris Luerss. Bot. Centralbl. **11**. 30. 1882.
subdecussatum Christ, Bull. Boiss. II. **4**. 960. 1904 = Dryopteris.
subdiaphanum Wall. List n. 343. 1828 = Dryopteris crenata.
subdigitatum Bl. Enum. 171. 1828 = Monachosorum.
(S) **subebeneum** Christ, Bull. Boiss. II. **5**. 256. 1905 — Costa Rica.
subelongatum Bl. Enum. 166. 1828 = Polystichum aristatum.
subexaltatum Christ, Bull. Boiss. II. **4**. 616. 1904 = Dryopteris.
subglandulosum Mett. (Kuhn, Fil. Afr. 138. 1868) = Dryopteris Helliana.
subinerme Kze. Linn. **24**. 290. 1851 = Polystichum aculeatum var. 36.
subintegerrimum Hk. et Arn. Bot. Beech. Voy. 52. 1832 = Polystichum aculeatum var. 10.
subjunctum Christ, Engl. Jahrb. **23**. 354. 1896 = Dryopteris.
sublanosum Wall. List n. 365. 1828 = Nephrolepis cordifolia.
sublobatum Bl. Enum. 166. 1828 = Polystichum aculeatum.
submite Christ, Nu. Giorn. Bot. It. n. s. **4**. 93. 1897 = Polystichum.
subobliquatum Fée, Cr. vasc. Br. **2**. 75. 1872-73 = Dryopteris.
subpectinatum Bl. Enum. 145. 1828 = Arthropteris obliterata.
subpectinatum Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 248 = Dryopteris [otaria.
(S) **subpedatum** (Harr.) Diels, Nat. Pfl. 1^4. 186. 1899 — Formosa. Nephrodium Harr. Journ. Linn. Soc. **16**. 30. 1877.
subpubescens Bl. Enum. 149. 1828 = Dryopteris parasitica *amboinensis.
subquinquefidum Pal. Beauv. Fl. d'Oware **1**. 34 t. 19. 1804 = Dryopteris protensa.
(T) **subrepandum** Bak. Ann. Bot. **5**. 315. 1891; Sod. Cr. vasc. quit. 639. 1893. NPfl. 185 — Ecuador. — A. repandum Sod. Rec. 42. 1883 (non Willd. 1810).
subsericeum Mett. Ann. sc. nat. IV. **15**. 74. 1861 = Dryopteris.
subspinulosum Christ, Bull. Boiss. II. **2**. 829. 1902 = Dryopteris.
(S) **subtriphyllum** (Hk. et Arn.) Hk. sp. **4**. 52. 1862. Chr. 227. NPfl. 185 — China. Birma. Ceylon. Malesia-Polynesia. — Polypodium Hk. et Arn. Bot. Beech. Voy. 256 t. 50. 1836-40; Sagenia Bedd. 1863; Nephrodium Bak. 1867. HB. 296; Bathmium Fourn. 1873; Dryopteris O. Ktze. 1891.
subtriphyllum auctt. quoad pl. mascaren. = A. puberulum.
subtripinnatum Miq. Ann. Lugd. Bat. **3**. 179. 1867 = Dryopteris.
sulcatum Klf.; Bory, Bél. Voy. Bot. **2**. 58. 1833 = Dryopteris crinita.
sulcinervium Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
(Ar) **sumatranum** C. Chr. Ind. 94. 1905 — Sumatra. — Polypodium Hancockii Bak. Ann. Bot. **8**. 127. 1894 (non 1885 nec Aspidium Bak. 1891).
syrmaticum Willd. sp. **5**. 237. 1810 = Dryopteris.
tacticopterum Kze. Linn. **24**. 290. 1851 = Polystichum aculeatum var. 35.
tanacetifolium Rupr. Distr. Cr. vasc. Ross. 37. 1845 = Dryopteris spinulosa *dilatata.

Tasmaniae Mett. Fil. Lips. 88. 1856 = Polystichum vestitum.
Ar) **Tatei** (Bak.) Diels, Nat. Pfl. **1**⁴. 184. 1899 — Nicaragua.
Polypodium Bak. Syn. 506. 1874; Ic. pl. t. 1670; Phegopteris Salom. 1883.
Tavoyanum Wall. List n. 1032. 1828 = Nephrolepis cordifolia.
taygetense Bory, Exp. sci. de Morée **3**². 288. 1832 = Cystopteris fragilis *regia.
tectaria Desv. Prod. 245. 1827 = ? A. repandum.
tectum Wall. List n. 394. 1828 = Dryopteris didymosora.
Telfairianum Wall. List n. 385. 1828 = Cyathea borbonica.
Teijsmannianum Hk. sp. **4**. 41 t. 236. 1862 = A. siifolium.
tenericaule Thwait. Enum. pl. Zeyl. 393. 1864 = Dryopteris setigera.
(Ar) **tenerifrons** (Hk.) Diels, Nat. Pfl. **1**⁴. 186. 1899 — Birma. — Polypodium Hk. sp. **4**. 104. 1862 (excl. pl. afric.). HB. 318; Dictyopteris Bedd. 1866.
tenerrimum Fée, Cr. vasc. Br. **1**. 134 t. 43 f. 1. 1869 = Dryopteris Balbisii.
tenerum Spr. Syst. **4**. 109. 1827 = Dryopteris.
tenerum Fée, Cr. vasc. Br. **1**. 147. 1869 = Polystichum denticulatum.
tenue Retz. Obs. **6**. 39. 1791 = Cheilanthes tenuifolia.
tenue Sw. Syn. 58. 1806 = Cystopteris fragilis.
tenuiculum Fée, Gen. 292. 1850-52 — Cuba?
Lastrea Moore 1858 — (Dryopteris sp. dub.).
tenuifolium Mett.; Kuhn, Linn. **36**. 122. 1869 = A. cicutarium var.
tenuifolium Luerss. Fil. Graeff. 190 (syn.). 1871 = Dryopteris tenuifrons.
tenuisectum Bl. Enum. 170. 1828 = Cystopteris.
terminans Wall. List n. 386. 1828 = Dryopteris pteroides.
(S) **ternatum** (Bak.) Diels, Nat. Pfl. **1**⁴. 184. 1899 — Borneo.
Nephrodium Bak. Syn. 296. 1867; Dryopteris O. Ktze. 1891.
tetragonum Sw. Schrad. Journ. **1800**². 33. 1801 = Dryopteris.
tetragonum Mett. Pheg. u. Asp. 95. n. 232. 1858 = Dryopteris pseudotetragona.
tetragonum Sturm, En. pl. vasc. cr. chil. —. 1858 = Polystichum.
Thelypteris Sw. Schrad. Journ. **1800**². 40. 1801 = Dryopteris.
thelypteroides Sw. Syn. 57. 1806 = Dryopteris noveboracensis.
thelypteroides Mett. Pheg. u. Asp. 81. n. 194. 1858 = Dryopteris tomentosa.
thibeticum Franch. Nouv. Arch. Mus. II. **10**. 118. 1887 = Dryopteris.
Thomsoni Hk. 2 Cent. t. 25. 1860 = Polystichum.
Thonningii Schum. Vid. Selsk. Afh. **4**. 229 (seors. 455). 1827 — Guinea.
(Pl) **Thwaitesii** Bedd. Ferns br. Ind. Suppl. 15. 1876. NPfl. 188 — Ceylon.
Sagenia Bedd. Ferns S. Ind. t. 244. 1863; Pleocnemia Bedd. 1883; Nephrodium simulans Bak. 1874. HB. ed. II. 300; Dryopteris O. Ktze. 1891.
tijucense Fée, Cr. vasc. Br. **2**. 72 t. 102. 1872-73 = Dryopteris itatiaiensis.
tomentosum Willd. sp. **5**. 234. 1810 = Nephrolepis hirsutula.
tomentosum Kuhn, v. Deck. Reisen **3**³. Bot. 38. 1879 = Dryopteris.
Tonduzii Christ, Prim. Fl. costaric. **3**. 34. 1901 = Dryopteris Leprieurii.
Torresianum Gaud. Freyc. Voy. Bot. 334 (syn.). 1827 = Polystichum.
totta Engl. Hochgbfl. trop. Afr. 99. 1892 = Dryopteris africana.
transitorium Christ, Bull. Boiss. **7**. 822. 1899 = Dryopteris tokyoensis.
trapezoides Sw. Schrad. Journ. **1800**². 31. 1801 = ? Polystichum viviparum.
trapezoides Schkuhr, Kr. Gew. **1**. 30 t. 29 b. 1806 = Nephrolepis pectinata.
? trapezoides Kze. Linn. **9**. 93. 1834 = Polystichum aculeatum var. 24.
Trejoi Christ, Bull. Boiss. II. **5**. 255. 1905 = Polystichum.
Trianae Mett. Ann. sc. nat. V. **2**. 243. 1864 = Dryopteris.
triangulare Desv. Prod. 247. 1827. } = Polystichum triangulum.
triangulum Sw. Schrad. Journ. **1800**². 31. 1801 }

tricholepis Christ, Bull. Boiss. II. **5**. 259. 1905 = Dryopteris.
trichophlebium Kuhn, v. Deck. Reisen **3**[3]. Bot. 65. 1879 = Dryopteris.
trichophorum Fée, 11 mém. 81 t. 23 f. 2. 1866 = Dryopteris.
trichotomum Fée, Gen. 295. 1850-52 — Cochinchina.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
(T) **tricuspe** Bedd. Handb. Suppl. 44. 1892 — Perak.
tridens Hk. sp. **4**. 15 t. 215. 1862 = Polystichum.
trifidum Sw. Schrad. Journ. **1800**[2]. 41. 1801 = Cystopteris fragilis.
(T) **trifoliatum** (L.) Sw. Schrad. Journ. **1800**[2]. 30. 1801. Mett. n. 297. HB. 258. Chr. 227. NPfl. 185 — Amer. trop. Florida. Texas.
Polypodium L. sp. **2**. 1087. 1753; Tectaria Cav. 1802; Bathmium Link 1841; Dryopteris O. Ktze. 1891 — Aspidium heracleifolium Willd. 1810; Bathmium Fée 1850-52. — vide et Aspidium fimbriatum Willd. 1810.
trifoliatum auctt. quoad pl. mascar. = A. pica.
trifoliatum Dcne. Nouv. Annal. Mus. **3**. 345. 1834 = ? A. Haenkei.
trifoliatum Hk. JoB. **9**. 340. 1857 = A. subtriphyllum.
(T) **trilobum** Sod. Rec. 41. 1883; Cr. vasc. quit. 638. 1893 — Ecuador.
Trimeni Diels, Nat. Pfl. **1**[4]. 188. 1899 = A. giganteum.
(S) **tripartitum** (Bak.) Diels, Nat. Pfl. **1**[4]. 185. 1899. — Fiji.
Nephrodium Bak. JoB. **1879**. 296; Ic. pl. t. 1666.
Tripteris Eat. in Perry, Narr. Exp. to China **2**. 330. 1856 } = Polystichum
tripteron Kze. Bot. Zeit. **1848**. 569 } tripteron.
triste Bl. Enum. 169. 1828 = Alsophila.
triste Kze.; Fée, Gen. 292. 1850-52 (nomen) — Africa austr.
Lastrea moesta Moore 1858. — (Dryopteris sp. dub.).
triste Mett. Pheg. u. Asp. 94. n. 229. 1858 = Dryopteris.
truncatulum Sw. Schrad. Journ. **1800**[2]. 36. 1801 = Didymochlaena.
truncatum Gaud. Freyc. Voy. Bot. (332) t. 10. 1827 = Dryopteris.
truncatum Trevis. Hedwigia **13**. 149. 1874 = Cyclopeltis Presliana.
Tsus-Simense Hk. sp. **4**. 16 t. 220. 1862 = Polystichum lobatum var. 2.
tuberosum Bory; Willd. sp. **5**. 234. 1810 = Nephrolepis cordifolia.
Tussaci Fée, Gen. 297. 1850-52 — San Domingo.
Lastrea Moore 1858 — (Dryopteris sp. dub.).
tylodes »Kze.« auctt. plur. vide xylodes.
uliginosum Kze. Linn. **20**. 6. 1847 = Dryopteris setigera.
uliginosum Nym. Consp. Fl. Eur. 866. 1884 = Dryopteris cristata × spinulosa.
umbilicatum Desv. Berl. Mag. **5**. 320. 1811 = Dryopteris filix mas.
umbrosum Sw. Schrad. Journ. **1800**[2]. 42. 1801 = Athyrium.
undulatum Afz.; Sw. Schrad. Journ. **1800**[2]. 32. 1801 = Nephrolepis cordifolia.
undulatum Thwait. En. pl. Zeyl. 444. 1864 = Dryopteris sparsa*.
unitum Sw. Schrad. Journ. **1800**[2]. 32. 1801 = Dryopteris gongylodes *propinqua.
unitum Sieb.; Mett. Pheg. u. Asp. 107 n. 257. 1858 = Dryopteris.
unitum Mett. Ann. Lugd. Bat. **1**. 230. 1864 = Dryopteris gongylodes.
urophyllum Christ, Farnkr. 248. 1897 = Dryopteris.
Van Heurckii Fourn. Mex. pl. **1**. 97. 1872 — Mexico. — (Dryopteris sp.).
(Ar) **varians** (Moore) C. Chr. Ind. 96. 1905 — Africa occ. trop.
Dictyopteris varians Moore, Gard. Chr. **1863**. 1108. Polypodium Cameroonianum Hk. sp. **5**. 104. 1864. HB. 318; Dictyopteris J. Sm. 1866; Aspidium Mett.; Kuhn 1868. NPfl. 186.

varians Mett.; Kuhn, Linn. **36**. 114. 1869 = Dryopteris imbricata.

(S) **variolosum** Wall. List n. 379. 1828; Hk. sp. **4**. 51. 1862. Chr. 228. NPfl. 185 — India bor. Birma-Malesia.
Sagenia (variolata) Moore 1858; Nephrodium Bak. 1867. HB. 298; Dryopteris O. Ktze. 1891 — Aspidium alienum Mett.; Kuhn 1869.

varium Sw. Schrad. Journ. **1800**[2]. 35. 1801 = Polystichum.

varium Willd. sp. **5**. 236. 1810 — India. China.
Sagenia (J. Sm. 1841?) Moore 1858.

(S) **vastum** Bl. Enum. 142. 1828. Chr. 230. NPfl. 186 — India bor. Birma-Malesia.
Sagenia Moore 1857; Nephrodium Bak. 1867. HB. 296; Dryopteris O. Ktze. 1891. Aspidium alatum Wall. 1828; Hk. et Grev. Ic. Fil. t. 184. 1830. Mett. n. 290; Bathmium Fée 1850-52; Sagenia Bedd. 1867 — Polypodium heterosorum Bak. 1874. HB. 506; Dictyopteris Bedd. 1876.

(S) **Veitchianum** C. Chr. Ind. 97. 1905 — »Polynesia«.
Nephrodium amblyotis Bak. Ann. Bot. **5**. 331. 1891 (non Aspidium Kze. 1848).

velatum Kze.; Mett. Pheg. u. Asp. 79. n. 190. 1858 = Dryopteris.

velleum Willd. sp. **5**. 255. 1810 = Dryopteris.

velutinum A. Rich. Fl. N. Zél. 70. 1832 = Dryopteris.

venulosum Bl. Enum. 151. 1828 = Dryopteris gongylodes.

venulosum Wall. List n. 352. 1828 = Dryopteris arida.

venustum Hew. Mag. Nat. Hist. II. **2**. 464. 1838 = Dryopteris.

venustum Hk.fil. Fl. antarct. **1**. 106. 1844 = Polystichum vestitum.

verrucosum Kze. Bot. Zeit. **1846**. 473 = Dryopteris immersa.

vestitum Sw. Schrad. Journ. **1800**[2]. 37. 1801 = Polystichum.

vestitum Fée, Cr. vasc. Br. **1**. 134. 1869 = Dryopteris deflexa.

Vieillardii Mett. Ann. sc. nat. IV. **15**. 75. 1861 = Dryopteris.

vile Kze. Bot. Zeit. **1846**. 474 = Dryopteris.

villosum Sw. Schrad. Journ. **1800**[2]. 39. 1801 = Dryopteris.

villosum Moore, Ind. 107. 1858 — Hort. — Bathmium Fée, Gen. 289. 1850-52.

violascens Link, Hort. Berol. **2**. 115. 1833 = Dryopteris parasitica var.

(Pl) **viridans** Mett.; Kuhn, Verh. zool. bot. Ges. **19**. 578. 1869 — N. Caledonia. N. Hebridae. Fiji. — Polypodium Milnei Hk. sp. **4**. 254. 1862 (non Nephrodium Hk. 1861); Pleocnemia Fourn. 1873. Nephrodium membranifolium Bak. HB. 282 (pt.) —? Hemitelia laciniata Spr. 1827.

viridescens Miquel, Cat. Mus. Lugd. Bat. 127. 1870 = Dryopteris.

viridulum Desv. Berl. Mag. **5**. 321. 1811 = Cystopteris fragilis.

viscidulum Mett. Pheg. u. Asp. 38. n. 85. 1858 = Polystichum.

viscosum Bedd. JoB. **1888**. 4 = Dryopteris.

(S) **vitis** (Racib.) C. Chr. Ind. 97. 1905 — Borneo.
Nephrodium Racib. Bull. int. Ac. Cracovie **1902**. 60.

viviparum Mett. Pheg. u. Asp. 44. n. 99. 1858 = Polystichum.

Vogelii Hk. Ic. pl. t. 921. 1854 = Dryopteris.

Volkensii Hier. in Engl. Pflw. Ostafr. C. 86. 1895 = Polystichum.

vulcanicum Bl. Enum. 163. 1828 = Polystichum aculeatum var. 31.

waikarense Col. Tasm. Journ. **2**. —. 1845 = Polystichum vestitum.

Wakefieldii Hier. in. Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.

Wallichianum Spr. Syst. **4**. 104. 1827 = Dryopteris filix mas.

Wallichianum Wall. List n. 373. 1828 = Oleandra Wallichii.

Wallichianum Bory, Bél. Voy. Bot. **2**. 56. t. 5. 1833 = Oleandra neriiformis.

Wallichianum Kze. Linn. **24**. 291. 1851 = Polystichum lobatum var. 3.

Wallichii Hk. Exot. Flora **1**. t. 5. 1823 = Oleandra.
Warburgii Kuhn et Christ in Warb. Monsunia **1**. 81. 1900 = Dryopteris.
Wardii Kuhn, v. Deck. Reis. **3**[2]. Bot. 69. 1879 = Dryopteris.
Wattii Bedd. JoB. **1888**. 234 = Polystichum.
Wawraeanum Szysz. in Beck, It. Princ. Cob. **2**. —. 1888 —?.
Webbianum A.Br. Flora **1841**. 711 = Polystichum.
Weigleanum Kze. Linn. **24**. 284. 1851 = Dryopteris sparsa.
(S) **Weinlandii** Christ, Bull. Boiss. II. **1**. 453. 1901 — N. Guinea.
Welwitschii Kuhn, Fil. Afr. 209. 1868 = Dryopteris.
Wigmanii Racib. Bull. int. Ac. Cracovie **1902**. 61 = Dryopteris.
Wilsoni Hier. in. Engl. Pflw. Ostafr. C. 84. 1895 = Dryopteris.
Wrightii Mett.; Eat. Mem. Amer. Acad. n. s. **8**. 210. 1860 = Dryopteris.
xiphophyllum Bak. JoB. **1888**. 227 = Polystichum.
xylodes Kze. Linn. **24**. 283. 1851 = Dryopteris ochthodes var.
yaeyamense Mak. Bot. Mag. Tokyo **11**. 18. 1897 = Dryopteris.
yunnanense Christ, Bull. Boiss. **6**. 965. 1898 = Dryopteris stenolepis.
zambesiacum Hier. in Engl. Pflw. Ostafr. C. 85. 1895 = Dryopteris.
zerophyllum Col. Tr. N. Zeal. Inst. **29**. 418. 1897 = Polystichum.
zeylanicum Fée, Gen. 297. 1850-52 — Ceylon.
Lastrea Moore 1858 — (an Dryopteris sagenioides?).
(Ar) **Zippelianum** C. Chr. Ind. 98. 1905 — N. Guinea. — Phegopteris ferruginea Mett. Ann. Lugd. Bat. **1**. 224 t. 6 f. 2. 1864; Polypodium Bak. 1867. HB. 318; Aspidium Diels 1899. NPfl. 186 (non Sw. 1806 nec Fée 1850-52).
Zollingerianum Kze. Bot. Zeit. **1846**. 462 = A. brachiatum.
Zollingerianum Bedd. Ferns br. Ind. t. 251. 1868 = A. variolosum.

Aspidopodium Diels, Nat. Pfl. **1**[4]. 321. 1899 *(Polypodium §)* = Polypodium.
ASPIDOTIS Nuttall msc. (Hk. sp. **2**. 70. 1852) = **Hypolepis** (californica).
ASPLENIDICTYUM J. Smith in Hook. Ic. plant. t. 937. 1854 (syn.); Hist. Fil. 333. 1875 = **Asplenium.**
Finlaysonianum J. Sm. Hist. Fil. 334. 1875.
Purdieanum J. Sm. Hist. Fil. 334. 1875.

ASPLENIOPSIS Mettenius; Kuhn, Chaetopterides 324. 1882 — (Polypodiaceae Gen. **70**).
decipiens Mett.; Kuhn, Chaetopt. 325. 1882. NPfl. 272 — Polynesia occ.
Gymnogramme Mett. Ann. sc. nat. IV. **15**. 60. 1861. HB. 381. Chr. 69.

Asplenites J. Smith, Hist. Fil. 317. 1875 *(Asplenium §)* = Asplenium.
Asplenitis J. Smith, Hist. Fil. 158. 1875 *(Gymnogramma §)* = Syngramma.

ASPLENIUM Linné (Gen. plant. 783. 1737) Spec. plant. **2**. 1078. 1753 — (Polypodiaceae Gen. **48**). — (rectius Asplenum).
abbreviatum Mak. Bot. Mag. Tokyo **13**. 12. 1899 — Japonia.
abrotanoides Pr. Rel. Haenk. **1**. 47 t. 8 f. 2. 1825 = A. fragrans *foeniculaceum.
abscissum Willd. sp. **5**. 321. 1810. Mett. n. 42. HB. 203. Chr. 195. NPfl. 237 — Amer. trop. Florida. Arizona. — A. laetum Schkuhr 1809 (non Sw. 1806); A. Schkuhrianum Pr. 1836 — A. firmum Kze. 1845 — A. polymorphum Mart. et Gal. 1842; Tarachia Pr. 1849 — Asplenium platychlamys Fée 1857.
abscissum Raddi, Opusc. sci. Bol. **3**. 291. 1819 = A. semicordatum.
abscissum Bl. Enum. 182. 1828 = A. unilaterale.
abyssinicum Fée, Gen. 199. 1850-52. Mett. n. 97 — Africa trop.
A. cicutarium β HB. 220.

achalense Hier. Engl. Jahrb. **22**. 378. 1896. NPfl. 239 — Argentina.
achilleaefolium Liebm. Vid. Selsk. Skr. V. **1**. 249 (seors. 97). 1849 = Athyrium.
achilleifolium (Lam.) C. Chr. Ind. 99. 1905 — Japonia. China. India bor. Ceylon. Fiji. N. Caledonia. Africa austr. cum ins. adjac. — Adiantum Lam. Enc. **1**. 43. 1783. Caenopteris furcata Berg. 1786 (vix Adiantum L.fil. 1781); Darea Willd. 1810. Darea stans Bory 1833; Asplenium Kze. 1839 — Caenopteris rutaefolia Berg. 1786; Darea Willd. 1810; Asplenium Kze. 1836. Mett. n. 60. HB. 222. Chr. 208. NPfl. 242 — Adiantum borbonicum Jacq. 1789; Asplenium Hk. 1860. HB. 222 — A. prolongatum Hk. 1860 (Asia) — ? Lonchitis bipinnata Forsk. Fl. Aeg. Arab. 184. 1775; Darea disticha Klf. 1824; Caenopteris Spr. 1827; Asplenium Mett.; Salom. 1883. (Arabia felix). (»bipinnatum« verisimiliter nom. optim.).
acrostichoides Sw. Schrad. Journ. **1800**^{2}. 54. 1801 = Athyrium.
acuminatum Wall. List n. 205. 1828 (nomen) = ? Diplazium Christii.
acuminatum Hk. et Arn. Bot. Beech. Voy. 106. 1832. HB. 218. NPfl. 241 — Ins. Hawaii. — A. polyphyllum Pr. 1836; Tarachia Pr. 1849.
acuminatum Willd.; Pr. Tent. 107. 1836 = A. Willdenowii Pr.
acutiusculum Bl. Enum. 178. 1828 — Java. Celebes. Ins. Philipp. Samoa. A. anisodontum Pr. 1849. Mett. n. 141. HB. 205.
acutum Bory; Willd. sp. **5**. 347. 1810 = A. adiantum nigrum *onopteris.
adiantoides (L.) C. Chr. Ind. 99. 1905 — Asia, Polynesia, Australia trop. N. Zealand. Afr. orient. cum ins. adjac. — Trichomanes L. sp. **2**. 1098. 1753. Asplenium falcatum Lam. Enc. **2**. 306. 1786. Mett. n. 156. HB. 208. Chr. 198. NPfl. 239; Tarachia Pr. 1849 — Asplenium polyodon Forst. 1786. Mett. n. 155; Tarachia Pr. 1849.
adiantoides Lam. Enc. **2**. 309. 1786 = A. praemorsum.
adiantoides Raddi, Opusc. sci. Bol. **3**. 291. 1819; Pl. Bras. **1**. 40 t. 51 f. 2 = A. scandicinum.
adiantoides Raoul, Ann. sc. nat. III. **2**. 115. 1844 = A. Hookerianum.
Adiantum lanceolatum Hoffm. Deutschl. Fl. **2**. 12. 1795 = A. adiantum nigrum.
adiantum nigrum L. sp. **2**. 1081. 1753. Mett. n. 128. HB. 214. Chr. 202. NPfl. 240 — Europa. Asia temp.-Himalaya. Ins. atlant. Montes Africae. Africa austr. Réunion. Ins. Hawaii. ? Puerto Rico.
Tarachia Pr. 1849; Asplenium nigrum Lam. 1778; Phyllitis lancifolia Moench 1802 — Asplenium tabulare Schrad. 1818 — A. argutum Klf. 1824; Tarachia Pr. 1849 — Asplenium obtusum Kit.; Willd. 1810; Tarachia Pr. 1849. — *A. cuneifolium Viv. 1806; A. Forsteri Sadl. 1820; A. Serpentini Tausch 1839. Chr. 203.
— *A. Onopteris L. 1753; A. acutum Bory; Willd. 1810; Tarachia Pr. 1849 — A. davallioides Tausch 1839 — A. silesiacum Milde 1855.
adiantum nigrum × trichomanes Milde, Verh. zool. bot. Ges. **14**. 165. t. 4. 1864 — Tirol. — A. dolosum Milde l. c.; A. trichomanes × adiantum nigrum Aschers. 1896.
adscensionis Bernh. Schr. Akad. Erfurt **1802**. 18 f. 8 = Pteris dentata var.
adulterinum Milde, Sporenpfl. 40. 1865. HB. 483. Chr. 192. NPfl. 235 — Europa centr.
adulterinum × viride Aschers. Syn. Mitteleur. Fl. **1**. 59. 1896 — Sachsen.
aequabile Bak. Syn. 485. 1874 — Ins. Diego Garcia (Mascarenae).
aequibasale Bak. Journ. Linn. Soc. **22**. 225. 1886 = Diplazium.

affine Sw. Schrad. Journ. **1800**[2]. 56. 1801. HB. 215. Chr. 205. NPfl. 240 — Ins. afric. orient. Ceylon. Malesia-Polynesia.
— A. spathulinum J. Sm. 1841 — A. pygmaeum Hk. 1860. HB. 196 (non L. 1759) — A. Gilpinae Bak. 1877 — A. herpetopteris Bak. 1877.
affine Mett.; Salom. Nom. 77. 1883 = Diplazium.
affine auctt. quoad pl. Hawaii. = A. insiticium.
africanum Desv. Berl. Mag. **5**. 322. 1811 — Africa trop. — A. sinuatum Pal. Beauv. post 1811. Mett. n. 8. HB. 192. Chr. 189. NPfl. 233 (non Salisb. 1796).
Aitoni Moore, Ind. XLIX. 1857 = Athyrium umbrosum.
alatum H.B. Willd. sp. **5**. 319. 1810. Mett. n. 76. HB. 200. Chr. 194. NPfl. 236 — Amer. trop.
alienum Mett. Fil. Lechl. **2**. 18. 1859; Aspl. 169 n. 181. 1859 = Diplazium sandwichianum.
alismaefolium Hk. sp. **3**. 267. 1860 = Diplazium Cumingii.
allantodioides Mett. Aspl. 191 n. 228. 1859 = Diplazium asplenioides.
alloeopteron Kze.; Kl. Linn. **20**. 353. 1847 = A. radicans var.
alpestre Bl. Enum. 176. 1828 = Diplazium crenato-serratum vel D. pallidum.
alpestre Mett. Aspl. 198 n. 245 t. 6 f. 1-6. 1859 = Athyrium.
alternans Wall. List n. 221. 1828 = Ceterach Dalhousiae.
alternifolium Jacq. Misc. austr. **2**. 51 t. 5 f. 2. 1781 = A. germanicum.
alternifolium Mett. Fil. Lips. 75 t. 12 f. 1-2. 1856 = Diplazium lineolatum.
alternifolium Mett. Ann. Lugd. Bat. **2**. 237. 1866 = Diplazium bantamense var.
alternifrons Dillwyn, Ref. Hort. Mal. 64. 1839 (Rheede **12** t. 16) — India.
altissimum Jenm. JoB. **1879**. 259 = Diplazium.
amabile Liebm. Vid. Selsk. Skr. V. **1**. 251 (seors. 99). 1849 = A. uniseriale.
ambiguum Sw. Schrad. Journ. **1800**[2]. 54. 1801 = Diplazium esculentum.
ambiguum Spr. Fl. Halensis, Mant. prima 54. 1807 = A. Sprengelii.
ambiguum Schkuhr, Kr. Gew. **1**. t. 75 (non 75 b) 1809 = Diplazium japonicum.
ambiguum Raddi, Opusc. sci. Bol. **3**. 291. 1819; Pl. Bras. **1**. 38 t. 54. 1825 = Diplazium Shepherdii.
amblyodon Fée, Gen. 191. 1850-52 (nomen) — Fiji.
amboinense Willd. sp. **5**. 303. 1810. Mett. n. 15. Chr. 189. NPfl. 235 — Malesia. Polynesia. — A. taeniosum Kze. 1848; Thamnopteris Pr. 1849; Neottopteris Fée 1850-52 — Asplenium feejeense Brack. 1854. Mett. n. 16. HB. 192; Micropodium J. Sm. 1875.
amboinense Brack. Expl. Exp. **16**. 147 t. 19 f. 2. 1854 = A. vittaeforme.
ammifolium Mett.; Kuhn, Fil. Afr. 96. 1868 = Athyrium.
amoenum Pr. Tent. 107. 1836; Mett. Aspl. 131 n. 101. 1859 = A. unilaterale.
amplissimum Bak. Ann. Bot. **5**. 310. 1891 = Diplazium.
amplum Willd.; Kuhn, Fil. Afr. 96. 1868 (nomen) — Mauritius.
amplum Fourn. Mex. pl. **1**. 109. 1872 = Diplazium striatum.
anauniense Loss, Voce Cattolica **1872** n. 90 = A. lepidum.
anceps L. v. Buch. Beschr. Canar. Ins. 189. 1825; Hk. et Grev. Ic. Fil. t. 195. 1830 = A. trichomanes*.
Andersoni Clarke, Journ. Linn. Soc. **15**. 154. 1876 = Athyrium fimbriatum.
angolense Bak. Syn. 485. 1874 — Angola.
angustatum Desv. Prod. 275. 1827 = A. mucronatum.
angustatum Bl. Enum. 187. 1828 = ? A. laserpitiifolium (vel A. cuneatum).
angustatum Pr. Tent. 108. 1836 = A. lineatum.
angustatum Kze. Flora **1839**[1] Beibl. 40; Mett. Aspl. 167. n. 177. 1859 = A. Martianum.
angustifolium Jacq. Coll. **1**. 121. 1786 = ? Polypodium Billardieri.
angustifolium Michx. Fl. bor. Am. **2**. 265. 1803 = Athyrium.

angustifolium Guss. in Tenore, Fl. Nap. Prod. App. V. 30. 1826; Pl. Rar. t. 65. 1826 = A. fissum.
angustifrons Mett.; Salom. Nom. 78. 1883 = Diplazium arboreum.
augustum Sw. Vet. Ak. Hdl. **1817**. 66 t. 4 f. 1. Mett. n. 10. HB. 191. Chr. 189 — Guiana. Brasilia. — A. loriforme Hk. 1854.
anisodontum Pr. Epim. 73. 1849 = A. acutiusculum.
anisophyllum Kze. Linn. **10**. 511. 1836. Mett. n. 36. HB. 204. Chr. 196. NPfl. 237 — Africa trop. *America trop.
— *A. sanguinolentum Kze.; Mett. n. 35. 1859 — A. nigrescens Hk. f. 1847 (non Bl. 1828); A. nubilum Moore 1859 (Galapagos).
anomalum Desv. Prod. 276. 1827 = ? Diplazium ambiguum.
anomodum Col. Tr. N. Zeal. Inst. **15**. 309. 1883 = A. obtusatum.
anthraxacolepis Mett.; Salom. Nom. 78. 1883 = Diplazium Shepherdii.
anthriscifolium Jacq. Coll. **2**. 103 t. 2 f. 3-4. 1788 = A. pumilum.
apicidens Moore, Gard. Chr. n. s. **15**. 267. 1881 = A. Vieillardii var.
apicidentatum Moritz; Hombr. et Jacq. Voy. Pôle Sud, Crypt. t. 1. f. A. 1852 = A. obtusatum *obliquum.
appendiculatum Pr. Tent. 106. 1836 = A. bulbiferum var.
appendiculatum var. angustilobum Müll. Linn. **25**. 718. 1852 = A. flaccidum.
approximatum Bl. Enum. 179. 1828 = A. pellucidum.
? aquaticum Kl. et Karst. Linn. **20**. 354. 1847 = A. obtusifolium.
aquilinum Bernh. Schrad. Journ. **1799**^{1}. 310 = Pteridium.
arborescens Mett. Fil. Lips. 78 t. 13 f. 19-20. 1856 = Diplazium.
arboreum Willd. sp. **5**. 320. 1810 = Diplazium.
arcuatum Liebm. Vid. Selsk. Skr. V. **1**. 241 (seors. 89). 1849. Mett. n. 112. HB. 197 — Mexico. Ecuador.
arcuatum Hk. sp. **3**. 219. 1860 = Athyrium.
argutans Fée, Gen. 194. 1850-52; 7 mém. 53 t. 24 f. 2 = Diplazium tomentosum.
argutum Klf. Enum. 176. 1824 = A. adiantum nigrum.
arifolium Burm. Fl. Ind. 231. 1768 = Hemionitis.
Arnottii Bak. Syn. 240. 1867 = Diplazium.
ascensionis Watson, Proc. Amer. Acad. **26**. 163. 1891 — Ascension.
asperum Bergsm. Cat. ht. Rheno-Traj. 1857 = Diplazium polypodioides.
aspidiiforme Fée, Gen. 199. 1850-52 = Diplazium Franconis var.
aspidioides Schlecht. Adumbr. 24 t. 13. 1825 = Athyrium scandicinum.
aspidioides Spr. Syst. **4**. 90. 1827 = Athyrium.
aspidioides Goldm. Nova Acta **19**. Suppl. I. 461. 1843 = Diplazium Meyenianum.
aspidioides Sod. Cr. vasc. quit. 182. 1893 = Athyrium reductum.
assimile Endl. Prod. Fl. Norf. 10. 1833 = Athyrium.
athyrioides Fée, 8 mém. 83. 1857 = Athyrium achilleifolium.
Athyrium Spr. Anleit. **3**. 113. 1804 = Athyrium filix femina.
Atkinsoni Clarke, Tr. Linn. Soc. II. Bot. **1**. 487. 1880 = Athyrium.
Atkinsoni var. Andersoni Clarke l. c. t. 57. 1880 = Athyrium fimbriatum.
atropurpureum Bernh. Schrad. neu. Journ. **1**^{2}. 10. 1806 = Pellaea.
atrovirens (Hk.) Bak. Syn. 484. 1874 — Ins. Salomonis.
A. persicifolium var. atrovirens Hk. sp. **3**. 109. 1860.
attenuatum R. Br. Prod. Fl. N. Holl. 150. 1810. Mett. n. 132. HB. 194 — Australia orient. — Tarachia Pr. 1849.
attenuatum Klf. Enum. 174. 1824 = A. Martianum.
attenuatum Pr. Rel. Haenk. **1**. 42. 1825 = Diplazium proliferum.

aureum Cav. (Anal. Cienc. **4**. 100. 1801) Descr. 256. 1802 = Ceterach.
?aureum Bl. Enum. 185. 1828 = Osmunda banksiifolia.
auricularium Desv. Prod. 273. 1827 = A. lunulatum.
auriculatum (Thbg.) Kuhn, Fil. Afr. 97. 1868 — Africa austr.
Caenopteris Thbg. Nova Acta Petrop. **9**. 159 t. E. f. 2. 1795; Darea Juss.; Willd. 1810. Asplenium Thunbergii Kze. 1836. Mett. n. 67. HB. 223. NPfl. 243.
auriculatum Sw. Vet. Ak. Hdl. **1817**. 68 = A. semicordatum.
auriculatum Mett. Aspl. 164. n. 172. 1859 = Diplazium arboreum.
auritum Sw. Schrad. Journ. **1800**^{2}. 52. 1801 = A. sulcatum var.
auritum Wall. List n. 222. 1828 = A. bipartitum.
australasicum Hk. Fil. exot. t. 88. 1859 = A. nidus var.
australe Sw. Schrad. Journ. **1800**^{2}. 50. 1801 = Actiniopteris.
australe Brack. Expl. Exp. **16**. 173. 1854 = Athyrium umbrosum*.
avicula Cord. Bull. Soc. sci. Réunion **1890-91** —; Fl. Réun. 63. 1895 — Réunion.
axillare J. Sm. Bot. Mag. **72**. Comp. 30. 1846 = Athyrium filix femina*.
Bakeri C. Chr. Ind. 102. 1905 — N. Guinea. — Davallia (Loxoscaphe) lanceolata Bak. Kew Bull. **1899**. 119 (non Asplenium Huds. 1762).
Baldwinii Hill. Fl. Haw. 618. 1888 = Athyrium.
Balfourii Bak. Ann. Bot. **5**. 303. 1891 — Réunion.
bantamense Bak. Syn. 231. 1867 = Diplazium.
Baptistii Moore, Gard. Chr. n. s. **15**. 235. 1881 = A. Vieillardii var.
barbadense Jenm. Gard. Chr. III. **15**. 134. 1894 — Barbados.
barisanicum Bak. Ann. Bot. **8**. 126. 1894 = Diplazium.
Barteri Hk. 2 Cent. t. 75. 1861 = A. lunulatum var.
basilare Moore, Ind. XLIX. 1857 = Athyrium silvaticum.
Baumgartneri Dörfler, Oest. Bot. Zeit. **45**. 169 t. 9. 1895 = A. septentrionale × trichomanes.
Beccarianum Cesati, Rend. Acad. Napoli **16**. 26, 29. 1877 — N. Guinea.
Beddomei Mett.; Kuhn, Linn. **36**. 93. 1869 = A. crinicaule.
Belangeri Bory, Bél. Voy. Bot. **2**. 47. 1833 = A. longissimum.
Belangeri Kze. Bot. Zeit. **1848**. 176 = A. tenerum*.
bellum Clarke, Tr. Linn. Soc. II. Bot. **1**. 496 t. 63 f. 2. 1880 = Athyrium umbrosum var.
Bernieri Cord. Bull. Soc. sci. Réunion **1890-91** —. 1891; Fl. Réun. 66. 1895 — [Réunion.
bianthemum Gilbert, Bull. Torr. Cl. **24**. 259. 1897 — Jamaica.
bicrenatum Liebm. Vid. Selsk. Skr. V. **1**. 245 (seors. 93). 1849 = A. semicor- [datum.
bidentatum Willd. sp. **5**. 318. 1810 = A. sulcatum.
bifidum Pr. Tent. 109 t. 3 f. 19. 1836 = A. lineatum var.
bifissum Fée, Gen. 199. 1850-52. (HB. 220) — Cuba.
bifolium L. sp. **2**. 1078. 1753 (Plum. t. 133) — Hispaniola.
bifrons Sod. Cr. vasc. quit. 184. 1893 = Diplazium.
bifurcatum Opiz, Kratos **4** —. 1819
bifurcum Opiz, Böh. phan. u. crypt. Gew. —. 1823 } = A. septentrionale.
Billetii Christ, Bull. sci. France et Belg. **28**. 267 t. 12 f. 4-5. 1898 — Tonkin.
Billotii F. Schultz, Flora **1845**. 738 = A. lanceolatum. [China austr.
bipartitum Bory; Willd. sp. **5**. 328. 1810. Mett. n. 65. HB. 212 — Ins. Comorae et Mascarenae. — Diplazium Pr. 1849.
bipartitum Spr. Syst. **4**. 85. 1827 = Diplazium semihastatum.
bipinnatifidum Bak. Syn. ed. I. 221. 1867 — Fiji.
A. furcatum Brack. Expl. Exp. **16**. 170. 1854 (non Thbg. 1800); A. bipinna-

tum Brack. l. c. 344. 1854. HB. ed. II. 221 (non Roxb. 1844). A. Seemanni Carr. 1873 (non Bak. 1867).
bipinnatum Roxb. Calc. Journ. **4**. 499. 1844 = Diplazium esculentum.
bipinnatum Brack. Expl. Exp. **16**. 344. 1854 = A. *bipinnatifidum*.
bipinnatum Hill. Fl. Haw. 595. 1888 = A. parallelum.
biseriale Bak. Tr. Linn. Soc. II. Bot. **4**. 252. 1894 = Diplazium.
biserratum Pr. Del. Prag. **1**. 177. 1822 = Diplazium.
bissectum Sw. Schrad. Journ. **1800**². 55. 1801; Syn. 82. 1806 = A. dissectum.
Blakistoni Bak. Syn. ed. I. 216. 1867 = A. Sarelii.
Blanchetii Mett.; Kuhn, Linn. **36**. 104. 1869 = Diplazium.
blandulum Fée, 7 mém. 51. 1857 = A. monanthes.
blechnoides Lag.; Sw. Syn. 76. 1806 = Blechnum.
blepharodes Eat. Zoë **1**. 197. 1890 — Lower California.
blepharophorum Bertol. Nov. Comment. Bonon. **4**. 443. 1840 — Guatemala.
Blumeanum Moore, Ind. 116. 1859 = A. vulcanicum.
Blumei Bergsm. Cat. ht. Rheno-Traj. 1857; Mett. Aspl. 189. n. 224 = Diplazium polypodioides.
Bodinieri Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 242 c. fig. — China.
bogotense Karst. Fl. Col. **2**. 77 t. 139. 1864-69 = Diplazium hians var.
Boivini Mett.; Kuhn, Fil. Afr. 98. 1868 — Réunion.
Boltoni Hk. Syn. 204. 1867 in obs. (sp. **3**. t. 166. 1860) = A. anisophyllum.
borbonicum Hk. sp. **3**. 207. 1860 = A. *achilleifolium*.
borneense Hk. sp. **3**. 135 t. 186. 1860. HB. 203 — Borneo. Perak.
Boryanum Moore, Ind. 117. 1859 — Ins. Mascarenae.
Darea asplenioides Bory, Bél. Voy. Bot. **2**. 53. 1833.
Bourgaei Boiss.; Milde, Bot. Zeit. **1866**. 384. HB. 216 — Syria.
brachycarpum (Mett.) Kuhn, Linn. **36**. 104. 1869 — N. Hebridae.
Davallia Mett. msc. (HB. 470); Loxoscaphe Kuhn 1879.
brachyopterum Moore et Houlst. Gard. Mag. Bot. **3**. 260. 1852 = A. bra- [chypteron.
brachyotus Kze. Linn. **10**. 512. 1836 = A. laetum.
brachyphyllum Gaspar. Rend. Acad. Napoli **4**. 108. 1845 = A. lepidum.
brachypodum Bak. Syn. 490. 1874 = Diplazium.
brachypteron Kze. Linn. **23**. 232. 1850 = A. Dregeanum.
brachysorum R. Br.; Moore, Ind. 117. 1859 (nomen) — Congo.
Brackenridgei Bak. Syn. 234. 1867 = Diplazium bulbiferum.
Bradleyi Eat. Bull. Torr. Cl. **4**. 11. 1873. HB. 487. NPfl. 240 — U. S. A. atlant.
brasiliense Sw. Vet. Ak. Handl. **1817**. 65 t. 3 f. 1 = Phyllitis.
brasiliense Raddi, Pl. Bras. **1**. 36 t. 51 f. 1. 1825 = A. lunulatum.
brasiliense Desv. Prod. 273. 1827 = A. semicordatum.
brasiliense Moore et Houlst. Gard. Mag. Bot. **3**. 258. 1852 = A. serratum.
brevipes Bak. Journ. Linn. Soc. **16**. 201. 1877 = Diplazium.
brevisorum Wall. List n. 220. 1828 = Athyrium.
brevisorum Mett. Aspl. 192 n. 229 b. 1859 = Athyrium silvaticum.
Breynii Retz. Obs. **1**. 32. 1779 = A. germanicum.
Brownii J. Sm. JoB. **4**. 174. 1841; Hk. Ic. pl. t. 978. 1854 = Athyrium um- [brosum *australe.
brunneo-viride Jenm. JoB. **1886**. 269 = Diplazium.
Brunonianum Mett. Fil. Lips. 71. 1856 = Diplaziopsis javanica.
bulbiferum Forst. Prod. 80. 1786. Mett. n. 52. HB. 218. Chr. 206. NPfl. 241 — N. Zealand. Australia. India bor. Penang. — Caenopteris Desv. 1827 — C. appendiculata Lab. 1806; Darea Willd. 1810; Asplenium Pr. 1836.

A. laxum R. Br. 1810 — ? A. Fabianum Hombr. et Jacq. 1852 (conf. A. lineatum) — A. bullatum Wall. 1828. Mett. n. 51 — A. canterburiense Armstr. 1882.

bulbiferum Bernh. Schr. Ak. Erfurt **1802**. 18 f. 7 = ? Diplazium sp. dubia.

bulbosum Lour. Fl. Cochinch. **2**. 833. 1790 = ? Nephrolepis cordifolia.

bullatum Wall. List n. 215. 1828 = A. bulbiferum.

caenopteroides Desv. Prod. 276. 1827 = A. praemorsum.

caespitosum Bl. Enum. 175. 1828 = A. vulcanicum f. indivisa.

caespitosum Wall. List n. 217. 1828 = A. laciniatum.

Callipteris Fée, Gen. 193. 1850-52 = A. vittaeforme.

Callipteris Bak. Syn. 231. 1867 = Diplazium.

calophyllum J. Sm. JoB. **3**. 408. 1841 = Diplazium pallidum.

camarinum Bak. Syn. ed. II. 230. 1874 = Diplazium.

Campbellii Jenm. Gard. Chr. n. s. **24**. 7. 1885 = Diplazium.

camptocarpon Fée, Cr. vasc. Br. **1**. 63 t. 16 f. 1. 1869 = A. oligophyllum.

camptocarpon Fourn. Mex. pl. **1**. 109. 1872 = Diplazium.

camptorachis Kze. Linn. **24**. 262. 1851 = A. lunulatum var.

camptosorum Mett.; Kuhn, Linn. **36**. 101. 1869 = A. serra var.

canaliculatum Bl. Enum. 180. 1828 = A. macrophyllum.

canaliculatum Christ in Warb. Monsunia **1**. 73. 1900 = A. impressum.

canariense Willd. sp. **5**. 339. 1810 = A. praemorsum.

canterburiense Armstr. Tr. N. Zeal. Inst. **14**. 361. 1882 = A. bulbiferum var.

capillipes Mak. Bot. Mag. Tokyo **17**. 77. 1903 — Japonia.

caput-serpentis Henriq. (Bak. Ann. Bot. **5**. 302. 1891) = A. trichomanes.

caracasanum Willd. sp. **5**. 338. 1810 = Diplazium.

cardiophyllum (Hance) Bak. Ann. Bot. **5**. 311. 1891 — Hainan. Formosa.
Micropodium Hance JoB. **1883**. 268.

caripense (Kl.) Fourn. Mex. pl. **1**. 106. 1872 — Mexico. Columbia.
A. cuneatum var. caripense Kl. Linn. **20**. 356. 1847.

Carruthersii Bak. Syn. 483. 1874 — Fiji.
A. salignum Carr. 1873 (non Bl. 1828).

caryaefolium Bak. Ann. Bot. **5**. 310. 1891 = Diplazium.

caryotoides Pr. Tent. 107. 1836 = ? A. dimidiatum.

castaneaefolium Hk. sp. **3**. 237. 1860 = Diplazium.

castaneo-viride Bak. Ann. Bot. **5**. 304. 1891 — China.

castaneum Schlecht. et Cham. Linn. **5**. 611. 1830. Mett. n. 114. (HB. 197) — Mexico-Venezuela.

cataractarum Bl. Enum. 177. 1828 = Diplazium pallidum vel D. crenato-serratum.

caucense Karst. Fl. Col. **1**. 155 t. 77. 1861 = Diplazium Roemerianum.

caudatum Forst. Prod. 80. 1786. Mett. n. 139. HB. 209. Chr. 199. NPfl. 239 Trop. — Tarachia Pr. 1849.

caudatum Cav. Descr. 265. 1802 = A. tenerum.

caudatum Bedd. Ferns S. Ind. 143. 1863 = A. adiantoides.

celtidifolium Mett. Fil. Lips. 75 t. 12 f. 3-4. 1856 = Diplazium.

centrifugale Bak. Syn. 485. 1874 — Peru.

centrifugale Bak. Journ. Linn. Soc. **25**. 360. 1889 = A. Listeri.

centripetale Bak. Syn. 490. 1874 = Diplazium celtidifolium.

centrochinense Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 111. c. fig. — China.

ceramicum Miq. Ann. Lugd. Bat. **4**. 168. 1868-69 = Diplazium.

ceratolepis Christ, Bull. Boiss. **4**. 658. 1896 = Diplazium.

Cesatianum Bak. Malesia **3**. 39. 1886 — N. Guinea.
Ceterach L. sp. **2**. 1080. 1753 = Ceterach.
Ceylonense Kl.; J. Sm. Ferns br. and for. 221. 1866 = Athyrium ceylanense.
Chamissonianum Pr. Tent. 107. 1836 (nomen) — Luzon. — Tarachia Pr. 1849.
chaerophylloides Fée, Cr. vasc. Br. **1**. 71 t. 16 f. 2. 1869 = A. scandicinum.
cheilosorum Kze.; Mett. Aspl. 133. n. 104. t. 5 f. 12-13. 1859 — Formosa. India bor.-Borneo. — A. heterocarpum Wall. List n. 218. 1828 (nomen); Hk. sp. **3**. 132 t. 175. 1860. HB. 210. Chr. 194. NPfl. 237.
chihuahuense Bak. Ann. Bot. **5**. 305. 1891. NPfl. 241 — Mexico.
A. dubiosum Dav. 1891.
chimborazense Spruce; Bak. Syn. 244. 1867 = Diplazium.
chinense Bak. Syn. 237. 1867 = Diplazium.
chlaenopteron Fée, Gen. 194. 1850-52; 7 mém. 47 t. 16 f. 1 — Réunion. (prob. = A. lineatum var.).
chlorophyllum Bak. JoB. **1885**. 104 = Diplazium.
chocoense Triana; Mett. Ann. sc. nat. V. **2**. 231. 1864 = Diplazium.
chondrophyllum Bert.; Colla, Mem. Ac. Torino **39**. 40 t. 68. 1836 = A. obtusatum.
Christii Hier. in Engl. Pflw. Ostafr. C. 82. 1895. NPfl. 237 — Afr. orient. trop.
cicutarium Sw. Prod. 130. 1788 = A. cristatum.
cicutarium Roxb. Calc. Journ. **4**. 500. 1844 = ? A. praemorsum.
cicutarium var. paleaceum Dav. Bot. Gaz. **19**. 390. 1894 = A. Eatoni.
cirrhatum Rich.; Willd. sp. **5**. 321. 1810 — Amer. trop.
A. mastigophyllum Fée 1857.
cladolepton Fée, 7 mém. 55 t. 22 f. 4. 1857 (HB. 220) — Columbia.
Clarkei Atkins.; Clarke, Tr. Linn. Soc. II. Bot. **1**. 489. 1880 = Athyrium nigripes [var.
Clutei Gilbert, Fern Bull. **8**. 62 c. ic. 1900 — Jamaica.
coarctatum Mett. Fil. Lips. 75. 1856 = ? Diplazium denticulosum.
coenobiale Hance, JoB. **1874**. 142 — Canton. — A. fuscipes Bak. 1879.
Colensoi Col. Tasm. Journ. **2**. 170. 1845 (Colensii). HB. 219 — N. Zealand.
columbiense Hier. Engl. Jahrb. **34**. 461. 1904 = A. sessilifolium var.
commutatum Mett.; Kuhn, Linn. **36**. 99. 1869 — Mexico-Guatemala.
comorense C. Chr. Ind. 105. 1905 — Ins. Comorae.
A. debile Mett.; Kuhn, Fil. Afr. 101. 1868 (non Fée 1865).
comosum Christ in Schum. et Laut. Fl. deut. Südsee 127. 1901 — N. Guinea.
compactum L. v. Buch, Beschr. Canar. Ins. 189. 1825 (nomen) — Madeira.
compressum Sw. Schrad. Journ. **1800**^{2}. 52. 1801. Mett. n. 39. HB. 206. NPfl. 239 — St. Helena. Ins. Trinidad (Atlant. austr.).
comptum Kze.; Moore et Houlst. Gard. Mag. Bot. **2**. 259. 1852 = A. Karstenianum.
comptum Hance, Ann. sc. nat. V. **5**. 255. 1866; Kuhn, Bot. Zeit. **1869**. 131. (HB. 486) — Canton.
conchatum Moore, Ind. XLIX. 1857 = Athyrium costale.
concinnum Wall. List n. 216. 1828 = A. tenuifolium.
concinnum Kuhn, Fil. Afr. 99. 1868 = A. theciferum*.
concisum Desv. Prod. 277. 1827 — »Amer. austr.«.
A. dareoides Desv. Berl. Mag. **5**. 322. 1811.
concolor Hk. sp. **3**. 88 t. 164 A. 1860 = A. diversifolium f. indivisa.
confertum Bak. Ann. Bot. **8**. 125. 1894 = Diplazium.
confluens Kze. Bot. Zeit. **1848**. 174 = A. pusillum.
congestum C. Chr. Ind. 105. 1905 — Ecuador.
A. debile Sod. Cr. vasc. quit. 141. 1893 (non Fée 1865 nec Mett. 1869).

congruum Christ, Engl. Jahrb. **23**. 348. 1896 = Diplazium.
Conilii Franch. et Sav. En. pl. Jap. **2**. 227. 1876, 625. 1879 = Diplazium japonicum.
consanguineum Gaud. Freyc. Voy. Bot. 315. 1827 = A. lunulatum.
consimile Remy; Fée, Gen. 191. 1850-52 (nomen); Gay, Fl. chil. **6**. 501. 1853
consimile ht. Veitch; Gartenfl. **13**. 246. 1864 — Hort. [= A. obtusatum.
contaminans Wall. List n. 2210. 1829 (nomen) — Singapore.
contiguum Klf. Enum. 172. 1824. Mett. n. 145. HB. 205. NPfl. 239 — Hawaii. Ins. Philipp. — Tarachia Pr. 1849 — Asplenium lepturus J. Sm. 1841. Mett. n. 146.
contiguum Bedd. Ferns S. Ind. t. 140. 1863 = A. caudatum.
cordatum Sw. Schrad. Journ. **1800**². 54. 1801 = Ceterach.
Cordemoyi C. Chr. Ind. 106. 1905 — Réunion.
A. inaequale Cord. 1891; Fl. Réun. 60. 1895 (non Kze. 1848).
Corderoi Sod. Rec. 39. 1883 = Diplazium.
cordifolium Spr. Syst. **4**. 88. 1827 = Llavea.
cordifolium Mett. Fil. Lips. 74 t. 12 f. 6. 1856 = Diplazium.
cordovense Bak. Ann. Bot. **8**. 125. 1894 = Diplazium.
coriaceum Desv. Prod. 275. 1827 = A. obtusifolium.
coriaceum Bory, Bél. Voy. Bot. **2**. 46. 1833 = A. Wightianum.
coriaceum Roxb. Calc. Journ. **4**. 497. 1844 = A. macrophyllum.
coriaceum Fée, Gen. 193. 1850-52 = A. pycnophyllum.
coriaceum Bak. Syn. 192. 1867 = A. subintegrum.
coriandrifolium Pr. Tent. 108. 1836 (nomen) —?
coriifolium Liebm. Vid. Selsk. Skr. V. **1**. 249 (seors. 97). 1849 = A. fragrans
costale Sw. Schrad. Journ. **1800**². 55. 1801 = Diplazium. [*foeniculaceum.
costale Moore, Ind. 43. 1857 = Athyrium.
crassidens Fée, 8 mém. 82. 1857 (crassides) = Diplazium Roemerianum var.
crassum Thouars, Fl. Trist. d'Ac. 33. 1804 = A. obtusatum *obliquum.
crenato-lobatum Bak. Ann. Bot. **5**. 309. 1891 = Diplazium.
crenato-serratum Bl. Enum. 177. 1828 = Diplazium.
crenato-serratum Bomm.; Christ, Bull. Boiss. **4**. 662. 1896 = Diplazium
crenatum Desv. Prod. 279. 1827 = A. cuneatum. [Bommeri.
crenatum Roxb. Calc. Journ. **4**. 498. 1844 = Diplazium.
crenatum Fries, Summ. Veg. **1**. 82, 253. 1845 = Athyrium.
crenatum Miq. Ann. Lugd. Bat. **3**. 174. 1867 = Dryopteris Miqueliana.
crenulatum Pr. Tent. 106. 1836 = A. serratum.
crenulatum Bak. Syn. ed. I. 224. 1867 = Athyrium crenatum.
crenulatum Bak. Syn. 236. 1867 = Diplazium striatum.
crinicaule Hance, Ann. sc. nat. V. **5**. 254. 1866. HB. 208. NPfl. 239 — China austr. India bor. et austr.
A. Hancei Bak. 1867; Kuhn, Bot. Zeit. **1869**. 130. A. Beddomei Mett. 1869.
crinitum Bak. Journ. Linn. Soc. **24**. 258. 1887 = Diplazium.
crinulosum Desv. Prod. 272. 1827 — Réunion.
cristatum Lam. Enc. **2**. 310. 1786 — Amer. trop.
A. cicutarium Sw. 1788. Mett. n. 70. HB. 220 (excl. β). Chr. 205. NPfl. 241; Caenopteris Thbg. 1795; Darea Willd. 1810.
cristatum Desv. Prod. 271. 1827 = A. obtusatum *obliquum.
cristatum Wall. List n. 211. 1828 = A. unilaterale.
cristatum Brack. Expl. Exp. **16**. 163 t. 21 f. 3. 1854 = A. cuneatum.
cristatum Mett.; Salom. Nom. 82. 1883 = Meniscium.

Cubense Hk. sp. **3**. 253 t. 207. 1860 = Diplazium semihastatum.
cultratum Sieb. Fl. Mart. exsicc. n. 365 = A. semicordatum.
cultratum Gaud. Freyc. Voy. Bot. 317. 1827 = A. adiantoides.
cultratum Roxb.; Wall. List n. 2214. 1829 (nomen) — India.
cultratum Hk. sp. **3**. 247. 1860 = Diplazium.
cultrifolium L. sp. **2**. 1081. 1753 = Diplazium.
cultrifolium Bory, Ess. Isles fort. 312. 1803 = A. monanthes.
cultrifolium Sieb. Syn. Fil. exs. n. 179 = A. semicordatum.
cultrifolium Roxb. Calc. Journ. **4**. 498. 1844 = A. adiantoides.
cultrifolium Gris. Fl. br. W. Ind. 680. 1864 = A. semicordatum.
Cumingii Mett. Fil. Lips. 74 t. 12 f. 8. 1856 = A. adiantoides.
cuneatiforme Christ, Bull. Boiss. II. **4**. 613. 1904 — Formosa.
cuneatum Lam. Enc. **2**. 309. 1786. Mett. n. 72. HB. 214. Chr. 204. NPfl. 240 — Trop. — Tarachia Pr. 1849 — Asplenium cristatum Brack. 1854 (non Lam. 1786).
cuneatum Hk. et Grev. Ic. Fil. t. 189. 1830 = A. praemorsum.
cuneatum Kze. Linn. **10**. 516. 1836 = A. lunulatum *erectum.
cuneatum auctt. quoad. pl. Hawaii = A. insiticium.
cuneatum F. Schultz, Flora **1844**. 807 = A. lanceolatum.
cuneatum var. caripense Kl. Linn. **20**. 356. 1847 = A. caripense.
cuneifolium Viv. Fragm. Fl. Ital. 16 t. 18. 1806 = A. adiantum nigrum*.
Currori Hk. sp. **3**. 82. 1860. HB. 192 — Guinea.
curtisorum Christ, Ann. Jard. Buit. II. **4**. 39. 1904 — Celebes.
curvatum Klf. Enum. 168. 1824 = A. sulcatum.
curvatum Fée, 9 mém. 16. 1857 = A. arcuatum.
cuspidatum Lam. Enc. **2**. 310. 1786 — Peru.
cyatheaefolium Rich. Sert. Astrol. XIX. 1834 = Diplazium.
cyathioides Bernh. Schrad. Journ. **1799**[1]. 313 = Athyrium filix femina.
cyrtopteron Kze. Linn. **23**. 233, 303. 1850 = A. radicans var.
cystopteroides Hk. sp. **3**. 220. 1860 = Athyrium.
Dalhousiae Hk. Ic. pl. t. 105. 1837 = Ceterach.
dareaefolium Bory; Willd. sp. **5**. 335. 1810 = A. praemorsum.
dareoides Desv. Berl. Mag. **5**. 322. 1811 = A. concisum.
dareoides Bory, Bél. Voy. Bot. **2**. 50. 1833 — Java.
dareoides Moritz, Verz. 110. 1845-46 = A. cuneatum.
daucifolium Lam. Enc. **2**. 310. 1786 — Ins. Mascarenae. Caenopteris Desv. 1827.
? **davallioides** Hk. JoB. **9**. 343. 1857. HB. 222. Chr. 209. NPfl. 242 — Japonia. Hongkong. — Humata dareoidea Mett. 1859.
davallioides Tausch, Flora **1839**. 479 = A. adiantum nigrum.
debile Fée, 10 mém. 28 t. 35 f. 2. 1865 — Bolivia.
debile Mett.; Kuhn, Fil. Afr. 101. 1868 = A. comorense.
debile Sod. Cr. vasc. quit. 141. 1893 = A. congestum.
decipiens Mett. Aspl. 195. n. 237. 1859 = Athyrium macrocarpum.
decipiens Kuhn, v. Deck. Reis. **3**[3]. Bot. 71. 1879 = A. protensum.
decorum Kze. Bot. Zeit. **1848**. 176 = A. tenerum *Belangeri.
decrescens Kze. Linn. **24**. 261. 1851. Mett. n. 144 — India austr. (Mont. Nilgir.).
decurrens Willd. sp. **5**. 316. 1810 = A. obtusatum *obliquum.
decurrens Wall. List n. 190. 1828 = A. unilaterale.
decurrens Bak. Syn. 484. 1874 = A. semiadnatum.

decurtatum Kze.; Link, Fil. sp. 94. 1841 = Athyrium.
decussatum Sw. Schrad. Journ. **1800**². 51. 1801 = Diplazium proliferum.
decussatum Wall. List n. 2208. 1829 = Diplazium japonicum.
deflexum Mett. Aspl. 184. n. 211. 1859 = Diplazium tomentosum.
deflexum Moore, Ind. 43, 123. 1859 = Diplazium Petersenii.
delicatulum Pr. Rel. Haenk. **1**. 47 t. 7 f. 3. 1825. Mett. n. 49. HB. 219. NPfl. 241 — Peru.
Delislei Bak. Ann. Bot. **5**. 301. 1891 — Réunion.
deltoideum Pr. Rel. Haenk. **1**. 47 t. 7 f. 2. 1825 = Diplazium.
densum Brack. Expl. Exp. **16**. 151 t. 20 f. 3. 1854 = A. trichomanes var.
dentato-serratum Fourn. Ann. sc. nat. V. **18**. 305. 1873 — N. Caledonia.
dentatum L. Syst. Nat. ed. X. **2**. 1323. 1759. Mett. n. 84. HB. 196. Chr. 192. NPfl. 235 — U. S. A. merid.-Ind. occ. Guatemala.
dentatum Krauss, Flora **1846**. 131; Pappe et Raws. Syn. Fil. Afr. austr. 19. 1858 = A. Kraussii.
dentex L. v. Buch, Beschr. Canar. Ins. 189. 1825 = A. monanthes.
dentex Kze. Linn. **23**. 233. 1850 = A. praemorsum.
denticulatum Bl. Enum. 186. 1828 — Java.
denticulatum J. Sm. Cat. cult. ferns 47. 1857 = Athyrium nigripes.
denticulosum Desv. Berl. Mag. **5**. 323. 1811 = Diplazium.
denudatum Mett.; Kuhn, Linn. **36**. 93. 1869. (HB. 483) — Peru.
deparioides Brack. Expl. Exp. **16**. 172. 1854 = Athyrium proliferum.
depauperatum Wall. List n. 234. 1828 (nomen) = A. laciniatum var.
depauperatum Fée, 7 mém. 52 t. 15 f. 3. 1857 — Bolivia.
depressum Loud.; Kze. Linn. **23**. 233. 1850 — Hort.
Desvauxii Mett. Aspl. 181. n. 205. t. 5 f. 4. 1859 = Diplazium costale.
dichotomum Hk. sp. **3**. 210. 1860; 2 Cent. t. 39. HB. 221 — Borneo.
dichroum Kze.; Pr. Tent. 108. 1836 = A. trichomanes.
Dietrichianum Luerss. Journ. Mus. Godeff. **3**. 16. 1873 = Diplazium.
difforme R. Br. Prod. Fl. Nov. Holl. 151. 1810. Mett. n. 53. (HB. 207); Chr. 197 — Australia. Ins. Norfolk. N. Zealand.
dilatatum Hk. sp. **3**. 258. 1860 = Diplazium maximum.
dimidiatum Sw. Prod. 129. 1788. Mett. n. 153. HB. 209. Chr. 198. NPfl. 239 — Amer. trop. Guinea. Tonkin. Perak. — Tarachia Pr. 1849 — ? Asplenium zamiaefolium Willd. 1810; Tarachia Pr. 1849.
dimidiatum Lowe, Ferns **5**. t. 13 A. 1858 = A. semicordatum.
diminutum Bak.; Jenm. JoB. **1881**. 53 = A. Fadyeni.
dimorphum Kze. Linn. **23**. 233. 1850. Mett. n. 55. HB. 223. Chr. 206. NPfl. 244 — Ins. Norfolk. — A. diversifolium A. Cunn.; Loud. Hort. Br. Suppl. 581. 1832 (non Bl. 1828).
diodon Fée, Gen. 195. 1850-52 — Ins. Philipp.
diplazioides Hk. et Arn. Bot. Beech. Voy. 107. 1832 = Diplazium Arnottii.
diplazioides Bory, Bél. Voy. Bot. **2**. 51. 1833 = Diplazium polypodioides.
diplotion Bak. Syn. 484. 1874 — N. Hebridae.
discolor Kze. Linn. **9**. 65. 1834 = A. semicordatum.
discolor Pappe et Raws. Syn. Fil. Afr. austr. 17. 1858 = A. gemmiferum.
dispersum Kze. Linn. **23**. 233, 304. 1850 = A. sulcatum.
dissectum Sw. Prod. 130. 1788 — Ind. occ. Columbia. Ecuador.
A. bissectum Sw. 1801. Mett. n. 138. HB. 211. NPfl. 239.
dissectum Gmel. Syst. Nat. **2**². 1302. 1791 (Plum. t. 46) = ? A. sulcatum.
dissectum Link, Hort. Berol. **2**. 68. 1833 = A. cristatum.

dissectum Brack. Expl. Exp. **16**. 170 t. 24. 1854 = A. schizophyllum.
distans Don, Prod. Fl. Nepal. 9. 1825 = Athyrium.
distans Fée, Gen. 198. 1850-52 — Mexico.
distans Brack. Expl. Exp. **16**. 155. 1854 = A. remotum.
distentum A. Br. Ind. Sem. Ht. Berol. 1860 = ? Diplazium expansum.
distichum Mett.; Salom. Nom. 84. 1883 = ? A. achilleifolium.
divaricatum Wall. List n. 2204. 1829 (nomen) — Singapore.
divaricatum Kze. Linn. **9**. 71. 1834. Mett. n. 68. HB. 219 — Peru. Chile.
divergens Mett.; Bak. Fl. bras. 1^2. 445. 1870. HB. 215 — Brasilia.
A. scandicinum Hk. sp. **3**. 183 t. 204. 1860 (non Klf. 1824).
diversifolium Bl. Enum. 175. 1828 — Java. — A. salignum Bak. 1867. HB. 199. (pt. ?). ? NPfl. 236 — A. concolor Hk. 1860. HB. 191 (f. indivisa).
diversifolium A. Cunn.; Loud. Hort. Brit. Suppl. 581. 1832 = A. dimorphum.
diversifolium Wall. List n. 203. 1828 = Diplazium maximum.
divisissimum Bak. Syn. 242. 1867 = Diplazium.
Döderleinii Luerss. Engl. Jahrb. **4**. 355. 1883 = Diplazium.
Dolabella Fée, Gen. 191. 1850-52 = A. lunulatum.
dolosum Milde, Verh. zool. bot. Ges. **14**. 165 t. 4. 1864 = A. adiantum nigrum × trichomanes.
Dombeyi Mett. Ann. sc. nat. V. **2**. 238. 1864 = Athyrium filix femina var.
Donianum Mett. Aspl. 177. n. 198 b. 1859 = Diplazium bantamense.
Donnell-Smithii Christ, Bot. Gaz. **20**. 544. 1895 = A. lunulatum.
Doreyi Kze. Anal. 23. 1837 = A. tenerum.
Douglasii Hk. et Grev. Ic. Fil. t. 150. 1829 = Phyllitis plantaginea.
Dregeanum Kze. Linn. **10**. 517. 1836. Mett. n. 66. HB. 221 — Africa trop. — A. brachypteron Kze. 1850. HB. 221. NPfl. 243 — (prob. = A. Sandersoni f. dareoidea).
drepanophyllum Kze. Linn. **9**. 66. 1834 = ? A. laetum.
drepanophyllum Bak. Syn. ed. II. 226. 1874 = Athyrium falcatum.
drepanopterum A. Br. Ind. Sem. ht. Berol. 1856; Mett. Aspl. 198. n. 244. 1859 = Athyrium.
Drueryii ht.; May; Garden **45**. 472. 1894; Gard. Chr. III. **15**. 665. 1894 — Hort.
duale Jenm. Gard. Chr. III. **13**. 10. 1893 (ex err. sub Adianto) — Jamaica.
dubiosum Dav. Gard. and Forest **4**. 483. 1891; Bot. Gaz. **21**. 264. 1896 = A. chihuahuense.
dubium Gaud. Freyc. Voy. Bot. 314. 1827 = Phyllitis brasiliensis.
dubium Brack. Expl. Exp. **16**. 172. 1854 (HB. 220) — Polynesia. N. Guinea.
dubium Mett. Aspl. 187. n. 216. 1859 = Diplazium ambiguum.
Dufourii Fée, 10 mém. 26 t. 35 f. 1. 1865 = A. prionites.
D'Urvillei Mett.; Kuhn, Linn. **36**. 95. 1869 — N. Zealand.
Duthiei Bedd. JoB. **1889**. 72 = Athyrium.
Eatoni Dav. Bot. Gaz. **21**. 258. 1896 — Mexico.
A. cicutarium var. paleaceum Dav. 1894 (non A. paleaceum R. Br. 1810).
ebeneum Ait. Hort. Kew. **3**. 462. 1789 = A. platyneuron.
ebeneum Sod. Cr. vasc. quit. 146. 1893 = A. Sodiroi.
ebenoides Scott; Berkeley, Journ. Hort. Soc. **1866**. 87 t. 2 f. 1. (HB. 198) — U. S. A. atlant.
ebenum Kuhn, Verh. zool. bot. Ges. **19**. 576. 1869 = Diplazium.
eburneum J. Sm. Cat. cult. ferns 47. 1857 = Athyrium drepanopterum.
efulense Bak. Kew Bull. **1901**. 137 — Kamerun.

Eggersii Sod. Cr. vasc. quit. 188. 1993 = Diplazium.
elasticum Fée, Gen. 196. 1850-52 — India.
elatius Link, Fil. sp. 94. 1841 = Athyrium filix femina var. angustum.
elatum Mett. Aspl. 180. n. 203. 1859 = Diplazium silvaticum.
elegans Mett. Fil. Lips. 74 t. 11 f. 5. 1856 = Diplazium fraxinifolium.
elegantulum Hk. sp. **3**. 190. 1860 = A. incisum.
elongatum Salisb. Prod. 402. 1796 = Phyllitis scolopendrium.
elongatum Sw. Syn. 79. 1806 = A. tenerum.
elongatum Mett.; Salom. Nom. 85. 1883 = Diplazium.
emarginato-dentatum Zenker; Kze. Linn. **24**. 263. 1851 = A. unilaterale.
emarginatum Pal. Beauv. Fl. d'Oware **2**. 6 t. 61 1807(?). HB. 200. Chr. 200. NPfl. 236 — Africa trop. Samoa.
enatum Brack. Expl. Exp. **16**. 153 t. 21 f. 1. 1854. HB. 205 — Ins. Hawaii.
ensiforme Wall. List n. 200. 1828; Hk. et Grev. Ic. Fil. t. 71. 1829. Mett. n. 130. HB. 191. Chr. 189. NPfl. 233 — India. Yunnan.
A. stenophyllum Bedd. 1866.
erectum Bory; Willd. sp. **5**. 328. 1810 = A. lunulatum*.
erectum Bl. Enum. 178. 1828 = A. longissimum.
eroso-dentatum Bl. Enum. 182. 1828 = A. obscurum.
erosum L. Syst. Nat. ed. X. **2**. 1324. 1759. Mett. n. 157. HB. 208. (NPfl. 239) India occ. — Tarachia Pr. 1849.
erythrocaulon Bl. Enum. 183. 1828 = A. unilaterale.
Escragnollei Fée, Cr. vasc. Br. **1**. 62 t. 15. 1869 = A. oligophyllum.
esculentum Pr. Rel. Haenk. **1**. 45. 1825 = Diplazium.
excisum Pr. Epim. 74. 1849 = A. unilaterale.
exiguum Bedd. Ferns S. Ind. t. 146. 1863 — China merid.-India austr. (conf. A. Glenniei) — (an A. fontanum var.?).
expansum Pr. Rel. Haenk. **1**. 46. 1825 = Diplazium.
extensum Fée, 7 mém. 51 t. 13 f. 2. 1857. HB. 197 — Columbia-Peru.
Fabianum Hombr. et Jacq. Voy. Pôle Sud t. 3 bis f. ζ. 1852 = A. bulbiferum var. vel A. lineatum var. vel sp. propr. (Ins. Mascarenae).
Fadyeni Hk. sp. **3**. 193. 1860; 2 Cent. t. 27. 1860. HB. 220 — Jamaica. — A. diminutum Bak. 1881.
falcatum Lam. Enc. **2**. 306. 1786; Retz. Obs. **6**. 37. 1791 = A. adiantoides.
falcatum Thbg. Prod. Fl. Cap. 172. 1800 = A. lunulatum.
falcatum Sw. Fl. Ind. occ. **3**. 1618. 1806 = A. erosum.
falcatum Bedd. Ferns S. Ind. t. 141. 1863 = A. crinicaule.
falcifolium Moore, Ind. 129. 1859 = ? A. adiantoides.
fallax Mett. Aspl. 194. n. 236. 1859 = Athyrium macrocarpum.
falsum Retz. Obs. **6**. 38. 1791 = A. praemorsum.
falx Desv. Prod. 274. 1827 = A. semicordatum.
fastigiatum Fourn. Ann. sc. nat. V. **18**. 305. 1873 — N. Caledonia.
Fauriei Christ, Bull. Boiss. **7**. 818. 1899 — Japonia.
Fawcettii Jenm. Gard. Chr. III. **26**. 121. 1899 — Jamaica.
Feei Kze.; Fée, Gen. 194. 1850-52; 7 mém. 49 t. 15 f. 2. 1857 — Mexico.
feejeense Brack. Expl. Exp. **16**. 147 t. 19 f. 1. 1854 = A. amboinense.
Fenzlianum Luerss. Flora **1875**. 434 = Diplazium.
Fenzlii Ettingsh. Denkschr. Akad. Wien **23**. 67 t. 11 f. 10-11. 1864 — Java.
fernandezianum Kze. Anal. 22. 1837 = A. lunulatum var.

ferulaceum Moore, Ind. 130. 1859 (nomen); Hk. sp. **3**. 216. 1860; 2 Cent. t. 38. HB. 224. Chr. 206. NPfl. 244 — Costa Rica-Ecuador.
Athyrium Christ 1904.
fibrillosum Pringle et Dav.; Dav. Bot. Gaz. **21**. 257 t. 18 f. 1-4. 1896 — Mexico.
ficifolium Goldm. Nova Acta **19**. Suppl. I. 461. 1843 = A. nidus var.
Fieldingianum Moore, Ind. 43. 1857 = Athyrium.
filamentosum Roxb. in Beats. St. Helena 299. 1816 = A. praemorsum.
filicaule Bak. JoB. **1881**. 204. Chr. 208 — Columbia. Costa Rica.
filiforme Klf. Enum. 172. 1824 = A. contiguum.
filipendulaefolium Thouars, Fl. Trist. d'Ac. 34 t. 4. 1804(?) = Gymnogramma cheilanthoides.
filix-femina Bernh. Schrad. neu. Journ. 1^2. 26, 48 t. 2 f. 7. 1806 = Athyrium.
fimbriatum Kze. Linn. **18**. 117. 1844 = A. varians.
fimbriatum Hk. sp. **3**. 234. 1860 = Athyrium.
Finckii Bak. Ann. Bot. **8**. 126. 1894. (Chr. 189) — Mexico.
(an = A. Ghiesbreghtii?).
Finlaysonianum Wall. List n. 191. 1828 (nomen) = A. macrophyllum.
Finlaysonianum Wall.; Hk. Ic. plant. t. 937. 1854. Mett. n. 149. HB. 245. Chr. 199. NPfl. 239 — India.
Hemidictyum Moore 1857; Asplenidictyum J. Sm. 1875. Asplenium Hookerianum Wall. 1829 (nomen); Hemidictyum Moore 1857.
firmum Kze. Bot. Zeit. **1845**. 283 = A. abscissum.
firmum Fée, Gen. 197. 1850-52 = A. platybasis.
fissidens Bory, Bél. Voy. Bot. **2**. 39. 1833 — Mauritius. Réunion.
fissum Kit.; Willd. sp. **5**. 348. 1810. Mett. n. 125. HB. 213. Chr. 202. NPfl. 240 — Europa centr. Italia. — Aspidium cuneatum Schkuhr 1809 (non Asplenium Lam. 1786); Athyrium Heufl. 1856. Asplenium angustifolium Guss. 1826.
flabellatum Kze. Bot. Zeit. **1845**. 285 (err.) = A. flabellulatum.
flabellifolium Cav. Descr. 257. 1802 (flavelifolium); Sw. Syn. 273. 1806. Mett. n. 118. HB. 195. Chr. 192. NPfl. 235 — N. Zealand. Tasmania. Australia.
flabellulatum Kze. Linn. **9**. 71. 1834 — Amer. trop.
flabellulatum Kl. Linn. **20**. 357. 1847 = A. partitum.
flabellulatum Mett. Aspl. 130. n. 99. 1859; a = A. radicans var. cyrtopteron
b = A. uniseriale.
flaccidum Forst. Prod. 80. 1786. Mett. n. 61. HB. 222. Chr. 208. NPfl. 241 — N. Zealand. Tasmania. Australia. Africa austr. — Caenopteris Thbg. 1795; Darea Willd. 1810 — Caenopteris odontites Thbg. 1795; Darea Willd. 1810; Asplenium R. Br. 1810. Caenopteris novae-zeelandiae Spr. 1799.
flagelliferum Wall. List n. 219. 1828 = A. longissimum.
flagelliferum Fée, 8 mém. 83. 1857 = A. partitum.
flavelifolium Cav. Descr. 257. 1802 = A. flabellifolium.
flavescens Mett. Ann. sc. nat. V. **2**. 234. 1864 = Diplazium.
flavidum Sod. Cr. vasc. quit. 148. 1893 — Ecuador.
flexuosum Schrad. Gött. gel. Anz. **1818**. 916 = A. gemmiferum var.
flexuosum Pr. Rel. Haenk. **1**. 46 t. 7 f. 1. 1825 = Diplazium Preslianum.
flexuosum Wikstr. Vet. Ak. Handl. **1825**. 438. 1826 — Guadeloupe. (Diplazium [sp. dub.).
foecundum Kze. Linn. **20**. 3. 1847 = A. compressum.
foeniculaceum HBK. Nov. Gen. et. Sp. **1**. 15. 1815 = A. fragrans*.
foliolosum Wall. List n. 2205. 1829 } = Athyrium foliolosum.
foliosum Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 255. 1902 } = Athyrium foliolosum.

fontanum (L.) Bernh. Schrad. Journ. **1799**[1]. 314. Mett. n. 119. HB. 216. Chr. 204. NPfl. 241 — Europa. Asia centr. — Polypodium L. sp. **2**. 1089. 1753; Athyrium Roth 1799; Aspidium Sw. 1801 — Athyrium Halleri Roth 1799, 1800; Aspidium Willd. 1810; Asplenium DC. 1815 — A. refractum Moore 1857. — (conf. A. exiguum).

fontanum × viride Christ, Farnkr. d. Schweiz 88 f. 11. 1900 — Alpes maritimes. A. Gautieri Christ in Burnat, Matér. p. fl. Alp. marit. 15. 1900 (non Hk. 1860).

Forbesii Bak. Ann. Bot. **5**. 312. 1891 = Diplazium.

foresiacum (Le Grand) Christ, Farnkr. d. Schweiz 84 f. 10. 1900 — Gallia centr. Italia. — A. Halleri var. foresiacum Le Grand, Bull. Soc. Fr. **16**. 61. 1869.

forisiense Le Grand, Statistique du Forez 252. 1873 = A. foresiacum.

Formosae Christ, Bull. Boiss. II. **4**. 613. 1904 = A. loriceum.

formosanum Bak. Ann. Bot. **5**. 305. 1891 — Formosa. A. Hancockii Bak. JoB. **1885**. 104 (non Maxim. 1883).

formosum Willd. sp. **5**. 329. 1810. Mett. n. 106. HB. 210. Chr. 193. NPfl. 236, 237 — Amer. trop. Afr. occ. trop. India austr. Ceylon. A. subalatum Hk. et Arn. 1840 — A. nanum Willd. 1810.

Forsteri Sadler, Descr. pl. epiph. Hung. 29. 1820 = A. adiantum nigrum *cuneifolium.

Forsterianum Col. Tasm. Journ. **2**. 171. 1845 = A. adiantoides.

Fournieri Kuhn; Fourn. Bull. Soc. Fr. **17**. 237. 1870 — Nicaragua.

fragile Pr. Tent. 108. 1836. Mett. n. 87. HB. 195. Chr. 191. NPfl. 235 — Mexico-Peru. Ins. Hawaii. — A. stoloniferum Pr. Rel. Haenk. **1**. 44 t. 6 f. 4. 1825 (non Bory 1804) — A. rhomboideum Brack. 1854.

fragrans Sw. Prod. 130. 1788. Mett. n. 48. HB. 216. (Chr. 203) — Amer. trop. Tarachia Pr. 1849 — Asplenium mexicanum Mart. et Gal. 1842. Mett. n. 47. — *A. foeniculaceum HBK. 1815; Caenopteris Desv. 1827. Asplenium abrotanoides Pr. 1825 — A. coriifolium Liebm. 1849.

fragrans Schkuhr, Kr. Gew. **1**. 199 t. 130 b. 1809 = A. praemorsum.

fragrans Hk. Ic. pl. t. 88. 1837 = A. hastatum.

Franconis Mett. Aspl. 166. n. 176. t. 5. f. 30. 1859 = Diplazium.

fraternum Pr. Epim. 74. 1849 = A. unilaterale.

fraternum Fée, 9 mém. 16. 1857 = A. laetum.

fraxinifolium Wall. List n. 194. 1828; Hk. 2 Cent. t. 19. 1860 = Diplazium bantamense.

frondosum Wall. List p. 63 (n. 230). 1829 = Diplazium maximum vel D. polypodioides.

frondosum Ettingsh. Denkschr. Ak. Wien **23**. 85. 1864 = Diplazium polypodioides.

fruticosum Vell. Fl. flum. **11** t 104. 1827; Arch. Mus. Nac. Rio Jan. **5**. 458. 1881 = Didymochlaena truncatula.

fugax Christ, Bull. Soc. Fr. **52** Mém. I. 53. 1905 — China merid.

fuliginosum Hk. sp. **3**. 120. 1860. HB. 203 — Borneo. A. lugubre Hk. 2 Cent. t. 3. 1860 (non Liebm. 1849).

fuliginosum Sod. Rec. 40. 1883 = Diplazium fuscum.

furcatum Thbg. Prod. Fl. Cap. 172. 1800 = A. praemorsum.

furcatum Brack. Expl. Exp. **16**. 170. 1854 = A. bipinnatifidum.

furcatum J. Sm. Cat. cult. ferns 45. 1857 = A. lineatum var. inaequale.

fuscipes Bak. JoB. **1879**. 304 = A. coenobiale.

fuscopubescens Hk. sp. **3**. 264. 1860 = Diplazium sandwichianum.

fuscum Bak. Ann. Bot. **5**. 312. 1891 = Diplazium.

Galeottii Fée, Gen. 192. 1850-52; 7 mém. 50 t. 16 f. 2 = A. monanthes.

galipanense Hier. Hedwigia **44**. 195 t. 6 f. 2. 1905 — Venezuela.
Gardneri Bak. Gard. Chr. **1873**. 712. HB. 485 — Ceylon. Sumatra. Khasia.
Gastonis Fée, Cr. vasc. Br. **1**. 70 t. 19 f. 2. 1869 — Brasilia.
Gaudichaudii Mett.; Kuhn, Linn. **36**. 103. 1869 = Diplazium.
Gautieri Hk. sp. **3**. 88 t. 144. 1860. HB. 193. NPfl. 234 — Ins. Nossi-Bé.
Gautieri Christ in Burnat, Matér. p. fl. Alp. marit. 15. 1900 = A. fontanum [× viride.
gedeanum Racib. Pterid. Buit. 223. 1898 = Athyrium.
geminaria Bory, Ess. Isles fort. 313. 1803 = Anogramma leptophylla.
gemmiferum Schrad. Gött. gel. Anz. **1818**. 916. Mett. n. 23. HB. 207. NPfl. 239 — Africa orient. et austr. cum ins. — A. lucidum Schlecht. 1825. Chr. 197 (pt.) (non Forst. 1786) — A. flexuosum Schrad. 1818. A. discolor Pappe [et Raws. 1858.
Geppii Carr. Cat. Welw. Pl. 2^2. 269. 1901 — Angola.
germanicum Weis, Pl. Crypt. Fl. Gött. 299. 1770. HB. 212. Chr. 201. NPfl. 235 — Europa. Kashmir. Hongkong. — Amesium Newm. 1844; Tarachia Pr. 1849; Asplenium Breynii Retz. 1779. Mett. n. 124. A. alternifolium Jacq. 1781; Scolopendrium Roth 1800; Phyllitis heterophylla Moench 1794; Asplenium trichomanes × septentrionale Aschers. 1896.
germanicum × trichomanes — Europa centr.
A. Heufleri Reichardt, Verh. zool. bot. Ges. **9**. 95 t. 4. 1859. HB. 196. A. pertrichomanes × septentrionale Aschers. Syn. Mitteleur. Fl. **1**. 77. 1896.
Ghiesbreghtii Fourn. Mex. pl. **1**. 111 t. 5. 1872. HB. 492 — Mexico. (conf. A. Finckii).
gibberosum (Forst.) Mett. Novara Exp. Bot. **1**. 211. 1870 — Polynesia.
Trichomanes Forst. Prod. 85. 1786; Davallia Sw. 1801. HB. 100, 470 (excl. 2); Loxoscaphe Moore 1853. Davallia glaucescens Hedw. 1803.
gibbosum Fée, Gen. 195. 1850-52 = A. semicordatum.
Gibertianum Hk. 2 Cent. t. 22. 1860; sp. **3**. 199. 1860. HB. 219 — Paraguay.
A. inciso-alatum Moore 1859 (nomen).
Gilliesianum Hk. et Grev. Ic. Fil. t. 73. 1829 = A. Gilliesii.
Gilliesii Hk. Exot. Flora **3** ad t. 208. 1827 — Peru. Bolivia.
A. Gilliesianum Hk. et Grev. 1829. HB. 195. NPfl. 235.
Gilpinae Bak. Journ. Linn. Soc. **16**. 200. 1877 = A. affine var.
glaberrimum Mett. Fil. Lips. 75 t. 11 f. 1, 2. 1856 = Diplazium crenato-serratum.
glandulosum Loisel. Notice 145. 1810 — Europa merid. occ. Marocco.
Polypodium Petrarchae Guérin 1804 (nomen); Asplenium DC. 1815. Mett. n. 116. HB. 197. Chr. 192. NPfl. 236. A. Vallis-clausae Requien 1813. A. pilosum Guss. 1844.
Glaziovii Bak. Fl. bras. 1^2. 455. 1870 = Diplazium herbaceum.
Glenniei Bak. Syn. 488. 1874 — Mexico. Arizona.
Athyrium gracile Fourn. Mex. pl. **1**. 102. 1872; Asplenium Hemsl. 1885. NPfl. 241 (non Don 1825). — vide Hope: Bull. Torr. Cl. **26**. 58. 1899.
glochidiatum Racib. Bull. int. Ac. Cracovie **1902**. 62 — Borneo.
Godmani Bak. Ann. Bot. **5**. 166 t. 11. 1891 = Diplazium.
Goeringianum Mett. Aspl. 198 n. 243 t. 6 f. 11-12. 1859 = Athyrium.
Goetzii Hier. Engl. Jahrb. **28**. 343. 1900 — Africa trop.
Gourlieanum Moore. Ind. 135. 1859 (nomen) — Penang.
gracile Don, Prod. Fl. Nepal. 8. 1825 = ? Athyrium nigripes.
gracile Fée, Gen. 198. 1850-52; 7 mém. 52 t. 27 f. 1. 1857 = ? A. cuneatum.
gracile Pappe et Raws. Syn. Fil. Afr. austr. 22. 1858 = A. lunulatum var.
gracile Hemsl. Biol. Centr. Amer. **3**. 634. 1885 = A. Glenniei.

gracilescens Mett. Ann. sc. nat. V. **2**. 237. 1864 = Diplazium.
gracillimum Col. Tr. N. Zeal. Inst. **22**. 453. 1890 — N. Zealand.
gracillimum Kuhn in Engl. Hochgbfl. trop. Afr. 103. 1892 = A. Kuhnianum.
gradatum Vell. Fl. flum. **11**. t. 105. 1827; Arch. Mus. Nac. Rio Jan. **5**. 458. 1881 = ? Diplazium denticulosum.
Graeffei Luerss. Fil. Graeff. 168. 1871 = Diplazium Brackenridgei.
graminoides Sw. Fl. Ind. occ. **3**. 1608. 1806 = Monogramma.
Grammitis Mett. Aspl. 181. n. 206. 1859 = Diplazium.
grammitoides Hk. Ic. pl. t. 913. 1854 = Diplazium.
Granatense Fée, Cr. vasc. Br. **1**. 64. 1869 = A. neo-granatense.
grande Sw. Syn. 77. 1806 — Ins. Mariannae.
A. macrophyllum Cav. Descr. 259. 1802 (non Sw. 1801).
grande Fée, 8 mém. 82. 1857 = Athyrium achilleifolium var.
grandifolium Sw. Prod. 130. 1788 = Diplazium.
grandifrons Christ, Bull. Boiss. **6**. 959. 1898 (nomen); **7**. 9. 1899 — Yunnan.
Gregoriae Bak. Kew Bull. **1896**. 40 — Madagascar.
Grevillei Wall. List n. 1036. 1828; Hk. et Grev. Ic. Fil. t. 228. 1831. Mett. n. 4. HB. 191. NPfl. 234 — Tavoy. Mishmee.
Neottopteris J. Sm. 1841; Thamnopteris Moore 1857.
Griffithianum Hk. Ic. pl. t. 928. 1854. Mett. n. 12. HB. 193. NPfl. 234 — [India bor. Hongkong.
Griffithii Bak. Syn. 239. 1867 = Diplazium.
guatemalense Hier. Engl. Jahrb. **34**. 461. 1904 = A. sessilifolium var.
Gueinzianum Mett.; Kuhn, Fil. Afr. 103. 1868. HB. 486 — Natal.
Guildingii Jenm. Gard. Chr. III. **15**. 70. 1894 — St. Vincent.
guineense Schum. Vid. Selsk. Afh. **4**. 232. (seors. 458). 1827 = A. africanum.
gymnogrammoides Kl.; Mett. Aspl. 193. n. 234 t. 16 f. 13-14. 1859 = Athyrium.
Hahnii Fourn. Mex. pl. **1**. 109. 1872 = Diplazium.
Halleri DC. Fl. franç. **5**. 240. 1815 = A. fontanum.
Halleri var. foresiacum Le Grand, Bull. Soc. Fr. **16**. 61. 1869 = A. foresiacum.
Hallii Hk. sp. **3**. 202. 1860; 2 Cent. t. 30. HB. 212. NPfl. 237 — Amazonas. Ecuador. Columbia. — A. pectinatum Moore 1859 (nomen).
Hancei Bak. Syn. ed. I. 208. 1867 = A. crinicaule.
Hancockii Maxim. Mél. Biol. **11**. 868. 1883 — Formosa.
Hancockii Bak. JoB. **1885**. 104 = A. formosanum.
Hanningtoni Bak. JoB. **1883**. 245 = A. Sandersoni var.
Hansii Aschers. Syn. Mitteleur. Fl. **1**. 78. 1896 = A. septentrionale × trichomanes.
Harovii Godr. Proc. Linn. Soc. **1**. 159. 1842 = A. trichomanes.
harpeodes Kze. Linn. **18**. 329. 1844 = A. lunulatum*.
Harrisii Jenm. Gard. Chr. III. **17**. 68. 1895. NPfl. 235 — Jamaica.
Harrisoni Bak. JoB. **1884**. 362 = Diplazium.
Hartianum Jenm. JoB. **1886**. 268 = Diplazium Klotzschii.
hastatum Bory; Munby, Fl. d'Alg. 111. 1847 (nomen) = A. hemionitis.
hastatum Kl.; Kze. Linn. **23**. 235, 305. 1850. Mett. n. 43. HB. 205 — Venezuela -Ecuador. — — A. fragrans Hk. 1837 (non Sw. 1788); A. odoratum Moore 1859.
hastile Christ, Engl. Jahrb. **24**. 108. 1897 = Diplazium.
Haussknechtii Godet et Reuter; Milde, Fil. Eur. 78. 1867. HB. 487 — Asia minor.
hemionitideum Bak. JoB. **1877**. 163 = Diplazium.
hemionitis L. sp. **2**. 1078. 1753. HB. 194. Chr. 190. NPfl. 235 — Portugal. Algérie. Marocco. Ins. Atlantidis.
A. palmatum Lam. 1786. Mett. n. 19; Tarachia Pr. 1849.

hemionitis Lam. Fl. franç. 1. 25. 1778 = Phyllitis.
hemionitis Lour. Fl. Cochinch. 832. 1790 = Polypodium (Selliguea) sp. dub.
hemionitoides Roxb. Calc. Journ. 4. 498. 1844 = Diplazium tomentosum.
Hendersoni Houlst. Gard. Mag. Bot. 3. 259. 1851 — Hort. (= ? A. ebenoides).
Henryi Bak. Ann. Bot. 5. 306. 1891 = Athyrium.
herbaceum Fée, 7 mém. 55 t. 22 f. 3. 1857 = A. sessilifolium.
herbaceum Bak. Syn. 492. 1874 = Diplazium.
Herbstii ht. May; Gard. Chr. III. 20. 470. 1896 — Hort.
herpetopteris Bak. Journ. Linn. Soc. 16. 200. 1877 = A. affine var.
heterocarpum Wall. List n. 218. 1828 = A. cheilosorum.
heterochroum Kze. Linn. 9. 67. 1834 = A. trichomanes var.
heterodon Bl. Enum. 179. 1828 = ? A. macrophyllum.
heterogenum Wenderoth, Schr. Ges. Nat. Marburg 2. (211). 1831 — ?
heterophlebium Mett.; Bak. Syn. 243. 1867 = Diplazium.
? heterophyllum Pr. Rel. Haenk. 1. 40. 1825 = A. pumilum.
heterophyllum A. Rich. Fl. N. Zél. 74. 1832 = A. flaccidum.
heteropteron Mett. Aspl. 174. n. 191. 1859 = Diplazium spinulosum.
Heufleri Reichardt, Verh. zool. bot. Ges. 9. 95 t. 4. 1859 = A. germanicum [× trichomanes.
hians Mett. Aspl. 188. n. 219. 1859 = Diplazium.
Hillebrandii C. Chr. Ind. 115. 1905 — Ins. Hawaii.
A. Mannii Hill. Fl. Haw. 594. 1888 (non Hk. 1860).
Hilsenbergii Sieb. Fl. mixt. n. 316 = A. pellucidum.
hirsutum Heyne; Wall. List n. 212. 1828 = A. praemorsum.
hirtum Klf. Enum. 169. 1824 = A. pellucidum.
Hohenackerianum Kze. Bot. Zeit. 1849. 771 = Athyrium.
holophlebium Bak. JoB. 1877. 163 — Ecuador. Costa Rica.
holophyllum Bak. Ann. Bot. 5. 301. 1891 — Formosa.
holosorum Christ, Bull. Boiss. 7. 10. 1898 — Yunnan. Hongkong.
Hookerianum Wall. List n. 2682 (nomen). 1829 = A. Finlaysonianum.
Hookerianum Wall. List n. 7090 (nomen). 1829 = Diplazium bantamense.
Hookerianum Col. Tasm. Journ. 2. 169. 1844. HB. 123. NPfl. 239 — N. Zealand. Australia. — A. adiantoides Raoul 1844. Mett. n. 94 (non Lam. 1786).
horizontale Bak. Ann. Bot. 8. 125. 1894 — Sumatra.
horridum Klf. Enum. 173. 1824. Mett. n. 140. HB. 211. NPfl. 239 — Malesia. [Ins. Hawaii.
humile Spr. Neu. Entdeck. 3. 6. 1822 = A. pumilum.
humile Bl. Enum. 185. 1828 (patria erronea) = A. adiantum nigrum.
humile Bak. Syn. ed. I. 230. 1867; Kuhn, Fil. Afr. 104. 1868 = Diplazium [silvaticum.
Huttoni Bak. Ann. Bot. 5. 311. 1891 = Diplazium.
hylophilum Hier. in Engl. Pflw. Ostafr. C. 84. 1895 = Diplazium.
hymenodes Mett. Aspl. 163. n. 169. t. 5 f. 18-19. 1859 = Diplazium semihastatum.
hymenophylloides Fée, 7 mém. t. 15 f. 4. 1857 = A. pumilum var.
hypomelas Kuhn, Fil. Afr. 104. 1868 — Fernando Po.
Davallia nigrescens Hk. 2 Cent. t. 93. 1861. HB. 101 (non Asplenium Bl. 1828); Loxoscaphe Moore 1861.
imbricatum Hk. et Grev. Ic. Fil. t. 165. 1829 = A. triphyllum var.
impressum Christ in C. Chr. Ind. 115. 1905 — Ins. Batjan.
A. canaliculatum Christ in Warb. Monsunia 1. 73. 1900 (non Bl. 1828).
inaequale Kze. Bot. Zeit. 1848. 176 = A. lineatum var.
inaequale Cord. Bull. soc. sci. et arts Réunion 1890-91 —. 1891; Fl. Réunion 60. 1895 = A. Cordemoyi.

inaequalidens Fée, 11 mém. 34 t. 9 f. 3. 1866 — Guadeloupe.
inaequilaterale Willd. sp. **5**. 322. 1810 = A. unilaterale (f. Cordemoy).
inaequilaterale Mart. et Gal. Mém. Ac. Brux. **15**. 57. 1842 = A. monanthes.
inaequilaterale Fée, 9 mém. 17. 1857 = Diplazium denticulosum.
inaequilaterum Mett. Aspl. 166. n. 175. 1859 = Diplazium denticulosum.
inciso-alatum Moore, Ind. 137. 1859 = A. Gibertianum.
incisuratum Fée, Cr. vasc. Br. **2**. 44 t. 94 f. 1. 1872-73 = A. semicordatum var.
incisum Thbg. Tr. Linn. Soc. **2**. 342. 1794. HB. 217. Chr. 203. NPfl. 241 — Japonia. China. — A. Trichomanes Thbg. Fl. Jap. 334. 1874 (non L. 1753) — A. elegantulum Hk. 1850.
incisum Opiz, Kratos **2**. 17. 1826 = A. adiantum nigrum *cuneifolium.
incisum J. Sm. JoB. **4**. 174. 1841 = Athyrium nigripes.
incisum ht. May; Gard. Chr. III. **15**. 665. 1894 — Hort.
incurvatum Fée, Cr. vasc. Br. **1**. 69 t. 18 f. 1. 1869 = A. serra.
induratum Hk. 2 Cent. t. 68. 1861 = A. pellucidum var.
induratum Christ, Bull. Soc. bot. Belg. **35**. 201. 1896 = Diplazium costari-[canum.
insigne Bl. Enum. 188. 1828 = A. nitidum.
insigne Liebm. Vid. Selsk. Skr. V. **1**. 246 (seors. 94). 1849 = A. serra.
insiticium Brack. Expl. Exp. **16**. 161 t. 22 f. 2. 1854 — Ins Hawaii. A. affine et cuneatum auctt. quoad pl. Haw.
insulare Carm. Tr. Linn. Soc. **12**. 512. 1818 = A. lunulatum var.
integerrimum Spr. Nova Acta **10**. 231. 1821 = A. juglandifolium.
integerrimum Wall.; Hk. et Grev. Ic. Fil. t. 136. 1829 = A. macrophyllum.
integrifolium Mett. Fil. Lips. 74 t. 12 f. 7. 1856 = Diplazium cordifolium.
integrum Fée, Gen. 193. 1850-52 = A. serratum.
interjectum Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 241 — China.
intermedium Pr. Del. Prag. **1**. 232. 1822 = A. viride.
intermedium Klf. in Sieb. Syn. Fil. exs. n. 68; Spr. Syst. **4**. 84. 1827 = A. [adiantoides.
intermedium Bl. Enum. 181. 1828 = A. macrophyllum.
irregulare Bak. Syn. 222. 1867 — Guinea. A. productum Mett.; Kuhn 1868.
jamaicense Jenm. JoB. **1886**. 268 — Jamaica. Guatemala.
Jamesoni Hk. sp. **3**. 184 t. 205. 1860. HB. 218. (Chr. 206). NPfl. 241 — Ecuador.
japonicum Thbg. Fl. Jap. 334. 1784 = Diplazium.
japonicum Kze. Ind. Sem. ht. Lips. .1845 = Onychium.
javanicum Bl. Enum. 175. 1828 = Diplaziopsis.
jucundum Fée, Cr. vasc. Br. **1**. 68 t. 17 f. 1. 1869 — Brasilia.
juglandifolium Lam. Enc. **2**. 307. 1786 — Jamaica. Venezuela-Columbia. Diplazium Cav.; Sw. 1806 — Asplenium integerrimum Spr. 1821 — A. Kapplerianum Kze. 1848 — A. salicifolium varr. Mett. n. 40. — A. Neogranatense Fée 1857.
juglandifolium Hk. Fil. Exot. t. 100. 1859; sp. **3**. 242. 1860 = Diplazium Roemerianum.
Kamchatkanum Gilbert, Fern Bull. **9**. 54. 1901 — Yeso. Kamtschatka. (Athyrium sp.?)
Kapplerianum Kze. Linn. **21**. 216. 1848 = A. juglandifolium.
Karstenianum Kl. Bot. Zeit. **1846**. 101; Linn. **20**. 353. 1847. (pt.); Hier. Hedwigia **44**. 194 t. 6 f. 1. 1905 — Columbia.
Katzeri A. Braun, (Ind. sem. ht. Berol.?) 1861 = Diplazium silvaticum.

Kaulfussii Schlecht. Adumbr. 29. 1825 — Ins. Hawaii.
A. protensum Klf. Enum. 167. 1824 (non Schrad. 1818).
Kaulfussii Pr. Tent. 106. 1836 = A. adiantoides.
Knudsenii Hill. Fl. Haw. 601. 1888 — Ins. Hawaii.
Klotzschii Mett. Fil. Lips. 79. 1856 = Diplazium.
Kohautianum Pr. Tent. 107. 1836 = A. lunulatum *pteropus.
Kraussii Moore, Ind. 139. 1859 (nomen); Hk. sp. **3**. 147 t. 180 A. 1860. HB. 195. NPfl. 235 — Africa austr. — A. dentatum Krauss, Flora **1846**. 131 (nomen); Pappe et Rawson, Syn. Fil. Afr. austr. 17. 1858 (non. L. 1753).
Kuhnianum C. Chr. Ind. 117. 1905 — Africa orient. trop.
A. gracillimum Kuhn in Engl. Hochgbfl. trop. Afr. 103. 1892 (non Col. 1890).
Kunzei Mett. Fil. Lips. 74. 1856 = Diplazium pinnatifidum.
Labillardieri Kze.; Mett. Aspl. 95. n. 29 t. 4 f. 11. 1859 — Ins. Bourou.
laceratum Desv. Prod. 278. 1827 = A. praemorsum.
lacerum Schlecht. et Cham. Linn. **5**. 612. 1830. Mett. n. 50 — Mexico.
Tarachia Pr. 1849.
laciniatum Don, Prod. Fl. Nepal. 8. 1825. HB. 211. NPfl. 239 — India bor.-Ceylon. Japonia. — A. caespitosum Wall. 1828; Tarachia Pr. 1849 — Asplenium planicaule Wall. 1828. Mett. n. 158. HB. 211. NPfl. 239. A. truncatum Don; Pr. 1836; Tarachia Pr. 1849.
lacinulatum Schrad. Gött. gel. Anz. **1824**. 870 — Brasilia.
laetum Sw. Syn. 79, 271. 1806. HB. 210. Chr. 194. NPfl. 237 — Amer. trop. Africa austr. Madagascar. — A. Schkuhrianum Pr. 1836. ? Mett. n. 105. — A. brachyotus Kze. 1836. Mett. n. 74. (Afr.).
laetum Schkuhr, Kr. Gew. **1**. 65 t. 70. 1809 = A. abscissum.
laetum Wall. List n. 209. 1828 = A. unilaterale.
laevigatum Dcne. Nouv. Ann. Mus. **3**. 347. 1834 = Diplazium microphyllum.
Laffanianum Bak. Gard. Chr. n. s. **17**. 673. 1882 = Diplazium.
Lamotteanum Héribaud, Note 1880; Le Grand, Bull. Soc. Fr. **30**. 74. 1883 = A. adiantum nigrum *cuneifolium.
lamprocaulon Fée, Gen. 197. 1850-52 = A. semicordatum var.
lanceolatum Huds. Fl. Angl. 454. 1762. Mett. n. 120. HB. 217. Chr. 203. NPfl. 241 — Europa occ. et merid. Mediterr. Ins. Atlantidis.
Tarachia Pr. 1849; Athyrium Heufl. 1856 — Asplenium obovatum Viv. 1824; Cystopteris Pr. 1836; Athyrium Fée 1850-52.
lanceolatum Forsk. Fl. Aeg.-Arab. 185. 1775 = A. praemorsum.
lanceolatum? var. elegans Hk. JoB. **9**. 342. 1857 = A. incisum.
lanceum Thbg. Fl. Jap. 333. 1784 = Diplazium.
lanciforme Fée, Gen. 192. 1850-52 = A. serratum.
laserpitiifolium Lam. Enc. **2**. 310. 1786. Mett. n. 163. HB. 215. Chr. 205. NPfl. 240 — Malesia. Polynesia. Australia trop. — Tarachia Pr. 1849.
laserpitiifolium Ham.; Don, Prod. Fl. Nepal. 9. 1825 = A. bulbiferum.
laserpitiifolium Bedd. Ferns S. Ind. t. 225. 1863 et auctt. quoad pl. ind. = [A. nitidum.
lasiopteris Mett. Fil. Lips. 78. 1856 = Diplazium Petersenii.
lassum Raddi, Pl. Bras. 37 t. 22 bis f. 4. 1825 (err. vide laxum) = A. mu- [cronatum.
Lastii C. Chr. Ind. 117. 1905 — Madagascar.
A. longisorum Bak. JoB. **1891**. 4 (non Bak. 1881).
lastreoides Bak. JoB. **1888**. 227 = Athyrium.
latifolium Bory, Ess. Isles fortunées 311 t. 6. 1803 = Ceterach aureum.
latifolium Don, Prod. Fl. Nepal. 8. 1825 = Diplazium maximum.

latifolium Sturm, Enum. pl. vasc. cr. chil. 28. 1858 = Athyrium.
Laurentii Bomm.; Christ, Bull. Boiss. **4**. 663. 1896. Chr. 199. NPfl. 239 — Congo.
Lauterbachii Christ in Schum. et Laut. Fl. deut. Südsee 130. 1901 — N. Guinea.
laxum R. Br. Prod. Fl. N. Holl. 151. 1810 = A. bulbiferum var.
laxum Raddi, Opusc. sci. Bol. **3**. 291. 1819 = A. mucronatum.
laxum Kuhn, Fil. Afr. 105. 1868 = Athyrium scandicinum var.
Lechleri Mett. Fil. Lechl. **1**. 16 t. 2 f. 10. 1856 = Diplazium.
lepidorachis Bak. JoB. **1890**. 264 = Diplazium.
lepidum Pr. Verh. Ges. Vaterl. Mus. **1836**. 63 t. 3 f. 1. HB. 487. Chr. 202. NPfl. 239 — Europa austr. Hungaria.
Tarachia Pr. 1849. Asplenium brachyphyllum Gasp. 1845.
leptochlamys Sod. Cr. vasc. quit. 200. 1893 = Diplazium leptogrammoides.
leptophyllum Sw. Obs. Bot. 403. 1791 = Anogramma.
leptophyllum Lag. Anal. Cienc. **5**. 155 t. 41 f. 3. 1802 — Hispania. (an A. fontanum var.?).
leptophyllum Fée, 7 mém. 50 t. 14 f. 2. 1857 = A. monanthes.
leptophyllum Schultz in Rabenh. Krypt. Fl. 2^3. 315. 1848 = A. ruta muraria.
lepturus J. Sm. JoB. **3**. 408. 1841; Pr. Epim. 72. 1849 = A. contiguum.
leucostegioides Bak. Ann. Bot. **5**. 302. 1891 — Ins. Maui (Hawaii).
Levyi Fourn. Bull. Soc. Fr. **19**. 257. 1872 — Nicaragua.
limbatum Willd. sp. **5**. 310. 1810 = Diplazium marginatum.
Linckii Kuhn, Fil. Deck. 22. 1867. HB. 487 — Africa orient. trop.
Lindbergii Mett. Ann. sc. nat. V. **2**. 236. 1864 = Diplazium.
Lindeni Hk. sp. **3**. 185. 1860 = A. pseudo-nitidum.
lineare Pr. Tent. 108. 1836 = Pteris tripartita.
lineatum Sw. Schr. Journ. **1800**^{2}. 51. 1801. HB. 202. Chr. 198 — Ins. Mascarenae.
Asplenium nodulosum Klf.; Spr. 1827. Mett. n. 31 — Darea inaequalis Willd. 1810; Caenopteris Bory 1810; Asplenium Kze. 1848 — Darea prolifera Willd. 1810 — D. violascens Bory 1833; Caenopteris Boj. 1837; Asplenium Moore 1860 — Darea bifida Bory 1833; Asplenium Pr. 1836. — vide et A. viviparum et A. Fabianum. Conf. Cordemoy: Rév. gén. de Bot. **8**. 85 ff.
lineatum Wawra, Maxim. Reise 195 t. 98. 1866 = A. oligophyllum.
lineolatum Mett. Ann. Lugd. Bat. **2**. 238. 1866 = Diplazium fraxinifolium.
linguaeforme Roxb. Calc. Journ. **4**. 497. 1844 = ? Polypodium Feei.
Linza Cesati, Rend. Ac. Napoli **16**. 26, 29. 1877 = Triphlebia.
Listeri C. Chr. Ind. 118. 1905 — Christmas Ins. (mare ind.).
A. centrifugale Bak. Journ. Linn. Soc. **25**. 360. 1889 (non Bak. 1874).
lividum Mett.; Kuhn, Linn. **36**. 100. 1869. HB. 486 — Venezuela. Angola.
lobatum Pappe et Raws. Syn. Fil. Afr. austr. 22. 1858 = A. lunulatum.
Lobbianum Hk. 2 Cent. t. 17. 1860 = Diplazium.
lobulatum Mett.; Kuhn, Linn. **36**. 100. 1869. HB. 486 — Ins. Hawaii.
lobulosum Wall. List n. 210. 1828 = Diplazium.
Loherianum Christ, Bull. Boiss. **6**. 152. 1898 — Luzon.
lomariaceum Christ, Verh. Nat. Ges. Basel **11**. 229. 1895 = Diplazium porphyrorachis.
lonchophyllum Mett. Aspl. 161 n. 165. 1859 = Diplazium subserratum.
longicauda Hk. 2 Cent. t. 69. 1861. HB. 200. Chr. 201. NPfl. 236 — Africa trop.
longifolium Schrad. Gött. gel. Anz. **1824**. 870 = ? A. serratum.
longifolium Don. Prod. Fl. Nepal. 7. 1825 = Diplazium lobulosum.

longipes Fée, Gen. 195. 1850-52; 7 mém. 49 t. 6 f. 3 = A. Wightianum vel A. vulcanicum.
longisorum Bak. JoB. **1881**. 204 = Diplazium.
longisorum Bak. JoB. **1891**. 4 = A. Lastii.
longissimum Bl. Enum. 178. 1828. Mett. n. 134. HB. 199. Chr. 197. NPfl. 236 — India bor.-Malesia. Mauritius. Juan Fernandez (t. Johow).
longissimum Bak. Ann. Bot. **5**. 301. 1891 (err.) = Diplazium longisorum.
Lorentzii Hier. Engl. Jahrb. **22**. 375. 1896. NPfl. 235 — Argentina.
loriceum Christ in C. Chr. Ind. 119. 1905 — Formosa. — A. Formosae Christ, Bull. Boiss. II. **4**. 613. 1904 (non A. formosanum Bak. 1891).
loriforme Hk. Ic. pl. t. 926. 1854 = A. angustum.
loxoscaphoides Bak. Tr. Linn. Soc. II. Bot. **2**. 354. 1887 — Africa orient. trop.
lucidum Burm. Fl. Ind. Fl. Cap. Prod. 28. 1768 = ? Polystichum pungens.
lucidum Forst. Prod. 80. 1786. Mett. n. 22. Chr. 197 (pt.). NPfl. 239 — N. Zealand. Tasmania. — A. obtusatum var. HB. 207 — A. Lyallii (Hk. fil. Fl. N. Zeal. 33 t. 77. 1854) Moore 1859.
lucidum Salisb. Prod. 403. 1796 = A. adiantum nigrum.
lucidum Schlecht. Adumbr. 25 t. 14. 1825 = A. gemmiferum.
ludens Bak. Journ. Linn. Soc. **19**. 294. 1882 — Ins. Salomonis.
lugubre Liebm. Vid. Selsk. Skr. V. **1**. 243 (seors. 91). 1849 = A. laetum.
lugubre Hk. **2** Cent. t. 3. 1860 = A. fuliginosum.
lunulatum Sw. Schrad. Journ. **1800²**. 52. 1801. Mett. n. 80. HB. 202. Chr. 193. NPfl. 237 — Trop. — (sp. collect.; typ. Africa austr.).
—* A. erectum Bory; Willd. 1810. Mett. n. 81 — A. mutilatum Klf. 1824 — A. harpeodes Kze. 1844 — A. lobatum Pappe et Raws. 1858 — A. pulchrum Thouars; Pr. 1836. Mett. n. 73.
—* A. pteropus Klf. 1824. Mett. n. 77 — A. stellatum Colla 1836; A. Fernandezianum Kze. 1837. Mett. n. 85 — A. Barteri Hk. 1860.
— A. tenellum Roxb. 1816 (St. Helena) — A. gracile Pappe et Raws. 1858 (non Don 1825 nec Fée 1850-52); A. Pappei Moore 1859.
— A. brasiliense Raddi 1825 (non Sw. 1817); A. Raddianum Gaud. 1827. A. auricularium Desv. 1827.
— A. Macraei Hk. et Grev. 1831 (Ins. Hawaii). A. strictum Brack. 1854.
— A. camptorachis Kze. 1851. Mett. n. 78 (India).
(conf. Lindm. Ark. för Bot. **1**. 213-218).
luzoniense Spr. Syst. **4**. 85. 1827 = Diplazium proliferum.
Lyallii Moore, Ind. 143. 1859 = A. lucidum var.
Lydgatei Hill. Fl. Haw. 596. 1888 — Ins. Hawaii.
Macdonelli Bedd. JoB. **1889**. 73 = Athyrium.
macilentum Kze.; Kl. Linn. **20**. 351. 1847 = A. sulcatum.
Mackinnoni Hope, JoB. **1896**. 124 = Athyrium.
Macraei Hk. et Grev. Ic. Fil. t. 217. 1831 = A. lunulatum var.
macrocarpon Moore, Ind. 143. 1859 = Athyrium (Fée).
macrocarpum Desv. Prod. 271. 1827 = A. monanthes.
macrocarpum Bl.; J. Sm. Cat. cult. ferns 47. 1857; Hk. sp. **3**. 222. 1860 = Athyrium.
macrochlaena Mett. Aspl. 195. n. 236 b. 1859 — Mexico. — (Athyrium sp.?).
macrodictyon Bak. JoB. **1877**. 163 = Diplazium.
macrodictyon Bak. Kew Bull. **1901**. 144 = A. Sanderi.
macrodon Fée, 10 mém. 28. 1865 — Quito. Brasilia.

macrophlebium Bak. Syn. 485. 1874; Ic. pl. t. 1646. NPfl. 237 — Fernando Po. Kamerun.

macrophyllum Sw. Schrad. Journ. **1800**². 52. 1801. HB. 209. Chr. 199. NPfl. 239 — India austr. Ceylon. Malesia. Polynesia. Ins. Comorae et Mascarenae.
Tarachia Pr. 1849 — Asplenium canaliculatum Bl. 1828; Tarachia Pr. 1849 — Asplenium Finlaysonianum Wall. 1828 (List n. 191); Tarachia Pr. 1849 — Asplenium integerrimum Wall.; Hk. et Grev. 1829 — A. platyphyllum J. Sm. 1841. Mett. n. 148 — A. oxyphyllum J. Sm. 1841. Mett. n. 152.

macrophyllum Cav. Descr. 259. 1802 = A. grande.

macrophyllum Mett. Fil. Lechl. **1**. 16. 1856 = Diplazium costale.

macropterum Sod. Cr. vasc. quit. 637. 1893 = Diplazium.

macrosorum Bertero; Colla, Mem. Ac. Torino **39**. 39 t. 67. 1836; Kze. Anal. 21. 1837. Mett. n. 90. HB. 206. NPfl. 237 — Juan Fernandez.

macrotis Bak. JoB. **1884**. 362 = Diplazium.

Mactierii Bedd. JoB. **1888**. 3 — Penang. — (HB. 193 sub A. Griffithiano).

madagascariense Bak. Journ. Linn. Soc. **15**. 417. 1876 = Diplazium arborescens.

maderense Penny; Loud. Ht. Brit. 494. 1850 = A. praemorsum.

magellanicum Klf. Enum. 175. 1824. Mett. n. 96. HB. 213. (Chr. 202). NPfl. 240 — Amer. austr. temp.

magnum Karst. Fl. Col. **1**. 69 t. 34. 1859 = A. squamosum.

malabaricum Mett. Fil. Lips. 74. 1856 = Diplazium esculentum.

manilense Spr. Syst. **4**. 89. 1827 = Diplazium.

Mannii Hk. 2 Cent. t. 60. 1861; HB. 221. Chr. 207. NPfl. 243 — Africa trop. [Madagascar.

Mannii Hill. Fl. Haw. 594. 1888 = A. Hillebrandii.

marginale Hill. Fl. Haw. 613. 1888 = Diplazium.

marginatum L. sp. **2**. 1082. 1753 = Diplazium.

marginatum Wall. List n. 2209. 1829 = Diplazium polypodioides.

marinum L. sp. **2**. 1081. 1753. Mett. n. 107. HB. 207. Chr. 196. NPfl. 237 — Europa occ. Ins. Atlantidis.

marinum Thouars, Fl. Trist. d'Ac. 34. 1804 = A. lunulatum.

Martensii Kze. Amer. Journ. Sci. II. **6**. 86. 1848 = Athyrium filix femina var.

Martensii Fée, 9 mém. 16. 1857 = A. salicifolium.

Martianum C. Chr. Ind. 120. 1905 — Brasilia.
A. angustatum Kze. 1839; Mett. Aspl. 167. n. 177. 1859. HB. 217. NPfl. 241 (non Desv. 1827 nec Bl. 1828).

martinicense Willd. sp. **5**. 344. 1810 = A. cuneatum.

martinicense Raddi, Opusc. sci. Bol. **3**. 291. 1819 = A. pseudo-nitidum.

mascareinense Desv. Prod. 278. 1827 = A. praemorsum.

mascarenhense Fée, Gen. 194. 1850-52. Mett. n. 30 — Mascarenae.

mastigophyllum Fée, 8 mém. 83. 1857 = A. cirrhatum.

Mathewsii Moore, Ind. 145. 1859 = A. vomeriforme.

Matthioli Gasp. Progr. Sci. Napoli **4**. 161. 1833 = A. ruta muraria.

maximum Don, Prod. Fl. Nepal. 8. 1825 = Diplazium.

Mayii ht. May; Gard. Chr. III. **15**. 665. 1894; **23**. 332 f. 140. 1898 — Hort.

Meanum Gay, Fl. chil. **6**. 500. 1853 (err.) = A. Neeanum.

medium Hk. sp. **3**. 227. 1860 = Athyrium.

megalophyllum Desv. Prod. 275. 1827 = A. macrophyllum.

megaphyllum Bak. JoB. **1890**. 264 = Diplazium.

meiotomum Hill. Fl. Haw. 596. 1888. NPfl. 242 — Ins. Hawaii.

melanocaulon Willd. Enum. 1072. 1809 = A. trichomanes var.
melanocaulon Bak. Syn. 240. 1867 = Diplazium.
melanochlamys Hk. sp. **3**. 259. 1860 = Diplazium.
melanolepis Franch. et Sav. Enum. pl. Jap. **2**. 226, 625. 1879 = Athyrium filix femina var.
melanolepis Col. Tr. N. Zeal. Inst. **20**. 227. 1888 = A. trichomanes var.
melanolepis Bak. JoB. **1890**. 264 = A. tonkinense.
melanopodium Mett.; Salom. Nom. 93. 1883 = Diplazium deltoideum.
melanopus Sod. Cr. vasc. quit. 189. 1893 = Diplazium.
melanorachis C. Chr. Ind. 121. 1905. — Mexico-Costa Rica.
A. nigricans Eat. Proc. Amer. Acad. **8**. 619. 1873. HB. 483 (non Kze. 1834).
Melleri Mett.; Kuhn, Fil. Afr. 106. 1868 = A. vagans.
membranaceum Mett.; Kuhn, Linn. **36**. 103. 1869 = Diplazium.
meniscioides Sod. Cr. vasc. quit. 188. 1893 = Diplazium.
Menziesii Hk. et Grev. Ic. Fil. t. 100. 1829 = A. monanthes var.
Mertensianum Kze. Bot. Zeit. **1848**. 525. Mett. n. 58 — Polynesia occ.
mesosorum Mak. Bot. Mag. Tokyo **12**. 120. 1898 = Athyrium.
Mettenianum Miq. Ann. Lugd. Bat. **3**. 174. 1867 = Diplazium.
Mettenii Kuhn, Fil. Deck. 20. 1867. HB. 488 — Ins. Comorae. Afr. orient. trop.
Mettenii Fourn. Ann. sc. nat. V. **18**. 311. 1873 = Diplazium proliferum.
mexicanum Mart. et Gal. Mém. Ac. Brux. **15**. 62 t. 15 f. 4. 1842 = A. fragrans.
Meyenianum Pr. Tent. 106. 1836; Epim. 73. 1849 = A. pellucidum.
Meyenianum Mett. Aspl. 189. n. 222. 1859 = Diplazium.
Michauxii Spr. Syst. **4**. 88. 1827 = Athyrium filix femina var.
microdonton Desv. Prod. 274. 1827 — ?
microphyllum Tineo in Guss. Fl. Sic. Syn. **2**². 884. 1844 = A. trichomanes.
microphyllum Mett. Ann. Lugd. Bat. **2**. 239. 1866 = Diplazium.
micropteron Bak. Syn. 488. 1874; Ic. pl. t. 1647. NPfl. 241 — Paraguay. Peru.
microsorum Mak. Bot. Mag. Tokyo —. 1892 = Athyrium.
microxiphion Bak. Kew Bull. **1896**. 40 — Ins. Natuna.
Mikani Pr. Del. Prag. **1**. 177. 1822 = Diplazium marginatum.
Mildei Kuhn, Linn. **36**. 95. 1869 = Diplazium.
millefolium Pr. Tent. 109. 1836 — Chile.
Darea? Fée 1850-52 — (prob. = A. myriophyllum).
Milnei Carr. in Seem. Fl. Vit. 353. 1873. HB. 484 — Ins. Lord Howe.
minimum Mart. et Gal. Mém. Ac. Brux. **15**. 55 t. 15 f. 1. 1842 = A. pumilum.
minus Bl. Enum. 183. 1828 — Java.
miradorense Liebm. Vid. Selsk. Skr. V. **1**. 243 (seors. 91). 1849 = A. laetum.
mixtum Roxb. Calc. Journ. **4**. 499. 1844 = Diplazium silvaticum.
Moccennianum Sod. Rec. 34. 1883 = Diplazium.
Mocquerysii Christ, Bull. Boiss. II. **1**. 553. 1901 — Madagascar.
modestum Maxon, Bull. Torr. Cl. **31**. 657 c. fig. 1904 — Mexico.
mohillense Kuhn, Fil. Afr. 107. 1868 = Diplazium.
monanthemoides Roxb. Calc. Journ. **4**. 497. 1844 = A. normale.
monanthemum Murray, Linn. Syst. Veg. ed. XIV. 933. 1784 = A. monanthes.
monanthes L. Mant. 130. 1767 — Africa cum ins. Atlantidis. Amer. trop. (Arizona-Chile). Ins. Hawaii. — A. monanthemum Murr. 1784. Mett. n. 111. HB. 197. Chr. 193. NPfl. 236 — A. Menziesii Hk. et Grev. 1829 (Hawaii) — A. leptophyllum Fée 1857 — A. Galeottii Fée 1850-52.
mongolicum Franch. Nouv. Arch. Mus. II. **7**. 161. 1883 = Athyrium.

monodon Liebm. Vid. Selsk. Skr. V. **1**. 247 (seors. 95). 1849 = A. sulcatum.
monstrosum Kze. Linn. **23**. 235, 305. 1850 = Athyrium filix femina.
montanum Willd. sp. **5**. 342. 1810. Mett. n. 129. HB. 213. Chr. 202. NPfl. 240 — U. S. A. orient.
Montbrisonis Fée, Gen. 198 t. 6 A. f. 3. 1850-52 — Réunion.
monteverdense Hk. 2 Cent. t. 41. 1860 = A. myriophyllum.
monticolum Jenm. JoB. **1882**. 326 = Diplazium.
Moorei Bak. Ann. Bot. **5**. 303. 1891 — Ins. Salomonis.
Moritzii Mett. Fil. Lips. 130 t. 11 f. 4. 1856 = Diplazium esculentum.
moupinense Franch. Nouv. Arch. Mus. II. **10**. 114. 1887 — Tibet.
mucronatum Pr. Del. Prag. **1**. 178. 1822. Mett. n. 82. HB. 212. Chr. 195. NPfl. 239 — Brasilia.

A. laxum Raddi 1819 (non R. Br. 1810); A. lassum Raddi 1825.

multicaudatum Wall. List n. 229. 1828 = Athyrium spectabile.
multicaule Wall. List n. 208. 1828 = A. normale.
multicaule Pr. Verh. Ges. Vaterl. Mus. **1836**. 65 t. 3 f. 2 = A. ruta muraria.
multicaule Scholtz, Enum. pl. Siles. 48. 1836 = A. adiantum nigrum *cuneifolium.
multifidum Brack. Expl. Exp. **16**. 171 t. 23 f. 2. 1854 = A. Shuttleworthianum.
multiflorum Roxb. Calc. Journ. **4**. 499. 1844 = Diplazium.
multiforme Krasser, Ann. Hofmus. Wien **15**. 4 t. 1. 1900 — Africa austr.
multijugum Wall. List n. 207. 1828; Mett. Aspl. 135 n. 109. 1859 = A. normale.
multilineatum Hk. sp. **3**. 102 t. 183. 1860 = A. remotum.
multisectum Bl. Enum. 185. 1828 = A. caudatum.
multisectum Brack. Expl. Exp. **16**. 174. 1854 = Athyrium scandicinum.
multisoratum Wall. List pag. 63 (n. 204). 1829 = Diplazium crenato-serratum.
murale Bernh. Schrad. Journ. **1799**[1]. 311 { α = A. ruta muraria. β = A. germanicum.
Murbeckii Dörfl. Oest. Bot. Zeit. **45**. 223. 1895 = A. ruta muraria × septentrionale.
muricatum Mett. Ann. Lugd. Bat. **2**. 239. 1866 = Athyrium umbrosum var.
murorum Lam. Fl. franç. **1**. 28. 1778 = A. ruta muraria.
musaefolium Mett. Aspl. 86. n. 3. 1859 = A. nidus var.
muticum Gilb. Amer. Botanist **4**. 86. 1903 — Bermuda. Florida.
mutilatum Klf. Enum. 171. 1824 = A. lunulatum.
mutilum Mett. Aspl. 180. n. 204. 1859 = Diplazium.
myapteron Fée, 8 mém. 82. 1857 — Mexico.
myriophyllum (Sw.) Pr. Rel. Haenk. **1**. 48. 1825 — Florida. Amer. trop.

Caenopteris Sw. Schrad. Journ. **1800**[2]. 60. 1801; Darea Willd. 1810. Caenopteris rhizophylla Thbg.; Sm. Pl. Ic. ined. **2** t. 50. 1790; Darea Willd. 1810; Asplenium Kze. 1834. HB. 220 (pt.). NPfl. 241. (non L. 1753 nec L. 1763). Darea membranacea Poir. 1811 — Asplenium monteverdense Hk. 1860. — (teste G. Hieronymus in litt.).

mysorense Roth, Nov. Pl. Spec. 395. 1821 = A. nitidum.
Naganumanum Mak. Bot. Mag. Tokyo —. 1892 = Diplazium.
nanum Willd. sp. **5**. 323. 1810 = A. formosum var.
natunae Bak. Kew Bull. **1896**. 41 — Ins. Natuna.
Neeanum Kze. Anal. 22. 1837. Mett. n. 151 — Chiloe.
neglectum Karst. Fl. Col. **1**. 106 t. 52. 1860 = Diplazium.
nemorale Bak. Journ. Linn. Soc. **15**. 417. 1876 = Diplazium arborescens var.
neogranatense Fée, 7 mém. 47 t. 14 f. 1. 1857 = A. juglandifolium var.
nephrodioides Bak. JoB. **1887**. 170 = Athyrium.

nervosum Mett. Ann. sc. nat. V. **2**. 235. 1864 = Diplazium.

Nesii Christ, Nu. Giorn. bot. It. n. s. **4**. 90. 1897 — China (Shen-si).

Newmani Bolle, Bonplandia 7. 106. 1859. (HB. 483) — Ins. Canarienses.

Newtoni Bak. Ann. Bot. **5**. 307. 1891 = Athyrium.

nicotianaefolium Mett. Ann. sc. nat. V. **2**. 234. 1864 = Diplazium.

nidus L. sp. **2**. 1079. 1753. Mett. n. 1. HB. 190. Chr. 188. NPfl. 233 — Asia, Polynesia, Australia trop. Africa orient. cum insulis.

Neottopteris J. Sm. 1842; Thamnopteris Pr. 1849; Neottopteris vulgaris J. Sm. 1841 — Asplenium australasicum Hk. 1859. Mett. n. 2; Neottopteris J. Sm. 1857 — Asplenium ficifolium Goldm. 1843. Neottopteris musaefolia J. Sm. 1841 (nomen); Thamnopteris Pr. 1849; Asplenium Mett. n. 3. 1859 — A. Phyllitidis Don 1825. Mett. n. 6; Neottopteris J. Sm. 1841; Thamnopteris Pr. 1849.

— vide et A. pachyhyllum et A. Simonsianum; conf. Thamnopteris et Neottopteris.

Nidus Raddi, Opusc. sci. Bol. **3**. 290. 1819; Pl. Bras. **1**. 34 t. 53. 1825 = A. serratum.

nigrescens Bl. Enum. 180. 1828. Mett. n. 142. ? Racib. Pterid. Buit. 217. 1898 — Ins. Moluccae. ? Java.

nigrescens Hk. fil. Tr. Linn. Soc. **20**. 170. 1847 = A. anisophyllum var.

nigricans Kze. Linn. **9**. 69. 1834 = A. praemorsum.

nigricans Eat. Proc. Amer. Acad. **8**. 619. 1873 = A. melanorachis.

nigripes Hk. JoB. **9**. 268 t. 9. 1857 = Phyllitis.

nigripes Bl.; Hk. sp. **3**. 222. 1860 = Athyrium.

nigritianum Hk. 2 Cent. t. 44. 1860 = A. pedicularifolium.

nigropaleaceum A. Br. Ind. Sem. ht. Berol. 1861. 2 = Diplazium.

nigrum Lam. Fl. franç. **1**. 28. 1778 = A. adiantum nigrum.

niponicum Mett. Ann. Lugd. Bat. **2**. 240. 1865 = Athyrium.

nitens Sw. Syn. 264, 421. 1806. Mett. n. 147. HB. 206. HB. 239. NPfl. 239 — [Ins. Mascarenae.

nitidulum Moore, Ind. 43. 1857 = Athyrium.

nitidulum Hill. Fl. Haw. 601. 1888 — Ins. Hawaii.

nitidum Sw. Syn. 84, 280. 1806. Mett. n. 162. HB. 215. Chr. 204. NPfl. 240 Ind. bor.-Malesia.

Tarachia Pr. 1849 — Asplenium insigne Bl. 1828; Tarachia Pr. 1849.

nodosum L. sp. ed. II. **2**. 1539. 1763 = Danaea.

nodosum Lour. Fl. Cochinch. **2**. 832. 1790 — Cochinchina.

nodulosum Klf.; Spr. Syst. **4**. 83. 1827 = A. lineatum.

normale Don, Prod. Fl. Nepal. 7. 1825. HB. 197. Chr. XII. NPfl. 236 — China austr.-India bor.-Ceylon. Celebes. Hawaii.

A. multijugum Wall. 1828. Mett. n. 109 — A. opacum Kze. 1851. Mett. n. 108 — A. pavonicum Brack. 1854 (Hawaii).

notabile Fée, 10 mém. 27 t. 36 f. 1. 1865 = A. lineatum.

novae-caledoniae Hk. Ic. pl. t. 911. 1854. HB. 223. Chr. 207. NPfl. 243 — N. Caledonia.

novum Sadl. Descr. pl. epiph. Hung. 8. 1820 = A. adiantum nigrum *cunei- [folium.

nubilum Moore, Ind. 150. 1859 = A. anisophyllum var.

obesum Bak. Syn. 209. 1867. NPfl. 239 — Guatemala.

obliquum Forst. Prod. 80. 1786 = A. obtusatum*.

obliquum Wall. List n. 2217. 1829 — Mauritius.

oblongatum Mett.; Kuhn, Linn. **36**. 97. 1869. HB. 484 — Panama.

oblongifolium Col. Tasm. Journ. **2**. 171. 1845 = A. obtusatum *obliquum.

obovatum Viv. Fl. Lib. 68. 1824 = A. lanceolatum var.
obovatum Christ, Bull. Boiss. II. **4**. 971. 1904 = A. obversum.
obscurum Bl. Enum. 181. 1828 — Java. — A. serraeforme Mett. n. 75 1859. (HB. 210).
obtusatum Forst. Prod. 80. 1786. Mett. n. 20. HB. 207 (excl. γ). Chr. 197. NPfl. 238 — Australia. N. Zealand. Ins. antarct. Chile austr. Tristan d'Acunha.
— A. chondrophyllum Bert.; Colla 1836 (Juan Fernandez).
—* A. obliquum Forst. 1786. Mett. n. 21. Chr. 197 — A. crassum Thouars 1804 (Tristan d'Acunha) — A. scleropium Hombr. et Jacq. 1852. — A. sphenoides Kze. 1834 (Chile).
obtusifolium L. sp. **2**. 1080. 1753. Mett. n. 38. HB. 202. Chr. 195. NPfl. 237 — Amer. trop. — A. aquaticum Kl. 1847 — A. riparium Liebm. 1849.
obtusilobum Desv. Berl. Mag. **5**. 323. 1811 = A. cuneatum.
obtusilobum Hk. Ic. pl. t. 1000. 1854 = A. oceanicum.
obtusissimum Fée, Gen. 197. 1850-52 = A. monanthes.
obtusum Kit.; Willd. sp. **5**. 341. 1810 = A. adiantum nigrum.
obtusum Mett. Fil. Lips. 78. 1856 = Diplazium Wageneri.
obtusum Mett. Fil. Lechl. 17. 1856; Aspl. 187 n. 217 = Diplazium expansum.
obtusum (Desv.) Kuhn, Fil. Afr. 109. 1868 — Mauritius.
Darea Desv. Berl. Mag. **5**. 323. 1811. (an A. affine?).
obversum Christ, Bull. Boiss. II. **5**. 260. 1905 — Costa Rica.
A. obovatum Christ, Bull. Boiss. II. **4**. 971. 1904 (non Viv. 1824).
ocanniense Karst. Fl. Col. **1**. 173 t. 86. 1861 — Columbia.
oceanicum C. Chr. Ind. 124. 1905 — Polynesia. Celebes.
A. obtusilobum Hk. Ic. pl. t. 1000. 1854. HB. 221 (non Desv. 1811).
ochraceum Sod. Cr. vasc. quit. 207. 1893 = Diplazium fuscum.
odontites R. Br. Prod. Fl. Nov. Holl. 151. 1810 = A. flaccidum.
odontophyllum Wall. List n. 2216. 1829 = A. formosum.
odoratum Moore, Ind. 151. 1859 = A. hastatum.
Oldhami Hance, Ann. sc. nat. V. **5**. 256. 1866 — Formosa.
oligolepidum C. Chr. Ind. 124. 1905 — N. Caledonia.
A. subnudum Bak. Syn. 483. 1874 (non Karst. 1865).
oligophlebium Bak. Gard. Chr. n. s. **14**. 494. 1880 — Japan.
oligophyllum Klf. Enum. 166. 1824. Mett. n. 28. HB. 201. Chr. 198 — Brasilia. Columbia. Ecuador. Madagascar.
A. Escragnollei Fée 1869 — A. camptocarpum Fée 1869.
oligophyllum Luerss. Fil. Graeff. 152. 1871 = ? A. sumatranum.
Onopteris L. sp. **2**. 1081. 1753 = A. adiantum nigrum*.
onustum Christ, Prim. Fl. costaric. **3**. 24. 1901 — Costa Rica.
oosorum Bak. JoB. **1876**. 343 = Athyrium.
opacum Kze. Linn. **24**. 261. 1851 = A. normale.
orientale Bernh. Schrad. Journ. **1801**[1]. 17. 1802 = Blechnum.
ornatum Col. Tr. N. Zeal. Inst. **22**. 452. 1890 — N. Zealand.
oroupouchense Prestoe; Jenm. Gard. Chr. III. **18**. 388. 1895 — Trinidad.
Otites Link, Hort. Berol. **2**. 60. 1833 = A. pulchellum var.
otophorum Miq. Ann. Lugd. Bat. **3**. 175. 1867 = Diplazium.
Ottonis Hk. sp. **3**. 243. 1860 = Diplazium.
ovalescens Fée, Cr. vasc. Br. **1**. 72 t. 18 f. 2. 1869 = A. pseudo-nitidum var.
ovatum Wall. List n. 195. 1828 = Syngramma alismifolia.
oxyphyllum Wall. List n. 223. 1828 = A. pellucidum.

oxyphyllum J. Sm. JoB. **3**. 408. 1841; Kze. Bot. Zeit. **1846**. 441 = A. macrophyllum.
oxyphyllum Hk. sp. **3**. 221. 1860 = Athyrium drepanopterum.
pachyphyllum Kze. Bot. Zeit. **1848**. 146. Mett. n. 7 — Java.
Thamnopteris Pr. 1849; Neottopteris Fée 1850-52. — (A. nidus*).
pachysorum Bak. JoB. **1891**. 4. NPfl. 236 — Madagascar.
paleaceum R. Br. Prod. Fl. N. Holl. 150. 1810. HB. 208 — Australia. Tahiti.
pallidum Bl. Enum. 177. 1828 = Diplazium.
palmatifidum Moore, Ind. 151. 1859 (nomen) — Mauritius.
palmatum Lam. Enc. **2**. 302. 1786 = A. hemionitis.
paludosum Moore, Ind. 152. 1859 = Athyrium nigripes.
Pappei Moore, Ind. 152. 1859 = A. lunulatum var.
paradoxum Bl. Enum. 179. 1828. HB. 209 — Penang. Sumatra. Java.
Tarachia Pr. 1849. Asplenium zamioides Hk. 1860.
parallelum Wall. List n. 228. 1828 (nomen) = Diplazium sorzogonense.
parallelum Bak. Syn. 486. 1874 — Ins. Hawaii. — A. bipinnatum Hill. 1888.
partitum (Kl.) C. Chr. Ind. 125. 1905 — Amer. trop. (Venezuela-Ecuador).
A. flabellulatum β partitum Kl. Linn. **20**. 357. 1847 (non A. flabellulatum Kze. 1834); A. radicans var. partita Hier. Engl. Jahrb. **34**. 464. 1904; A. flagelliferum Fée 1857; A. rhizophorum var. tripinnatum Mett. Ann. sc. nat. V. **2**. 229. 1864 — A. radicans Sw. Syn. 84. 1806 (non L. 1759); A. flabellulatum α dentatum Kl. Linn. **20**. 357. 1847. — (teste G. Hieronymus in litt.).
parvulum Wall. List. n. 2207. 1829 = A. varians.
parvulum Hk. Ic. pl. t. 222. 1840 = A. trilobum.
parvulum Mart. et Gal. Mém. Ac. Brux. **15**. 60 t. 15 f. 3. 1842 = A. resiliens.
patens Klf. Enum. 175. 1824 — Ins. Hawaii.
Diplazium Pr. 1836; Athyrium Milde 1870. (an Diplazium sp.?).
pavonicum Brack. Expl. Exp. **16**. 150 t. 20 f. 1. 1854 = A. normale.
Pearcei Bak. Syn. 483. 1874 — Peru.
Pechuëllii O. Ktze. Flora **1880**. 303 = A. trichomanes.
pectinatum Wall. List n. 231. 1828 = Athyrium filix femina*.
pectinatum Moore, Ind. 152. 1859 = A. Hallii.
pedatum Salom. Nom. 97. 1883 = Diplazium.
pedicularifolium St. Hil. Voy. Dist. Diam. **1**. 380. 1833 — Brasilia. Guinea.
A. nigritianum Hk. 1860. HB. 215; Athyrium J. Sm. 1875.
Pekinense Hance, JoB. **1867**. 262 = A. Sarelii var.
pellucidum Lam. Enc. **2**. 305. 1786 (excl. β). Mett. n. 135 — Malesia-Polynesia. Madagascar. Mascarenae. Sechellae.
A. hirtum Klf. 1824. HB. 205. Chr. 196. NPfl. 236 — A. induratum Hk. 1861 — A. stenopteris Kze. 1848. Mett. n. 136.
pellucidum β Lam. Enc. **2**. 305. 1786 = A. abscissum.
?penangianum Wall. List n. 196. 1828 = Blechnum Finlaysonianum.
pendulum Fée, Gen. 196. 1850-52 = A. lunulatum *harpeodes.
Perkinsi Jenm. Gard. Chr. III. **19**. 8. 1896 — Guiana.
persicifolium J. Sm. JoB. **3**. 408. 1841; Mett. Aspl. 97. n. 32. 1859. HB. 202 — Ins. Philipp. Celebes.
persicifolium Bedd. Ferns S. Ind. t. 127. 1863 = A. Zenkerianum.
persicifolium var. atrovirens Hk. sp. **3**. 109. 1860 = A. atrovirens.
peruvianum Desv. Prod. 271. 1827 — Peru.
Petersenii Kze. Anal. 24. 1837 = Diplazium.

petiolare Ces. Rend. Acad. Napoli **16**. 26. 1877 = Diplazium.
petiolulatum Mett.; Kuhn, Fil. Afr. 110. 1868. (HB. 486) — Réunion.
Petrarchae DC. Fl. franç. **5**. 238. 1815 = A. glandulosum.
Phyllitidis Don, Prod. Fl. Nepal. 7. 1825 = A. nidus var.
physosorum Sieb. Fl. Mixt. n. 268 = Athyrium umbrosum* australe.
pilosum Guss. Fl. Sic. Syn. 2^2. 661. 1844 = A. glandulosum.
pimpinellifolium Fée et Schaffn; Fée, 7 mém. 52 t. 25 f. 5. 1857 = A. semi-
pinnatifido-pinnatum Hk. sp. **3**. 238. 1860 = Diplazium. [cordatum.
pinnatifidum Nutt. Gen. **2**. 251. 1818. Mett. n. 91. HB. 194. Chr. 190 — U. S. A. atlant. — Scolopendrium Diels 1899. NPfl. 231.
planicaule Wall. List n. 189. 1828 = A. laciniatum var.
planicaule Lowe, Ferns **5**. t. 10. 1858 = A. fragrans.
plantagineum L. sp. ed. II. **2**. 1537. 1763 } = Diplazium plantagi-
plantaginifolium L. Syst. Nat. ed. X. **2**. 1323. 1759 } nifolium.
platybasis Kze.; Mett. Aspl. 155. n. 154. 1859. HB. 206. NPfl. 239 — St. Helena. A. firmum Fée 1850-52 (non Kze. 1845).
platychlamys Fée, 7 mém. 48 t. 14 f. 3. 1857 = A. abscissum.
platyneuron (L.) Oakes; Eat. Ferns N. Am. **1**. 24. 1878. NPfl. 236 — U. S. A. Jamaica. Africa austr. — Acrostichum L. sp. **2**. 1069. 1753; Asplenium ebeneum Ait. 1789. Mett. n. 110. HB. 198. Chr. 193. A. polypodioides Sw. 1801. A. trichomanoides Michx. 1803.
platyphyllum J. Sm. JoB. **3**. 408. 1841; Mett. Aspl. 153 n. 148. 1859 = A. macro-
platyphyllum Bak. JoB. **1890**. 264 = Diplazium stenochlamys. [phyllum.
plumbeum Christ, Bull. Boiss. II. **4**. 1089. 1904 — Costa Rica.
plumosum Bory; Willd. sp. **5**. 323. 1810 = A. pellucidum.
Poeppigii Pr. Tent. 106 t. 3 f. 21. 1836 = A. serra.
Poiretianum Gaud. Freyc. Voy. Bot. 321 t. 13. 1827 = Athyrium scandicinum.
polyanthes Sol.; Forst. Prod. 93. 1786 (nomen); Bak. Syn. 492. 1874 = Diplazium.
polydactylon Webb in Hk. Niger Fl. 193. 1849 = Actiniopteris australis.
polymeris Moore, Ind. 154. 1859 = A. polyphyllum.
polymorphum Wall. List n. 230. 1828 = Diplazium maximum.
polymorphum Mart. et Gal. Mém. Ac. Brux. **15**. 56 t. 15 f. 2. 1842 = A. abscissum.
polymorphum Mett. Aspl. 192. n. 229. 1859 = ? Athyrium Fieldingianum.
polyodon Forst. Prod. 80. 1786 = A. adiantoides var.
polyodon Wall. List pag. 63. 1829 = A. protensum.
polyphyllum Bert. Nov. Comment. Bonon. **4**. 443. 1840 — Guatemala. A. polymeris Moore 1859.
polyphyllum Pr. Tent. 108. 1836 (nomen); Goldm. Nova Acta **19**. Suppl. I. 462. 1843 = A. acuminatum.
polypodioides Sw. Schrad. Journ. **1800^2**. 53. 1801 = A. platyneuron.
polypodioides Mett. Fil. Lips. 78. 1856 = Diplazium.
Polypodium Bory ap. Willd. sp. **5**. 322. 1810 (syn.); Desv. Prod. 273. 1827 = A.
polyrhizon Bak. Syn. 490. 1874 = Diplazium japonicum var. [unilaterale.
polystichoides Bl.; Mett. Aspl. 160. 1859 = A. nitidum.
Poolii Bak. Journ. Linn. Soc. **15**. 416. 1876; Ic. pl. t. 1645. Chr. 201. NPfl. 236 —
porphyrocaulon Bl. Enum. 182. 1828 = A. unilaterale. [Madagascar.
porphyrorachis Bak. JoB. **1879**. 40 = Diplazium.
porrectum Wall. List n. 204. 1828 = Diplazium crenato-serratum.
porrectum Wall. List n. 224. 1828 = A. protensum.
Powellii Bak. Syn. 224. 1867 = A. Shuttleworthianum*.

praemorsum Sw. Prod. 130. 1788. Chr. 204. NPfl. 240 — Trop. et Subtrop. A. adiantoides Lam. 1786 (non supra). A. furcatum Thbg. 1800. Mett. n. 161. HB. 214, 487; Tarachia Pr. 1849; Asplenium canariense Willd. 1810 (Ins. Atlant.) — ? Acrostichum filare Forsk. 1775.
Preismanni Aschers. et Luerss. Verh. Bot. Ver. Brand. **37**. XLVI. 1896 = A. ruta muraria × trichomanes.
Prenticei Bail. Tr. Linn. Soc. N. S. Wales **4**. 37. 1880 = Diplazium pallidum.
Prescottianum Ces. Atti Ac. Napoli 7^8. 21. 1876 = Diplazium porphyrorachis.
Preslii Ettingsh. Farnkr. 148 t. 97 f. 3. 1865 — »Asia« — (Diplazium sp. dub.).
Pringlei Dav. Gard. and For. **4**. 449 f. 71. 1891 — Mexico.
prionites Kze. Linn. **10**. 511. 1836. Mett. n. 24. HB. 204 — Africa austr. Ins. Mascarenae. — A. Dufourii Fée 1865.
prionurus J. Sm. JoB. **3**. 408. 1841; Mett. Aspl. 97. n. 33. 1859. HB. 202 — Ins. Philipp.
procerum Bernh. Schrift. Akad. Erf. **1802**. 17 t. 1 f. 1 = Blechnum capense.
? procerum Wall. List n. 2203. 1829 = Athyrium.
productum Pr. Rel. Haenk. **1**. 42 t. 8 f. 1. 1825 = A. tenerum.
productum Lowe, Tr. Cambr. Phil. Soc. **6**. 524. 1838 = A. adiantum nigrum.
productum Mett.; Kuhn, Fil. Afr. 112. 1868 = A. irregulare.
progrediens Fée, 8 mém. 81. 1857 = A. serra.
projectum Kze. Linn. **9**. 68. 1834. Mett. n. 86. HB. 194. Chr. 192. NPfl. 235 — Peru.
proliferum Lam. Enc. **2**. 307. 1786 = Diplazium.
proliferum Sw. Prod. 129. 1788 = Fadyenia Fadyenii.
proliferum Wall. List n. 202. 1828 = Diplazium esculentum.
proliferum Hk. sp. **3**. 239. 1860 = Diplazium silvaticum var.
prolixum Schrad. Gött. gel. Anz. **1824**. 870 = A. sulcatum.
prolongatum Hk. 2 Cent. t. 42. 1860 = A. achilleifolium.
protensum Schrad. Gött. gel. Anz. **1818**. 916. Mett. n. 137. HB. 211. Chr. 195. NPfl. 237 — Afr. trop. et austr. Ins. Mascarenae.
protensum Klf. Enum. 167. 1824 = A. Kaulfussii.
pseudo-falcatum Hill. Fl. Haw. 597. 1888 — Ins. Hawaii.
pseudo-nitidum Raddi, Pl. Bras. **1**. 39 t. 55. 1825. Mett. n. 93. HB. 217. Chr. 205. NPfl. 241 — Brasilia. Venezuela-Ecuador. — A. martinicense Raddi 1819 (non Willd. 1810) — A. Lindeni Hk. 1860 — A. ovalescens Fée 1869.
psilopteron Salom. Nom. 99. 1883 = Diplazium.
pteridis (Bory) Kuhn, Fil. Afr. 113. 1868 — Mauritius.
Darea Bory, Bél. Voy. Bot. **2**. 52. 1833.
pteridoides Bak. JoB. **1873**. 17; Ic. pl. t. 1649. HB. 488. NPfl. 243 — Ins. [Lord Howe.
pterophorum Pr. Tent. 107. 1836 = A. alatum.
pteropus Klf. Enum. 170. 1824 = A. lunulatum*.
puberulum Wall. List n. 2212. 1829 = Diplazium esculentum.
pubescens Moore et Houlst. Gard. Mag. Bot. **3**. 262. 1851 = Athyrium decurtatum.
pubescens Mett. Fil. Lips. 78 t. 11 f. 3. 1856 = Diplazium esculentum.
pubigerum Mett.; Salom. Nom. 98. 1883 = Diplazium.
pulchellum Raddi, Opusc. sci. Bol. **3**. 291. 1819; Pl. Bras. **1**. 37 t. 52 f. 2. 1825. Mett. n. 83. HB. 210. NPfl. 237 — Brasilia. Columbia-Peru. Afr. orient. trop. — A. Serronii Glaz. 1869 — A. otites Link 1833. NPfl. 237.
pulchellum Wall. List n. 214. 1828 = A. nitidum.
pulchrum Thouars; Pr. Tent. 108. 1836 (Wall. List n. 2211) = A. lunulatum [*erectum.
pulchrum Loud. Hort. Brit. 494. 1850 — Jamaica.

pulicosum Hk. sp. **3**. 262. 1860 = Diplazium.
Pullingeri Bak. Gard. Chr. n. s. **4**. 484. 1875 = Diplazium.
Pululahuae Sod. Rec. 33. 1883; Cr. vasc. quit. 154. 1893 — Ecuador.
pumilum Sw. Prod. 129. 1788. Mett. n. 92. HB. 212 — Ind. occ. Mexico-Columbia. Africa orient. trop. India bor. Tonkin.
Tarachia Pr. 1849. Asplenium anthriscifolium Lam. 1788. A. humile Spr. 1822. A. mininum Mart. et Gal. 1842 — A. Schimperianum Hochst.: Fée 1850-52. A. hymenophylloides Fée 1857. A. tenerrimum Hochst. (Afr.).
pumilum Dav. Bull. Torr. Cl. **15**. 226. 1888 = A. chihuahuense.
punctatum Mett.; Kuhn, Fil. Afr. 114. 1868 = A. vagans.
puncticaule Mett. Ann. Lugd. Bat. **2**. 240. 1866 = Athyrium macrocarpum.
purdieanoides Karst. Fl. Col. **1**. 65. t. 32. 1859 = A. Purdieanum.
Purdieanum Hk. Ic. pl. t. 938. 1854. Mett. n. 150. HB. 245 — Venezuela-Peru.
Hemidictyum Moore 1857; Asplenidictyum J. Sm. 1875.
purpurascens Mett.; Kuhn, Linn. **36**. 102. 1869. HB. 486 — Ecuador.
pusillum Bl. Enum. 183. 1828 — Java.
Tarachia Pr. 1849. Asplenium confluens Kze. 1848. ? HB. 490.
pycnocarpon Spr. Anleit. **3**. 112. 1804 = Athyrium angustifolium.
pycnophyllum Moore, Ind. 159. 1859 — Mexico.
A. coriaceum Fée, Gen. 193. 1850-52; 7 mém. 46 t. 15 f. 1. (non Desv. 1827).
pygmaeum Hk. sp. **3**. 147 t. 180 B. 1860 = A. affine stat. juv.
pygmaeum Boj. Hort. Maur. 395. 1837 = ? A. stoloniferum.
pygmeum L. Syst. Nat. ed. X. **2**. 1323. 1759 — Jamaica.
(f. Moore = A. ruta muraria).
pyramidatum Desv. Prod. 271. 1827 — ? Tristan d'Acunha.
pyramidatum Liebm. Vid. Selsk. Skr. V. **1**. 249. (seors. 97). 1849 = A. sulcatum.
quitense Hk. 2 Cent. t. 20. 1860. HB. 196 — Ecuador.
rachirhizon Raddi, Pl. bras. **1**. 39 t. 56. 1825 = A. uniseriale.
Raddianum Gaud. Freyc. Voy. Bot. 316. 1827 = A. lunulatum var.
Raddii Fée, Gen. 190. 1850-52 = A. serratum.
radiatum Sw. Schrad. Journ. **1800**[2]. 50. 1801 = Actiniopteris australis.
radicans L. Syst. Nat. ed. X. **2**. 1323. 1759 — Amer. trop.
A. rhizophyllum L. 1763 (non 1753); A. rhizophorum L. 1764. HB. 204 (pt.). Chr. 196 (pt.). NPfl. 236 — A. alloeopteron Kze. 1847 — A. cyrtopteron Kze. 1850; A. flabellulatum a Mett. n. 99. 1859 — (teste G. Hieronymus in litt.).
radicans Sw. Syn. 84. 1806 = A. partitum var.
radicans Schkuhr, Kr. Gew. **1**. 70 t. 76. 1809 = Diplazium sp. dub. (arboreum ?).
radicans Pritch. List of St. Helena Pl. 6. 1836 = A. lunulatum var. tenellum.
radicans Bak. Syn. 241. 1867 = Diplaziuum ambiguum et sp. plur.
ramosum Spr.: Bernh. Schrad. Journ. **1801**[1]. 2. 1802 = Diplazium sp. dub.
ramosum Poir. Enc. Suppl. **2**. 515. 1811 = Didymochlaena truncatula.
Raouli Mett. Aspl. 118 n. 73 b. 1859 = A. Hookerianum.
Rawsoni Bak. JoB. **1872**. 362. HB. 487 — Africa austr.
reclinatum Houlst. Gard. Mag. Bot. **3**. 260. 1851 = A. lunulatum var. tenellum.
recognitum Kze. Linn. **22**. 577. 1849 = A. sulcatum.
recurvatum Don, Prod. Fl. Nepal. 7. 1825 — Nepal.
reflexum Bory, Bél. Voy. Bot. **2**. 48. 1833 = A. lunulatum.
reflexum Sod. Cr. vasc. quit. 205. 1893 = Diplazium angulosum.
refractum Moore, Ferns Gr. Br. Nat. Print. t. 35 A. 1857 = A. fontanum var.

regulare Sw. Vet. Ak. Handl. **1817**. 67; Lindm. Ark. för Bot. **1**. 219 t. 10 f. 5. 1903 — Brasilia.

Reicheliae Dörfl. et Aschers. Verh. bot. Ver. Brand. **37**. XLVII. 1896 = A. ruta muraria × trichomanes.

remotum Moore, Ind. 160. 1859 — Polynesia. N. Guinea. Celebes.

A. distans Brack. Expl. Exp. **16**. 155. 1854 (non Don 1825 nec Fée 1850-52); A. multilineatum Hk. sp. **3**. 102 t. 183. 1860. HB. 190. Chr. 200. NPfl. 236.

repandulum Kze. Linn. **9**. 65. 1834 = A. obtusifolium.

repandum Mett.; Kuhn, Fil. Afr. 114. 1868. HB. 482 — Guinea.

repens Hk. 2 Cent. t. 31. 1860; HB. 219 — Ecuador.

repens Döll, Mannh. Jahresber. 1861 —. = Athyrium Schimperi.

repente Desv. Prod. 271. 1827 — Réunion.

resectum Sm. Icon. ined. **3**. t. 72. 1791 = A. unilaterale.

resiliens Kze. Linn. **18**. 331. 1844 — Mexico. U. S. A. merid. Jamaica.

A. parvulum Mart. et Gal. Mém. Ac. Brux. **15**. 61 t. 13 f. 3. 1842. NPfl. 236 (non Hk. 1840); A. trichomanoides Kze. 1848. Mett. n. 113 (non alior.).

reticulatum Wall. List n. 188. 1828 = Diplaziopsis javanica.

reticulatum Roxb. Calc. Journ. **4**. 497. 1844 = Polypodium scolopendrinum.

retortum Klf. Enum. 171. 1824 = A. mucronatum.

retusum Poir. Enc. Suppl. **2**. 503. 1811 — Réunion.

Reuteri Milde, Bot. Zeit. **1867**. 148.; Fil. Eur. 62. 1867 — Cilicia.

rhaeticum Brügger, Naturg. Beitr. Chur 47. 1874 = Athyrium alpestre.

rhizophorum L. Gen. pl. ed. VI. **2**. emendat. 1764 = A. radicans.

rhizophorum Schkuhr, Kr. Gew. **1**. t. 76. 1809 = Diplazium sp. dub. (arboreum?).

rhizophyllum L. sp. ed. I. **2**. 1078. 1753; ed. II. **2**. 1536 = Camptosorus.

rhizophyllum L. sp. ed. II. **2**. 1540. 1763 = A. radicans.

rhizophyllum Kze. Linn. **9**. 71. 1834 = A. myriophyllum.

rhoifolium Mett. Aspl. 178. n. 199. 1859 = Diplazium.

rhomboidale Desv. Prod. 272. 1827. (Plum. t. 65) — St. Domingo.

rhomboideum Brack. Expl. Exp. **16**. 156 t. 21 f. 2. 1854 = A. fragile.

Richardi Hk.fil. Fl. N. Zeal. **2**. 35. 1854. HB. 222 — N. Zealand.

Riedelianum Bong.; Kuhn, Linn. **36**. 102. 1869 = Diplazium.

rigidum Sw. Vet. Ak. Handl. **1817**. 68 = A. sulcatum.

riparium Liebm. Vid. Selsk. Skr. V. **1**. 244 (seors. 92). 1849 = A. obtusifolium.

riparium Brack. Expl. Exp. **16**. 162. 1854 = A. laserpitiifolium.

rivale Spruce; Bak. Syn. 244. 1867 = Diplazium.

Robinsonii F. Muell. JoB. **1884**. 289 — Ins. Norfolk.

robustum Bl. Enum. 189. 1828 = A. laserpitiifolium.

Roemerianum Kze. Linn. **9**. 62. 1834 = Diplazium.

rotundatum Klf.; Holl, Flora **1830**. 374 = A. lanceolatum.

rotundifolium (Pr.) Ettingsh. Denkschr. Ak. Wien **23**. 73. 1864 — N. Zealand.

rubinum Dav. Bot. Gaz. **19**. 391. 1894 — Mexico. [Tarachia Pr. hb.

Ruizianum Kl. Linn. **20**. 354. 1847 = A. abscissum.

rupicola Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 531 t. 5. 1899 = Athyrium.

rusciforme »Wall.«; Salom. Nom. 101. 1883 — Himalaya.

Ruta Wall. List n. 233. 1828 = A. varians.

rutaceum (Willd.) Mett. Aspl. 129 n. 98. 1859. HB. 220. NPfl. 241 — Venezuela-Ecuador. Galapagos. ? Jamaica.

Aspidium Willd. sp. **5**. 266. 1810; Athyrium Pr. 1836.

rutaefolium Kze. Linn. **10**. 521. 1836 = A. achilleifolium.

ruta muraria L. sp. 2. 1081. 1753. Mett. n. 126. HB. 213. Chr. 202. NPfl. 240 — Europa. Asia septentr.-Himalaya. U. S. A. orient.
Acrostichum Lam.; Phyllitis Moench 1794; Scolopendrium Roth 1800; Amesium Newm. 1844; Tarachia Pr. 1849.
ruta muraria Pappe et Raws. Syn. Fil. Afr. austr. 20. 1858 = A. Rawsoni.
ruta muraria × septentrionale Murbeck, Bot. Centralbl. **31**. 322. 1887 (nomen); Lunds Univ. Årsskr. **27**[6]. 36 t. 2. 1892 — Suecia (Upsala).
A. Murbeckii Dörfl. 1895; A. septentrionale × ruta muraria Aschers. 1896.
ruta muraria × trichomanes C. Chr. Ind. 130. 1905 — Europa centr.
A. Preismanni Aschers. et Luerss. Verh. bot. Ver. Brand. **37**. XLVI. 1896; A. trichomanes × ruta muraria Aschers. l. c. — A. Reicheliae Dörfl. et Aschers. l. c.
ruwenzoriense Bak. Kew Bull. **1901**. 137 — Uganda.
salicifolium L. sp. 2. 1080. 1753. Mett. n. 40 (excl. varr.). HB. 200. Chr. 200. NPfl. 236 — Amer. trop.
salicinum J. Sm. JoB. **3**. 408. 1841 = A. persicifolium.
salignum Bl. Enum. 175. 1828. (? HB. 199 pt.). ? NPfl. 236 — Java.
salignum Mett. Fil. Lips. 72 t. 7. 1856 = ? A. Zenkerianum vel A. Carruthersii.
salignum Hk. Bak. Syn. 199. 1867 (pt.). ? NPfl. 236 = A. diversifolium.
salignum Carr. in Seem. Fl. Vit. 354. 1873 = A. Carruthersii.
sambucifolium Ettingsh. Farnkr. 145 t. 95 f. 7, 9. 1865 = Diplazium rhoifolium.
Sammatii Kuhn, v. Deck. Reis. **3**[3]. Bot. 34. 1879 = Diplazium.
Sampsoni Hance, Ann. sc. nat. V. **5**. 257. 1866. (HB. 223) — China.
Sancti-Christofori Christ in Schum. et. Laut. Fl. deut. Südsee 128. 1901 — N. Guinea. Ins. Salomonis.
Sanderi C. Chr. Ind. 130. 1905 — Columbia.
A. macrodictyon Bak. Kew Bull. **1901**. 144 (non Bak. 1877); Arcasplenium grande Moore msc.
Sandersoni Hk. sp. **3**. 147 t. 179. 1860. HB. 196. NPfl. 235 — Africa orient. trop. et austr. cum ins. — — A. Hanningtoni Bak. 1883. (conf. A. Dregeanum).
sandwichense (Pr.) Milde, Bot. Zeit. **1870**. 353; Hill. Fl. Haw. 610. 1888 — Ins. Hawaii. — Diplazium Pr. 1836 (nomen); Epim. 85. 1849.
Sandwichianum Hk. sp. **3**. 225. 1860 = Diplazium.
sanguinolentum Kze.; Mett. Aspl. 98 n. 35 t. 4 f. 10. 1859 = A. anisophyllum*.
Sarelii Hk. in Blakiston, Yang-tsze 363, 364. 1862 — Japonia, China. Himalaya.
A. Saulii Bak. 1874. HB. ed. II. 216. Chr. 203. NPfl. 240 (err. t. A. Gepp in litt.); A. Blakistoni Bak. 1867. HB. ed. I. 216 — A. Pekinense Hance 1867. HB. ed. II. 213; A. sepulchrale Hk. 1867. HB. ed. I. 213.
sarmentosum Willd. sp. **5**. 316. 1810 = A. obtusatum.
Saulii Bak. Syn. ed. II. 216. 1874 = A. Sarelii.
Saulii var. latius Christ, Nu. Giorn. bot. It. n. s. **4**. 89. 1897 = A. woodsioides.
saxatile Salisb. Prod. 403. 1796 = A. trichomanes.
scandens J. Sm. JoB. **3**. 408. 1841. (nomen); Mett. Aspl. 108 n. 57. 1859. HB. 224. Chr. 207 — Ins. Philipp. N. Guinea. Fiji. — Darea Fée 1850-52.
scandens Houlst. et Moore Gard. Mag. Bot. **3**. 260. 1851 = A. Veitchianum.
scandicinum Klf. Enum. 177. 1824. Mett. n. 71. HB. 217 — Jamaica. Brasilia.
A. adiantoides Raddi 1819 (non Lam. 1786).
scandicinum Hk. sp. **3**. 183 t. 204. 1860 = A. divergens.
scandicinum Heller, Minn. bot. St. **1**. 775. 1897 = Athyrium.
Schaffneri Fourn. Mex. pl. **1**. 110. 1872 = Diplazium Feei.
Schiedei Mett. Aspl. 165. n. 174. 1859 = Diplazium denticulosum.

Schimperi A. Br. in Schweinf. Beitr. **1**. 224. 1867 = Athyrium.
Schimperi J. Sm. Hist. Fil. 320. 1875 = A. theciferum *concinnum.
Schimperianum Hochst.; Fée, Gen. 191. 1850-52 = A. pumilum var.
schizodon Moore, Gard. Chr. **1871**. 1004 f. 223 = A. Vieillardii var.
schizophyllum C. Chr. Ind. 131. 1905 — Ins. Hawaii. — A. dissectum Brack. Expl. Exp. **16**. 170 t. 24. 1854. HB. 214. NPfl. 240 (non Sw. 1788).
Schkuhrianum Pr. Tent. 107. 1836 = A. laetum.
Schkuhrii Mett. Aspl. 182. n. 208. 1859 = Diplazium silvaticum.
Schkuhrii Hk. sp. **3**. 251. 1860; Mett. Ann. Lugd. Bat. **2**. 238. 1866 = Diplazium japonicum.
Schkuhrii Thwait. Enum. pl. Zeyl. 385. 1864; Bak. Syn. 491. 1874 = Diplazium [Beddomei.
Schomburgkianum Kl. Linn. **20**. 350. 1847 = A. serratum.
Schottii Pr. Del. Prag. **1**. 179. 1822 = A. sulcatum.
Schwackei Christ in Schwacke, Pl. Nov. Mineiras **2**. 28. 1900; Bull. Boiss. II. **2**. 556. 1902 — Brasilia.
Schweinfurthii Bak. in Balfour, Bot. of Socotra 328 t. 100. 1888. (Tr. Roy. Soc. Edinb. **31**). NPfl. 238 — Socotra. Africa orient.
scleropium Homb. et Jacq. Voy. au Pôle sud. t. 1 f. D. 1852 = A. obtusatum var.
scolopendrioides J. Sm. JoB. **3**. 408. 1841 (nomen); Hk. Ic. pl. t. 930. 1854. Mett. n. 11. HB. 192. NPfl. 235 — Ins. Philipp. — Micropodium J. Sm. 1875.
Scolopendrium L. sp. **2**. 1079. 1753 = Phyllitis.
Scolopendrium Lour. Fl. Cochinch. 832. 1790 = A. nidus.
scolopendropsis F. Muell. Pap. Plants **3**. 49. 1876 — N. Guinea. (an Triphlebia sp.?).
Scortechinii Bedd. JoB. **1887**. 322 — Perak.
sechellarum Bak. Syn. 491. 1874 = Diplazium.
Seelosii Leybold, Flora **1855**. 81, 348 t. 15. Mett. n. 123. HB. 198. Chr. 201. NPfl. 235 — Europa centr. — Acropteris Heufler 1856.
Seemannii Bak. Syn. 231. 1867 = Diplazium.
Seemannii Carr. in Seem. Fl. Vit. 355. 1873 = A. bipinnatifidum.
Selenopteris Mett. Aspl. 106. n. 238. 1859 = Athyrium solenopteris.
Sellowianum Pr. Tent. 107. 1836 (nomen) — Brasilia.
semiadnatum C. Chr. Ind. 131. 1905 — Samoa.
A. decurrens Bak. Syn. 484. 1874 (non Willd. 1810).
semicordatum Raddi, Pl. Bras. **1**. 36 t. 52 f. 1. 1825. Mett. n. 44 — Amer. trop. A. auriculatum Sw. Vet. Ak. Hdl. **1817**. 68; HB. 203. Chr. 195. NPfl. 237 (non supra); A. brasiliense Desv. 1827 — A. discolor Kze. 1834 — A. falx Desv. 1827. Mett. n. 41 — A. pimpinellifolium Schaffn.; Fée 1857.
semidentatum Fée, 10 mém. 27 t. 38 f. 1. 1865 — Réunion.
semihastatum Kze.; Mett. Aspl. 162. n. 168. 1859 = Diplazium.
septentrionale (L.) Hoffm. Deutschl. Fl. **2**. 12. 1795. Mett. n. 122. HB. 198. Chr. 201. NPfl. 235 — Europa. Asia bor. Himalaya. U. S. A. occ. (Rocky Mts.). Acrostichum L. sp. **2**. 1068. 1753: Scolopendrium Roth 1800; Belvisia Mirb. 1802; Blechnum Wallr. 1831; Acropteris Link 1833; Amesium Newm. 1844.
septentrionale × ruta muraria Aschers. Syn. Mitteleur. Fl. **1**. 75. 1896 = A. ruta muraria × septentrionale.
septentrionale × trichomanes C. Chr. Ind. 131. 1905 — Europa centr. A. Baumgartneri Dörfler, Oest Bot. Zeit. **45**. 169 t. 9. 1895 — A. Hansii Aschers. 1896.
sepulchrale Hk. Syn. ed. I. 213. 1867 = A. Sarelii var. pekinense.

serpentini Tausch, Flora **1839**². 477 = A. adiantum nigrum *cuneiforme.
serra Langsd. et Fisch. Ic. Fil. 16 t. 19. 1810. Mett. n. 143. HB. 206, 485. Chr. 199. NPfl. 234 — Amer. trop. Africa occ. trop. Natal.
— A. woodwardioideum Gardn. 1842 — A. camptosorum Mett.; Kuhn 1869.
serraeforme Mett. Aspl. 119 n. 75. t. 4 f. 1. 1859 = A. obscurum.
serratum L. sp. **2**. 1079. 1753. Mett. n. 9. HB. 193. Chr. 189. NPfl. 234 — Amer. trop. Congo. — A. Nidus Raddi 1819; A. Raddii Fée 1850-52. A. crenulatum Pr. 1836.
Serricula Fée, Gen. 196. 1850-52 = A. Wightianum.
Serronii Glaz.; Fée, Cr. vasc. Br. **1**. 68 t. 17 f. 2. 1869 = A. pulchellum.
serrulatum Cav. Anal. Cienc. **4**. 105. 1801 = A. adiantum nigrum var.
serrulatum Sw. Fl. Ind. occ. **3**. 1607. 1806 = Polypodium.
serrulatum Pr. Rel. Haenk. **1**. 46. 1825 = Diplazium manilense.
serrulatum Roxb. Calc. Journ. **4**. 498. 1844 = Diplazium.
sertularioides Bak. Tr. Linn. Soc. II. Bot. **2**. 354. 1887 — Kilima-njaro.
sessilifolium Desv. Berl. Mag. **5**. 322. 1811 — Guatemala-Peru.
— A. guatemalense et columbiense Hier. 1904. (conf. Hier. Engl. Jahrb. **34**. 460).
setisectum Bl. Enum. 187. 1828 — Java. — Tarachia Pr. 1849.
setosum Desv. Berl. Mag. **5**. 322. 1811 = ? A. lunulatum.
?setosum Pr. Rel. Haenk. **1**. 42. 1825 = Diplazium.
setulosum J. Sm. Cat. Kew Ferns 5. 1856 = ? Athyrium setiferum.
Shepherdi Mett. Aspl. 164 n. 171. 1859 = Diplazium arboreum.
Shepherdii Spr. Nova Acta **10**. 231 t. 17 f. 5-6. 1821 = Diplazium.
shepherdioides Bak. Ann. Bot. **8**. 126. 1894 = Diplazium.
sherburgense Bak. Ann. Bot. **5**. 304. 1891 — Ins. Sherburg (Guinea).
shikokianum Mak. Bot. Mag. Tokyo —. 1892 — Japonia.
Shuttleworthianum Kze. Farnkr. **1**. 26 t. 14. 1840. Mett. n. 59 — Polynesia.
— A. multifidum Brack. 1854. HB. 224. Chr. 207. NPfl. 242. — A. Powellii Bak. 1867. HB. 224. — (conf. Luerss. Farne d. Samoa 376).
sibiricum Turcz. Flora **1834**¹. Beibl. 28 (nomen); Kze. Anal. 25 t. 15. 1837 = Athyrium crenatum.
sikkimense Clarke, Tr. Linn. Soc. II. Bot. **1**. 500 t. 65 f. 1. 1880 = Diplazium asperum.
silesiacum Milde, 33 Jahrb. Schles. Ges. 93. 1855 = A. adiantum nigrum *cuneiforme.
silvaticum Pr. Rel. Haenk. **1**. 42. 1825 = Diplazium.
silvaticum Moore, Ind. 43. 1857 = Athyrium.
simile Bl. Enum. 181. 1828 = A. macrophyllum.
simillimum Kuhn, Hill. Pl. Madag. exs. n. 3773 = A. affine var.
Simonsianum Hk. Ic. pl. t. 925. 1854. HB. 191 — India bor.
Thamnopteris Moore 1857; Neottopteris J. Sm. 1875.
simplex Bl. Enum. 174. 1828 = A. amboinense.
simplex Mett. Aspl. 86. n. 5. 1859 = A. nidus.
simplicifrons F. Muell. Fragm. **5**. 74. 1865. HB. 193. (Chr. 190). NPfl. 235 — Queensland.
sinuatum Salisb. Prod. 403. 1796 = Ceterach officinarum.
sinuatum Pal. Beauv. Fl. d'Oware **2**. 33 t. 79. post 1811 = A. africanum.
Skinneri Bak. Syn. 226. 1867 = Athyrium.
Smithianum Bak. Syn. 245. 1867 = Diplazium.
soboliferum Wall. List n. 201. 1828 = Diplazium tomentosum.
Sodiroi Christ, Prim. Fl. Costar. **3**. 26. 1901 — Ecuador. Costa Rica.
A. ebeneum Sod. Cr. vasc. quit. 146. 1893 (non Ait. 1789).
solenopteris vide selenopteris.

solidum Kze. Linn. **10**. 520. 1836. Mett. n. 127. HB. 214. (Chr. 202). — Afr. austr.
Tarachia Pr. 1849.
Solmsii Bak.; Hemsl. Biol. Centr. Amer. **3**. 639. 1885. (Chr. 206). NPfl. 241 (Solmsianum) — Amer. centr.
sorbifolium Jacq. Coll. **2**. 106 t. 3 f. 2. 1788 = Dryopteris reticulata*.
sorbifolium Willd. sp. **5**. 312. 1810 = Diplazium.
sordidum Kze. Bot. Zeit. **1848**. 174 = A. longissimum.
sororium Mett. Ann. sc. nat. IV. **15**. 73. 1861 = Diplazium maximum.
sorzogonense Pr. Rel. Haenk. **1**. 45. 1825 = Diplazium.
Sorzogonense β majus Hk. sp. **3**. 252. 1860 = Diplazium crinitum.
spathulatum Bak. Ann. Bot. **8**. 124. 1894 — Sumatra.
spathulinum J. Sm. JoB. **3**. 408. 1841; Bedd. Ferns S. Ind. t. 226. 1863 = A. affine.
spathulinum (Hk. sp. **3**. 170. 1860 pt.) Hill. Fl. Haw. 604. 1888 = ? A. caudatum.
speciosum Mett. Aspl. 185 n. 213. t. 5. f. 5. 1859 = Diplazium sorzogonense.
spectabile Wall. List n. 237. 1828 = Athyrium.
speluncae Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 113 c. fig. — China.
sphenoides Kze. Linn. **9**. 63. 1834 = A. obtusatum* var.
sphenotomum Hill. Fl. Haw. 599. 1888 — Ins. Hawaii.
Spicant Bernh. Schrad. Journ. **1799**[1]. 309 = Blechnum.
spinescens Mett.; Kuhn, Linn. **36**. 98. 1869. HB. 488 — Bolivia.
spinulosum Mett. Aspl. 172. n. 189. 1859 = Diplazium proliferum.
spinulosum Mett. Ann. Lugd. Bat. **2**. 237. 1866 = Diplazium.
spinulosum Miq. Ann. Lugd. Bat. **3**. 174. 1867; Bak. Syn. 225. 1867 = Athyrium.
splendens Kze. Linn. **10**. 516. 1836. Mett. n. 160. HB. 487 — Africa austr.
Tarachia Pr. 1849.
Sprengelii Wikstr. Vet. Ak. Hdl. **1825**. 443. 1826. —? Ind. occ.
A. ambiguum Spr. Fl. Halensis, Mant. prima 54. 1807.
Sprucei Bak. Syn. 234. 1867 = Diplazium.
squamigerum Mett. Ann. Lugd. Bat. **2**. 239. 1866 = Diplazium.
squamosum L. sp. **2**. 1082. 1753. HB. 218. Chr. 205. NPfl. 241 — Amer. trop.
squamulatum Bl. Enum. 174. Mett. n. 17. HB. 192. NPfl. 233 — Malesia.
Thamnopteris Pr. 1849; Neottopteris Fée 1850-52.
squamulosum Moore, Ind. 169. 1860 = Athyrium costale.
stans Kze. Linn. **10**. 521. 1836 = A. achilleifolium.
Steerei Harr. Journ. Linn. Soc. **16**. 28. 1877 — Luzon.
stellatum Colla, Mem. Ac. Torino **39**. 41 t. 69. 1836 = A. lunulatum* var.
stenocarpon Fée, Cr. vasc. Br. **1**. 64 t. 19 f. 1. 1869 = A. anisophyllum*.
stenocarpum Mett. Ann. sc. nat. V. **2**. 232. 1864 = Diplazium.
stenolobum C. Chr. Ind. 133. 1905 — Fiji.
Davallia *foeniculacea* Hk. 2 Cent. t. 54. 1861. HB. 101. Chr. 303 (non Asplenium HBK. 1815); Loxoscaphe Moore 1861.
stenophyllum Bedd. Ferns br. Ind. t. 147. 1866 = A. ensiforme.
stenopteris Kze. Bot. Zeit. **1848**. 174 = A. pellucidum.
stereophyllum Kze. Bot. Zeit. **1848**. 175. Mett. n. 159 — Java. — Tarachia Pr. 1849.
stipitatum J. Sm. Ferns br. and for. 210. 1866 = A. squamulatum.
Stoliczkae Clarke, Tr. Linn. Soc. II. Bot. **1**. 500. 1880 = Diplazium.
stoloniferum Bory, Voy. **1**. 329. 1804 — Ins. Mascarenae.
stoloniferum Pr. Rel. Haenk. **1**. 44 t. 6 f. 4. 1825 = A. fragile.
striatum L. sp. **2**. 1082. 1753 = Diplazium.

striatum Mett. Fil. Lips. 78. 1856 = Diplazium expansum.
striatum Hk. sp. **3**. 245. 1860 = Diplazium Shepherdii.
strictum Brack. Expl. Exp. **16**. 168 t. 23 f. 1. 1854 = A. lunulatum var. [Macraei.
strigillosum Lowe, Ferns **5** t. 36. 1858 = Athyrium nigripes.
Stuhlmannii Hier. in Engler, Pflanzw. Ostafr. C. 83. 1895 — Afr. orient. trop.
subalatum Hk. et Arn. Bot. Beech. Voy. 312. t. 71. 1840 = A. formosum.
subaquatile Ces. Atti Ac. Napoli 7^{8}. 20 t. 3 f. 5. 1876 — Borneo.
subavenium Hk. sp. **3**. 143 t. 181 B. 1860. HB. 198 — Madagascar.
Obs. Pl. Penangiana est A. pellucidum (f. Bedd.).
subdigitatum Mett.; Kuhn, Fil. Afr. 117. 1868 — Mauritius.
Darea palmata Klf. Enum. 181. 1824 (non Asplenium Lam. 1786); Caenopteris Spr. 1827.
subhastatum Hk. Ic. pl. t. 929. 1854. Mett. n. 18. HB. 193 — Caracas. Peru.
subintegrum C. Chr. Ind. 134. 1905 — Kamerun.
A. coriaceum Bak. Syn. 192. 1867 (non Desv. 1827 nec alior.).
subnudum Karst. Fl. Col. **2**. 93 t. 148. (1865) = Diplazium Lindbergii.
subnudum Bak. Syn. 483. 1874 = A. oligolepidum.
subserratum Bl. Enum. 174. 1828 = Diplazium.
subserratum Mett. Aspl. 90. n. 14. 1859 = A. vittaeforme.
subsessile Cav. Descr. 254. 1802 — Ins. Mariannae.
subsinuatum Wall.; Hk. et Grev. Ic. Fil. t. 27. 1827 = Diplazium lanceum.
subtriangulare Hk. Syn. 225. 1867 = Athyrium spinulosum var.
succulentum Clarke, Tr. Linn. Soc. II. Bot. **1**. 502 t. 64 f. 4. 1880 = Diplazium.
sulcatum Lam. Enc. **2**. 308. 1786 — Amer. trop. Madagascar. Réunion. India austr. — A. rigidum Sw. 1817. A. recognitum Kze. 1849 — A. auritum Sw. 1801. Mett. n. 45. HB. 208. Chr. 203 — A. bidentatum Willd. 1810. A. macilentum Kl.; Kze. 1847 — A. umbrosum Klf. 1824 — A. dispersum Kze. 1850. Mett. n. 46.
sulcatum Mett. Ann. Lugd. Bat. **2**. 236. 1866 = A. setisectum.
sumatranum Hk. sp. **3**. 110 t. 168. 1860. HB. 199 — Sumatra. ? Fiji.
sundense Bl. Enum. 175. 1828 = A. vittaeforme.
Surinamense Fée, Gen. 192. 1850-52 = A. angustum.
Swartzianum Kze. Linn. **19**. 404. 1847 = Onychium japonicum.
Swartzii Mett. Fil. Lips. 74 t. 11 f. 6. 1856 = Diplazium proliferum.
sylvaticum vide silvaticum.
symmetricum Col. Tr. N. Zeal. Inst. **31**. 264. 1899 — N. Zealand.
tabulare Schrad. Gött. gel. Anz. **1818**. 916 = A. adiantum nigrum.
taeniosum Kze. Bot. Zeit. **1848**. 145 = A. amboinense.
Tavoyanum Wall. List n. 1035. 1828 = A. adiantoides.
Taylorianum Jenm. JoB. **1886**. 269 = Diplazium altissimum var.
tenebrosum Jenm. Gard. Chr. III. **16**. 690. 1894 = Diplazium.
tenellum Roxb. in Beats. St. Helena 299. 1816 = A. lunulatum var.
tenellum Fée, Gen. 198. 1850-52 = A. fragrans *foeniculaceum.
tenellum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 259. t. 4. 1899 = Athyrium setiferum.
tenerrimum Hochst.; Schimp. pl. abyss. 2064 (nomen) = A. pumilum.
tenerrimum Mett.; Kuhn, Linn. **36**. 97. 1869. HB. 488 — Venezuela.
tenerum Forst. Prod. 80. 1786. Mett. n. 64. HB. 201. Chr. 194. NPfl. 237 — Ind. austr. Ceylon. Malesia-Polynesia. Ins. Sechellae. *Asia trop.

Darea Spr. 1799 — A. caudatum Cav. 1802 (non Forst. 1786); A. elongatum Sw. 1806. Mett. n. 63. A. productum Pr. 1825.

— *Darea Belangeri Bory 1833; Asplenium Kze. 1848. Mett. n. 62. HB. 223. Chr. 208. NPfl. 242 (non Bory 1833); A. Veitchianum Moore 1857. Darea furcata Bl. 1828 — D. appendiculata Bl. 1828; Asplenium decorum Kze. 1848.

tenerum Raddi, Opusc. sci. Bol. **3**. 290. 1819 = A. lunulatum.

tenerum Gaud. Freyc. Voy. Bot. 315. 1827 = ? A. semicordatum.

tenue Pr. Rel. Haenk. **1**. 44 t. 6 f. 5. 1825 — Peru. Quito. (poss. = A. fragile var.).

tenuifolium Don, Prod. Fl. Nepal. 8. 1825. Mett. n. 95. HB. 220. Chr. 205. NPfl. 241 — India.

tenuifolium Guss. Pl. rar. 377 t. 65. 1826 = A. fissum.

tenuifrons Wall. List n. 206. 1828 = Athyrium nigripes.

tenuilobum Moore, Ind. 172. 1860 = A. fragrans *foeniculaceum.

tenuisectum Hk. sp. **3**. 222. 1860 = Cystopteris.

ternatum Pr. Rel. Haenk. **1**. 45. 1825; Fée 7 mém. 54 t. 16 f. 4. = A. triphyllum var.

ternatum Hk. sp. **3**. 265. 1860 = Diplazium.

Textori Miq. Cat. Mus. Bot. Lugd. Bat. 126. 1870 = Diplazium Mettenianum.

? *thalictroides* (Loud.) Kze. Linn. **23**. 238. 1850 — Jamaica (Hort.).

Caenopteris Loud. 1850. (= ? A. cristatum).

theciferum (HBK.) Mett. Ann. sc. nat. V. **2**. 227. 1864. Chr. 209. NPfl. 244 — Amer. austr. trop. *Afr. trop. et austr.

Davallia HBK. Nov. Gen. et Sp. **1**. 23. 1815. HB. 100; Loxoscaphe Moore 1853; Darea Fourn. 1872 — Davallia Lindeni Hk. 1846; Loxoscaphe Moore 1853; Microlepia Mett. 1856.

— *Davallia concinna Schrad. 1818; Loxoscaphe Moore 1853; Asplenum Kuhn 1868. Davallia campyloptera Kze. 1836 — D. Schimperi Hk. 1846; Loxoscaphe Moore 1853; Microlepia Mett. 1856; Asplenium J. Sm. 1875.

thelypteroides Michx. Fl. bor. Am. **2**. 265. 1803 = Athyrium acrostichoides.

Thunbergii Kze. Linn. **10**. 517. 1836 = A. auriculatum.

Thwaitesianum Szysz. ap. Beck, It. Princ. Coburg. **2**. —. 1888 = Diplazium polypodioides var.

Thwaitesii A. Br. Ind. sem. ht. Ber. 1857; Mett. Aspl. 183. n. 209 b. 1859 = Diplazium.

tomentosum Lam. Enc. **2**. 308. 1786 = Gymnopteris.

tomentosum Mett. Aspl. 182. n. 209. 1859 = Diplazium Petersenii.

tomentosum Hk. sp. **3**. 249. 1860 = Diplazium.

tonkinense C. Chr. Ind. 135. 1905 — Tonkin.

A. melanolepis Bak. JoB. **1890**. 264 (non Franch. et Sav. 1879).

Toramanum Mak. Bot. Mag. Tokyo —. 1892 — Japonia.

torrentium Clarke, Tr. Linn. Soc. II. Bot. **1**. 500. t. 64. f. 2-3. 1880 = Diplazium maximum.

Torresianum Gaud. Freyc. Voy. Bot. 317. 1827 = A. pellucidum.

trapeziforme Wall. List n. 2213. 1829; Roxb. Calc. Journ. **4**. 497. 1844 — India austr. — (an A. lunulatum var.?).

trapezoides Sw. Syn. 76. 1806 = A. trilobum.

travancoricum Bak. Ann. Bot. **5**. 310. 1891 = Diplazium.

tremulum Hombr. et Jacq. Voy. an Pôle sud t. 3 bis f. △ 1852 = A. Fabianum.

Tretteneriauum Jan, Flora **1835**[1]. 32 = A. fissum.

Trianae Mett. Ann. sc. nat. V. **2**. 233. 1864 = Diplazium.

trichomanes L. sp. **2**. 1080. 1753. Mett. n. 115. HB. 196. Chr. 192. NPfl. 235 — Reg. temp. bor. et austr. Montes trop.

A. trichomanoides Houtt. 1783: Trichomanes crenatum Gilib. 1792; Phyllitis rotundifolia Moench 1794 — Asplenium melanocaulon Willd. 1809 — A. anceps L. v. Buch 1825 — A. densum Brack. 1854 — A. melanolepis Col. 1888 (non Franch. et Sav. 1879).

Trichomanes Thbg. Fl. Jap. 334. 1784 = A. incisum.

trichomanes × adiantum nigrum Aschers. Syn. Mitteleur. Fl. **1**. 80. 1896 = A. adiantum nigrum × trichomanes.

trichomanes × Cystopteris fragilis vide Cystopteris Blindi.

Trichomanes dentatum L. sp. **2**. 1080. 1753 = A. dentatum.

trichomanes × germanicum = A. germanicum × trichomanes.

Trichomanes ramosum L. sp. **2**. 1082. 1753 = A. viride.

trichomanes × ruta muraria Aschers. Verh. Bot. Ver. Brandenb. **37**. XLVI. 1896 = A. ruta muraria × trichomanes.

trichomanes × septentrionale Aschers. Syn. Mitteleur. Fl. **1**. 75. 1896 = A. germanicum.

Trichomanes × viride Milde, Sporenpfl. 40. 1865 = A. adulterinum.

trichomanoides Houtt. Pfl. Syst. **13**[1]. 145. 1786 = A. trichomanes.

trichomanoides Lour. Fl. Cochinch. **2**. 833. 1790 — Cochinchina.

trichomanoides Lumn. Fl. Poson. 1020. 1791 = A. adiantum nigrum.

trichomanoides Michx. Fl. bor. Am. **2**. 265. 1803 = A. platyneuron.

trichomanoides Kze. Am. Journ. Sci. II. **6**. 85. 1848 = A. resiliens.

trigonopterum Kze. Bot. Zeit. **1848**. 524. Mett. n. 54 — Ins. Bonin-Sima.

trilobum Cav. Descr. 255. 1802. HB. 193. Chr. 190 — Chile. Brasilia austr. A. trapezoides Sw. 1806. Mett. n. 131; Tarachia Pr. 1849 — Asplenium parvulum Hk. 1840.

trilobum Fourn. Mex. pl. **1**. 110. 1872 = ? A. cristatum.

tripartitum Bl. Enum. 185. 1828 = A. praemorsum.

triphyllum Pr. Rel. Haenk. **1**. 45. 1825. Mett. n. 88. HB. 219. Chr. 193. NPfl. 241 — Columbia-Peru.
— A. ternatum Pr. 1825. Mett. n. 89 — A. imbricatum Hk. et Grev. 1829.

tripinnatum Roxb. Calc. Journ. **4**. 500. 1844 = A. laserpitiifolium.

triste Klf. Enum. 170. 1824 = A. lunulatum vel A. regulare.

triste Raoul, Ann. sc. nat. III. **2**. 115. 1844 = A. bulbiferum var.

truncatilobum Fée, Gen. 191. 1850-52 = A. caudatum.

truncatum Bl. Enum. 184. 1828 = A. caudatum.

truncatum »Don«; Pr. Tent. 107. 1836 = A. laciniatum var.

tucumanense Hier. Engl. Jahrb. **22**. 380. 1896. NPfl. 241 — Argentina.

Tussaci Moore, Ind. 175. 1860 = Athyrium costale var.

umbrosum Vill. Hist. pl. Dauph. **1**. 281. 1786 = A. viride.

umbrosum Schrad. Gött. gel. Anz. **1824**. 870; Klf. Enum. 168. 1824 = A. [sulcatum.

umbrosum J. Sm. JoB. **4**. 174. 1841 = Athyrium.

umbrosum Mett. Fil. Lips. 74. 1856 = Diplazium esculentum.

umbrosum var. cristovalense Bak. Journ. Linn. Soc. **19**. 294. 1882 = Athyrium assimile.

unilaterale Lam. Enc. **2**. 305. 1786 (α) — Japonia-Asia trop.-Polynesia. Afr. occ. Ins. afric. orient. — A. resectum Sm. 1791. Mett. n. 103. HB. 210. Chr. 194. NPfl. 237 — A. amoenum Pr. 1836. Mett. n. 101 — A. emarginato-dentatum Zenker; Kze. 1851. Mett. n. 102.

unilaterale β Lam. Enc. **2**. 305. 1786 = A. monanthes.

unilobum Poir. Enc. Suppl. **2**. 505. 1811 = Diplazium arboreum.

uniseriale Raddi, Opusc. sci. Bol. 3. 291. 1819 — Amer. trop.
A. rachirhizon Raddi, Pl. Bras. 1. 39. t 56. 1825; A. flabellulatum b. Mett. n. 99. 1859. A. rhizophorum pt. HB. 204 — A. amabile Liebm. 1849.
Urbani Christ, Engl. Jahrb. 24. 108. 1897 = Diplazium.
urophyllum Wall. List n. 192. 1828 = A. adiantoides vel A. macrophyllum.
uropteron Miq. Ann. Lugd. Bat. 3. 174. 1867 = Athyrium.
vacillans Kze. Bot. Zeit. 1848. 172 = Diplazium.
vagans Bak. Syn. 195. 1867. NPfl. 235 — St. Thomé (Guinea). Madagascar.
A. Melleri Mett. 1868 — A. punctatum Mett. 1868.
Vallisclausae Req. ap. Guérin, Descr. Vaucl. ed. II. 239. 1813 = A. glandulosum.
variabile Hk. sp. 3. 93 t. 185. 1860. HB. 194 — Fernando Po.
varians Wall.; Hk. et Grev. Ic. Fil. t. 172. 1830. HB. 216. Chr. 204. NPfl. 240 — Japonia. China-Ceylon. Afr. austr. Ins. Hawaii.
A. fimbriatum Kze. 1844. Mett. n. 121.
varium Roxb. Calc. Journ. 4. 498. 1844 = Diplazium Roxburghii.
varium Mett.; Salom. Nom. 106. 1833 = Diplazium.
vastum Mett. Ann. sc. nat. V. 2. 237. 1864 = Diplazium.
Veitchianum Moore, Ind. XLIX. 1857 = A. tenerum*.
venosum Hk. sp. 3. 83. 1860 = A. africanum.
venulosum Bak. Syn. 238. 1867 = Diplazium.
venustum Und. et Maxon, Bull. Torr. Cl. 29. 581. 1902 — Cuba.
verapax Donn. Smith, Bot. Gaz. 13. 77 t. 2. 1888 = Diplazium Riedelianum.
vespertinum Maxon, Bull. Torr. Cl. 27. 197. 1900 — California.
vestitum Hk. 2 Cent. t. 46. 1860 = Diplazium.
verecundum Chapm.; Fourn. Mex. pl. 1. 111. 1872 = ? A. myriophyllum.
vexans Und. ap. Heller, Minn. bot. St. 1. 776. 1897 — Ins. Hawaii.
Vidalii Franch. et Sav. En. pl. Jap. 2. 229. 624. 1879 — Japonia.
Vieillardii Mett. Ann. sc. nat. IV. 15. 72. 1861. HB. 200 — N. Caledonia. Fiji. — A. schizodon Moore 1871 — A. apicidens Moore 1881 — A. Baptistii Moore 1881.
?villosum Pr. Rel. Haenk. 1. 46. 1825 = Diplazium.
Vincentis Christ, Engl. Jahrb. 24. 109. 1897 = Diplazium.
Vinsoni Cord. Fl. Réun. 68. 1895 — Réunion.
violascens Moore, Ind. 177. 1860 = A. lineatum var.
Virchowii Kuhn in Hill. Pl. Madag. exsicc. n. 4168; Bak. Ann. Bot. 5. 264. 1891 = Diplazium.
virens Pr. Rel. Haenk. 1. 41 t. 6 f. 3. 1825. HB. 201 — Amer. trop. occ.
virens Desv. Prod. 273. 1827 = A. abscissum.
virescens Mett. Aspl. 191. n. 227. 1859 = Diplazium.
Virgilii Bory, Exp. de la Morée 3^2. 289. 1832 = A. adiantum nigrum.
viridans Lab. Sert. Austr. Cal. 2 t. 2. 1824 — N. Caledonia. Australia.
Caenopteris Spr. 1827; Darea Bory 1833.
viride Huds. Fl. Angl. 385. 1762. Mett. n. 117. HB. 195. Chr. 191. NPfl. 235 — Europa. Himalaya. Amer. bor.
A. Trichomanes ramosum L. sp. 2. 1082. 1753 (prob. nom. opt.).
viridifrons Mak. Bot. Mag. Tokyo —. 1892 = Athyrium.
virillae Christ, Bull. Boiss. II. 4. 1090. 1904 — Costa Rica. Columbia.
vitiense Bak. Syn. 245. 1867 = Diplazium esculentum var.
vittaeforme Cav. Descr. 255. 1802. Mett. n. 13. HB. 192. Chr. 190. NPfl. 234 — Malesia-Polynesia. — Diplazium Kze. 1848. Micropodium J. Sm. 1875; Eremopodium Trevis. 1877. Asplenium sundense Bl. 1828; Micropodium

Mett. 1866. ? Eremopodium Trevis. 1877. Asplenium amboinense Brack. 1854 (non Willd. 1810).

viviparioides Kuhn in Hill. Pl. Madag. exsicc. n. 4148. NPfl. 243 — Madagascar. (f. Bak. = A. affine*).

viviparum (L. fil.) Pr. Tent. 109 t. 3 f. 20. 1836. Mett. n. 56. HB. 223. NPfl. 243 — Ins. Mascarenae. — Acrostichum L. fil. Suppl. 444. 1781; Caenopteris Berg. 1786; Darea Willd. 1810. — conf. A. lineatum.

viviparum Bl. Enum. 176. 1828 = A. vulcanicum.

vogesiacum F. Schultz, Jahresber. d. Pollichia XXII-XXIV. 188. 1866 = A. viride.

Volkensii Hier. in Engl. Pflw. Ostafr. C. 83. 1895 — Afr. orient. trop.

vomeriforme Hk. sp. **3**. 109 t. 162. 1860. HB. 206 — Peru.
A. Mathewsii Moore 1859 (nomen).

vulcanicum Bl. Enum. 176. 1828. Mett. n. 25. HB. 201. Chr. 200. NPfl. 236 — Malesia. Ceylon. — A. Walkerae Hk. 1860 — A. caespitosum Bl. 1828 — A. heterodon Kze. 1850. Mett. n. 26 (vix Bl. 1828).

Wageneri Mett. Ind. Sem. ht. Berol. 1856; Aspl. 188. n. 220. 1859 = Diplazium.

Wagneri Mett.; Kuhn, Linn. **36**. 96. 1869. (HB. 483) — Quito.

Walkerae Hk. sp. **3**. 108 t. 163. 1860 = A. vulcanicum var.

Wallichianum Pr. Tent. 107. 1836 (nomen) — India.

Wallisii Bak. Kew Bull. **1901**. 145 — Columbia. — (an Diplazium?).

Warburgianum Christ in Warb. Monsunia **1**. 72. 1900 — Java.

Wardii Hk. sp. **3**. 189. 1860 = Athyrium.

Welwitschii Hk. Syn. 235. 1867; Mett.; Kuhn, Fil. Afr. 119. 1868 = Diplazium.

Wheeleri Bak. Ann. Bot. **5**. 309. 1891 = Diplazium.

Wichurae Mett. Ann. Lugd. Bat. **2**. 237. 1866 = Diplazium.

Wightianum Wall. List n. 2215. 1829; Hk. sp. **3**. 105 t. 167. 1860. HB. 199. NPfl. 236 — Ceylon. Madras.
A. coriaceum Bory 1833 (non Desv. 1827) — ? A. longipes Fée 1850-52.

Wildii Bail. Queensl. Bot. Bull. **4**. 20 t. 1, 2. 1891; Lithogr. t. 108. 1892 — Queensland.

Wilfordii Mett.; Kuhn, Linn. **36**. 94. 1869. HB. 487 — Japonia.

Willdenowii Pr. Tent. 290. 1836 —?

Willdenowii A. Br.; Mett. Ann. sc. nat. V. **2**. 236. 1864 = Diplazium expansum.

Willdenowii Jenm. Gard. Chr. III. **19**. 8. 1896 = ? A. rutaceum.

Wilsoni Bak. Syn. 242. 1867 = Diplazium.

woodsioides Christ, Bull. Soc. Bot. It. 1900. 261 — China.

woodwardioides Bernh. Schrad. Journ. **1801**[1]. 17. 1802 = Lomaria.

woodwardioides Roxb. Calc. Journ. **4**. 500. 1844 — Chittagong.

woodwardioides Bak. Syn. 229. 1867 = Athyrium silvaticum.

woodwardioideum Gardn. Lond. JoB. **1**. 547. 1842 = A. serra var.

Wrightii Eaton; Hk. sp. **3**. 113 t. 182. 1860. HB. 204 — Japonia.

wrightioides Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 238 — China.

xiphophyllum Bak. JoB. **1879**. 40 = Diplazium.

Yokoscense Franch. et Sav. Enum. pl. Jap. **2**. 225, 622. 1879 = Athyrium.

yunnanense Franch. Bull. Soc. Fr. **32**. 28. 1885. NPfl. 241 — China merid.

zamiaefolium Willd. sp. **5**. 325. 1810 = ? A. dimidiatum.

zamiaefolium Pr. Rel. Haenk. **1**. 43. 1825 = A. adiantoides.

zamioides Hk. sp. **3**. 114 t. 170. 1860 = A. paradoxum.

zanzibaricum Bak. Ann. Bot. **5**. 311. 1891 = Diplazium.

Zenkerianum Kze. Linn. **24**. 259. 1851. Mett. n. 34. HB. 484 — India austr. Ceylon.

Zeyheri Pappe et Raws. Syn. Fil. Afr. austr. 18. 1858 = A. lunulatum.
zeylanicum Hk. sp. **3**. 237. 1860 = Diplazium.
zeylanicum Ces. Atti Acad. Napoli 7^{8}. 21. 1876 = Diplazium porphyrorachis.
zoliense Kit.; Sadler, Fil. ver. Hung. 29. 1830 = A. ruta muraria.

ASTEROGLOSSUM J. Smith msc. (Moore, Ind. 179. 1860) = **Drymoglossum** (carnosum).

Atactosia Blume, Enum. pl. Javae 134. 1828 (*Polypodium §*) = Polypodium, Dipteris, Drynaria.

ATHYRIUM Roth, Röm. Mag. 2^{1}. 105. 1799; Tent. Fl. Germ. **3**. 58. 1800 — (Polypodiaceae Gen. **41**).

accedens Milde, Bot. Zeit. **1870**. 353 = Diplazium proliferum.
achilleifolium (Mart. et Gal.) Fée, Gen. 186. 1850-52. Chr. 225. NPfl. 224 — Mexico. Guatemala. — Caenopteris Mart. et Gal. Mém. Ac. Brux. **15**. 63 t. 16. 1842; Asplenium Liebm. 1849. HB. 227, 489 — A. grande Fée 1857 (non Sw. 1806); Athyrium Fourn. 1872.
acrostichoides (Sw.) Diels, Nat. Pfl. 1^{4}. 223. 1899 — Amer. bor. atlant. Amur-China-Himalaya. — Asplenium Sw. Schrad. Journ. **1800**^{2}. 54. 1801. A. thelypteroides Michx. 1803. HB. 226, 489; Athyrium Desv. 1827. Chr. 223; Diplazium Pr. 1836 — Athyrium allantodioides Bedd. 1867.
acrostichoideum Bory; Mérat, Fl. Paris. ed. IV. 1. 372. 1836 = A. filix femina.
affine Milde, Bot. Zeit. **1870**. 353 = Diplazium.
alatum Christ, Bull. Boiss. **6**. 963. 1898 — China merid.
allantodioides Bedd. Ferns br. Ind. t. 221. 1867 = A. acrostichoides.
alpestre (Hoppe) Rylands in Moore, Ferns Gr. Br. Nat. Pr. **1** t. 7. 1857. NPfl. 224 — Europa. Asia minor. ? Amer. bor. pacif.
Aspidium Hoppe, Neu. bot. Taschenb. **1805**. 216; Polypodium (Hoppe 1821) Spenner 1825. HB. 311; Pseudathyrium Newm. 1851; Phegopteris Mett. 1856; Asplenium Mett. n. 245. 1859; Polypodium rhaeticum pt. L. 1753; Aspidium Sw. 1801 (pt.); Phegopteris Pérard 1869; Asplenium Brügger 1874; Athyrium Dalla Torre 1882. Chr. 224 — Pseudathyrium flexile Newm. 1851; Polypodium Moore 1853; Phegopteris J. Sm. 1857; Athyrium? Moore 1860.
alpinum Spr. Anleit. **3**. 143. 1804 = Cystopteris fragilis *regia.
alternifolium Milde, Bot. Zeit. **1870**. 353 = Diplazium bantamense var.
ambiguum Milde, Bot. Zeit. **1870**. 353 = 1) Diplazium esculentum; 2) D. ambiguum.
ammifolium (Mett.) C. Chr. Ind. 139. 1905 — Fernando Po. Kamerun. Asplenum Mett.; Kuhn, Fil. Afr. 96. 1868.
amplum Milde, Bot. Zeit. **1870**. 350 = Diplazium striatum.
angustifolium (Michx.) Milde, Bot. Zeit. **1866**. 376. NPfl. 223 — Amer. bor. atlant. Asplenium Michx. Fl. bor. Am. **2**. 265. 1803. HB. 199. Chr. 200.
angustum Pr. Rel. Haenk. **1**. 39. 1825 = A. filix femina var.
anisopterum Christ, Bull. Boiss. **6**. 962. 1898 — China merid.
arborescens Milde, Bot. Zeit. **1870**. 353 = Diplazium.
arboreum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
arcuatum Liebm. Vid. Selsk. Skr. V. **1**. 278 (seors. 126). 1849 — Mexico. Asplenium Hk. 1860 (sp. **3**. 219 non 142).
Arnottii Milde, Bot. Zeit. **1870**. 371 = Diplazium.
asperum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
aspidioides (Willd.) Pr. Tent. 98. 1836 — India.
Darea Willd. Enum. 1072. 1809; sp. **5**. 301. 1810; Asplenium Spr. 1827; Caenopteris Desv. 1827. (conf. A. scandicinum).

aspidioides Christ. Farnkr. 224. 1897 = A. scandicinum.
asplenioides Desv. Prod. 266. 1827 = A. filix femina.
asplenioides Milde, Bot. Zeit. **1870**. 353 = Diplazium.
assimile (Endl.) Pr. Tent. 98. 1836 — Polynesia. Ceylon.
Asplenium Endl. Prod. Fl. Norf. 10. 1833; Diplazium Bedd. 1868; Asplenium umbrosum var. HB. 489.
Atkinsoni Bedd. Ferns br. Ind. Suppl. 11 t. 359. 1876 — Himalaya.
Asplenium Clarke 1880.
atomarium Pr. Rel. Haenk. **1**. 40. 1825 = Cystopteris bulbifera.
atratum Bedd. Handb. Suppl. 33. 1892 — Munipore.
australe Pr. Tent. 98. 1836 = A. umbrosum var.
axillare Pr. Tent. 98. 1836 = A. filix femina*.
Azoricum Fée, Gen. 186. 1850-52 = A. filix femina.
Baldwinii (Hill.) C. Chr. Ind. 140. 1905 — Ins. Hawaii.
Asplenium Hill. Fl. Haw. 618. 1888.
bantamense Milde, Bot. Zeit. **1870**. 353 = Diplazium.
basilare Fée, Gen. 186. 1850-52 = A. silvaticum.
Biondii Christ, Nu. Giorn. bot. It. n. s. **4**. 91. 1897 — China.
biserratum Milde, Bot. Zeit. **1870** 353 = Diplazium.
Bourgaei Fourn. Mex. pl. **1**. 102. 1872 — Mexico.
brevisorum (Wall.) Moore, Ind. 117. 1859 — India bor.
Asplenium Wall. List n. 220. 1828; Hk. sp. **3**. 229. 1860. HB. 228.
Brownii J. Sm. Hist. Fil. 328. 1875 = A. umbrosum*.
Brunonianum Milde, Bot. Zeit. **1870**. 353 = Diplaziopsis javanica.
Caracasanum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
celtidifolium Milde, Bot. Zeit. **1870**. 353 = Diplazium.
ceylanense (Kl.) Moore, Ind. 181. 1860 — (Ceylon) Hort.
Asplenium Kl.; ht. van Houtte 1858; J. Sm. 1866; Diplazium Moore 1859.
Clarkei Bedd. Ferns br. Ind. Suppl. 11 t. 360. 1876 = A. nigripes var.
comosum Pr. Tent. 98. 1836 = Alsophila squamulata.
conchatum Fée, Gen. 188 t. 17 C. f. 1. 1850-52 = A. costale.
convexum Newm. Phytol. **4**. app. XIII. 1851 = A. filix femina.
cordatum Opiz ? = A. filix femina.
coreanum Christ, Bull. Boiss. II. **2**. 827. 1902 — Korea.
Corsicum Fée, Gen. 186. 1850-52 = A. filix femina.
costale (Desv.) C. Chr. Ind. 140. 1905 — Ind. occ.
Allantodia Desv. Prod. 265. 1827; Asplenium Moore 1857 (non Sw. 1801). Athyrium conchatum Fée 1850-52. NPfl. 224; Asplenium Moore 1857; HB. 228; Diplazium J. Sm. 1866 — Hypochlamys pectinata, squamulosa, Tussaci Fée 1850-52; Athyrium Fée 1866; Asplenium squamulosum et [Tussaci Moore 1860.
costale Moore, Ind. XLIX. 1857 = A. nigripes.
costale Milde, Bot. Zeit. **1870**. 353 = Diplazium.
costaricense Christ, Bull. Boiss. II. **5**. 252. 1905 — Costa Rica.
crenato-serratum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
crenatum (Sommerf.) Rupr.; Nyland. Spicil. pl. fenn. **2**. 14. 1844. Chr. 225. NPfl. 224 — Europa bor. Asia bor. et temp.
Aspidium Sommerf. Vet. Ak. Handl. **1834**. 104. 1835; Cystopteris Fr. 1842; Asplenium Fr. 1845. Mett. n. 232. HB. 224; Aspidium sibiricum Turcz. 1834 (nomen); Asplenium Turcz.; Kze. 1837.
crenulato-serrulatum Mak. Bot. Mag. Tokyo **13**. 26. 1899 = Dryopteris.

crenulatum Milde, Bot. Zeit. **1870**. 371 = Diplazium striatum.
cultratum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
Cumingianum Pr.; Milde, Bot. Zeit. **1870**. 353 (nomen) —?
Cumingii Milde, Bot. Zeit. **1870**. 353 = Diplazium.
cuneatum Heufl. Verh. zool. bot. Ges. **6**. 346. 1856 = Asplenium fissum.
cyatheaefolium Milde, Bot. Zeit. **1870**. 353 = Diplazium.
cyclosorum Rupr. Distr. Crypt. vasc. Ross. 41. 1845 = A. filix femina var.
cystopteroides Eat. Proc. Amer. Acad. **4**. 110. 1858 — Japonia. Liu Kiu.
Asplenium Hk. 1860. HB. 225; Hypodematium Kuhn 1889.
Davidi (Franch.) Christ, Bull. Soc. Fr. **52** Mém. I. 50. 1905 — Thibet orient. Polypodium Franch. Nouv. Arch. Mus. II. **10**. 119. 1887; Phegopteris Bedd. 1892.
decursivum Yabe, Bot. Mag. Tokyo **17**. 65. 1903 — Japonia.
decurtatum (Kze.) Pr. Tent. 98 t. 3 f. 3. 1836. Chr. 223. NPfl. 224 — Brasilia. Argentina. — Asplenium Kze.: Link, Fil. sp. 94. 1841. HB. 226; (Allantodia Kze. 1851).
decussatum Milde, Bot. Zeit. **1870**. 353 = Diplazium Petersenii.
Delavayi Christ, Bull. Soc. Fr. **52** Mém. I. 47. 1905 — China.
deltoideum Newm. Phytol. **4**. app. XI. 1851 = A. crenatum.
deltoideum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
denticulatum J. Sm. Hist. Fil. 328. 1875 = A. nigripes.
deparioides Christ, Farnkr. 223. 1897 = A. proliferum.
depauperatum Schum. En. pl. Saell. **2**. 17. 1803 = A. filix femina.
dilatatum Milde, Bot. Zeit. **1870**. 353 = Diplazium maximum.
distans (Don) Moore, Ind. 125, 181. 1860 — Nepal.
Asplenium Don, Prod. Fl. Nepal. 9. 1825.
distentum Milde, Bot. Zeit. **1870**. 353 = Diplazium expansum var.
Dombei Desv. Prod. 266. 1827 (Dombeyi auctt.) = A. filix femina var.
drepanophyllum Bedd. Handb. Suppl. 32. 1892 = A. falcatum.
drepanopterum (Kze.) A. Br.; Milde, Fil. Eur. 49. 1867 — Japonia-India bor. Yunnan. — Polypodium Kze. Linn. **23**. 278, 318. 1850; Asplenium A. Br. 1856. Mett. n. 244; Aspidium Mett. Fil. Lips. 93 t. 19 f. 1-4. 1856. Polypodium oxyphyllum Wall. 1828 (nomen, non Kze. 1848); Athyrium Moore 1857. NPfl. 224; Asplenium Hk. 1860. HB. 228. — Aspidium eburneum part. Wall. 1828 (nomen); Lastrea J. Sm. 1846 (nomen); Asplenium J. Sm. 1857. Mett. n. 235. Athyrium J. Sm. 1875.
Duthiei Bedd. Handb. Suppl. 34. 1892 — Himalaya.
Asplenium Bedd. JoB. **1889**. 72.
ebeneum Milde, Bot. Zeit. **1870**. 353 = ? Diplazium asperum.
eburnum J. Sm. Hist. Fil. 328. 1875 = A. drepanopterum.
expansum (Willd.) Moore, Ind. 91, 181. 1860 — Amer. merid.
Aspidium Willd. sp. **5**. 284. 1810.
expansum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
falcatum Bedd. Ferns S. Ind. t. 151. 1863. NPfl. 223 — India.
Asplenium drepanophyllum Bak. 1874. HB. 226 (non Kze. 1834); Athyrium Bedd. 1892.
fallaciosum Milde, Fil. Eur. 54. 1867 — China bor.
*fallax Milde, Fil. Eur. 49. 1867 = A. macrocarpum.
Fargesii Christ, Bull. Soc. Fr. **52** Mém. I. 49. 1905 — China.

Fauriei (Christ) Mak. Bot. Mag. Tokyo **17**. 160. 1903 — Japonia. China. Luzon.
Nephrodium Christ, Bull. Boiss. **4**. 671. 1896; Aspidium Christ 1898; Athyrium nikkoense Mak. 1903.

ferulaceum Christ, Bull. Boiss. II. **4**. 968. 1904 = Asplenium.

Fieldingianum (Kze.) Milde, Bot. Zeit. **1870**. 370 — India austr.
Allantodia Kze. Linn. **24**. 268. 1851; Asplenium Moore 1857. A. polymorphum Mett. n. 229. 1859 (non Wall. 1828).

filix femina (L.) Roth, Röm. Mag. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 65. 1900. Chr. 223. NPfl. 224 — Europa. Afr. bor. Ins. Atlant. Asia occ. et bor.-India. Java. Amer. bor.-Mexico-Peru-Argentina.
Polypodium L. sp. **2**. 1090. 1753; Aspidium Sw. 1801; Tectaria Cav. 1802; Nephrodium Michx. 1803; Asplenium Bernh. 1806. Mett. n. 246. HB. 227; Cyathea Bertol. 1819; Cystopteris Coss. et Germ. 1845; Lastrea Colomb 1888 — Athyrium rhaeticum Roth 1800; Polypodium pt. L. 1753.
— Asplenium pectinatum Wall. 1828. Mett. n. 241; Athyrium Pr. 1836 — India. — (an sp. propr.?).
— Asplenium melanolepis Franch. et Sav. 1879; Athyrium Christ 1896 Japonia.
— Cystopteris retusa Dcne. 1844 — India bor.
— Athyrium cyclosorum Rupr. 1845 — Asia et Amer. bor.
— Aspidium angustum Willd. 1810; Athyrium Pr. 1825. Asplenium Michauxii Spr. 1827; Athyrium Fée 1850-52 — Amer. bor.-Mexico.
— Nephrodium asplenioides Michx. 1803; Aspidium Sw. 1806; Athyrium Desv. 1827; Asplenium Athyrium Spr. 1804 — Amer. bor.
— Asplenium Martensii Kze. 1848. Mett. n. 248; Athyrium Moore 1860. A. Galeottii Fée 1850-52 — Mexico.
— Athyrium Dombei Desv. 1827; Asplenium Dombeyi Mett. 1864 — Mexico-Peru.
— Polypodium axillare Ait. 1789; Aspidium Sw. 1801; Allantodia Klf. 1824; Athyrium Pr. 1836; Asplenium J. Sm. 1846. Mett. n. 247 — Ins. Atlant. (f. auctt. plur. = A. umbrosum*).
— de formibus chinensibus vide Christ, Bull. Soc. Fr. **52** Mém. I. 47-49. 1905.
— conf. A. Bourgaei Fourn. et A. Schimperi Moug.

fimbriatum (Wall.) Moore, Ind. 185. 1860. NPfl. 224 — Himalaya. Yunnan.
Aspidium Wall. 1828 (nomen); ? Polystichum Pr. 1849; Asplenium Hk. sp. **3**. 234. 1860. HB. 229 — A. Andersoni Clarke 1876 — Davallia athyriifolia Bak. 1891 — (conf. A. lastreoides).

?flexile Moore, Ind. 185. 1860 = A. alpestre.

flexuosum Milde, Bot. Zeit. **1870**. 353 = Diplazium Preslianum.

foliolosum (Wall.) Moore, Cat. Hort. Sim 1859; Ind. 185. 1860 (pt.; nomen); Bedd. Handb. Suppl. 37. 1892 — India bor.
Asplenium Wall. 1829 (pt.; nomen).

fontanum Roth, Röm. Mag. **2**[1]. 105. 1799; Tent. Fl. Germ. **3**. 59. 1800 = [Asplenium.

fragile Spr. Anleit. **3**. 136. 1804 = Cystopteris.

fraxinifolium Milde, Bot. Zeit. **1870**. 353 = Diplazium.

fumaroides Pr. Rel. Haenk. **1**. 39 t. 6 f. 2. 1825 = Cystopteris fragilis.

Galeottii Fée, Gen. 187. 1850-52 = A. filix femina var. Martensii.

Gaudichaudii Fée, Gen. 188. 1850-52 — Ins. Hawaii.

gedeanum (Rac.) Christ, Journ. de Bot. **19**. 68. 1905 — Java. Annam.
Asplenium Racib. Pterid. Buitenz. 223. 1898.

Giraldii Christ, Nu. Giorn. bot. It. n. s. **4**. 91. 1897; Bull. Soc. bot. It. **1898**. 29. NPfl. 224 — China.

Göringianum (Kze.) Moore, Ind. 185. 1860 — Japonia. — Aspidium Kze. Bot. Zeit. **1848**. 557; Lastrea Moore 1858; Asplenium Mett. n. 243. 1859.

gracile Fourn. Mex. pl. **1**. 102. 1872 = Asplenium Glenniei.

grammitoides Milde, Bot. Zeit. **1870**. 353 = Diplazium.

grammitoides Bedd. Ferns br. Ind. Correct. II. 1870 = A. falcatum.

grande Fourn. Mex. pl. **1**. 101. 1872 = A. achilleifolium var.

grandifolium Milde, Bot. Zeit. **1870**. 353 = Diplazium.

Griffithii Milde, Bot. Zeit. **1870**. 354; Bedd. Ferns br. Ind. Suppl. 12. 1876 = Diplazium.

gymnogrammoides (Kl.) Bedd. Ferns S. Ind. t. 156. 1863 — Ceylon. Asplenium Kl.; Mett. Aspl. 193 n. 234. t. 16 f. 13-14. 1859.

Haenkeanum Pr. Tent. 98. 1836; Epim. 66. 1849 = Asplenium cristatum.

Halleri Roth, Röm. Mag. **2**[1]. 105. 1799; Tent. Fl. Germ. **3**. 60. 1800 = Asplenium fontanum.

Henryi (Bak.) Diels, Nat. Pfl. **1**[4]. 224. 1899 — China. Asplenium Bak. Ann. Bot. **5**. 306. 1891.

hians Milde, Bot. Zeit. **1870**. 354 = Diplazium.

Hohenackerianum (Kze.) Moore, Ind. XLIX. 1857 — India. Ceylon. Asplenium Kze. Bot. Zeit. **1849**. 711 (nomen). Mett. n. 233. HB. 225; Allantodia Kze. Farnkr. **2**. 63 t. 126. 1850.

Hookerianum Moore, Ind. 185. 1860 = A. spinulosum.

incisum Newm. Phytol. **4**. app. XIII. 1851 = A. filix femina.

incisum Fée, Gen. 187. 1850-52 = A. filix femina.

integrifolium Milde, Bot. Zeit. **1870**. 353 = Diplazium cordifolium.

irriguum Gray, Nat. Arr. br. Pl. **2**. 10. 1821 = A. filix femina.

juglandifolium Milde, Bot. Zeit. **1870**. 371 = Diplazium flavescens.

Klotzschii Milde, Bot. Zeit. **1870**. 354 = Diplazium.

Kunzeanum Milde; Salom. Nom. 110. 1883 = ? Diplazium asperum.

laetum Gray, Nat. Arr. br. Pl. **2**. 10. 1821 = A. filix femina.

lanceolatum Heufl. Verh. zool. bot. Ges. **6**. 345. 1856 = Asplenium.

lanceum Moore, Ind. 185. 1860 = A. macrocarpum.

lanceum Milde, Bot. Zeit. **1870**. 354 = Diplazium.

lastreoides (Bak.) Diels, Nat. Pfl. **1**[4]. 224. 1899 — China. Korea. Asplenium Bak. JoB. **1888**. 227. (f. Bedd. = A. fimbriatum).

latifolium Pr. Tent. 98 t. 3 f. 4. 1836; Epim. 66. 1849 — Chile. Asplenium Sturm 1858.

latifolium Milde, Bot. Zeit. **1870**. 354 = Diplazium maximum var.

laxum Schum. Enum. Pl. Saell. **2**. 16. 1803 = A. filix femina.

laxum Pappe et Raws. Syn. Fil. Afr. austr. 16. 1858 = A. scandicinum var.

Lechleri Milde, Bot. Zeit. **1870**. 353 = Diplazium.

longifolium Milde, Bot. Zeit. **1870**. 354 = Diplazium lobulosum.

longipes Christ, Bull. Soc. Fr. **52** Mém. I. 48. 1905 — Yunnan.

Mac Donellii Bedd. Handb. Suppl. 34. 1892 (McDonelli) — Himalaya. China merid. — Asplenium Bedd. JoB. **1889**. 73.

Mackinnoni (Hope) C. Chr. Ind. 143. 1905 — India bor.-occ. Asplenium Hope, JoB. **1896**. 124; Journ. Bomb. Nat. Hist. Soc. **14**. 122 t. 23. 1902.

macrocarpon Fée, Gen. 188. 1850-52 — Mexico. Asplenium Moore 1859 — (an Asplenium sp. aff. A. bulbifero?).

macrocarpum (Bl.) Bedd. Ferns S. Ind. t. 152, 153. 1863. Chr. 224. NPfl. 224 —

Japonia. China. Asia trop. — Aspidium Bl. Enum. 162. 1828; Asplenium Bl.; Hk. 1860. HB. 227; Lastrea? Moore 1858 — Aspidium puncticaule Bl. 1828; Athyrium Moore 1860; Asplenium Mett. 1866 — A. fallax Mett. n. 236. 1859; Athyrium Milde 1867 — Asplenium decipiens Mett. n. 237. 1859 — Aspidium lanceum Kze. 1846; Nephrodium Moore 1858; Athyrium Moore 1860 — Nephrodium sparsum var. membranaceum Arnott; HB. 498.

marginatum Milde, Bot. Zeit. **1870**. 353 = Diplazium.

Martensii Moore, Ind. 186. 1860 = A. filix femina var.

medium (Carm.) Moore, Ind. 186. (NPfl. 224) — Tristan d'Acunha. Aspidium Carm. Tr. Linn. Soc. **12**. 511. 1818; Asplenium Hk. 1860. HB. 225.

melanocaulon Milde, Bot. Zeit. **1870**. 371 = Diplazium.

melanolepis Christ, Bull. Boiss. **4**. 668. 1896 = A. filix femina.

mesosorum Mak. Bot. Mag. Tokyo **13**. 82. 1899 — Japonia. Asplenium Mak. Bot. Mag. Tokyo **12**. 120. 1898.

Meyenianum Milde, Bot. Zeit. **1870**. 354 = Diplazium.

Michauxii Fée, Gen. 186. 1850-52 = A. filix femina var.

microsorum Mak. Bot. Mag. Tokyo **13**. 25. 1899 — Japonia. Asplenium Mak. Bot. Mag. Tokyo — 1892.

molle Roth, Röm. Mag. 2^{1}. 106. 1799; Tent. Fl. Germ. **3**. 61. 1800 = A. filix femina.

molliusculum Baumg. Enum. stirp. Transs. **4**. 31. 1846 = A. filix femina.

mongolicum (Franch.) Diels, Nat. Pfl. 1^{4}. 224. 1899 — China. Asplenium Franch. Nouv. Arch. Mus. II. **7**. 161. 1883.

montanum Röhl; Spr. Anleit. **3**. 143. 1804 = Cystopteris.

Moritzianum (Kl.) Fourn. Mex. pl. **1**. 102. 1872 (nomen) — Mexico. Caracas. Puerto Rico. — Asplenium Kl. hb.

multicaudatum Pr. Tent. 98. 1836 = A. spectabile.

mutilum Milde, Bot. Zeit. **1870**. 354 = Diplazium.

nephrodioides (Bak.) Christ, Bull. Soc. Fr. **52** Mém. I. 47. 1905 — China merid. Asplenium Bak. JoB. **1887**. 170.

Newtoni (Bak.) Diels, Nat. Pfl. 1^{4}. 224. 1899 — Ins. S. Thomé (Guinea). Asplenium Bak. Ann. Bot. **5**. 307. 1891.

nigripes (Bl.) Moore, Ind. XLIX. 1857. Chr. 224. NPfl. 224 — China. Asia trop. Madagascar. — Aspidium Bl. Enum. 162. 1828; Asplenium Bl.; Hk. 1860. HB. 227 — A. tenuifrons Wall. 1828; Athyrium Moore 1857. ? Asplenium gracile Don 1825 — Allantodia paludosa Zippel; Kze. 1848; Asplenium Moore 1858. Mett. n. 179 — Athyrium Clarkei Bedd. 1876; Asplenium Atk.; Clarke 1880.

nigripes Bedd. Ferns S. Ind. t. 157. 1863 = A. solenopteris.

Nigritianum J. Sm. Hist. Fil. 328. 1875 = Asplenium pedicularifolium.

nikkoense Mak. Bot. Mag. Tokyo **17**. 43. 1903 = A. Fauriei.

niponicum (Mett.) Hance, Journ. Linn. Soc. **13**. 92. 1873. NPfl. 224 — Japonia. China. — Asplenium Mett. Ann. Lugd. Bat. **2**. 240. 1866. HB. 227.

nitidulum (Kze.) Milde, Bot. Zeit. **1870**. 370 — Java. Allantodia Kze. Bot. Zeit. **1848**. 191; Asplenium Moore 1857. Mett. n. 180.

obovatum Fée, Gen. 186. 1850-52 = Asplenium lanceolatum var.

obtusum Milde, Bot. Zeit. **1870**. 354 = Diplazium.

Okuboanum Mak. Bot. Mag. Tokyo **13**. 16. 1899 — Japonia. Aspidium Mak. Bot. Mag. Tokyo — 1892.

oosorum (Bak.) Christ, Engl. Jahrb. **23**. 349. 1896 — Samoa. Asplenium Bak. JoB. **1876**. 343.

ordinatum Christ, Bull. Boiss. II. **4**. 967. 1904 — Costa Rica.
oshimense Christ, Bull. Boiss. II. **1**. 1017. 1901 — Liu Kiu.
Ottonis Milde, Bot. Zeit. **1870**. 371 = Diplazium.
ovatum Roth, Röm. Mag. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 64. 1800 = A. [filix femina.
oxyphyllum Moore, Ind. XLIX. 1857 = A. drepanopterum.
pallidum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
patens Milde, Bot. Zeit. **1870**. 354 = Asplenium.
pectinatum Pr. Tent. 98. 1836 = A. filix femina var.
pectinatum Bedd. Ferns S. Ind. t. 155. 1863 = A. solenopteris.
pectinatum Fée, 11 mém. 31. 1866 = A. costale.
pedatum Milde, Bot. Zeit. **1870**. 371 = Diplazium.
pentagonum Moore, Ind. 186. 1860 (nomen) — Moulmein.
petiolare Milde, Bot. Zeit. **1870**. 354 = Diplazium.
physosorum Sieber; Milde, Fil. Eur. 49. 1867 (nomen) = A. umbrosum.
pinnatifidum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
plantagineum Milde, Bot. Zeit. **1870**. 354 = Diplazium plantaginifolium.
Poiretianum Pr. Tent. 98. 1836 = A. scandicinum.
polypodioides Schur, Oest. Bot. Zeit. **1858**. 194; Enum. Pl. Trans. 837. 1866. [= A. alpestre.
polypodioides Milde, Bot. Zeit. **1870**. 354 = Diplazium.
Pontederae Desv. Prod. 266. 1827 = Cystopteris fragilis.
porrectum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
procerum (Wall.) Milde, Bot. Zeit. **1866**. 376 — India bor.
  Asplenium? Wall. 1829 (nomen); Clarke, Tr. Linn. Soc. II. Bot. **1**. 495. 1880; A. umbrosum var. HB. 489.
proliferum (Klf.) C. Chr. Ind. 145. 1905 — Ins. Hawaii. ? N. S. Wales.
  Dicksonia Klf. Enum. 225. 1824; Deparia Hk. et Grev. 1831. HB. 55; Cibotium Pr. 1836; ? Davallia F. Muell. 1865; Deparia Macraei Hk. et Grev. 1829 — Asplenium deparioides Brack. 1854. HB. 226; Athyrium Chr. 1897. 223.
proliferum Moore, Gard. Chr. **1866**. 778 — Ceylon.
proliferum Milde, Bot. Zeit. **1870**. 353 = Diplazium.
pterorachis Christ, Bull. Boiss. **4**. 668. 1896 — Japonia. Korea.
pumilio Christ, Bull. Boiss. II. **3**. 147. 1903 — Mexico.
puncticaule Moore, Ind. 186. 1860 = A. macrocarpum.
purpureum Lowe; Milde, Fil. Eur. 49. 1867 = ? A. nigripes.
pycnosorum Christ, Bull. Boiss. II. **2**. 827. 1902 — Korea.
radicans Milde, Bot. Zeit. 1870. 354 = Diplazium sp. dub.
reductum Christ, Bull. Boiss. II. **4**. 966. 1904 — Ecuador. Costa Rica.
  Asplenium aspidioides Sod. Cr. vasc. quit. 182. 1893 (non Schlecht. 1825).
regium Spr. Anleit. **3**. 139. 1804 = Cystopteris alpina*.
rhaeticum Roth, Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 67. 1800 = A. filix femina.
rhaeticum Sadler, Descr. pl. epiph. Hung. 22. 1820 = Cystopteris fragilis.
rhaeticum Dalla Torre, Anl. wiss. Beob. Alpenr. **2**. 348. 1882 = A. alpestre.
rigescens Mak. Bot. Mag. Tokyo **13**. 27. 1899 — Japonia.
Roemerianum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
roseum Christ, Bull. Boiss. **6**. 961. 1898 — China merid.
rupicola (Hope) C. Chr. Ind. 145. 1905 — India bor.
  Asplenium Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 531 t. 5. 1899.
rutaceum Pr. Tent. 98. 1836 = Asplenium.

sandwichianum Pr. Tent. 98. 1836 = Diplazium.

Sarasinorum Christ, Verh. Nat. Ges. Basel **11.** 434. 1896; Ann. Jard. Buit. **15.** 123 t. 15 f. 6. 1897 — Celebes. Luzon.

?scabrum Pr. Epim. 67. 1849 = Alsophila tristis.

scandicinum (Willd.) Pr. Tent. 98. 1836 — Africa austr. cum ins. Ins. Hawaii. Aspidium Willd. sp. **5.** 285. 1810; Allantodia Klf. 1824; Cystopteris Desv. 1827; Nephrodium Bory 1833; Asplenium Heller 1897. A. Poiretianum Gaud. 1827; Athyrium Pr. 1836. Asplenium aspidioides Schlecht. 1825. Mett. n. 239. HB. 228; Allantodia Kze. 1848; Athyrium Christ 1897. 224. NPfl. 224 (vix Pr. 1836) — A. laxum Pappe et Raws. 1858 (non Schum. 1803); Asplenum Kuhn 1868 (non R. Br. 1810) — A. multisectum Brack. 1854. Mett. n. 242 (non Bl. 1828).

Schimperi Moug.; Fée, Gen. 187. 1850-52 — Abyssinia. India bor. Asplenium A. Br. 1867. HB. 489.

?scorpioideum Chauvin, Mém. Soc. Linn. Calvados **1825.** 402 = ? A. filix femina.

selenopteris Bedd. Handb. 166. 1883. Chr. 225 = A. solenopteris.

serrulatum Milde, Bot. Zeit. **1870.** 353 = Diplazium manilense.

setiferum C. Chr. Ind. 146. 1905 — India bor. Allantodia tenella Wall. hb.; Asplenium Hope, Journ. Bomb. Nat. Hist. Soc. **12.** 529 t. 4. 1899 (non Roxb. 1816 nec Fée 1850-52).

setigerum Hort.; Gard. Chr. III. **12.** 249. 1892 (nomen) — Hort.

silvaticum (Bl.) Milde, Bot. Zeit. **1866.** 376 — Java. Celebes. Ins. Philipp. Allantodia Bl. Enum. 173. 1828; Asplenium Moore 1857 (non Pr. 1825). Brachysorus woodwardioides Pr. 1849; Asplenium Bak. 1867. HB. 229. (non A. woodwardioideum Gardn. 1842); Athyrium Christ 1895. Diplazium brevisorum J. Sm. 1841. Athyrium basilare Fée 1850-52; Asplenium Moore 1857.

silvaticum Milde, Bot. Zeit. **1870.** 354 = Diplazium.

sinense Rupr. Distr. Cr. vasc. Ross. 41. 1845 — China bor.

sitchense Rupr. Distr. Cr. vasc. Ross. 41. 1845 = A. filix femina var.

Skinneri Moore, Ind. 187. 1860 (nomen). NPfl. 224 — Guatemala. Asplenium Bak. Syn. 226. 1867.

solenopteris (Kze.) Moore, Ind. 43, 187. 1857. Bedd. 1883. Chr. 225 (selenopteris) — India austr. Ceylon. — Allantodia Kze. Linn. **24.** 266. 1851; Asplenium Mett. n. 238. 1859. (HB. 227) (selenopteris).

solutum Christ, Bull. Boiss. II. **4.** 967. 1904 — Costa Rica.

Sorzogonense Milde, Bot. Zeit. **1870.** 354 = Diplazium.

speciosum Milde, Bot. Zeit. **1870.** 354 = ? Diplazium sorzogonense.

spectabile (Wall.) Pr. Tent. 98. 1836 — India bor. Asplenium Wall. 1828 (nomen); Mett. Aspl. 196 n. 240. 1859 — A. multicaudatum Wall. 1828 (nomen); Clarke, Tr. Linn. Soc. II. Bot. **1.** 502. 1880; Athyrium Pr. 1836 (nomen); Asplenium umbrosum var. HB. 489 — Diplazium Jerdoni Bedd. 1869 — (an potius Diplazium?).

sphaerocarpon Fée, Gen. 186. 1850-52 = Dryopteris athyrioides.

spinulosum (Maxim.) Milde, Bot. Zeit. **1866.** 376. Chr. 226. NPfl. 224 — Japonia. Korea. China-Himalaya. — Cystopteris Maxim. Mém. Ac. St. Pétersb. **9.** 340. 1859; Asplenium Bak. 1867. HB. 225 — Athyrium Hookerianum Moore 1860 (nomen); Asplenium subtriangulare Hk. 1867. HB. 225; Athyrium Bedd. 1868.

spinulosum Milde, Bot. Zeit. **1870.** 354 = Diplazium.

squamulosum Fée, 11 mém. 32 t. 9 f. 2. 1866 = A. costale var.
striatum Milde, Bot. Zeit. **1870**. 354 = ? Diplazium.
strigillosum Moore; Salom. Nom. 112. 1883 = A. nigripes.
subserratum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
subsimile Christ, Bull. Soc. Bot. It. **1898**. 29 — China.
subtriangulare Bedd. Ferns br. Ind. t. 242. 1868 = A. spinulosum var.
sylvaticum vide silvaticum.
tenerum Fée, Gen. 186. 1850-52 = A. umbrosum.
tenuifrons Moore, Ind. 43, 188. 1857 = A. nigripes.
tenuisectum Moore, Ind. 188. 1860 = Cystopteris.
Thelypteris Spr. Anleit. **3**. 134. 1804 = Dryopteris.
thelypteroides Desv. Prod. 266. 1827 = A. acrostichoides.
Thwaitesii Milde, Bot. Zeit. **1870**. 354 = Diplazium.
tomentosum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
trifidum Roth, Röm. Mag. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 63. 1800 = A. filix femina.
Tussaci Fée, 11 mém. 32 t. 9 f. 1. 1866 = A. costale.
umbrosum (Ait.) Pr. Tent. 98. 1836. Chr. 225. NPfl. 224 — Ins. atlant. Afr., Asia, Austr. trop. N. Zealand. — Polypodium Ait. Hort. Kew. **3**. 466. 1789; Aspidium Sw. 1801; Allantodia Klf. 1824; Asplenium J. Sm. 1841. Mett. n. 231. HB. 229; Diplazium Bedd. 1883. Asplenium Aitoni Moore 1857. Tectaria caudata Cav. 1801; Aspidium Sw. 1806.
—* Allantodia australis R. Br. 1810; Athyrium Pr. 1836; Asplenium Brack. 1854. Asplenium Brownii J. Sm. 1841. Mett. n. 230; Athyrium J. Sm. 1875 — Allantodia tenera R. Br. 1810; Athyrium Fée 1850-52 — Asplenium muricatum Mett. 1866 (Java) — A. bellum Clarke 1880 (Himalaya). — conf. A. filix femina *axillare, A. assimile, A. spectabile, A. procerum, omnia f. auctt. plur. subspecies A. umbrosi.
umbrosum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
uropteron (Miq.) C. Chr. Ind. 147. 1905 — Japonia.
Asplenium Miq. Ann. Lugd. Bat. **3**. 174. 1867.
vestitum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
villosum Milde, Bot. Zeit. **1870**. 354 = Diplazium.
violascens Diels, Engl. Jahrb. **29**. 196. 1900 — China.
viridifrons Mak. Bot. Mag. Tokyo **13**. 15. 1899 — Japonia.
Asplenium Mak. Bot. Mag. Tokyo — 1892.
Wagnerianum Milde, Bot. Zeit. **1870**. 354 = Diplazium Wageneri.
Wardii (Hk.) Mak. Bot. Mag. Tokyo **13**. 15. 1899 — Japonia. Korea. China.
Asplenium Hk. sp. **3**. 189. 1860; 2 Cent. t. 33. HB. 217.
Wilsoni Christ, Bull. Boiss. II. **3**. 512. 1903 — China.
woodwardioides Christ, Verh. Nat. Ges. Basel **11**. 233. 1895 = A. silvaticum.
yokoscense (Franch. et Sav.) Christ, Bull. Boiss. **4**. 668. 1896 — Japonia.
Asplenium Franch. et Sav. Enum. pl. Jap. **2**. 225, 622. 1879.
Zeylanicum Milde, Bot. Zeit. **1870**. 354 = Diplazium.

AUSTROGRAMME Fournier, Ann. sc. nat. V. **18**. 278. 1873.
Deplanchei Fourn. Ann. sc. nat. V. **18**. 279. 1873 = Polypodium.
marginata Fourn. Ann. sc. nat. V. **18**. 279. 1873 = Syngramma.

AZOLLA Lamarck, Encycl. méth. **1**. 343. 1783 — (Salviniaceae Gen. **1**).
africana Desv. Prod. 178. 1827 = A. pinnata.
arbuscula Desv. Prod. 178. 1827 = A. filiculoides.

bonariensis Bertol. Misc. **21**. t. 5 f. 2 a, b. 1860 (Rend. Sess. Acc. Sci. Bol. **1859-60**. **64**. 1860) = A. caroliniana.

caroliniana Willd. sp. **5**. 541. 1810. Bak. Hdb. 138. NPfl. 401 — Amer. septentr. Brasilia. — A. microphylla Klf. 1824. A. cristata Klf. 1824. Salvinia Azolla Raddi 1825. Azolla mexicana Schlecht. 1830. ? Carpanthus axillaris Rafin. 1808.

cristata Klf. Enum. 274. 1824 = A. caroliniana.

decomposita Zoll. Syst. Verz. 51, 52. 1854 = A. pinnata.

densa Desv. Prod. 178. 1827 = A. caroliniana.

filiculoides Lam. Enc. **1**. 343. 1783. Bak. Hdb. 137. NPfl. 401 — California -Chile. Brasilia-Argentina. — A. magellanica Willd. 1810.

guineensis Schum. Vid. Selsk. Afh. **4**. 236 (seors 462). 1827 = A. pinnata.

japonica Franch. et Sav. En. pl. Jap. **2**. 195. 1876 = A. pinnata.

magellanica Willd. sp. **5**. 541. 1810 = A. filiculoides.

mexicana Schlecht. et Cham. Linn. **5**. 625. 1830 = A. caroliniana.

microphylla Klf. Enum. 273. 1824 = A. caroliniana.

nilotica Dcne.; Mett. Pl. Tinn. 54 t. 25. 1867. Bak. Hdb. 138. NPfl. 401 — Africa orient.

pinnata R. Br. Prod. Fl. N. Holl. 167. 1810. Bak. Hdb. 138. NPfl. 401 — Japonia. Asia, Austr., Africa trop.
Rhizosperma Meyen 1836. Azolla africana Desv. 1827.

portoricensis Spr. Syst. **4**. **9**. 1827 = A. caroliniana.

rubra R. Br. Prod. Fl. N. Holl. 167. 1810. Bak. Hdb. 137 — Australia. N. Zealand.

squamosa Molina, Saggio s. storia nat. Chile ed. II. 125. 1810 — A. filiculoides.

BAKERIOPTERIS O. Ktze. Rev. Gen. Pl. **2**. 807. 1891.

gleicheniodes O. Ktze. Rev. Gen. Pl. **2**. 807. 1891 = Pellaea.

? petiolata O. Ktze. Rev. Gen. Pl. **2**. 807. 1891 } = Pellaea pinnata.
pinnata O. Ktze. Rev. Gen. Pl. **2**. 807. 1891 }

triphylla O. Ktze. Rev. Gen. Pl. **2**. 807. 1891 = Doryopteris.

BALANTIUM Kaulfuss, Enum. Fil. 228. 1824 — (Cyatheaceae Gen. **1**).

antarcticum Pr. Tent. 134. 1836 = Dicksonia.

arborescens Hk. Gen. t. 20. 1838 = Dicksonia.

auricomum Klf. Enum. 228 t. 1 f. 13. 1824 = Dicksonia arborescens.

Berteroanum Kze. Anal. 40. 1837 = Dicksonia.

Beyrichii Roem.; Kze. Anal. 40. 1837 (nomen) —?

Blumei Kze. Bot. Zeit. **1848**. 214 = Dicksonia.

Brownianum Pr. Tent. 134 t. 5 f. 4. 1836 = Davallia dubia.

chrysotrichum Hassk. Obs. Fil. Jav. **1**. 53. 1856 = Dicksonia Blumei.

coniifolium (Hk.) J. Sm. Hist. Fil. 258. 1875. NPfl. 119 — Jamaica. Mexico-Ecuador-Brasilia. — Dicksonia Hk. sp. **1**. 70 t. 24 A. 1844. HB. 51. Chr. 314; Microlepia Fée 1866. Dicksonia Martiana Kl.; Hk. 1844; Balantium Fée 1869.

culcita (L'Hérit.) Klf. Enum. 229. 1824. NPfl. 119 — Ins. Azorae. Madeira. Teneriffa. — Dicksonia L'Hérit. Sert. Angl. 31. 1788. HB. 51. Chr. 313; Culcita macrocarpa Pr. 1836.

fibrosum Fée, Gen. 341. 1850-52 = Dicksonia.

glaucescens Link, Fil. sp. 40. 1841 = Cibotium barometz.

Karstenianum Kl. Linn. **20**. 444. 1847 = Dicksonia.

lanatum Fée, Gen. 341. 1850-52 = Dicksonia.

magnificum de Vriese; Jungh. Java ed. germ. **1**. 476. 1852 = Dicksonia Blumei.

Martianum Fée, Cr. vasc. Br. **1**. 155. 1869 = B. coniifolium.

Sellowianum Pr. Tent. 134. 1836 = Dicksonia.

squarrosum Kze. Linn. **23**. 239. 1850 = Dicksonia.

stramineum (Labill.) Diels, Nat. Pfl. 1^4. 119. 1899 — Polynesia.

Dicksonia Labill. Sert. Austr. Cal. 7 t. 10. 1824. HB. 51; Sitolobium Brack. 1854; Dennstaedtia J. Sm. 1875 — Dicksonia Torreyana Brack. 1854.

thyrsopteroides Fourn. L'Ill. Hort. **23**. 99. 1876 = Dicksonia.

BATHMIUM (Presl) Link, Fil. spec. 99, 114. 1841 = **Aspidium.**

Aspidium § Bathmium Pr. Tent. 88. 1836.

alatum Fée, Gen. 287. 1850-52 = A. vastum.

Billardieri Fée, Gen. 287. 1850-52 = A. sinuatum.

ebeneum Fée, Gen. 288. 1850-52 = A. pica.

fraxinifolium Link, Fil. sp. 115. 1841 = A. martinicense.

heracleifolium Fée, Gen. 287. 1850-52 = A. trifoliatum.

kanakorum Fourn. Ann. sc. nat. V. **18**. 301. 1873.

macrocarpon Fée, Gen. 287. 1850-52 = A. plantagineum.

macrophyllum Link, Fil. sp. 114. 1841 = A. martinicense.

nicaraguense Fourn. Bull. Soc. Fr. **19**. 256. 1872.

plantagineum Fourn. Bull. Soc. Fr. **19**. 254. 1872.

Poeppigii Fourn. Bull. Soc. Fr. **19**. 256. 1872.

repandum Fée, Gen. 287. 1850-52.

Seemanni Fourn. Ann. sc. nat. V. **18**. 300. 1873 = ? A. latifolium.

sinuatum Fée, Gen. 287. 1850-52 = A. plantagineum.

sinuatum Fourn. Ann. sc. nat. V. **18**. 300. 1873.

? subfalcatum Fée, Gen. 287. 1850-52 = Polypodium Zippelii.

subtriphyllum Fourn. Ann. sc. nat. V. **18**. 291. 1873.

trifoliatum Link, Fil. sp. 114. 1841.

? undulatum Fée, Gen. 287. 1850-52 = ? Polypodium membranaceum.

villosum Fée, Gen. 289. 1850-52.

BELVISIA Mirbel, (Hist. nat. vég. ed. Déterv. **5**. 473. 1802?) Hist. nat. des plantes **4**. 65. 1803.

australis Mirb. (Hist. nat. vég. **5**. 473. 1802?) = Actiniopteris.

digitata Mirb. (l. c.) = Schizaea.

septentrionalis Mirb. (l. c.) = Asplenium.

siliquosa Mirb. (l. c.) = Ceratopteris thalictroides.

spicata Mirb. (l. c.) = Hymenolepis.

BERGERA Schaffner (Fée, 9 mém. 30. 1857) = **Trichomanes.**

BLECHNIDIUM Moore, Ferns Gr. Brit. Nat. Pr. oct. ed. **2**. 210; Ind. Fil. CLV. 1860 (?) = **Blechnum.**

melanopus Moore, l. c.

BLECHNOPSIS Presl, Epim. bot. 115. 1849 = **Blechnum.**

brasiliensis Pr. l. c. 115 — cartilaginea Pr. l. c. 116 — Finlaysoniana Pr. l. c. 116 — orientalis Pr. l. c. 117 — serrulata Pr. l. c. 119 = Blechnum sp. homonym.

? adnata Pr. l. c. 119 — Cumingiana Pr. l. c. 116 — elongata Pr. l. c. 117 — ? imbricata Pr. l. c. 118, 262 — ? javanica Pr. l. c. 118 — latifolia Pr. l. c. 116 — ? longifolia Pr. l. c. 118 — pectinata Pr. l. c. 118 — pyrophila Pr. l. c. 117 — salicifolia Pr. l. c. 116 — stenophylla Pr. l. c. 118 = Blechnum orientale formae.

denticulata Pr. Epim. 116. 1849 = B. serrulatum.

malaccensis Pr. Epim. 120. 1849 = B. serrulatum.

nitida Pr. Epim. 116. 1849 = B. elongatum.

striata Pr. Epim. 119. 1849 = B. serrulatum.

BLECHNOPTERIS Trevisan, Atti dell'Ist. Veneto II. **2**. 166. 1851 = **Blechnum.**
ambigua Trev. l. c. = B. laevigatum.
australis Trev. l. c. = Blechnum.
hastata Trev. l. c. = B. auriculatum.
procera Trev. l. c. = B. capense.
punctulata Trev. l. c. = Blechnum.

BLECHNUM Linné, Spec. plant. **2**. 1077. 1753 — (Polypodiaceae Gen. **51**).

Subgenera: B. = **Eublechnum.**
L. = **Lomaria.**
S. = **Salpichlaena.**

Obs. Vide sub Lomaria dubias species subgeneris Lomariae. — Numerus cit. post »Spicanta O. Ktze.« paginam in O. Ktze. Rev. Gen. Pl. **2**. significat.

(L) **acrodontum** (Fée) C. Chr. Ind. 150. 1905 — Mexico.
Lomaria Fée, 8 mém. 70. 1857.
acuminatum Fée, Gen. 75. 1850-52 = B. occidentale.
acuminatum Sturm, Flora **1853**. 362 = B. arcuatum.
(L) **acutum** (Desv.) Mett. Ann. sc. nat. V. **2**. 225. 1864 — Columbia-Peru-Brasilia.
Lomaria Desv. Prod. 290. 1827. HB. 180; Spicanta O. Ktze. 821. 1891. Lomaria cuspidata Kze. 1834.
adnatum Reinw.; de Vriese, Ned. Kr. Arch. **1**. 10. 1846 = B. orientale.
aduncum Liebm. Vid. Selsk. Skr. V. **1**. 237 (seors. 85). 1849 = B. occidentale.
agrostifolium Goldm. Nova Acta **19** Suppl. I. 459. 1843 = B. orientale.
alpinum Mett. Fil. Lips. 64. 1856 = B. penna marina.
(L) **alternans** (Col.) C. Chr. Ind. 150. 1905 — N. Zealand.
Lomaria Col. Tr. N. Zeal. Inst. **28**. 616. 1896.
(L) **amabile** Mak. Bot. Mag. Tokyo **11**. 83. 1897 — Japonia.
Lomaria crenulata Moore (HB. 180 nomen); Blechnum Salom. 1883 (nomen); Mak. 1901.
ambiguum Klf.; Pr. Tent. 103. 1836 = B. laevigatum.
(L) **andicola** (Phil.) C. Chr. Ind. 150. 1905 — Chile.
Lomaria Phil. Anal. Univ. Chil. **94**. 353. 1896.
(L) **andinum** (Bak.) C. Chr. Ind. 150. 1905 — Andes.
Lomaria Bak. Syn. 482. 1874; Spicanta O. Ktze. 821. 1891.
angustatum Schrad. Gött. gel. Anz. **1824**. 872 = B. serrulatum.
angustifolium Willd. sp. **5**. 414. 1810 = B. serrulatum.
angustifolium Poir. Enc. Suppl. **1**. 644. 1810 = Woodwardia areolata.
angustifolium Roxb. Calc. Journ. **4**. 501. 1844 = Taenitis blechnoides.
angustifolium Hier. Engl. Jahrb. **34**. 472. 1904 = B. polypodioides.
angustifrons Fée, 7 mém. 25 t. 9 f. 2. 1857 = B. asplenioides.
(B) **appendiculatum** Willd. sp. **5**. 410. 1810 — Columbia.
Mesothema Pr. 1849; Struthiopteris Trev. 1869.
(L) **araucanum** (Phil.) C. Chr. Ind. 150. 1905 — Chile.
Lomaria Phil. Anal. Univ. Chil. **94**. 354. 1896.
arborescens Noronha; ex Thouars, Mél. bot et voy. 1811 — Madagascar.
(B) **arcuatum** Remy; Fée, Gen. 73. 1850-52 (nomen); Gay, Fl. Chil. **6**. 477. 1853. HB. 185. NPfl. 247 — Chile.
Spicanta O. Ktze. 821. 1891. Blechnum acuminatum Sturm 1853 (non Fée 1850-52); Lomaria Bibreae Sturm —.; Blechnum Mett. (1856).
(L) **areolare** (Harr.) C. Chr. Ind. 150. 1905 — Luzon.
Lomaria Harr. Journ. Linn. Soc. **16**. 28. 1877; Spicanta O. Ktze. 822. 1891.

(L) **asperum** (Kl.) Sturm, Enum. pl. vasc. cr. chil. 22. 1858. NPfl. 249 — Chile. Lomaria Kl. Linn. **20**. 344. 1847. HB. 177; Spicanta O. Ktze. 821. 1891.

(B) **asplenioides** Sw. Vet. Ak. Handl. **1817**. 72 t. 3 f. 3. HB. 183. Chr. 181. NPfl. 246 — Amer. trop. — Spicanta O. Ktze. 821. 1891. Blechnum ceteraccinum Raddi 1819.

Atherstoni Pappe et Raws. Syn. Fil. Afr. austr. 16. 1858 = B. punctulatum var.

(L) **attenuatum** (Sw.) Mett. Fil. Lips. 64 t. 3 f. 1-6. 1856. Chr. 179. NPfl. 247 — Africa austr. cum ins. N. S. Wales. Polynesia. ? Chile. Juan Fernandez. — Onoclea Sw. Schrad. Journ. **1800**². 73. 1801; Lomaria Willd. 1810. HB. 176 (pt.); Spicanta O. Ktze. 821. 1891 — Lomaria gigantea Klf. 1824; Blechnum Schlecht. 1827 — Lomaria heterophylla Desv. 1811; Blechnum Schlecht. 1827 — ? Lomaria bella Phil. 1857 (Chile. J. Fernandez).

attenuatum auctt. quoad pl. amer. trop. = B. meridense.

(B) **auriculatum** Cav. Descr. 262. 1802 — Amer. austr. temp. Mesothema Pr. 1849. Blechnum hastatum Klf. 1824. HB. 185. Chr. 182. NPfl. 246; Lomaria Kze. 1836; Mesothema Pr. 1849; Blechnopteris Trev. 1851; Struthiopteris Trev. 1869; Spicanta O. Ktze. 1891 — Blechnum trilobum Pr. 1825; Mesothema Pr. 1849; Lomaria Fée 1850-52; Struthiopteris Trev. 1869 — Blechnum remotum Pr. 1836. NPfl. 246; Mesothema Pr. 1849; Struthiopteris Trev. 1869. Blechnum pubescens Hk. 1837 (non Desv. 1827); Lomaria Kze. 1843 — Taenitis sagittaefera Bory 1828. — (f. auctt. plur. B. australe*).

auritum Goldm. Nova Acta **19** Suppl. I. 459. 1843 = A. auriculatum.

(B) **australe** L. Mant. 130. 1767. HB. 186. Chr. 182. NPfl. 246 — Ins. Caboverd. Africa austr. cum ins. Ins. antarct. — Lomaria Link 1841; Mesothema Pr. 1849; Blechnopteris Trev. 1851; Struthiopteris Trev. 1869; Spicanta O. Ktze. 821. 1891. Blechnum rigidum Sw. 1801; Mesothema Pr. 1849; Struthiopteris Trev. 1869 — Lomaria pumila Klf. 1824. — (conf. B. auriculatum).

(L) **Bakeri** C. Chr. Ind. 151. 1905 — Madagascar. Lomaria pubescens Bak. Journ. Linn. Soc. **15**. 415. 1876 (non Kze. 1843 nec Blechnum Desv. 1827 nec Hk. 1837); Spicanta O. Ktze. 1891.

Banisterianum Poir. Enc. Suppl. **1**. 644. 1810 = Woodwardia virginica.

(L)]**Banksii** (Hk. fil.) Mett.; Diels, Nat. Pfl. **1**⁴. 249. 1899 — N. Zealand. Lomaria Hk. fil. Fl. N. Zeal. **2**. 31 t. 76. 1854. HB. 178; Spicanta O. Ktze. 821. 1891.

Bibreae Mett. Fil. Lechl. exs. (Fil. Lechl. **1**. 14. 1856. syn.) = B. arcuatum.

(L) **biforme** (Bak.) Christ, Farnkr. 180. 1897. NPfl. 250 — Madagascar. Lomaria Bak. Journ. Linn. Soc. **15**. 415. 1876; Ic. pl. t. 1643; Spicanta O. Ktze. 822. 1891.

(B) **blechnoides** (Lag.) C. Chr. Ind. 151. 1905 — Amer. trop. Asplenium Lag.; Sw. Syn. 76. 1806; Blechnum unilaterale Sw. Berl. Mag. **4**. 79 t. 3 f. 1. 1810. HB. 184. Chr. 181. NPfl. 246; Spicanta O. Ktze. 1891 — Blechnum polypodioides Raddi 1819 (non infra).

blechnoides Keys. Pol. Cyath. Hb. Bung. 65. 1873 = B. valdiviense.

(B) *Blumei* Moore, Ind. 193. 1860 — Java. Lomaria auriculata Bl. Enum. 202. 1828 (non Desv. 1811); Mesothema javanicum Pr. 1849 (non Blechnum Bl. 1828); Struthiopteris Trev. 1869.

boreale Sw. Schrad. Journ. **1800**². 75. 1801 = B. spicant.

Boryanum Schlecht. Adumbr. 35 t. 19. 1827 = B. tabulare.

(B) **brasiliense** Desv. Berl. Mag. **5**. 330. 1811. HB. 184. Chr. 181. NPfl. 246 — Brasilia. Peru. — Blechnopsis Pr. 1849; Salpinchlaena Trevis 1869; Spicanta O. Ktze. 821. 1891 — Blechnum corcovadense Raddi 1819.

calophyllum Langsd. et Fisch. Ic. Fil. 20 t. 23. 1810 = B. serrulatum.

campylotis J. Sm. Ferns br. and for. 198. 1866 = B. occidentale var.

(L) **capense** (L.) Schlecht. Adumbr. 34 t. 18. 1825. Chr. 178. NPfl. 249 — Africa austr. Perak. (? Malesia). Polynesia. Ins. reg. austr. ? Amer. trop. Osmunda L. Mant. 306. 1771; Onoclea Thbg. 1800; Lomaria Willd. 1810; Spicanta O. Ktze. 821. 1891 — Osmunda procera Forst. 1786; Onoclea Spr. 1799; Blechnum Sw. 1801; Asplenium Bernh. 1802; Stegania R. Br. 1810; Lomaria Spr. 1827. HB. 179 (pt.); Parablechnum Pr. 1849; Blechnopteris Trev. 1851.

capense Burm. Fl. Ind. Fl. Cap. Prod. 28. 1768 (nomen) = B. australe.

carolinianum Walt. Fl. Carol. 257. 1788 = Woodwardia virginica.

(B) **cartilagineum** Sw. Syn. 114, 312. 1806. HB 184. NPfl. 247 — Australia. Luzon. Mishmee. — Blechnopsis Pr. 1849; Salpinchlaena Trev. 1869; Spicanta O. Ktze. 821. 1891. Blechnum nitidum var. contracta Hk. 1860. *Obs.* t. Sw. Adnot. 69 est planta Swartzii non in Australia sed in Cayenne lecta et verisimiliter non eadem ac planta Australiae ab auctoribus »cartilagineum« nominata.

cartilagineum Schkuhr, Kr. Gew. **1**. 101 t. 108 b. 1809 = B. occidentale.

caudatum Cav. Descr. 262. 1802 = B. occidentale var.

ceteraccinum Raddi, Opusc. sci. Bol. **3**. 294. 1819; Pl. Bras. **1**. 52 t. 60 f. 1. 1825 = B. asplenioides.

(L) **chilense** (Klf.) Mett. Fil. Lechl. **1**. 14. 1856 — Chile. Lomaria Klf. Enum. 154. 1824. (HB. 179) — L. Reedei Phil. 1873.

(L) **Christii** C. Chr. Ind. 152. 1905 — Costa Rica. Lomaria spissa Christ, Bull. Boiss. II. **4**. 1092. 1904 (non Fée 1850-52).

(B) *ciliatum* Pr. Rel. Haenk. **1**. 50. 1825 — Chile. Peru. — Parablechnum Pr. 1849.

ciliatum Mart. et Gal. Mém. Ac. Brux. **15**. 50. 1842 = B. occidentale.

ciliatum Christ, Farnkr. 177. 1897 = B. Moorei.

cognatum Pr. Epim. 107. 1849 = B. occidentale.

colubrinum Blanco, Fl. Filip. 834. 1837 — Ins. Philipp.

(L) **confine** (Cord.) C. Chr. Ind. 152. 1905 — Réunion. Lomaria Cord. Fl. Réun. 58. 1895.

confluens Schlecht. et Cham. Linn. **5**. 613. 1830 = B. occidentale.

(L) **contiguum** Mett. Ann. sc. nat. IV. **15**. 70. 1861 — N. Caledonia. Lomaria Carr. 1873; Fourn. 1873.

corcovadense Raddi, Opusc. sci. Bol. **3**. 294. 1819; Pl. bras. **1**. 54 t. 61. 1825 = B. brasiliense.

(L) **costaricensis** (Christ) C. Chr. Ind. 152. 1905 — Costa Rica. Lomaria Christ, Bull. Boiss. II. **4**. 1092. 1904 (non Bak. 1874).

crenulatum Salom. Nom. 115. 1883; Mak. Bot. Mag. Tokyo **15**. 96. 1901 = B. amabile.

crispum Hartm. Handb. Skand. Fl. ed. I. 372. 1820 = Cryptogramma.

Cumingianum Trevis. Atti Ist. Veneto II. **2**. 165. 1851 = B. serrulatum.

Cunninghamii Moore, Ind. 195. 1860 (descr.) = B. occidentale.

(L) **cycadifolium** (Colla) Sturm, Enum. pl. vasc. cr. chil. —. 1858 — Juan Fernandez. — Lomaria Colla, Mem. Ac. Torino **39**. 43 t. 71. 1836; L. lanuginosa Kze. 1837; Blechnum Sturm 1858.

(L) **cycadoides** (Pappe et Raws.) Kuhn, Fil. Afr. 91. 1868 — Natal. Madagascar.
Lomaria Pappe et Raws. Syn. Fil. Afr. austr. 28. 1858.

(L) **Dalgairnsiae** (Pappe et Raws.) Kuhn, Fil. Afr. 92. 1868 — Africa austr.
Lomaria Pappe et Raws. Syn. Fil. Afr. austr. 27. 1858.

(L) **danaeaceum** (Kze.) Christ, Prim. Fl. Costar. **3**. 22. 1901 — Mexico-Peru. Brasilia. — Lomaria Kze. Linn. **18**. 326. 1844. (HB. 179).

decurrens Roxb. Calc. Journ. **4**. 502. 1844 = ? B. orientale.

deflexum Diels, Engl. Jahrb. **29**. 198. 1900 = B. Faberi.

(L) **dendrophilum** (Sod.) C. Chr. Ind. 153. 1905 — Ecuador.
Lomaria Sod. Cr. vasc. quit. 118. 1893.

(L) **dentatum** (Kuhn) Diels in Schum. et Laut. Fl. deut. Südsee 130. 1901 — N. Hannover. — Lomaria Kuhn, Forschr. Gazelle **4** Farne 6. 1889.

denticulatum Sw. Syn. 113, 311. 1806 = B. serrulatum.

(L) **Deplanchei** Lenorm. msc. (Hk. Bak. Syn. 481. 1874) — N. Caledonia.
Lomaria Bak. Syn. 481. 1874; Spicanta O. Ktze. 821. 1891.

diplotaxicum Fée, Cr. vasc. Br. **1**. 25 t. 8 f. 1. 1869 = B. fraxineum.

(L) **discolor** (Forst.) Keys. Pol. Cyath. Hb. Bung. 66. 1873. Chr. 177. NPfl. 247 — Australia. N. Zealand. — Osmunda Forst. Prod. 78. 1786. Hemionitis Schkuhr 1806; Onoclea Sw. 1806; Gymnopteris Bernh. 1806; Lomaria Willd. 1810. HB. 175; Stegania A. Rich. 1832; Spicanta O. Ktze. 821. 1891. Stegania falcata R. Br. 1810; Lomaria Spr. 1827 — Onoclea nuda Lab. 1806; Lomaria Willd. 1810 (HB. 481); Stegania R. Br. 1810; Blechnum Mett.; Luerss. 1876.

distans Pr. Tent. 103. 1836; Epim. 105. 1849 = B. occidentale.

divergens Mett. Ann. sc. nat. V. **2**. 225. 1864 = ? B. Plumieri.

(L) **diversifolium** Mett. Ann. sc. nat. IV. **15**. 71. 1861. NPfl. 249 — N. Caledonia.
Lomaria Bak. 1867. HB. 181; Spicanta O. Ktze. 821. 1891.

doodioides Hk. Fl. bor. Am. **2**. 263. 1840 = B. spicant var.

drepanophyllum Trev. Atti Ist. Veneto II. **2**. 166. 1851 = B. occidentale.

(L) **durum** (Moore) C. Chr. Ind. 153. 1905 — Chatham Ins.
Lomaria Moore, Gard. Chr. **1866**. 290. HB. 177; Spicanta O. Ktze. 821. 1891; Lomaria rigida J. Sm. 1866.

(L) **eburneum** Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 233 c. fig. — China.

(L) **egenolfioides** (Bak.) C. Chr. Ind. 153. 1905 — Borneo.
Lomaria Bak. Kew Bull. **1894**. 7.

(B) **elongatum** Gaud. Freyc. Voy. Bot. 395. 1827 — Ins. Philipp. Polynesia. ? Brasilia. — Spicanta O. Ktze. 821. 1891; Blechnum nitidum Pr. 1825 (non 1822). HB. 184. Chr. 181. NPfl. 246; Blechnopsis Pr. 1849; Salpinchlaena Trev. 1869. — (an B. cartilagineum var.?).

elongatum Pr. Tent. 103. 1836 = B. orientale.

elongatum Mett. Fil. Lips. 61 t. 4 f. 18. 1856 = B. Patersoni*.

(L) **ensiforme** (Liebm.) C. Chr. Ind. 153. 1905 — Mexico.
Lomaria Liebm. Vid. Selsk. Skr. V. **1**. 234 (seors 82). 1849.

(B) **extensum** Fée, Gen. 75. 1850-52 — Brasilia.

(L) **Faberi** C. Chr. Ind. 153. 1905 — China.
Lomaria deflexa Bak. JoB. **1888**. 226 (non Col. 1844 nec Liebm. 1849); Spicanta O. Ktze. 1891; Blechnum Diels 1900.

falcatum Lodd.; Link, Hort. Berol. **2**. 77. 1833 = B. occidentale.

falciculatum Pr. Epim. 106. 1849 = B. occidentale.

(L) **falciforme** (Liebm.) C. Chr. Ind. 154. 1905 — Mexico.
Lomaria Liebm. Vid. Selsk. Skr. V. **1**. 234 (seors. 82). 1849.

(L) **Feei** (Jenm.) C. Chr. Ind. 154. 1905 — Jamaica.
Lomaria Jenm. Bull. Dep. Jam. I. n. 43. 7. 1893; L. Plumieri Fée 1866 (non Desv. 1811).

(B) **Fendleri** Hk. sp. **3**. 48 t. 158. 1860. HB. 185 — Venezuela. Brasilia.
Spicanta O. Ktze. 821. 1891.

(L) **filiforme** (A. Cunn.) Ettingsh. Denkschr. Ak. Wien **24**. 21 t. 6 f. 5. 1864 — N. Zealand. Fiji. ? N. Caledonia. — Lomaria A. Cunn. in Hk. Comp. Bot. Mag. **2**. 363. 1836. HB. 180; Spicanta O. Ktze. 821. 1891; Osmunda reptans Banks et Sol.; Forst. 1786 (nomen); Blechnum Luerss. 1871. Chr. 180. NPfl. 249. Stenochlaena heteromorpha J. Sm. 1841; Lomariopsis Moore 1857 — Stenochlaena feejeensis Brack. 1854.

(B) **Finlaysonianum** Wall. List n. 2172. 1829; Hk. et Grev. Ic. Fil. t. 225. 1831. HB. 187. Chr. 182. NPfl. 247 — Penang. Singapore. Borneo.
Blechnopsis Pr. 1849; Salpichlaena Fée 1850-52; Spicanta O. Ktze. 821. 1891.

flabellatum Pr. Tent. 103. 1836 = Actiniopteris australis.

(L) **Floresii** (Sod.) C. Chr. Ind. 154. 1905 — Ecuador.
Lomaria Sod. Cr. vasc. quit. 115. 1893.

fluminense Vell. Fl. flum. **11** t. 106. 1827; Arch. Mus. Nac. Rio Jan. **5**. 458. 1881 = B. brasiliense.

(L) **fluviatile** (R. Br.) Lowe; Salom. Nom. 115. 1883. NPfl. 249 — Australia. Tasmania. N. Zealand. — Stegania R. Br. Prod. Fl. N. Holl. 152. 1810; Lomaria Spr. 1827. HB. 181; Spicanta O. Ktze. 821. 1891; Lomaria rotundifolia Col. 1841, 1844; Raoul 1844 (non Bl. 1828).

Fontanesianum Gaud. Freyc. Voy. Bot. 397 t. 15. 1827 = Sadleria cyatheoides.

(L) **Fraseri** (A. Cunn.) Luerss. Flora **1876**. 292. Chr. 180. NPfl. 250 — N. Zealand. Ins. Philipp. — Lomaria A. Cunn. in Hk. Comp. Bot. Mag. **2**. 364. 1836. HB. 182; Spicanta O. Ktze. 821. 1891.

(B) **fraxineum** Willd. sp. **5**. 413. 1810 — Amer. trop.
Distaxia Pr. 1849; Spicanta O. Ktze. 821. 1891 — Blechnum Schlimense Fée 1857 — B. longifolium H.B. Willd. 1810 (non Cav. 1802). HB. 185. Chr. 183. NPfl. 247.

fraxinifolium Desv. Prod. 284. 1827 = B. fraxineum.

(L) **Fullagari** (F. Muell.) C. Chr. Ind. 154. 1905 — Ins. Lord Howe.
Lomaria F. Muell. Fragm. **8**. 157. 1874; L. auriculata Bak. 1874. HB. 481 (non Desv. 1811); Spicanta O. Ktze. 1891.

Galeottii Moore, Ind. 196. 1860 = B. occidentale.

Gayanum Sturm, Enum. pl. cr. vasc. chil. —. 1858 = B. penna marina.

(L) **Germainii** (Hk.) Christ, Bull. Boiss. II. **2**. 554. 1902 — Chile. Brasilia austr.
Lomaria Hk. sp. **3**. 32 t. 152. 1860. HB. 180; Spicanta O. Ktze. 821. 1891.

(L) **Ghiesbreghtii** (Bak.) C. Chr. Ind. 154. 1905 — Mexico.
Lomaria Gheisbreghtii Bak. Syn. 481. 1874; Spicanta O. Ktze. 821. 1891.

(L) **gibbum** (Lab.) Mett. Ann. sc. nat. IV. **15**. 68. 1861. Chr. 177. NPfl. 247 — N. Caledonia. Aneiteum.
Lomaria Lab. Sert. Austr.-Cal. 3 t. 4, 5. 1824. HB. 175.

giganteum Schlecht. Adumbr. 36 t. 20, 22 f. 1. 1827 = B. attenuatum.

(L) **Gilliesii** (Hk. et Grev.) Mett. Fil. Lips. 64. 1856 — Chile.
Lomaria Hk. et Grev. Ic. Fil. t. 207. 1831; Orthogramma Pr. 1849.

glabrum Roxb. Calc. Journ. **4**. 502. 1844 = Taenitis blechnoides.

(B) **glandulosum** Link, Enum. alt. 2. 462. 1822; ?Kze. Farnkr. 1. 132 t. 58 f. 2. 1843 — Amer. trop. — B. meridionale Pr. 1822 — B. pectinatum Hk. 1837.
glandulosum Wall. List n. 56. 1828 = B. occidentale.
(L) **Glaziovii** Christ, Ann. Jard. Cons. bot. Genève 3. 42. 1899 (syn.) — Brasilia. Lomaria Christ, l. c. 1899.
(B) **gracile** Klf. Enum. 158. 1824 (HB. 185) — Amer. trop.
Gueinzii Moore, Ind. 197. 1860 = B. tabulare.
Haenkeanum Trev. Atti Ist. Veneto II. 2. 166. 1851 = B. occidentale.
(L) **Hamiltonii** C. Chr. Ind. 155. 1905 — N. Zealand. — Lomaria intermedia Col. Tr. N. Zeal. Inst. 19. 274. 1887 (non Blechnum Link 1833).
(L) **Hancockii** Hance, JoB. 1883. 267 — Formosa. — Lomaria Bak. 1891; Spicanta O. Ktze. 822. 1891; Lomaria apodophylla Bak. 1885.
hastatum Klf. Enum. 161. 1824 = B. auriculatum.
helveolum Fée, Gen. 75. 1850-52 = B. blechnoides.
Herminieri vide L'Herminieri.
heterocarpum Fée, Gen. 74. 1850-52 = B. glandulosum.
heterophyllum Schlecht. Adumbr. 37. 1827 = B. attenuatum.
heterophyllum Opiz ? = B. spicant.
(L) **Hillii** C. Chr. Ind. 155. 1905 — N. Zealand. — Lomaria distans Col. Tr. N. Zeal. Inst. 28. 615. 1896 (non Blechnum Pr. 1836, 1849).
Houttuynii Poir. Enc. Suppl. 1. 644. 1810 = Woodwardia radicans var. orientalis.
(L) **Humblotii** C. Chr. Ind. 155. 1905 — Madagascar. — Lomaria stenophylla Bak. JoB. 1884. 142 (non Kl. 1847); Spicanta O. Ktze. 1891.
humile Salisb. Prod. 402. 1796 = B. occidentale.
imbricatum Bl. Enum. 198. 1828 = B. orientale.
(L) **imperiale** (Fée et Glaz.) Christ in Schwacke, Pl. Nov. Mineiras 2. 27. 1900 — Brasilia. — Lomaria Fée et Glaz.; Fée, Cr. vasc. Br. 1. 21 t. 7 f. 1. 1869.
(B) *impressum* Fée, Gen. 75. 1850-52 — Columbia.
indicum Burm. Fl. Ind. 231. 1768 = ? B. serrulatum.
(L) **inflexum** (Kze.) Kuhn, Fil. Afr. 92. 1868 — Natal. — Lomaria Kze. Linn. 18. 117. 1844. L. discolor var. natalensis Bak. 1874. HB. 481.
(B) *integerrimum* Spr. Syst. 4. 93. 1827 — Brasilia.
(B) **intermedium** Link, Hort. Berol. 2. 75. 1833. (HB. 185) — Amer. trop.
(B) **irregulare** Carr. in Seem. Fl. Vit. 351. 1873 — N. Caledonia.
japanense Moore, Ind. 198. 1860 = B. spicant.
japonicum L.fil. Suppl. 445. 1781 = Woodwardia.
japonicum Houtt. Nat. Hist. 14. t. 97 f. 1. 1783 = Woodwardia radicans [var. orientalis.
javanicum Bl. Enum. 197. 1828 = B. orientale.
Kaulfussianum Gaud. Voy. Bonite Bot. t. 78. 1846 = Sadleria cyatheoides.
(B) **laevigatum** Cav. Descr. 263. 1802. HB. 186 — N. S. Wales. Orthogramma Pr. 1849; Spicanta O. Ktze. 821. 1891 — Blechnum ambiguum Klf.; Pr. 1836; Parablechnum Pr. 1849; Lomaria Fée 1850-52; Blechnopteris Trev. 1851.
(B) **lanceola** Sw. Vet. Akad. Handl. 1817. 71 t. 3 f. 2. HB. 183. Chr. 183. NPfl. 246 — Amer. trop. — Spicanta O. Ktze. 821. 1891; Blechnum lanceolatum Raddi 1819 — Mesothema plantagineum Pr. 1849; Blechnum Hk. 1860; Struthiopteris Trev. 1869 — Blechnum trifoliatum Klf. 1824.
(L) **lanceolatum** (R. Br.) Sturm, Enum. pl. cr. chil. 25. 1858. Chr. 177. NPfl. 249 — Australia. Tasmania. N. Zealand. Polynesia. ? Chile.

Stegania R. Br. Prod. Fl. N. Holl. 152. 1810; Lomaria Spr. 1827. HB. 177; Spicanta O. Ktze. 821. 1891 — Lomaria aggregata Col. 1888.

lanceolatum Raddi, Opusc. sci. Bol. **3**. 294. 1819; Pl. Bras. **1**. 52 t. 60 f. 3. 1825 = B. lanceola.

lanuginosum Sturm, Enum. pl. cr. chil. —. 1858 = B. cycadifolium.

latifolium Pr. Tent. 103. 1836 = B. orientale.

latifolium Moritz, Bot. Zeit. **1854**. 855; Ettingsh. Farnkr. 119 t. 77 f. 5. 1865 = B. fraxineum.

(B) **Lechleri** Mett. Fil. Lechl. **2**. 17. 1859 — Peru. Brasilia.

(L) **Lehmannii** Hier. Engl. Jahrb. **34**. 473. 1904 — Columbia.

(L) **Lenormandi** (Bak.) Diels, Nat. Pfl. **1**[4]. 249. 1899 — N. Caledonia.
Lomaria Bak. Syn. 181. 1867; Spicanta O. Ktze. 821. 1891.

(L) **Leyboldtianum** (Phil.) C. Chr. Ind. 156. 1905 — Chile.
Lomaria Phil. (Anal. Univ. Chil. **1861**. 68); Linn. **33**. 303. 1864; Spicanta O. Ktze. 822. 1891.

(L) **L'Herminieri** (Bory) Mett. Fil. Lips. 64 t. 4 f. 14-15. 1856. Chr. 178. NPfl. 249 — Amer. trop. — Lomaria Bory; Kze. Farnkr. **1**. 173 t. 73. 1845. HB. 176; Lomaridium Pr. 1849; Spicanta O. Ktze. 821. 1891 — Lomaria exaltata Fée 1866.

(L) **lineatum** (Sw.) Hier. Engl. Jahrb. **34**. 473. 1904 — Amer. trop.
Osmunda Sw. Prod. 127. 1788; Onoclea Sw. 1801; Lomaria Willd. 1810 — L. arborescens Kl. et Karst. 1844 — L. Regnelliana Kze. 1847.

linguifolium Stokes (Bot. Mat. Med. **4** — 1812?) = Phyllitis scolopendrium.

lomarioides Gaud. Freyc. Voy. Bot. 396. 1827 = B. orientale.

lomarioides Mett. Fil. Lechl. **1**. 14. 1856 = B. valdiviense.

lomarioides Sod. Cr. vasc. quit. 128. 1893 = B. Sodiroi.

longifolium Cav. Descr. 263. 1802 = B. orientale.

longifolium H.B. Willd. sp. **5**. 413. 1810 = B. fraxineum.

lyratum Moritz, Bot. Zeit. **1854**. 855 (nomen) — Columbia.

macrophyllum Goldm. Nova Acta **19**. Suppl. I. 459. 1843; Trev. Atti Ist. Veneto II. **2**. 166. 1851 = B. orientale.

magellanicum Mett. Fil. Lechl. **1**. 14. 1856 = B. tabulare.

malaccense Fée, Gen. 74. 1850-52 = B. serrulatum.

marginatum Kuhn, Fil. Afr. 92. 1868 = B. Montbrisonis.

(B) **melanopus** Hk. sp. **3**. 64 t. 161. 1860. HB. 186. Chr. 183. NPfl. 247 — Khasia.
Blechnidium Moore, Ind. CLV. 1860(?); Spicanta O. Ktze. 821. 1891.

(L) **membranaceum** (Col.) Mett.; Diels, Nat. Pfl. **1**[4]. 249. 1899 — N. Zealand.
Lomaria Col.; Hk. sp. **3**. 34 t. 145. 1860. HB. 181; Spicanta O. Ktze. 821. 1891 — Lomaria oligoneuron Col. 1884.

meridense Kl. Linn. **20**. 349. 1847 = B. fraxineum.

(L) **meridense** (Kl.) Mett. Fil. Lips. 61. 1856 — Amer. trop.
Lomaria Kl. Linn. **20**. 345. 1847; L. vel Blechnum attenuatum auctt. quoad pl. amer. trop. HB. 176.

meridionale Pr. Del. Prag. **1**. 186. 1822 = B. glandulosum var.

(L) **microbasis** (Bak.) C. Chr. Ind. 156. 1905 — Madagascar.
Lomaria Bak. JoB. **1880**. 328; Spicanta O. Ktze. 822. 1891.

(L) **Milnei** (Carr.) C. Chr. Ind. 156. 1905 — Fiji.
Lomaria Carr. in Seem. Fl. Vit. 351. 1873.

minus Ettingsh. Denkschr. Ak. Wien **23**. 63 t. 8 f. 5, 12. 1864 = B. capense.

(B) **minutulum** Christ in Schwacke, Pl. Nov. Mineiras **2**. 27. 1900; Bull. Boiss. II. **2**. 552. 1902 — Brasilia.

moluccanum Desv. Berl. Mag. **5**. 325. 1811 = B. serrulatum.
moluccanum Roxb. Calc. Journ. **4**. 502. 1844 = B. orientale.
(L) **Montbrisonis** C. Chr. Ind. 157. 1905 — Réunion. — Lomaria marginata Fée, Gen. 71. 1850-52 (non Schrad. 1824); Blechnum Kuhn 1868.
(L) **Moorei** C. Chr. Ind. 157. 1905 — N. Caledonia.
Lomaria ciliata Moore, Gard. Chr. **1866**. 290. HB. 175; Spicanta O. Ktze. 1891; Blechnum Christ 177. 1897. NPfl. 247 (non Pr. 1825).
(B) **mucronatum** Fée, Cr. vasc. Br. **2**. 17. 1872-73 — Brasilia.
(L) **neocaledonicum** (Lind. & Fourn.) Salom. Nom. 117. 1883 — N. Caledonia.
Lomaria Lind. & Fourn. L'Ill. Hort. **23**. 132 t. 251. 1876.
(B) **nigro-squamatum** Gilbert, Bull. Torr. Cl. **24**. 258. 1897 — Bolivia.
(L) **nigrum** (Col.) Mett. Ann. sc. nat. IV. **15**. 69. 1861. NPfl. 249 — N. Zealand.
Lomaria Col. Tasm. Journ. **1**. 375. 1841; **2**. 176. 1844; Hk. Ic. pl. t. 960. 1854. HB. 181; Spicanta O. Ktze. 822. 1891.
nitidum Pr. Del. Prag. **1**. 187. 1822 = B. brasiliense.
nitidum Pr. Rel. Haenk. **1**. 49. 1825. HB. 184 = B. elongatum.
nitidum var. contracta Hk. sp. **3**. 44 t. 156. 1860 = B. cartilagineum var.
(L) **norfolkianum** (Hew.) C. Chr. Ind. 157. 1905 — Ins. Norfolk. N. Zealand. Samoa.
Lomaria Hew. Lond. JoB. **1**. 122. 1842. NPfl. 247; L. acuminata Bak. 1874. HB. 481 (non Desv. 1811); Spicanta O. Ktze. 1891.
nudum Mett.; Luerss. Flora **1876**. 292 = B. discolor var.
(L) **obtusatum** (Lab.) Mett. Ann. sc. nat. IV. **15**. 68. 1861 — N. Caledonia. N. Hebridae. — Lomaria Lab. Sert. Austr. Cal. 4 t. 6. 1824. HB. 178; Spicanta O. Ktze. 822. 1891.
obtusifolium Ettingsh. Denkschr. Ak. Wien **23**. 59. 9 t. 8 f. 4. 1864 = B. Schomburgkii.
(B) **occidentale** L. (sp. **2**. 1077. 1753; orientale ex err.); ed. II. **2**. 1524. 1763. HB. 185. Chr. 182. NPfl. 247 — Amer. trop.-Chile.
Spicanta O. Ktze. 822. 1891 — Blechnum caudatum Cav. 1802 — B. confluens Schlecht. et Cham. 1830. B. triangulare Link 1833. B. aduncum Liebm. 1849 — B. cognatum Pr. 1849 — Lomaria campylotis Kze. 1843; Mesothema Pr. 1849: Blechnum J. Sm. 1866.
onocleoides Sw. Schrad. Journ. **1800**^{2}. 75. 1801 = B. polypodioides.
(L) **opacum** Mett. Ann. sc. nat. IV. **15**. 69. 1861. NPfl. 247 — Polynesia.
Lomaria Bak. 1867. HB. 176; Spicanta O. Ktze. 821. 1891 — Lomaria emarginata Carr. 1873.
(B) **orientale** L. (sp. **2**. 1077. 1753; occidentale ex err.); ed. II. **2**. 1535. 1763. HB. 186. Chr. 182. NPfl. 247 — Asia, Austr., Polynesia trop.
Asplenium Bernh. 1802; Blechnopsis Pr. 1849; Salpichlaena Fée 1850-52; Spicanta O. Ktze. 821. 1891 — Blechnum longifolium Cav. 1802; Blechnopsis? Pr. 1849; Salpinchlaena Trev. 1869 — (conf. Blechnopsis et Salpichlaena).
(L) **ornifolium** (Pr.) Ettingsh. Denkschr. Ak. Wien **23**. 61 t. 8. f. 7. 1864 — Mexico-Peru-Brasilia. — Lomaria Pr. Rel. Haenk. **1**. 51. 1825 (HB. 179). L. longifolia Schlecht.; Mart. et Gal. 1842 (non Klf. 1824) — L. spectabilis Liebm. 1849.
pallidum Brack. Expl. Exp. **16**. 133. 1854 = Sadleria.
(L) **parvifolium** (Col.) C. Chr. Ind. 157. 1905 — N. Zealand.
Lomaria Col. Tr. N. Zeal. Inst. **20**. 224. 1888 — L. pumila Raoul, Ann. sc. nat. III. **2**. 115. 1844 (non Klf. 1824) HB. 178; Spicanta O. Ktze. 1891.

(B) **parvulum** Phil. Anal. Univ. chil. **1873**. 580 — Juan Fernandez.

(L) **Patersoni** (R. Br.) Mett. Fil. Lips. 64 t. 4 f. 4-10. Chr. 176. NPfl. 247 — Ins. Philipp. Australia. Tasmania. — *India austr.-Malesia-Fiji-N. Zealand. Stegania R. Br. Prod. Fl. N. Holl. 152. 1810; Lomaria Spr. 1827. HB. 174; Salpichlaena Fée 1850-52; Spicanta O. Ktze. 822. 1891. —* Lomaria elongata Bl. 1828; Blechnum Mett. 1856 (non Gaud. 1827); — L. punctata Bl. 1828 — L. Colensoi Hk.fil. 1844 — L. melanocaulon Brack. 1854 — L. coriacea Brack. 1854 (non Schrad. 1818).

pectinatum Pr. Rel. Haenk. **1**. 51. 1825 = B. orientale.

pectinatum Hk. Ic. pl. t. 95. 1837 = B. glandulosum var.

(L) **penna marina** (Poir.) Kuhn, Fil. Afr. 92. 1868. Chr. 180. NPfl. 249 — Regiones temp. austr. et antarct. — Polypodium Poir. Enc. **5**. 520. 1804; Lomaria Trev. 1869. Stegania alpina R. Br. 1810; Lomaria Spr. 1827. HB. 178; Blechnum Mett. 1856. Acrostichum polypodioides Thouars 1804 (polytrichoides) — Lomaria Gayana Fée 1853; Blechnum Sturm 1858 (Chile).

(L) **petiolare** (Sod.) C. Chr. Ind. 158. 1905 — Ecuador. Lomaria Sod. Cr. vasc. quit. 114. 1893.

plantagineum Hk. sp. **3**. 47. 1860 = B. lanceola.

(L) **Plumieri** (Desv.) Mett. Fil. Lips. 61 t. 4. f. 19-20. 1856. Chr. 178. NPfl. 248 — Amer. trop. — Lomaria Desv. Berl. Mag. **5**. 325. 1811. HB. 176; Lomaridium Pr. 1849; Spicanta O. Ktze. 822. 1891 — Lomaria divergens Kze. 1834; Blechnum Mett. 1864 — Lomaria mucronata Fée 1869.

Poeppigianum Sturm, Enum. pl. cr. chil. —. 1858 = B. penna marina.

Pohlianum Pr. Tent. 103 t. 11 f. 11. 1836 = B. blechnoides.

(L) **polypodioides** (Sw.) Kuhn, Fil. Afr. 92. 1868 (reduct.) — Amer. trop. Osmunda Sw. Prod. 127. 1788; Onoclea Sw. 1806 (non L. 1771); Lomaria Desv. 1827; Struthiopteris Trev. 1869; Spicanta O. Ktze. 821. 1891; Blechnum onocleoides Sw. 1801; NPfl. 249; Lomaria Spr. 1827. HB. 178; Spicanta Pr. 1849 — Lomaria angustifolia HBK. 1815; Blechnum Hier. 1904 (non Willd. 1810).

polypodioides Raddi, Opusc. sci. Bol. **3**. 294. 1819; Pl. Bras. **1**. 53 t. 60 f. 2. 1825 = B. blechnoides.

polystichoides Brack. Expl. Exp. **16**. 134. 1854 = Sadleria.

procerum Sw. Schrad. Journn. **1800**². 75. 1801 = B. capense.

productum Moritz, Bot. Zeit. **1854**. 855 (nomen) — Columbia.

(L) **pteropus** (Kze.) Mett. Fil. Lips. 61. 1856 — Brasilia. Lomaria Kze. Farnkr. **1**. 97 t. 46. 1842. Acrostichum heterophyllum Raddi 1825 (non L. 1753 nec Lomaria Desv. 1811).

(B) *pubescens* Desv. Prod. 284. 1827 — Amer. merid.

pubescens Hk. Ic. pl. t. 97. 1837 = B. auriculatum var.

(L) **punctulatum** Sw. Schrad. Journ. **1800**². 74. 1801. Chr. 178. NPfl. 249 — Afr. austr. — Lomaria Kze. 1836. HB. 179; Mesothema Pr. 1849; Blechnopteris Trev. 1851; Struthiopteris Trev. 1869; Spicanta O. Ktze. 822. 1891 — Lomaria auriculata Desv. 1811 — Blechnum Atherstoni Pappe et Raws. 1858 — Onychium Krebsii Kze. 1836; Scolopendrium Kze. 1844; Schismatosorus Kl. msc.; Lomaria J. Sm. 1875.

(L) **pygmaeum** (Col.) C. Chr. Ind. 158. 1905 — N. Zealand. Lomaria Col. Tr. N. Zeal. Inst. **25**. 322. 1893.

pyrophyllum Bl. Enum. 197. 1828 = B. orientale.

radiatum Pr. Tent. 103. 1836 = Actiniopteris australis.

radicans L. Mant. 307. 1771 = Woodwardia.
remotum Pr. Tent. 103. 1836 = B. auriculatum var.
reptans Luerss. Fil. Graeff. 142. 1871 = B. filiforme.
rigidum Sw. Schrad. Journ. **1800**². 75. 1801 = B. australe.
(L) **Rimbachii** (Sod.) C. Chr. Ind. 159. 1905 — Ecuador.
Lomaria Sod. Cr. vasc. quit. 117. 1893.
(B) **rugosum** Moore, Gard. Chr. n. s. **21**. 408. 1884 — ? (Hort.).
salicifolium Klf. Enum. 160. 1824 = B. orientale.
salicifolium Ettingsh. Denkschr. Ak. Wien **23**. 63 t. 7 f. 8. 1864 = B. tabulare.
(B) **scaberulum** Sod. Cr. vasc. quit. 130. 1893 — Ecuador.
scabrum Liebm. Vid. Selsk. Skr. V. **1**. 236 (seors. 84). 1849 = B. blechnoides.
scandens Bory, Dup. Voy. Bot. **1**. 272 t. 36. 1828 = B. volubile.
Schlimense Fée, 8 mém. 71. 1857 = B. fraxineum.
(L) **Schomburgkii** (Kl.) C. Chr. Ind. 159. 1905 — Amer. trop.
Lomaria Kl. Linn. **20**. 346. 1847 — L. obtusifolia Pr. 1836 (nomen); Christ, Bull. Boiss. II. **4**. 1091; Blechnum Ett. 1864. Lomaria Boryana vel tabularis vel magellanica auctt. quoad pl. amer. trop. HB. 180. — an adhuc: Lomaria Moritziana Kl. 1847 et L. aurata Fée 1857? — an L. rufa Spr. 1821. L. Ryani Klf. 1824? (Ind. occ.).
Schottianum Mett. Ann. sc. nat. IV. **15**. 71. 1861 = B. tabulare.
seminudum Willd. Phyt. 13 t. 8 f. 2. 1794 = Monogramma.
septentrionale Wallr. Fl. Cr. Germ. **1**. 24. 1831 = Asplenium.
(B) **serrulatum** Rich. Act. Soc. Hist. Nat. Paris **1**. 114. 1792. HB. 186. Chr. 182. NPfl. 247 — Amer., Asia, Austr. trop. N. Caledonia. — Blechnopsis Pr. 1849; Salpinchlaena Trev. 1869; Spicanta O. Ktze. 822. 1891. Blechnum denticulatum Sw. 1806 (patria err.); Blechnopsis Pr. 1849 — Blechnum striatum R. Br. 1810; Blechnopsis Pr. 1849; Salpinchlaena Trev. 1869 — Blechnum angustifolium Willd. 1810 — B. calophyllum Langsd. et Fisch. 1810 — B. stramineum Lab. 1824 — B. stagninum Raddi 1819 — ? B. indicum Burm. 1768 (an nomen optim.?).
(L) **sessilifolia** (Kl.) C. Chr. Ind. 159. 1905 — Costa Rica — ?
Lomaria Kl.; Christ, Bull. Boiss. II. **4**. 1092. 1904.
(L) **simillimum** (Bak.) Diels, Nat. Pfl. **1**⁴. 248. 1899 — Madagascar.
Lomaria Bak. JoB. **1884**. 141; Spicanta O. Ktze. 822. 1891.
(L) **sociale** Sod. Rec. 30. 1883. Chr. 179 — Ecuador. Brasilia.
Lomaria Sod. Cr. vasc. quit. 122. 1893.
(B) **Sodiroi** C. Chr. Ind. 159. 1905 — Ecuador. — B. lomarioides Sod. Cr. vasc. quit. 128. 1893 (non Gaud. 1827 vix Mett. 1856).
Souleytianum Gaud. Voy. Bonite Bot. t. 2. 134. 1846 = Sadleria.
(L) **spicant** (L.) Wither. Bot. Arr. ed. III. 765. 1796. (Sm. Mém. Ac. Turin **5**. 411. 1793). Chr. 179. NPfl. 249 — Europa. Madeira. Asia septent. Japonia. Amer. bor. pacif. — Osmunda L. sp. **2**. 1066. 1753; Struthiopteris Weis 1770; Acrostichum Willd. 1787; Onoclea Hoffm. 1795; Asplenium Bernh. 1799; Lomaria Desv. 1811. HB. 178. Osmunda borealis Salisb. 1796; Blechnum Sw. 1801; Lomaria Link. 1833; Spicanta Pr. 1849 — Acrostichum lineatum Cav. 1799. Lomaria crenata Pr. 1825 — Blechnum doodioides Hk. 1840; Struthiopteris Trev. 1869. (Amer. pacif.) — Lomaria niponica Kze. 1848 (Japonia).
spinulosum Poir. Enc. Suppl. **1**. 644. 1810 = Doodia caudata.

(L) **Sprucei** C. Chr. Ind. 160. 1905 — Ecuador. — Lomaria caudata Bak. Syn. 179. 1867; Spicanta O. Ktze. 1891 (non Blechnum Cav. 1802).

squamosum Stokes, Bot. Mat. Med. **4**. 617. 1812 = Ceterach officinarum.

squamulosum Klf. in Sieb. Fl. mixt. n. 242 = B. serrulatum.

(L) **squamulosum** (Desv.) Mett. Ann. sc. nat. V. **2**. 225. 1864 — Columbia. Lomaria Desv. Prod. 290. 1827. (HB. 179).

squarrosum Gaud. Voy. Bonite Bot. t. 2. 1846 = Sadleria polystichoides.

stagninum Raddi, Opusc. sci. Bol. **3**. 294. 1819; Pl. bras. **1**. 54 t. 62. 1825 = B. serrulatum.

stans Poir. Enc. Suppl. **1**. 644. 1810 = Woodwardia radicans.

(L) **stenophyllum** (Kl.) Mett. Fil. Lips. 64 t. 3 f. 10-11. 1856 — Amer. austr. Lomaria Kl. Linn. **20**. 346. 1847.

stenophyllum Fée, Gen. 74. 1850-52 = B. orientale.

stenopterum Hance, JoB. **1883**. 268 = Plagiogyria.

(L) **stipitellatum** (Sod.) C. Chr. Ind. 160. 1905 — Ecuador. Lomaria Sod. Rec. 29. 1883; Cr. vasc. quit. 121. 1893.

(L) **stoloniferum** Mett. (Fourn. Mex. pl. **1**. 113. 1872. syn.) — Mexico. Lomaria Fourn. Mex. pl. **1**. 113. 1872.

stramineum Lab. Sert. Austr. Cal. 2 t. 3. 1824 = B. serrulatum.

(L) **striatum** (Sw.) C. Chr. Ind. 160. 1905 — Amer. trop. Onoclea Sw. Syn. 304, 422. 1806; Lomaria Willd. 1810 — L. tuberculata J. Sm. 1856 —? L. brasiliensis Raddi 1825. L. procera pt. HB. 179.

striatum R. Br. Prod. Fl. N. Holl. 152. 1810 = B. serrulatum.

suburbicum Vell. Fl. flum. **11** t. 107. 1827; Arch. Mus. Nac. Rio Jan. **5**. 458. 1881 = B. glandulosum.

(L) **tabulare** (Thbg.) Kuhn, Fil. Afr. 94. 1868. Chr. 179. NPfl. 249 — Africa austr. cum ins. Amer. austr. temp. et antarct. — Pteris Thbg. Prod. Fl. cap. 171. 1800; Lomaria Mett.; Bak. 1870. Pteris osmundoides Bory 1804; Spicanta O. Ktze. 1891; Onoclea Boryana Sw. 1806; Lomaria Willd. 1810. HB. 180; Blechnum Schlecht. 1827 — Pteris palmaeformis Thouars 1804; Lomaria Desv. 1827; L. robusta Carm. 1818 (Tristan d'Acunha) — Parablechnum salicifolium Pr. 1849; Lomaria Fée 1850-52; Blechnum Ett. 1864; Lomaria Gueinzii Fée 1850-52; Blechnum Moore 1860 (Natal) — Lomaria magellanica Desv. 1811; Blechnum Mett. 1856 (Amer. antarct.).

triangulare Link, Hort. Berol. **2**. 76. 1833 = B. occidentale var.

triangulatum J. Sm. Cat. cult. ferns 38. 1857 = B. occidentale var.

tricuspe Klf.; Sieb. Syn. Fil. exs. n. 5 = ? B. australe.

trifoliatum Klf. Enum. 157. 1824 = B. lanceola.

trilobum Pr. Rel. Haenk. **1**. 50 t. 49 f. 2. 1825 = B. auriculatum var.

(L) **uliginosum** (Phil.) C. Chr. Ind. 160. 1905 — Chile. Lomaria Phil. Linn. **29**. 105. 1827.

(L) **unguiculatum** (Fée) Kuhn, Fil. Afr. 94. 1868 — Madagascar. Lomaria Fée, Gen. 70. 1850-52.

unilaterale Sw. Berl. Mag. **4**. 79 t. 3 f. 1. 1810 = B. blechnoides.

(L) **valdiviense** C. Chr. Ind. 160. 1905 — Chile. Juan Fernandez. Lomaria blechnoides Bory, Dup. Voy. Bot. **1**. 273. 1828. HB. 177 (non Desv. 1827); Blechnum Keys. 1873 (non supra); Spicanta O. Ktze. 1891; B. lomarioides Mett. 1856. NPfl. 249 (non Gaud. 1827) — Lomaria Lechleri Moore 1866 (non Blechnum Mett. 1856).

validum Fée, Gen. 74. 1850-52 = B. brasiliense.
(L) **varians** (Fourn.) C. Chr. Ind. 161. 1905 — Mexico.
Lomaria Fourn. Mex. pl. **1**. 113. 1872.
(L) **vestitum** (Bl.) Kuhn, Ann. Lugd. Bat. **4**. 284. 1869 — Malesia.
Lomaria Bl. Enum. 203. 1828. (HB. 179).
(B) *vestitum* Moore, Ind. 207. 1860 (descr.) — Venezuela.
(L) **Vieillardii** Mett. Ann. sc. nat. IV. **15**. 70. 1861. NPfl. 247 — N. Caledonia.
Lomaria Bak. 1867. HB. 175; Spicanta O. Ktze. 822. 1891.
(L) **violaceum** (Fée) Hier. Engl. Jahrb. **34**. 474. 1904 — Ind. occ. (Guadeloupe). Columbia. Costa Rica. — Lomaria Fée, 11 mém. 11 t. 5. 1866.
virginicum L. Mant. 307. 1771 = Woodwardia.
(B) **vittatum** Brack. Expl. Exp. **16**. 131 t. 16. 1854. HB. 482 — Fiji. ? N. Caledonia.
? Lomaria Fourn. 1873; Spicanta O. Ktze. 822. 1891.
(S) **volubile** Klf. Enum. 159. 1824. HB. 187. Chr. 183. NPfl. 247 — Amer. trop.
Salpichlaena J. Sm. 1841; Spicanta O. Ktze. 822. 1891. Blechnum scandens Bory 1828; Salpichlaena Pr. 1849 — Lomaria volubilis Hk. 1860. HB. 182; Spicanta Hookeriana O. Ktze. 1891.
(L) **vulcanicum** (Bl.) Kuhn, Ann. Lugd. Bat. **4**. 284. 1869. Chr. 179. NPfl. 248 — Java. Polynesia. Queensland. Tasmania. N. Zealand.
Lomaria Bl. Enum. 202. 1828. HB. 176; Spicanta O. Ktze. 822. 1891. Lomaria pilosa Brack. 1854.
(L) **Werckleanum** (Christ) C. Chr. Ind. 161. 1905 — Costa Rica.
Lomaria Christ, Bull. Boiss. II. **4**. 1091. 1904.
(B) **Whelani** Bailey, Rep. Govt. Sci. Exp. Bellenden-Ker 77. 1889; Queensl. Fl. Suppl. III. 92. 1890; Lithogr. t. 91 — Queensland.
(L) **xiphophyllum** (Bak.) C. Chr. Ind. 161. 1905 — Madagascar.
Lomaria Bak. JoB. **1884**. 142; Spicanta O. Ktze. 822. 1891.

BOLBITIS Schott, Gen. Fil. t. 14. 1834 = **Leptochilus.**
diversifolia Schott, Gen. ad t. 14. 1834.
flagellifera Schott, Gen. ad t. 14. 1834 = L. heteroclitus.
repanda Schott, Gen. ad t. 14. 1834 = L. cuspidatus.
serratifolia Schott, Gen. t. 14. 1834.
virens Schott, Gen. ad t. 14. 1834.

BOMMERIA Fournier in Baillon, Dict. de Bot. **1**. 448. 1876: Bull. Soc. Fr. **27**. 327. 1880 = **Gymnopteris.**
Ehrenbergiana Und. Bull. Torr. Cl. **29**. 633. 1902.
hispida Und. Bull. Torr. Cl. **29**. 633. 1902.
pedata Fourn. Bull. Soc. Fr. **27**. 327. 1880.
podophylla Fourn. Bull. Soc. Fr. **27**. 328. 1880 = G. Ehrenbergiana.
Schaffneri Fourn. Bull. Soc. Fr. **27**. 327. 1880 = G. hispida.

BOTRYCHIUM Swartz, Schrader's Journ. **1800**[2]. 8. 110. 1801 — (Ophioglossaceae Gen. **2**).
anthemoides Pr. Abh. böhm. Ges. V. **5**. 323. 1848 = B. virginianum.
australe R. Br. Prod. Fl. N. Holl. 164. 1810. NPfl. 471 — Australia. Tasmania. N. Zealand. Argentina.
Baeckeanum Brockmüller, Arch. Ver. Fr. Nat. Mecklb. **1863**. 170 = B. matricariae.
biforme Col. Tr. N. Zeal. Inst. **18**. 223. 1886 — N. Zealand.
biternatum (Sav.) Und. Bot. Gaz. **22**. 407 t. 21. 1896 — U. S. A. merid. Costa Rica.
Osmunda Sav. in Lam. Enc. **4**. 650. 1797; Botrypus lunarioides Michx. 1803; Botrychium Sw. 1806. NPfl. 471. B. fumarioides Willd. 1810.

boreale (Fr.) Milde, Bot. Zeit. **1857**. 478, 880. Chr. 365. NPfl. 470 — Europa, Asia bor. Alaska. — B. Lunaria v. boreale Fr. Herb. normale **16**. 85 —. — B. crassinervium Rupr.; Milde 1858.

brachystachys Kze. Linn. **18**. 305. 1844 = B. virginianum.

brevifolium Ångstr. Bot. Notis. **1866**. 40 = ? B. boreale.

Breynii Fr. Summ. Veg. **1**. 252. 1845 = B. matricariae.

chamaeconium Bitter et Hier. Nat. Pfl. **1**[4]. 471. 1900 — Kamerun.

charcoviense Portenschl.; Pr. Suppl. 47. 1845 = B. virginianum.

cicutarium Sw. Syn. 171. 1806 (Plum. t. 159) — San Domingo.

Coulteri Und. Bull. Torr. Cl. **25**. 537. 1898 — Rocky Mts.

crassinervium Rupr.; Milde, Nova Acta **26**. 763 t. 55 f. 10-11. 1858 = B. boreale var.

cuneatum Desv. Prod. 195. 1827 = B. obliquum.

daucifolium Wall.; Hk. et Grev. Ic. Fil. t. 161. 1829. HB. 448. Chr. 366. NPfl. 471 — India bor.-Ceylon-Java. — B. subcarnosum Wall. 1828 (nomen).

daucifolium *β* japonicum Prantl, Jahrb. Bot. Gart. Berl. **3**. 340. 1884 = B. japonicum.

decompositum Mart. et Gal. Mém. Ac. Brux. **15**. 15 t. 1. 1842 — Mexico.

dichronum Und. Bull. Torr. Cl. **30**. 45. 1903 — Jamaica.

dissectum Spr. Anleit. **3**. 172. 1804 — U. S. A. atlant. — Osmunda Poir. 1816.

erosum Milde, Bot. Zeit. **1864**. 102 = B. australe.

Fumariae Spr. Syst. **4**. 23. 1827 = B. biternatum.

fumarioides Willd. sp. **5**. 63. 1810 = B. biternatum.

gracile Pursh, Fl. Amer. sept. **2**. 656. 1814 = B. virginianum.

japonicum (Prantl) Und. Bull. Torr. Cl. **25**. 538. 1898 — Japonia.
B. daucifolium *β* japonicum Prantl, Jahrb. Bot. Gart. Berl. **3**. 340. 1884.

Jenmani Und. Fern Bull. **8**. 59. 1900 — Jamaica.

Kannenbergii Klinsm. Bot. Zeit. **1852**. 379 t. 6 = B. simplex.

lanceolatum (Gmel.) Ångstr. Bot. Notis. **1854**. 68. Chr. 366. NPfl. 470 — Europa, Asia, Amer. bor. et arct. Europa centr.
Osmunda Gmel. Nov. Comm. Ac. Petr. **12**. 516 t. 11 f. 2. 1768. Botrychium rutaceum *β* HB. 447. B. palmatum Pr. 1845.

lanceolatum Rupr. Distr. Cr. vasc. Ross. 33. 1845 = B. ramosum.

lanuginosum Wall. List n. 48. 1828; ? Hk. et Grev. Ic. Fil. t. 79. 1829. Chr. 366. NPfl. 471 — India. Ceylon. — B. virginianum *β* HB. 448.

lunaria (L.) Sw. Schrad. Journ. **1800**[2]. 110. 1801. HB. 447. Chr. 365. NPfl. 470 — Europa. Asia occ. et bor.-Himalaya. Amer. bor. Patagonia. Australia. Tasmania. N. Zealand. — Osmunda L. sp. **2**. 1064. 1753; Botrypus Rich. 1801.

Lunaria v. boreale Fr. Herb. normale **16**. 85 —. = B. boreale.

Lunaria v. cordatum Fr. Summ. Veg. **1**. 251. 1845 = B. simplex.

lunarioides Schkuhr, Kr. Gew. **1**. 158 t. 157. 1809 = B. obliquum.

lunaroides Sw. Syn. 172. 1806 = B. biternatum.

lunatum Gray, Nat. Arr. br. Pl. **2**. 19. 1821 = B. lunaria.

matricariae (Schrank) Spr. Syst. **4**. 23. 1827. NPfl. 471 — Europa, Asia, Amer. septentr. China austr. — Osmunda Schrank, Bair. Flora **2**. 419. 1789. Botrychium rutaceum Sw. 1801. B. matricarioides Willd. 1810; Osmunda Poir. 1816. B. rutaefolium A. Br. 1843. Chr. 366.

matricariaefolium A. Br.; Döll, Rhein. Flora 24. 1843 (syn.) = B. ramosum.

matricariaefolium Fr. Summ. Veg. **1**. 252. 1845 = B. lanceolatum.

matricarioides Willd. sp. **5**. 62. 1810 = B. matricariae.

millefolium Hochst.; Milde, Verhdl. zool. bot. Ges. Wien **19**. 158. 1869 = B. australe.

multifidum Trevis. Atti Soc. Ital. sci. nat. **17**. —. 1874 = B. matricariae.

neglectum Wood, Class-book ed. II. 816. 1860; Und. Bull. Torr. Cl. **30**. 1903 — Amer. bor. — B. matricariaefolium auctt. quoad pl. amer.

obliquum Mühl.; Willd. sp. **5**. 63. 1810. NPfl. 471 — Amer. bor. atlant. China merid. (Su-tchuen). — Osmunda Poir. 1816.

occidentale Und. Bull. Torr. Cl. **25**. 538. 1898 — Amer. bor. pacif.

onondagense Und. Bull. Torr. Cl. **30**. 47 f. 1-2. 1903 — New York.

palmatum Pr. Suppl. 43. 1845 = B. lanceolatum.

pumicola Coville; Und. Our Nat. Ferns ed. VI. 69. 1900; Coville, Bull. Torr. Cl. **28**. 109 t. 7. 1901 — Oregon.

pusillum Und. Bull. Torr. Cl. **30**. 50. 1903 — Mexico.

ramosum (Roth) Aschers. Fl. Brandenb. **1**. 906. 1864. NPfl. 470 — Europa.
Osmunda Roth, Tent. Fl. Germ. **1**. 444. 1788. Botrychium rutaceum Willd. 1810 (non Sw. 1801). HB. 447. (excl. β); Osmunda Poir. 1816. Botrychium matricariaefolium A. Br. 1843. Chr. 366.

Reuteri Payot, Cat. Foug. Montblanc 15. 1860 = B. ramosum.

robustum (Rupr.) Und. Bull. Torr. Cl. **30**. 51. 1903 — Kamtschatka. Unalaska. B. rutaefolium v. robustum Rupr.; Milde, Nova Acta **26**. 763 t. 55. f. 9. 1858.

rutaceum Sw. Schrad. Journ. **1800²**. 110. 1801 = B. matricariae.

rutaceum Willd. sp. **5**. 62. 1810. HB. 447 = B. ramosum.

rutaceum β tripartitum Ledeb. Fl. ross. **4**. 505. 1853 = B. lanceolatum.

rutaefolium A. Br. in Döll, Rhein. Flora 24. 1843 = B. matricariae.

rutaefolium v. robustum Rupr.; Milde, Nova Acta **26**. 763 t. 55 f. 9. 1858 [= B. robustum.

Schaffneri Und. Bull. Torr. Cl. **30**. 51. 1903 — Mexico.

silaifolium Pr. Rel. Haenk. **1**. 76. 1825. NPfl. 471 — Amer. bor. pacif.

silesiacum Kirschleger, Flora d'Alsace 401. 1855 = B. matricariae.

simplex Hitch. Amer. Journ. Sci. **6**. 103 t. 8. 1823. HB. 447. Chr. 366. NPfl. 470 — Amer. bor. Eur. bor. Japonia. — B. Kannenbergii Klinsm. 1852.

strictum Und. Bull. Torr. Cl. **30**. 52. 1903 — Japonia.

subbifoliatum Brack. Expl. Exp. **16**. 317 t. 44 f. 2. 1854. NPfl. 471 — Hawaii.

subcarnosum Wall. List n. 49. 1828 = B. daucifolium.

tenebrosum A. A. Eat. Fern Bull. **7**. 8. 1899 — U. S. A. atlant.

tenellum Ångstr. Bot. Notis. **1854**. 69 f. 1-4 = B. ramosum.

tenuifolium Und. Bull. Torr. Cl. **30**. 52. 1903 — U. S. A. merid. orient.

ternatum (Thbg.) Sw. Schrad. Journ. **1800²**. 111. 1801. HB. 449 (pt.). Chr. 366 (pt.). NPfl. 471 — Japonia. Himalaya. ? Argentina.
Osmunda Thbg. Fl. Jap. 329 t. 32. 1784.

ternatum var. daedaleum Christ, Prim. Fl. Costar. **3**. 46. 1901 = B. biternatum.

Underwoodianum Maxon. Bull. Torr. Cl. **32**. 220 t. 6. 1905 — Jamaica.

virginianum (L.) Sw. Schrad. Journ. **1800²**. 111. 1801. HB. 449 (excl. β). Chr. 366. NPfl. 472 — Amer. bor.-Ecuador-Brasilia. Europa. Asia temp.-Yunnan. — Osmunda L. sp. **2**. 1064. 1753. O. multifida Gmel. 1768. O. virginica Houtt. 1786; Botrypus Michx. 1803; Botrychium Willd. 1810 — B. gracile Pursh 1814 — B. anthemoides Pr. 1848.

virginianum Hk.fil. Fl. Tasm. **2**. 154 t. 169. 1850 = B. australe.

virginicum Willd. sp. **5**. 64. 1810 = B. virginianum.

zeylanicum Sw. Schrad. Journ. **1800²**. 111. 1801 = Helminthostachys.

BOTRYOGRAMME Fée, Gen. Fil. 166. 1850-52 = **Llavea.**
Karwinskii Fée, Gen. 166 t. 15 C. 1850-52 = L. cordifolia.
BOTRYOPTERIS Presl, Rel. Haenk. **1**. 76. 1825 = **Helminthostachys.**
crenata Pr. Abh. böhm. Ges. V. **5**. 324. 1848 } = H. zeylanica.
mexicana Pr. Rel. Haenk. **1**. 76 t. 12 f. 1. 1825 } = H. zeylanica.
BOTRYOTHALLUS Kotzsch, Bot. Zeit. **1846**. 104 = **Polybotrya.**
Kunzei Kl. Bot. Zeit. **1846**. 104 = P. serratifolia.
BOTRYPUS Richard, Cat. Hort. Med. Paris 120. 1801 = **Botrychium.**
Lunaria Rich. Cat. Ht. Med. Paris 120. 1801.
lunarioides Michx. Fl. bor. Am. **2**. 274. 1803 = B. biternatum.
virginicus Michx. Fl. bor. Am. **2**. 274. 1803 = B. virginianum.
BOWRINGIA Hooker, Journ. Bot. **5**. 237. 1853 = **Brainea.**
insignis Hk. JoB. **5**. 237 t. 2. 1853.
BRACHYSORUS Presl, Epim. Bot. 70. 1849 = **Athyrium.**
woodwardioides Pr. Epim. 70. 1849 = A. silvaticum.
BRAINEA J. Smith, Catal. Kew Ferns 5. 1856 — (Polypodiaceae Gen. **53**).
insignis (Hk.) J. Sm. Cat. Kew Ferns 5. 1856. HB. 390. Chr. 185. NPfl. 251 — China austr.-Siam. — Bowringia Hk. JoB. **5**. 237 t. 2. 1853.
CAENOPTERIS Bergius, Acta Acad. Petrop. **1782²**. 249. 1786 = **Asplenium.**
achilleaefolia Mart. et Gal. Mém. Ac. Brux. **15**. 63 t. 16. 1842 = Athyrium.
appendiculata Labill. Nov. Holl. Pl. Spec. **2**. 94 t. 243. 1806 = A. bulbiferum var.
auriculata Thbg. Nova Acta Petr. **9**. 159 t. E. f. 2. 1795.
aspidioides Desv. Prod. 268. 1827 = Athyrium.
bulbifera Desv. Prod. 268. 1827.
canariensis Willd. Phyt. **1**. 14. 1794 = Davallia.
cicutaria Thbg. Nova Acta Petr. **9**. 158 t. C. f. 1. 1795 = A. cristatum.
cuneata Desv. Prod. 267. 1827 = A. affine.
daucifolia Desv. Prod. 268. 1827.
disticha Spr. Syst. **4**. 91. 1827 = ? A. achilleifolium.
Fabiana Bory in Willd. sp. **5**. 299. 1810 (syn.); Spr. Syst. **4**. 91. 1827.
flaccida Thbg. Nova Acta Petr. **9**. 158 t. D. f. 1-2. 1795.
foeniculacea Desv. Prod. 268. 1827 = A. fragrans var.
fumarioides Desv. Prod. 268. 1827 = Odontosoria meifolia.
furcata Berg. Acta Petr. **1782²**. 249 t. 7 f. 1. 1786 = A. achilleifolium.
graminea Schkuhr, Kr. Gew. **1**. t. 87. 1809 = Monogramma.
inaequalis Bory in Willd. sp. **5**. 298. 1810 (syn.); Spr. Syst. **4**. 91. 1827 = A. lineatum var.
japonica Willd. Phytogr. 14 t. 8 f. 1. 1794 = Microlepia speluncae.
japonica Thbg. Nova Acta Petr. **9**. 161 t. G. f. 2. 1795 = Onychium.
myriophylla Sw. Schrad. Journ. **1800²**. 60. 1801.
novae-zeelandiae Spr. Schrad. Journ. **1799²**. 269 = A. flaccidum.
odontites Thbg. Nova Acta Petr. **9**. 159 t. E. f. 1. 1795 = A. flaccidum.
palmata Spr. Syst. **4**. 91. 1827 = A. subdigitatum.
quadripinnata Blanco, Fl. Filip. 833. 1837 — Ins. Philipp.
rhizophylla Thbg. Sm. Pl. Ic. ined. **2** t. 50. 1790 = A. myriophyllum.
thalictroides Loud. Hort. Brit. Suppl. 504. 1850 = ? A. cristatum.
triloba Desv. Prod. 267. 1827 = A. cristatum.
violascens Boj. Hort. Maur. 394. 1837 = A. lineatum var.
viridans Spr. Syst. **4**. 91. 1827.
vivipara Berg. Acta Petr. **1782²**. 250 t. 7 f. 3. 1786.

Cafraria Presl. Epim. Bot. 166. 1849 *(Stenochlaena §)* = Stenochlaena.
CALAMISTRUM (L.) O. Kuntze, Revisio Gen. Plant. **2**. 822. 1891 = **Pilularia.**
americanum - globuliferum - Mandonii - minutum - Novehollandiae - Novezelandiae O. Ktze. Rev. Gen. Pl. **2**. 822. 1891 = P. sp. hymonym.
Callipteridastrum Fée, Gen. Fil. 213. 1850-52 *(Diplazium §)* = Diplazium.
CALLIPTERIS Bory, Voyage **1**. 282. 1804 = **Diplazium.**
accedens J. Sm. JoB. **4**. 179. 1841 = D. proliferum.
alismaefolia J. Sm. JoB. **3**. 409. 1841 = D. Cumingii.
ambigua Moore, Ind. LV. 1857 = D. esculentum.
arborescens Bory, Voy. **1**. 283. 1804.
attenuata Pr. Epim. 89. 1849 = D. proliferum.
castaneifolium Bory, Voy. **1**. 282. 1804.
elegans J. Sm. JoB. **3**. 409. 1841 = D. fraxinifolium.
esculenta J. Sm.; Moore et Houlst. Gard. Mag. Bot. **3**. 265. 1851.
ferox J. Sm. Bonpl. **9**. 262. 1861 = ? D. proliferum.
fraxinifolia J. Sm.; Moore, Ind. 217. 1861.
heterophylla Moore, Ind. 217. 1861 = Angiopteris evecta.
malabarica J. Sm. JoB. **3**. 409. 1841 = D. esculentum.
microphylla Fée, Gen. 219. 1850-52.
ovata J. Sm. JoB. **3**. 409. 1841 = D. cordifolium.
paradoxa Moore, Ind. 217. 1861.
pinnatifida Fée, Gen. 219. 1850-52.
prolifera Bory, Voy. **1**. 283. 1804.
repanda Pr. Epim. 260. 1849 = D. proliferum.
Serampurensis Fée, Gen. 219. 1850-52 = D. esculentum.
serrulata Fée, Gen. 219. 1850-52 = D. manilense.
Smithiana Bedd. Ferns br. Ind. t. 332. 1870.
spinulosa J. Sm. JoB. **3**. 409. 1841 = D. proliferum.
Swartzii Pr. Epim. 260. 1849 = D. proliferum.
sylvatica Bory, Voy. **1**. 282. 1804 = D. silvaticum.
ternata Moore, Ind. 218. 1861.
undulosa Pr. Epim. 90. 1849.
vittaeformis J. Sm. JoB. **3**. 409. 1841 = Syngramma.
Wallichii J. Sm. JoB. **4**. 179. 1841 = D. esculentum.
Zollingeri Fée, Gen. 219. 1850-52 = D. cordifolium.
CALLOGRAMME Fée, Gen. Fil. 169. 1850-52 = **Syngramma.**
Caeciliae Fée, Gen. 169 t. 15 A. 1850-52; 7 mém. 41. t. 18 f. 1 = S. alismifolia.
Calomelanos Presl, Tent. 218. 1836 *(Gymnogramma §)* = Ceropteris.
CALYMELLA Presl, Tent. Pterid. 48. 1836 = **Gleichenia.**
alpina Pr. Tent. 49. 1836.
dicarpa Pr. Abh. böhm. Ges. V. **5**. 338. 1848.
microphylla Pr. Tent. 49 t. 1 f. 3. 1836 = G. circinnata.
vulcanica Pr. Abh. böhm. Ges. V. **5**. 338. 1848.
CALYMMODON Presl, Tent. Pterid. 203. 1836 = **Polypodium.**
clavifer Moore, Ind. 219. 1861.
cucullatus Pr. Tent. 204. 1836.
? denticulatus Moore, Ind. LXIII. 1857.
hirtum Brack. Expl. Exp. **16**. 2. 1854 = P. cucullatum.
CALYPTERIUM Bernhardi, Schrad. Journ. **1801**[1]. 22. 1802; neu. Journ. 1[2]. 25. 1806 = **Onoclea.**
sensibile Bernh. Schrad. Journ. **1801**[1]. 22 t. 1 f. 7-11. 1802.

CAMPIUM Presl, Tent. Pterid. 238. 1836 = **Leptochilus.**
costatum Pr. Tent. 239 t. 10. f. 23. 1836 = L. scalpturatus.
crispatulum Pr. Epim. 171. 1849 = L. virens var.
proliferum Pr. Epim. 170. 1849 = L. subcrenatus.
punctulatum Pr. Tent. 239 t. 10 f. 22. 1836 = L. Preslianus.
repandum Pr. Tent. 239. 1836 = L. cuspidatus.
subcrenatum Pr. Tent. 239. 1836.
undulatum Pr. Epim. 170. 1849 = L. scalpturatus var.
virens Pr. Tent. 239. 1836.
CAMPTERIA Presl, Tent. Pterid. 146. 1836 = **Pteris.**
anamallayensis Bedd. Ferns S. Ind. t. 45. 1863 = P. geminata.
Balbisiana Pr. Tent. 147. 1836 —? — Litobrochia Fée 1850-52.
biaurita Hk. Gen. t. 65. A. 1841.
Galeottii Moore, Ind. 220. 1861 = P. biaurita.
Gardneri Moore, Ind. XLIII. 1857.
geminata J. Sm. Hist. Fil. 294. 1875.
heterophlebia Moore, Ind. XLIII. 1857.
Kleiniana Pr. Tent. 147 t. 5. f. 19. 1836 = P. geminata.
laurea Moore, Ind. 220. 1861.
Luschnathiana Kl. Linn. **14**. 286. 1840 (nomen).
nemoralis J. Sm. Bot. Mag. **72** Comp. 23. 1846 = P. biaurita.
pectinata Presl, Tent. 147. 1836 = P. Wallichiana.
Pseudo-Lonchitis Pr. Tent. 147. 1836 = P. tripartita var.
Rottleriana Pr. Tent. 147. 1836 = P. biaurita.
triplicata Moore, Ind. 221. 1861 = P. heteroclita.
Wallichiana Moore, Ind. 221. 1861.
Wightiana Pr. Tent. 147. 1836 = P. biaurita.
CAMPTODIUM Fée, Gen. Fil. 298. 1850-52 = **Dryopteris.**
pedatum Fée, Gen. 298. 1850-52.
CAMPTOSORUS Link, Hort. Berol. **2**. 69. 1833 — (Polypodiaceae Gen. 47).
rhizophyllus (L.) Link, Hort. Berol. **2**. 69. 1833 — Amer. bor. atlant.
Asplenium L. sp. **2**. 1078. 1753; ed. II. **2**. 1536 (non 1540); Scolopendrium Endl. 1840. HB. 248. Chr. 213. NPfl. 232; Antigramma J. Sm. 1841; Phyllitis O. Ktze. 1891.
rumicifolius Link, Fil. sp. 83. 1841 = Phyllitis plantaginea.
sibiricus Rupr. Distr. Crypt. vasc. Ross. 45. 1845 — Sibiria. China. Japonia.
Scolopendrium Hk. 1860. HB. 248. Chr. 213. NPfl. 231; Antigramma J. Sm. 1875; Phyllitis O. Ktze 1891.
CAMPYLONEURUM Presl, Tent. Pterid. 189. 1836 = **Polypodium.**
(Campylonevron Fée).
acrocarpon Fée, Cr. vasc. Br. **1**. 115 t. 35 f. 3. 1869 = P. laevigatum.
amphostenon Fée, Gen. 258. 1850-52 = P. angustifolium var.
angustifolium Fée, Gen. 257. 1850-52.
aphanophlebium Moore, Ind. 223. 1861 = P. laevigatum.
brevifolium Link, Fil. sp. 124. 1841 = P. repens.
caespitosum Link, Fil. sp. 125. 1841 = P. repens.
caudatum Fée, 8 mém. 96. 1857 — Mexico. Cuba. (an P. phyllitidis var.?).
chrysopodum Fée, Gen. 258. 1850-52 = P. repens.
coarctatum Fée, Gen. 258. 1850-52.
costatum Pr. Tent. 190. 1836.

crispum Fée, Gen. 259. 1850-52 = P. nitidum.
cubense Fée, Gen. 259. 1850-52; 6 mém. 14 t. 3 f. 2 = P. vexatum.
decumanum Fée, Cr. vasc. Br. **1**. 115. 1869.
decurrens Pr. Tent. 190. 1836.
difforme Moore, Ind. 224. 1861.
dimorphum Fourn. Mex. pl. **1**. 84. 1872 = P. angustifolium.
ensifolium J. Sm. Cat. cult. ferns 12. 1857 = P. angustifolium var.
fallax Fée, Cr. vasc. Br. **1**. 114 t. 35 f. 2. 1869 = P. leuconeuron.
fasciale Pr. Tent. 190. 1836 = P. laevigatum.
Fendleri Moore, Ind. 224. 1861. (descr.) = P. laevigatum.
Fendleri J. Sm. Hist. Fil. 97. 1875.
immersum J. Sm. Bot. Voy. Herald 231. 1854 = P. costatum.
jalapense Fourn. Mex. pl. **1**. 85. 1872 = P. xalapense.
Jamesoni Fée, Gen. 259. 1850-52; 6 mém. 14 t. 2 f. 5. 1853 = P. angustifolium var.
juglandifolium Fée, Cr. vasc. Br. **1**. 117 t. 45 f. 1. 1869 = P. decurrens.
laevigatum Pr. Tent. 190. 1836.
lanciforme Pr. Tent. 190. 1836 = P. laevigatum.
latum Moore, Ind. 225. 1861.
leuconevron Fée, Cr. vasc. Br. **1**. 103 t. 35 f. 1. 1869.
leucorhizon Fée, Gen. 258. 1850-52 = P. angustifolium.
loreum Fée, 8 mém. 129. 1857 = P. angustifolium.
lucidum Moore, Ind. 225. 1861 = P. angustifolium*.
macrosorum Fée, 8 mém. 96. 1857 = P. phyllitidis.
magnificum Moore, Ind. 226. 1861 = P Fendleri.
minus Fée, Gen. 258. 1850-52; 7 mém. 64 t. 24 f. 3 = P. laevigatum.
Moritzianum Fée, Gen. 258. 1850-52 = P. phyllitidis.
nitidum Pr. Tent. 190. 1836.
nodosum Kl.; Fée, Gen. 258. 1850-52 = P. angustifolium.
oligophlebium Fée, Gen. 258. 1850-52 = P. repens.
ophiocaulon Fée, Gen. 258. 1850-52.
Phyllitidis Pr. Tent. 190 t. 7 f. 18-20. 1836.
polyanthum Pr. Tent. 190 t. 7 f. 16. 1836 = P. nitidum.
repens Pr. Tent. 190. 1836.
rigidum J. Sm. Cat. Kew Ferns 2. 1856 = P. angustifolium*.
Sieberianum Pr. Tent. 190. 1836 = P. phyllitidis.
solutum Fée, Gen. 258. 1850-52 = P. angustifolium.
sphenodes Fée, Gen. 258. 1850-52.
taeniosum Fée, Gen. 257. 1850-52 = P. angustifolium.
undulatum Pr. Tent. 190. 1836 = P. phyllitidis.
xalapense Fée, Gen. 258. 1836.

CANDOLLEA Mirbel Hist. nat. Vég. **5**. 86. 1803; Hist. nat. Plant. **4**. 69. 1803 = **Cyclophorus.**
heterophylla Mirb. Hist. nat. Vég. **5**. 87. 1803 = C. spissus.
lanceolata Mirb. l. c. 89 = ? C. acrostichoides.
longifolia Mirb. l. c. 88.
polypodioides Mirb. l. c. 89 = Polypodium.

Cardinitis J. Smith, Hist. Fil. 151. 1875 *(Hemionitis §)* = Hemionitis.

CARDIOCHLAENA Fée, Gen. Fil. 314. 1850-52 = **Aspidium.**
alata Fée, Gen. 315. 1850-52 = A. decurrens.
ampla Fée, Gen. 315. 1850-52 — Guadeloupe.
Sagenia Moore 1861 — (prob. = A. martinicense.

confluens Fée, Gen. 316. 1850-52 — Guadeloupe.
Sagenia Moore 1861 — (prob. = A. martinicense).
?fimbriata Fée, 11 mém. 91. 1866 = Aspidium fimbriatum Willd.
laevis Fée, Gen. 316. 1850-52 = A. grandifolium.
macrophylla Fée, Gen. 315. 1850-52 = A. martinicense.
Menyanthidis Fée, Gen. 315. 1850-52.
?*pentaphylla* Fée, 11 mém. 91. 1866 (Plum. t. 91) — S. Domingo.
pilosa Fée, 10 mém. 45. 1865; Cr. vasc. Br. **1**. 45 t. 10 f. 4. 1869 = A. martinicense.
sinuosa Fée, Gen. 316. 1850-52 = A. grandifolium.
subpinnatifida Fée, Gen. 315. 1850-52 = A. melanocaulon.
trilobata Fée, Gen. 316. 1850-52 = A. Plumieri.
CARDIOMANES Presl, Hymenoph. 12. 1843 = **Trichomanes.**
reniforme Pr. Hym. 12. 1843.
Cardiostegia Moore, Ind. XC. 1857 (*Nephrolepis* §) = Nephrolepis.
CARPANTHUS Rafinesque, New York Med. Repository II. **5**. 356. 1808; Journ. de Bot. **1**. 221. 1808 = **Azolla.**
axillaris Rafin. l. c. 1808; Desv. Prod. 376. 1827 = ? A. caroliniana.
CASSEBEERA Kaulfuss, Enum. Fil. 216. 1824 (Cassebeeria auctt.).
argentea J. Sm. JoB. **4**. 159. 1841 = Cheilanthes.
auriculata J. Sm. Bot. Mag. **72**. Comp. 20. 1846 = Pellaea.
cuneata J. Sm. JoB. **4**. 159. 1841 = Cheilanthes angustifolia var.
farinosa J. Sm. JoB. **4**. 159. 1841 = Cheilanthes.
gleichenioides Gardn. in Hk. Ic. pl. t. 507. 1843 = Pellaea.
hastata J. Sm. JoB. **4**. 159. 1841 = Pellaea viridis.
inframarginalis J. Sm. JoB. **4**. 159. 1841 = Pellaea intramarginalis.
microphylla J. Sm. JoB. **4**. 159. 1841 = Cheilanthes.
microphylla Christ in Schwacke, Pl. Nov. Mineiras **2**. 25. 1900 = Doryopteris.
paradoxa Fée, 7 mém. 30 t. 20 f. 2. 1857 = Doryopteris.
pedata J. Sm. JoB. **3**. 404. 1841 = Doryopteris.
pedatifida Christ in Schwacke, Pl. Nov. Mineiras **2**. 25. 1900 = Doryopteris.
petiolata Fée, 7 mém. 30 t. 12 f. 4. 1857 } = Pellaea pinnata.
pinnata Klf. Enum. 217 t. 1 f. 11. 1824 }
pteroides Pr. Tent. 155 t. 6 f. 7. 1836 = Pellaea.
tenuifolia J. Sm. JoB. **4**. 159. 1841 = Cheilanthes.
triphylla Klf. Enum. 216. 1824 = Doryopteris.
CASSIOPTERIS Karsten msc. (Kl. Linn. **20**. 437. 1847) = **Ophioglossum.**
Catapodium J. Smith, Cat. cult. ferns 16. 1857 (*Phegopteris* §) = Dryopteris.
CATENULARIA Zippel msc. (Mett. Polyp. 37. 1857) = **Polypodium.**
(CELANTHERA Thouin, Mém. Act. sci. Paris — 1786 = **Marattia.**)
CEPHALOMANES Presl, Hymenoph. 17. 1843 = **Trichomanes.**
alatum Pr. Abh. böhm. Ges. V. **5**. 334. 1848 = T. javanicum var.
asplenioides Pr. Abh. böhm. Ges. V. **5**. 334. 1848 = T. javanicum.
atrovirens Pr. Hym. 18 t. 5. 1843 = T. javanicum var.
auriculatum v.d.B. Ned. kr. Arch. **4**. 352. 1859.
Australicum v.d.B. Ned. kr. Arch. **5**². 139. 1861 = T. javanicum var.
Boryanum v.d.B. Ned. kr. Arch. **4**. 351. 1859 = T. javanicum var.
curvatum v.d.B. Ned. kr. Arch. **4**. 351. 1859 = T. javanicum var.
dissectum v.d.B. Ned. kr. Arch. **4**. 352. 1859 = T. auriculatum.
javanicum Pr. Abh. böhm. Ges. V. **5**. 334. 1848.
Madagascariense v.d.B. Ned. kr. Arch. **4**. 351. 1859.

oblongifolium Pr. Abh. böhm. Ges. V. **5**. 334. 1848; Epim. 19 t. 10. 1849 = T. javanicum.

rhomboideum v.d.B. Ned. kr. Arch. **4**. 350. 1859 = T. javanicum var.

Singaporianum v.d.B. Ned. kr. Arch. **4**. 351. 1859 = T. javanicum var.

Wilkesii v.d.B. Ned. kr. Arch. **5**². 140. 1861 = T. javanicum var.

Zollingeri v.d.B. (Pl. Jungh. **1**. 552. 1856); Ned. kr. Arch. **4**. 351. 1859 = T. javanicum var.

Cephalosorium Moore, Ind. Fil. LXXV. 1857 (*Campyloneurum §*) = Polypodium.

CERATODACTYLIS J. Smith in Hook. Gen. Fil. t. 36. 1839 = **Llavea.**

osmundioides J. Sm. in Hk. Gen. t. 36. 1839 = L. cordifolia.

CERATOPTERIS Brongniart, Bull. Soc. Philom. **1821**. 186 — (Parkeriaceae).

cornuta Le Prieur, Ann. sc. nat. **19**. 103 t. 4 A. 1830 = C. thalictroides.

Gaudichaudii Brongn. Bull. Soc. Philom. **1821**. 187 = C. thalictroides.

Lockharti Kze. Linn. **23**. 241. 1850 = C. thalictroides.

Parkeri J. Sm. JoB. **4**. 70. 1841 = C. thalictroides.

Richardi Brongn. Dict. Class. d'Hist. Nat. **3**. 351. 1823 = C. thalictroides.

thalictroides (L.) Brongn. Bull. Soc. Philom. **1821**. 186 c. tab. HB. 174. Chr. 357. NPfl. 342 — Trop. et Subtrop.

Acrostichum L. sp. **2**. 1070. 1753; Pteris Sw. 1801; Teleozoma R.Br. 1823; Furcaria Desv. 1827; (Cladostachys Wall. hb.). Acrostichum siliquosum L. 1753; Belvisia Mirb. 1803. Pteris cornuta Pal. Beauv. 1807; Ellobocarpus Klf. 1824; Furcaria Desv. 1827; Ceratopteris Le Prieur 1830; Onychium Hassk. 1843. Ellobocarpus oleraceus Klf. 1824 — Cryptogenis ferulacea Rich. msc.; Pteris Rich.; Klf. 1824 — Parkeria pteridioides Hk. 1825; Ceratopteris Parkeri J. Sm. 1841.

Cerogramme Diels, Nat. Pflanzenfam. **1**⁴. 260. 1899 (*Gymnogramme §*) = Gymnogramma.

CEROPTERIS Link, Fil. sp. 141. 1841 — (Polypodiaceae Gen. **65**).

argentea Kuhn, v. Deck. Reisen **3**³. Bot. 18. 1879 = Gymnogramma.

calomelaena Link, Fil. sp. 141. 1841 = C. calomelanos.

calomelanos (L.) Und. Bull. Torr. Cl. **29**. 632. 1902 — Amer. trop. Afr. occ. Natal

Acrostichum L. sp. **2**. 1072. 1753; Gymnogramma Klf. 1824. HB. 385 Chr. 70; Neurogramme Diels 1899. NPfl. 264; Ceropteris calomelaena Link 1841 — C. Herminieri Link 1841; Gymnogramme Bory; Kze. 1850 — Ceropteris Martensii Link 1841; Gymnogramme Bory; Kze. 1850 — Ceropteris Massoni Link 1841; Gymnogramme Loud.; Kze. 1850 — Ceropteris intermedia Fée 1866 — Acrostichum chrysophyllum Sw. 1801 Gymnogramma Klf. 1824; Ceropteris Link 1841.

chrysophylla Link, Fil. sp. 143. 1841 = C. calomelanos var.

distans Link, Fil. sp. 142. 1841 = C. calomelanos.

Herminieri Link, Fil. sp. 164. 1841 = C. calomelanos var.

intermedia Fée, 11 mém. 30. 1866 = C. calomelanos var.

Martensii Link, Fil. sp. 143. 1841 = C. calomelanos var.

Massoni Link, Fil. sp. 143. 1841 = C. calomelanos var.

monosticha Fée, 7 mém. 44 t. 22 f. 2. 1857 = Notholaena sulphurea.

obtusa Fée, 8 mém. 80. 1857 — Columbia. — Gymnogramma Moore 1861.

pallescens Fée, 9 mém. 14. 1857 = C. tartarea.

peruviana (Desv.) Link, Fil. sp. 142. 1841 — Mexico-Peru.

Gymnogramma Desv. Berl. Mag. **5**. 329. 1811; Neurogramme Hier. 1904; Gymnogramme calomelanos β HB. 385 — Ceropteris Schaffneri Fée 1857 Gymnogramma Moore 1861.

plicata Fée, 8 mém. 81. 1857 = C. peruviana.
Schaffneri Fée, 8 mém. 80. 1857 = C. peruviana.
serrata Fée, 8 mém. 81. 1857 = C. calomelanos.
sulfurea Fée, Gen. 183. 1850-52 = Gymnogramma sulphurea.
tartarea (Cav.) Link, Fil. sp. 142. 1841 — Amer. trop. ? Samoa.
Acrostichum Cav. Descr. 242. 1802; Gymnogramma Desv. 1811. HB. 384; Neurogramme Diels 1899. NPfl. 264; Hemionitis dealbata Willd. 1809; Gymnogramma Pr. 1825 — G. ochracea Pr. 1825 — G. ornithopteris Kl. 1847. Allosorus farinosus Kze. 1848 — Gymnogramme adiantoides Karst.; Mett. 1864 — G. Brackenridgei Carr. 1873 (Samoa) — G. Stelzneri Koch 1859.
triangularis (Klf.) Und. Bull. Torr. Cl. **29**. 630. 1902 — Amer. bor. pacif.-Ecuador.
Gymnogramma Klf. Enum. 73. 1824. HB. 384. Chr. 69; Neurogramme Diels 1899. NPfl. 264; Gymnopteris Und. 1900.
triangulata (Jenm.) Und. Bull. Torr. Cl. **29**. 632. 1902 — Jamaica.
Gymnogramme Jenm. Bull. Dep. Jam. **4**. 206. 1897.
trifoliata Kuhn; Hier. Engl. Jahrb. **22**. 398. 1896 = Trismeria.
viscosa (Eat.) Und. Bull. Torr. Cl. **29**. 631. 1902 — California.
Gymnogramme triangularis v. viscosa Eat. Ferns N. Am. **2**. 16 t. 48 f. 5. 1881.
Cerosora Baker, Journ. Linn. Soc. **24**. 260. 1887 *(Gymnogramme §)* = Gymnogramma.
CETERAC Adanson, Fam. des plantes 2. 20. 1763 = Asplenium et Ceterach.
CETERACH Lam. et DC. Fl. franç. 2. 566. 1805; Willdenow, Enum. 1068. 1809; sp. pl. **5**. 136. 1810 — (Polypodiaceae Gen. **49**).
alpina DC.; Lam. et DC. Fl. franç. 2. 567. 1805 = Woodsia.
alternans Kuhn, v. Deck. Reis. **3**[3]. Bot. 36. 1879 = C. Dalhousiae.
aspidioides Willd. sp. **5**. 137. 1810 = Dryopteris.
aureum (Cav.) L. v. Buch, Abh. Ak. Wiss. Berlin **1816-17**. 361. 1819. (Chr. 210). NPfl. 245 — Ins. Canar. Madeira. — (Acrostichum Cav. 1801 non L. 1753); Asplenium Cav. Descr. 256. 1802; Grammitis Sw. 1806; Asplenium latifolium Bory 1803; Ceterach Fée 1850-52; C. canariensis Willd. 1810. Asplenium Ceterach β HB. 245.
barbatum Pr. Del. Prag. **1**. 162. 1822 = Gymnopteris tomentosa.
canariensis Willd. sp. **5**. 137. 1810 = C. aureum.
capensis Kze. Linn. **10**. 496. 1836 = C. cordatum.
ceterach Newm. Phytol. **1851**. app. V. in obs.; ante pag. 105 = C. officinarum.
cordatum (Thbg.) Desv. Prod. 223. 1827. NPfl. 245 — Africa austr. St. Helena.
Acrostichum Thbg. Prod. Fl. Cap. 171. 1800; Asplenium Sw. 1801; Grammitis Sw. 1806; Cincinalis Desv. 1811; Notholaena Desv. 1813; Gymnogramma Schlecht. 1825. HB. 379. Chr. 67. G. capensis Spr.; Klf. 1831; Ceterach Kze. 1836; Grammitis Moore 1857 — G. cordata var. subbipinnata Hk. 2 Cent. t. 7. 1860; Gymnogramme Haughtoni Hk. 1868. HB. 381 (St. Helena).
crenata Klf. Enum. 85. 1824 = C. cordatum.
Dalhousiae (Hk.) C. Chr. Ind. 170. 1905 — Himalaya. Abyssinia.
Asplenium Hk. Ic. pl. t. 105. 1837; A. alternans Wall. 1828 (nomen); Hk. sp. **3**. 92. 1860. HB. 194. Chr. 190; Ceterach Kuhn 1879. NPfl. 244.
hispanicum Mett. Fil. Lips. 80. 1856 = Pleurosorus Pozoi.
immersum Sturm, Enum. pl. cr. vasc. chil. 29. 1858 = Pleurosorus papaverifolius.
latifolium Fée, Gen. 206, 363 t. 30 f. 1. 1850-52 = C. aureum.
lobatum Pr. Del. Prag. **1**. 162. 1822 = Gymnopteris tomentosa.

Marantae DC.; Lam. et DC. Fl. franç. **2**. 567. 1805 = Notholaena.

officinarum DC.; Lam. et DC. Fl. franç. **2**. 566. 1805. Chr. 210. NPfl. 244 — Europa centr., atlant. et merid. Madeira. Asia occ. et temp.-Himalaya. Africa austr. — Asplenium Ceterach L. sp. **2**. 1080. 1753. HB. 245 (excl. β); Scolopendrium Symons 1798; Vittaria Bernh. 1799 (α); Grammitis Sw. 1806; Gymnopteris Bernh. 1806; Gymnogramme Spr. 1827; Notolepeum Newm. 1844; Ceterach Newm. 1851; Hemidictyum Bedd. 1876.

papaverifolium Mett. Fil. Lips. 80 t. 13 f. 14. 1856 = Pleurosorus.

pedunculata Hk. et Grev. Ic. Fil. t. 5. 1827 = Polypodium.

polypodioides Raddi, Opusc. sci. Bol. **3**. 284. 1819 = Dryopteris.

Pozoi A.Br.: Milde, Bot. Zeit. **1866**. 310 = Pleurosorus.

rutaefolium Mett. Fil. Lips. 80. 1856 = Pleurosorus.

Ceterachopsis J. Smith, Hist. Fil. 317. 1875 *(Asplenium §)* = Ceterach.

Charitophyllum v. d. Bosch, Pl. Jungh. **1**. (18). 1856 *(Hymenophyllum §)* = Hymenophyllum.

Cheilanthastrum Fée, Gen. 149. 1850-52 *(Myriopteris §)* = Cheilanthes.

Cheilanthastrum Fée, 7 mém. 115. 1857 *(Adiantopsis §)* = Adiantopsis.

CHEILANTHES Swartz, Syn. Fil. 5, 126. 1806 — (Polypodiaceae).

acrostica Todaro, Syn. Pl. Acot. vasc. Sic. 10. 1866 = C. pteridioides.

aculeata Klf. Flora **1828**[1]. 367 = Hypolepis repens.

affinis Mett. Cheil. 19. n. 2. 1859 = Notholaena.

alabamensis (Buckl.) Kze. Linn. **20**. 4. 1847. Mett. n. 37. NPfl. 275 — U. S. A. merid. — Pteris Buckl. Am. Journ. Sci. **45**. 177. 1843; Pellaea Bak. 1867. HB. 148; Allosorus O. Ktze. 1891.

albida Bak. Ann. Bot. **5**. 212. 1891 — Mexico.

albofusca Bak. Kew Bull. **1895**. 54 — China merid.

albomarginata Clarke, Tr. Linn. Soc. II. Bot. **1**. 456 t. 52. 1880 — Himalaya. India austr.

allosuroides Mett. Cheil. 32. n. 35. 1859. HB. 137 — Mexico. Cheilosoria Trev. 1877.

alpina Bl. Enum. 138. 1828 = Hypolepis.

amaurorachis Kze. Linn. **23**. 242, 306. 1850 = Dryopteris punctata.

ambigua A. Rich. Fl. N. Zél. 84. 1832 = Hypolepis tenuifolia vel Dryopteris punctata *rugosula.

ambigua Brack. Expl. Exp. **16**. 91. 1854 = Notholaena distans.

ambigua Mett. Cheil. 49. n. 78. 1859 = Pellaea.

amoena A. A. Eat. Fern Bull. **5**. 44. 1897 = Hypolepis californica.

anceps Blanford, Simla Nat. Hist. Soc. $^{25}/_{6}$ 1886; Hope, Journ. Bomb. Nat. Hist. Soc. **13**. 249. 1900 = C. farinosa var.

andina Hk. sp. **2**. 115. 1852. (HB. 134) — Peru.

andina Gris. Abh. Ges. Wiss. Gött. **24**. 342. 1879 = Woodsia montevidensis.

angustifolia HBK. Nov. Gen. et. Sp. **1**. 21. 1815. Mett. n. 66. Chr. 146 — Mexico-Venezuela. — Allosorus Pr. 1836; Onychium Kze. 1848; Pellaea Bak. 1867. HB. 150; Pteris F. Muell. 1870. Cheilanthes decomposita Willd. 1813 (nomen); Fée 1857; Allosorus Mart. et. Gal. 1842; Pellaea Hk. 1858 — Cheilanthes cuneata Link 1833; Allosorus Pr. 1836; Cassebeera J. Sm. 1841. (Othonoloma Link 1833). (an sp. propr.? cum syn. C. chaerophylla Kze.? conf. C. marginata.)

anthriscifolia Willd. sp. **5**. 461. 1810 = Dennstaedtia.

anthriscifolia Schlecht. (Adumbr. 52 t. 32. 1832 ined.) = Hypolepis sparsisora.

aquilinaris Fée, Cr. vasc. Br. **2**. 37 t. 91 f. 1. 1869 = Hypolepis.
arabica »Dcne.«; Buchinger, Flora **1846**. 303 = C. coriacea.
arborescens Sw. Syn. 129, 336. 1806 = Hypolepis tenuifolia.
argentea (Gmel.) Kze. Linn. **23**. 242. 1850. Mett. n. 69. HB. 142. Chr. 149. NPfl. 277 — Sibiria-Malesia. Ins. Philipp. — Pteris Gmel. Nov. Comm. Petr. **12**. 519 t. 12 f. 2. 1768; Allosorus Pr. 1836; Cassebeera J. Sm. 1841; Aleuritopteris Fée 1850-52; Doryopteris Christ 1902 — Pellaea Tamburii Hk. 1858. HB. 146. (f. Bedd.); Cheilanthes Moore 1861; Allosorus O.Ktze. 1891.
argyrostigma J. Sm. Hist. Fil. 283. 1875 (nomen) — (Mexico?). Nothochlaena J. Sm. JoB. **4**. 50. 1841 (nomen).
Aschenborniana Mett. Cheil. 21. n. 7. 1859 = Notholaena.
aspera Klf. Linn. **6**. 186. 1831 = Hypolepis sparsisora var.
aspera Hk. sp. **2**. 111 t. 108 A. 1852 = Pellaea scabra.
aspidioides Fée, Gen. 157. 1850-52 = C. microphylla var.
Atherstonii Hk. sp. **2**. 107. 1852 = Pellaea quadripinnata.
aurantiaca (Cav.) Moore, Ind. XXXVIII. 1857. Mett. n. 74. HB. 141. Chr. 149 — Mexico. — ? Pteris Cav. Descr. 266. 1802; Allosorus Pr. 1836. A. ochraceus Hk. 1840; Cheilanthes Hk. 1852. Aleuritopteris lutea Fourn. 1872 (an Pteris lutea Cav. 1802?; conf. Cheilanthes lutea).
aurea Bak. Syn. ed. I. 453. 1868; ed. II. 476. 1874; Ic. pl. t. 1637 — Guatemala.
auriculata Link, Hort. Berol. **2**. 36. 1833 = Pellaea.
Belangeri (Bory) C. Chr. Ind. 172. 1905 — Asia trop. — Pteris Bory, Bél. Voy. Bot. **2**. 44. 1833; Pteridella Mett.; Kuhn 1879. Pteris varians Wall. 1828 (nomen); Cheilanthes Hk. 1852. HB. 136. Chr. 145. NPfl. 275.
Bergiana Schlecht. (Adumbr. 51 t. 31 ined.); Kze. Linn. **10**. 541. 1836 = [Hypolepis.
Bockii Diels, Engl. Jahrb. **29**. 199. 1900 — China.
Boivini Mett.; Kuhn, Fil. Afr. 69. 1868. HB. 475 — Ins. Comorae et Mascarenae.
Bojeri Moore, Ind. 236. 1861 = Pellaea quadripinnata var.
Bolusii Bak. in Hk. Ic. pl. t. 1636. 1886. NPfl. 277 — Africa austr.
Borsigiana Rchb. fil. et Wswz.; Koch, Wochenschr. Gärtn. Pflk. Berlin **1858**[1]. 2 = Notholaena sulphurea.
brachypus Kze. Linn. **23**. 243, 307. 1850 = Notholaena.
Bradburii Hk. sp. **2**. 97 t. 109 B. 1852 = C. tomentosa.
Brandegii Eat. Bull. Torr. Cl. **17**. 215 t. 104. 1890 — Lower California.
Brandtii Franch. et Sav. Enum. pl. Jap. **2**. 212, 620. 1879 = C. farinosa var.
brasiliensis Raddi, Opusc. sci. Bol. **3**. 295. 1819; Pl. Bras. **1**. 60 t. 75 f. 2. 1825 = Adiantopsis chlorophylla.
bullata Bak. Syn. ed. I. 137. 1867 = C. bullosa.
bullosa Kze. Linn. **24**. 274. 1851. Mett. n. 79. HB. 137 — India austr. China.
bullosa Bedd. Ferns S. Ind. t. 192. 1863 = C. farinosa.
Cadieri Christ, Journ. de Bot. **19**. 70. 1905 — Annam.
caespitosa Mett. Fil. Lips. 51. 1856 = C. marginata.
californica Mett. Cheil. 44. n. 64. 1859 = Hypolepis.
candida Mart. et Gal. Mém. Ac. Brux. **15**. 73 t. 20 f. 1 b. 1842 = Notholaena sulphurea var.
candida Zoll. Nat. Gen. Arch. Neerl. Ind. **2**. 203. 1845 = C. farinosa.
canescens Kze. Linn. **13**. 143. 1839 = Woodsia mollis var.
canescens Mett. Cheil. 25. n. 18. 1859 = Notholaena.
capensis Sw. Syn. 128. 1806 = Adiantopsis.
cartilaginea Gris. Abh. Ges. Wiss. Gött. **19**. 275. 1874 = C. Poeppigiana.

caudata R. Br. Prod. Fl. N. Holl. 156. 1810 — Austr. trop.
(Dicksonia Poir. 1811 ex err.) — conf. Pellaea nitida.
chaerophylla Kze. Linn. **23**. 243, 307. 1850 = C. marginata.
chaerophylloides Fée, 8 mém. 117. 1857 = C. marginata.
cheiloglyphis (Fée) Moore, Ind. 236. 1861 — Mexico.
Myriopteris Fée, 8 mém. 77. 1857.
chilensis Fée, Gen. 156. 1850-52; 7 mém. 37 t. 18 f. 2 = C. glauca.
chlorophylla Sw. Vet. Ak. Handl. **1817**. 76 = Adiantopsis.
chrysophylla Hk. sp. **2**. 113. 1852 = C. farinosa var.
Chusana Hk. sp. **2**. 95 t. 106 B. 1852 = C. mysurensis.
ciliata Fée, 9 mém. 11. 1857 = C. marginata.
cinnamomea Eat. Proc. Amer. Ac. **18**. 186. 1883 = C. Schaffneri.
Clelandi F. Muell. et Tate, Proc. Roy. Soc. S. Austr. **10**. 81. 1887 — S. Australia.
Clevelandii Eat. Bull. Torr. Cl. **6**. 33. 1875. NPfl. 277 — California.
commutata Kze. Linn. **10**. 542. 1836 = Hypolepis sparsisora.
contigua Wall. List n. 72. 1828 = Onychium japonicum.
contigua Bak. Syn. 476. 1874 = C. tenuifolia var.
contracta Kze. Linn. **23**. 243, 307. 1850 = Pellaea quadripinnata var.
contracta Mett.; Kuhn, Fil. Afr. 70. 1868 = C. hirta var.
Cooperae Eat. Bull. Torr. Cl. **6**. 33. 1875 — California.
coriacea Dcne. Arch. Mus. **2**. 190. 1841 — Arabia. Abyssinia.
C. arabica »Dcne.« Buching. 1846. Mett. n. 82. HB. 134. NPfl. 275: Pellaea? Fée 1850-52. Cheilanthes Decaisnii Kze. (1850).
cornuta Kze. Linn. **10**. 534 (pt.) 1836. Mett. Cheil. 31. n. 33. 1859 — Afr. austr.
crenata Kze. Linn. **9**. 84. 1834 = Woodsia.
crenulata Link, Hort. Berol. **2**. 41. 1833 = C. marginata.
cretacea Mett. Cheil. 20. n. 5. 1859 = Notholaena.
cucullans Fée, 7 mém. 39 t. 25 f. 4. 1857 = C. microphylla var.
culcita Desv. Prod. 305. 1827 = Ochropteris pallens.
cuneata Link, Hort. Berol. **2**. 40. 1833 = C. angustifolia var.
Dalhousiae Hk. sp. **2**. 80 t. 78 B. 1852 = C. farinosa var.
davallioides Bory; Willd. sp. **5**. 461. 1810 = Ochropteris pallens.
dealbata Pursh, Fl. Am. sept. **2**. 671. 1814 = Pellaea.
dealbata Don, Prod. Fl. Nepal. 16. 1825 = C. farinosa var.
Decaisnii Kze. Ind. Sem. ht. Lips. —. (Linn. **23**. 307. 1850) = C. coriacea.
decomposita Willd. Enum. Suppl. 70. 1813 (nomen); Fée, 9 mém. 11. 1857 = C. angustifolia.
Delavayi Bak. Ann. Bot. **5**. 211. 1891. NPfl. 275 — Yunnan.
deltoidea Kze. Linn. **10**. 535. 1836 = Doryopteris.
? densa Fée, Gen. 156. 1850-52 = C. hirsuta var.
depauperata Bak. Ann. Bot. **5**. 210. 1891 — Africa austr.
dichotoma Sw. Syn. 129, 335. 1806 = Adiantopsis.
dicksonioides Endl. Prod. Fl. Norf. 15. 1833 = Hypolepis.
digitata Pr. Tent. 160. 1836 (nomen) —? (= ? C. pruinata).
discolor Griff. Galc. Journ. **4**. 466. 1844 = C. farinosa.
dissecta Hk. et Arn. Bot. Beech. Voy. 75. 1832 = Hypolepis tenuifolia.
distans Mett. Cheil. 25. n. 19. 1859 = Notholaena.
dubia Hope, Journ. Bomb. Soc. Nat. Hist. **12**. 528 t. 2. 1899 — India bor.
? **dura** Moore, Ind. 44. 1857 = Pellaea.
Duthiei Bak. Ann. Bot. **5**. 210. 1891 — India bor.

Eatoni Bak. Syn. 140. 1867. NPfl. 277 — Arizona. Texas.
Eckloniana Mett. Cheil. 22. n. 10. 1859 = Notholaena.
elata Kze. Linn. **10**. 542. 1836 = Hypolepis Bergiana.
elegans Desv. Berl. Mag. **5**. 328. 1811 = C. myriophylla.
Ellisiana ht. = C. hirta.
elongata Willd.; Klf. Enum. 213. 1824 = C. microphylla var.
erecta Col. Tr. N. Zeal. Inst. **28**. 619. 1896 — N. Zealand.
eriophora Mett. Cheil. 23. n. 14. 1859 = Notholaena.
farinosa (Forsk.) Klf. Enum. 212. 1824. Mett. n. 71. HB. 142. Chr. 148. NPfl. 276 — Japonia. China. Asia, Africa, America trop. Fiji.
Pteris Forsk. Fl. Aegypt.-Arab. 187. 1775; Allosorus Pr. 1836; Cassebeera J. Sm. 1841; Aleuritopteris Fée 1850-52. Pteris argyrophylla Sw. 1806; Allosorus Pr. 1836; Aleuritopteris Fée 1850-52 — Cheilanthes dealbata Don 1825 (non Pursh 1814); Allosorus Pr. 1836; Aleuritopteris Fée 1850-52 — Cheilanthes Dalhousiae Hk. 1852. HB. 137 — Ch. chrysophylla Hk. 1852. Mett. n. 73; C. argentea var. HB. 142 — C. Brandtii Franch. et Sav. 1879 — C. anceps et C. grisea Blanford 1886; Hope 1900.
fasciculata Goldm. Nova Acta **19**. Suppl. I. 456. 1843 = C. pruinata.
Feei Moore, Ind. XXXVIII. 1857 — U. S. A.
C. vestita Hk. Fl. bor. Am. **2**. 264. 1840; sp. **2**. 98 t. 108 B. 1852 (non Sw. 1806); Myriopteris gracilis Fée, Gen. 150 t. 29 f. 6. 1850-52; Cheilanthes Riehl.; Mett. n. 43. 1859 (non Klf. 1824); C. lanuginosa Nutt. (1852). HB. 139. NPfl. 277 (non Mart. et Gal. 1842); Myriopteris J. Sm. 1875.
Fendleri Hk. sp. **2**. 103 t. 107 B. 1852. HB. 139. NPfl. 277 — U. S. A. occ. et merid. Mexico. — Myriopteris Fourn. 1872.
ferruginea Willd.; Link, Enum. alt. **2**. 463. 1822 = Notholaena bonariensis.
fibrillosa Dav. Bull. Torr. Cl. **15**. 225. 1888 — California.
C. lanuginosa var. fibrillosa Dav. Bull. Torr. Cl. **12**. 21. 1885.
fimbriata Vis. Fl. Dalm. **1**. 42 t. 1 f. 1. 1842 = C. persica.
firma Moore, JoB. **5**. 225. 1853 = Pellaea quadripinnata.
flexuosa Kze. Linn. **22**. 578. 1849. Mett. n. 61. HB. 476 — Brasilia centr.
flexuosa Bak. Syn. ed. I. 138. 1867 = C. Regnelliana.
flexuosissima Bak. Ann. Bot. **5**. 212. 1891 — Brasilia.
Fordii Bak. JoB. **1879**. 304 = Adiantopsis.
Fournieri C. Chr. Ind. 174. 1905 — Mexico. — Myriopteris intermedia Fourn. Bull. Soc. Fr. **27**. 328. 1880 (non Fèe 1850-52); Cheilanthes Bak. 1891.
fragillima F. Muell. Fragm. **5**. 123. 1866 = Notholaena fragilis.
fragilis Hk. Fil. Exot. t. 96. 1859. HB. 135 — Moulmein. Yunnan.
fragrans Webb et Berth. Hist. nat. Canar. **3**². pars 3. 452. 1847 = C. pteridioides.
fragrans Sw. Syn. 127, 325. 1806 = C. mysurensis.
Fraseri Mett.; Kuhn, Linn. **36**. 83. 1869 = Notholaena.
frigida Linden, Cat. 1856; Moore, Gard. Chr. **1857**. 772 = C. lendigera.
furfuracea Pr.; Koch, Wochenschr. Gärtn. Pflzk. **1858**. 3 = Notholaena sulphurea var. candida.
fuscata Bl. Enum. 116. 1828 (patria erronea) = Mohria caffrorum.
gigantea Ces. Rend. Acad. Napoli **16**. 25, 29. 1877 = Dryopteris brunneovillosa.
glaberrima Fée, Cr. vasc. Br. **1**. 54 t. 13 f. 2. 1869 — Brasilia
glandulifera Liebm. Vid. Selsk. Skr. V. **1**. 258 (seors. 106). 1849 = C. Kaulfussii.
glandulifera Moore, Ind. 242. 1861 = ? Paesia viscosa.

glandulifera Fée, Cr. vasc. Br. **2**. 36 t. 88 f. 3. 1872-73 = Gymnogramma myriophylla.
glandulosa Sw. Vet. Ak. Handl. **1817**. 77 = Gymnogramma.
glandulosa Fée, Gen. 158. 1850-52 = Paesia viscosa.
glandulosa Pappe et Raws. Syn. Fil. Afr. austr. 35. 1858 = C. hirta var.
glauca (Cav.) Mett. Cheil. 31. n. 32. t. 3 f. 18-19. 1859 — Mexico. Chile.
Acrostichum Cav. Anal. Hist. Nat. **1**. 107. 1799; Pteris Cav. Descr. 269. 1802; Pellaea J. Sm. 1866. HB. 150; Allosorus J. Sm. 1875 — A. hirsutus Pr. 1825; Pellaea Hk. 1858 — Cheilanthes chilensis Fée 1850-52.
globuligera Christ, Ann. Cons. Jard. Bot. Genève **3**. 40. 1899 — Brasilia.
gracilis Klf. Enum. 209. 1824 = Cryptogramma Stelleri.
gracilis Riehl.; Mett. Cheil. 36. n. 43. 1859 = C. Feei.
gracillima Eat. Rep. U. S. et Mex. Bound. Surv. 2^1. 234. 1859. HB. 139. Chr. 145. NPfl. 277 — Amer. bor. pacif. Guatemala. — Myriopteris J. Sm. 1875.
Griffithiana Fée, Gen. 157. 1850-52 = C. Belangeri.
grisea Blanford, Simla Nat. Hist. Soc. $^{25}/_{6}$ 1886; Hope, Journ. Bomb. Nat. Hist. Soc. **13**. 250. 1900 = C. farinosa var.
guanchica Bolle, Bonplandia **7**. 107. 1859 = C. pulchella.
Hancockii Bak. Kew Bull. **1895**. 54 — Yunnan.
hastaefolia Kze. Linn. **23**. 243. 1850 = Pellaea viridis.
hastata Kze. Linn. **10**. 532. 1836 = Pellaea viridis.
heterophylla Willd.; Klf. Enum. 210. 1824 = Doryopteris pilosa.
heterotricha Anders. Vet. Ak. Handl. **1853**. 129. 1855 — Ins. Galapagos.
hirsuta Link, Hort. Berol. **2**. 42. 1833 = C. marginata.
hirsuta Mett. Cheil. 25. n. 17. 1859 = Notholaena.
hirta Sw. Syn. 128, 329. 1806. Mett. n. 26. HB. 136. Chr. 144. NPfl. 275 — Afr. austr. Réunion. — Adiantum Poir. 1810; Nothochlaena J. Sm. 1841; Myriopteris J. Sm. 1866. Adiantum caffrorum Sw. 1801 (non L.fil. 1781) — Myriopteris intermedia Fée 1850-52 — Cheilanthesolivacea Fée 1850-52 — Adiantum parvilobum Sw. 1801; Cheilanthes Sw. 1806 — Myriopteris contracta Fée 1850-52; Cheilanthes Mett. 1868 — C. glandulosa Pappe et Raws. 1858 (non Sw. 1817).
hispanica Mett. Cheil. 30. n. 29. 1859. HB. 136. Chr. 145. NPfl. 276 — Portugal. Hispania. ? Sicilia. — ? C. Tinaei Todaro 1866.
hispidula Kze. Bot. Zeit. **1848**. 212 = C. tenuifolia.
hostilis Kze. Linn. **9**. 86. 1834 = Hypolepis.
hypoleuca Mett. Cheil. 22. n. 11. 1859 = Notholaena.
imbricata Desv. Prod. 303. 1827 = Jamesonia.
? inaequalis Mett. Cheil. 24. n. 24. t. 3. f. 4. 1859 = Notholaena.
incisa Kze.; Mett. Cheil. 44 n. 65 t. 3 f. 28-31. 1859 = Hypolepis.
induta Kze. Linn. **10**. 538. 1836. Mett. n. 51. HB. 138. NPfl. 277 — Africa austr. Myriopteris Fée 1850-52.
intermedia Bak. Ann. Bot. **5**. 213. 1891 = C. Fournieri.
interrupta Klf. Wes. d. Farnkr. 116. 1827 = Notholaena distans.
intramarginalis Hk. sp. **2**. 112. 1852 = Pellaea.
javanica Kze. Bot. Zeit. **1848**. 211 = Notholaena hirsuta.
javensis Moore, Ind. 244. 1861 = C. tenuifolia vel Notholaena hirsuta.
Kaulfussii Kze. Linn. **13**. 145. 1839 — Mexico-Venezuela.
C. viscosa Link, Hort. Berol. **2**. 43. 1833. Mett. n. 31. HB. 136. Chr. 142 (non Carm. 1818).
Kirkii Hk. 2 Cent. t. 81. 1861 = Doryopteris concolor var.

Kirkii Armstr. Tr. N. Zeal. Inst. **13**. 360. 1881 = C. tenuifolia.
Kleinhofii Bl. Enum. 137. 1828 = Notholaena hirsuta.
Krameri Franch. et Sav. Enum. pl. Jap. **2**. 212, 619. 1879 — Japonia
Kuhnii Milde, Bot. Zeit. **1867**. 149. HB. 476 — China bor.
Kunzei Mett. Cheil. 27. n. 22. t. 3. f. 6, 7. 1859 — Africa austr.
Cheilosoria Trev. 1877. Cheilanthes profusa var. minor Kze. Linn. **10**. 535. 1836.
laciniata Sod. Cr. vasc. quit. 79. 1893 — Ecuador. (an Adiantopsis?).
lanosa (Michx.) Watt, JoB. **1874**. 48. Chr. 144. NPfl. 275 — U. S. A.
Nephrodium Michx. Fl. bor. Am. **2**. 270. 1803; Polypodium Poir. 1804; Aspidium Sw. 1806; Adiantum vestitum Spr. 1804; Cheilanthes Sw. 1806. Mett. n. 27. HB. 134 (non Hk. 1840); Cincinalis Desv. 1811; Notholaena Desv. 1813; Myriopteris J. Sm. 1857.
lanosa Moore, Ind. 245. 1861 = C. Feei.
lanuginosa Mart. et Gal. Mém. Ac. Brux. **15**. 75 t. 20 f. 2. 1842 = C. lendigera.
lanuginosa Nutt.; Hk. sp. **2**. 99. 1852 (syn.). HB. 139 = C. Feei.
lanuginosa var. fibrillosa Dav. Bull. Torr. Cl. **12**. 21. 1885 = C. fibrillosa.
laxa Moore, Ind. 245. 1861 = C. Thwaitesii.
lendigera (Cav.) Sw. Syn. 128, 328. 1806. Mett. n. 44. HB. 141. Chr. 147 — Arizona-Ecuador. — Pteris Cav. Descr. 268. 1802; Adiantum Poir. 1810; Nothochlaena J. Sm. 1846; Myriopteris J. Sm. 1857; Cheilanthes lentigera Willd. 1810; Nothochlaena J. Sm. 1841; Myriopteris Fée 1850-52 — Cheilanthes frigida Linden; Moore 1857; Myriopteris J. Sm. 1857 — Cheilanthes minor Mart. et Gal. 1842; Myriopteris Fée 1850-52.
lentigera Willd. sp. **5**. 460. 1810 = C. lendigera.
lepida Phil. Anal. Univ. Chil. **94**. 355. 1896 — Chile.
leptophylla R.Br. in H. Salt, Voy. to Abyss. App. IV. LXV. 1814 (nomen) = Onychium melanolepis.
leucopoda Link, Fil. sp. 66. 1841 (HB. 136). Chr. 148. NPfl. 277 — Texas. [Mexico-Venezuela.
Lidgatii Bak. Syn. 475. 1874 = Pteris Lydgatii.
Lindheimeri Hk. sp. **2**. 101 t. 107 A. 1852. Mett. n. 41 c. HB. 140 — Texas-Arizona-Mexico. — Myriopteris J. Sm. 1854.
Lindigii Mett. Ann. sc. nat. V. **2**. 218. 1864 = Adiantopsis.
linearis Moore, JoB. **5**. 226. 1853 = Pellaea quadripinnata.
longipila Bak. Ann. Bot. **5**. 211. 1891 — Mexico.
lucida Wall. List n. 69. 1828 = Onychium japonicum var.
lutea (Cav.) Moore, Ind. 246. 1861 — Amer. trop.
Pteris Cav. Descr. 267. 1802. (an C. aurantiaca?).
Mac Leanii Hk. sp. **2**. 93 t. 110 B. 1852 = C. pilosa.
macrophylla Kze. Linn. **23**. 244, 307. 1850 = Pellaea viridis.
madagascariensis Bak. Journ. Linn. Soc. **16**. 198. 1877 = Adiantopsis.
maderensis Lowe JoB. **1**. 26. 1834 (nomen); Tr. Cambr. Phil. Soc. **6**. 528. 1838 = [C. pteridioides var.
Malaccensis Fée, Gen. 157. 1850-52 = C. Belangeri.
marginata HBK. Nov. Gen. et Sp. **1**. 22. 1815; **7**. t. 669. Mett. n. 67. Chr. 146 — Arizona-Argentina. — Allosorus J. Sm. 1854; Pellaea Bak. 1867. HB. 151 Allosorus ciliatus Pr. 1825; Cheilanthes Fée 1857; Pellaea J. Sm. 1866 — Allosorus chaerophyllus Mart. et Gal. 1842; Cheilanthes Kze. 1850: Onychium Fée 1850-52. (an sp. propr.? an subsp. C. cuneatae Link?) — Cheilanthes pyramidalis Fée 1857 — C. (Othonoloma) hirsuta Link 1833.
Mathewsii Kze. Farnkr. **1**. 50 t. 25. 1840 = C. pruinata.

meifolia Eat. Proc. Amer. Ac. **18**. 185. 1883 = Hypolepis.
melanocoma Bory, Bél. Voy. Bot. **2**. 71. 1833 = C. mysurensis.
mexicana Dav. Bull. Torr. Cl. **15**. 227. 1888 — Mexico.
Meyeniana Pr. Tent. 160. 1836 (nomen) = C. pilosa.
micrantha Wall. List n. 68. 1828 = C. tenuifolia.
micromera Link, Hort. Berol. **2**. 36. 1833 = C. microphylla var.
microphylla Sw. Syn. 127. 1806. Mett. n. 36. HB. 135. Chr. 144. NPfl. 275 — Amer. trop. — Adiantum Sw. Prod. 135. 1788; Allosorus Liebm. 1849; Nothochlaena Keys. 1873 — Adiantum pubescens Poir. 1810 (non Schkuhr 1809); Cheilanthes HBK. 1815 — C. micromera Link 1833. C. aspidioides Fée 1850-52 — C. Moritziana Kze. 1847. Mett. n. 39 — C. elongata Kze. 1850. Mett. n. 38; Cassebeera microphylla J. Sm. 1841. ? Pteris nigricans Willd. 1810; Notholaena Desv. 1827 — Cheilanthes cucullans Fée 1857.
microphylla L. v. Buch, Abh. Ak. Wiss. Berlin. **1816-17**. 360. 1819 = C. pteridioides.
micropteris Sw. Syn. 126, 324 t. 3 f. 5. 1806. Mett. n. 23. HB. 134. Chr. 145 (pt.). NPfl. 275 — Ecuador-Argentina. — Nothochlaena Keys. 1873.
minor Mart. et Gal. Mém. Ac. Brux. **15**. 75 t. 21 f. 1. 1842 = C. lendigera.
mollis Pr. Tent. 160. 1836 = Notholaena.
moluccana Bl. Enum. 136. 1828 = Notholaena hirsuta.
moluccana Kze. Bot. Zeit. **1848**. 445 = C. tenuifolia.
moncloviensis Bak. Ann. Bot. **5**. 210. 1891 — Mexico.
monosticha Mett. Cheil. 20. n. 4. 1859 = Notholaena sulphurea.
monticola Gardn. in Hk. Ic. pl. t. 487. 1842 = Adiantopsis.
Moritziana Kze. Linn. **23**. 244. 1850 (Kl. Linn. **20**. 338. 1847. syn.) = C. microphylla var.
mucronata Moore, Ind. 45, 248. 1861 = Pellaea.
multifida Sw. Syn. 129, 334. 1806. Mett. n. 52. HB. 138. Chr. 145. NPfl. 275 — Afr. austr. St. Helena. — Adiantum Sw. Schrad. Journ. **1800²**. 85. 1801.
myriophylla Desv. Berl. Mag. **5**. 328. 1811. HB. 140. Chr. 147. NPfl. 277 — Mexico-Chile. — Nothochlaena J. Sm. 1841; Myriopteris J. Sm. 1854 — Cheilanthes elegans Desv. 1811. Mett. n. 42; Myriopteris J. Sm. 1857. Cheilanthes paleacea Mart. et Gal. 1842; Myriopteris Fée 1850-52.
mysurensis Wall. List n. 66. 1828 (nomen); Hk. sp. **2**. 94 t. 100 A. 1852. Mett. n. 24. HB. 135. Chr. 144. NPfl. 275 — Japonia. China. Asia trop. C. fragrans Sw. Syn. 127, 325 t. 3 f. 6. 1806 (non Polypodium L. 1771); C. Swartzii Webb et Berth. 1847 — C. Chusana Hk. 1852 — ? C. opposita Klf. 1824 (an nomen opt. ?).
nitidula Hk. sp. **2**. 112. 1852 = Pellaea.
nudiuscula Moore, Ind. 249. 1861 = Notholaena hirsuta.
obducta Mett.; Kuhn, Linn. **36**. 83. 1869 = Notholaena.
obtusata Pr. Rel. Haenk. **1**. 64 t. 11 f. 1. 1825 — Peru. (an Hypolepis ob. Kuhn ?).
ochracea Hk. sp. **2**. 114. 1852 = C. aurantiaca.
odora Sw. Syn. 127, 327. 1806 = C. pteridioides.
olivacea Fée, Gen. 156. 1850-52 = C. hirta var.
opposita Klf. Enum. 211. 1824 = ? C. mysurensis.
Orbignyana Mett.; Kuhn, Linn. **36**. 82. 1869 — Bolivia.
ornithopus Moore, Ind. 250. 1861 = Pellaea.
paleacea Mart. et Gal. Mém. Ac. Brux. **15**. 76 t. 21 f. 2. 1842 = C. myriophylla v. elegans.
pallens Desv. Prod. 306. 1827 = Ochropteris.
pallida Bl. Enum. 139. 1828 = Hypolepis tenuifolia var.
Palmeri Eat. Proc. Amer. Ac. **22**. 464. 1887 — Mexico.

parallelogramma Kze. Linn. **9**. 85. 1834 = Hypolepis.
Parishii Dav. Bull. Torr. Cl. **8**. 61 t. 8. 1881 — California.
parviloba Sw. Syn. 128, 331. 1806 = C. hirta var.
patula Bak. JoB. **1888**. 225 — China.
paupercula Mett. Fil. Lips. 52. 1856 = Adiantopsis.
pedata A. Br. Ind. Sem. ht. Berol. 1857; Mett. Cheil. 42. n. 59. 1859 = Adiantopsis
pellita Spr. Nov. prov. hort. 10. 1819 — Aegyptus.
pellucida Col. Tasm. Journ. **2**. 173. 1844 = Hypolepis tenuifolia.
persica (Bory) Mett.; Kuhn, Fil. Afr. 73. 1868. Chr. 148. NPfl. 277 — Himalaya-Italia. Algeria. — Notholaena Bory, Bél. Voy. Bot. **2**. 23. 1833. Cheilanthes Szovitsii Fisch. et Mey. 1833 (nomen); 1838. Mett. n. 47. HB. 139; Oeosporangium Vis. 1872; Nothochlaena Keys. 1873; Myriopteris J. Sm. 1875. Cheilanthes fimbriata Vis. 1842; Oeosporangium Vis. 1867.
? peruviana Moore, Ind. 250. 1861 = Notholaena scariosa.
peruviana Bak. Ann. Bot. **5**. 213. 1891 = Plecosorus.
pilosa Goldm. Nova Acta **19**. Suppl. I. 455. 1843. Mett. n. 50. HB. 135 — Peru. C. Mac Leanii Hk. 1852; Myriopteris J. Sm. 1875.
Poeppigiana Mett.; Kuhn, Linn. **36**. 84. 1869 — Peru. Bolivia. Argentina. Allosorus rigidus Kze. 1834 (non Pteris Sw. 1806). Cheilanthes cartilaginea Gris. 1874.
Pohliana Mett. Cheil. 23. n. 13. 1859 = Notholaena.
Pohliana Keys. Pol. Cyath. Hb. Bung. 25. 1873 = Doryopteris concolor.
polymorpha Desv. Prod. 305. 1827 = Doryopteris pilosa.
polypodioides Bl. Enum. 139. 1828 = Dryopteris punctata.
praetexta Klf. Enum. 212. 1824 = Adiantopsis capensis.
Preissiana Kze. in Lehm. Pl. Preiss. **2**. 112. 1847 = C. Sieberi.
Prenticei Luerss. Bot. Centralbl. **9**. 442. 1882 = Notholaena.
Pringlei Dav. Bull. Torr. Cl. **10**. 61 t. 34. 1883. NPfl. 277 — Arizona.
profusa Kze. Linn. **10**. 535. 1836 = ? Pellaea involuta.
profusa var. minor Kze. l. c. = C. Kunzei.
propinqua Mett. Ann. sc. nat. V. **2**. 218. 1864 = Adiantopsis.
pruinata Klf. Enum. 210. 1824. NPfl. 275 — Peru. Bolivia. Argentina. C. Mathewsii Kze. 1840. Mett. n. 25. HB. 134; Nothochlaena Gris. 1874.
pruinosa Kze. Linn. **23**. 245. 1850 = C. pruinata.
pteridioides (Reich.) C. Chr. Ind. 178. 1905 — Ins. Atlant.-Reg. Mediterr.-Himalaya. — Polypodium Reich. 1780; P. fragrans L. Mant. 307. 1771 (non 1753); Adiantum L.fil. 1781; Pteris Lag. 1802; Cheilanthes Webb et Berth. 1847. Mett. n. 48. HB. 134. Chr. 144. NPfl. 275. Pteris acrosticha Balbis 1801; Cheilanthes Todaro 1866 — C. suaveolens Sw. 1806; Adiantum Poir. 1810 — Cheilanthes odora Sw. 1806; Adiantum DC. 1815 — Cheilanthes maderensis Lowe 1834. Mett. n. 49.
pteroides Sw. Syn. 128. 1806 = Pellaea.
pubescens HBK. Nov. Gen. et Sp. **1**. 21. 1815 = C. microphylla var.
pulchella Bory; Willd. sp. **5**. 456. 1810. Mett. n. 83. HB. 136. Chr. 146. NPfl. 275 — Madeira. Ins. Canar. — Allosorus Pr. 1836.
pulveracea Pr. Rel. Haenk. **1**. 64. 1825 = Notholaena sulphurea.
pulveracea Spr. Syst. **4**. 115. 1827 = C. farinosa.
pumilio F. Muell. Cens. 138. 1882 = Notholaena.
pygmaea Kl. Linn. **20**. 338. 1847 = C. microphylla.
pyramidalis Fée, 7 mém. 38 t. 25 f. 3. 1857 = C. marginata var.

quadripinnata Kuhn, Fil. Afr. 74. 1868 = Pellaea.
radiata R. Br.; J. Sm. JoB. **4**. 159. 1841 = Adiantopsis.
ramentacea Wahlenb. Fl. Carpat. 331. 1814 = Pedicularis palustris.
Rawsoni Mett.; Kuhn, Fil. Afr. 75. 1868 = Notholaena.
recurvata Bak. JoB. **1878**. 299. NPfl. 277 — Paraguay.
Reesii Jenm. JoB. **1886**. 267 = Adiantopsis.
refracta Pappe et Raws. Syn. Fil. Afr. austr. 34. 1858 — Africa austr.
Regnelliana Mett.; Bak. Fl. bras. 1^2. 391 t. 43 f. 1. 1870. HB. ed. II. 138. NPfl. 277 — Brasilia. — Pellaea Prantl 1882; Gymnogramme Christ 1900; Cheilanthes flexuosa Bak. 1867. HB. ed. I. 138 (non Kze. 1849).
regularis Mett. Cheil. 41. n. 56. 1859 = Adiantopsis.
remota Kze. Anal. 36. 1837 = Dryopteris punctata.
repens Klf. Enum. 215. 1824 = Hypolepis.
resinifera Bl. Enum. 138. 1828 = Dennstaedtia.
resistens Kze. Linn. **24**. 275. 1851 = Dryopteris punctata.
rigescens Kze. Flora **1839**[1]. Beibl. 51 = Hypolepis.
rigida Moore in Fée, 7 mém. 34. 1857; Mett. Cheil. 48. n. 77. 1859 = Doryopteris.
rigidula Wall. List n. 2175. 1829 = C. farinosa.
rudis Kze. Bot. Zeit. **1848**. 213 = Hypolepis tenuifolia vel Dryopteris setigera.
rufa Don, Prod. Fl. Nepal. 16. 1825. Mett. n. 72. HB. 141. Chr. 149. NPfl. 276 [— India bor.
rufescens Link, Hort. Berol. **2**. 39. 1833 = C. marginata.
rugosula Fourn. Ann. sc. nat. V. **18**. 326. 1873 = Paesia.
rupestris Wall. List n. 67. 1828 = C. tenuifolia.
scabra Karst. Bot. Zeit. **1854**. 855 = Hypolepis parallelogramma.
scariosa Pr. Rel. Haenk. **1**. 65. 1825. Mett. n. 34. HB. 140. NPfl. 277 — Peru. Mexico. — Myriopteris Fée 1850-52 (certe non Acrostichum Sw. 1806; nomen malum).
scariosa Desv. Prod. 305. 1827 = Notholaena.
scariosa Mart. et Gal. Mém. Ac. Brux. **15**. 74. 1842 = Plecosorus speciosissimus.
Schaffneri Moore, Ind. 252. 1861 — Mexico. — Myriopteris rufa Fée, 8 mém. 77. 1857 (non Cheilanthes Don 1825). Cheilanthes cinnamomea Eat. 1883.
Schimperi Kze. Farnkr. **1**. 52 t. 26. 1840 = Hypolepis.
scita ht. v. d. Heede; Gard. Chr. III. **11**. 813. 1892 — Hort.
Seemanni Hk. sp. **2**. 85 t. 97 A. 1852. HB. 133 — Mexico.
Sellowiana Pr. Tent. 160. 1836 = Gymnogramma.
semiglabra Fée, Gen. 156. 1850-52 = C. tenuifolia.
setigera Bl. Enum. 138. 1828 = Dryopteris.
Sieberi Kze. Ind. Sem. ht. Lips. 1839; Pl. Preiss. **2**. 112. 1847. HB. 137 — Australia. N. Zealand. N. Caledonia.
Skinneri Moore, Ind. 253. 1861 = Doryopteris.
sparsisora Schrad. Gött. gel. Anz. **1818**. 918 = Hypolepis.
speciosissima A.Br.; Kze. Anal. 35 t. 23. 1837 = Plecosorus.
spectabilis Klf. Enum. 214. 1824 = Adiantopsis chlorophylla.
spinulosa Link, Hort. Berol. **2**. 39. 1833 = Pteris leptophylla.
squamosa Gill.; Hk. et Grev. Ic. Fil. t. 151. 1829 = Notholaena scariosa.
squamosa Heer et Regel, Linn. **24**. 174. 1851 = C. pteridioides.
squamosa var.? brachypus Kze. Linn. **18**. 340. 1844 = Notholaena brachypus.
stenophylla Kze. Bot. Zeit. **1848**. 212 = Dryopteris setigera.
Streetiae Bak. JoB. **1880**. 327 — Madagascar.
Nothochlaena Bak. Journ. Linn. Soc. **16**. 204. 1877.

suaveolens Sw. Syn. 127. 1806 = C. pteridioides.

subvillosa Hk. sp. **2**. 87 t. 98 B. 1852. Mett. n. 76. HB. 137 — Himalaya.

Swartzii Webb et Berth. Hist. nat. Canar. 3^2. pars 3. 453. 1847 = C. mysurensis.

Szovitsii Fisch. et Mey. Bull. Soc. Mosc. **6**. 260. 1833 (nomen); **1838**. 241 = C. persica.

taliensis Christ, Bull. Soc. Fr. **52** Mém. I. 58. 1905 — Yunnan.

Tamburii Moore, Ind. 254. 1861 = C. argentea var.

tenuifolia (Burm.) Sw. Syn. 129, 332. 1806. Mett. n. 21. HB. 138. Chr. 144. NPfl. 275 — Asia, Australia, Polynesia trop.-Tasmania-N. Zealand.
Trichomanes Burm. Fl. Ind. 237. 1768; Adiantum Sw. 1801; Cassebeera J. Sm. 1841; Cheilosoria Trev. 1877. Acrostichum tenue Retz. 1791. Adiantum cicutaefolium Lam. 1783 — Nothochlaena semiglabra Kze. 1850; Cheilanthes Fée 1850-52 — C. contigua Bak. 1874. HB. 476 — C. hispidula Kze. 1848. Mett. n. 20.

tenuis Pr. Rel. Haenk. **1**. 65. 1825 — Mexico.

tenuis Lowe, Ferns **4**. t. 23. 1859 = C. lendigera.

ternifolia Moore, Ind. 255. 1861 = Pellaea.

Thwaitesii Mett.; Kuhn, Linn. **36**. 82. 1869. HB. 475 — India austr. Ceylon. C. laxa Moore 1861 (nomen).

Tinaei Todaro, Syn. pl. acot. vasc. Sicil. 12. 1866 = ? C. hispanica.

tomentosa Link, Hort. Berol. **2**. 42. 1833. Mett. n. 45. HB. 140. Chr. 147. NPfl. 277 — U. S. A. merid.-Mexico. — Nothochlaena J. Sm. 1842; Myriopteris Fée 1850-52; Nothochlaena Keys. 1873 — Cheilanthes Bradburii Hk. 1852. Mett. n. 46; Myriopteris J. Sm. 1875.

triangula Kze. Linn. **10**. 536. 1836 = Pellaea quadripinnata.

trichomanoides Mett. Cheil. 19 n. 1. 1859 = Notholaena.

trichophylla Bak. Ann. Bot. **5**. 211. 1891. NPfl. 276 — Yunnan.

trifurcata Bak. Kew Bull. **1901**. 144 = Adiantopsis monticola var.

Tweediana Hk. sp. **2**. 84 t. 96 B. 1852 — Brasilia austr.

valdiviana Phil. Linn. **29**. 106. 1857 — Chile.

varians Hk. sp. **2**. 89 t. 103 A. 1852 = C. Belangeri.

vellea F. Muell. Fragm. **5**. 123. 1866 = Notholaena Brownii.

venosa Col. Tr. N. Zeal. Inst. **25**. 321. 1893 — N. Zealand.

venusta Fée, 9 mém. 12. 1857 = C. angustifolia.

vestita Sw. Syn. 128, 338. 1806 = C. lanosa.

vestita Hk. Fl. bor. Am. **2**. 264. 1840 = C. Feei.

vestita Brack. Expl. Exp. **16**. 91. 1854 = C. gracillima.

villosa Dav. Cat. Dav. Herb. Suppl. 45. 1883 = C. myriophylla.

viridis Sw. Syn. 127. 1806 = Pellaea.

viscida Dav. Bull. Torr. Cl. **6**. 191. 1877 — California.

viscosa Carm. Trans. Linn. Soc. **12**. 511. 1818 = Dryopteris punctata.

viscosa Link, Hort. Berol. **2**. 43. 1833 = C. Kaulfussii.

viscosa Karst. Bot. Zeit. **1854**. 855 = Hypolepis.

Weddelliana Moore, Ind. 257. 1861 = Pellaea ternifolia.

Welwitschii Hk. Syn. 142. 1867. NPfl. 275 — Angola.

Wrightii Hk. sp. **2**. 87 t. 110 A. 1852. HB. 138 — Texas-New Mexico.

CHEILOGRAMMA Maxon, Proc. U. S. Nat. Mus. **23**. 630. 1901 (an Und. Our nat. ferns ed. VI. 1900?) = **Paltonium** — (§ Chilogramme Bl.).

lanceolata Maxon l. c. (Und. l. c.?)

CHEILOLEPTON Fée, Hist. Acrost. 19. 1845 = **Leptochilus.**

Blumeanum Fée, Acrost. 89 t. 51. 1845 = L. lomarioides.

CHEILOPLECTON Fée, 7 mémoire 33. 1857. (Cheilopecton ex err.) = **Doryopteris.**
rigidum Fée, 7 mém. 34 t. 20 f. 3. 1857.

CHEILOSORIA Trevisan, Atti dell'Istit. Veneto V. 3. 579. 1877 = **Cheilanthes.**
allosuroides Trev. l. c. 579. 1877.
javensis Trev. l. c. 580. 1879 = C. tenuifolia.
Kunzei Trev. l. c. 580. 1877.
tenuifolia Trev. l. c. 579. 1877.

Cheilosorum J. Smith, Hist. Fil. 317. 1875 (*Asplenium §*) = Asplenium.

CHEIROGLOSSA Presl, Suppl. Tent. Pterid. 56. 1845 = **Ophioglossum.**
palmata Pr. Suppl. 57. 1845.

CHEIROPLEURIA Presl, Epim. Bot. 189. 1849 — (Polypodiaceae Gen. **110**).
bicuspis (Bl.) Pr. Epim. 189. 1849. Chr. 128. NPfl. 336 — Java. Liu Kiu. Formosa. Ins. Philipp. N. Guinea. — Polypodium Bl. Enum. 125. 1828; Anapausia Moore 1857; Acrostichum Hk. 1864. HB. 421. A. trinerve Hassk. 1844 — Gymnopteris Vespertilio Hk. 1846; Cheiropleuria Pr. 1849; Acrostichum Mett. 1856; Anapausia Moore 1857.
tricuspis J. Sm. Hist. Fil. 139. 1875 = Leptochilus.
Vespertilio Pr. Epim. 190. 1849 = C. bicuspis.
(*Obs.* vide Polypodium cantoniense Bak.).

CHEIROPTERIS Christ, Bull. Boiss. 6. 876. 1898 = **Neocheiropteris.**
Ebertii Christ, Bull. Boiss. 6. 878. 1898 (nomen) — Canton.
Henryi Christ, Bull. Boiss. 6. 876. 1898 = N. palmatopedata.
palmatopedata Christ, Bull. Boiss. 7. 21 t. 1. 1899.

Chilodium Presl, Hymenoph. 23. 1843 (*Didymoglossum §*) = Trichomanes et Hymenophyllum.

Chilogramme Blume, Fl. Javae Fil. 70. 1828 (*Antrophyum §*) = Eschatogramme, Paltonium, Vittaria (angustifolia).

Chilopteris Presl, Tent. Pterid. 208. 1836 (*Grammitis §*) = Polypodium.

Chlamydia Moore, Ind. Fil. CIV. 1857 (*Amphicosmia §*) = Hemitelia.

CHNOOPHORA Kaulfuss, Enum. Fil. 250. 1824 = **Alsophila.**
excelsa Mart. Ic. Cr. Bras. t. 27, 29 f. 1, 2, 37. 1828-34 = A. corcovadensis.
glauca Bl. Enum. 243. 1828.
Humboldtii Klf. Enum. 250. 1824 = A. villosa.
lanuginosa Jungh. in Hoev. Tijdschr. 8. 372. 1841; Flora **1847**. 522 = A. [tomentosa.
lurida Bl. Enum. 244. 1828.
?tomentosa Bl. Enum. 244. 1828.

CHONTA Molina, Saggio storia nat. Chile 156. 1782 (1809?) = **? Thyrsopteris.**

CHORISTOSORIA Mett.; Kuhn, v. Deck. Reisen 3^3. Bot. 13. 1879 = **Pellaea.**
pteroides Mett.; Kuhn, v. Deck. Reis. 3^3. Bot. 13. 1879.

CHORIZOPTERIS Moore, Gard. Chron. **1855**. 854 = **Leptochilus.**
bipinnata Moore, l. c. (nomen) } = L. lomarioides f. monstr.
pinnata Moore, l. c. (nomen) }

Chrysochosma J. Smith, Hist. Fil. 279. 1875 (*Nothochlaena §*) = Notholaena.

Chrysodia Fée, Gen. Fil. 183. 1850-52 (*Ceropteris §*) = Ceropteris.

CHRYSODIUM Fée, Hist. d'Acrost. 22. 1845.
acuminatum Mett. Fil. Lips. 22. 1856 = Leptochilus nicotianifolius.
alienum Mett. Fil. Lips. 21 t. 10 f. 5. 1856 = Leptochilus.
aureo-nitens Christ, Farnkr. 47. 1897 = Trachypteris pinnata.
aureum Mett. Fil. Lips. 21. 1856 = Acrostichum.
bipinnatifidum Mett.; Kuhn, Fil. Afr. 50. 1868 = Leptochilus cuspidatus var.

Boivini Mett.; Kuhn, Fil. Afr. 51. 1868 = Leptochilus.
Cayennense Fée, Acrost. 100 t. 59. 1845 = Acrostichum aureum.
crinitum Mett.; Fil. Lips. 21. 1856 = Elaphoglossum.
cuspidatum Kuhn, Ann. Lugd. Bat. **4**. 293. 1869 = Leptochilus.
danaeaefolium Fée, Acrost. 101. 1845 = Acrostichum aureum.
decrescens Salom. Nom. 135. 1883 = Stenochlaena.
diversifolium Ettingsh. Denkschr. Ak. Wien **22**. 59 t. 9 f. 8. 1864 = Leptochilus
fasciculatum Fourn. Ann. sc. nat. V. **18**. 272. 1873 = Acrostichum.
flagelliferum Mett. Fil. Lips. 21. 1856 = Leptochilus heteroclitus.
fluviatile Kuhn, Fil. Afr. 51. 1868 = Leptochilus.
fraxinifolium Fée, Acrost. 101. t. 62. 1845 = Acrostichum aureum.
fraxinifolium Mett. Fil. Lips. 21. 1856 = Leptochilus serratifolius.
Gaboonense Kuhn, Fil. Afr. 51. 1868 = Leptochilus.
Harlandii Hance, Journ. Linn. Soc. **13**. 138. 1873 = Leptochilus.
heterocliton Kuhn, Ann. Lugd. Bat. **4**. 294. 1869 = Leptochilus.
Heudelotii Kuhn, Fil. Afr. 51. 1868 = Leptochilus.
hirsutum Fée, Acrost. 99 t. 62 f. 2. 1845 = Acrostichum aureum.
inaequale Fée, Acrost. 100. 1845 = Acrostichum aureum.
Lindigii Mett. Ann. sc. nat. V. **2**. 204. 1864 = Leptochilus.
Linnaeanum Kuhn, Ann. Lugd. Bat. **4**. 293. 1869 = Leptochilus.
lomarioides Jenm. Timehri **4**. 314. 1885 = Acrostichum aureum var.
maracaybense Karst. Fl. Col. **1**. 115 t. 57. 1860 = Leptochilus alienus var.
nicotianaefolium Mett. Fil. Lips. 22. 1856 = Leptochilus.
oligarchicum Salom. Nom. 136. 1883 = Leptochilus.
opacum Mett. Ann. sc. nat. V. **2**. 204. 1864 = Leptochilus.
palustre Luerss. Fil. Graeff. 73. 1871 = Stenochlaena.
pellucens Mett. Ann. sc. nat. V. **2**. 205. 1864 — Columbia. (Leptochilus sp. dub.).
polyphyllum Luerss. Fil. Graeff. 69. 1871 = Polybotrya.
praestantissimum Mett. Fil. Lips. 21. 1856 = Acrostichum.
punctatum Mett.; Kuhn, Fil. Afr. 51. 1868 } = Leptochilus auriculatus.
punctulatum Mett. Fil. Lips. 21. 1856 }
Quoyanum Ettingsh. Denkschr. Ak. Wien **22**. 59 t. 8 f. 1. 1864 = Leptochilus cuspidatus.
repandum Mett. Novara Exp. Bot. **1**. 202. 1870 = Leptochilus cuspidatus.
sagenioides Kuhn, Linn. **36**. 63. 1869 = Leptochilus.
salicinum Kuhn, Fil. Afr. 52. 1868 = Leptochilus.
scalpturatum Fée, Acrost. 100 t. 61. 1845 = Acrostichum aureum.
scalpturatum Mett. Fil. Lips. 21. 1856 = Leptochilus.
serratifolium Ettingsh. Denkschr. Ak. Wien **22**. 57 t. 8 f. 4, 7. t. 10 f. 5. 1864 [= Leptochilus.
serratum Kuhn, Linn. **36**. 63. 1869 = Leptochilus.
sorbifolium Luerss. Fil. Graeff. 71. 1871 = Stenochlaena.
speciosum Fée, Acrost. 101. 1845 = Acrostichum aureum.
Urvillei Fée, Acrost. 100 t. 60. 1845 = Acrostichum aureum.
vulgare Fée, Acrost. 97. 1845 = Acrostichum aureum.
Zollingeri Kuhn, Ann. Lugd. Bat. **4**. 293. 1869 = Leptochilus.

CHRYSOPTERIS Link, Fil. spec. 116, 120. 1841 = **Polypodium.**
araneosa Fourn. Mex. pl. **1**. 86. 1872 } = P. aureum *areolatum.
areolata Fée, 9 mém. 27. 1857 }
aurea Link, Fil. sp. 121. 1841.

Billardieri Link, Fil. sp. 123. 1841 = P. diversifolium.
decumana Fée, Gen. 265. 1850-52.
dictyocallis Fée, Gen. 265. 1850-52 = P. decumanum var.
dulcis Fée, Gen. 265. 1850-52 = ? P. aureum.
glauca Fée, Gen. 265. 1850-52 } = P. aureum *areolatum.
glaucina Fourn. Mex. pl. **1**. 86. 1872 }
grandis Fée, 8 mém. 96. 1857 = P. aureum.
lanosa Fée, 8 mém. 97. 1857 = P. aureum *areolatum.
lepidopoda Link, Fil. sp. 123. 1841 = P. diversifolium.
longipes Link, Fil. sp. 122. 1841 = P. phymatodes.
martinicensis Fée, 8 mém. 130. 1867 = P. aureum.
microdictya Fée, 8 mém. 97. 1857 = P. aureum.
multiserialis Fée, 11 mém. 71. 1866 = P. decumanum var.
peltidea Link, Fil. sp. 123. 1841 = P. phymatodes.
Phymatodes Link, Fil. sp. 122. 1841.
pulvinata Link, Fil. sp. 121. 1841 = P. aureum*.
Raddiana Fée, Cr. vasc. Br. **1**. 120. 1869 = P. catharinae.
sporadocarpa Link, Fil. sp. 121. 1841 = P. aureum *areolatum.
terminalis Link, Fil. sp. 122. 1841 = P. phymatodes.
trilobata Fée, Gen. 266. 1850-52 = P. aureum *areolatum.

CIBOTIUM Kaulfuss, Jahrb. d. Pharmacie **1820** —; Enum. Fil. 229. 1824 — (Cyatheaceae Gen. **8**).

adiantoides Pr. Tent. 69. 1836 = Saccoloma domingense.
assamicum Hk. sp. **1**. 83 t. 29 B. 1844 = C. barometz.
barometz (L.) J. Sm. Lond. JoB. **1**. 437. 1842. Chr. 315. NPfl. 121 — China austr.-Assam-Malesia. — Polypodium? L. sp. **2**. 1092. 1753; Aspidium Willd. 1810; Nephrodium Sweet 1827; Dicksonia Link 1841 (Baranetz). HB. 49 — Balantium glaucescens Link 1841; Cibotium Kze. 1841. C. glaucum J. Sm. 1841 — C. assamicum Hk. 1844; Dicksonia Griff. 1849 — Cibotium Cumingii Kze. 1841; 1850 — C. djambianum Hassk. 1856.
Billardieri Klf. Enum. 230. 1824 = Dicksonia antarctica.
Chamissoi Klf. Enum. 230 t. 1 f. 14. 1824. Chr. 316. NPfl. 121 — Ins. Hawaii. Dicksonia Hk. Bak. 1866. HB. 50. Pinonia splendens Gaud. 1824; Dicksonia Desv. 1827.
Cumingii Kze. Farnkr. **1**. 64. 1841; Miq. Verh. Nederl. Inst. III. **4**. 46. t. 5. 1851 = C. barometz var.
djambianum Hassk. Obs. Fil. Jav. **1**. 61. 1856 = C. barometz.
glaucescens Kze. Farnkr. **1**. 63 t. 31. 1841 = C. barometz.
?glaucophyllum Pr. Tent. 69. 1836 = C. barometz.
glaucum (Sm.) Hk. et Arn. Bot. Beech. Voy. 108. 1832. NPfl. 121 — Ins. Hawaii. Dicksonia Sm. in Rees, Cycl. **11** —. 1808. HB. 49; Hiatea Menzies msc.
glaucum J. Sm. JoB. **3**. 418. 1841; Bedd. Ferns br. Ind. t. 83. 1866 = C. barometz.
guatemalense Reichb.; Kuhn, Linn. **36**. 152. 1869. NPfl. 121 — Guatemala. Costa Rica. — Dicksonia Bak. 1874. HB. 461.
Hasskarlianum Miq.; Salom. Nom. 138. 1883 (nomen) — Sumatra.
horridum Liebm. Vid. Selsk. Skr. V. **1**. 279 (seors. 127). 1849 — Mexico.
magnificum de Vriese; Kze. Bot. Zeit. **1848**. 214 = Dicksonia javanica.
Menziesii Hk. sp. **1**. 84 t. 29 C. 1844. NPfl. 121 — Ins. Hawaii. Dicksonia Hk. Bak. 1866. HB. 49, 460 — Cibotium pruinatum Mett.; Kuhn 1869.

proliferum Pr. Tent. 69 t. 11. f. 10. 1836 = Athyrium.
pruinatum Mett.; Kuhn, Linn. **36**. 150. 1869 = C. Menziesii var.
regale Linden; J. Sm. Hist. Fil. 256. 1875 — Mexico.
Dicksonia Bak. Syn. 461. 1874.
Schiedei Schlecht. et Cham. Linn. **5**. 616. 1830. Chr. 316. NPfl. 121 — Mexico-Guatemala. — Dicksonia Bak. 1866. HB. 50.
Siempay Teysm. et Binn.; Salom. Nom. 138. 1883 (nomen) — Sumatra.
Wendlandi Mett.; Kuhn, Linn. **36**. 151. 1869. NPfl. 121 — Guatemala. Mexico.
Dicksonia Bak. 1874. HB. 460.

CINCINALIS Gleditsch, Syst. Plant. 290. 1764 = **Pteridium.**
aquilina Gled. Syst. pl. 290. 1764; Trevis. Syll. Sporoph. Ital. **1** —. 1874.
Gleditschii Güldenst. Reise durch Russl. **1**. 284. 1787 — Russia. (prob. = P aquilinum).

CINCINALIS Desvaux, Berl. Mag. **5**. 311. 1811.
chilensis Fée in Gay, Fl. chil. **6**. 497. 1853 = Pellaea.
ciliata Desv. Berl. Mag. **5**. 312. 1811 = Notholaena hirsuta.
cordata Desv. Berl. Mag. **5**. 311. 1811 = Ceterach.
dealbata Fée, Gen. 160. 1850-52 = Pellaea.
Fendleri Fée, Gen. 160. 1850-52 = Pellaea.
ferruginea Desv. Berl. Mag. **5**. 311. 1811 = Notholaena bonariensis.
?flavens Desv. Berl. Mag. **5**. 329. 1811 = Pellaea.
hirsuta Desv. Berl. Mag. **5**. 313. 1811 = Notholaena.
Hookeri J. Sm. Ferns br. and for. 178. 1866 = Pellaea nivea.
javensis Desv. Berl. Mag. **5**. 329. 1811 = Notholaena hirsuta.
Maranthae Desv. Berl. Mag. **5**. 312. 1811 = Notholaena Marantae.
nivea Desv. Berl. Mag. **5**. 313. 1811 = Pellaea.
pulchella J. Sm. Ferns br. and. for. 178. 1866 = Pellaea.
setigera Desv. Berl. Mag. **5**. 311. 1811 = Woodsia ilvensis.
subcordata Desv. Berl. Mag. **5**. 312. 1811 = Notholaena Marantae.
tenera Fée, Gen. 160. 1850-52 = Pellaea.
tomentosa Desv. Berl. Mag. **5**. 312. 1811 = Notholaena bonariensis.
trichomanoides Desv. Berl. Mag. **5**. 311. 1811 = Notholaena.
vellea Desv. Berl. Mag. **5**. 312. 1811 = Notholaena.
vestita Desv. Berl. Mag. **5**. 312. 1811 = Cheilanthes lanosa.

CIONIDIUM Moore, Gard. Comp. **1852**. 143; Proc. Linn. Soc. **2**. 212. 1854 = [**Aspidium.**
Moorei Moore, Gard. Comp. **1852**. 143.

Clathropeltis Kuhn, v. Decken's Reisen **3**[3]. Bot. 51. 1879 *(Polypodium §)* = Polypodium.

CLEMENTEA Cavanilles, Descripc. plant. 553. 1803 = **Angiopteris.**
palmiformis Cav. Descr. 554. 1803 = ? A. evecta.

CNEMIDARIA Presl, Tent. Pter. 56. 1836 = **Hemitelia.**
horrida Pr. Tent. 57. 1836.
Kohautiana Pr. Tent. 57 t. 1 f. 18. 1836 = H. grandifolia.
munita Pr. Tent. 57. 1836.
obtusa Pr. Tent. 57. 1836 = H. grandifolia.
speciosa Pr. Tent. 57 t. 1 f. 16, 17. 1836 = H. subincisa.

CNEMIDOPTERIS Reichb. Nomencl. Ind. 148, 235. 1841 = Cnemidaria.

COCHLIDIUM Kaulfuss, Jahrb. d. Pharmacie. Berlin **1820**. 36; Enum. Fil. 86. [1824 = **Monogramma.**
graminoides Klf. Enum. 86. 1824.

COELOPTERIS A. Braun msc. (Mett. Polyp. 50. 1857) = **Polypodium** (obliquatum).

Colposoria Presl, Tent. Pter. 128. 1836 *(Davallia §)* = Davallia.
COLYSIS Presl, Epim. Bot. 146. 1849 = **Polypodium.**
affine J. Sm. Hist. Fil. 101. 1875 = P. commutatum.
dilatatum J. Sm. Hist. Fil. 101. 1875 = P. euryphyllum.
erioides J. Sm. Hist. Fil. 101. 1875 (err. : irioides) = P. punctatum.
hemionitidea Pr. Epim. 147. 1849.
insigne J. Sm. Hist. Fil. 101. 1875.
Labrusca J. Sm. Hist. Fil. 101. 1875 = Aspidium labrusca.
longissima J. Sm. Hist. Fil. 101. 1875.
macrophylla Pr. Epim. 147. 1849.
marginata Pr. Epim. 148. 1849 = P. selliguea.
membranacea Pr. Epim. 147. 1849 = P. selliguea.
membranacea J. Sm. Cat. cult. ferns 11. 1857.
normale J. Sm. Hist. Fil. 100. 1875.
ovata J. Sm. Hist. Fil. 100. 1875 = P. phyllomanes var.
pothifolia Pr. Epim. 148. 1849 = P. ellipticum.
Samarense J. Sm. Hist. Fil. 101. 1875 = P. tenuilore.
Spectra J. Sm. Ferns br. and for. 98. 1866.
superficiale J. Sm. Hist. Fil. 101. 1875.
tenuiloris J. Sm. Hist. Fil. 101. 1875.
tridactylis Fée, Gen. 176. 1850-52 (nomen) = P. digitatum.
tridactylon J. Sm. Hist. Fil. 101. 1875 = P. pteropus.
Wallichiana Pr. Epim. 147. 1849 = P. scolopendrinum.
Zippelii J. Sm. Hist. Fil. 101. 1875.
zosteraeforme J. Sm. Hist. Fil. 100. 1875.
Compteris Brazzaiana Linden, Gard. Chr. III. **29**. 25 may, Suppl. 2. 1901 — Hort. (Africa occ. trop.) — ? Pteris sp.
Condyloneura Christ, Mon. Elaph. 22. 1899 *(Elaphoglossum §)* = Elaphoglossum.
CONIOGRAMME Fée, Gen. Fil. 167. 1850-52 — (Polypodiaceae Gen. **62**).
falcata Salom. Nom. 139. 1883 = C. fraxinea.
fraxinea (Don) Diels, Nat. Pfl. **1**^{4}. 262. 1899 — Japonia. China. Asia, Australia, Polynesia, Africa trop. Madagascar.
Diplazium Don, Prod. Fl. Nepal. 12. 1825; Gymnogramme Bedd. 1876; Syngramme Bedd. 1883; Neurogramme Christ 63. 1897. Gymnogramma javanica Bl. 1828. HB. 381; Coniogramme Fée 1850-52; Neurosorus Trev. 1851 — Gymnogramme serrulata Bl. 1828; Coniogramme Fée 1850-52; Grammitis Thwait. 1864 — Gymnogramme pilosa Brack. 1854 (Hawaii).
indica Fée, 10 mém. 22. 1865 = C. fraxinea.
japonica (Thbg.) Diels, Nat. Pfl. **1**^{4}. 262. 1899 — Japonia Formosa.
Hemionitis Thbg. Fl. Jap. 333. 1784; Gymnogramma Desv. 1811. HB. 390; Dictyogramme (Pr. 1849) Fée 1850-52; Notogramme Pr. 1849 (Dyctiogramme).
javanica Fée, Gen. 167 t. 14 B. f. 1. 1850-52 = C. fraxinea.
procera Fée, 10 mém. 22. 1865 = C. fraxinea.
Serra Fée, Gen. 167. 1850-52 = C. fraxinea.
serrulata Fée, Gen. 167 t. 14 B. f. 2. 1850-51 = C. fraxinea var.
COPTOPHYLLUM Gardner, Lond. Journ. Bot. **1**. 133. 1842 — **Aneimia.**
buniifolium Gardn. Lond. JoB. **1**. 133. 1842.
cicutarium Kl. Linn. **18**. 527. 1844.
millefolium Gardn. Lond. JoB. **1**. 133. 1842.

CORMOPHYLLUM Newman, Phytol. **5**. 237. 1854.
arboreum Newm. Phytol. **5**. 238. 1854 = Cyathea.
capense Newm. Phytol. **5**. 238. 1854 = Hemitelia.
horridum Newm. Phytol. **5**. 238. 1854 = Hemitelia.
COSENTINIA Todaro, Syn. pl. acot vasc. Sicil. 14. 1866 = **Notholaena**.
catanensis Todaro, l. c. 16. 1866 } = N. vellea.
vellea Todaro, l. c. 15. 1866 }
CRASPEDARIA Link, Fil. spec. 117. 1841 = **Polypodium.**
auriseta Fée, Gen. 264. 1850-52 = P. tectum.
borbonica Fée, 10 mém. 34 t. 39 f. 2. 1865 = P. lycopodioides.
calva Fée, Gen. 264. 1850-52 = P. neglectum.
camptocarpa Fée, Cr. vasc. Br. **2**. 65 t. 98 f. 1. 1872-73.
cardiophylla Fourn. Bull. Soc. Fr. **19**. 252. 1872 = ? P. cordatum Desv.
chinensis Link, Fil. sp. 118. 1841 = Cyclophorus lingua.
ciliata Link, Fil. sp. 117. 1841.
cordifolia Fée, Cr. vasc. Br. **1**. 118 t. 36 f. 1. 1869 = P. vacciniifolium.
crispata Fée, Cr. vasc. Br. **1**. 119 t. 36 f. 2. 1869 = P. galatheae.
cuspidiflora Fée, Gen. 264. 1850-52 = P. accedens.
Gestasiana Fée, 6 mém. 15 t. 4 f. 2. 1853 = P. tectum.
grandis Fée, Cr. vasc. Br. **1**. 119 t. 37 f. 2. 1869 = P. lycopodioides.
Javanensis Fée, 7 mém. 65. 1857; 10 mém. 33 t. 39 f. 1 = P. accedens.
lagopodioides Fée, Gen. 264. 1850-52 = P. lycopodioides.
lanceolata Fée, 7 mém. 65. 1857 = P. ciliatum.
nummularia Fée, Gen. 264. 1850-52; 6 mém. 16 t. 5 f. 3 = P. pyrolifolium.
ovalis Pr. Epim. 263. 1849 = Cyclophorus serpens.
pertusa Link, Fil. sp. 118. 1841 = Cyclophorus adnascens.
piloselloides Fée, Gen. 264. 1850-52.
? *Plumierii* Fée, 11 mém. 71. 1866 (Plum. t. 120) — Hispaniola.
rupestris Link, Fil. sp. 118. 1841 = Cyclophorus r.
serpens Pr. Epim. 263. 1849 = Cyclophorus s.
serpens Fée, Gen. 264. 1850-52 = P. runcinatum.
surinamensis Fée, Gen. 264. 1850-52; 7 mém. 64 t. 17 f. 2 = P. lycopodioides*.
vacciniifolia Link, Fil. sp. 117. 1841.
veronicaefolia Fée, Gen. 264. 1850-52 = P. piloselloides.
Craspedoglossa Christ, Mon. Elaphogl. 20. 1899 (*Elaphoglossum §*) = Elaphoglossum.
CRASPEDONEURON Van den Bosch, Hymenoph. Jav. 12 t. 10. 1861 = **Trichomanes.**
Trichomanes § Craspedoneuron v.d.B. Ned. Kr. Arch. **4**. 361. 1859.
album v.d.B. Hym. Jav. 12 t. 7. 1861 }
Braunii v.d.B. Hym. Jav. 15 t. 9. 1861 } = T. pallidum.
glaucofuscum v.d.B. Hym. Jav. 13. 1861 }
pallidum v.d.B. Hym. Jav. 14 t. 8. 1861 }
Craspedophyllum Presl, Hymenoph. 33. 1843 (*Hymenophyllum §*) = Hymenophyllum.
Crepidium Presl, Hymenoph. 23. 1843 (*Didymoglossum §*) = Trichomanes.
CREPIDOMANES Presl, Epim. Bot. (17) 258. 1849 = **Trichomanes.**
humile v.d.B. Hym. Jav. 16. t. 11. 1861.
intramarginale Pr. Epim. 258. 1849.
CRYPSINUS Presl, Epim. Bot. 123. 1849 = **Polypodium.**
nummularius Pr. Epim. 123. 1849 = P. pyrolifolium.
CRYPTERIS Nuttall msc. (Hk. sp. **2**. 149. 1858) = **Pellaea** (andromedifolia).

CRYPTOGENIS Richard msc. (Brongn. Dict. Class. d. hist. nat. **3**. 350. 1823) = **Ceratopteris.**

CRYPTOGRAMMA R. Brown apud Richards. in Franklin, Narr. of a journey 767. 1823 — (Polypodiaceae Gen. **76**).

acrostichoides R. Br. ap. Richards. in Frankl. Narr. of a journ. 754. 767. 1823. NPfl. 279 — Amer. bor. subarct. et temp. — Allosorus Spr. 1827; Gymnogramma Pr. 1836; Phorolobus Fée 1850-52; Cryptogramme crispa var. auctt. HB. 144.

aurata Prantl, Engl. Jahrb. **3**. 413. 1882 = Onychium siliculosum.

Brunoniana Wall. List n. 396. 1828 (nomen); Hk. et Grev. Ic. Fil. t. 158. 1829. NPfl. 280 — Himalaya. China austr. Japania. Gymnogramma Pr. 1836; Allosorus J. Sm. 1841; Phorolobus Fée 1850-52; Cryptogramme crispa var. auctt. HB. 144.

cordifolia Prantl, Engl. Jahrb. **3**. 414. 1882 = Llavea.

crispa (L.) R. Br.; Hk. Gen. Fil. t. 115B. 1842. HB. 144 (excl. varr.) Chr. 156. NPfl. 279. — Europa. Asia occ. ? Chile. — Osmunda L. sp. **2**. 1067. 1753; Pteris All. 1785; Acrostichum Vill. 1789; Polypodium Roth 1794; Onoclea Hoffm. 1795; Allosorus Bernh. (1806); Spr. 1827; Blechnum Hartm. 1820; Phorolobus Desv. 1827; Struthiopteris Wallr. 1831; Stegania onocleoides Gray 1821 — adhuc teste Christ: Pellaea fumariaefolia Phil.; Bak. 1891; Allosorus O. Ktze. 1891 (Chile).

densa Diels, Nat. Pfl. **1**[4]. 280. 1899 = Pellaea.

heterophylla Prantl, Engl. Jahrb. **3**. 414. 1882 = Anopteris hexagona.

Jamesoni Hk. et Grev. Ic. Fil. ad t. 158. 1829 = Cheilanthes marginata.

japonica Prantl, Engl. Jahrb. **3**. 413. 1882 = Onychium.

melanolepis Prantl, Engl. Jahrb. **3**. 413 1882 = Onychium.

pallens Prantl, Engl. Jahrb. **3**. 414. 1882 = Ochropteris.

retrofracta Grev. et Hk. Bot. misc. **3**. 385 t. 112. 1833 = Gymnogramma flexuosa.

robusta Pappe et Raws. Syn. Fil. Afr. austr. 32. 1858 = Doryopteris.

sitchensis (Rupr.) Moore, Ind. LXVII. 1857 — Ins. Sitcha. — Allosorus Rupr. Distr. cr. vasc. Ross. **3**. 47. 1845.

Stelleri (Gmel.) Prantl, Engl. Jahrb. **3**. 413. 1882. NPfl. 280 — Sibiria. Himalaya. Amer. bor. temp. — Pteris Gmel. Nov. Comm. Petr. 12. 519 t. 12 f. 1. 1768; Allosorus Rupr. 1845; Pellaea Bak. 1868; Chr. 157. Pteris gracilis Michx. 1803; Cheilanthes Klf. 1824; Allosorus Pr. 1836; Pellaea Hk. 1858. HB. 145. Allosorus minutus Turcz. 1844.

stricta Prantl; Salom. Nom. 141. 1883 = Onychium.

Cryptophyllum Bot. Zeit. **1**. 316. 1843 (err.) = Coptophyllum.

CRYPTOSORUS Fée, Congr. Sci. France X. sess. **1**. 178. 1843; Gen. 231. 1850-52 = **Polypodium.**

blechnoides J. Sm. Hist. Fil. 87. 1875.

Blumei Fée, Gen. 232. 1850-52 = P. obliquatum.

celebicus J. Sm. Hist. Fil. 87. 1875.

Dionaea Fée, Gen. 232 t. 19 C. f. 1. 1850-52 } = P. venulosum.

elasticus Fée, Gen. 231 t. 19 C. f. 2. 1850-52 } = P. venulosum.

khasianus Fée, 10 mém. 31. 1865.

obliquatus J. Sm. Hist. Fil. 87. 1875.

Seemanni J. Sm. Bonplandia **9**. 262. 1861 = P. blechnoides.

CRYPTOSTIGMA A. Braun msc. (Mett. Fil. Lips. 80. 1856) = **Plecosorus.**

CTEISIUM Michaux. Flor. bor. Amer. **2**. 275. 1803 = **Lygodium.**

paniculatum Michx. Fl. bor. Am. **2**. 275. 1803 = L. palmatum.

CTENOPTERIS Blume, Flora Javae. Fil. 132. 1828; Kunze, Bot. Zeit. **1846**. 425 [= **Polypodium.**

achillaefolia J. Sm. Hist. Fil. 185. 1875.

Adenophorus J. Sm. Hist. Fil. 185. 1875.

confusa J. Sm. Hist. Fil. 184. 1875 = P. jubaeforme.

crispata J. Sm. Bot. Voy. Herald 227 t. 48. 1854 = P. goniopteroides.

decipiens J. Sm. Hist. Fil. 185. 1875 = P. capillare.

delicatula J. Sm. Bot. Voy. Herald 227. 1854.

discolor J. Sm. Hist. Fil. 185. 1875.

firma J. Sm. Hist. Fil. 184. 1875.

flabelliformis J. Sm. Hist. Fil. 184. 1875.

funicula J. Sm. Hist. Fil. 185. 1875.

fuscata Kze. Bot. Zeit. **1846** 425.

glandulosa J. Sm. Hist. Fil. 185. 1875 = P. corticolum.

gracilis J. Sm. Hist. Fil. 185. 1875.

grammitidis J. Sm. Hist. Fil. 185. 1875.

jubaeformis J. Sm. Hist. Fil. 184. 1875.

leucosticta J. Sm. Hist. Fil. 185. 1875 = P. longiusculum.

Lobbiana J. Sm. Hist. Fil. 185. 1875 = P. pediculatum.

mollicoma Kze. Bot. Zeit. **1846**. 425 (mollicola J. Sm. Hist. Fil. 185. 1875).

moniliformis J. Sm. Hist. Fil. 184. 1875.

nutans J. Sm. Hist. Fil. 185. 1875.

parvula J. Sm. Hist. Fil. 185. 1875.

pendula J. Sm. Hist. Fil. 185. 1875.

peruviana J. Sm. Hist. Fil. 184. 1875.

pilipes J. Sm. Hist. Fil. 185. 1875.

platynotus Kze. Zoll. Syst. Verz. 37. 1854 — Java.

plumula J. Sm. Hist. Fil. 185. 1875.

pteropus J. Sm. Hist. Fil. 185. 1875 = P. subsessile.

repandula Kze. Zoll. Syst. Verz. 37. 1854.

rigescens J. Sm. Hist. Fil. 184. 1875.

rufescens Kze. Bot. Zeit. **1846**. 425 = P. obliquatum.

Smithiana Kze.; Zoll. Syst. Verz. 37. 1854 = P. minutum.

solida Kze.; Zoll. Syst. Verz. 37. 1854.

subfalcata Kze. Bot. Zeit. **1848**. 120.

subscabra J. Sm. Hist. Fil. 185. 1875.

subtilis J. Sm. Hist. Fil. 184. 1875.

tenuisecta J. Sm. Hist. Fil. 185. 1875.

trichomanoides J. Sm. Hist. Fil. 184. 1875.

trichosora J. Sm. Hist. Fil. 184. 1875.

truncicola J. Sm. Hist. Fil. 184. 1875.

venulosa Kze. Bot. Zeit. **1846**. 425.

vulgaris Newm. Phytol. **2**. 274. 1846.

CULCITA Presl, Tent. Pterid. 135. 1836 = **Balantium.**

macrocarpa Pr. Tent. 135 t. 5 f. 5. 1836 = B. culcita.

schlimensis Fée, 10 mém. 47 t. 36 f. 3. 1865 = B. conifolium.

CUSPIDARIA Fée, 3 mémoire 8, 25. 1851-52 = **Eschatogramme.**

furcata Fée, 3 mém. 25. 1851-52
subpinnatifida Fée, 3 mém. 26. 1851-52 } = Eschatogramme furcata.
tricuspis Fée, 3 mém. 26. 1851-52

CYATHEA Smith, Mém. Acad. Turin **5**. 416. 1793 — (Cyatheaceae Gen. **5**).
abruptecaudata Fée, Cr. vasc. Br. **1**. 183 t. 62 f. 2. 1869 = C. vestita.
acanthomelas Fée, Cr. vasc. Br. **1**. 177 t. 64 f. 1. 1869 — Brasilia.
aculeata Willd.; Klf. Enum. 255. 1824 = Alsophila armata.
affinis (Forst.) Sw. Schrad. Journn. **1800**[2]. 94. 1801; Carr. in Seem. Fl. Vit. 333. 1873. HB. 27 — Polynesia.
Polypodium Forst. Prod. 83. 1786. Cyathea bisulca Schkuhr 1809.
affinis Brack. Expl. Exp. **16**. 283. 1854 = C. propinqua.
albifrons Vieill.; Fourn. Ann. sc. nat. V. **18**. 351. 1873. HB. 453. NPfl. 129 — N. Caledonia.
alpina Roth, Tent. Fl. Germ. **3**. 99. 1800 = Cystopteris fragilis *regia.
alternans (Wall.) Pr. Abh. böhm. Ges. V. **5**. 347. 1848; Bedd. Handb. Suppl. 2. 1892 — Penang. Borneo. — Polypodium Wall. 1828 (nomen); Hemitelia Hk. Ic. pl. t. 622. 1844. NPfl. 132; Amphicosmia? Moore 1857; Alsophila Hk. 1866. HB. 41. Chr. 327; Schizocaena J. Sm. 1875 — Cyathea sarawakensis Hk. 1865. HB. 23. NPfl. 128 — C. Lobbiana Hk. 1865. HB. 24.
aneitensis Hk. Syn. 26. 1865. NPfl. 129 — Aneiteum. Ternate.
angolensis Welw. »Apont. in. Ann. Cons. Ultram. 358. 1858«; Hk. Syn. 22. 1865. NPfl. 127 — Angola.
anthriscifolia Roth, Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 98. 1800 = Cystopteris fragilis.
appendiculata Bak. Journ. Linn. Soc. **15**. 411. 1876. NPfl. 126 — Madagascar.
arachnoidea Hk. Syn. 24. 1865. NPfl. 128 — Ternate. Queensland.
arborea (L.) Sm. Mém. Ac. Turin **5**. 417. 1793. HB. 16 (pt.). Chr. 319. NPfl. 124 — India occ. Brasilia. Ecuador. — Polypodium L. sp. **2**. 1092. 1753; Disphenia Pr. 1836; Hemitelia Fée 1850-52; Cormophyllum Newm. 1854 — Cyathea Grevilleana Mart. 1834.
arborea α nigriscens Hk. sp. **1**. 17. 1844 = C. nigrescens.
arborea var. concinna Bak.; Jenm. JoB. **1881**. 52 = C. concinna.
arborea β pallida Hassk. Obs. Fil. Jav. **1**. 15. 1856 = C. orientalis.
articulata Fée, 8 mém. 111. 1857 — Mexico.
aspera Sw. Schrad. Journ. **1800**[2]. 93. 1801 = Alsophila.
aspidioides Moritz, Verz. 108. 1845 = Diacalpe.
aspidioides Sod. Rec. 14. 1883; Cr. vasc. quit. 505. 1893. NPfl. 125 — Ecuador.
assimilis Hk. Syn. 24. 1865. NPfl. 128 — Borneo. Celebes.
C. Beccariana Ces. 1876.
attenuata Fée, Cr. vasc. Br. **1**. 178 t. 66 f. 1. 1869 = C. Gardneri.
aurea Kl. Bot. Zeit. **1846**. 101 (nomen); Karst. Fl. Col. **2**. 149 t. 178. 1869. HB. 451. NPfl. 126 — Columbia.
aureo-nitens Christ, Bull. Boiss. II. **4**. 948. 1904 — Costa Rica.
azuayensis Sod. Cr. vasc. quit. 644. 1893 — Ecuador.
balanocarpa Eat. Mem. Amer. Ac. n. s. **8**. 215. 1860. HB. 17. NPfl. 124 — Cuba.
barbata Bory; Kuhn, Linn. **36**. 164. 1869, HB. 451 — Guadeloupe.
basilaris Christ, Bull. Boiss. II. **4**. 949. 1904 — Costa Rica.
Beccariana Ces. Atti Ac. Napoli **7**[8]. 3. 1876 = C. assimilis.
Beyrichiana Pr. Tent. 55. 1836 = Hemitelia setosa.
bicrenata Liebm. Vid. Selsk. Skr. V. **1**. 289 (seors. 137). 1849 = Alsophila.
bisulca Schkuhr, Kr. Gew. **1**. sub t. 132 b. 1809 = C. affinis.
boconensis Karst. Linn. **28**. 458. 1856; Fl. Col. **2**. 171 t. 190. 1869. HB. 452. NPfl. 125 — Columbia.

Boivini Mett.; Kuhn, Fil. Afr. 162. 1868. HB. 452. NPfl. 127 — Madagascar. Ins. Comorae.

Bonii Christ, Journ. de Bot. **4**. 410. 1890 = Alsophila podophylla.

borbonica Desv. Berl. Mag. **5**. 328. 1811 — Madagascar, Ins. Mascarenae. Congo. C. canaliculata Willd.; Klf. 1824 (nomen); Spr. Syst. **4**. 26. 1827. HB. 23. Chr. 321. NPfl. 126. C. mascarena »Sw.« Desv. 1827. — var. latifolia Hk. sp. **1**. 24. t. 13 A. 1844; C. Commersoniana Fée 1850-52 — C. laevigata Willd.; Klf. 1824; Schizocaena J. Sm. 1842; Cyathea Pervilliana Fée 1850-52 — C. madagascariensis Klf. 1824. C. discolor Fée 1850-52. — var. Congi Christ, Bull. Soc. bot. Belg. **36**[2]. 92. 1897 (Congo).

Borjae Sod. Cr. vasc. quit. 504. 1893. NPfl. 125 — Ecuador.

Bourgaei Fourn. Mex. pl. **1**. 135. 1872 = C. insignis.

Boylei F. Muell. South. Science Record **1881**. 33 = C. Cunninghami.

Brackenridgei Mett. Ann. Lugd. Bat. **1**. 56. 1863. HB. 452. NPfl. 129 — Ins. Salomonis. Samoa.

brevipinna Bak.; Benth. Fl. austr. **7**. 709. 1878. NPfl. 129 — Ins. Lord Howe.

Brunei Christ, Bull. Boiss. II. **4**. 947. 1904 — Costa Rica.

Brunoniana (Wall.) Clarke et Bak. Journ. Linn. Soc. **24**. 409. 1888 — India bor. Alsophila Wall. List n. 7073. 1829 (nomen); Hk. sp. **1**. 52. 1844. HB. 41; Hemitelia Clarke 1880. NPfl. 132; Amphicosmia Bedd. 1883; Alsophila latebrosa var. hemitelioides Scott 1874.

Brunonis Wall. List n. 179. 1828 = C. moluccana.

bulbifera Bernh. Schrad. Journ. **1801**[1]. 1. 1802 = Cystopteris.

Burkei Hk. sp. **1**. 23 t. 17 B. 1844 = C. Dregei.

calva Karst. Fl. Col. **2**. 175 t. 192. 1869. HB. 19. NPfl. 125 — Columbia. — C. firma Mett. 1869.

camerooniana Hk. Syn. 21. 1865. NPfl. 126 — Kamerun.

canaliculata Willd.; Klf. Enum. 260. 1824 = C. borbonica.

capensis Sm. Mém. Ac. Turin **5**. 417. 1793 = Hemitelia.

caribaea Jenm. W. Ind. and Guiana Ferns 57. 1898 — St. Vincent.

castanea Bak. Syn. 451. 1874 — Peru.

?celebica Bl. Enum. 245. 1828 = Alsophila.

Commersoniana Fée, Gen. 352. 1850-52 = C. borbonica v. latifolia.

commutata Spr. Anleit. **3**. 146 t. 4 f. 32. 1804 = Hemitelia horrida.

compta Mart. Denkschr. bot. Ges. Regensb. **2**. 146 t. 2 f. 1, 2. 1822 = Alsophila.

concinna (Bak.) Jenm. Bull. Dep. Jam. n. 26. 4. 1891. NPfl. 125 — Jamaica. C. arborea var. concinna Bak.; Jenm. JoB. **1881**. 52 — C. jamaicensis Bak. 1891 (non Jenm. 1882).

conquisita Jenm. JoB. **1882**. 324. NPfl. 126 — Jamaica. Amer. centr.

conspersa Christ, Bull. Boiss. II. **5**. 260. 1905 — Costa Rica. C. furfuracea Christ, Bull. Boiss. II. **4**. 950. 1904 (non Bak. 1874).

Copelandii Kuhn et Luerss.; Luerss. Abh. Nat. Ver. Bremen **7**. 278. 1882 — Ins. Trinidad (Atlant. austr.)

corallifera Sod. Rec. 11. 1883; Cr. vasc. quit. 508. 1893. NPfl. 125 — Ecuador.

cordata Mett.; Diels, Nat. Pfl. **1**[4]. 126. 1899 = C. marattioides.

crassipes Sod. Rec. 10. 1883; Cr. vasc. quit. 506. 1893. NPfl. 125 — Ecuador.

crenata Christ, Farnkr. 323. 1897 = Hemitelia.

crenulata Bl. Enum. 244. 1828. HB. 23. NPfl. 128 — Java.

Cumingii Bak. Syn. 453. 1874. NPfl. 129 — Ins. Pitcairn. Coral Island.

Cunninghami Hk.fil. in Hk. Ic. pl. t. 985. 1854. HB. 25. NPfl. 129 — N. Zealand. Australia. — C. Boylei F. Muell. 1881.
cuspidata Kze. Linn. **9**. 101. 1834. HB. 17 — Peru.
cynapifolia Roth, Röm. Arch. 2^1. 106. 1799; Tent. Fl. Germ. **3**. 98. 1800 = [Cystopteris fragilis.
cystolepis Sod. Rec. 15. 1883 = Hemitelia.
dealbata (Forst.) Sw. Schrad. Journ. **1800**^{2}. 94. 1801. HB. 26. Chr. 321. NPfl. 129 — N. Zealand. Ins. Chatham. Ins. Lord Howe.
Polypodium Forst. Prod. 83. 1786 — Cyathea tricolor Col. 1883.
decipiens (Scott) Clarke et Bak. Journ. Linn. Soc. **24**. 409. 1888 — India bor. Alsophila Scott; Bedd. Ferns br. Ind. t. 311. 1869; Hemitelia Scott 1874. HB. 455; Amphicosmia Bedd. 1876.
Deckenii Kuhn, v. Deck. Reis. **3**^{3}. Bot. 57. 1879 — Kilimanjaro. (an C. Manniana forma?). — C. sp. indet. Kuhn, Fil. Deck. 27. 1867.
decrescens Mett.; Kuhn, Fil. Afr. 164. 1868. HB. 452 — Madagascar.
Delgadii Pohl; Sternb. Fl. d. Vorwelt **1**. t. B. 1820 = C. vestita.
dentata Sm. Fl. Brit. **3**. 1141. 1804 = Cystopteris fragilis.
denticulata Goldm. Nova Acta **19** Suppl. I. 466. 1843 — Rio Janeiro.
denudans Kze. Linn. **18**. 349. 1844 = C. mexicana.
discolor Bory, Dup. Voy. Bot. **1**. 281. 1828 = Alsophila quadripinnata.
discolor Fée, Gen. 353. 1850-52 = C. borbonica v. madagascariensis.
discolor Bak. Journ. Linn. Soc. **15**. 412. 1876 = C. similis.
dissoluta Bak.; Jenm. JoB. **1881**. 52. NPfl. 126 — Jamaica.
divergens Kze. Linn. **9**. 100. 1834. HB. 18. (ed. I. 17) Chr. 318. NPfl. 125 — Costa Rica. Peru. — C. globularis Pr. 1849.
Dregei Kze. Linn. **10**. 551. HB. 21. Chr. 322. NPfl. 127 — Africa orient. et austr. C. Burkei Hk. 1844.
dulitensis Bak. Kew Bull. **1896**. 40 — Borneo.
Dyeri Sod. Cr. vasc. quit. 515. 1893. NPfl. 125 — Ecuador.
ebenina Karst. Linn. **28**. 461. 1856. HB. 20. Chr. 320. NPfl. 126 — Columbia.
Eggersii Hier. Engl. Jahrb. **34**. 438. 1904 — Ecuador.
elegans Hew. Mag. Nat. Hist. II. **2**. 466. 1838. NPfl. 125 — Jamaica. Cuba. C. minor Eat. 1860. C. arborea v. pallida HB. 16.
elongata Mett.; Karst. Fl. Col. **2**. 159 t. 183 f. 2. 1869 — Columbia.
equestris Kze. Linn. **9**. 100. 1834. HB. 19. Chr. 320. NPfl. 125 — Costa Rica-Peru.
erinacea Karst. Linn. **28**. 453. 1856; Fl. Col. **2**. 159 t. 183 f. 1. 1869 — Columbia.
excelsa Sw. Schrad. Journ. **1800**^{2}. 93. 1801. HB. 22. Chr. 321. NPfl. 127 — [Ins. Mascarenae.
excelsa Kze. Bot. Zeit. **1848**. 284 = C. crenulata.
extensa Sw. Schrad. Journ. **1800**^{2}. 93. 1801 = Alsophila.
Feei Glaziou; Fée, Cr. vasc. Br. **1**. 179 t. 66 f. 2. 1869 — Brasilia.
Filix femina Bertol. Amoen. Ital. 429. 1819 = Athyrium.
firma Mett.; Kuhn, Linn. **36**. 163. 1869 = C. calva.
flaccida Spr. Schrad. Journ. **1799**^{2}. 271 = Dennstaedtia.
flaccida Mett. Ann. sc. nat. V. **2**. 265. 1864 (nomen) — Columbia.
flavovirens Kuhn; Diels, Nat. Pfl. **1**^{4}. 127. 1899 — Madagascar.
fragilis Sm. Mém. Ac. Turin **5**. 417. 1793 = Cystopteris.
frondosa Karst. Fl. Col. **1**. 149 t. 74, 100. 1860. HB. 18. NPfl. 125 — Columbia.
fulva Fée, 9 mém. 33 (25). 1857 = C. schanschin var.
fulva Sod. Rec. 13. 1883 = C. Sodiroi.
funebris Linden; Cat. 1871 c. fig. — N. Caledonia.
Fourniera Fourn. L'Ill. Hort. **23**. 99. 1876.

furfuracea Bak. Syn. 450. 1874. NPfl. 126. — Jamaica.
C. muricata Gris. 1864 (non Willd. 1810).
furfuracea Christ, Bull. Boiss. II. **4**. 950. 1904 = C. conspersa.
fusca Bak. Malesia **3**. 31. 1886. NPfl. 129 — N. Guinea.
Gardneri Hk. sp. **1**. 21 t. 10 A. 1844. HB. 20. Chr. 320. NPfl. 125 — Brasilia.
— C. incurvata Kze. 1849 — C. attenuata, mamillata, Taunaysiana Fée 1869.
glauca Bory, Voy. **2**. 206. 1804. Chr. 321 — Réunion.
glauca Fourn. Mex. pl. **1**. 135. 1872 = ? C. mexicana vel ? C. arborea.
globularis Pr. Epim. 30. 1849 = C. divergens.
Goudotii Kze. Bot. Zeit. **1844**. 283 — Madagascar.
gracilis Gris. Fl. br. W. Ind. 704. 1864. HB. 18. NPfl. 125 — Jamaica. Columbia.
grandifolia Willd. sp. **5**. 490. 1810 = Hemitelia.
grenadensis Trevis. Atti Ist. Veneto II. **2**. 164. 1851 = C. schanschin.
Grevilleana Mart. Ic. Cr. Bras. 78. 1834 = C. arborea.
guadalupensis Spr. Nova Acta **10**. 233. 1821 = C. arborea.
hastulata Christ, Bull. Boiss. II. **4**. 945. 1904 — Costa Rica.
Havilandii Bak. Tr. Linn. Soc. II. Bot. **4**. 249. 1894. NPfl. 128 — Borneo.
hexagona Fée et Schaffn.; Fée, 8 mém. 111. 1857 = C. mexicana.
Hildebrandtii Kuhn, Ind. Sem. Ht. Berol. **1875**. 20. NPfl. 127 — Ins. Comorae.
hirsuta Pr. Del. Prag. **1**. 190. 1822 = Alsophila armata.
hirsuta Bak. JoB. **1884**. 140 = C. hirsutifrons.
hirsutifrons C. Chr. Ind. 192. 1905 — Madagascar.
C. hirsuta Bak. JoB. **1884**. 140 (non Pr. 1822).
hirtula Mart. Ic. Cr. Bras. 76 t. 53. 1834 = C. vestita.
Holstii Hier. in Engl. Pflw. Ostafr. C. 88. 1895. NPfl. 126 — Africa orient.
Hookeri Thwait. En. pl. Zeyl. 396. 1864. HB. 16. Chr. 318. NPfl. 127 — Ceylon.
horrida Sm. Mém. Ac. Turin **5**. 416. 1793 = Hemitelia.
Humblotii Bak. JoB. **1884**. 140 — Madagascar.
humilis Hier. in Engl. Pflw. Ostafr. C. 88. 1895. NPfl. 126 — Africa orient.
hymenodes Mett. Ann. Lugd. Bat. **1**. 57. 1863. HB. 24. NPfl. 128 — Sumatra.
hypotricha Christ, Bull. Boiss. II. **4**. 947. 1904 — Costa Rica.
Imrayana Hk. sp. **1**. 18 t. 9 B. 1844 = C. Tussacii.
incana Karst. Fl. Col. **1**. 75 t. 37. 1859. HB. 18. NPfl. 126 — Columbia-Ecuador.
incisa Sm. Engl. Bot. **3** t. 163. 1794 = Cystopteris fragilis *regia.
inciso-crenata Bak. Syn. 452. 1874. NPfl. 129 — N. Caledonia.
incurvata Kze. Linn. **22**. 579. 1849 = C. Gardneri.
inquinans Christ, Verh. Nat. Ges. Basel. **11**. 422. 1896; Ann. Buit. **15**. 83 t. 13 f. 5. 1897. NPfl. 128 — Celebes.
insignis Eat. Mem. Amer. Ac. n. s. **8**. 215. 1860. HB. 17. NPfl. 125 — Jamaica. Cuba. Mexico. — C. princeps J. Sm. 1866. C. Bourgaei Fourn. 1872.
integra J. Sm. JoB. **3**. 419. 1841 (nomen); Hk. Ic. pl. t. 638. 1844. HB. 23. NPfl. 129 — Ins. Philipp. Amboina. — C. petiolata J. Sm. 1841.
interjecta Bak. Syn. 453. 1874. NPfl. 129 — N. Caledonia.
jamaicensis Jenm. JoB. **1882**. 323. NPfl. 124 — Jamaica.
jamaicensis Bak. Ann. Bot. **5**. 184. 1891 = C. concinna.
javanica Bl. Enum. 245. 1828. HB. 23. NPfl. 128 — Java. Sumatra.
Jurgensenii Fourn. Mex. pl. **1**. 135. 1872 — Mexico.
Kirkii Hk. Syn. 22. 1865. NPfl. 127 — Ins. Comorae.
Korthalsii Mett. Ann. Lugd. Bat. **1**. 57. 1863. HB. 25. NPfl. 128 — Sumatra. [? Java.
laevigata Willd.; Klf. Enum. 256. 1824 = C. borbonica var.

lanuginosa Jungh. Reisen durch Java 484. 1845; Flora **1847**. 525 = Alsophila tomentosa.
Lastii Bak. JoB. **1891**. 3 — Madagascar.
Lechleri Mett. Fil. Lechl. **2**. 32. 1859 — Peru.
Lenormandi Vieill.; Bak. Syn. 454. 1874. NPfl. 129 — N. Caledonia.
lepidopoda C. Chr. Ind. 193. 1905 — Ecuador.
C. squamipes Sod. Cr. vasc. quit. 514. 1893. NPfl. 126 (non Kl.; Karst. 1861).
leptochlamys Bak. Journ. Linn. Soc. **22**. 535. 1887 — Madagascar.
leucolepis Mett. Ann. Lugd. Bat. **1**. 56. 1863. HB. 26. NPfl. 129 — Aneiteum. Samoa.
leucophaës Hassk. JoB. **7**. 323. 1855; Obs. Fil. Jav. **1**. 26. 1856 — Java.
leucosticta Fée, Cr. vasc. Br. **1**. 182 t. 65. 1869 — Brasilia. Paraguay.
ligulata Bak. JoB. **1884**. 140 — Madagascar.
Lindeniana Pr. Epim. 30. 1849 = C. schanschin vel C. mexicana. (Columbia).
Lindsayana Hk. Syn. 25. 1865 — Australia bor.
Lobbiana Hk. Syn. 24. 1865 = C. alternans.
Macarthurii (F. Muell.) Bak. JoB. **1874**. 280. NPfl. 129 — Ins. Lord Howe.
Hemitelia F. Muell. Fragm. 8. 176. 1874; Cyathea Moorei Bak. 1874. HB. 453.
Macgregori F. Muell. Tr. Roy. Soc. Victoria **1**. 40. 1889; F. Muell. et Bak. JoB. **1890**. 104. NPfl. 129 — N. Guinea.
madagascariensis Klf. Enum. 257. 1824 = C. borbonica var.
mamillata Fée, Cr. vasc. Br. **1**. 176 t. 63 f. 2. 1869 = C. Gardneri.
Manniana Hk. Syn. 21. 1865. Chr. 322. NPfl. 127 — Africa occ. trop.
marattioides Willd.; Klf. Enum. 256. 1824 — Madagascar.
Hemitelia cordata Desv. 1827; Cyathea Mett.; Diels. NPfl. 126. 1899; C. canaliculata var. marattioides Bak. HB. 452.
marianna Gaud. Freyc. Voy. Bot. 74. 1827 = Alsophila extensa.
mascarena »Sw.« Desv. Prod. 322. 1827 = C. borbonica.
Mastersiana L. Linden, Gard. Chr. III. **15**. 663. 1894 — Hort.
medullaris (Forst.) Sw. Schrad. Journ. **1800²**. 94. 1801. HB. 25. Chr. 321. NPfl. 129. — N. Zealand. Tasmania. Australia temp.
Polypodium Forst. Pl. escul. 74. 1786; Prod. 82. 1786; Sphaeropteris Bernh. 1801 — Cyathea polyneuron Col. 1879.
melanocaula Desv. Prod. 322. 1827 = C. borbonica.
meridensis Karst. Fl. Col. **2**. 161 t. 184. 1869. HB. 450. NPfl. 125 — Columbia.
Mettenii Karst. Fl. Col. **1**. 113 t. 56. 1860. HB. 18. NPfl. 125 — Columbia.
mexicana Schlecht. et Cham. Linn. **5**. 616. 1830. HB. 20. NPfl. 125 — Mexico. — C. hexagona Schaffn.; Fée 1857.
microphylla Mett. Fil. Lechl. **1**. 23 t. 3 f. 1-6. 1856. HB. 21. Chr. 320. NPfl. 125 — Ecuador. Peru. — Diacalpe Moore 1857.
Milnei Hk.; Hk.fil. Hdb. Fl. N. Zeal. 349. 1864. HB. 26. NPfl. 129 — Ins. Kermadec.
minor Eat. Mem. Amer. Ac. n. s. **8**. 215. 1860 = C. elegans.
moluccana R. Br.; Desv. Prod. 322. 1827 — Malesia. — C. Brunonis Wall. 1828 (nomen); Hk. sp. **1**. 15. 1844. HB. 16. Chr. 318. NPfl. 127: Schizocaena J. Sm. 1838.
moniliformis Jenm. W. Ind. and Guiana Ferns 59. 1898 — Amer. centr.
monstrabila Jenm. JoB. **1881**. 275. NPfl. 126 (monstrabella) — Jamaica.
montana Sm. Mém. Ac. Turin **5**. 417. 1793 = Cystopteris.
monticola Pr. Abh. böhm. Ges. V. **5**. 347. 1848 = Alsophila quadripinnata var.
Moorei Bak. Syn. 453. 1874 = C. Macarthurii.
Moorei F. Muell. Cens. **1**. 137. 1882 = Hemitelia.
Moricandiana Kze.; Moore, Ind. 272. 1861 = Hemitelia macrocarpa.

Moseleyi Bak. Journ. Linn. Soc. **15**. 104. 1876. NPfl. 129 — Ins. Admiralitatis.
mossambicensis Bak. Ann. of Bot. **5**. 185. 1891. NPfl. 126 — Mossambique.
Muelleri Bak. JoB. **1890**. 104. NPfl. 129 — N. Guinea.
multiflora Sm. Mém. Ac. Turin **5**. 416. 1793 = Hemitelia.
(conf. Alsophila phalerata).
munita Willd.; Klf. Enum. 260. 1824 = Hemitelia.
muricata Willd. sp. **5**. 497. 1810. HB. 450. NPfl. 125 — Ind. occ. Costa Rica.
Alsophila Desv. 1827; Disphenia Pr. 1848; Hemitelia Fée 1850-52.
muricata Gris. Fl. br. W. Ind. 704. 1864 = C. furfuracea.
?*nigra* Linden; Fourn. L'Ill. Hort. **23**. 73 t. 242. 1876 — N. Caledonia.
nigrescens (Hk.) J. Sm. Ferns br. and for. 242. 1866; Jenm. Bull. Dep. Jam. n. 26. 4. 1891. NPfl. 125 — Jamaica. Cuba.
C. arborea α nigrescens Hk. sp. **1**. 17. 1844. C. arborea pt. HB. 16.
nigricans Mett. Ann. Lugd. Bat. **1**. 56. 1863. HB. 26. NPfl. 129 — Ins. Ualan.
Nockii Jenm. JoB. **1879**. 257. NPfl. 124 — Jamaica. Trinidad.
ocanensis Bak. Ann. Bot. **5**. 184. 1891 — Columbia.
oinops Hassk. JoB. **7**. 322. 1855; Obs. Fil. Jav. **1**. 23. 1856 — Java.
C. sinops Racib. 1898.
oligocarpa Kze. Linn. **9**. 101. 1834 = C. schanschin.
oligocarpa Jungh. Hoev. Tijdschr. 8. 373. 1841; Flora **1847**. 522 — Java.
onusta Christ, Bull. Boiss. II. **4**. 950. 1904 — Costa Rica.
orientalis (Kze.) Moore, Ind. 272. 1861; Mett. Ann. Lugd. Bat. **1**. 58. 1863. HB. 24. NPfl. 127 — Java. Celebes. — Disphenia Kze. Bot. Zeit. **1848**. 283 (pt.).
oyapoka Jenm. W. Ind. and Guiana Ferns 58. 1898 — Guiana.
papyracea Christ, Bull. Boiss. II. **4**. 946. 1904 — Costa Rica.
patellaris Christ, Ann. Cons. Jard. bot. Genève **4**. 207. 1900 — Costa Rica.
patens Karst. Fl. Col. **2**. 173 t. 191. 1869. HB. 451. NPfl. 125 — Costa Rica. [Columbia.
pauciflora Kze. Bot. Zeit. **1846**. 101 = Alsophila.
pelliculosa Christ, Bull. Boiss. II. **4**. 946. 1904 — Costa Rica.
pendula Jenm. JoB. **1882**. 324. NPfl. 126 — Jamaica. Amer. centr.
percussa Cav. Descr. 548. 1803 — Ins. Mariannae.
Polypodium cyathoides Sw. 1806 — (Dryopteris sp. dubia).
Pervilliana Fée, Gen. 352. 1850-52 = C. borbonica var. laevigata.
petiolata J. Sm. JoB. **3**. 419. 1841 = C. integra var.
petiolulata Karst. Fl. Col. **2**. 163 t. 185. 1869. NPfl. 125 — Columbia.
phalerata Mart. Denkschr. bot. Ges. Regensb. **2**. 146 t. 2 f. 3. 1822 = Alsophila.
phanerophlebia Bak. Ann. Bot. **8**. 122. 1894. NPfl. 126 — Madagascar.
philippinensis Bak. Ann. Bot. **5**. 186. 1891 — Ins. Philipp.
pilosa Bak. Syn. ed. II. 19. 1874. NPfl. 126 — Peru.
pinnata Roxb. Calc. Journ. **4**. 517. 1844 = C. moluccana.
polycarpa Jungh. Nat. Geneesk. Arch. **2**. 40. 1845; Flora **1847**. 522 = C. spinulosa.
polyneuron Col. Tr. N. Zeal. Inst. **11**. 429. 1879 = C. medullaris var.
polyphlebia Bak. Journ. Linn. Soc. **20**. 303. 1883. NPfl. 127 – Madagascar.
polypoda Bak. Tr. Linn. Soc. II. Bot. **4**. 250. 1894 — Borneo.
polypodioides Sw. Vet. Ak. Handl. **1817**. 78 = Hemitelia capensis.
portoricensis Spr.; Kuhn, Linn. **36**. 163. 1869. HB. 451. NPfl. 124 — Puerto Rico.
portoricensis Christ, Prim. Fl. costar. **3**. 40. 1901 = C. aureonitens.
Preussii Diels, Nat.Pfl. **1**[4]. 127. 1899 — Kamerun.
princeps J. Sm. Ferns br. and for. 291. 1866; E. Mayer, Gartenfl. **17**. 10. 1868 = C. insignis.

propinqua Mett. Ann. Lugd. Bat. **1**. 56. 1863. HB. 452. NPfl. 129 — Fiji. Samoa.
C. affinis Brack. 1854 (non Sw. 1801).
puberula Sod. Rec. 16. 1883; Cr. vasc. quit. 513. 1893. NPfl. 126 — Ecuador.
pubescens Mett.; Kuhn, Linn. **36**. 164. 1869. HB. 450. NPfl. 124 — Jamaica. Haiti. Puerto Rico.
punctifera Christ, Prim. Fl. costaric. **3**. 40. 1901 — Costa Rica.
Purdiaei Jenm. W. Ind. and Guiana Ferns 58. 1898 — Trinidad.
purpurascens Sod. Cr. vasc. quit. 503. 1893. NPfl. 125 — Ecuador.
pygmaea L. Linden, Gard. Chr. III. **15**. 663. 1894 — Hort.
quadrata Bak. Journ. Linn. Soc. **15**. 411. 1876. NPfl. 126 — Madagascar.
quindiuensis Karst. Linn. **28**. 454. 1856; Fl. Col. **2**. 151 t. 179. HB. 450. NPfl. 125 — Columbia.
regia Forst. in Sym. Syn. 194. 1798 = Cystopteris fragilis*.
regularis Bak. Journ. Linn. Soc. **25**. 349. 1889 — Madagascar.
reticulata Werckle; Christ, Bull. Boiss. II. **5**. 251. 1905 — Costa Rica.
rigidula Bak. Journ. Linn. Soc. **22**. 534. 1887 — Madagascar.
riparia Willd. sp. **5**. 493. 1810 = Hemitelia capensis.
robusta Boj.; Wall. List n. 182. 1828 = C. borbonica.
Ruiziana Kl. Linn. **20**. 439. 1847; Bak. Ann. Bot. **5**. 184. 1891. NPfl. 126 — Peru.
Rumphii Desv. Prod. 323. 1827 (Rumph. t. 27) — India. Amboina. Cochinchina.
Polypodium arboreum Lour. Fl. Cochinch. **2**. 831. 1790 (non L. 1753).
saccata Christ, Ann. Jard. Buit. II. **4**. 42. 1904 — Celebes.
samoensis Bak. JoB. **1876**. 9 — Samoa.
sarawakensis Hk. Syn. 23. 1865 = C. alternans.
Sartorii »Liebm.« Salom. Nom. 144. 1883 (nomen) — Mexico.
scabra Bak. JoB. **1876**. 343. NPfl. 129 — Samoa.
schanschin Mart. Ic. Cr. Bras. 77 t. 29 f. 3, 4, t. 54. 1834. HB. 19. Chr. 319. NPfl. 126 — Amer. trop. — Alsophila fulva Mart. et Gal. 1842; Cyathea Fée 1857 — C. sphaerocarpa Fée 1869.
Schenckii Christ, Hedwigia **35**. 155. 1896. NPfl. 125 — Brasilia.
schizochlamys Bak. JoB. **1880**. 209. NPfl. 128 — Sumatra.
Schlechtendalii Kze. Bot. Zeit. **1845**. 288 — Caracas.
Schottiana Kze. Linn. **23**. 246. 1850 (nomen) — Mexico.
sclerolepis Bak. Syn. 453. 1874. NPfl. 129 — N. Caledonia.
sechellarum Mett. Ann. Lugd. Bat. **1**. 58. 1863. HB. 22. NPfl. 127 — Ins. Sechellae.
segregata Bak. Journ. Linn. Soc. **20**. 303. 1883. NPfl. 127. — Madagascar.
Sellowiana Pr. Tent. 55. 1836 = ? Alsophila paleolata. (t. Mett. = Hemitelia setosa).
serra Willd. sp. **5**. 491. 1810. HB. 17. Chr. 318. NPfl. 125 — Ind. occ. Mexico. Ecuador. — Hemitelia Desv. 1827.
serratifolia Bak. JoB. **1884**. 139 — Madagascar.
similis C. Chr. Ind. 195. 1905 — Madagascar.
C. discolor Bak. Journ. Linn. Soc. **15**. 412. 1876 (non Bory 1828, sed fortasse Fée 1850-52; vide C. borbonica).
sinops Racib. Pterid. Buitenz. 36. 1898 = C. oinops.
sinuata Hk. et Grev. Ic. Fil. t. 106. 1828. HB. 16. Chr. 317. NPfl. 127 — Ceylon.
Schizocaena J. Sm. 1842.
Smithii Hk.fil. Fl. N. Zeal. **2**. 8 t. 72. 1854 = Hemitelia.
societarum Bak. Syn. 453. 1874. NPfl. 129 — Ins. Societatis.
Sodiroi C. Chr. Ind. 195. 1905 — Ecuador. — C. fulva Sod. Rec. 13. 1883; Cr. vasc. quit. 516. 1893. NPfl. 126 (non Fée 1857).

speciosa H.B. Willd. sp. **5**. 490. 1810 = Hemitelia.
sphaerocarpa Fée, Cr. vasc. Br. **1**. 180 t. 53 f. 2. 1869 = C. schanschin.
spinulosa Wall. List n. 178. 1828; Hk. sp. **1**. 25 t. 12 C. 1844. HB. 23. Chr. 321. NPfl. 127. — Japonia. Formosa. Ins. Philipp. China austr. India bor.-Malesia.
Sprucei Hk. Syn. 20. 1865. NPfl. 126 — Ecuador.
squamipes Kl.; Karst. Fl. Col. **1**. 199 t. 99, 100. 1861. HB. 18 — Columbia.
squamipes Sod. Cr. vasc. quit. 514. 1893 = C. lepidopoda.
Sternbergii Pohl; Sternb. Fl. d. Vorwelt **1**. 47 t. G. 1820 = C. arborea.
straminea Karst. Linn. **28**. 457. 1856; Fl. Col. **2**. 157 t. 182, 183 f. 3. 1869. HB. 451. NPfl. 125 — Columbia.
strigosa Christ, Ann. Jard. Buit. **15**. 84 t. 13 f. 6. 1897. NPfl. 129 — Celebes.
Stuhlmanni Hier. Engl. Jahrb. **28**. 340. 1899 — Africa trop.
suluensis Bak. JoB. **1879**. 65. NPfl. 129 — Sulu Archip.
sumatrana Bak. JoB. **1880**. 209. NPfl. 128 — Sumatra.
superfusa Kze.; Ettingsh. Farnkr. 224 t. 160 f. 2, 3. 1865 = C. Gardneri.
Taunaysiana Glaz. et Fée, Cr. vasc. Br. **1**. 178 t. 64 f. 2. 1869 = C. Gardneri.
tenera (J. Sm.) Moore, Ind. 274. 1861; Gris. Fl. br. W. Ind. 704. 1864. HB. 450. NPfl. 124 — St. Vincent. — Alsophila J. Sm. 1842 (nomen); Hk. sp. **1**. 49. 1844.
Thomsoni Bak. JoB. **1881**. 180 — Africa trop. (Nyassa).
Thurstoni Bak. JoB. **1886**. 182 — Fiji.
tomentosa Zoll.; Moritz, Syst. Verz. —. 1845-46 = Alsophila.
Trejoi Christ, Bull. Boiss. II. **5**. 733. 1905 — Mexico.
tricolor Col. Tr. N. Zeal. Inst. **15**. 304. 1883 = C. dealbata.
tripinnatifida Wall. List n. 7076. 1829; Roxb. Calc. Journ. **4**. 518. 1844 — Ins. Moluccae.
Tussacii Desv. Prod. 323. 1827. HB. 17. Chr. 319. NPfl. 125 — Jamaica. Columbia. C. Imrayana Hk. 1844.
usambarensis Hier. in. Engl. Pflw. Ostafr. C. 88. 1895. NPfl. 127 — Africa [orient.
venulosa Wall. List n. 180. 1828 = Alsophila glabra.
vestita Mart. Denkschr. bot. Ges. Regensb. **2**. 146. 1822; Ic. Cr. Bras. 75 t. 52. 1834. HB. 19. NPfl. 126 — Guiana. Brasilia. Columbia. — C. Delgadii Pohl 1820 (nomen?). C. hirtula Mart. 1834 — C. abruptecaudata Fée 1869.
Vieillardii Mett. Ann. sc. nat. IV. **15**. 82. 1861. HB. 27. NPfl. 129 — N. Caledonia.
villosa H.B. Willd. sp. **5**. 495. 1810 = Alsophila.
Walkerae Hk. Ic. pl. t. 647. 1844 = Hemitelia.
Welwitschii Hk. Syn. 21. 1865. NPfl. 126 — Africa occ. trop.
Whitmeei Bak. JoB. **1876**. 343 — Samoa.
woodwardioides Klf. Enum. 255. 1824 —?
zambesiaca Bak. Ann. Bot. 8. 121. 1894. NPfl. 127 — Africa trop. (Nyassa).
Zollingeriana Mett. Ann. Lugd. Bat. **1**. 57. 1863. NPfl. 128 — Java.

Cyatheopsis Karst. Fl. Columb. **1**. 3. 1858 *(Cyathea §)* = Cyathea.

CYCLODIUM Presl, Tent. Pter. 85. 1836 — (Polypodiaceae Gen. **20**).
abbreviatum Pr. Epim. 260. 1849 = Polystichum.
acrostichoides J. Sm. Lond. JoB. **1**. 199. 1842 = Dryopteris canescens.
confertum Pr. Tent. 85. 1836 = C. meniscioides.
Cumingianum Moore, Ind. LXXXIII. 1857 = Dryopteris otaria.
glandulosum Pr. Tent. 85 t. 2 f. 21. 1836 = Dryopteris.
heterodon Moore, Ind. 275. 1861 = ? C. meniscioides.
meniscioides (Willd.) Pr. Tent. 85 t. 2 f. 20. 1836. NPfl. 195 — Trinidad-Brasilia bor. Peru-Ecuador. — Aspidium Willd. sp. **5**. 218. 1810. Mett. n. 70.

HB. 257; Chr. 234; Nephrodium J. Sm. 1841; Dryopteris O. Ktze. 1891. Aspidium confertum Klf. 1824; Mett. n. 69; Cyrtomium Pr. 1836; Nephrodium J. Sm. 1841; Aspidium Hookeri Kl. 1847 — ? Aspidium heterodon Schrad. 1824. Mett. n. 71; Polystichum Pr. 1849; Cyrtomium Moore 1861.

rigidissimum C. Chr. Bot. Tidsskr. **25**. 79. 1902 — Brasilia.
Aspidium C. Chr. l. c.

Serra A. Brongn. Voy. aut. d. Monde de Vénus. Bot. t. 5. 1846 = ? Dryopteris.

Cycloglossum Presl, Hym. 32. 1843 *(Hymenophyllum §)* = Hymenophyllum.

CYCLOPELTIS J. Smith, Bot. Mag. **72**. Comp. 36. 1846 — (Polypodiaceae Gen. **14**).

Presliana (J. Sm.) Berkeley, Introd. Crypt. Bot. 517. 1857. NPfl. 183 — Birma-Malesia-Ins. Philipp. N. Guinea.
Lastrea J. Sm. 1841 (nomen); Polystichum Moore 1857; Apidium Mett. Pheg. u. Asp. 36. n. 81. 1858. Hemicardion crenatum et Cumingianum Fée 1850-52 — Aspidium Kingii Hance 1862.

semicordata (Sw.) J. Sm. Bot. Mag. **72**. Comp. 36. 1846. NPfl. 183 — America trop.
Polypodium Sw. Prod. 132. 1788; Aspidium Sw. 1801. Mett. n. 80. HB. 249 (pt.); Chr. 232; Nephrodium Pr. 1825; Lastrea Pr. 1836; Polystichum Moore 1857; Dryopteris O. Ktze. 1891. Hemicardion Nephrolepis Fée 1850-52. Polypodium caducum Willd. 1810; Aspidium HBK. 1815; Lastrea Pr. 1836.

CYCLOPHORUS Desvaux, Berl. Mag. **5**. 300. 1811 — (Polypodiaceae Gen. **102**).

abbreviatus (Zoll.) C. Chr. Ind. 197. 1905 — Java. Sumatra.
Niphobolus Zoll. Nat. en Gen. Arch. **1**. 397. 1844; Gies. Niph. 160; Sphaerostichum Pr. 1849; Polypodium Mett. 1856. Pol. n. 266.

acrocarpus (Christ et Gies.) C. Chr. Ind. 197. 1905 — China.
Niphobolus Christ et Gies. Bull. Ac. Géogr. Bot. Mans **1902**. 221.

acrostichoides (Forst.) Pr. Epim. 130. 1849 — Ceylon-Malesia-Polynesia. Queensland. — Polypodium Forst. Prod. 81. 1786. Mett. n. 258. HB. 350; Niphobolus Rich. 1834; Gies. Niph. 213. NPfl. 325. Cyclophorus glaber Desv. 1811. C. Scolopendrium Desv. 1827; Niphobolus Moore 1861. N. puberulus Bl. 1828 — N. fissus Bl. 1828 (non alior.); ? Gyrosorium Pr. 1849. — an Acrostichum lanceolatum L. 1753; Candollea Mirb. 1803? an Acrostichum longifolium Burm. 1768? (vide Cyclophorus infra).

adnascens (Sw.) Desv. Berl. Mag. **5**. 300. 1811 — Asia trop. Polynesia.
Polypodium Sw. Syn. 25, 222 t. 2 f. 2. 1806. HB. 349 (pt.). Chr. 98; Niphobolus Klf. 1824. NPfl. 325. Gies. Niph. 196 — Polypodium pertusum Roxb.; Hk. 1825. Mett. n. 252; Niphobolus Spr. 1827; Craspedaria Link 1841 — Niphobolus caudatus Klf. 1824; Polypodium Mett. n. 255. 1857 (non Raddi 1819) — Niphobolus carnosus Bl. 1828; Polypodium Mett. n. 249. 1857 — Niphobolus elongatus Bl. 1828.

africanus (Kze.) C. Chr. Ind. 197. 1905 — Africa austr.
Niphobolus Kze. Linn. **10**. 501. 1839. NPfl. 325. Gies. Niph. 177; Gyrosorium Pr. 1849; Polypodium Mett. n. 268. 1857. HB. 351. Chr. 98 (non Desv. 1827).

albicans (Bl.) Pr. Epim. 131. 1849 — Java.
Niphobolus Bl. Enum. 107. 1828; Fl. Javae, Fil. 60 t. 25. 1828; Gies. Niph. 185; Polypodium Mett. n. 257. 1857. HB. 352. P. acrostichoides Willd. 1810 (non Forst. 1786) — Niphobolus glaber Klf. 1824 (non Cyclophorus Desv. 1811); N. Blumeanus Kze. 1848; Cyclophorus Pr. 1849. — (an adhuc C. flocciger ?).

?**americanus** (Hk.) C. Chr. Ind. 198. 1905 — Ecuador.
Polypodium Hk. sp. **5**. 54. 1863. HB. 352. Chr. 98; Niphidium J. Sm. 1875; Niphobolus Diels 1899. NPfl. 325.

anetioides (Christ) C. Chr. Ind. 198. 1905 — Celebes.
Niphobolus Christ, Ann. Jard. Buit. II. **4**. 38. 1904.

angustatus (Sw.) Desv. Berl. Mag. **5**. 300. 1811 — India bor.-Malesia-Polynesia.
Polypodium Sw. Syn. 27, 224. 1806. HB. 356. Chr. 101; Niphobolus Spr. 1827. NPfl. 326. Gies. Niph. 193; Pleopeltis Pr. 1836; Drynaria? Fée 1850-52; Niphopsis J. Sm. 1857. Polypodium sphaerocephalum Wall. 1828; Mett. n. 244; Pleopeltis Bl. 1828; Niphobolus Hk. et Grev. 1829; Phymatodes Pr. 1836 — Niphobolus macrocarpus Hk. et Arn. 1832.

annamensis (Christ) C. Chr. Ind. 198. 1905 — Annam.
Niphobolus Christ, Journ. de Bot. **19**. 74. 1905.

assimilis (Bak.) C. Chr. Ind. 198. 1905 — China.
Polypodium Bak. JoB. **1875**. 201; Niphobolus Diels 1900. Gies. Niph. 129.

asterosorus (Bak.) C. Chr. Ind. 198. 1905 — Sumatra.
Polypodium Bak. JoB. **1880**. 214.

Beddomeanus (Gies.) C. Chr. Ind. 198. 1905 — China austr. India bor.
Niphobolus Gies. Niph. 101. 1901. Polypodium costatum Wall. 1828 (pt., nomen); Mett. n. 267. 1857 (non Kze. 1834); Niphobolus Pr. 1836; Apalophlebia Pr. 1849. Polypodium stigmosum pt. auctt. — ? Pyrrosia chinensis Mirb. 1803.

blepharolepis C. Chr. Ind. 198. 1905 — Tahiti.
Polypodium tricholepis Mett.; Kuhn, Linn. **36**. 139. 1869 (non Schrad. 1824); Niphobolus Carr. 1873; Gies. Niph. 175. 1901; Polypodium serpens pt. HB. 349.

Blumeanus Pr. Epim. 130. 1849 = C. albicans.

Bonii (Christ) C. Chr. Ind. 198. 1905 — Tonkin.
Niphobolus Christ; Gies. Niph. 120. 1901.

Boothii (Hk.) C. Chr. Ind. 198. 1905 — India bor. — Polypodium Hk. sp. **5**. 53. 1863. HB. 352; Niphobolus Bedd. 1868. Gies. Niph. 115.

calvatus (Bak.) C. Chr. Ind. 198. 1905 — China.
Polypodium Bak. JoB. **1879**. 304; Niphobolus Diels 1900.

Cavalerianus (Christ) C. Chr. Ind. 198. 1905 — China.
Niphobolus Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 107.

ceylanicus (Gies.) C. Chr. Ind. 198. 1905 — Ceylon.
Niphobolus Gies. Niph. 216 f. 19. 1901.

Christii (Gies.) C. Chr. Ind. 198. 1905 — Borneo.
Niphobolus Gies. Niph. 140. 1901.

confluens (R. Br.) C. Chr. Ind. 198. 1905 — Australia. Ins. Norfolk. Ins. Lord Howe. N. Caledonia. N. Zealand. — Polypodium R. Br. Prod. Fl. N. Holl. 146. 1810; Niphobolus Spr. 1827. NPfl. 326. Gies. Niph. 172; N. acrostichoides Klf. 1824; Scytopteris Pr. 1849; Cyclophorus Scytopteris Fourn. 1873. Polypodium glabrum Mett. n. 245. 1857. HB. 356. Chr. 101 (non Burm. 1768); Pleopeltis Fourn. 1869; Drymoglossum Cunninghami Moore 1857.

?**cuneatus** (Kuhn) C. Chr. Ind. 198. 1905 — Ecuador. — Niphobolus Kuhn, Bot. Zeit. **1868**. 40. NPfl. 325; Polypodium Kuhn 1868. HB. 513.

distichocarpus (Mett.) C. Chr. Ind. 198. 1905 — Sumatra. — Polypodium Mett. Ann. Lugd. Bat. **2**. 231. 1866. HB. 352; Niphobolus Gies. 1899. NPfl. 325.

Drakeanus (Franch.) C. Chr. Ind. 198. 1905 — China. — Polypodium Franch. Nouv. Arch. Mus. II. **7**. 165. 1883; Niphobolus Diels 1900. Gies. Niph. 117.

flocciger (Bl.) Pr. Epim. 131. 1849 — India bor.-Malesia. Ins. Philipp.
Niphobolus Bl. Enum. 107. 1828. NPfl. 325; Polypodium Mett. n. 262. 1857. HB. 351 — Antrophyum niphoboloides Kze. 1848; Taenitis Moore 1857 — (an ad C. albicantem pertinens?).

flocculosus (Don) C. Chr. Ind. 199. 1905 — India bor.
Polypodium Don, Prod. Fl. Nepal. 1. 1825. HB. 351; Niphobolus Spr. 1827. Gies. Niph. 125; Apalophlebia? Pr. 1849; Polypodium detergibile Hk. 1863; Niphobolus Bedd. 1866.

Gardneri (Kze.) C. Chr. Ind. 199. 1905 — Ceylon. — Niphobolus Kze. hb.; J. Sm. 1857. Gies. Niph. 145; Polypodium Mett. Pol. 129. n. 263. 1857. HB. 352.

Giesenhagenii (Christ) C. Chr. Ind. 199. 1905 — Bhotan.
Niphobolus Christ, Ann. Cons. Jard. bot. Genève 7-8. 330. 1904.

glaber Desv. Berl. Mag. **5**. 301. 1811 = C. acrostichoides.

gralla (Gies.) C. Chr. Ind. 199. 1905 — Yunnan.
Niphobolus Gies. Niph. 128. 1901.

hastatus (Thbg.) C. Chr. Ind. 199. 1905 — Japonia. Korea.
Acrostichum Thbg. Fl. Jap. 331 t. 34. 1784; Niphobolus Kze. 1848; Polycampium Pr. 1849. Polypodium tricuspe Sw. 1801. HB. 350. Chr. 97; Niphobolus J. Sm. 1875. NPfl. 325. Gies. Niph. 165.

heteractis (Mett.) C. Chr. Ind. 99. 1905 — India bor.
Polypodium Mett.; Kuhn, Linn. **36**. 140. 1869. (HB. 512). (Chr. 97); Niphobolus J. Sm. 1877. Gies. Niph. 153. Polypodium Lingua pt. auctt.

heterophyllus Desv. Berl. Mag. **5**. 301. 1811 = C. spissus.

inaequalis (Christ) C. Chr. Ind. 199. 1905 — China.
Niphobolus Christ, Bull. Soc. Fr. **52** Mém. I. 25. 1905.

laevis (J. Sm.) C. Chr. Ind. 199. 1905 — India bor.
Niphobolus J. Sm.; Bedd. Ferns br. Ind. t. 161. 1866; Gies. Niph. 141; Polypodium Mett.; Kuhn, Linn. **36**. 139. 1869. P. jaintense Clarke 1880.

lanuginosus (Gies.) C. Chr. Ind. 199. 1905 — Ins. Philipp.
Niphobolus Gies. Niph. 190. 1901.

Lauterbachii (Christ) C. Chr. Ind. 199. 1905 — N. Guinea. — Niphobolus Christ in Schum. et Laut. Fl. deut. Südsee 142. 1901; Gies. Niph. 221 f. 20.

linearifolius (Hk.) C. Chr. Ind. 199. 1905 — Japonia. Korea. China. Formosa.
Niphobolus Hk. 2 Cent. t. 58. 1861. NPfl. 326. Gies. Niph. 171; Polypodium Hk. 1863. HB. 356.

lingua (Thbg.) Desv. Prod. 224. 1827 — Japonia. China. Tonkin. Formosa.
Acrostichum Thbg. Fl. Jap. 330 t. 33. 1784; Polypodium Sw. 1806; Mett. n. 264. HB. 350 (pt.). Chr. 97; Niphobolus Spr. 1827. NPfl. 325. Gies. Niph. 156; Polycampium Pr. 1849.

longifolius (Burm.) Desv. Berl. Mag. **5**. 301. 1811 — Java.
Acrostichum Burm. Fl. Ind. 228. 1768; Candollea Mirb. 1803; Niphobolus Spr. 1827. (an = C. acrostichoides?).

macropodus (Bak.) C. Chr. Ind. 199. 1905 — Ins. Aru.
Polypodium Bak. Journ. Linn. Soc. **15**. 108. 1876.

malacophyllus (Christ) C. Chr. Ind. 199. 1905 — Yunnan. — Polypodium Christ in litt.; P. mollissimum Christ. Bull. Boiss. **7**. 5. 1899 (non Fée 1866).

Martini (Christ) C. Chr. Ind. 199. 1905 — China. Thibet orient.
Niphobolus Christ, Bull. Soc. Fr. **52** Mém. I. 23. 1905.

mollis (Kze.) Pr. Epim. 131. 1849 — Java. Timor. Ad flum. Iravaddi.
Niphobolus Kze. Bot. Zeit. **1848**. 121; Polypodium Mett. n. 261. 1857.

nudus (Gies.) C. Chr. Ind. 200. 1905 — India bor. China austr.
Niphobolus Gies. Niph. 149. 1901.

nummularifolius (Sw.) C. Chr. Ind. 200. 1905 — India bor.-Malesia-Ins. Philipp.
Acrostichum Sw. Syn. 191, 419 t. 2 f. 1. 1806; Pteropsis Desv. 1827; Gymnopteris Pr. 1836; Niphobolus J. Sm. 1841. NPfl. 325. Gies. Niph. 179; Galeoglossa Pr. 1849; Polypodium Mett. n. 247. 1857; HB. 351; Chr. 99 — Galeoglossa rotundifolia Pr. 1849; Niphobolus Fée 1850-52 — Acrostichum obovatum Bl. 1828; Niphobolus Kze. 1848; Galeoglossa Pr. 1849; Polypodium Mett. n. 248. 1857.

pachydermus (Bak.) C. Chr. Ind. 200. 1905 — Ins. Moluccae.
Polypodium Bak. Journ. Linn. Soc. **15**. 108. 1876.

pannosus (Mett.) C. Chr. Ind. 200. 1905 — Ceylon. Birma.
Polypodium Mett.; Kuhn, Linn. **36**. 141. 1869. HB. 512; Niphobolus Bedd. 1876; Gies. Niph. 105; N. Lingua Bedd. 1863-65.

pekinensis C. Chr. Ind. 200. 1905 — China.
Polypodium Davidii Bak. Ann. Bot. **5**. 472. 1891 (non Franch. 1887); Niphobolus Gies. 1899. Niph. 132. NPfl. 325.

penangianus (Hk.) C. Chr. Ind. 200. 1905 — Birma. Perak. Penang. Sumatra. Java. — Niphobolus Hk. Ic. pl. t. 203. 1840. Gies. Niph. 97; Polycampium Pr. 1849; Polypodium Hk. 1863. HB. 352.

petiolosus (Christ) C. Chr. Ind. 200. 1905 — Korea. Amur. China.
Polypodium Christ, Nu. Giorn. bot. It. n. s. **4**. 96 t. 1 f. 2. 1897; Niphobolus Diels 1900. Gies. Niph. 146.

polydactylus (Hance) C. Chr. Ind. 200. 1905 — Formosa. — Polypodium Hance, JoB. **1883**. 269; Niphobolus Gies. 1899. Niph. 167. NPfl. 325.

porosus (Wall.) Pr. Epim. 130. 1849 — India bor. China.
Polypodium Wall. 1828 (nomen); Mett. Pol. 128. n. 259. 1857; Niphobolus Pr. 1836. Polypodium fissum pt. auctt. HB. 351. Chr. 98; ? Niphobolus Bedd. 1870. NPfl. 325 (non Bl. 1828). N. Mannii Gies. 1901. Niph. 107.

princeps (Mett.) C. Chr. Ind. 200. 1905 — N. Guinea. — Polypodium Mett. Ann. Lugd. Bat. **2**. 232. 1866. (HB. 351); Niphobolus Gies. 1899. NPfl. 325.

Rasamalae (Racib.) C. Chr. Ind. 200. 1905 — Java.
Polypodium Racib. Pterid. Buit. 99. 1898; Niphobolus Gies. Niph. 187. 1901.

rupestris (R. Br.) C. Chr. Ind. 200. 1905 — Australia. ? N. Zealand.
Polypodium R. Br. Prod. Fl. N. Holl. 146. 1810. Mett. n. 250; Niphobolus Spr. 1827. Gies. Niph. 169; Craspedaria Link. 1841. Polypodium serpens pt. HB. 349. P. confluens Hk. 1863. HB. 349 (non R. Br. 1810); Niphobolus Bail. 1874 (non Spr. 1827).

samarensis (Pr.) C. Chr. Ind. 200. 1905 — Ins. Philipp.
Gyrosorium Pr. Epim. 140. 1849 (conf. C. varius); Niphobolus Fée 1850-52. Gies. Niph. 191; Polypodium Mett. n. 246. 1587. HB. 356.

Sarasinorum (Gies.) C. Chr. Ind. 200. 1905 — Celebes.
Niphobolus Gies. Niph. 162. 1901.

Schimperianus (Mett.) C. Chr. Ind. 200. 1905 — Africa trop.
Polypodium Mett.; Kuhn, Fil. Afr. 152. 1868 (HB. 513); Niphobolus Gies. 1899. Niph. 111. NPfl. 325.

? Schmidianus Pr. Epim. 132. 1849 = ? C. sticticus.

scolopendrium Desv. Prod. 225. 1827 = C. acrostichoides.

Scytopteris Fourn. Ann. sc. nat. V. **18**. 286. 1873 = C. confluens.

serpens (Forst.) C. Chr. Ind. 201. 1905 — N. Zealand. N. Caledonia. Ins. Norfolk. Polypodium Forst. Prod. 81. 1786 (non Sw. 1788). HB. 349 (pt.). Chr. 99 (pt.); Niphobolus Endl. 1833(?); Carr. 1873. NPfl. 325. Gies. Niph. 219; Craspedaria Pr. 1849; Polypodium stoloniferum Gmel. 1791; P. stellatum Vahl 1794. P. eleagnifolium Bory 1828 — ? Niphobolus bicolor Klf. 1824; Polypodium Mett. n. 251. 1857.

Sheareri (Bak.) C. Chr. Ind. 201. 1905 — China. Polypodium Bak. JoB. **1875**. 201; Niphobolus Diels 1899. NPfl. 325.

sphaerostichus (Mett.) C. Chr. Ind. 201. 1905 — Luzon. Celebes. Polypodium Mett. Pol. 130. n. 265 t. 3 f. 13. 1857. HB. 350. Sphaerostichum acrostichoides Pr. Epim. 135. 1849 (non Cyclophorus supra).

spissus (Bory) Desv. Berl. Mag. **5**. 301. 1811 — Africa austr. cum ins. adjac. China austr. India. Ceylon. — Polypodium Bory; Willd. sp. **5**. 146. 1810; Mett. n. 253; Niphobolus Klf. 1824. Gies. Niph. 204. Candollea heterophylla Mirb. 1803 (non Acrostichum L. 1753); Cyclophorus Desv. 1811; Niphobolus Spr. 1827. Acrostichum dubium Poir. 1810. Polypodium adnascens pt. auctt. HB. 349 — Polypodium vittarioides Wall. 1828. Mett. n. 256. 1857; Niphobolus Pr. 1836. NPfl. 325; Cyclophorus Pr. 1849.

splendens (J. Sm.) C. Chr. Ind. 201. 1905 — Ins. Philipp. Niphobolus J. Sm. JoB. **3**. 396. 1841 (nomen); Gies. Niph. 99; Apalophlebia Pr. Epim. 138. 1849; Polypodium Hk. 1863. HB. ed. I. 353; P. nitens Bak. 1874. HB. ed. II. 353 (non Desv. 1827).

stenophyllus (Bedd.) C. Chr. Ind. 201. 1905 — India bor. Niphobolus Bedd.; Gies. Niph. 131. 1901.

sticticus (Kze.) C. Chr. Ind. 201. 1905 — China austr.-Ceylon. Ins. Philipp. Niphobolus Kze. Linn. **24**. 257. 1851. Gies. Niph. 135; Polypodium Mett. n. 260. 1857. P. fissum pt. auctt.

stigmosus (Sw.) Desv. Berl. Mag. **5**. 301. 1811 — India austr. Ceylon. Malesia. Polypodium Sw. Schrad. Journ. **1800**[2]. 21. 1801. HB. 350 (pt.); Niphobolus Moore 1861. NPfl. 325. Gies. Niph. 113. N. venosus Bl. 1828; Goniopteris J. Sm. 1842; Apalophlebia Pr. 1849.

subfurfuraceus (Hk.) C. Chr. Ind. 201. 1905 — China austr. India bor. Polypodium Hk. sp. **5**. 52. 1863. HB. 351. Chr. 98; Niphobolus Bedd. 1868. NPfl. 325. Gies. Niph. 122.

subvelutinus (Christ) C. Chr. Ind. 201. 1905 — Bhotan. Niphobolus Christ, Ann. Cons. Jard. bot. Genève **7-8**. 331. 1904.

taeniodes C. Chr. Ind. 201. 1905 — China. Polypodium angustissimum Bak. Ann. Bot. **5**. 472. 1891 (non Fée 1869); Niphobolus Gies. 1899. Niph. 183. NPfl. 326.

taiwanensis (Christ) C. Chr. Ind. 201. 1905 — Formosa. Polypodium Christ in Warb. Monsunia **1**. 60. 1901.

tener (Gies.) C. Chr. Ind. 201. 1905 — Réunion. Niphobolus Gies. Niph. 211. 1901.

tonkinensis (Gies.) C. Chr. Ind. 201. 1905 — Tonkin. Niphobolus Gies. Niph. 144. 1901.

varius (Klf.) Gaud. Freyc. Voy. Bot. 364. 1827 — Malesia-Polynesia. Niphobolus Klf. Enum. 125. 1824. (HB. 350). Gies. Niph. 208; Polypodium Mett. n. 254. 1857. (adhuc pertinet f. Bak. Gyrosorium samarense Pr. 1849; conf. C. samarensis).

vittarioides Pr. Epim. 129. 1849 = C. spissus.

Warburgii (Gies.) C. Chr. Ind. 202. 1905 — Celebes.
Niphobolus Gies. Niph. 163. 1901.
xiphioides (Christ) C. Chr. Ind. 202. 1905 — China.
Niphobolus Christ, Bull. Soc. Fr. **52** Mém. I. 25. 1905.
CYCLOPTERIS Gray, Nat. Arr. br. Plants **2**. 9. 1821 = **Cystopteris.**
dentata Gray, Nat. Arr. br. Pl. **2**. in corr. 1821 = C. fragilis.
fragilis Gray, Nat. Arr. br. Pl. **2**, 9. 1821.
regia Gray, Nat. Arr. br. Pl. **2**. in corr. 1821 = C. fragilis*.
CYCLOSORUS Link, Hort. Berol. **2**. 128. 1833 = **Dryopteris.**
gongylodes Link, Hort. Berol. **2**. 128. 1833.
CYRTOGONIUM J. Smith, Journ. Bot. **3**. 402. 1841; **4**. 154. 1842 = **Leptochilus.**
acuminatum Brack. Expl. Exp. **16**. 86. 1854 = L. cuspidatus.
costatum J. Sm. JoB. **4**. 154. 1841 = L. scalpturatus.
crispatulum J. Sm. JoB. **4**. 154. 1841 = L. virens var.
diversifolium J. Sm. JoB. **3**. 403. 1841.
flagelliferum J. Sm. JoB. **4**. 154. 1841 = L. heteroclitus.
laciniatum J. Sm. JoB. **3**. 403. 1841 = L. cuspidatus.
palustre Brack. Expl. Exp. **16**. 86 t. 12. 1854 = L. cuspidatus.
pinnatum J. Sm. Hist. Fil. 142. 1875 = L. lomarioides.
punctulatum J. Sm. JoB. **4**. 154. 1841 = L. auriculatus.
repandum J. Sm. JoB. **3**. 403. 1841 = L. cuspidatus.
rivulare Brack. Expl. Exp. **16**. 85 t. 11 f. 3. 1854.
scandens J. Sm. JoB. **4**. 154. 1841 = L. guianensis.
serratifolium J. Sm. JoB. **4**. 154. 1841.
sinuosum J. Sm. JoB. **3**. 403. 1841 = L. cuspidatus.
subcrenatum J. Sm. JoB. **4**. 154. 1841.
virens J. Sm. JoB. **4**. 154. 1841.
Cyrtomiphlebium Hooker, Spec. Fil. **5**. 15. 1863 *(Polypodium §)* = Polystichum.
CYRTOMIUM Presl, Tent. Pter. 86. 1836 = **Polystichum.**
abbreviatum J. Sm. Ferns br. and for. ed. II. 304. 1877.
caducum Moore, Ind. 276. 1861 = P. Hookerianum.
caryotideum Pr. Tent. 86 t. 2 f. 26. 1836 = P. falcatum*.
falcatum Pr. Tent. 86. 1836.
Fortunei J. Sm. Ferns br. and for. 286. 1866 = P. falcatum var.
fraxinellum Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 264.
juglandifolium Moore, Ind. LXXXIII. 1857 = Phanerophlebia j.
lonchitoides Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 264.
nobile Moore, Ind. LXXXIII. 1857 = Phanerophlebia n.
vittatum Christ, Bull. Soc. Fr. **52** Mém. I. 33. 1905.
CYRTOPHLEBIUM (R. Br.) J. Smith, Journ. Bot. **4**. 58. 1841 = **Polypodium.**
Polypodium § Cyrtophlebium R. Brown in Horsf. Pl. Jav. rar. 4. 1838.
angustifolium J. Sm. Bot. Mag. **72** Comp. 12. 1846.
costatum J. Sm. Lond. JoB. **1**. 196. 1842.
decurrens J. Sm. JoB. **4**. 58. 1841.
nitidum J. Sm. JoB. **4**. 58. 1841.
Phyllitidis J. Sm. JoB. **4**. 58. 1841.
repens J. Sm. JoB. **4**. 58. 1841.
CYSTEA Smith, Engl. Flora **4**. 260, 284. 1828 = **Cystopteris.**
alpina Sm. Engl. Fl. **4**. 291. 1828 = C. fragilis *regia.
angustata Sm. Engl. Fl. **4**. 288. 1828 = C. fragilis.

dentata Sm. Engl. Fl. **4**. 287. 1828 = C. fragilis.
fragilis Sm. Engl. Fl. **4**. 289. 1828.
regia Sm. Engl. Fl. **4**. 289. 1828 = C. fragilis*.
CYSTIDIUM Lindl. Veg. Kingdom 80. 1847 (err.) = Cystodium.
CYSTIDIUM J. Smith msc. (Moore, Ind. 278. 1861) = **Polypodium** (obliquatum).
CYSTODIUM J. Smith in Hook. Gen. Fil. t. 96. 1841 = **Saccoloma.**
sorbifolium J. Sm. in Hk. Gen. t. 96. 1841.
CYSTOPTERIS Bernhardi, Schrad. Neu. Journal **1**². 5, 26. 1806 — (Polypodiaceae Gen. **5**).
abyssinica Fée, Gen. 300. 1850-52 = C. fragilis.
acuta Fée, Gen. 300. 1850-52 = C. fragilis *regia.
affinis Fée, Gen. 299. 1850-52 = Davallia hymenophylloides.
albescens Link, Fil. sp. 47. 1841 = Woodsia obtusa.
Allioni Newm. Phytol. **1851**. app. XXV = C. montana.
alpina Desv. Prod. 264. 1827 = C. fragilis *regia.
aspidioides Pr. Tent. 93. 1836 = Microlepia speluncae.
atomaria Pr. Tent. 93. 1836 = C. bulbifera.
atrovirescens Pr. Tent. 93. 1836; Epim. 66. 1849 (atrovirens) = C. fragilis *regia.
azorica Fée, Gen. 300. 1850-52 = C. fragilis var. diaphana.
Baenitzii Dörfler in Baenitz, Herb. Eur. n. 6510, Prospect. 4. 1891. = C. fragilis.
Blindi Parmentier, Bull. Ac. Géogr. Bot. Mans **9**. 40 c. tab. 1900 — Jura.
(? Cystopteris fragilis × Asplenium trichomanes).
?brasiliana Pr. Tent. 93. 1836 = Hypolepis.
brevinervis Fée, Gen. 300. 1850-52; 7 mém. 65 t. 26 f. 2 = Dryopteris hirta.
bulbifera (L.) Bernh. Schrad. neu. Journ. **1**². 10. 1806. HB. 103. Chr. 281. NPfl. 164.
Polypodium L. sp. **2**. 1091. 1753; Aspidium Sw. 1801; (? Cyathea Bernh. 1802); Nephrodium Michx. 1803; Filix Und. 1900. Aspidium atomarium Mühl.; Willd. 1810; Athyrium Pr. 1825; Cystopteris Pr. 1836.
canariensis Pr. Tent. 93. 1836 = C. fragilis var. diaphana.
Caucasica Fée, Gen. 299. 1850-52 = C. fragilis.
Chilensis Fée, Gen. 300. 1850-52 = C. fragilis.
comosa Pr. Tent. 93. 1836 = Alsophila squamulata.
crenata Fries; Novit. Mantissa **3**. 195. 1842 = Athyrium.
Dalhousiana Fée, 8 mém. 108. 1857 = Davallia hymenophylloides.
dentata Desv. Prod. 263. 1827 = C. fragilis.
Dickieana Sim, Gard. Chr **1848**. 308 = C. fragilis *regia.
dimidiata Dcne. Jacq. Voy. 4. 177 t. 178. 1844 = Davallia bullata vel immersa.
Douglasii Hk. sp. **1**. 200. 1846 — Ins. Hawaii. — C. Sandwicensis Brack. 1854.
?elata Desv. Prod. 264. 1827 = { Dryopteris Boryana (f. Mett.). / Dryopteris punctata (f. Kuhn).
emarginata Hk. sp. **1**. 201. 1846 } C. fragilis var. diaphana.
emarginulata Pr. Tent. 93. 1836: Epim. 65. 1849 } C. fragilis var. diaphana.
Filix femina Coss. et Germ. Fl. Paris 676. 1845 = Athyrium.
Filix fragilis Chiovenda, Ann. di Bot. **1**. 210. 1903 = C. fragilis.
fragilis (L.) Bernh. Schrad. neu. Journ. **1**². 26 t. 2 f. 9. 1806. HB. 103. Chr. 280. NPfl. 163 — Reg. arct., antarct., temp. bor. et austr. Montes trop.
Polypodium L. sp. **2**. 1091. 1753; Cyathea Sm. 1793; Aspidium Sw. 1801; Athyrium Spr. 1804; Cyclopteris Gray 1821; Cystea Sm. 1828; Filix Und. 1900. — Polypodium diaphanum Bory 1804. Aspidium viridulum Desv. 1811; Cystopteris Desv. 1827. C. canariensis Pr. 1836 — Europa austr. Ins. Atlant. Ins. Mascar. Amer. trop.

— *Polypodium regium L. 1753; Cyathea Forst. 1798; Aspidium Sw. 1801; Athyrium Spr. 1804; Cyclopteris Gray 1821; Cystopteris Desv. 1827. NPfl. 164; Cystea Sm. 1828; Polypodium alpinum Jacq. 1788; Cyathea Roth 1800; Aspidium Sw. 1801; Athyrium Spr. 1804; Cystopteris Desv. 1827. HB. 103. Chr. 281; Cystea Sm. 1828; Nephrodium Opiz (?1823) — Europa.
— Cystopteris Tasmanica Hk. 1846 — Tasmania.
— Woodsia laetevirens Prentice 1891 — N. S. Wales.

fragilis × **montana** Christ, Farnkr. d. Schweiz 162 f. 28. 1900 — Mts. Jura.
fragilis × Asplenium trichomanes vide C. Blindi.
fumarioides Schott, Gen. ad t. 8. 1834 = C. fragilis *regia.
gigantea Pr. Tent. 93. 1836 = ? Acrophorus stipellatus.
hirta Kl. Linn. **20**. 361. 1847 = Dryopteris.
jamaicensis Desv. Prod. 263. 1827 = C. fragilis.
japonica Luerss. Engl. Jahrb. **4**. 363. 1883 — Japonia.
javensis Desv. Prod. 265. 1827 = Dryopteris punctata.
laciniata Col. Tr. N. Zeal. Inst. **31**. 265. 1899 — N. Zealand.
leptophylla Pr. Tent. 93. 1836 = Hypolepis incisa.
leucosoria Schur, Oest. Bot. Zeit. **8**. 328. 1858 = C. sudetica.
mexicana? Fée, Gen. 300. 1850-52 = C. fragilis.
montana (Lam.) Bernh.; Desv. Prod. 264. 1827. HB. 104. Chr. 281. NPfl. 164 — Europa. Kamtschatka. Alaska-Colorado-Labrador.
Polypodium Lam. Fl. franç. **1** 23. 1778; Cyathea Sm. 1793; Aspidium Sw. 1801; Athyrium Spr. 1804; Filix Und. 1900.
moupinensis Franch. Nouv. Arch. Mus. II. **10**. 111. 1887 = C. sudetica.
myrrhidifolia Newm. Hist. br. ferns ed. III. 97. 1854 = C. montana.
nodosa Mett. Ann. Lugd. Bat. **1**. 241. 1864 = Acrophorus stipellatus.
novae-zealandicae Armstr. Tr. N. Zeal. Inst. **13**. 360. 1881 — N. Zealand.
obovata Pr. Tent. 93. 1836 = Asplenium lanceolatum var.
obtusa Pr. Tent. 93. 1836 = Woodsia.
odorata Desv. Prod. 264. 1827 = Dryopteris crenata.
orientalis Desv. Prod. 264. 1827 = C. fragilis.
Perriniana Link, Hort. Berol. **2**. 131. 1833 = Woodsia obtusa.
Pontederae Link, Fil. sp. 45. 1841 = C. fragilis.
regia Desv. Prod. 264. 1827 = C. fragilis*.
retusa Dcne. Jacq. Voy. **4**. 176 t. 177. 1844 = Athyrium filix femina var.
rhaetica Link, Hort. Berol. **2**. 131. 1833 = C. fragilis.
rufescens Fée, Gen. 300. 1850-52; 6 mém. 22 t. 6 f. 5 = Dryopteris hirta.
Sandwicensis Brack. Expl. Exp. **16**. 234. 1854 = C. Douglasii.
scandicina Desv. Prod. 264. 1827 = Athyrium.
setosa Bedd. Ferns br. Ind. t. 312. 1869. Chr. 281 — India bor. Birma. China.
Lastrea Bedd. 1868; Davallia Bak. 1874. HB. 468.
spinulosa Maxim. Mém. Ac. St. Pétersb. **9**. 340. 1859 = Athyrium.
squamata Dcne. Jacq. Voy. **4**. 178. 1844 = Davallia pulchra.
sudetica A. Br. et Milde, 33 Jahresber. schles. Ges. **1855**. 92. HB. 103. Chr. 281. NPfl. 164 — Europa orient. Sibiria orient. Thibet. Yunnan.
C. leucosoria Schur 1858 — C. moupinensis Franch. 1887 — Davallia triangularis Bak. 1891.
Tasmanica Hk. sp. **1**. 199. 1846 = C. fragilis var.
taygetensis Heldr. & Sart. in Heldr. Herb. norm. n. 35; Salom Nom. 147. 1883 = C. fragilis *regia.

tenuis Desv. Prod. 263. 1827 = C. fragilis.

?**tenuisecta** (Bl.) Mett. Ann. Lugd. Bat. **1**. 241. 1864 — Java.
Aspidium Bl. Enum. 170. 1828; Alsophila Bl.; Moore 1857. HB. 44. Athyrium Moore 1860; Asplenium Hk. 1860.

translucens Desv. Prod. 264. 1827 = C. fragilis.

tristis Mett. Ann. Lugd. Bat. **1**. 241. 1864 = Alsophila.

Ulei Christ in Schwacke, Pl. Nov. Mineiras **2**. 30. 1900; Bull. Boiss. II. **2**. 637. 1902 — Brasilia austr.

vestita Pr. Tent. 93. 1836 = Woodsia obtusa.

villosa Desv. Prod. 263. 1827 = Dryopteris cruciata.

villosa Fée, Gen. 299. 1850-52 (nomen) —?.

viridula Desv. Prod. 264. 1827 = C. fragilis var. diaphana.

DANAEA Smith, Mém. Acad. Turin **5**. 420. 1793 — (Marattiaceae Gen. **5**).

alata Sm. Mém. Ac. Turin **5**. 420. 1793; Tracts 261. 1798; Sw. Syn. 167. 1806. HB. 443. Chr. 360. NPfl. 444 — India occ.

angustifolia Pr. Suppl. 35. 1845 = D. nodosa.

Augustii Karst.; Kze. Linn. **20**. 2. 1847. HB. 443. NPfl. 444 — Columbia. Venezuela.

cordata Fée, Cr. vasc. Br. **1**. 216 t. 71 f. 5. 1869 — Brasilia.

crispa Endres; Reichb.fil. Bot. Zeit. **1872**. 489. HB. 525. Chr. 361. NPfl. 444 — Costa Rica.

cuspidata Liebm. Vid. Selsk. Skr. V. **1**. 307 (seors. 155). 1849 — Mexico.

dubia Pr. Suppl. 36. 1845 — Brasilia.

elata Liebm. Vid. Selsk. Skr. V. **1**. 306 (seors. 154). 1849 — Mexico.

elliptica Sm. in Rees, Cycl. **11**. —. 1809; Hk. et Grev. Ic. Fil. t. 52. 1828. HB. 444. Chr. 360. NPfl. 444 — Amer. trop. — D.geniculata Raddi 1819.

evecta Spr. Schrad. Journ. **1799**². 272 = Angiopteris.

Fendleri Und. Bull. Torr. Cl. **29**. 673. 1902 — India occ.

geniculata Raddi, Opusc. Bol. **3**. 281. 1819; Pl. Bras. **1**. 75 t. 5 f. 1. 1825 = D. elliptica.

humilis Moore, Ind. 286. 1861. HB. 443. NPfl. 444 — Ecuador.

intermedia J. Sm. Lond. JoB. **2**. 393. 1843 (nomen) —?.

jamaicensis Und. Bull. Torr. Cl. **29**. 675. 1902 — Jamaica.

Jenmani Und. Bull. Torr. Cl. **29**. 677. 1902 — Jamaica.

Leprieurii Kze. Farnkr. **1**. 137 t. 60. 1843. HB. 443. NPfl. 444 — Guiana. [Amazonas.

longifolia Desv. Berl. Mag. **5**. 307. 1811 = D. nodosa.

Mazeana Und. Bull. Torr. Cl. **29**. 676. 1902 — Guadeloupe.

media Liebm. Vid. Selsk. Skr. V. **1**. 306 (seors. 154). 1849 = D. elliptica.

Moritziana Pr. Suppl. 35. 1845. HB. 443. NPfl. 444 — Costa Rica-Peru. Jamaica.

nigrescens Jenm. Gard. Chr. III. **24**. 413. 1898 — Guiana.

nodosa (L.) Sm. Mém. Ac. Turin **5**. 420 t. 9 f. 11. 1793. HB. 444. Chr. 360. NPfl. 444 — Amer. trop. — Acrostichum L. sp. **2**. 1070. 1793; Asplenium L. 1763 — Danaea longifolia Desv. 1811.

oligosora Fourn.; Bak. Ann. Bot. **5**. 499. 1891; Christ, Hedwigia **44**. 369. 1905 — Guadeloupe.

paleacea Raddi, Pl. Bras. **1**. 76 t. 5 f. 2. 1825 — Brasilia. — Danaeopsis Pr. 1845.

polymorpha Leprieur; Bak. Ann. Bot. **5**. 499. 1891 — India occ.

Sellowiana Pr. in Corda, Beitr. Fl. d. Vorwelt t. 51 f. 18-23. 1845: Pr. Suppl. [37. 1845 — Brasilia.

serrulata Bak. JoB. **1881**. 208. NPfl. 444 — Columbia.

simplicifolia Rudge, Pl. Guian. **1**. 24 t. 36. 1805. HB. 442. Chr. 360. NPfl. 443 — Guiana. Brasilia bor.

stenophylla Kze. Farnkr. **1**. 55 t. 28. 1840. HB. 443. NPfl. 444 — Guadeloupe. Brasilia. — Heterodanaea Pr. 1845.
trichomanoides Spruce; Moore, Ind. 288. 1861. HB. 442 — Peru.
trifoliata Reichb.; Kze. Anal. 4 t. 2. 1837. HB. 442. Chr. 360. NPfl. 443 — Guiana.
Ulei Christ, Hedwigia **44**. 368. 1905 — Amazonas.
Wendlandi Reichb.fil. Bot. Zeit. **1872**. 490. HB. 526. NPfl. 444 — Costa Rica.
Wrightii Und. Bull. Torr. Cl. **29**. 676. 1902 — Cuba. Puerto Rico.

DANAEOPSIS Presl, Suppl. Tent. Pterid. 39. 1845 = **Danaea.**
paleacea Pr. Suppl. 39. 1845.

DAREA Jussieu, Genera plant. 15. 1789; Smith, Mém. Acad. Turin **5**. 409. 1793 [= **Asplenium.**
appendiculata Willd. sp. **5**. 296. 1810 = A. bulbiferum var.
appendiculata Bl. Enum. 206. 1828 = A. tenerum *Belangeri.
aspidioides Willd. Enum. 1072. 1809; sp. **5**. 301. 1810 = Athyrium a.
asplenioides Bory, Bél. Voy. Bot. **2**. 53. 1833 = A. Boryanum.
auriculata Willd. sp. **5**. 296. 1810.
Belangeri Bory, Bél. Voy. Bot. **2**. 51. 1833 = A. tenerum*.
bifida Bory, Bél. Voy. Bot. **2**. 54. 1833 = A. lineatum var.
cicutaria Willd.; sp. **5**. 300. 1810 = A. cristatum.
cuneata Desv. Berl. Mag. **5**. 323. 1811 = A. affine.
disticha Klf. Enum. 180. 1824 = ? A. achilleifolium.
flaccida Willd. sp. **5**. 295. 1810.
foecunda Fée, Gen. 333. 1850-52 = A. compressum.
fumarioides H.B. Willd. sp. **5**. 299. 1810 = Odontosoria meifolia.
? furcans Bory, Dup. Voy. Bot. **1**. 269 t. 25 f. 2. 1828 = Elaphoglossum furcatum.
furcata Willd. sp. **5**. 297. 1810 = A. achilleifolium.
furcata Bl. Enum. 207. 1828 = A. tenerum *Belangeri.
inaequalis Willd. sp. **5**. 298. 1810 } = A. lineatum var.
intermedia Klf.; Bory, Bél. Voy. **2**. 55. 1833 } = A. lineatum var.
japonica Willd. sp. **5**. 302. 1810 = Onychium j.
membranacea Poir. Enc. Suppl. **2**. 451. 1811 = A. myriophyllum.
? Millefolium Fée, Gen. 333. 1850-52.
myriophylla Willd. sp. **5**. 301. 1810.
obtusa Desv. Berl. Mag. **5**. 323. 1811 = A. achilleifolium.
odontites Willd. sp. **5**. 296. 1810 = A. flaccidum.
palmata Klf. Enum. 181. 1824 = A. subdigitatum.
polypodioides Willd.; Klf. Enum. 181. 1824 — ?
prolifera Willd. sp. **5**. 299. 1810 = A. lineatum var.
Pteridis Bory, Bél. Voy. Bot. **2**. 52. 1833.
rhizophylla Willd. sp. **5**. 300. 1810 = A. myriophyllum.
rutaefolia Willd. sp. **5**. 298. 1810 = A. achilleifolium.
scandens Fée, Gen. 332. 1850-52.
stans Bory, Bél. Voy. Bot. **2**. 53. 1833 = A. achilleifolium.
tenera Spr. Schrad. Journ. **1799**^{2}. 269.
? thalictroides Fée, Gen. 333. 1850-52 = ? A. cristatum.
thecifera Fourn. Mex. pl. **1**. 132. 1872.
trilobata Desv. Berl. Mag. **5**. 323. 1811 = A. cristatum.
tripinnata Cav. Descr. 259. 1802 = A. myriophyllum.
violascens Bory, Bél. Voy. Bot. **2**. 55. 1833 = A. lineatum var.
viridans Bory, Bél. Voy. Bot. **2**. 54. 1833.
vivipara Willd. sp. **5**. 302. 1810.

Dareastrum Fée, Gen. Fil. 190. 1850-52 *(Asplenium §)* = Asplenium.

DAVALLIA Smith, Mém. Acad. Turin **5**. 414. 1793 — (Polypodiaceae Gen. **32**).

Subgenera: D = **Eudavallia.**
L = **Leucostegia.**
P = **Prosaptia.**
S = **Scyphularia.**

achilleaefolia Wall. List n. 248. 1828; Hk. sp. **1**. 195 t. 56 D. 1846 = Stenochlaena aculeata.
aculeata Sm. Mém. Ac. Turin **5**. 415. 1793 = Odontosoria.
aculeata Hedw. Fil. Gen. et Sp. t. 23. 1803 = Odontosoria fumarioides.
aculeata Spr. Anleit. **3**. 150 t. 5 f. 37. 1804 = D. flexuosa.
adiantifolia Hk. sp. **1**. 176. 1846 = Lindsaya adiantoides.
adiantifolia Kze. Bot. Zeit. **1848**. 216 = Lindsaya pectinata.
adiantoides Sw. Syn. 131. 1806 = Saccoloma domingense.
aemula Mett.; Kuhn, Linn. **36**. 144. 1869 = Humata.
affinis Hk. sp. **1**. 158 t. 52 B. 1846 = D. hymenophylloides.
(P) **alata** Bl. Enum. 230. 1828. NPfl. 214 — Asia trop.-Samoa.
Pachypleuria Pr. 1849. Davallia serraeformis Wall. 1828; Polypodium J. Sm. 1841; Prosaptia Christ 1898. Davallia Emersoni Hk. et Grev. 1829. HB. 94. Chr. 306; Prosaptia Pr. 1836.
alata Hew. Mag. Nat. Hist. II. **2**. 465. 1838 = Saccoloma inaequale.
alata J. Sm. JoB. **3**. 417. 1841 = D. decurrens.
alpina Bl. Enum. 231. 1828 = Humata repens var.
Amboynensis Hk. sp. **1**. 178 t. 56 C. 1846 = Tapeinidium.
Andersonii Bak. Ann. Bot. **5**. 200. 1891 = Leptolepia.
Anderssonii Mett.; Kuhn, Linn. **36**. 143. 1869 = Humata.
angustata Wall. List n. 242. 1828 } = Humata heterophylla.
angustifolia Roxb. Calc. Journ. **4**. 513. 1844 } = Humata heterophylla.
arborescens Willd. sp. **5**. 470. 1810 = Dennstaedtia obtusifolia.
(D) **arctotheca** Fourn. Ann. sc. nat. V. **18**. 339. 1873 — N. Caledonia.
asperrima Ces. Rend. Ac. Napoli **16**. 25, 28. 1877 = Saccoloma moluccanum.
aspidioides Bak. Ann. Bot. **5**. 200. 1891 = Leptolepia.
asplenioides Moore, Ind. 290. 1861 = Humata.
(L) **assamica** (Bedd.) Bak. Syn. ed. I. 452. 1868; ed. II. 467. 1874 — India bor.
Acrophorus Bedd. Ferns br. Ind. t. 94. 1866; Leucostegia J. Sm. 1875; Davallia micans Mett.; Bak. 1867; HB. ed. I. 97; Humata Diels 1889. NPfl. 209.
(L) **athamantica** Christ, Bull. Soc. Fr. **52** Mém. I. 65. 1905 — Yunnan.
athyriifolia Bak. Ann. Bot. **5**. 200. 1891 = Athyrium fimbriatum.
attenuata Lodd. Cat. 1849 = D. denticulata.
Beccariana Ces. Atti Ac. Napoli **7**[8]. 14 t. 3 f. 6. 1876 = D. pallida.
(L) **Beddomei** Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 527 t. 1. 1899 — India bor.
Belangeri Bory, Bél. Voy. Bot. **2**. 72 t. 7 f. 1. 1833 = Humata vestita.
Berteriana Colla, Mem. Ac. Torino **39**. 37 t. 65. 1836 = Dicksonia.
bidentata Schkuhr, Kr. Gew. **1**. 119 t. 127. 1809 = D. denticulata.
bifida Klf. Enum. 222. 1824 = Odontosoria.
biflora Klf. Enum. 221. 1824 = Odontosoria.
bipinnata Hk. sp. **1**. 161. 1846 = Odontosoria uncinella.
bipinnata Bak. Ann. Bot. **5**. 200. 1811 = Leptolepia tripinnata

bipinnatifida Bl. Enum. 232. 1828 = Humata vestita.
bipinnatifida Bak. Kew Bull. **1899**. 119 = Humata neoguineensis.
biserrata Bl. Enum. 232. 1828 = Tapeinidium pinnatum.
Blumeana Hk. sp. **1**. 117 (pt.) t. 54 A. 1846 = Lindsaya.
Blumei Mett.; Kuhn, Ann. Lugd. Bat. **4**. 289. 1869 = D. contigua var.
borneensis J. Sm.; Kuhn, Ann. Lugd. Bat. **4**. 286. 1869 = Dryopteris.
Boryana Pr. Rel. Haenk. **1**. 66. 1825 = Lindsaya repens var.
botrychioides Bak. Syn. 90. 1867 = Humata.
brachycarpa Mett. msc. (Bak. Syn. 470. 1874) = Asplenium.
brachypoda Bak. Syn. 468. 1874 = Lindsaya cultrata var.
Brasiliensis Hk. sp. **1**. 185. 1846 = Saccoloma inaequale.
(D) **bullata** Wall. List n. 258. 1828 (nomen); Hk. sp. **1**. 169 t. 50 B. 1846. HB. 97. Chr. 301. NPfl. 214 — Japonia. China. Asia trop. — D. decora Moore 1859.
calobodon Bak. Syn. 470. 1874 = Microlepia madagascariensis.
calvescens Hk. sp. **1**. 172 t. 48 B. 1846 = Microlepia marginata var.
campyloptera Kze. Linn. **10**. 544. 1836 = Asplenium theciferum*.
campylura Kze. Bot. Zeit. **1850**. 132 = Saccoloma moluccanum var.
(D) **canariensis** (L.) Sm. Mém. Ac. Turin **5**. 414 t. 9 f. 6. 1793. HB. 97. Chr. 301. NPfl. 214 — Ins. Atlant. Marocco. Portugal. Hispania.
Trichomanes L. sp. **2**. 1099. 1753; Caenopteris Willd. 1794. Polypodium lusitanicum L. 1753.
capillacea Willd. sp. **5**. 479. 1810 = Trichomanes.
caudata Cav. Descr. 279. 1802 = D. solida var.
caudata Wall. List n. 2220. 1829 = D. denticulata.
celebica Christ, Verh. Nat. Ges. Basel **11**. 426. 1896 = Dennstaedtia scabra.
chaerophylla Wall. List n. 259. 1828; Hk. sp. **1**. 157 t. 61 A. 1846 = D. pulchra.
(D) **chaerophylloides** (Poir.) Steud. Nom. **2**. 146. 1824. Chr. 301. NPfl. 214 — Africa trop. cum ins. — Trichomanes Poir. Enc. **8**. 80. 1808; Humata Desv. 1827. Davallia nitidula Kze. 1836. HB. 97.
chinensis Sm. Mém. Ac. Turin **5**. 414. 1793 = Odontosoria.
chusana Willd. sp. **5**. 475. 1810 = Odontosoria chinensis.
(L) **cicutarioides** Bak. JoB. **1890**. 106 — N. Guinea.
Humata Diels 1899. NPfl. 210.
(?) ?*ciliata* Pr. Del. Prag. **1**. 188. 1822 — Brasilia.
ciliata Hk. sp. **1**. 184 t. 60 A. 1846 = Microlepia hirsuta.
(L) **Clarkei** Bak. Syn. ed. II. 91. 1874. Chr. 302 — Himalaya. Yunnan.
Acrophorus Hookeri Moore 1857 (nomen); Leucostegia Bedd. 1883; Humata Diels 1899. NPfl. 209 (non Davallia Hookeriana Wall. 1828; Hk. 1846) — D. Delavayi Bedd. 1888.
clavata Sm. Mém. Ac. Turin **5**. 415. 1793 = Odontosoria.
concinna Schrad. Gött. gel. Anz. **1818**. 918 = Asplenium theciferum*.
concinna Pr. Rel. Haenk. **1**. 66. 1825 = Dennstaedtia Mathewsii.
coniifolia Wall. List n. 252. 1828 = D. denticulata.
(P) **contigua** (Forst.) Spr. Schrad. Journ. **1799**². 271. HB. 94. Chr. 306. NPfl. 214 — Malesia. Polynesia. — Trichomanes Forst. Prod. 84. 1786; Humata Desv. 1827; Prosaptia Pr. 1836; Polypodium J. Sm. 1841 — Davallia pectinata Bl. 1828 (non Sm. 1793); Prosaptia Moore 1861; Davallia Blumei Mett.; Kuhn 1869.
corcovadensis Lodd.; Kze. Linn. **23**. 258. 1850 — Brasilia.
cordifolia Roxb. Calc. Journ. **4**. 514. 1844 = Humata repens.

(D) **corniculata** Moore, Ind. 292. 1861 — Malesia.
D. epiphylla Bl. Enum. 235. 1828. HB. 96 (non Spr. 1799).
Cumingii Hk. sp. **1**. 155 t. 45 B. 1846 = Humata lepida.
cuneata Spr. Schrad. Journ. **1799**². 271 = Lindsaya.
?cuneifolia Hk. sp. **1**. 176. 1846 = Lindsaya adiantoides.
cuneiformis Sw. Schrad. Journ. **1800**². 87. 1801 = Odontosoria chinensis.
dareaeformis Levinge; Clarke, Tr. Linn. Soc. II. Bot. **1**. 443. 1880 = D. Clarkei et Polypodium dareiforme.
decipiens Ces. Rend. Ac. Napoli **16**. 25, 29. 1877 = Odontosoria retusa.
decomposita Bak. JoB. **1884**. 141 = Odontosoria.
decora Moore in Sim, Cat. Ferns 39. 1859 = D. bullata var.
decora ht. Bull.; Gard. Chr. III. **13**. 415. 1893 — Hort. — (an praec.?).
(D) **decurrens** Hk. sp. **1**. 167 t. 44 B. 1846. HB. 95. Chr. 300. NPfl. 214 — Celebes. Ins. Philipp.
Microlepia Fée 1850-52; Davallia alata J. Sm. 1841 (non Bl. 1828).
Delavayi Bedd.; (Clarke et Bak. Journ. Linn. Soc. **24**. 410. 1888) = D. Clarkei.
delicatula Christ, Verh. Nat. Ges. Basel **11**. 224 t. 3 f. 1-3. 1895 = Lindsaya.
Denhami Hk. 2 Cent. t. 47. 1860 = Tapeinidium.
(D) **denticulata** (Burm.) Mett.; Kuhn, Fil. Deck. 27. 1867 — Asia, Austr., Polynesia trop. Madagascar. Fernando Po.
Adiantum Burm. Fl. Ind. 236. 1768; Trichomanes Houtt. 1783; Davallia elegans Sw. 1801. HB. 95. Chr. 300. NPfl. 214; Trichomanes Poir. 1808; Humata Desv. 1827; Parestia Pr. 1849 — Davallia patens Sw. 1806 — Trichomanes elatum Forst. 1786; Davallia Spr. 1799; Wibelia Bernh. 1801; Humata Desv. 1827; Parestia Pr. 1849 — Trichomanes epiphyllum Forst. 1786; Davallia Spr. 1799; Humata Desv. 1827; Wibelia multifida Spr. 1801 — Davallia Vogelii Hk. 1846 (Fernando Po).
deparioides Ces. Atti Ac. Napoli **7**⁸. 13 t. 4 f. 8. 1876 = Lecanopteris.
didyma Hedw. Fil. Gen. et Sp. t. 22. 1803 = Odontosoria chinensis.
(D) **dissecta** J. Sm.; Moore et Houlst. Gard. Mag. Bot. **3**. 325. 1851; Moore, Gard. Chr. **1855**. 469 c. fig. HB. 469 — Java.
distans Klf. Enum. 222. 1824 = ? Saccoloma inaequale.
(D) **divaricata** Bl. Enum. 237. 1828. HB. 96. Chr. 300. NPfl. 214 — Asia trop.
D. polyantha Hk. 1846; Microlepia Fée 1850-52.
divaricata Schlecht. et Cham. Linn. **5**. 617. 1830 = Odontosoria Schlechtendalii.
divergens Kze. Linn. **19**. 404; **20**. 5. 1847 = Microlepia trichosticha.
domingensis Spr. Anleit. **3**. 149 t. 4 f. 33. 1804 = Saccoloma.
(L) ?**dubia** R. Br. Prod. Fl. N. Holl. 157. 1810. HB. 468 — Australia. ? Fiji.
Woodsia? Desv. 1827; Dicksonia Gaud. 1827; Sitolobium Brack. 1854; Balantium Brownianum Pr. 1836.
dumosa Sw. Syn. 135, 353. 1806 = Odontosoria aculeata.
elata Spr. Schrad. Journ. **1799**². 271 = D. denticulata var.
elegans Sw. Schrad. Journ. **1800**². 87. 1801 = D. denticulata.
elegans Hedw. Fil. Gen. et Sp. 1803 = D. solida.
elegans β pulchra Hk. sp. **1**. 165. 1846 = D. corniculata.
Emersoni Hk. et Grev. Ic. Fil. t. 105. 1829 = D. alata.
epiphylla Spr. Schrad. Journ. **1799**². 271 = D. denticulata var.
epiphylla Bl. Enum. 235. 1828 = D. corniculata.
falcata Sm. Mém. Ac. Turin **5**. 415. 1793 = Nephrolepis exaltata.
falcata Sw. Schrad. Journ. **1800**². 87. 1801 = ? Nephrolepis cordifolia.

(D?) **falcinella** Pr. Rel. Haenk. **1**. 66 t. 11 f. 2. 1825. HB. 91 — Ins. Philipp.
Leucostegia J. Sm. 1841; Acrophorus Moore 1857.
Fejeensis Hk. sp. **1**. 166 t. 55 D. 1846 = D. solida var.
ferruginea Cav. Descr. 277. 1802 = Odontosoria chinensis.
ferruginea Desv. Prod. 315. 1827 = Odontosoria Melleri.
(L) **ferulacea** Moore, Ind. 294. 1861. HB. 92. Chr. 303 — Fiji.
Schizoloma Kuhn 1882; Humata Diels 1899. NPfl. 210; Davallia trichomanoides Hk. 2 Cent. t. 64. 1861 (non Bl. 1828).
fijiensis Diels, Nat. Pfl. 1^4. 214. 1899 = D. solida var.
firmula Bak. Ann. Bot. **8**. 123. 1894 = Microlepia.
flabellifolia Bak. Journ. Linn. Soc. **15**. 414. 1876 = Odontosoria.
flaccida R. Br. Prod. Fl. N. Holl. 157. 1810 (Don. Prod. 10. 1825) = Microlepia speluncae.
flagellifera Wall. List n. 243. 1828 = Tapeinidium pinnatum.
flexuosa Spr.; Sieb. Fl. Mart. Suppl. n. 23; Pr. Tent. 129. 1836; Kze. Bot. Zeit. **1850**. 213 — India occ. — Microlepia Ettingsh. 1865; Davallia aculeata Spr. Anleit. **3**. 150 t. 5 f. 37. 1804 (non Sm. 1793) — (Odontosoria sp.).
foeniculacea Hk. 2 Cent. t. 54. 1861 = Asplenium stenolobum.
Forsteri Carr. in Seem. Fl. Vit. 339. 1873 = Odontosoria angustifolia.
(P) **Friderici et Pauli** Christ, Verh. Nat. Ges. Basel **11**. 202 t. 2 f. 1-4. 1895; Ann. Jard. Buit. **15**. 94 t. 14 f. 10. 1897. NPfl. 214 — Celebes.
Polypodium Christ 1904 (non 1896).
(D) **fructuosa** Christ in Warb. Monsunia **1**. 86. 1900 — Java.
fumarioides Sw. Schrad. Journ. **1800**^{2}. 89. 1801 = Odontosoria.
?Gaimardiana Pr. Tent. 128. 1836 = Humata.
gibberosa Sw. Schrad. Journ. **1800**^{2}. 88. 1801 = Asplenium.
glauca Cav. Descr. 278. 1802 — Peru.
glaucescens Hedw. Fil. Gen. et Sp. 1803 = Asplenium gibberosum.
Goudotiana Kze. Anal. 35 t. 22 f. 2. 1837 = Lindsaya.
gracilis Bl. Enum. 233. 1828 = Tapeinidium pinnatum var.
(D) **Graeffei** Luerss. Fil. Graeff. 211 t. 18. 1871 — Samoa. Aneiteum.
(D) **Griffithiana** Hk. sp. **1**. 168 t. 49 B. 1846. HB. 96. Chr. 301. NPfl. 214 — China. India bor. Formosa. — Leucostegia J. Sm. 1875.
Griffithii Christ, Bull. Boiss. II. **4**. 616. 1904 — D. Griffithiana.
hemiptera Bory, Bél. Voy. Bot. **2**. 73 t. 7 f. 2. 1833 = Lindsaya repens.
heterophylla Sm. Mém. Ac. Turin **5**. 415. 1793 = Humata.
hirsuta Sw. Schrad. Journ. **1800**^{2}. 87. 1801 = Microlepia pilosella.
hirta Klf. Enum. 223. 1824 = Microlepia.
Hookeriana Wall. List n. 2684. 1829 = Microlepia.
Hornei Bak. Fl. Maur. 470. 1877 = Lindsaya.
(L) **Hosei** Bak. JoB. **1888**. 323 — Borneo.
humilis Hk. sp. **1**. 185. 1846 = Cystopteris fragilis.
(I) **hymenophylla** (Parish) Bak. Syn. ed. I. 452. 1868; ed. II. 90. 1874 — Moulmein.
Acrophorus Parish; Bedd. Ferns br. Ind. t. 96. 1865; Leucostegia Bedd. 1876; Davallia Parishii Hk. 1867. HB. ed. I. 90.
(L) **hymenophylloides** (Bl.) Kuhn, Ann. Lugd. Bat. **4**. 286. 1869. Chr. 302 — Asia, Polynesia trop. — Aspidium Bl. Enum. 172. 1828; Leucostegia Bedd. 1876. L. affinis J. Sm. 1841 (nomen); Davallia Hk. 1846. HB. 92; Microlepia Pr. 1849; Cystopteris Fée 1850-52; Acrophorus Moore 1854; Humata Mett. 1856.

hymenophylloides Bak. Syn. 93. 1867 = Lindsaya.
illustris ht; Gard. Chr. III. **26**. 17. 1899 — Hort.
(L) **immersa** Wall. List n. 256. 1828 (nomen): Hk. sp. **1**. 156. 1846. HB. 91. Chr. 302 — India bor.-Java. Celebes. — Leucostegia Pr. 1836; Acrophorus Moore 1854; Humata Mett. 1856. NPfl. 209. ? Cystopteris dimidiata Dcne. 1844.
Imrayana Hk. sp. **1**. 171 t. 49 A. 1846 = Saccoloma.
inaequalis Kze. Linn. **9**. 87. 1834 = Saccoloma.
inaequalis γ minor Hk. sp. **1**. 180 t. 58 A. 1846 = Saccoloma moluccanum.
intermarginalis Bl. Enum. 230. 1828 = Humata Gaimardiana.
intermedia ht.; Gard. Chr. III. **25**. 31. 1899 (hybr.) — Hort.
intramarginalis Ces. Rend. Ac. Napoli **16**. 25, 29. 1877 = Tapeinidium pinnatum.
Jamaicensis Hk. sp. **1**. 183. 1846 = Microlepia speluncae.
Japonica Kze. Bot. Zeit. **1848**. 542 = Microlepia strigosa.
Khasiyana Hk. sp. **1**. 173 t. 47 A, 57 A. 1856 = Microlepia strigosa.
(L) **Kingii** Bak. in Hk. Ic. pl. t. 1622. 1886 — Java.
Humata Diels 1899. NPfl. 210.
(D) **Korthalsii** Kuhn, Linn. **36**. 142. 1869 — Sumatra.
Kunzeana Hk. sp. **1**. 177. 1846 = Lindsaya davallioides.
Kurzii Clarke, Tr. Linn. Soc. II. Bot. **1**. 446. 1880 = Microlepia.
lanceolata Steud. Nom. **2**. 146. 1824 = Trichomanes.
lanceolata Bak. Kew Bull. **1899**. 119 = Asplenium Bakeri.
Lapeyrousii Hk. 2 Cent. t. 56. 1861 = Lindsaya hymenophylloides.
Lenormandi Bak. Syn. 471. 1874 = Odontosoria.
lenta (Poir.) Steud. Nom. **2**. 146. 1824 — Madagascar.
Trichomanes Poir. Enc. **8**. 80. 1808.
lepida Pr. Tent. 128. 1836 = Humata.
(D) **leptocarpa** Mett.; Kuhn, Linn. **36**. 143. 1869. HB. 469 — Aneiteum.
L'Herminieri Kze. Bot. Zeit. **1850**. 132 — Guadeloupe.
Microlepia Moore 1861 — (Saccoloma sp.).
ligulata Wall. List n. 254. 1828 = Acrophorus stipellatus.
Lindeni Hk. sp. **1**. 193 t. 56 B. 1846 = Asplenium theciferum var.
Lindleyi Hk. sp. **1**. 163 t. 58 B. 1846 = D. solida var.
?lobata Desv. Prod. 315. 1827 = Lindsaya decomposita.
(D) **Lobbiana** Moore, Ind. 296. 1861. HB. 95. NPfl. 214 — Borneo.
lobulosa Wall. List n. 241. 1828 = Humata heterophylla.
lonchitidea Wall. List n. 240. 1828; Hk. sp. **1**. 173 t. 46 B. 1846 = Microlepia platyphylla.
(D) **longicauda** Christ, Engl. Jahrb. **23**. 339. 1896 — Samoa.
longifolia Roxb. Calc. Journ. **4**. 514. 1844 = D. alata.
longipinnula Ces. Rend. Ac. Napoli **16**. 25, 29. 1877 = Tapeinidium pinnatum.
longula Kze. Bot. Zeit. **1848**. 215 = Humata lepida.
(D) **Lorrainii** Hance, Ann. sc. nat. V. **5**. 254. 1866. HB. 469. NPfl. 214 — Malacca.
Luzonica Hk. sp. **1**. 174 t. 60 B. f. 2, 3, 5. 1846 = Tapeinidium pinnatum var.
(L) **Macgillivrayi** (Fourn.) Bak. Ann. Bot. **5**. 201. 1891 — N. Caledonia.
Leucostegia Fourn. Ann. sc. nat. V. **18**. 344. 1873.
(D) **Macleayi** Bak. Syn. 469. 1874 — »Polynesia«.
Macraeana Hk. et Arn. Bot. Beech. Voy. 108. 1832 = Lindsaya repens.
madagascariensis Kze. Bot. Zeit. **1850**. 173 = Microlepia.
magellanica Desv. Berl. Mag. **5**. 328. 1811 = D. solida.
majuscula Lowe, Ferns 8 t. 33. 1859(?) = Microlepia.

Manilensis Goldm. Nova Acta **19** Suppl. I. 465. 1843 = Microlepia.
?**Mannii** (Eat.) Bak. Syn. 471. 1874 — Ins. Hawaii.
Microlepia Eat. ap. Mann, Proc. Amer. Ac. 7. 212. 1868; Loxoscaphe Kuhn 1879; Lindsaya Hill. 1888; Humata? Diels 1899. NPfl. 210.
marginalis Bak. Syn. ed. I. 452. 1868; ed. II. 98. 1874 = Microlepia marginata.
marginalis Wall. List pag. 64 (n. 322). 1829 = Polypodium marginale Wall.
(D) **Mariesii** Moore, Bak. Ann. Bot. **5**. 201. 1891 — Japonia.
(D) **mauritiana** Hk. sp. **1**. 164 t. 55 B. 1846. HB. 96 — Mauritius. Ins Comorae.
Stenolobus Pr. 1849.
(L) **maxima** (Fourn.) Bak. Ann. Bot. **5**. 201. 1891 — N. Caledonia.
Leucostegia Fourn. Ann. sc. nat. V. **18**. 344. 1873.
meifolia HBK. Nov. Gen. et Sp. **1**. 23. 1815 = Odontosoria.
Melleri Hk.; Bak. Syn. ed. I. 102. 1867 = Odontosoria.
(L) **membranulosa** Wall. List n. 255. 1828 (nomen); Hk. sp. **1**. 159 t. 53 A. 1846. HB. 91 — Yunnan. Nepal. — Acrophorus Moore 1854; Leucostegia J. Sm. 1875; Humata Diels 1899. NPfl. 209.
micans Mett.; Bak. Syn. ed. I. 95. 1867 = D. assamica.
microcarpa Sm. in Rees, Cycl. **11**. —. 1808 = Odontosoria chinensis.
mitis Kze. Bot. Zeit. **1850**. 214 — Guiana. — (Odontosoria sp.).
mollis Kze. Bot. Zeit. **1850**. 172 = Microlepia speluncae.
moluccana Bl. Enum. 237. 1828 = Saccoloma.
moluccana Roxb.; Wall. List n. 2219. 1829 = Tapeinidium amboynense.
Mooreana Masters; Moore, Gard. Chr. **1869**. 964 c. fig. = D. pallida.
(D?) **Moorei** Hk. 2 Cent. t. 53. 1861. HB. 95 — N. Caledonia. Fiji.
Lindsaya Fourn. 1873; Odontosoria Kuhn 1882.
Moritziana Kl.; Kze. Bot. Zeit. **1850**. 54 = Humata.
mucronata Bl. Enum. 235. 1828 = D. decurrens vel D. divaricata.
mucronata Kze. Bot. Zeit. **1848**. 216 = Humata vestita.
(L) **multidentata** (Wall.) Hk. Syn. 91. 1867 — India bor.
Aspidium Wall. 1828 (nomen); Leucostegia Bedd. 1876; Humata Diels 1899. NPfl. 209. Acrophorus Thomsoni Moore 1857 (nomen). Microlepia pteropus Bedd. 1869.
multifida Spr. (Schrad. Journ. **1799**[2]. 271) = Hymenophyllum.
multiflora Roxb. Calc. Journ. **4**. 515. 1844 = Humata Gaimardiana vel Nephrolepis sp. (hirsutula?).
nepalensis Spr. Syst. **4**. 121. 1827 = Microlepia speluncae.
nephrodioides F. Muell. Fragm. **10**. 104. 1877 = ? Dryopteris decomposita var.
(L) **nephrodioides** Bak. Journ. Linn. Soc. **24**. 257. 1887 — Borneo. Formosa.
nigrescens Kze. Bot. Zeit. **1850**. 132 = Saccoloma inaequale var.
nigrescens Hk. 2 Cent. t. 93. 1861 = Asplenium hypomelas.
nipponica Miq. Ann. Lugd. Bat. **3**. 180. 1867 = Microlepia.
nitidula Kze. Linn. **10**. 545. 1836 = D. chaerophylloides.
?nodosa Hk. sp. **1**. 157. 1846 = Acrophorus stipellatus.
Novae Zelandiae Col. Tasm. Journ. **2**. 182. 1844 = Leptolepia.
odontolabia Bak. JoB. **1884**. 140 = Odontosoria.
(L) **oligophlebia** Bak. JoB. **1888**. 323 — Borneo.
ornata Wall. List n. 246. 1828 = D. solida.
padioides Zoll. Syst. Verz. 45. 1854 = Microlepia speluncae.
(D) **pallida** Mett.; Kuhn, Linn. **36**. 142. 1869. HB. 469 — Borneo. Aneiteum. Samoa.
D. Mooreana Masters 1869 — D. Beccariana Ces. 1876.
parallela Wall. List n. 251. 1828 = Humata Gaimardiana.

Parishii Hk. Syn. ed. I. 90. 1867 = D. hymenophylla

Parkeri Hk. sp. 1. 176 t. 53 C. 1846 = Lindsaya.

(L) **parvula** Wall. List n. 247. 1828 (nomen); Hk. et Grev. Ic. Fil. t. 138. 1829. HB. 92 — Malesia. — Leucostegia J Sm. 1842; Acrophorus Moore 1854; Humata Mett. 1856. NPfl. 209.

patens Sw. Syn. 132, 348. 1806 = D. denticulatum.

pectinata Sm. Mém. Ac. Turin 5. 415. 1793 = Humata.

pectinata Bl. Enum. 229. 1828 = D. contigua var.

pectinata Meyen; Goldm. Nova Acta 19 Suppl. I. 465. 1843 = D. alata.

pedata Sm. Mém. Ac. Turin 5. 415. 1793 = Humata repens.

pellucida Desv. Prod. 316. 1827 = ? Humata repens.

(S) **pentaphylla** Bl. Enum. 232. 1828. HB. 94. Chr. 299. NPfl. 214 — Malesia. Polynesia. — Stenolobus Pr. 1849; Scyphularia Fée 1850-52. Humata pinnata Desv. 1827; Davallia Mett.: Kuhn 1869 (non Cav. 1802) — Davallia pycnocarpa Brack. 1854.

(D) **perdurans** Christ, Bull. Boiss. 6. 970. 1898. NPfl. 214 — Yunnan.

phanerophlebia Bak. JoB. 1890. 262 = Microlepia.

philippinensis Harr. Journ. Linn. Soc. 16. 27. 1877 = Microlepia.

pilosa Roxb. Calc. Journ. 4. 515 t. 32. 1844 = Microlepia hirta.

pilosella Hk. 2 Cent. t. 96. 1861 = Microlepia.

pilosiuscula Sm. in Rees, Cycl. 11. —. 1808 — ?

pilosula Wall. List n. 263. 1828 = Microlepia.

Pinkneyi Col. Tr. N. Zeal. Inst. 29. 415. 1897 = Microlepia.

pinnata Cav. Descr. 277. 1802 = Tapeinidium.

pinnata Mett.; Kuhn, Ann. Lugd. Bat. 4. 288. 1869 = D. pentaphylla.

pinnatifida Sw. Syn. 130. 1806 = Humata heterophylla.

pinnatifida Bak. Journ. Linn. Soc. 24. 257. 1887 = Humata intermedia.

(D) **platylepis** Bak. Kew Bull. 1898. 229 — Yunnan.

platyphylla Don, Prod. Fl. Nepal. 10. 1825 = Microlepia.

(D) **plumosa** Bak. JoB. 1876. 10 — Samoa.

polyantha Hk. sp. 1. 168 t. 59 A. 1846 = D. divaricata.

polypodioides Don, Prod. Fl. Nepal. 10. 1825 = Microlepia strigosa.

polypodioides Hk. sp. 1. 181. 1846 = Microlepia speluncae.

polypodioides Benth. Fl. Hongk. 461. 1861 = Microlepia Hancei.

polypodioides γ hispida Hk. sp. 1. 182. 1846 = Microlepia pilosula.

polysperma Steud. Nom. 2. 147. 1824 = Odontosoria chinensis.

Preslii Hk. sp. 1. 161. 1846 = D. contigua.

procera Hedw. Fil. Gen. et Sp. t. 24. 1803 = D. solida.

proxima Bl. Enum. 238. 1828 = Microlepia.

proxima β remota Bl. Enum. 238. 1828 = Saccoloma moluccanum.

proxima Thwait. Enum. pl. Zeyl. 438. 1864 = Microlepia majuscula.

(L) **pseudo-cystopteris** Kze. Bot. Zeit. 1850. 68 — India bor. Acrophorus Moore 1857; Leucostegia Bedd. 1876.

? puberula Wall. List n. 262. 1828 = Microlepia hirta.

pulchella Hk. sp. 1. 175 t. 53 B. 1846 = Lindsaya adiantoides.

(L) **pulcherrima** Bak. Kew Bull. 1895. 53 — Yunnan. Humata Diels 1899. NPfl. 210.

(L) **pulchra** Don, Prod. Fl. Nepal. 11. 1825. Chr. 303 — Yunnan. Himalaya-Ceylon. Leucostegia J. Sm. 1842; Acrophorus Moore 1857; Humata Diels 1899. NPfl. 209. Davallia chaerophylla Wall. 1828. HB. 92; Leucostegia

J. Sm. 1842; Acrophorus Moore 1854; Humata Mett. 1856; Microlepia Ettingsh. 1865.
pumila Willd.; Moore, Ind. 299. 1861 = Dicksonia.
pusilla Mett. Ann. sc. nat. IV. **15**. 79. 1861 = Humata.
pycnocarpa Brack. Expl. Exp. **16**. 242 t. 35 f. 2. 1854 = D. pentaphylla.
pyramidata Wall. List n. 261. 1828 = Microlepia hirta.
(D) **pyxidata** Cav. Descr. 278. 1802. HB. 96. Chr. 301. NPfl. 214 — Australia. Humata Desv. 1827.
pyxidata Klf. Enum. 221. 1824 = D. solida.
(P) **Reineckei** Christ, Engl. Jahrb. **23**. 341 t. 5 f. 1. 1896. Chr. 306. NPfl. 214 [— Samoa.
remota Klf. Enum. 223. 1824 = Odontosoria chinensis.
repens Desv. Prod. 314. 1827 = Lindsaya.
repens Kuhn, Fil. Deck. 27. 1867 = Humata.
retusa Cav. Descr. 278. 1802 = Odontosoria.
rhomboidea Wall. List n. 257. 1828 (nomen); Kze. Bot. Zeit. **1850**. 158 = Microlepia trapeziformis.
rhomboidea Hk. 2 Cent. t. 48. 1860 = Microlepia Wilfordii.
rigida Sol.; Salom. Nom. 155. 1883 = Humata.
Roxburghii Wall. List n. 2218. 1829 = Microlepia hirta.
Saccoloma Spr. Syst. **4**. 119. 1827 = Saccoloma elegans.
Sarasinorum Christ, Verh. Nat. Ges. Basel **11**. 425. 1896 = Microlepia ma- [nilensis.
scabra Don, Prod. Fl. Nepal. 9. 1825 = Microlepia marginata.
scandens Sw. Schrad. Journ. **1800**². 89. 1801 = Lygodium japonicum.
scandens Moore, Ind. 299. 1861 = Odontosoria.
Schimperi Hk. sp. **1**. 193 t. 50 A. 1846 = Asplenium theciferum*.
schizophylla Bak. Syn. 468. 1874 = Lindsaya orbiculata var.
Schlechtendalii Pr. Tent. 129. 1836 = Odontosoria.
scoparia Hk. Syn. 101. 1867 = Odontosoria.
serraeformis Wall. List n. 249. 1828 = D. alata.
serrata Willd. sp. **5**. 467. 1810 = Humata repens.
serrata Roxb. Calc. Journ. **4**. 514. 1844; Hk. sp. **1**. 174. 1846 (= biserrata Bl.) = Tapeinidium pinnatum.
sessilifolia Bl. Enum. 231. 1828 = Humata.
setosa Sm. in Rees, Cycl. **11**. —. 1808 —?
setosa Bak. Syn. 468. 1874 = Cystopteris.
Sieboldiana Miq. Ann. Lugd. Bat. **3**. 180. 1867 = Microlepia marginata.
Sloanei Jenm. JoB. **1886**. 37 = Saccoloma.
(D) **solida** (Forst.) Sw. Schrad. Journ. **1800**². 87. 1801. HB. 95. Chr. 300. NPfl. 214 — Malesia. Polynesia. Queensland. — Trichomanes Forst. Prod. 86. 1786; Humata Desv. 1827; Stenolobus Pr. 1836 — Davallia caudata Cav. 1802; Parestia Pr. 1849 — Davallia ornata Wall. 1828; Stenolobus Pr. 1836 — Davallia Lindleyi Hk. 1846 — D. Fejeensis Hk. 1846. HB. 97 NPfl. 214 (fijiensis); Stenolobus Pr. 1849; Odontoloma J. Sm. 1875.
(D) **speciosa** Mett.; Kuhn, Linn. **36**. 145. 1869. HB. 469 — Moulmein.
Speluncae Bak. Syn. 100. 1867 = Microlepia.
splendens Bl. Enum. 234. 1828 = D. solida.
Sprucei Bak. Fl. bras. **1**². 346. 1870 = Odontosoria meifolia.
stenocarpa Kl. Allg. Gartenzeit. **1855**. 65 — ? India occ. (Hort.).
(D) **stenoloba** Bak. Malesia **3**. 35. 1886 — Amboyna.
stenomera Kze. Bot. Zeit. **1848**. 216 = D. trichomanoides.

stipellata Wall. List n. 260. 1828 = Acrophorus.
stolonifera Bak. Syn. 468. 1874 = Lindsaya.
strigosa (Sw. Adnot. 69. 1829) Kze. Bot. Zeit. **1848**. 542 = Microlepia.
stylosa Steud. Nomencl. **2**. 147. 1824 = Trichomanes.
subimbricata Bl. Enum. 231. 1828 = Humata repens.
subvolubilis Kze. Bot. Zeit. **1850**. 158 = Microlepia speluncae.
tahitensis Brack. Expl. Exp. **16**. 245. 1854 = D. solida.
(D) **Tasmani** Field, Ferns N. Zeal. 75. 1890; Cheeseman, Tr. N. Zeal. Inst. **23**. 416. 1891 — Ins. Kermadec.
tegularis Desv. Prod. 316. 1827. (Plum. t. 99 A) — Hispaniola.
Stenoloma? Fée 1866.
tenuifolia Sw. Schrad. Journ. **1800**^{2}. 88. 1801 = Odontosoria chinensis.
thalictroides Pr. Tent. 129. 1836 = Odontosoria Melleri.
thecifera HBK. Nov. Gen. et Sp. **1**. 23. 1815 = Asplenium.
Thwaitesii Bak. Syn. ed. I. 99. 1867 = Microlepia majuscula.
trapeziformis Roxb. Calc. Journ. **4**. 516. 1844 = Microlepia.
trapezoides Lodd. Cat. 1849 — Hort.
triangularis Bak. Ann. Bot. **5**. 202. 1891 = Cystopteris sudetica.
(D) **trichomanoides** Bl. Enum. 238. 1828. HB. 95 — Malesia.
D. stenomera Kze. 1848.
trichomanoides Hk. 2 Cent. t. 64. 1861 = D. ferulacea.
trichomanoides Bedd. Ferns br. Ind. t. 178. 1866 = Lindsaya orbiculata.
trichosticha Hk. sp. **1**. 183. 1846 = Microlepia.
trifoliata (L.) Sw. Syn. 133. 1806. (Plum. t. 99 B.) — Hispaniola.
Adiantum L. sp. **2**. 1095. 1753; Stenoloma? Fée 1866.
triloba (L.) Willd. sp. **5**. 468. 1810 (Plum. t. 99 C.) — Hispaniola.
Adiantum L. sp. **2**. 1095. 1753; Odontosoria? Fée 1866.
(S) **triphylla** Hk. sp. **1**. 162 t. 46 A. 1846. HB. 94. NPfl. 214 — Singapore.
Stenolobus Pr. 1849; Scyphularia Fée 1850-52.
tripinnata F. Muell.; Benth. Fl. austr. **7**. 717. 1878 = Leptolepia.
triquetra Bak. Syn. 93. 1867 = Lindsaya tenuifolia.
tristis Racib. Pterid. Buit. 131. 1898 = Alsophila.
Truffautiana ht.; L'Ill. Hort. **1895**. 186; Gard. Mag. **1896**. 352 — Hort.
truncata Don, Prod. Fl. Nepal. 10. 1825 — Nepal.
Tyermanni Moore, Gard. Chr. **1871**. 871 (syn.) = Humata.
uncinella Kze. Bot. Zeit. **1850**. 213 = Odontosoria.
urophylla Wall. List n. 2683. 1829 = Microlepia marginata.
urophylla Hk. Fil. exot. t. 19 in nota. 1859 = Microlepia caudigera.
(D) **Veitchii** Bak. JoB. **1879**. 39 — Borneo.
venusta Schkuhr, Kr. Gew. **1**. 122 t. 128. 1809 = Odontosoria chinensis.
vestita Bl. Enum. 233. 1828 = Humata.
villosa Don, Prod. Fl. Nepal. 10. 1825 = Microlepia hirta.
villosa Wall. List n. 244. 1828; Hk. sp. **1**. 172 t. 48 A. = Microlepia marginata.
virens Wall. List n. 264. 1828 = Microlepia speluncae.
(L) **viscidula** Mett.; Kuhn, Linn. **36**. 145. 1869. HB. 467 — Java.
Vogelii Hk. sp. **1**. 168 t. 59 B. 1846 = D. denticulata var.
Wilfordii Bak. Syn. 98. 1867 = Microlepia.
(L) **yaklaensis** (Bedd.) C. Chr. Ind. 215. 1905 — Sikkim.
Leucostegia Bedd. Handb. Suppl. 13. 1892.
(L) **yunnanensis** Christ, Bull. Boiss. **6**. 970. 1898 — Yunnan.

Davalliastrum Fournier, Ann. sc. nat. V. **18**. 334. 1873 *(Lindsaya §)* = Lindsaya et Odontosoria.

DAVALLIOPSIS Van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 323. 1861 = **Trichomanes.**

Definitifolium Hieron. Engl. Jahrb. **34**. 479. 1904 *(Gymnogramme §)* = Gymnogramma.

DENDROGLOSSA Presl, Epim. Bot. 149. 1849 = **Leptochilus.**

lanceolata Pr. Epim. 149. 1849 = L. Linnaeanus.

latifolia Fée, Gen. 81. 1850-52.

normalis Pr. Epim. 149. 1849 = L. lanceolatus var.

quercifolia Fée, Gen. 80 t. 7 B. f. 2. 1850-52 = L. zeylanicus.

subquinquefida Fée, Gen. 80 t. 7 B. f. 1. 1850-52 = L. latifolius.

taccaefolia Fée, Gen. 80 t. 7 B. f. 3. 1850-52 = L. latifolius.

DENNSTAEDTIA Bernhardi, Schrader's Journ. **1800**². 124. 1801 — (Polypodiaceae Gen. **36**).

adiantoides (H. B. Willd.) Moore, Ind. XCVII. 1857 — Amer. trop. Dicksonia H. B. Willd. sp. **5**. 488. 1810. HB. 52 (pt.); Sitolobium J. Sm. 1842 — Dicksonia apiifolia Hk. 1844. HB. 53 (non Sw. 1801); Sitolobium J. Sm. 1842; Dennstaedtia Moore 1857 (pt). ? NPfl. 218. — an Dicksonia bipinnata Cav. 1802? an Polypodium globuliferum Poir. 1804? — conf. Kuhn, Bot. Zeit. **1869**. 164.

ampla (Bak.) Bedd. JoB. **1893**. 227 — Borneo. Perak. Dicksonia Bak. Journ. Linn. Soc. **22**. 223. 1886.

anthriscifolia (Bory) Moore, Ind. 303. 1861; Kuhn, Bot. Zeit. **1869**. 391 — Ins. Mascarenae. — Lonchitis Bory in Willd. 1810 (syn.); Cheilanthes Willd. sp. **5**. 461. 1810; Dicksonia Klf. 1824; ? Sitolobium J. Sm. 1866; Dicksonia rubiginosa β HB. 53.

antillensis (Jenm.) C. Chr. Ind. 216. 1905 — Jamaica. Dicksonia Jenm. JoB. **1886**. 267.

apiifolia Moore, Ind. XCVII. 1857 = D. adiantoides et D. dissecta.

appendiculata (Wall.) J. Sm. Hist. Fil. 265. 1875. Chr. 312. NPfl. 217 — Himalaya. — Dicksonia Wall. 1828 (nomen); Hk. sp. **1**. 79 t. 27 C. 1844. HB. 54; Patania Bedd. 1876. Sitolobium glutinosum J. Sm. 1842 (nomen); Dicksonia Hk. 1844 — D. Elwesii Bak. 1874; Patania Bedd. 1876; Dennstaedtia Bedd. 1883. NPfl. 217.

cicutaria (Sw.) ? Moore, Ind. XCVII. 1857; Kuhn, Chaetopt. 348. 1882. ? Chr. 311. ? NPfl. 218 — Amer. trop. — Dicksonia Sw. Schrad. Journ. **1800**². 91. 1801 (vix HB. 53); Sitolobium J. Sm. 1842.

cicutarioides (Fée) Hier. Engl. Jahrb. **34**. 454. 1904 — India occ. Ecuador. Dicksonia Fée, 11 mém. 95 t. 25 f. 2. 1866.

concinna Moore, Ind. XCVII. 1857 = D. Mathewsii.

consanguinea (Kl.) Moore, Ind. 305. 1861 — Venezuela-Ecuador. Dicksonia Kl. Linn. **20**. 445. 1847; Patania Pr. 1849. Dicksonia adiantoides pt. HB. 52.

cornuta (Klf.) Mett. Ann. sc. nat. V. **2**. 260. 1864 — Amer. centr.-Brasilia. Dicksonia Klf. Enum. 227. 1824.

coronata (Sod.) C. Chr. Ind. 216. 1905 — Ecuador. — Dicksonia Sod. Cr. vasc. quit. 48. 1893; D. adiantoides β coronata Sod. 1883.

cuneata (J. Sm.) Moore, Ind. XCVII. 1857 — Ins. Philipp. Sitolobium J. Sm. 1841 (nomen); Dicksonia Hk. sp. **1**. 80 t. 28 C. 1844. HB. 462.

davallioides (R. Br.) Moore, Parker's Cat. 1858; Ind. 305. 1861. Chr. 312. NPfl. 218 — Australia. Tasmania. — Dicksonia R. Br. Prod. Fl. N. Holl. 158. 1810. HB. 54; Sitolobium J. Sm. 1842. Dicksonia nitidula Kze. 1850; Dennstaedtia Moore 1857.

decomposita Christ, Bull. Boiss. II. **5**. 732. 1905 — Costa Rica. Mexico. Dicksonia Christ, Bull. Soc. bot. Belg. **35**. 180. 1896.

deltoidea Moore, Ind. XCVII. 1857 = D. scabra.

dissecta (Sw.) Moore, Ind. 305. 1861 — Amer. trop. Polypodium Sw. Prod. 134. 1788; Dicksonia Sw. 1801; Sitolobium J. Sm. 1842 — ? Dicksonia rubiginosa pt. HB. 53.

distenta (Kze.) Moore, Ind. 306. 1861 — Mexico. Dicksonia Kze. Anal. 39. 1837.

divaricata (Sod.) C. Chr. Ind. 217. 1905 — Ecuador. Dicksonia Sod. Cr. vasc. quit. 48. 1893.

D'Orbignyana Kuhn, Linn. **36**. 146. 1869 — Bolivia.

erosa (Kze.) Moore, Ind. 306. 1861 — Peru. Dicksonia Kze. Linn. **9**. 88. 1834. D. adiantoides pt. HB. 52.

Elwesii Bedd. Handb. 26. 1883 = D. appendiculata var.

erythrorachis (Christ) Diels, Nat. Pfl. 1^4. 218. 1899 — Celebes. Dicksonia Christ, Ann. Jard. Buit. **15**. 86 t. 13 f. 7. 1897.

exaltata (Kze.) Hier. Engl. Jahrb. **34**. 454. 1904 — Venezuela. Dicksonia Kze. Bot. Zeit. **1850**. 59. Microlepia? oedematosa Fée 1850-52.

flaccida (Forst.) Bernh. Schrad. Journ. **1800**^{2}. 124 t. 1 f. 3. 1801. NPfl. 218 — Polynesia. — Trichomanes Forst. Prod. 85. 1786; Cyathea Spr. 1799; Dicksonia Sw. 1801. HB. 53; Microlepia Pr. 1849.

fluminensis (Fée) C. Chr. Ind. 217. 1905 — Brasilia. Microlepia Fée, Cr. vasc. Br. **1**. 151 t. 51 f. 1. 1869.

formosae Christ, Bull. Boiss. II. **4**. 617. 1904 — Formosa.

glabrata (Ces.) C. Chr. Ind. 217. 1905 — N. Guinea. Dicksonia Ces. Rend. Ac. Napoli **16**. 24, 28. 1877; Bak. Ann. Bot. **5**. 198. 1891.

Godeffroyi Luerss. Fil. Graeff. 222 t. 19. 1871 = Aspidium.

gomphophylla (Bak.) C. Chr. Ind. 217. 1905 — Borneo. Dicksonia Bak. Journ. Linn. Soc. **22**. 223. 1886.

grandifrons Christ, Prim. Fl. Costar. **3**. 38. 1901 — Costa Rica. Mexico.

Henriettae (Bak.) Diels, Nat. Pfl. 1^4. 218. 1899 — Madagascar. Dicksonia Bak. Syn. 462. 1874; Ic. pl. t. 1606; Microlepia Kuhn 1879.

hirsuta Mett.; Miq. Ann. Lugd. Bat. **3**. 181. 1867 = Microlepia pilosella.

incisa (Fée) Kuhn, Linn. **36**. 146. 1869 — Guadeloupe. Brasilia. Dicksonia Fée, 11 mém. 94 t. 25 f. 1. 1866.

incurvata (Bak.) C. Chr. Ind. 217. 1905 — Fiji. Dicksonia Bak. JoB. **1879**. 294.

javanica Christ, Bull. Boiss. II. **4**. 617. 1904 = Dicksonia.

Kingi Bedd. JoB. **1893**. 227 — Perak. — Dicksonia Bedd. Handb. Suppl. 6. 1892.

Lagerheimii (Sod.) C. Chr. Ind. 217. 1905 — Ecuador. Dicksonia Sod. Cr. vasc. quit. 50. 1893.

Lambertieana (Remy) Christ, Farnkr. 312. 1897 (Lambertyana) — Chile Bolivia. Dicksonia Remy; Gay, Fl. chil. **6**. 523. 1853.

lindsayiformis (Fée) C. Chr. Ind. 217. 1905 — Brasilia. Microlepia Fée, Cr. vasc. Br. **1**. 152 t. 51 f. 2. 1869.

macrophylla Moore, Ind. XCVII. 1857 = D. obtusifolia.

Mathewsii (Hk.) C. Chr. Ind. 218. 1905 — Peru. Brasilia. Paraguay.
Deparia Hk. sp. 1. 85 t. 30 B. 1844; Dicksonia Fée 1850-52. Davallia concinna Pr. 1825 (non Schrad. 1818); Patania Pr. 1836; Dicksonia Hk. 1844; Dennstaedtia Moore 1857; Deparia Bak. 1866. HB. 55.

moluccana (Bl.) Moore, Ind. XCVII. 1857. NPfl. 218 — Malesia. Formosa. Fiji.
Dicksonia Bl. Enum. 239. 1828. HB. 53; Sitolobium J. Sm. 1857.

Munchii Christ, Bull. Boiss. II. 5. 732. 1905 — Mexico.

nitidula Moore, Ind. XCVII. 1857 = D. davallioides.

Novae Zelandiae Keys. Pol. Cyath. Hb. Bung. 22. 1873 = Leptolepia.

obtusifolia (Willd.) Moore, Ind. 306. 1861. Chr. 311. NPfl. 218 — Amer. trop.
Dicksonia Willd. sp. 5. 483. 1810; Patania Pr. 1836; Sitolobium J. Sm. 1842. Dicksonia adiantoides pt. HB. 52 — ? Davallia arborescens Willd. 1810.

Orbignyana Kuhn, Chaetopt. 348. 1882 = D. D'Orbignyana.

ordinata (Klf.) Moore, Ind. 306. 1861 — Amer. trop.
Dicksonia Klf. Enum. 226. 1824.

Pavoni (Hk.) Moore, Ind. 307. 1861 — Peru.
Dicksonia Hk. sp. 1. 74 t. 26 A. 1844; Sitolobium J. Sm. 1866.

Pearcei (Bak.) C. Chr. Ind. 218. 1905 — Ecuador
Dicksonia Bak. Ann. Bot. 5. 197. 1891.

producta Mett. Ann. sc. nat. V. 2. 260. 1864 — Columbia.
Deparia Bak. 1868. HB. 463.

pubescens (Bak.) C. Chr. Ind. 218. 1905 — Columbia.
Dicksonia Bak. JoB. **1881**. 203.

punctiloba J. Sm. Hist. Fil. 265. 1875 = D. punctilobula.

punctilobula (Michx.) Moore, Ind. XCVII. 1857 — Amer. bor.
Nephrodium Michx. Fl. bor. Am. 2. 268. 1803; Aspidium Sw. 1806; Dicksonia Kze. 1848; Litolobium Newm. 1854; Polypodium punctilobum Poir. 1804; Aspidium Willd. 1810; Sitobolium Desv. 1827; Dicksonia Hk. 1844. HB. 54; Dennstaedtia J. Sm. 1875. Chr. 312. NPfl. 217; Polypodium pilosiusculum Mühl.; Dicksonia Willd. 1809; Adectum Link 1841; Sitolobium J. Sm. 1842; Dicksonia pubescens Schkuhr 1809.

remota (Christ) Diels, Nat Pfl. 1^4. 218. 1899 — Celebes. — Dicksonia Christ, Verh. Nat. Ges. Basel **11**. 423. 1896; Ann. Jard. Buit. **15**. 86 t. 4 f. 8. 1897.

?**resinifera** (Bl.) Mett.; Kuhn, Ann. Lugd. Bat. 4. 290. 1869 — Java.
Cheilanthes Bl. Enum. 138. 1828; Hypolepis Hk. 1852.

rhombifolia (Bak.) C. Chr. Ind. 218. 1905 — N. Guinea.
Dicksonia Bak. JoB. **1890**. 105.

rubicaulis Christ, Bull. Boiss. II. 5. 258, 732. 1905 — Mexico. Costa Rica

rubiginosa (Klf.) Moore, Ind. XCVII. 1857. Chr. 311. NPfl. 218 — Amer. trop.
Dicksonia Klf. Enum. 226. 1824. HB. 53 (pt excl. β); Sitolobium J. Sm. 1842.

samoensis (Brack.) Moore, Ind. 307. 1857 — Polynesia. N. Guinea.
Sitolobium Brack. Expl. Exp. **16**. 274 t. 38 f. 1. 1854; Dicksonia Bak. 1874. HB. 462 — D. delicata F. Muell. 1876.

scabra (Wall.) Moore, Ind. 307. 1861. Chr. 312. NPfl. 218 — China. Asia trop.
Dicksonia Wall. 1829 (nomen); Hk. sp. **1**. 80 t. 28 B. 1844. HB. 54; Patania Bedd. 1876 — Dicksonia deltoidea Hk. 1844; Dennstaedtia Moore 1857.

scandens (Bl.) Moore, Parker's Cat. 1858; Ind. 307. 1861. Chr. 312. NPfl. 218 Malesia. Samoa. — Dicksonia Bl. Enum. 240. 1828. HB. 53; Sitolobium? Brack. 1854. Dicksonia Zippeliana Kze. 1845; Dennstaedtia Moore 1857 — Hypolepis Smithii Fée 1850-52.

Smithii (Hk.) Moore, Ind. 308. 1861 — Ins. Philipp. — Dicksonia Hk. sp. **1.** 80 t. 28 D. 1844. HB. 54; Sitolobium J. Sm. 1877. Dicksonia neglecta Fée [1850-52.
Sodiroi Diels, Nat. Pfl. 1^4. 218. 1899 — Ecuador.
Dicksonia scandens Bak. JoB. **1877**. 162; Ic. pl. t. 1605 (non Bl. 1828).
solida (Lodd.) Moore, Ind. 308. 1861 — Hort.
Sitolobium Lodd. 1849; Dicksonia Kze. Linn. **23**. 249. 1850 (nomen).
Sprucei Moore, Ind. 308. 1861 — Ecuador. — Dicksonia Bak. 1877.
stenochlaena (Fée) Moore, Ind. 308. 1861 — Réunion.
Dicksonia Fée, Gen. 336. 1850-52.
straminea J. Sm. Hist. Fil. 265. 1875 = Balantium.
strigosa J. Sm. Hist. Fil. 265. 1875 = Microlepia.
tenera (Pr.) Mett. Ann. sc. nat. V. **2**. 261. 1864 — Amer. trop.
Dicksonia Pr. Del. Prag. **1**. 189. 1822; Mart. Ic. Cr. Br. 96 t. 66, 72 f. 1; Sitolobium Brack. 1854. Dicksonia flaccida Raddi 1819 (non Sw. 1801).
vagans (Bak.) Diels, Nat. Pfl. 1^4. 218. 1899 — Ecuador.
Dicksonia Bak. JoB. **1877**. 162.
Zippeliana Moore, Ind. XCVII. 1857 = D. scandens.

DEPARIA Hooker et Greville, Icon. Fil. t. 154. 1829.
concinna Bak. Syn. 55. 1866 = Dennstaedtia Mathewsii.
Godefroyi Bak. Ann. Bot. **5**. 199. 1891 = Aspidium Godeffroyi.
Macraei Hk. et Grev. Ic. Fil. t. 154. 1829 = Athyrium proliferum.
Mathewsii Hk. sp. **1**. 85 t. 30 B. 1844 = Dennstaedtia.
Moorii Hk. JoB. **4**. 55 t. 3. 1852 = Aspidium.
nephrodioides Bak. Gard. Chr. **1872**. 253 = ? Dryopteris decomposita var.
producta Bak. Syn. ed. I. 451. 1868; ed. II. 463. 1874 = Dennstaedtia.
prolifera Hk. et Grev. Ic. Fil. addend. 1831 = Athyrium.
triangularis Und. ap. Heller, Minn. bot. St. **1**. 778. 1897 — Ins. Hawaii. (prob. = Athyrium proliferum var.).

Depariopsis Baker, Ann. Bot. **5**. 462. 1891 *(Polypodium §)* = Aspidium.

DERMATOPHLEBIUM Presl, Epim. Bot. 258. 1849 = **Hymenophyllum.** (Sphaerocionium § Pr. l. c. 17).
cristatum Pr. Epim. 258. 1849 = H. fucoides var.
Jamesoni Pr. Epim. 258. 1849.
tomentosum Pr. Epim. 258. 1849.

DESMOPODIUM J. Smith msc. (Moore, Ind. 308. 1861) = **Dryopteris.**
Phegopteris § Desmopobium (err.) J. Sm. Hist. Fil. 233. 1875.

DIACALPE Blume, Enum. pl. Javae 241. 1828 — (Polypodiaceae Gen. **1**).
aspidioides Bl. Enum. 241. 1828. HB. 45. Chr. 286. NPfl. 159 — Asia trop.
Physematium Kze. 1837; Cyathea Moritz 1845-46; Peranema Mett. 1859. Sphaeropteris Hookeriana Wall. 1829; Diacalpe Moore 1854 — D. pseudocaenopteris Kze. 1846.
Burgessiana Trev. Nu. Giorn. bot. It. **7**. 161. 1875 = Woodsia.
foeniculacea Clarke, Tr. Linn. Soc. II. Bot. **1**. 434. 1880 = Polystichum.
fragilis Trev. Nu. Giorn. bot. It. **7**. 160. 1875 = Woodsia.
guatemalensis Trev. Nu. Giorn. bot. It. **7**. 160. 1875 = Woodsia mollis
madagascariensis Fée, Gen. 339. 1850-52 — Ins. Comorae.
manchuriensis Trev. Nu. Giorn. bot. It. **7**. 160. 1875 = Woodsia.
microphylla Moore, Ind. C. 1857 = Cyathea.
peruviana Trev. Nu. Giorn. bot. It. **7**. 160. 1875 = Woodsia crenata var.
pseudo-caenopteris Kze. Bot. Zeit. **1846**. 457 = D. aspidioides.

Diafnia Presl. Epim. Bot. 119. 1849 *(Blechnopsis §)* = Blechnum.

Diagramma Blume, Enum. pl. Jav. 114. 1828 *(Grammitis §)* = Polypodium.
DIBLEMMA J. Smith, Journ. Bot. **3**. 399. 1841; **4**. 65. 1841 = **Polypodium.**
samarensis J. Sm. JoB. **3**. 399. 1841 = P. tenuilore.
DICHASIUM (A. Br.) Fée, Gen. Fil. 302. 1850-52 = **Dryopteris.**
Aspidium § Dichasium A. Braun, Flora **1841**. 710.
parallelogrammum Fée, Gen. 302 t. 23 B. f. 1. 1850-52 = D. filix mas*.
patentissimum Fée, Gen. 302 t. 23 B f. 2. 1850-52 = D. filix mas*.
DICHOREXIA Presl, Abh. böhm. Ges. Wiss. V. **5**. 344. 1848 = **Alsophila.**
gigantea Pr. l. c. 344. 1848 = A. glabra.
latebrosa Pr. l. c. 344. 1848.
DICKSONIA L'Héritier, Sertum anglicum 30. 1788 — (Cyatheaceae Gen. **2**).
abrupta Bory, Voy. **2**. 187 t. 30. 1804 = Nephrolepis.
aculeata Spr. Neu. Entdeck. **3**. 7. 1822 = Hypolepis repens.
adiantoides H. B. Willd. sp. **5**. 488. 1810 = Dennstaedtia.
adiantoides β coronata Sod. Rec. 23. 1883 = Dennstaedtia coronata.
altissima Sm. in Rees, Cycl. **11**. —. 1808 = ? Dennstaedtia adiantoides.
ampla Bak. Journ. Linn. Soc. **22**. 223. 1886 = Dennstaedtia.
andina Phil. Anal. Univ. Chil. **94**. 309. 1896 — Chile. — (an Dennstaedtia?).
angustidens Pr. Tent. 136. 1836 = ? Dennstaedtia dissecta.
antarctica Lab. Nov. Holl. Pl. Spec. **2**. 100 t. 249. 1806. HB. 50. Chr. 314. NPfl. 120 — Australia. Tasmania — Balantium Pr. 1836; Cibotium Billardierii Klf. 1824; Dicksonia F. Muell. 1882.
antarctica Hk.fil. Fl. N. Zeal. **2**. 9. 1854 = D. fibrosa.
anthriscifolia Klf. Enum. 227. 1824 = Dennstaedtia.
antillensis Jenm. JoB. **1886**. 267 = Dennstaedtia.
apiifolia Sw. Schrad. Journ. **1800**[2]. 91. 1801 = Polystichum.
apiifolia Spr. Syst. **4**. 123. 1827 = Dennstaedtia cornuta.
apiifolia Hk. sp. **1**. 77 t. 26 C. 1844. HB. 53 = ? Dennstaedtia adiantoides.
appendiculata Wall. List n. 65. 1828 = Dennstaedtia.
arborescens L'Hérit. Sert. angl. 31 t. 43. 1788. HB. 50. Chr. 313. NPfl. 120 — St. Helena. — Balantium Hk. 1838; Dicksonia integra Sw. 1801; Balantium auricomum Klf. 1824; Dicksonia Spr. 1827.
asperula Fée, Gen. 335. 1850-52 = Dryopteris.
assamica Griff. Notul. **2**. 607. 1849; Ic. pl. as. **2**. t 136 f. 2. 1849 = Cibotium barometz.
auricoma Spr. Syst. **4**. 122. 1827 = D. arborescens.
Baranetz Link, Fil. sp. 166. 1841 } = Cibotium barometz.
Barometz Hk. & Bak. Syn. 49. 1866 } = Cibotium barometz.
Baudouini Fourn. Ann. sc. nat. V. **18**. 347. 1873. NPfl. 120 — N. Caledonia.
Berteriana (Colla) Hk. sp. **1**. 67 t. 23 A. 1844. HB. 51. NPfl. 120 (Berteroana) — Juan Fernandez. ? Polynesia. — Davallia Colla, Mem. Ac. Torino **39**. 37 t. 65. 1836; Balantium Berteroanum Kze. Anal. 40. 1837.
Berteroana Brack. Expl. Exp. **16**. 277. 1854 = D. Brackenridgei.
Billardieri F. Muell. Cens. **1**. 137. 1882 = D. antarctica.
bipinnata Cav. Descr. 274. 1802 = ? Dennstaedtia adiantoides.
Blumei (Kze.) Moore, Ind. 190. 1860 — Java. Celebes.
Balantium Kze. Bot. Zeit. **1848**. 214. B. chrysotrichum Hassk. 1856; Dicksonia Moore 1860. HB. 50. NPfl. 221.
Brackenridgei Mett. Ann. sc. nat. IV. **15**. 81. 1861. HB. 462. NPfl. 120 — Fiji. Samoa. — D. Berteroana Brack. Expl. Exp. **16**. 277. 1854 (non Hk. 1844).
caudata Poir. Enc. Suppl. **2**. 475. 1811 = Cheilanthes.
Chamissoi Hk. & Bak. Syn. 50. 1866 = Cibotium.

chrysotricha Moore, Ind. 190, 311. 1860 = D. Blumei.
cicutaria Sw. Schrad. Journ. **1800**². 91. 1801 = Dennstaedtia.
cicutaria Hk. & Bak. Syn. 53. 1866 = Dennstaedtia cornuta, ordinata et tenera.
cicutarioides Fée, 11 mém. 95 t. 25 f. 2. 1866 = Dennstaedtia.
concinna Hk. sp. **1**. 74. 1844 = Dennstaedtia Mathewsii.
coniifolia Hk. sp. **1**. 70 t. 24 A. 1844 = Balantium.
consanguinea Kl. Linn. **20**. 445. 1847 = Dennstaedtia.
cornuta Klf. Enum. 227. 1824 = Dennstaedtia.
coronata Sod. Cr. vasc. quit. 48. 1893 = Dennstaedtia.
culcita L'Hérit. Sert. angl. 31. 1788 = Balantium.
cuneata Hk. sp. **1**. 80 t. 28 C. 1844 = Dennstaedtia.
davallioides R. Br. Prod. Fl. N. Holl. 158. 1810 = Dennstaedtia.
decomposita Christ, Bull. Soc. bot. Belg. **35**. 180. 1896 = Dennstaedtia.
decurrens Kl.; Kze. Bot. Zeit. **1850**. 58 = Dennstaedtia consanguinea.
delicata F. Muell. Descr. Pap. Pl. **4**. 77. 1876 = Dennstaedtia samoensis.
deltoidea Hk. sp. **1**. 80 t. 28 A. 1844 = Dennstaedtia scabra.
Deplanchei Vieill.; Bak. Syn. 462. 1874 — N. Caledonia.
dissecta Sw. Schrad. Journ. **1800**². 91. 1801 = Dennstaedtia.
distenta Kze. Anal. 39. 1837 = Dennstaedtia.
divaricata Sod. Cr. vasc. quit. 48. 1893 = Dennstaedtia.
domingensis Desv. Prod. 317. 1827 — Hispaniola. — Microlepia Fée 1866.
dubia Gaud. Freyc. Voy. Bot. 367. 1827 = Davallia.
dubia Hk. & Bak. Syn. ed. I. 51. 1866 (pt.) = Balantium stramineum.
Eggersii Prantl, Engl. Jahrb. **24**. 84. 1897 = Saccoloma domingense var.
elegans Mett. Ann. sc. nat. IV. **15**. 81. 1861 = Thyrsopteris.
Elwesii Bak. Syn. ed. II. 54. 1874 = Dennstaedtia appendiculata var.
erosa Kze. Linn. **9**. 88. 1834 = Dennstaedtia.
erythrorachis Christ, Ann. Jard. Buit. **15**. 86 t. 13 f. 7. 1897 = Dennstaedtia.
exaltata Kze. Bot. Zeit. **1850**. 59 = Dennstaedtia.
expansa Klf.; Fée, Gen. 335. 1850-52 = Dennstaedtia dissecta.
fallax Klf.; Sieb. Syn. n. 111 = Davallia dubia.
fibrosa Col. Tasm. Journ. **2**. 179. 1844; Hk. sp. **1**. 68 t. 23 B. 1844. HB. 461. Chr. 315 — N. Zealand. Ins. Chatham. — Balantium Fée 1850-52 — Dicksonia microcarpa Col. 1888 — D. Sparmanniana Col. 1880.
flaccida Sw. Schrad. Journ. **1800**². 90. 1801 = Dennstaedtia.
flaccida Raddi, Opusc. sci. Bol. **3**. 296. 1819 = Dennstaedtia tenera.
flaccida Hk. et Arn. Bot. Beech. Voy. 108. 1832 (pt.) = Leptolepia Andersoni.
fragilis Trevir. Berl. Mag. **7**. 155 t. 3 f. 18-19. 1816 = Woodsia.
gigantea Karst. Fl. Col. **2**. 177 t. 193. 1869 = D. Karsteniana.
glabrata Ces. Rend. Ac. Napoli **16**. 24, 28. 1877 = Dennstaedtia.
glauca Sm. in Rees, Cycl. **11**. —. 1808 = Cibotium.
globulifera O. Ktze. Rev. Gen. Pl. **3**². 378. 1898 = Dennstaedtia Lambertieana.
globuligera Desv. Prod. 317. 1827 = Dennstaedtia cicutaria.
glutinosa Wall.; Hk. sp. **1**. 81. 1844 = Dennstaedtia appendiculata.
gomphophylla Bak. Journ. Linn. Soc. **22**. 223. 1886 = Dennstaedtia.
gracilis Col. Tr. N. Zeal. Inst. **15**. 306. 1883 = D. squarrosa var.
guatemalensis Bak. Syn. 461. 1874 = Cibotium.
Henriettae Bak. Syn. 462. 1874 = Dennstaedtia.
hypolepidoides Bak. Journ. Linn. Soc. **16**. 197. 1877 = Microlepia madagascariensis.
incisa Fée, 11 mém. 94 t. 25 f. 1. 1866 = Dennstaedtia.

incurvata Bak. JoB. **1879**. 294 = Dennstaedtia.
integra Sw. Schrad. Journ. **1800**². 90. 1801 = D. arborescens.
japonica Sw. Schrad. Journ. **1800**². 92. 1801 = Microlepia strigosa.
javanica Bl. Enum. 240. 1828 — Java.
Dennstaedtia Christ 1904 — (an Dennstaedtia sp.?).
Karsteniana (Kl.) Moore, Ind. 190. 1860. Chr. 314. NPfl. 120 — Guatemala. Columbia. — Balantium Kl. Linn. **20**. 444. 1847; Dicksonia Sellowiana var. HB. 461 — D. gigantea Karst. 1869.
Kaulfussiana Gaud. Freyc. Voy. Bot. 368. 1827 = Microlepia hirta.
Kingi Bedd. Handb. Suppl. **6**. 1892 = Dennstaedtia.
laevis Hew. Lond. JoB. **1**. 283. 1842 (nomen) = D. lanata.
Lagerheimii Sod. Cr. vasc. quit. 50. 1893 = Dennstaedtia.
Lambertieana Remy in Gay, Fl. chil. **6**. 523. 1853 = Dennstaedtia.
lanata Col. Tasm. Journ. **2**. 181. 1844; Hk. sp. **1**. 69 t. 23 C. 1844. HB. 461. Chr. 315. NPfl. 120 — N. Zealand. — Balantium Fée 1850-52.
Lathamii Moore, Gard. Chr. n. s. **24**. 584, 689 f. 154. 1885 — Hort.
Lessonii Bory, Bél. Voy. Bot. **2**. 75. 1833 = Microlepia speluncae
Lindeni Hk. sp. **1**. 72 t. 25 B. 1844 = Saccoloma domingense var.
linearis Cav. Descr. 274. 1802 = Tapeinidium pinnatum var.
macrophylla Desv. Prod. 317. 1827 = Dennstaedtia obtusifolia.
?madagascariensis Kze. Anal. 38. 1837 = Microlepia.
marginalis Sw. Schrad. Journ. **1800**². 92. 1801 = Microlepia marginata.
Martiana Kl.; Hk. sp. **1**. 70 t. 24 B. 1844 = Balantium coniifolium.
Mathewsii Fée, Gen. 335. 1850-52 = Dennstaedtia.
Menziesii Hk. & Bak. Syn. 49. 1866 = Cibotium.
microcarpa Col. Tr. N. Zeal. Inst. **20**. 214. 1888 = D. fibrosa var.
millefolium Desv. Prod. 318. 1827 = Microlepia speluncae.
moluccana Bl. Enum. 239. 1828 = Dennstaedtia.
moluccana Roxb.; Wall. List n. 2174. 1829; Roxb. Calc. Journ. **4**. 517. 1844 = Saccoloma sorbifolium.
montevidensis Spr. Syst. **4**. 122. 1827 = Woodsia.
multifida Sw. Schrad. Journ. **1800**². 92. 1801 = Microlepia speluncae.
neglecta Fée, Gen. 335. 1850-52 = Dennstaedtia Smithii.
nephrodioides F. Muell. Cens. **1**. 137. 1882 = ? Dryopteris decomposita var.
nephrolepioides Christ, Verh. Nat. Ges. Basel **11**. 241. 1895 = Nephrolepis dicksonioides.
nitidula Kze. Linn. **23**. 249, 308. 1850 = Dennstaedtia davallioides.
obtusa Moritz; Fée, Gen. 335. 1850-52 = ? Dennstaedtia obtusifolia.
obtusifolia Willd. sp. **5**. 483. 1810 = Dennstaedtia.
ordinata Klf. Enum. 226. 1824 = Dennstaedtia.
organensis Miers; J. Sm. Lond. JoB. **1**. 435. 1842 = D. Sellowiana.
Papuana F. Muell. Descr. Pap. Pl. **4**. 76. 1876 = Saccoloma sorbifolium.
Pavoni Hk. sp. **1**. 74 t. 26 A. 1844 = Dennstaedtia.
Pearcei Bak. Ann. Bot. **5**. 197. 1891 = Dennstaedtia.
pilosiuscula Willd. Enum. 1076. 1809 = Dennstaedtia punctilobula.
pilosiuscula Raddi, Opusc. sci. Bol. **3**. 296 1819 = Dennstaedtia tenera
Plumieri Hk. sp. **1**. 72. 1844 = Saccoloma domingense.
polypodioides Sw. Schrad. Journ. **1800**² 91. 1801 = Microlepia strigosa vel [M. speluncae.
prolifera Klf. Enum. 225. 1824 = Athyrium.
pubescens Sw.; Schkuhr, Kr. Gew. **1**. 125 t. 131. 1809 = Dennstaedtia punctilobula.

pubescens Bak. JoB. **1881**. 203 = Dennstaedtia.
pumila (Willd.) Kze. Bot. Zeit. **1850**. 254 (nomen) — Amer. austr. temp.
Davallia Willd. hb.; Moore 1861 (nomen).
punctiloba Hk. sp. **1**. 79. 1844. } = Dennstaedtia
punctilobula Kze. Amer. Journ. Sci. II. **6**. 88. 1848 } punctilobula.
punctulata Poir. Enc. Suppl. **2**. 475. 1811 — Amer. austr.
recognita Kze. Bot. Zeit. **1850**. 57 = Dennstaedtia obtusifolia.
regalis Bak. Syn. 461. 1874 = Cibotium.
remota Christ, Verh. Nat. Ges. Basel **11**. 423. 1896 = Dennstaedtia.
repens Bory, Voy. **2**. 323. 1804 = Lindsaya.
rhombifolia Bak. Journ. Bot. **1890**. 105 = Dennstaedtia.
rubiginosa Klf. Enum. 226. 1824 = Dennstaedtia.
samoensis Bak. Syn. 462. 1874 = Dennstaedtia.
scabra Wall. List n. 2173. 1829 = Dennstaedtia.
scandens Bl. Enum. 240. 1828 = Dennstaedtia.
scandens Bak. JoB. **1877**. 162 = Dennstaedtia Sodiroi.
Schiedei Bak. Syn. 50. 1866 = Cibotium.
Sellowiana (Pr.) Hk. sp. **1**. 67 t. 22 B. 1844. HB. 51 (excl. varr. 461). Chr. 314. NPfl. 120 — Brasilia. Ecuador. — Balantium Pr. 1836 (nomen). Dicksonia organensis Miers; J. Sm. 1842 (nomen).
siifolia Kze. Bot. Zeit. **1845**. 815 (nomen); Ettingsh. Farnkr. 214 t. 152 f. 3-5. 1865 = Dennstaedtia cicutaria.
Smithii Hk. sp. **1**. 80 t. 28 D. 1844 = Dennstaedtia.
solida Kze. Linn. **23**. 249. 1850 = Dennstaedtia.
sorbifolia Sm. in Rees, Cycl. **7**. —; Hk. sp. **1**. 72 t. 25 A. 1844 = Saccoloma.
Sparmanniana Col. Tr. N. Zeal. Inst. **12**. 363. 1880 = D. fibrosa.
splendens Desv. Prod. 318. 1827 = Cibotium Chamissoi.
Spruceana Mett.; Kuhn, Linn. **36**. 153. 1869. NPfl. 120 — Peru.
D. Sellowiana var. HB. 461.
Sprucei Bak. JoB. **1877**. 162 = Dennstaedtia.
squarrosa (Forst.) Sw. Schrad. Journ. **1800**[2]. 90. 1801. HB. 51. Chr. 314. NPfl. 120 — N. Zealand. — Trichomanes Forst. Prod. 86. 1786; Balantium Kze. 1850 — Dicksonia gracilis Col. 1883.
stenochlaena Fée, Gen. 336. 1850-52 = Dennstaedtia.
straminea Lab. Sert. Austr. Cal. 7 t. 10. 1824 = Balantium.
strigosa Thbg. Tr. Linn. Soc. **2**. 341. 1794 = Microlepia.
tenera Pr. Del. Prag. **1**. 189. 1822 = Dennstaedtia.
thyrsopteroides Mett. Ann. sc. nat. IV. **15**. 81. 1861. HB. 462. NPfl. 121 — N. Caledonia. — Balantium Fourn. 1876.
Torreyana Brack. Expl. Exp. **16**. 278 t. 38 f. 2. 1854 = Balantium stramineum.
umbrosa Liebm. Vid. Selsk. Skr. V. **1**. 262 (seors. 110). 1849 = Dennstaedtia [rubiginosa.
vagans Bak. JoB. **1877**. 162 = Dennstaedtia.
Wendlandi Bak. Syn. 460. 1874 = Cibotium.
Youngiae C. Moore; Moore, Proc. Hort. Soc. —; J. Sm. Ferns br. and for. 291. 1866; Bak. Syn. 461. 1874. NPfl. 120 — N. S. Wales. Queensland.
zeylanica Sw. Schrad. Journ. **1800**[2]. 91. 1801 = Dennstaedtia flaccida.
Zippeliana Kze. Bot. Zeit. **1845**. 838 = Dennstaedtia scandens.

DICLIDOPTERIS Brackenridge. Explor. Exped. **16**. 135. 1854 = **Monogramma.**
angustissima Brack. Expl. Exp. **16**. 135 t. 17. 1854 } = M. paradoxa.
paradoxa Carr. in Seem. Fl. Vit. 370. 1873 }

DICLISODON Moore, Ind. Filicum XCV. 1857 = **Dryopteris.**
deparioides Moore, Ind. XCVI. 1857.
DICRANODIUM Newman, Hist. Brit. Ferns ed. III. 13. 1854 = **Anogramma** (leptophylla).
DICRANOGLOSSUM J. Smith, Bot. Voy. Herald 232. 1854 = **Eschatogramme.**
furcatum J. Sm. Bot. Voy. Herald 232. 1854 } = E. furcata.
subpinnatifidum Moore, Ind. XXXIII. 1857 }
Dicranophlebia Martius, Icon. Sel. Cr. Bras. 63. 1834 *(Alsophila §)* = Alsophila.
DICRANOPTERIS Bernhardi, Schrad. neu. Journ. 1^2. 26, 38. 1806 = **Gleichenia.**
dichotoma Bernh. Schrad. neu. Journ. 1^2. 38, 49 t. 3 f. 13. 1806 = G. linearis.
Dicranopteris Bl.; Endl. Gen. Suppl. I. 1346. 1841 *(Polypodium §)* = Polypodium.
DICTYMIA J. Smith, Bot. Mag. **72** Comp. 16. 1846 = **Polypodium.**
attenuata J. Sm. Bot. Mag. **72** Comp. 16. 1846 = P. Brownii.
lanceolata Hk.fil. Fl. N. Zeal. **2**. 43. 1854 = P. dictyopteris.
DICTYOCLINE Moore, Gard. Chron. **1855**. 854; Index LIX. 1857 = **Aspidium.**
Griffithii Moore, Ind. LIX. 1857 } = A. Griffithii.
Wilfordii J. Sm. Hist. Fil. 149. 1875 }
DICTYOGLOSSUM J. Smith, Bot. Mag. **72** Comp. 18. 1846 = **Elaphoglossum.**
crinitum J. Sm. l. c.
Dictyogramma Presl, Epim. Bot. 148. 1849 *(Colysis §)* = Polypodium.
DICTYOGRAMME Fée, Gen. Fil. 170. 1850-52 (Dyctiogramme Pr.).
Boryana Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
elongata Moore, Ind. LX. 1857 = Syngramma pinnata.
ensiformis Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
gigantea Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
Griffithii Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Aspidium.
hederaefolia Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Hemionitis.
japonica Fée, Gen. 171 t. 15 A. f. 2. 1850-52 = Coniogramme.
lanceolata Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
latifolia Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
latipes Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum semicostatum.
Manniana Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
obtusa Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Antrophyum.
obtusifolia Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Syngramma.
pinnata Moore, Ind. 317. 1861 = Syngramma.
pinnatifida Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Hemionitis.
podophylla Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Gymnopteris Ehrenbergiana.
quinata Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Syngramma.
Smithii Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Hemionitis.
Wallichii Trev. Atti Ist. Veneto V. **3**. 592. 1877 = Syngramma alismifolia var.
Wilfordi Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Aspidium Griffithii var.
Dictyophlebia Gris. Abh. Ges. Gött. **7**. 283, 286. 1857 *(Polypodium* et *Aspidium §)* = Polypodium et Aspidium.
DICTYOPTERIS Presl, Tent. Pterid. 194. 1836 = **Aspidium.**
attenuata Pr. Tent. 194 t. 8 f. 8. 1836 = Polypodium Brownii.
attenuata Hk. Gen. t. 71 B. 1841 = Polypodium dictyopterts.
Barberi Bedd. Ferns br. Ind. t. 322. 1869.
Camerooniana (is) J. Sm. Ferns br. and for. 285. 1866 = A. varians.
chattagrammica Bedd. Handb. 299. 1883 = A. membranifolium.
Cumingiana Pr. Epim. 61. 1849 = A. macrodon.

difformis Moore, Ind. 90. 1858 = A. irregulare.
heterosora Bedd. Ferns br. Ind. Suppl. 20. 1876 = A. vastum.
irregularis Pr. Tent. 194 t. 8. f. 7. 1836.
lanceolata J. Sm. JoB. **4**. 64. 1841 = Polypodium dictyopteris.
lifuensis Fourn. Ann. sc. nat. V. **18**. 291. 1873.
macrodonta Pr. Tent. 194. 1836.
megalocarpa J. Sm. Hist. Fil. 195. 1875.
petrophya Moore, Ind. 318. 1861 = A. saxicola.
polycarpa Bedd. Ferns br. Ind. Suppl. 20. 1876 = A. Kuhnii.
pteroides Pr. Tent. 194 t. 8 f. 6, 13. 1836 = A. Brongniartii.
tenerifrons Bedd. Ferns br. Ind. t. 4. 1865.
varians Moore, Gard. Chr. **1863**. 1108.

DICTYOXIPHIUM Hooker, Gen. Fil. t 62. 1840 — (Polypodiaceae Gen. **39**).
Michlerianum Moore, Ind. 319. 1861 = Lindsaya.
panamense Hk. Gen. t. 62. 1840. HB. 113. Chr. 298. NPfl. 219 — Guatemala-Columbia. — Lindsaya Mett. 1856. Amphiblestra simplex Fourn. 1873.

DIDYMOCHLAENA Desvaux, Berl. Mag. **5**. 303. 1811 — (Polypodiaceae Gen. **13**).
dimidiata Kze. Linn. **18**. 122. 1844 = D. truncatula.
lunulata Desv. Prod. 282. 1827 = D. truncatula.
polycarpa Bak. Syn. 248. 1867 = Mesochlaena.
sinuosa Desv. Berl. Mag. **5**. 303 t. 7. f. 6. 1811 = D. truncatula.
squamata Desv. Prod. 282. 1827 = D. truncatula.
truncatula (Sw.) J. Sm. JoB. **4**. 196. 1841 — Malesia. Polynesia. Africa trop. Madagascar. Amer. trop. — Aspidium Sw. Schrad. Journ. **1800**[2]. 36. 1801; Adiantum lunulatum Houtt. Nat. Hist. **14**. 209. 1783 (non Burm. 1768); Didymochlaena Desv. 1827. HB. 248. Chr. 287. NPfl. 183; Nephrolepis Keys. 1873 — Didymochlaena sinuosa Desv. 1811; Monochlaena Gaud. 1827. Aspidium squamatum Willd. 1810; Didymochlaena Desv. 1827. Diplazium pulcherrimum Raddi 1819 — Aspidium cultratum Pr. 1822 — Tegularia adiantifolia Reinw. 1824 — Didymochlaena dimidiata Kze. 1844.

Didymochlamys Moore, Index LV. 1857 (*Diplazium §*) = Diplazium.

DIDYMOGLOSSUM Desvaux, Prodrome 330. 1827 = **Trichomanes.**
acanthoides v.d.B. Pl. Jungh. **1**. 560. 1856 } = Hymenophyllum aculeatum.
aculeatum v.d.B. Pl. Jungh. **1**. 559. 1856 } = Hymenophyllum aculeatum.
affine v.d B. Pl. Jungh. **1**. 562. 1856 = Hymenophyllum holochilum.
alatum Desv. Prod. 331. 1827 = T. radicans.
alatum Pr. Hym. 23. 1843 = T. bipunctatum.
angustifrons Fée, 11 mém. 113 t. 28 f. 5. 1866 = T. pusillum.
anomalum v.d.B. Ned. kr. Arch. **5**[3]. 140. 1863 = T. bipunctatum.
apodum v.d.B. Ned. kr. Arch. **4**. 380. 1859 = T. hymenoides.
bilabiatum v.d.B. Pl. Jungh. **1**. 557. 1856.
bipunctatum Fourn. Ann. sc. nat. V. **18**. 263. 1873.
Braunii v.d.B. Pl. Jungh. **1**. 560. 1856 = Hymenophyllum brachyglossum.
brevipes Pr. Hym. 23, 47. 1843 = T. bilabiatum.
capillatum Pr. Hym. 65. 1843 = T. bipunctatum var.
cordifolium Fée, 11 mém. 113 t. 28. f. 4. 1866 = ? T. hymenoides.
crispulum Fée, 11 mém. 112. 1866 — India occ.
decipiens Desv. Prod. 330 t. 7 f. 3. 1827 = T. bipunctatum.
denticulatum Hassk. Obs. Bot. Fil. **2**. 16. 1857-58 = Hymenophyllum d.
dilatatum v.d.B. Ned. kr. Arch. **5**[3]. 144. 1863 = T. bipunctatum.

euphlebium v.d.B. Ned. kr. Arch. **5**^{3}. 142. 1863 = T. bipunctatum.
ferox Hassk. Obs. Bot. Fil. **2**. 17. 1857-58 — Java.
filicula Desv. Prod. 331. 1827 = T. bipunctatum.
fructuosum Fée, 11 mém. 112 t. 28 f. 3. 1866 = T. Kraussii.
fuscum Hassk. Obs. Bot. Fil. **2**. 19. 1857-58 = Hymenophyllum f.
Griffithii v d.B. Ned. kr. Arch. **5**^{3}. 141. 1863 = T. bipunctatum.
Hedwigii Pr. Abh. böhm. Ges. V. **5**. 335. 1848 = ? T. reptans.
holochilum v.d.B. Pl. Jungh. **1**. 561. 1856 = Hymenophyllum h.
Hookeri Pr. Hym. 23. 1843 = T. sphenoides.
humile Pr. Hym. 23. 1843.
insigne v.d.B. Ned. kr. Arch. **5**^{3}. 143. 1863 = T. bipunctatum.
intramarginale Pr. Abh. böhm. Ges. V. **5**. 335. 1848.
Kraussii Pr. Hym. 23. 1843.
laceratum Fée, 11 mém. 113 t. 32 f. 1. 1866 = T. lineolatum.
latealatum v.d.B. Ned. kr. Arch. **5**^{3}. 138. 1863 = T. bipunctatum var.
laxum v.d.B. Hym. Jav. 37 t. 27. 1861 = T. bilabiatum.
lineare Desv. Prod. 331. 1827 = Hymenophyllum l.
lineolatum v.d.B. Ned. kr. Arch. **5**^{3}. 136. 1863.
longisetum Pr. Hym. 23, 49. 1843 = T. cupressoides.
magellanicum Desv. Prod. 331. 1827 = Hymenophyllum m.
minutulum Pr. Hym. 23. 1843 = T. humile.
muscoides Desv. Prod. 330. 1827 = T. hymenoides.
Neesii Pr. Hym. 23. 1843 = Hymenophyllum denticulatum.
nummularium v.d.B. Ned. kr. Arch. **5**^{3}. 135. 1863.
obscurum Hassk. Obs. Bot. Fil. **2**. 22. 1857-58 = T. cupressoides.
ovale Fourn. Bull. Soc. Fr. **19**. 240. 1872 = T. punctatum.
Pabstianum v.d.B. Ned. kr. Arch. **4**. 380. 1859.
plicatum v.d.B. Ned. kr. Arch. **5**^{3}. 139. 1863 = T. bipunctatum.
pumilum Hassk. Obs. Bot. Fil. **2**. 14. 1857-58 — Java.
punctatum Desv. Prod. 330. 1827.
pusillum Desv. Prod. 330. 1827.
quercifolium Pr. Hym. 23. 1843 = T. montanum.
racemulosum v.d.B. Ned. kr. Arch. **5**^{3}. 137. 1863 = T. bipunctatum.
reptans Pr. Hym. 23. 1843.
Schaffneri v.d.B. Ned. kr. Arch. **4**. 380. 1859 = T. hymenoides.
serrulatum Pr. Hym. 23, 48. 1843 = Hymenophyllum s.
Simonsianum v.d.B. Ned. kr. Arch. **5**^{3}. 145. 1863 = Hymenophyllum S.
sociale Fée, Cr. vasc. Br. **2**. 85 t. 85 f. 3. 1872-73.
sphenoides Pr. Hym. 23. 1843.
undulatum Pr. Hym. 23, 48. 1843 = T. bipunctatum.
Zollingeri Hassk. Obs. Bot. Fil. **2**. 20. 1857-58 = Hymenophyllum Zollingerianum.

DIELLIA Brackenridge, U. S. Expl. Exp. **16**. 217. 1854 — (Polypodiaceae Gen. **30**).
Alexandri (Hill.) Diels. Nat. Pfl. **1**^{4}. 212 f. 114 G-K. 1899 — Ins. Hawaii
Lindsaya Hill Fl. Haw. 622 1888.
centifolia (Hill.) Diels, Nat. Pfl. **1**^{4}. 212 f. 114 D. 1899 — Ins. Hawaii.
Lindsaya Hill. Fl. Haw. 621. 1888.
erecta Brack. Expl. Exp. **16**. 218 t. 31 f. 2. 1854. NPfl. 212 — Ins. Hawaii.
Schizoloma Moore 1857; Lindsaya Mett. 1861. HB. 113.
falcata Brack. Expl. Exp. **16**. 219 t. 31 f. 1. 1854. NPfl. 212 — Ins. Hawaii.
Schizoloma Moore 1857; Lindsaya Hk. 1867. HB. 113.

Knudsenii (Hill.) Diels, Nat. Pfl. **1**⁴. 212 f. 114 f. L-N. 1899 — Ins. Hawaii.
Lindsaya Hill. Fl. Haw. 623. 1888.
laciniata (Hill.) Diels, Nat. Pfl. **1**⁴. 212 f. 114 E-F. 1899 — Ins. Hawaii.
Lindsaya Hill. Fl. Haw. 621. 1888.
pumila Brack. Expl. Exp. **16**. 219. 1854. NPfl. 212 — Ins. Hawaii.
Schizoloma Moore 1861; Lindsaya Hk. 1867. HB. 112. Chr. 298.
Digramma Kunze, Bot. Zeit. **1848**. 209 (*Antrophyum §*) = Cyclophorus.
DIGRAMMARIA Presl, Tent. Pterid. 116. 1836.
ambigua Pr. Tent. 117 t. 4 f. 12. 1836 = Aspidium.
ambigua Hk. Gen. t. 56 C. 1840 } = Diplazium esculentum.
esculenta Fée, Gen. 217. 1850-52 }
robusta Fée, Gen. 218. 1850-52 = Diplazium proliferum.
DIPLAZIOPSIS C. Chr. Index Fil. 227. 1905 — (Polypodiaceae Gen. **43**).
Allantodia Wallich, Pl. asiat. rar. **1**. 44. 1830 (non R. Brown 1810).
javanica (Bl.) C. Chr. Ind. 227. 1905 — China austr. India bor.-Malesia-Polynesia. — Asplenium Bl. Enum. 175. 1828; Allantodia Trevis. 1875. Chr. 213. NPfl. 229; A. Brunoniana Wall. Pl. asiat. rar. **1**. 44 t. 52. 1830. HB. 426; Asplenium Mett. 1856; Athyrium Milde 1870; Hemidictyum? Brunonis Pr. 1836 —? H. Drummondii Fourn. 1873.
DIPLAZIUM Swartz, Schrad. Journ. **1800**². 61. 1801 — (Polypodiaceae Gen. **42**).
accedens Bl. Enum. 192. 1828 = D. proliferum.
acuminatum Raddi, Pl. Bras. **1**. 41 t. 57 f. 2. 1825 = D. plantaginifolium.
acuminatum Bl. Enum. 193. 1828 = D. Christii.
acuminatum Pr. Tent. 113. 1836 = D. silvaticum.
acutale Fée, Gen. 215. 1850-52 = D. denticulosum.
aemulum Und. et Maxon, Bull. Torr. Cl. **29**. 583. 1902 — Cuba.
aequibasale (Bak.) C. Chr. Ind. 227. 1905 — Borneo.
Asplenium Bak. Journ. Linn. Soc. **22**. 225. 1886.
affine J. Sm. JoB. **3**. 407. 1841 (nomen) — Ins. Philipp.
Athyrium Milde 1870 (nomen); Asplenium Mett.; Salom. 1883 (nomen).
alismaefolium Pr. Rel. Haenk. **1**. 49 t. 8 f. 3. 1825 = Syngramma.
alternifolium Bl. Enum. 190. 1828 = D. bantamense var.
altissimum (Jenm.) C. Chr. Ind. 227. 1905 — Jamaica.
Asplenium Jenm. JoB. **1879**. 259 — A. Taylorianum Jenm. 1886.
ambiguum Raddi, Opusc. sci. Bol. **3**. 292. 1819; Pl. Bras. **1**. 41 t. 58. 1825 — Amer. trop. — Athyrium Milde 1870. Asplenium dubium Mett. n. 216. 1859 (non Gaud. 1827 nec Brack. 1854). A. radicans pt. HB. 241 (non L. 1759, vix Schkuhr 1809); Diplazium Chr. 1897. 220. NPfl. 228 (pt.?) — D. leptocarpon, leptochlamys, remotum, rostratum Fée 1869 — Gymnogramme expansa Fée 1869.
ambiguum Hk. JoB. **9**. 343. 1857 = D. esculentum.
ambiguum J. Sm. Cat. cult. ferns 47. 1857 = D. Shepherdii.
amplissimum (Bak.) Diels, Nat. Pfl. **1**⁴. 227. 1899 — Amboyna.
Asplenium Bak. Ann. Bot. **5**. 310. 1891.
amplum Liebm. Vid. Selsk Skr. V. **1**. 256 (seors. 104). 1849 = D. striatum.
angulosum C. Chr. Ind. 227. 1905 — Ecuador.
Asplenium reflexum Sod. Cr. vasc. quit. 205. 1893 (non Bory 1833).
angustatum Amann, Nat. Tijdschr. Ned. Ind. **23**. 406. 1861 — Bangka.
angustifrons Pr. Tent. 114 1836; Epim. 85. 1849 = D. arboreum.
anthraxacolepis Fée, 8 mém. 84. 1857 = D. Shepherdii.

Apollinaris L'Herm.: Fée, 11 mém. 40 t. 10 f. 1. 1866 = D. costale.

arborescens (Bory) Sw. Syn. 92. 1806. Chr. 220. NPfl. 226 — Ins. afric. orient. Afr. orient. S. Thomé. — Callipteris Bory, Voy. **1**. 283. 1804; Asplenium Mett. 1856; Aspl. n. 215. HB. 240; Athyrium Milde 1870 — Diplazium serrulatum Desv. 1827 — Asplenium madagascariense Bak. 1876 — A. nemorale Bak. 1876.

arboreum (Willd.) Pr. Tent. 114. 1836. NPfl. 226 — Ind. occ. Venezuela-Ecuador. — Asplenium Willd. sp. **5**. 320. 1810. HB. 233; Athyrium Milde 1870. Diplazium auriculatum Klf. 1824; Asplenium Mett. n. 172. 1859

Arnottii Brack. Expl. Exp. **16**. 144. 1854. NPfl. 226 — Ins Hawaii. Asplenium Bak. 1867. HB. 240; Athyrium Milde 1870; Asplenium diplazioides Hk. et Arn. Bot. Beech. Voy. 107. 1832 (non Bory 1833).

asperum Bl. Enum. 195. 1828. (HB. 238). Chr. 219 — India bor.-Java. Microstegia Pr. 1849; Athyrium Milde 1870 — Asplenium sikkimense Clarke 1880.

aspidioides Klf.; Pr. Tent. 114. 1836 (nomen) —?

asplenioides (Kze.) Pr. Tent. 114 t. 4 f. 4. 1836 — Peru. Allantodia Kze. Linn. **9**. 72. 1834; Athyrium Milde 1870 (non Desv. 1827); Asplenium allantodioides Mett. n. 228. 1859.

assimile Bedd. Ferns br. Ind. t. 294. 1868 = Athyrium.

athyrioides Moore, Ind. LV. 1857 = D. expansum.

auriculatum Klf. Enum. 183. 1824 = D. arboreum.

bantamense Bl. Enum. 191. 1828. Chr. 216. NPfl. 226 — Asia trop. Asplenium Bak. 1867. HB. 231; Athyrium Milde 1870. Asplenium fraxinifolium Wall. 1828; A. Donianum Mett. n. 198 b. 1859 — Diplazium alternifolium Bl. 1828; Asplenium Mett. 1866 (non 1856); Athyrium Milde 1870.

barbadense hort.; Kze. Linn. **23**. 250. 1850 — Hort. (Ind occ.).

barisanicum (Bak.) C. Chr. Ind. 228. 1905 — Sumatra. Asplenium Bak. Ann. Bot. **8**. 126. 1894.

Beddomei C. Chr. Ind. 228. 1905 — Ceylon. — Asplenium Schkuhrii Thwaites, Enum. pl. Zeyl. 385. 1864. HB. 491 (non Mett. 1859); Diplazium Bedd. 1863-65 (non J. Sm. 1841). — (an = D. microphyllum Desv.?).

bifrons (Sod.) C. Chr. Ind. 228. 1905 — Ecuador. Asplenium Sod. Cr. vasc. quit. 184. 1893.

bipartitum Pr. Epim. 88. 1849 = Asplenium.

biseriale (Bak.) C. Chr. Ind. 228. 1905 — Borneo. Asplenum Bak. Tr. Linn. Soc. II. Bot. **4**. 252. 1894.

biserratum Pr. Tent. 114 t. 4 f. 2. 1836 — Brasilia. Asplenium Pr. Del. Prag. **1**. 177. 1822; Athyrium Milde 1870.

Blanchetii Mett. Linn. **36**. 104. 1869 (syn.) — Brasilia. Asplenum Mett.; Kuhn, Linn. **36**. 104. 1869. (HB. 233).

Bommeri Christ in C. Chr. Ind. 228. 1905 — Congo. — Asplenium crenato-serratum Bomm.; Christ, Bull. Boiss. **4**. 662. 1896 (non Bl. 1828).

?Boryanum Pr. Tent 113. 1836 = D. cordifolium.

brachypodum (Bak) C. Chr. Ind. 228. 1905 — Luzon. Asplenium Bak. Syn. 490. 1874.

brachysorum »J. Sm.« Mett. Fil. Lips. 68 in obs. 1856 = D. brevisorum J. Sm. 1841.

Brackenridgei Moore, Ind. 324. 1861 — Samoa. Diplazium speciosum Brack. Expl. Exp. **16**. 145. 1854 (non Bl. 1828); Asplenium Graeffei Luerss. 1871.

Brackenridgei Diels in Schum. et Laut. Fl. deut. Südsee 124. 1901 = D. bulbiferum.
brevifolium Kze. Linn. **23**. 250, 309. 1850 = D. grandifolium.
brevipes (Bak.) C. Chr. Ind. 229. 1905 — Madagascar.
Asplenium Bak. Journ. Linn. Soc. **16**. 201. 1877.
brevisorum J. Sm. JoB. **3**. 408. 1841 = Athyrium silvaticum.
brevisorum J. Sm. Cat. Kew ferns 6. 1856 = D. expansum.
brunneo-viride (Jenm.) C. Chr. Ind. 229. 1905 — Jamaica.
Asplenium Jenm. JoB. **1886**. 269.
bulbiferum Brack. Expl. Exp. **16**. 141 t. 18 f. 1. 1854. NPfl. 226 — Luzon. Polynesia. — D. extensum J. Sm. 1841 (pt.; nomen); D. tenerum Pr. 1849 (nomen); Asplenium Brackenridgei Bak. 1867. HB. 234; Diplazium Diels 1901 (non Moore 1861). — (an Allantodia pinnata Blanco 1845?).
callipteris Fée, Gen. 214. 1850-52. Chr. 216. NPfl. 226 — India occ. Peru. Brasilia. — Asplenium Bak. 1867. HB. 231 (non Fée 1850-52).
camarinum (Bak.) C. Chr. Ind. 229. 1905 — South Camarines.
Asplenium Bak. Syn. ed. II. 230. 1874.
Campbellii (Jenm.) C. Chr. Ind. 229. 1905 — Guiana.
Asplenium Jenm. Gard. Chr. n. s. **24**. 7. 1885.
camptocarpon Fée, 8 mém. 84. 1857 — Mexico.
Asplenium Fourn. 1872 (non Fée 1869).
caracasanum (Willd.) Kze.; Moore, Ind. 324. 1861 — Venezuela-Peru.
Asplenium Willd. sp. **5**. 338. 1810. Mett. n. 173; Athyrium Milde 1870; Asplenium Shepherdi γ HB. 233.
caryifolium (Bak.) C. Chr. Ind. 229. 1905 — Columbia.
Asplenium Bak. Ann. Bot. **5**. 310. 1891.
castaneifolium (Bory) Sw. Syn. 91. 1806 — India occ.
Callipteris Bory, Voy. **1**. 282. 1804; Asplenium Hk. 1860.
caudatum J. Sm. JoB. **3**. 408. 1841 = D. cyatheifolium.
Cavalerii Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 114 — China.
celtidifolium Kze. Bot. Zeit. **1845**. 285. Chr. 217. NPfl. 225 — Amer. trop.
Asplenium Mett. 1856; Aspl. n. 200. HB. 232; Athyrium Milde 1870 — Asplenium centripetale Bak. 1874. HB. 490.
ceramicum (Miq.) C. Chr. Ind. 229. 1905 — Ceram.
Asplenium Miq. Ann. Lugd. Bat. **4**. 168. 1868-69.
ceratolepis Christ, Farnkr. 221. 1897. NPfl. 228 — Costa Rica. Guatemala.
Asplenium Christ, Bull. Boiss. **4**. 658. 1896; Gymnogramme Donn. Smith 1897 (var. Atirrensis).
ceylanense Moore, Ind. 119. 1859 = Athyrium.
Ceylanicum Christ, Farnkr. 216. 1897 = D. zeylanicum.
chimborazense (Spruce) Christ, Farnkr. 221. 1867. NPfl. 228 — Ecuador.
Asplenium Spruce; Bak. Syn. 244. 1867.
chinense (Bak.) C. Chr. Ind. 229. 1905 — Shanghai.
Asplenium Bak. Syn. 237. 1867.
chlorophyllum (Bak.) Bedd. Handb. Suppl. 39. 1892 — Formosa. Penang.
Asplenium Bak. JoB **1885**. 104.
chlororachis Kze.; Fée, Gen. 213. 1850-52 = D. caracasanum.
chocoense (Triana) Hier. Engl. Jahrb. **34**. 459. 1904 — Columbia.
Asplenium Triana; Mett. Ann. sc. nat. V. **2**. 231. 1864. (HB. 244).
Christii C. Chr. Ind. 229. 1905 — Malesia. — D. acuminatum Bl. Enum. 193. 1828; Christ, Ann. Jard. Buit. **12**. 220 (non Raddi 1825); ? Asplenium Wall. 1828 (HB. 235)

coarctatum Link, Hort. Berol. **2**. 70. 1833 = ? D. denticulosum.
?**Cominsii** (Bak.) C. Chr. Ind. 230. 1905 -- Ins. Salomonis.
Gymnogramme Bak. Journ. Linn. Soc. **19**. 296. 1882.
comorensis Bojer; Bak. Syn. 240 in obs. 1867 = D. arborescens.
conchatum J. Sm. Ferns br. and for. 223. 1866 = Athyrium costale.
confertum (Bak.) C. Chr. Ind. 230. 1905 — Sumatra.
Asplenium Bak. Ann. Bot. **8**. 125. 1894.
congruum Brack. Expl. Exp. **16**. 141 t. 18 f. 2. 1854 (HB. 235) — Fiji. Samoa.
Asplenium Christ 1896.
conterminum Christ, Journ. de Bot. **19**. 67. 1905 — Annam.
Corderoi (Sod.) Diels, Nat. Pfl. 1^4. 228. 1899 — Ecuador.
Asplenium Sod. Rec. 39. 1883; Cr. vasc. quit. 206. 1893.
cordifolium Bl. Enum. 190. 1828. Chr. 221. NPfl. 228 — Malesia. Ins. Philipp. (Polynesia. Austr. trop. Africa trop. Madagascar, t. Diels).
Asplenium Mett. 1856; Aspl. n. 186. HB. 243; Anisogonium Bedd. 1870; Oxygonium J. Sm. 1875 — Diplazium integrifolium Bl. 1828; Anisogonium Pr. 1836; Asplenium Mett. 1856; Oxygonium Moore 1857; Athyrium Milde 1870. Callipteris ovata J. Sm. 1841; Anisogonium Fée 1844 (non Asplenium Wall 1828).
cordovense (Bak.) C. Chr. Ind. 230. 1905 — Mexico.
Asplenium Bak. Ann. Bot. **8**. 125. 1894.
coriaceum Carr. in Seem. Fl. Vit. 357. 1873 = D. melanocaulon.
costale (Sw.) Pr. Tent. 114. 1836. Chr. 219. NPfl. 226 — India occ.-Peru.
Asplenium Sw. Schrad. Journ. **1800**^{2}. 55. 1801. HB. 236; Athyrium Milde 1870; Diplazium macrophyllum Desv. 1827; Asplenium Desvauxii Mett. n. 205. 1859 — Diplazium Apollinaris L'Herm.; Fée 1866.
costaricanum C. Chr. Ind. 230. 1905 — Costa Rica.
Asplenium induratum Christ, Bull. Soc. Bot. Belg. **35**. 201. 1896 (non Hk. 1860); Diplazium Diels 1899. NPfl. 226.
crenato-lobatum (Bak.) C. Chr. Ind. 230. 1905 — Ins Hawaii.
Asplenium Bak. Ann. Bot. **5**. 309. 1891.
crenato-serratum (Bl.) Moore, Ind. 121, 325. 1859. Chr. 217. NPfl. 225 — Malesia.
Asplenium Bl. Enum. 177. 1828. Mett. n. 197; Athyrium Milde 1870; Asplenium glaberrimum Mett. 1856; Diplazium Moore 1859 — Asplenium porrectum Wall. 1828. Mett. n. 196. HB. 230; Diplazium Pr. 1836.
crenatum Poir. Enc. Suppl. **2**. 488. 1811 = Asplenium cuneatum.
crenatum (Roxb.) Moore, Ind. 121, 325. 1859 — Malesia.
Asplenium Roxb. Calc. Journ. **4**. 498. 1844.
crenulans Fée; Bak. Syn. 232 in obs. 1867 = ? D. celtidifolium.
crenulatum Liebm. Vid. Selsk. Skr. V. **1**. 254 (seors. 103). 1849 = D. striatum.
crinitum (Bak.) C. Chr. Ind. 230. 1905 — Borneo.
Asplenium Bak. Journ. Linn. Soc. **24**. 258. 1887.
cubense Moore, Ind. 325. 1861 = D. semihastatum.
cultratum Pr. Epim. 84. 1849 — Luzon.
Asplenium Hk. 1860. HB. 231; Athyrium Milde 1870.
cultrifolium (L.) (Kze. Bot. Zeit. **1845**. 286) Moore, Ind. 326. 1861 — Amer. trop.
Asplenium L. sp. **2**. 1081. 1753. HB. 203.
Cumingii (Pr.) C. Chr. Ind. 230. 1905 — Luzon. Celebes.
Ochlogramma Pr. Epim. 94. 1849; Asplenium Mett. 1856; Aspl. n. 185; Athyrium Milde 1870; Callipteris alismaefolia J. Sm. 1841 (nomen, non

Diplazium Pr. 1825); Oxygonium J. Sm. 1841; Asplenium Hk. 1860. HB. 243; Anisogonium Christ 1895; Pteriglyphis elegans Fée 1850-52.
curvatum Desv. Prod. 280. 1827 = D. arboreum.
cyatheifolium (Rich.) Pr. Epim. 84. 1849. NPfl. 226 — Luzon. Polynesia. Asplenium Rich. Sert. Astrol. XIX. 1834. Mett. n. 223. HB. 238; Athyrium Milde 1870. Diplazium caudatum J. Sm. 1841.
decipiens (Kze. Bot. Zeit. **1848**. 193) Moore, Ind. 123. 1859 = D. subserratum.
decurrens Bedd. Ferns S. Ind. t. 229. 1863 65 = D. polypodioides var.
decussatum J. Sm. Bot. Mag. **72** Comp. 28. 1846 = D. Petersenii.
decussatum Bedd. Ferns br. Ind. t. 292. 1868 = D. japonicum.
decussatum Christ, Farnkr. 221. 1897 = D. proliferum.
deflexum J. Sm. JoB. **3**. 407. 1841 = D. tomentosum.
delicatulum Fée, Cr. vasc. Br. **2**. 49. 1872-73 = D. herbaceum.
deltoideum Pr. Tent. 114. 1836 — Luzon. — Asplenium Pr. Rel. Haenk. **1**. 47 t. 7 f. 2. 1825. HB. 237; Athyrium Milde 1870.
denticulosum Gaud. Freyc. Voy. Bot. 322. 1827 = D. Shepherdii.
denticulosum (Desv.) C. Chr. Ind. 231. 1905 — Amer. trop. Asplenium Desv. Berl. Mag. **5**. 323. 1811. Diplazium lonchophyllum Kze. 1839. D. inaequilaterum Liebm. 1849; Asplenium Mett. n. 175. 1859; A. Shepherdi β HB. 233. Diplazium acutale Fée 1850-52 — Asplenium Schiedei Mett. n. 174. 1859.
Dietrichianum (Luerss.) C. Chr. Ind. 231. 1905 — Queensland. Asplenium Luerss. Journ. Mus. Godeffr. **3**. 16. 1873 — (an D. esculentum* ?).
dilatatum Bl. Enum. 194. 1828 = D. maximum.
dissimile Fée, Cr. vasc. Br. **1**. 76 t. 21 f. 1. 1869 = D. Riedelianum.
distentum Moore, Ind. 327. 1861 = D. expansum.
diversifolium J. Sm. Ferns br. and for. 222. 1866 = D. maximum.
diversifrons Bory, Bél. Voy. Bot. **2**. 39. 1833 — India.
divisissimum (Bak.) Christ, Farnkr. 220. 1897. NPfl. 228 — Ecuador. Asplenium Bak. Syn. 242. 1867.
Döderleinii (Luerss.) Mak. Bot. Mag. Tokyo **13**. 14. 1899 — Liu Kiu. Asplenium Luerss. Engl. Jahrb. **4**. 355. 1883.
dubium Link, Hort. Berol. **2**. 70. 1833 = ? D. Shepherdii.
ebenum J. Sm. JoB. **3**. 408. 1841 = ? D. asperum.
elatum Fée, Gen. 214. 1850-52 = D. silvaticum var.
Eggersii (Sod.) C. Chr. Ind. 231. 1905 — Ecuador. Asplenium Sod. Cr. vasc. quit. 188. 1893.
elegans Hk. Ic. pl. t. 939-940. 1854 = D. fraxinifolium.
elegans Linden, Cat. 1859 — Columbia.
elongatum Fée, Gen. 215. 1850-52 — Mexico. — Asplenium Mett.; Salom. 1883.
epirachis Christ, Bull. Soc. Fr. **52** Mém. I. 51. 1905 — China.
esculentum (Retz.) Sw. Schrad. Journ. **1801**[2]. 312. 1803. Chr. 222. NPfl. 228. — Asia, Polynesia trop. — Hemionitis Retz. Obs. **6**. 38. 1791; Asplenium Pr. 1825. Mett. n. 192. HB. 244; Anisogonium Pr. 1836; Microstegia Pr. 1849; Digrammaria Fée 1850-52; Callipteris J. Sm.; Moore et Houlst. 1851. Asplenium ambiguum Sw. 1801: Digrammaria Hk. 1840; Microstegia Pr. 1849; Callipteris Moore 1857; Diplazium Hk. 1857 (non alior.); Athyrium Milde 1870 — Diplazium serampurense Spr. 1821; Anisogonium Pr. 1836; Callipteris Fée 1850-52 — Diplazium malabaricum Spr. 1827; Callipteris J. Sm. 1841; Asplenium Mett. 1856 — Diplazium pubescens Link 1833;

Microstegia Pr. 1849; Asplenium Mett. 1856 — Diplazium umbrosum Moritz 1845-46 (non Willd. 1810); Asplenium Mett. 1856; A. Moritzii Mett. 1856 — A. vitiense Bak. 1867. HB. 245; Diplazium Carr. 1873 — Gymnogramme edulis Ces. 1876.

expansum Willd. sp. **5**. 354. 1810 — Amer. trop.
Asplenium Pr. 1825. Mett. n. 221; Athyrium Milde 1870. Diplazium obtusum Klf.: Link 1833; Asplenium Mett. 1856; Aspl. n. 217 (non Desv. 1827) — Diplazium Haenkeanum Pr. 1836 — ? Asplenium distentum A.Br. 1860; Diplazium Moore 1861; Athyrium Milde 1870; Diplazium athyrioides Moore 1857 — Asplenium radicans pt. HB. 241 (vix Schkuhr 1809); Diplazium pt. Chr. 220. NPfl. 228.

extensum J. Sm. JoB. **3**. 407. 1841 = D. bulbiferum, cultratum, maximum, petiolare, vestitum.

falcatum Don, Prod. Fl. Nepal. 13. 1825 = Coniogramme fraxinea.

falcatum Liebm. Vid. Selsk. Skr. V. **1**. 253 (seors. 101). 1849 — Mexico. — [Hemitelia sp.?).

falcatum Brack. Expl. Exp. **16**. 143. 1854 = D. harpeodes.

Fauriei Christ, Bull. Boiss, II. **1**. 1015. 1901 — Liu Kiu.

Feei Schaffn.; Fée, 8 mém. 85. 1857 — Mexico.
Asplenium Schaffneri Fourn. 1872.

Fenzlianum (Luerss.) C. Chr. Ind. 232. 1905 — Ins. Hawaii.
Asplenium Luerss. Flora **1875**. 434.

firmum Fée, 10 mém. 30 t. 38 f. 2. 1865 — Ceylon.

flavescens (Mett.) Christ, Farnkr. 217. 1897. NPfl. 226 — India occ. Mexico-Peru.
Asplenium Mett. 1864. HB. 232; A. juglandifolium Hk. Fil. exot. t. 100. 1859 (non Lam. 1786); Diplazium J. Sm. 1866 (non Sw. 1806); Athyrium Milde 1870.

flexuosum Pr. Tent. 114. 1836 = D. Preslianum.

Forbesii (Bak.) C. Chr. Ind. 232. 1905 — Java.
Asplenium Bak. Ann. Bot. **5**. 312. 1891.

Franconis Liebm. Vid. Selsk. Skr. V. **1**. 256 (seors. 104). 1849. Chr. 218. NPfl. 226 — Mexico-Ecuador. — Asplenium Mett. n. 166. 1859. HB. 236 — A. aspidiiforme Fée 1850-52.

fraternum Pr. Tent. 114. 1836 = ? D. denticulosum.

fraxineum Don, Prod. Fl. Nepal. 12. 1825 = Coniogramme.

fraxinifolium Pr. Rel. Haenk. **1**. 49. 1825 — Malesia-Japonia.
Anisogonium Pr. 1836; Callipteris J. Sm.; Moore 1861; Athyrium Milde 1870. Diplazium luzoniense Spr. 1827. D. lineolatum Bl. 1828. NPfl. 228; Asplenium Mett. 1866. HB. 243; Anisogonium Bedd. 1869 — Callipteris elegans J. Sm. 1841; Oxygonium J. Sm. 1841; Anisogonium Pr. 1849; Diplazium Hk. 1854; Asplenium Mett. 1856; Aspl. n. 188. A. alternifolium Mett. 1856; Aspl. n. 198 (non 1866).

fraxinifolium Lodd.; Pr. Tent. 113. 1836 = D. grandifolium.

frondosum J. Sm. JoB. **3**. 408. 1841 = D. polypodioides.

fuscopubescens Moore, Ind. 329. 1861 = D. sandwichianum.

fuscum (Bak.) C. Chr. Ind. 232. 1905 — Ecuador. Costa Rica.
Asplenium Bak. 1891; A. fuliginosum Sod. Rec. 40. 1883 (non Hk. 1860); A. ochraceum Sod. Cr. vasc. quit. 207. 1893; Diplazium Diels 1899. NPfl. 228.

Gaudichaudii (Mett.) C. Chr. Ind. 232. 1905 — Ins. Rawak.
Asplenum Mett.; Kuhn, Linn. **36**. 103. 1869. HB. 491.

giganteum Karst.; Kze. Linn. **23**. 251. 1850 (nomen) = ? D. marginatum.

glaberrimum Moore, Ind. 135. 1859 = D. crenato-serratum.
Godmani (Bak.) C. Chr. Ind. 233. 1905 — St. Vincent. Grenada.
Asplenium Bak. Ann. Bot. **5**. 166 t. 11. 1891.
gracilescens Moore, Ind. 329. 1861 (nomen) — Columbia.
Asplenium Mett. Ann. sc. nat. V. **2**. 237. 1864. HB. 242.
grammitis (»Wall.« Mett.) Moore, Ind. 329. 1861 — Nepal.
Asplenium »Wall.« Mett. Aspl. 181. n. 206. 1859.
grammitoides Pr. Epim. 84. 1849 — Malesia. Ins. Philipp.
Asplenium Hk. 1854. Mett. n. 209 c. HB. 491; Athyrium Milde 1870.
grammitoides Fée, 11 mém. 43 t. 11. 1866 = D. striatum var.
grandifolium Sw. Schrad. Journ. **1800^2**. 62. 1801. Chr. 217. NPfl. 225 — Amer. trop. — Asplenium Sw. Prod. 130. 1788. Mett. n. 201. HB. 231; Hemionitis Sw. 1806; Athyrium Milde 1870.
Griffithii Moore, Ind. 330. 1861. NPfl. 227 — India bor.
Asplenium Bak. 1867. HB. 239; Athyrium Milde 1870; Bedd. 1876.
Haenkeanum Pr. Tent. 114. 1836 = D. expansum.
Hahnii (Fourn.) C. Chr. Ind. 233. 1905 — Mexico.
Asplenium Fourn. Mex. pl. **1**. 109. 1872.
harpeodes Moore, Ind. 330. 1861 — Samoa. — D. falcatum Brack. Expl. Exp. **16**. 143. 1854 (non Don 1825 nec Liebm. 1849).
Harrisonii (Bak.) Christ, Prim. Fl. costaric. **3**. 27. 1901 — Amer. centr.
Asplenium Bak. JoB. **1884**. 362.
hastile (Christ) C. Chr. Ind. 233. 1905 — Cuba. Haïti.
Asplenium Christ, Engl. Jahrb. **24**. 108. 1897.
hemionitideum (Bak.) C. Chr. Ind. 233. 1905 — Ecuador.
Asplenium Bak. JoB. **1877**. 163.
hemionitideum Christ, Bull. Boiss. **7**. 12. 1899 = D. rude.
herbaceum Fée, Cr. vasc. Br. **1**. 80 t. 23 f. 1. 1869 — Brasilia.
Asplenium Bak. 1874. HB. 492 (non Fée 1857); A. Glaziovii Bak. 1870 — Diplazium delicatulum Fée 1872-73.
heterophlebium (Mett.) Diels, Nat. Pfl. **1^4**. 228. 1899 — Himalaya.
Asplenium Mett.; Bak. Syn. 243. 1867; Anisogonium Bedd. 1869.
heterophyllum Bl. Enum. 190. 1828 = Angiopteris sp.
hians Kze.; Kl. Linn. **20**. 361. 1847. NPfl. 228 — India occ. Costa Rica. Ecuador.
Asplenium Mett. n. 219. 1859. HB. 241; Athyrium Milde 1870 — Asplenium *bogotense* Karst. 1864-69.
Hilsenbergianum Pr. Epim. 84. 1849 = D. silvaticum.
hirtipes Christ, Bull. Boiss. **7**. 11. 1899 — Yunnan.
horridum Kze. Anal. 26. 1837 = D. proliferum.
Hosei Christ, Bull. Soc. Bot. Belg. **33^2**. 93. 1894 — Borneo.
humile Boj. Hort. Maur. 398. 1837 = D. silvaticum.
Huttoni (Bak.) C. Chr. Ind. 233. 1905 — »Malay Isles«.
Asplenium Bak. Ann. Bot. **5**. 311. 1891.
hylophilum (Hier.) C. Chr. Ind. 233. 1905 — Africa orient. trop.
Asplenium Hier. in Engl. Pflw. Ostafr. C. 84. 1895.
inaequilaterum Liebm. Vid. Selsk. Skr. V. **1**. 255 (seors. 103). 1849 = D. denticulosum.
incisum Schum. Vid. Selsk. Afh. **4**. 232 (seors. 458). 1827 = D. proliferum.
induratum Diels, Nat. Pfl. **1^4**. 226. 1899 = D. costaricanum.
ingens Christ, Bull. Boiss. II. **4**. 970. 1904 — Costa Rica. Ecuador.

integrifolium Bl. Enum. 190. 1828 = D. cordifolium.

intercalatum Christ in Schwacke, Pl Nov. Mineiras **2**. 29. 1900; Bull. Boiss. II. **2**. 558. 1902 — Brasilia.

isobasis Christ, Bull. Boiss. II. **4**. 618. 1904 — Japonia.

japonicum (Thbg.) Bedd. Ferns br. Ind. Suppl. 12. 1876. Chr. 218. NPfl 226 . — Japonia-China-Asia trop. — Asplenium Thbg. Fl. Jap. 334. 1784. HB. 234. A. ambiguum Schkuhr 1809 (non Sw. 1801); Diplazium Schkuhrii J. Sm. 1841; Asplenium Hk. 1860; Mett. 1866 (non 1859) — Diplazium Oldhami Christ 1899 — D. decussatum Bedd. 1868 (non J. Sm. 1846); Asplenium polyrhizon Bak. 1874. HB. 490 — A. Conilii Franch. et Sav. 1876. Conf. D. Petersenii et D. Thwaitesii, f. Bedd. formae D. japonici.

Jerdoni Bedd. Ferns br. Ind. t. 327. 1869 = Athyrium spectabile.

juglandifolium Sw. Syn. 91, 282. 1806 = Asplenium.

juglandifolium J. Sm. Ferns br. and for. 222. 1866 = D. flavescens.

Katzeri Regel, Gartenfl. **6**. 35 t. 282. 1860 = D. silvaticum.

Klotzschii (Mett.) Moore, Ind. LV. 1857 — Columbia-Venezuela. Jamaica. Asplenium Mett. 1856; Aspl. n. 218. HB. 241; Athyrium Milde 1870; Lotzea diplazioides Kl. et Karst. Linn. **20**. 358. 1847 (non Asplenium Hk. et Arn. 1832) — Asplenium Hartianum Jenm. 1886.

Laffanianum (Bak.) C. Chr. Ind. 234. 1905 — Bermuda. Asplenium Bak. Gard. Chr. n. s. **17**. 673. 1882.

lanceum (Thbg.) Pr. Tent. 113. 1836. Chr. 215. NPfl. 225 — Japonia. China-India-Ceylon. — Asplenium Thbg. Fl. Jap. 333. 1784. Mett. n. 164. HB. 229: Athyrium Milde 1870; Micropodium J. Sm. 1875; Scolopendrium dubium Don 1825; Asplenium subsinuatum Wall.; Hk. et Grev. 1827.

lanceum Bory, Bél. Voy. Bot. **2**. 37 t. 4. 1833 = D. cordifolium.

larutense Bedd. Handb. Suppl. 38. 1892 — Perak.

Lasiopteris Kze. Linn. **17**. 568. 1843 = D. Petersenii.

latifolium Moore, Ind. 141. 1859 = D. maximum.

Lechleri (Mett.) Moore, Ind. 141, 331. 1859. Chr. 216. NPfl. 225 — Amer. contin. trop. — Asplenium Mett. Fil. Lechl. **1**. 16 t. 2 f. 10. 1856; Aspl. n. 184. HB. 231; Athyrium Milde 1870. Diplazium parallelogrammum Fée 1869.

Lehmannii Hier. Engl. Jahrb. **34**. 458. 1904 — Columbia.

lepidorachis (Bak.) C. Chr. Ind. 234. 1905 — Tonkin. Asplenium Bak. JoB. **1890**. 264.

leptocarpon Fée, Cr. vasc. Br. **1**. 80 t. 23 f. 2. 1869 = D. ambiguum.

leptochlamys Fée, Cr. vasc. Br. **1**. 79 t 22 f. 1. 1869 = D. ambiguum.

leptochlamys Diels, Nat. Pfl. 1^4. 228. 1899 = D. leptogrammoides.

leptogrammoides C. Chr. Ind. 234. 1905 — Ecuador. Asplenium leptochlamys Sod. Cr. vasc. quit. 200. 1893; Diplazium Diels 1899. NPfl. 228 (non Fée 1869).

leptophyllum Bak.; Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 245 (nomen) — (? China).

Lindbergii (Mett.) Christ, Prim. Fl. costar. **3**. 27. 1901 — Mexico-Brasilia. Asplenium Mett. Ann. sc. nat. V. **2**. 236. 1864. HB. 236. A. subnudum Karst. (1865).

lineatum Pr. Tent. 113. 1836 = D. crenato-serratum.

lineolatum Bl. Enum. 191. 1828 = D. fraxinifolium.

Lobbianum (Hk.) Moore, Ind. 331. 1861 — Java. Ins. Philipp. Asplenium Hk. 2 Cent. t. 17. 1860.

lobulosum (Wall.) Pr. Tent. 114. 1836 — Himalaya. Asplenium Wall. 1828 (nomen); Mett. Aspl. 163. n. 170. 1859; A. longi-

folium Don, Prod. Fl. Nepal. 7. 1825. HB. 234 (non Schrad. 1824): Diplazium Moore 1859. NPfl. 226; Athyrium Milde 1870.
Loddigesii J. Sm.; Diels, Nat. Pfl. 1⁴. 226. 1899 = D. nigropaleaceum.
lonchophyllum Kze. Linn. **13**. 141. 1839 = D. denticulosum.
lonchophyllum Kze. Bot. Zeit. **1848**. 192 = D. subserratum.
longifolium Moore, Ind. 141, 332. 1859 = D. lobulosum.
longipes Fée, Cr. vasc. Br. **1**. 77 t. 21 f. 2. 1869 = { D. celtidifolium (f. Bak.). D. grandifolium (f. Christ).
longisorum (Bak.) C. Chr. Ind. 235. 1905 — Columbia.
Asplenium Bak. JoB. **1881**. 204.
lucidum Poir. Enc. Suppl. **2**. 487. 1811 = D. nitidum.
luzoniense Spr. Syst. **4**. 68. 1827 = D. fraxinifolium.
macrodictyon (Bak.) Diels, Nat. Pfl. 1⁴. 228. 1899 — Ecuador. Costa Rica.
Asplenium Bak. JoB. **1877**. 163.
macrophyllum Desv. Prod. 280. 1867 = D. costale.
macropterum (Sod.) C. Chr. Ind. 235. 1905 — Ecuador.
Asplenium Sod. Cr. vasc. quit. 637. 1893.
macrotis (Bak.) Christ, Prim. Fl. costar. **3**. 28. 1901 — Costa Rica.
Asplenium Bak. JoB. **1884**. 362.
malabaricum Spr. Nova Acta **10**. 231 t. 17 f. 1. 1821 = D. esculentum.
malaccense Pr. Epim. 86. 1849 = D. maximum.
manilense (Spr.) C. Chr. Ind. 235. 1905 — Ins. Philipp.
Asplenium Spr. Syst. **4**. 89. 1827; A. serrulatum Pr. Rel. Haenk. **1**. 46 1825 (non Cav. 1801); Anisogonium Pr. 1836; Microstegia Pr. 1849; Callipteris Fée 1850-52; Athyrium Milde 1870.
marginale (Hill.) C. Chr. Ind. 235. 1905 — Ins. Hawaii.
Asplenium Hill. Fl. Hawaii. 613. 1888.
marginatum (L.) Diels, Nat. Pfl. 1⁴. 229. 1899 — Amer. trop.
Asplenium L. sp. **2**. 1082. 1753. Mett. n. 183. HB. 246; Hemidictyum Pr. 1836. Chr. 214; Athyrium Milde 1870. Asplenium limbatum Willd. 1810; Hemidictyum Pr. 1849. Asplenium Mikani Pr. 1822.
marginatum Bl. Enum. 195. 1828 = ? D. polypodioides.
maximum (Don) C. Chr. Ind. 235. 1905 — China. Asia, Austr., Polynesia trop.
Asplenium Don, Prod. Fl. Nepal. 8. 1825. HB. 239 (pt.) — A. latifolium Don 1825. HB. 239 (non Cav. 1803); Diplazium Moore 1859. Chr. 220. NPfl. 226; Athyrium Milde 1870 — Diplazium dilatatum Bl. 1828; Microstegia Pr. 1849; Asplenium Hk. 1860; Athyrium Milde 1870 — Asplenium sororium Mett. 1861; Diplazium Carr. 1873 — Gymnogramme gigantea Bak. 1889; Nephrodium Diels 1900.
megaphyllum (Bak.) Christ, Bull. Boiss. **6**. 961. 1898 — Tonkin. China.
Asplenium Bak. JoB. **1890**. 264.
melanocaulon Brack. Expl. Exp. **16**. 144. 1854 — Fiji. Aneiteum.
Asplenium Bak. 1867. HB. 240 (non Willd. 1809); Athyrium Milde 1870; Diplazium coriaceum Carr. 1873. NPfl. 226.
melanochlamys (Hk.) Moore, Ind. 332. 1861. NPfl. 227 — Ins. Lord Howe.
Asplenium Hk. sp. **3**. 259. 1860. HB. 239.
melanopodium Fée, 8 mém. 85. 1857 = D. deltoideum.
melanopus (Sod.) Hier. Engl. Jahrb. **34**. 456. 1904 — Ecuador. Columbia.
Asplenium Sod. Cr. vasc. quit. 189. 1893.

membranaceum Mett. Linn. **36**. 103. 1869 (syn.) — Ins. Polynesiae.
Asplenum Mett.; Kuhn, Linn. **36**. 103. 1869. HB. 491.
meniscioides (Sod.) Diels. Nat. Pfl. **1**[4]. 225. 1899 — Ecuador.
Asplenium Sod. Cr. vasc. quit. 188. 1893.
Mettenianum (Miq.) C. Chr. Ind. 236. 1905 — Japonia. China.
Asplenium Miq. Ann. Lugd. Bat. **3**. 174. 1867. A. Textori Miq. 1870
Diplazium Mak. 1899.
Meyenianum Pr. Tent 108. 1836; Epim. 88. 1849 — Ins. Philipp.
Asplenium Mett. n. 202. 1859. HB. 238; Athyrium Milde 1870.
microphyllum Desv. Prod. 281 t. 7 f. 1. 1827 — India. Malesia.
Microstegia Pr. 1849; Callipteris Fée 1850-52; Asplenium Mett. 1866.
Mildei (Kuhn) C. Chr. Ind. 236. 1905 — Ecuador.
Asplenum Kuhn, Linn. **36**. 95. 1869. HB. 491.
Moccennianum (Sod.) C. Chr. Ind. 236. 1905 — Ecuador.
Asplenium Sod. Rec. 34. 1883; Cr. vasc. quit. 194. 1893.
mohillense Fée, 8 mém. 85. 1857 — Madagascar. — Asplenum Kuhn 1868.
monticolum (Jenm.) C. Chr. Ind. 236. 1905 — Jamaica.
Asplenium Jenm. JoB. **1882**. 326.
multiflorum (Roxb.) Moore, Ind. 147, 333. 1859 — Malesia.
Asplenium Roxb. Calc. Journ. **4**. 499. 1844.
mutilum Kze. Flora **1839**[1]. Beibl. 37 — Amer. trop.
Asplenium Mett. n. 204. 1859. (HB. 233); Athyrium Milde 1870.
Naganumanum Mak. Bot. Mag. Tokyo **13**. 14. 1899 — Japonia.
Asplenium Mak. Bot. Mag. Tokyo **6**. 46. 1892 (nomen).
neglectum (Karst.) C. Chr. Ind. 236. 1905 — Columbia. Mexico.
Asplenium Karst. Fl. Col. **1**. 106 t. 52. 1860. HB. 490.
nervosum (Mett.) Diels, Nat. Pfl. **1**[4]. 226. 1899 — Columbia.
Asplenium Mett. Ann. sc. nat. V. **2**. 235. 1864. HB. 237.
nicotianifolium (Mett.) C. Chr. Ind. 236. 1905 — Columbia.
Asplenium Mett. Ann. sc. nat. V. **2**. 235. 1864. HB. 490.
nigropaleaceum Kze. Linn. **24**. 270. 1851 — St. Helena. — Asplenium A.Br. 1861. HB. 240; Diplazium Loddigesii J. Sm.; Diels 1899. NPfl. 226.
nitidum Cav. Anal. Cienc. (fasc.?) **19**. 67 t. 48 —; Sw. Syn. 92, 283. 1806 — Ins. Mariannae.
obtusum Desv. Prod. 281. 1827; Hier. Engl. Jahrb. **34**. 457. 1904 — Peru-Columbia.
Athyrium Milde 1870.
obtusum Klf.; Link, Hort. Berol. **2**. 73. 1833 = D. expansum.
ochraceum Diels, Nat. Pfl. **1**[4]. 228. 1899 = D. fuscum.
Oldhami Christ, Bull. Boiss. **7**. 819. 1899 = D. japonicum var.
Otites Moore et Houlst. Gard. Mag. Bot. **3**. 231. 1851 = D. arboreum.
otophorum (Miq.) C. Chr. Ind. 236. 1905 — Japonia.
Asplenium Miq. Ann. Lugd. Bat. **3**. 175. 1867.
Ottonis Kl. Linn. **20**. 360. 1847 — Amer. trop. Africa occ. trop.
Asplenium Hk. 1860. (HB. 233); Athyrium Milde 1870.
pallidum (Bl.) Moore, Ind. 333. 1861 — Malesia. Queensland.
Asplenium Bl. Enum. 177. 1828. Mett. n. 195. HB. 230; Athyrium Milde 1870.
Asplenium calophyllum J. Sm. 1841. Mett. n. 194 — A. Prenticei Bail. 1880.
paradoxum Fée, Gen. 214. 1850 52 — Ceylon. — Callipteris Moore 1861.
parallelogrammum Fée, Cr. vasc. Br. **1**. 76 t. 20 f. 2. 1869 = D. Lechleri.
patens Pr. Tent. 114. 1836 = Asplenium.

pedatum Kl. Linn. **20**. 360. 1847 — Columbia.
Athyrium Milde 1870; Asplenium Salom. 1883.
Petersenii (Kze.) Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 245 — China. India. Java
Asplenium Kze. Anal. 24. 1837. Diplazium Lasiopteris Kze. 1843. Chr. 218. NPfl. 226; Asplenium Mett. 1856. HB. 235. Diplazium decussatum J. Sm. 1846; Athyrium Milde 1870. Asplenium tomentosum Mett. n. 209. 1859 (non Lam. 1786 nec Diplazium Bl. 1828).
petiolare Pr. Epim. 86. 1849 — Ins. Philipp.
Athyrium Milde 1870; ? Asplenium Ces. 1877.
phanerotis Kze. Bot. Zeit. **1846**. 443 = D. crenato-serratum.
pinnatifido-pinnatum (Hk.) Moore, Ind. 334. 1861 — India.
Asplenium Hk. sp. **3**. 238. 1860. HB. 231.
pinnatifidum Kze. Linn. **9**. 72. 1834; Anal. 25 t. 16 — Peru. Ecuador.
Anisogonium Pr. 1836; Callipteris Fée 1850-52; Athyrium Milde 1870; Asplenium Kunzei Mett. 1856; Aspl. n. 187. HB. 243.
pinnatifidum Fée, 10 mém. 29 t. 35 f. 3. 1865 = D. zeylanicum.
plantagineum Sw. Schrad. Journ. **1800**². 62. 1801 = D. plantaginifolium.
plantaginifolium (L.) Urban, Symb. Antill. **4**. 31. 1903 — Amer. trop.
Asplenium L. Syst. Nat. ed. X. **2**. 1323. 1759; A. plantagineum L. 1763. Mett. n. 166. HB. 230; Hemionitis Sm. 1793; Diplazium Sw. 1801. Chr. 215. NPfl. 229; Athyrium Milde 1870. Diplazium acuminatum Raddi 1825.
polyanthes (Sol.) C. Chr. Ind. 237. 1905 — Otaheite.
Asplenium Sol.; Forst. 1786 (nomen); Bak. Syn. 492. 1874.
polymorphum Pr. Tent. 114. 1836 = D. polypodioides.
polypodioides Bl. Enum. 194. 1828. Chr. 219. NPfl. 227 — India bor.-Malesia-Austral. trop. — Microstegia Pr. 1849; Asplenium Mett. 1856; Aspl. n. 225 HB. 238; Athyrium Milde 1870 — Diplazium decurrens Bedd. 1863-65 — Asplenium diplazioides Bory 1833 (non Hk. et Arn. 1832); A. Blumei Bergsm. 1857. Mett. n. 224.
polypodioides Bedd. Ferns S. Ind. t. 163. 1863-65 = D. asperum.
popayanense Hier. Engl. Jahrb. **34**. 457. 1904 — Columbia.
porphyrorachis (Bak.) Diels. Nat. Pfl. 1⁴. 225. 1899 — Borneo. Celebes.
Asplenium Bak. JoB. **1879**. 40; Ic. pl. t. 1650. Polypodium subserratum Hk. 1862 (non Diplazium Moore 1861). Asplenium lomariaceum Christ 1895.
porrectum Pr. Tent. 113. 1836 = D. crenato-serratum.
praelongum Fée, 10 mém. 30 t. 37 f. 1. 1865 = D. silvaticum.
Prescottianum (Wall.) Moore, Ind. 156. 334. 1859 — Malesia.
Asplenium Wall. 1828 (nomen); Hk. sp. **3**. 251. 1860. (HB. 233).
Preslianum C. Chr. Ind. 237. 1905 — Peru.
Asplenium flexuosum Pr. Rel. Haenk. **1**. 46 t. 7 f. 1. 1825. Mett. n. 226. HB. 241 (non Schrad. 1818); Diplazium Pr. 1836; Athyrium Milde 1870.
prionophyllum Kze. Bot. Zeit. **1848**. 194 — Java. — Microstegia Pr. 1849.
proliferoides Bory, Bél. Voy. Bot. **2**. 38. 1833 = D. proliferum.
proliferum (Lam.) Thouars. Fl. Trist. d'Ac. 35. 1804; Klf. Enum. 182. 1824 — Malesia-Polynesia. Austral. trop. Africa trop. Madagascar. Mascarenae.
Asplenium Lam. Enc. **2**. 307. 1786; Callipteris Bory 1804; Athyrium Milde 1870; Asplenium decussatum Sw. 1801. Mett. n. 190. HB. 243; Anisogonium Pr. 1836; Diplazium Chr. 1897. 221. NPfl. 228 — D. Swartzii Bl. 1828; Callipteris Pr. 1849; Asplenium Mett. 1856 — Diplazium accedens Bl. 1828; Callipteris J. Sm. 1841; Athyrium Milde 1870 —

Callipteris spinulosa J. Sm. 1841; Asplenium Mett. 1859. ? A. Mettenii Fourn. 1873 (non Kuhn 1868) — Digrammaria robusta Fée 1850-52.
proliferum Brack. Expl. Exp. **16**. 140. 1854 = D. silvaticum.
pseudo-porrectum Hier. Engl. Jahrb. **28**. 342. 1900 — Africa trop.
psilopteron Kl. msc. — Ceylon. — Asplenium Salom. Nom. 99. 1883 (nomen).
pubescens Link, Hort. Berol. **2**. 72. 1833 = D. esculentum.
pubescens Lowe, Ferns **5** t. 52. 1858 = Athyrium decurtatum.
pubigerum J. Sm. Cat. Kew Ferns 1856 — Hort. — Asplenium Mett; Salom. 1883.
pulcherrimum Raddi, Opusc. sci. Bol. **3**. 292 t. 12 f. 2. 1819 = Didymochlaena truncatula.
pulicosum (Hk.) Moore, Ind. 335. 1861. NPfl. 228 — Ecuador.
Asplenium Hk. sp. **3**. 262 1860. HB. 240.
Pullingeri (Bak.) J. Sm. Ferns br. and for. ed. II. 315. 1877 — Hongkong.
Asplenium Bak. Gard. Chr. n. s. **4**. 484. 1875.
radicans Desv. Prod. 281. 1827 = ? D. arboreum.
radicans Moore, Ind. 335. 1861 = D. Shepherdii et D. denticulosum.
radicans Christ, Farnkr. 220. 1897 = D. ambiguum et D. expansum.
remotum Fée, Cr. vasc. Br. **1**. 81 t. 24 f. 1. 1869 = D. ambiguum.
repandum Bl. Enum. 191. 1828 = D. proliferum.
repandum Fée, Gen. 215. 1850-52 — Cuba.
repens Raddi, Opusc. sci. Bol. **3**. 292. 1819 = D. plantaginifolium.
rhoifolium Kze. msc. (Mett. Aspl. 178. syn. 1859) — Columbia-Ecuador.
Asplenium Mett. Aspl. 178. n. 199. 1859. HB. 232.
Riedelianum Kze. Linn. **36**. 103. 1869. (syn.) — Brasilia. Amer. centr.
Asplenum Bong.; Kuhn, Linn. **36**. 102. 1869. HB. 489; Diplazium dissimile Fée 1869. Asplenium verapax Donn. Smith 1888.
rigescens Kze. Bot. Zeit. **1845**. 283 = D. arboreum.
rivale (Spruce) Diels, Nat. Pfl. **1**[4]. 228. 1899 — Ecuador.
Asplenium Spruce; Bak. Syn. 244. 1867.
Roemerianum (Kze.) Pr. Tent. 113 (t. 4 f. 5). 1836; Kl. Linn. **20**. 359. 1847 — Cuba-Peru. — Asplenium Kze. Linn. **9**. 62. 1834. Mett. n. 167; Athyrium Milde 1870. Asplenium caucense Karst. 1861 — A. crassidens Fée 1857.
rostratum Fée, Cr. vasc. Br. **1**. 81 t. 24 f. 2. 1869 = D. ambiguum.
Roxburghii Moore, Ind. 176, 337. 1859 — Amboyna. — Asplenium varium Roxb. Calc. Journ. **4**. 498. 1844 (non Diplazium Gaud. 1827).
rude Christ in C. Chr. Ind. 238. 1905 — Yunnan.
D. hemionitideum Christ, Bull. Boiss. **7**. 12. 1899. NPfl. 228 (non supra).
Sammatii (Kuhn) C. Chr. Ind. 238. 1905 — Africa trop.
Asplenum Kuhn, v. Deck. Reis. **3**[3]. Bot. 34. 1879.
Sandwichense Pr. Tent. 114. 1836; Epim. 85. 1849 = Asplenium.
sandwichianum (Pr.) Diels, Nat. Pfl. **1**[4]. 228. 1899 — Ins. Hawaii. Peru.
Athyrium Pr. 1836 (nomen); Epim. 67. 1849; Asplenium Hk. 1860. HB. 242 — A. alienum Mett. n. 181. 1859. A. fuscopubescens Hk. 1860; Diplazium Moore 1861. (Peru).
Schkuhrii J. Sm. JoB. **3**. 407. 1841 = D. silvaticum.
Schkuhrii Bedd. Ferns S. Ind. t. 230. 1863-65 = D. Beddomei.
Schlimense Fée, 8 mém. 84. 1857 = D. grandifolium.
sechellarum (Bak.) C. Chr. Ind. 238. 1905 — Ins. Sechellae. Madagascar.
Asplenium Bak. Syn. 491. 1874.
Seemannii Moore, Ind. 337. 1862 — Panama. — Asplenium Bak. 1867. HB. 231.

semihastatum (Kze.) C. Chr. Ind. 239. 1905 — Cuba. Jamaica. Puerto Rico.
Asplenium Kze.; Mett. Aspl. 162. n. 168. 1859. HB. 233 — A. hymenodes Mett. 1859. n. 169. A. Cubense Hk. 1860; Diplazium Moore 1861.

serampurense Spr. Nova Acta 10. 231 t. 17 f. 1-2. 1821 = D. esculentum.

serratum Schum. Vid. Selsk. Afh. 4. 223 (seors. 459). 1827 = D. proliferum.

serrulatum Desv. Prod. 282. 1827 = D. arborescens.

serrulatum (Roxb.) Moore, Ind. 167, 337. 1860 — Amboyna.
Asplenium Roxb. Calc. Journ. 4. 498. 1844 (non Cav. 1801 nec Pr. 1825).

setosum Pr. Tent. 113. 1836 — Ins. Philipp.
Asplenium? Pr. Rel. Haenk. 1. 42. 1825 (non Desv. 1811).

Shepherdii (Spr.) Link, Hort. Berol. 2. 70. 1833. Chr. 218. NPfl. 226 (pt.?) — Amer. trop. — Asplenium Spr. Nova Acta 10. 231 t. 17 f. 5-6. 1821. Mett. n. 171. HB. 233 (excl. β et γ). A. ambiguum Raddi 1819 (non Sw. 1801 nec alior.); Diplazium J. Sm. 1857; Asplenium striatum L. hb.: Hk. 1860 — Diplazium anthraxacolepis Fée 1857.

shepherdioides (Bak.) C. Chr. Ind. 239. 1905 — Sumatra.
Asplenium Bak. Ann. Bot. 8. 126. 1894.

Sieberianum Pr. Tent. 113. 1836 = Asplenium salicifolium.

silvaticum (Bory) Sw. Syn. 92. 1806. Chr. 217. NPfl. 225 — Asia trop. Queensland. Polynesia. Mauritius. ? Amer. trop. — Callipteris Bory, Voy. 1. 282. 1804; Asplenium Pr. 1825. Mett. n. 202. HB. 232; Anisogonium Pr. 1836; Microstegia Pr. 1849; Athyrium Milde 1870 — Asplenium Schkuhrii Mett. n. 208. 1859 (non 1866) — Diplazium elatum Fée 1850-52; Asplenium Mett. n. 203. 1859 — Diplazium proliferum Brack. 1854 (non Thouars 1804) — D. acuminatum Pr. 1836 (non Raddi 1825 nec Bl. 1828); Asplenium Mett. Aspl. 181 n. 207. 1859 (vix Wall.). — vide D. Blanchetii, D. mutilum, D. Ottonis, subspecies americanae D. silvatici.

Smithianum Kze. Bot. Zeit. 1848. 194 (nomen) = D. sorzogonense.

Smithianum (Bak.) Diels. Nat. Pfl. 1[4]. 228. 1899 — Ceylon. Celebes. Batjan.
Asplenium Bak. Syn. 245. 1867; Callipteris Bedd. 1870; Anisogonium Bedd. 1876.

Solandri Carr. in Seem. Fl. Vit. 356. 1873. (HB. 491) — Ins. Societatis.

sorbifolium (Willd.) Pr. Tent. 113. 1836 — India.
Asplenium Willd. sp. 5. 312. 1810 (non Jacq. 1788).

sororium Carr. in Seem. Fl Vit. 356. 1873 = D. maximum var.

sorzogonense Pr. Tent. 114. 1836 — Asia trop.
Asplenium Pr. Rel. Haenk. 1. 45. 1825. Mett. n. 212. HB. 236; Hypochlamys Fée 1850-52; Athyrium Milde 1870 — Diplazium speciosum Bl. 1828. Chr. 218. NPfl. 226; Asplenium Mett. n 213. 1859. HB. 235; Athyrium Milde 1870. — Conf. Christ. Ann. Jard. Buit. 12. 217.

speciosum Bl. Enum. 193. 1828 = D. sorzogonense.

speciosum Brack. Expl. Exp. 16. 145. 1854 = D. Brackenridgei.

spinosum Bory, Dup. Voy. Bot. 1. 271. 1828 = D. proliferum.

spinulosum Bl. Enum. 193. 1828 — Moluccae. N. Guinea.
Asplenium Mett. 1866 (non 1859); Athyrium Milde 1870 Diplazium heteropteron Kze. hb; Asplenium Mett. n. 191. 1859.

Sprucei (Bak.) C. Chr. Ind. 239. 1905 — Ecuador.
Asplenium Bak. Syn 234. 1867.

squamigerum (Mett.) Christ. Bull. Soc. Fr. 52 Mém. I. 51. 1905 — Japonia. China. India bor. — Asplenium Mett. Ann. Lugd. Bat. 2. 239. 1866. HB. 237.

stenocarpum (Mett.) C. Chr. Ind. 240. 1905 — Columbia.
Asplenium Mett. Ann. sc. nat. V. **2**. 232. 1864. HB. 244.
stenochlamys C. Chr. Ind. 240. 1905 — Tonkin.
Asplenium platyphyllum Bak. JoB. **1890**. 264 (non J. Sm.: Mett. 1859).
Stoliczkae Bedd. Ferns br. Ind. Suppl. 13 t. 361. 1876 — India bor.
Asplenium Clarke 1880 — Diplazium thelypteroides Bedd. 1865 (non Pr. 1836).
striatum Desv. Prod. 281. 1827 = D. proliferum.
striatum Hk. Gen. t. 55 B. f. 3. 1838 = ? D. denticulosum.
striatum (L.) Pr. Tent. 114. 1836(?); Christ, Farnkr. 219. 1897. NPfl. 226 — Amer. trop. — Asplenium L. sp. **2**. 1082. 1753; Mett. Aspl. 185. n. 214. 1859; ? Athyrium Milde 1870. Diplazium crenulatum Liebm. 1849; Asplenium Bak. 1867. HB. 236; Athyrium Milde 1870 — Diplazium amplum Liebm. 1849; Asplenium Fourn. 1872. (HB. 491): Athyrium Milde 1870 — Diplazium grammitoides Fée 1866 (non Pr. 1849) — Gymnogramme grandis Bak. 1868. HB. 377; Phegopteris Salom. 1883 — Gymnogramme attenuata Fée 1869.
subserratum (Bl.) Moore, Ind. 338. 1862. NPfl. 225 — Java. Penang.
Asplenium Bl. Enum. 174. 1828. HB. 230; Athyrium Milde 1870. Diplazium lonchophyllum Kze. 1848 (non 1839); Asplenium Mett. n. 165. 1859.
subsilvaticum Christ, Prim. Fl. costar. **3**. 29. 1901; Bull. Boiss. II. **4**. 970. 1904 [— Costa Rica.
succulentum (Clarke) C. Chr. Ind. 240. 1905 — India bor.
Asplenium Clarke, Tr. Linn. Soc. II. Bot. **1**. 502. t. 64 f. 4. 1880.
sundense Hassk. Tijdschr. Nat. Gesch. **10**. 115. 1843 = D. subserratum.
Swartzii Bl. Enum. 191. 1828 = D. proliferum.
sylvaticum vide silvaticum.
tenebrosum (Jenm.) C. Chr. Ind. 240. 1905 — Jamaica.
Asplenium Jenm. Gard. Chr. III. **16**. 690. 1894.
tenerum Pr. Epim. 84. 1849 (nomen) = D. bulbiferum.
tenue Desv. Prod. 281. 1827 — Amer. austr. (an = D. striatum?).
ternatum Liebm. Vid. Selsk. Skr. V. **1**. 252 (seors. 100). 1849. NPfl. 228 — Mexico. — Asplenium Hk. 1860. HB. 242; Callipteris Moore 1861.
Textori Mak. Bot. Mag. Tokyo **13**. 31. 1899 = D. Mettenianum.
thelypteroides Pr. Tent. 114. 1836 = Athyrium acrostichoides.
thelypteroides Bedd. Ferns br. Ind. t. 68. 1865 = D. Stoliczkae.
Thwaitesii Kl. Cat. hort. Augustin. 4. 1857. NPfl. 226 — Ceylon.
Asplenium A.Br. 1857; Mett. Aspl. 183. n. 209 b. 1859. HB. 235; Athyrium Milde 1870. (D. japonicum*).
tomentosum Bl. Enum. 192. 1828. NPfl. 226 — India bor. Malesia.
Asplenium Hk. 1860. HB. 234; Athyrium Milde 1870. Asplenium argutans Fée 1850-52. Diplazium deflexum J. Sm. 1841; Asplenium Mett. n. 211. 1859.
travancoricum Bedd. Handb. 188. 1883 — India austr. — Asplenium Bak. 1891.
Trianae (Mett.) C. Chr. Ind. 240. 1905 — Columbia.
Asplenium Mett. Ann. sc. nat. V. **2**. 233. 1864 (HB. 232).
truncatum Pr. Tent. 114. 1836 = D. expansum.
tumulosum Moore, Ind. LV. 1857 = D. hians.
Tussacii Fée, Gen. 216. 1850-52 = D. costale.
umbrosum Willd. sp. **5**. 353. 1810 — Amer. austr. trop.
Athyrium Milde 1870. Asplenium radicans pt. HB. 211.
umbrosum Bl. Enum. 194. 1828 = D. maximum.
umbrosum Moritz, Syst. Verz. 111. 1845-46 = D. esculentum.

umbrosum Bedd. Handb. 188. 1883 = Athyrium.
undulatum Poir. Enc. Suppl. 2. 487. 1811 = D. undulosum.
undulosum Sw. Syn. 92, 284. 1806 (Plum. t. 107) — Martinique. — Callipteris Pr. 1849.
unilaterale Mett.; Christ, Prim. Fl. costar. 3. 29. 1901 (nomen) — Costa Rica.
Urbani (Christ) C. Chr. Ind. 241. 1905 — Cuba. Haïti.
Asplenium Christ, Engl. Jahrb. **24**. 108. 1897.
urticifolium Christ, Prim. Fl. costar. 3. 29. 1901 — Costa Rica.
vacillans (Kze.) C. Chr. Ind. 241. 1905 — Java. — Asplenium Kze. Bot. Zeit. **1848**. 172; Tarachia Pr. 1849.
varium Gaud. Freyc. Voy. Bot. 322. 1827 — Ins. Moluccae.
Microstegia Pr. 1849; Asplenium Mett.; Salom. 1883.
vastum (Mett.) Diels, Nat. Pfl. 1⁴. 228. 1899 — Columbia. Ecuador.
Asplenium Mett. Ann. sc. nat. V. 2. 237. 1864. HB. 241.
venulosum (Bak.) Diels, Nat. Pfl. 1⁴. 226, 228. 1899 — Ecuador.
Asplenium Bak. 238. 1867.
vestitum Pr. Epim. 87. 1849 — Ins. Philipp.
Asplenium Hk. 1860. HB. 239; Athyrium Milde 1870.
villosum Pr. Tent. 114. 1836 — Mexico.
Asplenium? Pr. Rel. Haenk. 1. 46. 1825; Athyrium Milde 1870.
vincentis (Christ) C. Chr. Ind. 241. 1905 — St. Vincent (Ind. occ.).
Asplenium Christ, Engl. Jahrb. **24**. 109. 1897.
Virchowii (Kuhn) Diels, Nat. Pfl. 1⁴. 228. 1899 — Madagascar.
Asplenium Kuhn, Hill. Pl. Madag. exsicc. n. 4168; Bak. 1891 (nomen).
virescens Kze. Bot. Zeit. **1848**. 537. NPfl. 226 — China. Japonia.
Asplenium Mett. n. 227. 1859. HB. 237.
Vitiense Carr. in Seem. Fl. Vit. 357. 1873 = D. esculentum var.
Wageneri Kl.; A.Br. Ind. Sem. ht. Berol. 1855; Moore, Ind. 340. 1862 (Wagenerianum; an jure?) — Amer. trop. — Asplenium Mett. 1856; Aspl. n. 220. 1859; Athyrium Wagnerianum Milde 1870; Asplenium obtusum Mett. Fil. Lips. 78. 1856 (non Willd. 1810).
Weinlandii Christ, Bull. Boiss. II. **1**. 452. 1901 — N. Guinea.
Welwitschii (Hk.) Diels, Nat. Pfl. 1⁴. 226. 1899 — Angola.
Asplenium Hk. Syn. 235. 1867; Mett.; Kuhn, Fil. Afr. 119. 1868.
Wercklearum Christ, Bull. Boiss. II. **4**. 969. 1904 — Mexico-Columbia.
Wheeleri (Bak.) Diels, Nat. Pfl. 1⁴. 227. 1899 — Japonia.
Asplenium Bak. Ann. Bot. **5**. 309. 1891.
Wichurae (Mett.) Diels, Nat. Pfl. 1⁴. 226. 1899 — Japonia. China.
Asplenium Mett. Ann. Lugd. Bat. **2**. 237. 1866. HB. 233.
Wilsoni (Bak.) Diels, Nat. Pfl. 1⁴. 228. 1899 — Jamaica.
Asplenium Bak. Syn. 242. 1867.
xiphophyllum (Bak.) C. Chr. Ind. 241. 1905 — Borneo.
Asplenium Bak. JoB. **1879**. 40.
zanzibaricum (Bak.) C. Chr. Ind. 241. 1905 — Zanzibar.
Asplenium Bak. Ann. Bot. **5**. 311. 1891.
zeylanicum (Hk.) Moore, Ind. 340. 1862. Chr. 216. NPfl. 225 (ceylanicum) — Ceylon. — Asplenium Hk. sp. **3**. 237. 1860; 2 Cent. t. 16. HB. 230; Athyrium Milde 1870. Diplazium pinnatifidum Fée 1865 (non Kze. 1839).
Zollingeri Kze. Bot. Zeit. **1848** 193 = D. cordifolium.

DIPLOÖPHYLLUM Van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 322. 1861. = **Hymenophyllum.**

Diplopterygium Diels, Nat. Pflanzenfam. 1⁴. 353. 1900 *(Gleichenia §)* = Gleichenia.

DIPLORA Baker, Journ. Bot. **1873**. 235 — (Polypodiaceae Gen. **45**).

Cadieri Christ, Journ. de Bot. **19**. 69. 1905 — Annam.

integrifolia Bak. JoB. **1873**. 235. HB. 492. NPfl. 230 — Ins. Salomonis.

DIPTERIS Reinwardt, Sylloge plant. **2**. 3. 1824 -- (Polypodiaceae Gen. **24**).

chinensis Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 109 c. fig et tab. — China.

conjugata Reinw. Syll. pl. **2**. 3. 1824. Chr. 123. NPfl. 203 — Asia, Polynesia trop. Polypodium Klf. 1827. Mett. Pol. n. 237 (non Poir. 1804); Phymatodes Pr. 1836; Polypodium Dipteris Bl. 1828. HB. 362; P. Horsfieldii R.Br.; Wall. 1828; Drynaria J. Sm. 1841; Dipteris Bedd. 1869.

crassifolia J. Sm. Lond. JoB. **1**. 196. 1842 = Polypodium.

Horsfieldii Bedd. Ferns br. Ind. t. 321. 1869 = D. conjugata.

Lobbiana (Hk.) Moore, Ind. LXXX. 1857. Chr. 123. NPfl. 203 — Malesia. Polypodium Hk. JoB. **5**. 309 t. 11. 1853. HB. ed. I. 362; P. bifurcatum Bak. 1874. HB. ed. II. 362.

quinquefurcata (Bak.) Christ, Farnkr. 123. 1897. NPfl. 203 — Borneo. Polypodium Bak. Journ. Linn. Soc. **24**. 260. 1887.

Wallichii (R.Br.) Moore, Ind. LXXX. 1857. Chr. 122. NPfl. 203 — India bor. Polypodium R.Br.; Wall. 1828 (nomen); Hk. et Grev. Ic. Fil. t. 168-169. 1829. Mett. Pol. n. 236. HB. 362; Drynaria J. Sm. 1841.

DISCOSTEGIA Presl, Suppl. Tent. Pterid. 11. 1845 = **Marattia.**

alata Pr. Suppl. 12. 1845.

laevis Pr. Suppl. 12. 1845 = M. alata.

microphylla Fée, 9 mém. 43. 1857.

DISPHENIA Presl, Tent. Pter. 55. 1836.

aculeata Pr. Tent. 56 t. 1 f. 13. 1836 = Alsophila armata.

arborea Pr. Tent. 56. 1836 = Cyathea.

aspera Pr. Abh. böhm. Ges. V. **5**. 349. 1848 = Alsophila.

Grevilleana Kze. Linn. **23**. 252. 1850 = Cyathea arborea.

muricata Pr. Abh. böhm. Ges. V. **5**. 349. 1848 = Cyathea.

orientalis Kze. Bot. Zeit. **1848**. 283 = Cyathea orientalis et javanica.

DISTAXIA Presl, Epim. Bot. 110. 1849 = **Blechnum.**

fraxinea Pr. Epim. 110, 261. 1849.

DOODIA R. Brown, Prod. Fl. Nov. Holl. 151. 1810 — (Polypodiaceae Gen. **56**).

aspera R.Br. Prod. Fl. N. Holl. 151. 1810. HB. 189. NPfl. 254 — Austr. temp. Woodwardia Fée 1850-52.

blechnoides Desv. Prod. 285. 1827 — Amer. austr. Woodwardia Desvauxii Moore 1862.

blechnoides A. Cunn. in Hk. Comp. Bot. Mag. **2**. 365. 1836 = D. maxima.

Brackenridgei Carr. in Seem. Fl. Vit. 352. 1873 = D. media var.

caudata (Cav.) R.Br. Prod. Fl. N. Holl. 151. 1810. HB. 190. Chr. 188. NPfl. 254 — Australia. N. Caledonia. N. Zealand. — Woodwardia Cav. Descr. 264. 1802 — D. rupestris Klf.; Link 1833 — D. linearis C. Moore; J. Sm. 1866 — D. squarrosa Col. 1881 — D. Harryana Moore 1884.

connexa Kze. Farnkr. **1**. 11 t. 6. 1840 = D. media var.

dives Kze. Bot. Zeit. **1848**. 144; Farnkr. **2**. 12 t. 105. HB. 189. Chr. 187. NPfl. 254 — Ceylon. Java. — Woodwardia Fée 1850-52.

duriuscula Moore, Gard. Chr. **1868**. 1114 = D. media.

Harryana Moore, Gard. Chr. n. s. **21**. 408. 1884 = D. caudata.

Kunthiana Gaud. Freyc. Voy. Bot. 401 t. 14. 1827 = D. media var.

linearis C. Moore; J. Sm. Ferns br. and for. 199. 1866 = D. caudata var.
lunulata R.Br. Prod. Fl. N. Holl. 151. 1810 = D. media.
maxima J. Sm. Bot. Mag. **72** Comp. 27. 1846 (syn.) — N. S. Wales.
D. blechnoides A. Cunn. in Hk. Bot. Mag. **2**. 365. 1836. HB. 189. NPfl. 254 (non Desv. 1827); Woodwardia Fée 1850-52.
media R.Br. Prod. Fl. N. Holl. 151. 1810. HB. 190, 482. Chr. 188. NPfl. 254 — Australia. N. Zealand. Polynesia. — Woodwardia Fée 1850-52 — Doodia lunulata R.Br. 1810; Woodwardia Fée 1850-52 — Doodia Kunthiana Gaud. 1827; Woodwardia Fée 1850-52 — Doodia connexa Kze. 1840; Woodwardia Fée 1850-52 — Doodia Brackenridgei Carr. 1873 — D. Milnei Carr. 1873 — D. duriuscula Moore 1868; Woodwardia Moore 1868. (N. Caledonia).
Milnei Carr. in Seem. Fl. Vit. 352. 1873 = D. media var.
polysora Terracino, Rend. Ac. Napoli **25**. 50. 1886 (polisora) — N. Caledonia.
rupestris Klf.; Link, Hort. Berol. **2**. 83. 1833 = D. caudata.
squarrosa Col. Tr. N. Zeal. Inst. **13**. 382. 1881 = D. caudata var.
virginica Pr Tent. 99. 1836 = Woodwardia.

DORCAPTERIS Presl, Epim. Bot. 166. 1849 = **Polybotrya.**
cervina Pr. Epim. 167. 1849.

Doryopteridastrum Fée, Cr. vasc. Br. **1**. 42. 1869 *(Pellaea §)* = Doryopteris.

DORYOPTERIS J. Smith, Journ. Bot. **3**. 404. 1841; **4**. 162. 1841 — (Polypodiaceae Gen. **68**).

acutiloba (Prantl) Diels, Nat. Pfl. 1^4. 269. 1899 — Brasilia.
Pellaea Prantl, Engl. Jahrb. **3**. 425. 1882.
alcicornis Diels, Nat. Pfl. 1^4. 270. 1899 = D. ornithopus.
angularis Fée, Cr. vasc. Br. **2**. 29 t. 88 f. 2. 1872-73 = D. varians.
argentea Christ, Bull. Boiss. II. **2**. 831. 1902 = Cheilanthes.
arifolia Christ in Schwacke, Pl. Nov. Mineiras **2**. 25. 1900; Bull. Boiss. II. **2**. 548. 1902 — Brasilia.
articulata Fée, Gen. 133. 1850-52 = Pellaea angulosa.
collina (Raddi) J. Sm. JoB. **4**. 163. 1841. NPfl. 270 — Brasilia. Guiana.
Pteris Raddi, Opusc. sci. Bol. **3**. 292. 1819; Pl. Bras. **1**. 44 t. 65 f. 1-2. 1825; Pellaea Prantl, Engl. Jahrb. **3**. 428. 1882 — Doryopteris euchlora Kl. 1847.
columbina (Hk.) Diels, Nat. Pfl. 1^4. 269. 1899 — Brasilia.
Pellaea lomariacea b columbina Hk. sp. **2**. 133 t. 112 A. 1858; P. columbina Hk. Syn. 146. 1867; Prantl, Engl. Jahrb. **3**. 424.
concolor (Langsd. et Fisch.) Kuhn, v. Deck. Reis. 3^3. Bot. 19. 1879. NPfl. 269 — Trop. — Pteris Langsd. et Fisch. Ic. Fil. 19 t. 21. 1810; Pellaea Bak. 1870. Chr. 157; Allosorus O. Ktze. 1891. Pteris geraniifolia Raddi 1819; Doryopteris Kl. 1847; Pellaea Fée 1850-52. HB. 146; Platyloma Lowe 1857; Pteris Pohliana Pr. 1822; Cheilanthes Keys. 1873 — C. Kirkii Hk. 1861. HB. 132.
cordifolia J. Sm. JoB. **4**. 163. 1841 (nomen) — ? — Litobrochia Moore 1862 (nomen).
cordifolia (Bak.) Diels, Nat. Pfl. 1^4. 270. 1899 — Madagascar.
Pteris Bak. JoB. **1891**. 4. Chr. 163.
crenulans Christ in Schwacke, Pl. Nov. Mineiras **2**. 26. 1900; Bull. Boiss. II. **2**. 549. 1902 = D. lomariacea var.
crispatula (Bak.) C. Chr. Ind. 243. 1905 — Brasilia.
Pellaea Bak. Ann. Bot. **5**. 215. 1891; Allosorus O. Ktze. 1891.
decipiens (Hk.) J. Sm. Hist. Fil. 289. 1875 — Ins. Hawaii.
Pteris Hk. sp. **2**. 209. 1858. HB. 167; Litobrochia Moore 1862.

decora Brack. Expl. Exp. **16**. 103 t. 13 f. 1. 1854 — Ins. Hawaii.
Pteris Hk. 1858. HB. 167; Litobrochia Moore 1862.

deltoidea (Kze.) Diels, Nat. Pfl. 1[4]. 269. 1899 — Africa austr. — Cheilanthes Kze. Linn. **10**. 535. 1836; Pellaea Bak. 1867. HB. 146; Allosorus O. Ktze. 1891.

Ducloux ii Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 231 — China.

Duvalii Bellair, Rev. Hort. **1897**. 563 — Hort. — (D. sagittifolia × palmata).

elegans Christ, Bull. Boiss. II. **1**. 429. 1901 = D. nobilis.

euchlora Kl. Linn. **20**. 342. 1847 = D. collina.

?**Fournieri** (Bak.) C. Chr. Ind. 244. 1905 — Mexico.
Pellaea Bak. Syn. 476. 1874; Allosorus O. Ktze. 1891; Pellaea flavescens Fourn. Mex. pl. **1**. 119. 1872 (non Fée 1869).

geraniifolia Kl. Linn. **20**. 343. 1847 = D. concolor.

Glaziovii Diels, Nat. Pfl. 1[4]. 269. 1899 = D. microphylla.

Harrisonae (Jenm.) C. Chr. Ind. 244. 1905 — Guiana.
Pteris Jenm. Gard. Chr. III. **24** 414. 1898.

hastata J. Sm. JoB. **4**. 163. 1841 = D. sagittifolia.

hederacea Diels, Nat. Pfl. 1[4]. 270. 1899 = D. varians.

itatiaiensis Christ, Bull. Boiss. II. **2**. 549. 1902 = D. lomariacea var.

lomariacea (Kze.) Kl. Linn. **20**. 343. 1847. NPfl. 269 — Guiana. Brasilia. Peru.
Pteris Kze. msc. 1849. HB. 164. Chr. 162; Pellaea Hk. 1858 (α); Prantl. Engl. Jahrb. **3**. 424; Allosorus O. Ktze. 1891 — Pteris lomariacea var. actinophylla Bak. Fl. bras. 1[2]. 406 t. 60. 1870. Pellaea crenulans Fée 1872-73; Doryopteris Christ 1900 — Pellaea itatiaiensis Fée 1872-73; Pteris Szysz. 1888; Doryopteris Christ 1902. — (an Pteris paradoxa Schrad. 1824?).

lonchophora (Römer) J. Sm. Hist. Fil. 289. 1875. NPfl. 269 — Brasilia.
Cheilanthes Römer msc.; Pteris Mett. Cheil. 4 t. 3 f. 1-3. 1859. HB. 166; Pellaea Bak. 1870; Prantl, Engl. Jahrb. **3**. 425; Litobrochia Moore 1861. Heteropteris Dryopteris Fée 1869.

Lorentzii (Hier.) Diels, Nat. Pfl. 1[4]. 270. 1899 — Argentina.
Pellaea Hier. Engl. Jahrb. **22**. 392. 1896.

ludens (Wall.) J. Sm. Hist. Fil. 289. 1875. NPfl. 270 — India. Ins. Philipp.
Pteris Wall. 1828 (nomen); Hk. sp. **2**. 210. 1858. HB. 166. Chr. 163; Litobrochia Bedd. 1865; Pellaea Prantl 1882. Pteris multifida Roxb. 1844 (non Poir. 1804). Doryopteris Wallichii J. Sm. 1841 (nomen); Litobrochia Smithii Moore 1862.

microphylla (Fée) Christ, Bull. Boiss. II. **2**. 546. 1902 — Brasilia.
Pellaea Fée, Cr. vasc. Br. **1**. 43 t. 4 f. 2. 1869. HB. 476 (pt.) (non Mett. 1869); Allosorus O. Ktze. 1891; Cassebeera Christ 1900. Pellaea Glaziovii Bak. 1870 (β); Prantl, Engl. Jahrb. **3**. 422; Doryopteris Diels 1899. NPfl. 269.

muralis Christ, Bull. Ac. Géogr. Bot. Mans. **1904**. 111 — China.

nobilis (Moore) J. Sm.; Bak. Syn. ed. II. 167. 1874 (syn.) — Brasilia.
Litobrochia Moore, Gard. Chr. **1862**. 932. Pteris elegans Vell. Fl. flum. **11**. t. 81. 1827; Bak. Syn. ed. II. 167. 1874 (non Poir. 1804 nec Sw. 1817); Doryopteris Christ 1902.

ornithopus (Mett.) J. Sm. Hist. Fil. 289. 1875 — Brasilia.
Pteris Mett.; Bak. Syn. 166. 1867; Pellaea alcicornis Prantl, Engl. Jahrb. **3**. 429. 1882; Doryopteris Diels 1899. NPfl. 270.

palmata J. Sm. JoB. **4**. 163. 1841 = D. pedata var.

paradoxa (Fée) Christ, Bull. Boiss. II. **2**. 546. 1902 — Brasilia.
Cassebeera Fée, 7 mém. 30 t. 20 f. 2. 1857; Pellaea Fée 1869 (non Hk. 1858); Pteris Kuhn 1896. Pellaea columbina β vestita Bak. 1867. HB. 146; Pellaea vestita Prantl, Engl. Jahrb. **3**. 423. 1882; Doryopteris Diels 1899. NPfl. 269.

patula Fée, Cr. vasc. Br. **2**. 30 t. 89 f. 2. 1872-73 — Brasilia.
Pellaea Prantl, Engl. Jahrb. **3**. 426. 1882.

pedata (L.) Fée, Gen. Fil. 133. 1850-52. NPfl. 270 — Amer. trop. India austr.
Pteris L. sp. **2**. 1075. 1753. HB. 167 (pt.). Chr. 162; Litobrochia Pr. 1836; Cassebeera J. Sm. 1841; Pellaea Prantl, Engl. Jahrb. **3**. 427. 1882 — Pteris palmata Willd. 1810. HB. 166; Doryopteris J. Sm. 1841; Litobrochia Moore 1862.

pedatifida Christ, Bull. Boiss. II. **2**. 546. 1902 — Brasilia.
Cassebeera Christ in Schwacke, Pl. Nov. Mineiras **2**. 25. 1900.

pedatoides (Desv.) Kuhn, v. Deck. Reis. **3**^{3}. Bot. 63. 1879 — Ins. Mascar. Madagascar. — Pteris Desv. Prod. 293 t. 7 f. 2. 1827.

phanerophlebia (Bak.) Diels, Nat. Pfl. **1**^{4}. 269. 1899 — Madagascar.
Pteris Bak. JoB. **1881**. 367; Ic. pl. t. 1640.

pilosa (Poir.) Kuhn, v. Deck. Reis. **3**^{3}. Bot. 63. 1879. NPfl. 269 — Ins. Mascarenae.
Pteris Poir. Enc. **5**. 717. 1804; Pellaea Hk. 1858. HB. 145. Chr. 157; Allosorus O. Ktze. 1891. Cheilanthes heterophylla Willd.; Klf. 1824; Allosorus Pr. 1836.

praealta Fée, Gen. 133. 1850-52 (nomen) —? — Litobrochia Moore 1862.

quinquelobata (Fée) Diels, Nat. Pfl. **1**^{4}. 269. 1899 — Brasilia.
Pellaea Fée, Cr. vasc. Br. **1**. 42 t. 10 f 1. 1869; Prantl, Engl. Jahrb. **3**. 422. P. Glaziovii Bak. 1870 (excl. var. minor). P. microphylla pt. HB. 476.

Raddiana (Pr.) Fée, Gen. 133. 1850-52 — Brasilia — Litobrochia Pr. 1836 (nomen); Pellaea Prantl, Engl. Jahrb. **3**. 427. 1882; Pteris pedata β et γ Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 45 t. 65 f. 2; 66 bis. 1825.

Raddiana β patula Fée, Cr. vasc. Br. **1**. 45 1869 = D. varians.

rediviva Fée, Cr. vasc. Br. **2**. 30 t. 89 f. 1. 1872-73 = D. varians.

?**rigida** (Sw.) Diels, Nat. Pfl. **1**^{4}. 269. 1899 — Mexico-Guatemala.
Pteris Sw. Syn. 104, 299. 1806; Allosorus Pr. 1836 (non Kze. 1834); Cheiloplecton Fée 1857; Pellaea Hk. 1858. HB. 147; Cheilanthes Moore: Fée 1857.

?**robusta** (Kze.) Diels, Nat. Pfl. **1**^{4}. 269. 1899 — Africa austr. (Namaqualand).
Allosorus Kze. Linn. **10**. 502. 1836; Onychium? Fée 1850-52; Pellaea Hk. 1858. HB. 149; Cryptogramme Pappe et Raws. 1858.

sagittifolia (Raddi) J. Sm. JoB. **4**. 163. 1841. NPfl. 269 — Venezuela Brasilia.
Pteris Raddi, Opusc. sci. Bol. **3**. 292. 1819; Pl. Bras. **1**. 43 t. 63 f. 1. 1825. HB. 166. Chr. 162; Litobrochia Pr. 1836; Pellaea Prantl, Engl. Jahrb. **3**. 426. 1882 — Pteris hastata Raddi 1825 (non Thbg. 1800 nec Sw. 1801); Litobrochia Pr. 1836; Doryopteris J. Sm. 1841.

?**Skinneri** (Hk.) C. Chr. Ind. 245. 1905 — Guatemala.
Pellaea Hk. sp. **2**. 141 t. 118 B. 1858. HB. 147; Allosorus O. Ktze. 1891.

squamosa (Hope et Wright) C. Chr. Ind. 245. 1905 — Yunnan.
Pellaea Hope et Wright, Journ. Linn. Soc **35**. 518. 1903.

subsimplex (Fée) Diels, Nat. Pfl. **1**^{4}. 269. 1899 — Brasilia.
Pellaea Fée, Cr. vasc. Br. **1**. 44 t. 4 f. 3. 1869; Prantl, Engl. Jahrb. **3**. 421; Pteris triphylla Bak. 1870 (non Ag. 1839); P. longula Bak. 1874. HB. 478 (an Mett.; Kuhn 1869?).

trifoliata (Bory) Fée, Gen. 133. 1850-52 (nomen) — ?
Pteris Bory hb.; Litobrochia Moore 1862.

triphylla (Lam.) Christ, Bull. Boiss. II. **2**. 546. 1902 — Brasilia austr.-Argentina.
Adiantum Lam. Enc. **1**. 41. 1783; Cassebeera Klf. 1824. HB. 142. Chr. 154. NPfl. 288; Pteris Mett. 1856; Pellaea Prantl 1882; Bakeriopteris O. Ktze. 1891.

varians (Raddi) J. Sm. JoB. **4**. 163. 1841 — Brasilia.
Pteris Raddi, Opusc. sci. Bol. **3**. 292. 1819; Pl. Bras. **1**. 44 t. 64. 1825; Litobrochia Pr. 1836. Pteris hederacea Pr. 1822; Litobrochia Pr. 1836; Pellaea Prantl, Engl. Jahrb. **3**. 428. 1882; Doryopteris Diels 1899. NPfl. 270 — D. angularis Fée 1872-73 — D. rediviva Fée 1872-73.

vestita Diels, Nat. Pfl. **1**[4]. 269. 1899 = D. paradoxa.

Wallichii J. Sm. JoB. **3**. 404. 1841 = D. ludens.

DRYMOGLOSSUM Presl, Tent. Pterid. 227. 1836 — (Polypodiaceae Gen. 93).

abbreviatum Fée, 7 mém. 26 t. 10 f. 2. 1857 — Cochinchina.
Hymenolepis Fée 1865.

acrostichoides Moore, Ind. XXXI. 1857 = Elaphoglossum conforme var.

Beddomei Clarke, Tr. Linn. Soc. II. Bot. **1**. 576. 1880 = D. heterophyllum.

carnosum (Wall.) J. Sm. in Hk. Gen. t. 78 A. 1841. HB. 397 (excl. var.). NPfl. 303 — China. India bor. — Nothochlaena Wall. 1828 (nomen); Lemmaphyllum Pr. Epim. 158. 1849; Taenitis Mett. 1856; Elaphoglossum Keys. 1873; Oetosis O. Ktze. 1891; (Asteroglossum J. Sm. msc.).

carnosum var. minor Hk. sp. **5**. 189. 1864 = D. microphyllum.

carnosum var. obovatum Harr. Journ. Linn. Soc. **16**. 33. 1877 = D. obovatum.

Cunninghami Moore, Ind. XXXI. 1857 = Cyclophorus confluens.

ellipticum Moore, Ind. XXXI. 1857 = ? D. heterophyllum.

heterophyllum (L.) C. Chr. Ind. 246. 1905 — Asia trop.
Acrostichum L. sp. **2**. 1067. 1753. Pteris piloselloides L. sp. ed. II. **2**. 1530. 1763; Taenitis (R.Br. 1810) Mett. 1856; Nothochlaena Klf. 1824; Pteropsis Desv. 1827; Drymoglossum Pr. 1836. HB. 398. Chr. 131. NPfl. 303; Elaphoglossum Keys. 1873; Lemmaphyllum Luerss. 1882; Oetosis O. Ktze. 1891 — ? Pteris elliptica Willd. 1810 (non Poir. 1804); Pteropsis Desv. 1827; Drymoglossum Moore 1857; Pteris ceilanica Wikstr. 1826 — Drymoglossum spatulatum Pr. 1836; Lemmaphyllum Pr. 1849.

lanceolatum J. Sm. JoB. **4**. 66. 1841 = Paltonium.

martinicense Christ, Engl. Jahrb. **24**. 137. 1897. Chr. 131. NPfl. 303 — Martinique.

microphyllum (Pr.) C. Chr. Ind. 246. 1905 — Japonia. China. Réunion.
Lemmaphyllum Pr. Epim. 263. 1849; Taenitis Mett. 1868; Pteris piloselloides Thbg. Fl. Jap. 331. 1784 (non L. 1763). Drymoglossum subcordatum Fée 1851-52. Chr. 131. NPfl. 303; D. carnosum β HB. 397.

niphoboloides (Luerss.) Bak in Hk. Ic. pl. t. 1686. 1887. Chr. 131. NPfl. 303 — Madagascar. — Taenitis Luerss. Abh. Nat. Ver. Bremen **7**. 49 t. 1 f. 3-5. 1882.

novo-guineae Christ in Schum. et Laut. Fl. deut. Südsee 137. 1901 — N. Guinea.

obovatum (Harr.) Christ, Journ. de Bot. **19**. 73. 1905 — Formosa. Annam.
D. carnosum var. obovatum Harr. Journ. Linn. Soc. **16**. 33. 1877.

piloselloides Pr. Tent. 227 t. 10 f. 5. 6. 1836 = D. heterophyllum.

rigidum Hk. Ic. pl. t. 996. 1854. HB. 398. Chr. 131. NPfl. 303 — Borneo.
Schizolepton Moore 1857; Oetosis O. Ktze. 1891.

rotundifolium Pr. Epim. 157. 1849 = D. heterophyllum.

spatulatum Pr. Tent. 227. 1836 = D. heterophyllum.

subcordatum Fée, 3 mém. 29. 1851-52; Gen. 94 t. 9 A. f. 1. 1850-52 = D. [microphyllum.
Wiesbaurii Sod. Cr. vasc. quit. 419. 1893 — Ecuador.

DRYMOTAENIUM Makino, Bot. Mag. Tokyo **15**. 102. 1901 — (Polypodiaceae Gen. **94**).
Miyoshianum Mak. Bot. Mag. Tokyo **15**. 109. 1901; Phan. et Pter. Jap. icon. **1**. t. 56-58. 1901 — Japonia. — Taenitis Mak. Bot. Mag. Tokyo **12**. 26. 1898.

DRYNARIA (Bory) J. Smith, Journ. Bot. **4**. 60. 1841 — (Polypodiaceae Gen. **106**).
Polypodium § Drynaria Bory, Ann. sc. nat. **5**. 464 t. 12-14. 1825.
acuminata Brack. Expl. Exp. **16**. 42. 1854 = Polypodium accedens.
acuminata Brack. Expl. Exp. **16**. 47. 1854 = Polypodium euryphyllum.
acuminata Fée, Cr. vasc. Br. **1**. 122 t. 37 f. 3 = Polypodium lycopodioides.
adfinis J. Sm. JoB. **3**. 398. 1841 } Polypodium commutatum.
affinis Fée, Gen. 271. 1850-52 }
alata Brack. Expl. Exp. **16**. 48 t. 6 f. 1. 1854 = Polypodium Wilkesii.
albido-squamata J. Sm. JoB. **3**. 397. 1841 = Polypodium.
alternifolia Brack. Expl. Exp. **16**. 53. 1854 = Polypodium.
angusta Fée, Gen. 270. 1850-52 = Polypodium.
?angustata Fée, Gen. 270. 1850-52 = Cyclophorus.
angustiloba (Pr.) Moore, Ind. 344. 1862 - ? — Phymatodes Pr. 1836 (nomen).
araneosa Fée, Gen. 270. 1850-52 = Polypodium aureum *areolatum.
Baronii Diels, Nat. Pfl. 1^4. 330. 1899 = D. reducta.
Baudouini Fourn. Ann. sc nat. V. **18**. 285. 1873. NPfl. 330 — N. Caledonia.
Polypodium Bak. 1891.
Billardieri J. Sm. Bot. Mag. **72** Comp. 14. 1846 = Polypodium diversifolium.
brancaefolia Moore, Ind. 345. 1862 = D. quercifolia.
Browniana Fée, Gen. 270. 1850-52 = Polypodium Brownii.
capitellata J. Sm. Bot. Mag. **72** Comp. 14. 1846 = Polypodium Wallichianum.
coadunata Brack. Expl. Exp. **16**. 49. 1854 = Aspidium macrodon.
compacta Fée, Gen. 270. 1850-52 = Polypodium lanceolatum.
conjugata Moore, Ind. 345. 1862 = Polypodium (Poir.).
conjugata Bedd. Ferns br. Ind. correct. 1870 = Polypodium coronans
cordifolia Fée, Gen. 270. 1850-52 = Aspidium trifoliatum.
coronans J. Sm. JoB. **4**. 61. 1841 = Polypodium.
crassifolia J. Sm. JoB. **4**. 61. 1841 = Polypodium.
crassinervata Fée, 8 mém. 97. 1857 = Polypodium lanceolatum*.
curvivenia (Wall.) Moore, Ind. 345. 1862 — Singapore.
Polypodium Wall. 1828 (nomen).
cuspidiflora J. Sm. JoB. **3**. 397. 1841 = Polypodium accedens.
decurrens Brack. Expl. Exp. **16**. 48. 1854 = Polypodium insigne.
Delavayi Christ, Bull. Soc. Fr. **52** Mém. I. 22. 1905 — China.
diversifolia J. Sm. JoB. **3**. 397. 1841 = D. rigidula.
dubia J. Sm. JoB. **3**. 397. 1841 = Polypodium pteropus.
elastica Fée, 11 mém. 72 t. 20 f. 2. 1866 = Polypodium petraefolium
elongata Fée, Gen. 271. 1850-52 = Polypodium lanceolatum*.
elongata Brack. Expl. Exp. **16**. 42. 1854 = Polypodium lineare.
excavata Fée, Gen. 270. 1850-52 = Polypodium.
exsul Fée, Cr. vasc. Br. **2**. 67. 1872-73 = Polypodium.
Fortunei (Kze.) J. Sm. Bot. Voy. Herald. 425. 1857. NPfl. 330 — China.
Polypodium Kze.; Mett. Pol. 121. n. 241. t. 3 f. 42-45. 1857. HB. 367. Chr. 119.
Fortuni Moore, Gard. Chr. **1855**. 708 = Polypodium lineare.

fulva Fée, Gen. 270. 1850-52 = Polypodium aureum *areolatum.
Gaudichaudii Bory; (Gaud. Freyc. Voy. Bot. 355. 1827) = D. rigidula.
glauca J. Sm. JoB. **3**. 397. 1841 = Polypodium glauco-pruinatum.
glaucina Fée, Gen. 270. 1850-52 = Polypodium aureum.
glaucistipes J. Sm. Hist. Fil. 108. 1875 = D. rigidula.
grossa Fée, Gen. 270. 1850-52 = Polypodium phymatodes.
hastata Fée, Gen. 271. 1850-52 = Polypodium.
hemionitidea J. Sm. JoB. **4**. 61. 1841 = Polypodium.
Heracleum Moore, Ind. 346. 1862 = Polypodium.
Horsfieldii R.Br.; J. Sm. JoB. **3**. 398. 1841 = Dipteris conjugata.
irioides J. Sm. JoB. **3**. 398. 1841 = Polypodium punctatum.
iteophylla J. Sm. Bot. Mag. **72** Comp. 14. 1846 = Polypodium geminatum.
latifolia Fée, Gen. 271. 1850-52 = Aspidium.
leiorhiza J. Sm. JoB. **4**. 61. 1841 = Polypodium.
(*lenticularis* Fée, Gen. 375 (index) 1850-52 —?).
lepidota Fée, Gen. 270. 1850-52 = Polypodium lanceolatum.
Lindbergii Fée, Cr. vasc. Br. **2**. 67. 1872-73 = Polypodium.
Linnaei Bedd. Ferns br. Ind. t. 315. 1869 = D. sparsisora.
lomarioides J. Sm. JoB. **3**. 397. 1841 = Polypodium.
longifolia Brack. Expl. Exp. **16**. 45. 1854 = Polypodium punctatum.
longifrons J. Sm. JoB. **3**. 397. 1841 = Polypodium normale var.
longipes J. Sm. Bot. Mag. **72** Comp. 14. 1846 = Polypodium phymatodes.
longissima J. Sm. JoB. **3**. 397. 1841 = Polypodium myriocarpum.
loriformis J. Sm. JoB. **4**. 61. 1841 = Polypodium excavatum.
lycopodioides Fée, Gen. 270. 1850-52 = Polypodium.
macrocarpa Fée, Gen. 270. 1850-52 = Polypodium lanceolatum.
?macrosora Fée, Gen. 270. 1850-52 = Cyclophorus angustatus.
maxima Brack. Expl. Exp. **16**. 51 t. 7. 1854 = Polypodium.
menisciicarpon J. Sm. JoB. **3**. 421. 1841 = Aspidium siifolium.
mexicana Fée, 8 mém. 97. 1857 = Polypodium lanceolatum*.
mollis Bedd. Ferns br. Ind. t. 216. 1867. NPfl. 330 — Himalaya. China merid.
Polypodium rivale Mett.; Bak. 1868. HB. 367. Chr. 121; Drynaria Christ 1899.
morbillosa J. Sm. JoB. **3**. 398. 1841 = Polypodium heracleum.
morbillosa Moore, Ind. 347. 1862 = D. quercifolia.
Moricandii Fée, Cr. vasc. Br. **1**. 121. 1869 = Polypodium.
musaefolia J. Sm. Cat. cult. ferns 14. 1857 = Polypodium.
nectarifera (Becc.) Diels, Nat. Pfl. 1^4. 330. 1899 — N. Guinea.
Polypodium Beccari; Bak. Malesia **2**. 247 t. 65. 1886.
neglecta J. Sm. JoB. **3**. 397. 1841 = Polypodium pyrolifolium.
normalis J. Sm. JoB. **4**. 61. 1841 = Polypodium.
nuda Fée, Gen. 270. 1850-52 = Polypodium lineare.
obtusata Brack. Expl. Exp. **16**. 44. 1854 = Polypodium punctatum.
oodes Fée, Gen. 270. 1850-52; 6 mém. 19 t. 7 f. 4. 1853 = Polypodium.
Ottoniana Fée, Gen. 270. 1850-52 = Phymatodes.
ovata Fée, Gen. 270. 1850-52 = Polypodium phyllomanes var.
palmata J. Sm. JoB. **3**. 397. 1841 = Polypodium.
Parishii Bedd. Ferns br. Ind. Suppl. 24. 1876 = D. propinqua.
Parkinsoni (Bak.) Diels in Schum. et Laut. Fl. deut. Südsee 143 1901 — N. Lauenburg. — Polypodium Bak. Ann. Bot. **5**. 481. 1891.
percussa Fée, Gen. 270. 1850-52 = Polypodium.

persicariaefolia Fée, Gen. 270. 1850-52 = Polypodium.
phlebodes Fée, Gen. 270. 1850-52 = Polypodium excavatum.
Phymatodes Fée, Gen. 270. 1850-52 = Polypodium.
pinnata Fée, Gen. 272. 1850-52 = D. rigidula.
plantaginea J. Sm. JoB. **4**. 61. 1841 = Aspidium.
pleuridioides Pr. Ann. Lugd. Bat. **2**. 229. 1866 (syn.). NPfl. 330 — Java. Polypodium Mett. Ann. Lugd. Bat. **2**. 229. 1866. (HB. 367); P. Willdenowii Bl. Fl. Jav. Fil. 156 t. 66. 1828. Mett. Pol. n. 238 (non Bory 1825).
polycarpa Brack. Expl. Exp. **16**. 44. 1854 = Polypodium punctatum.
Prieurii Fée, Gen. 271. 1850-52; 6 mém. 17 t. 2 f. 6 = Polypodium lanceolatum*.
propinqua (Wall.) J. Sm. JoB. **4**. 61. 1841. NPfl. 330 — China austr. India bor. Java. Fernando Po. — Polypodium Wall. 1828 (nomen); Mett. Pol. 120. n. 239 t. 3. f. 30. 1857. HB. 367. Chr. 119; Phymatodes Pr. 1836 — Pleopeltis Parishii Bedd. 1866; Drynaria Bedd. 1876.
Proustiana Gaud. Voy. Bonite, Bot. t. 3. 1846 = Polypodium Meyenianum.
pulverulenta Brack. Expl. Exp. **16**. 43. 1854 = Polypodium punctatum.
pustulata J. Sm. JoB. **4**. 61. 1841 = Polypodium.
quercifolia (L.) J. Sm. JoB. **3**. 398. 1841. NPfl. 329 — China austr. Asia, Austr. trop. Fiji. — Polypodium L. sp. **2**. 1087. 1753. Mett. Pol. n. 243. HB. 367. Chr. 118; Phymatodes Pr. 1836. Polypodium sylvaticum Schkuhr 1806; Phymatodes Pr. 1836; Polypodium Schkuhrii Bory 1825 — P. morbillosum Pr. 1825; Phymatodes Pr. 1836; Drynaria Moore 1862 (non J. Sm. 1841).
Raddiana Fée, Gen. 270. 1850-52 = Polypodium angustum.
reducta Christ in C. Chr. Ind. 249. 1905 — China. Polypodium Baronii Christ, Nu. Giorn. bot. It. n. s. **4**. 100 t. 2. 1897; Farnkr. 120 (non Bak. 1886); Drynaria Diels 1899. NPfl. 330.
revoluta J. Sm. JoB. **3**. 421. 1841 = Polypodium.
rigidula (Sw.) Bedd. Ferns br. Ind. t. 314. 1869. NPfl. 330 — Asia, Austr., Polynesia trop. — Polypodium Sw. Schrad. Journ. **1800**[2]. 26. 1801. HB. 368. Chr. 119; Goniophlebium Moore 1862. Polypodium diversifolium R.Br. 1810 (non Willd. 1810); Drynaria J. Sm. 1841. Polypodium Gaudichaudii Bory 1825. Mett. Pol. n 240; Drynaria Bory 1827; Phymatodes Pr. 1836.
rivalis Christ, Bull. Boiss. **7**. 6. 1899 = D. mollis.
rubida J. Sm. JoB. **3**. 397. 1841 = Polypodium longissimum.
rupestris J. Sm. JoB. **3**. 397. 1841 = Polypodium triquetrum var.
scandens Fée, Gen. 271. 1850-52 = Polypodium pustulatum.
Schomburgkiana Fée, Gen. 270. 1850-52 = Polypodium megalophyllum.
serpens Fée, Cr. vasc. Br. **2**. 66. 1872-73 = Polypodium Swartzii.
sesquipedalis J. Sm. Bot. Mag. **72** Comp. 13. 1846 = Polypodium excavatum.
sinica Diels, Engl. Jahrb. **29**. 208. 1900 — China.
sparsisora (Desv.) Moore, Ind. 348. 1862 — Ceylon. Malesia. Austr. trop. Fiji. Polypodium Desv. Berl. Mag. **5**. 315. 1811. P. Linnei Bory, Ann. sc. nat. **5**. 464 t. 12. 1825. Mett. Pol. n. 243 b. HB. 368. Chr. 118 (Linnaei); Drynaria Bedd. 1869. NPfl. 329.
spectrum J. Sm. JoB. **4**. 61. 1841 = Polypodium.
splendens Bedd. Ferns br. Ind. t. 316. 1869 = Dryostachyum.
squamata Fée, 11 mém. 72. 1866 = Polypodium.
stenoloma Fée, Gen. 272. 1850-52; 6 mém. 18 t. 4 f. 4 = Polypodium angustum.
stenophylla J. Sm. JoB. **3**. 397. 1841; Fée, 6 mém. 18 t. 8 f. 3. 1853 = Polypodium.

subfalcata J. Sm. JoB. **3**. 397. 1841 = Polypodium Zippelii.
subspathulata Hk. JoB. **9**. 356. 1857 = Polypodium lineare.
sylvatica Brack. Expl. Exp. **16**. 343. 1854 = Polypodium euryphyllum.
tenuiloris J. Sm. JoB. **3**. 397. 1841 = Polypodium.
Thouiniana Fée, Gen. 270. 1850-52 = Polypodium spectrum.
torulosa Fée, Gen. 271. 1850-52; 6 mém. 17 t. 5 f. 4 = Polypodium angustum.
tridactylis Fée, Gen. 271. 1850-52 = Polypodium pteropus.
undulata J. Sm. JoB. **3**. 397. 1841 = Polypodium Zollingerianum.
vestita Fée, Gen. 271. 1850-52; 6 mém. 16 t. 4 f. 3. = Polypodium polylepis.
vulgaris J. Sm. JoB. **3**. 397. 1841 = Polypodium phymatodes.
Wallichii J. Sm. JoB. **4**. 61. 1841 = Dipteris.
Willdenowii (Bory) Moore, Ind. LXXIX. 1857. NPfl. 330 — Ins. Mascarenae
Polypodium Bory, Ann. sc. nat. **5**. 468 t. 13. 1825. HB. 367; Phymatodes Thouarsii Pr. 1836; Polypodium Willd.; Mett. Pol. n. 238 b. 1857.
Zeylanica Fée, Gen. 272. 1850-52 = Polypodium lineare.

DRYOMENIS Fée, Genera Filicum 225. 1850-52 = **Aspidium.**
menisciicarpon Moore, Ind. XLVII. 1857 = A. siifolium.
Phymatodes Fée, Gen. 225 t. 18 A. f. 1. 1850-52 = A. siifolium.
plantaginea J. Sm. Bot. Voy. Herald 229. 1854.
Purdiei J. Sm. Bot. Voy. Herald 229. 1854 = A. draconopterum.

DRYOPTERIS Adanson, Fam. des plantes **2**. 20. 1763 — (Polypodiaceae Gen. **9**).

Subgenera: C = **Cyclosorus** (Nephrodium auctt.).
D = **Eudryopteris** (Lastrea auctt.).
G = **Goniopteris.**
L = **Leptogramma.**
M = **Meniscium.**
P = **Phegopteris.**
S = **Stegnogramma.**

Obs. Vide dubias species ad hoc genus pertinentes sub Aspidio, Goniopteride, Lastrea, Nephrodio, Phegopteride, Polypodio, Polysticho et aliis.

abbreviata Newm. Phytol. **1851**. app. XXI = D. filix mas.
abbreviata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.
(C) **abortiva** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Malesia. — Aspidium Bl. Enum. 154. 1828. Mett. n. 262; Nephrodium J. Sm.; Moore 1858. HB. 292.
abrupta O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. Kunzeana.
(D) **achalensis** (Hier.) C. Chr. Ind. 250. 1905 — Argentina.
Aspidium Hier. Engl. Jahrb. **22**. 371. 1896.
(D) **aciculata** (Bak.) C. Chr. Ind. 250. 1905 — Borneo.
Nephrodium Bak. Journ. Linn. Soc. **22**. 226. 1886.
(D) **acrosora** (Hier.) C. Chr. Ind. 250. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. **34**. 446. 1904.
acrostichoides O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.
aculeata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.
(D) **acuta** (Kl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Brasilia.
Lastrea Kl. hb.; Nephrodium Hk. sp. **4**. 147. 1862 (excl. t. 171). HB. 286; Aspidium Fée 1872-73.
(P) **acutidens** C. Chr. Ind. 250. 1905 — Ins. Hawaii. — Phegopteris spinulosa Hill. Fl. Haw. 566. 1888; Polypodium Bak. 1891 (non Dryopteris infra).

(C) **adenophora** C. Chr. Ind. 251. 1905 — Luzon. Celebes. — Nephrodium hirsutum J. Sm. 1841 (nomen); Pr. Epim. 48. 1849. HB. 291 (non Don 1825 nec Bory 1828); Aspidium Kze.; Mett. n. 256. 1858; Dryopteris O. Ktze. 1891.

adiantiformis O. Ktze. Rev. Gen. Pl. 3^2. 378. 1898 = Polystichum.

(D) **aemula** (Ait.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Eur. occ. Madeira. Azor. Polypodium Ait. Hort. Kew. **3**. 466. 1789; Aspidium Sw. 1801. Mett. n. 138. Chr. 262; Allantodia Desv. 1827; Polystichum Pr. 1836; Lastrea Brack. 1854; Nephrodium Bak. 1867. HB. 279. NPfl. 174. N. Foenisecii Lowe 1830; Aspidium Lowe 1834; Lastrea Wats. 1846; Lophodium Newm. 1851.

affinis Newm. Hist. br. Ferns ed. III. 187. 1854 = D. filix mas.

(L) **africana** (Desv.) C. Chr. Ind. 251. 1905 — Azor. Madeira-Afr. austr. Japonia. Korea. China-India bor.-Malesia. — Polypodium Desv. 1827; P. tottum Willd. sp. **5**. 201. 1810 (non Thbg. 1800); Gymnogramma Schlecht. 1825. HB. 376; Grammitis Pr. 1836; Leptogramma J. Sm. 1841; Phegopteris Mett. n. 31. 1858. Chr. 272; Aspidium Engl. 1892; Nephrodium Diels 1899. NPfl. 170; Gymnogramma Lowei Hk. et Grev. 1829; Leptogramma J. Sm. 1841. Gymnogramma aspidioides Bl. 1828 (non Desv. 1827); Grammitis Blumeana Pr. 1836.

(D) **alatella** (Christ) C. Chr. Ind. 251. 1905 — N. Guinea. — Nephrodium Christ in Schum. et Laut. Fl. deut. Südsee 112. 1901; Aspidium Christ 1901.

albopunctata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. orientalis.

(D) **Alfarii** (Christ) C. Chr. Ind. 251. 1905 — Costa Rica. — Aspidium Christ 1905; A. strigosum Christ, Bull. Soc. bot. Belg. **35**. 210. 1896 (non Willd. 1810).

(P) **alloëoptera** (Kze.) C. Chr. Ind. 251. 1905 — Columbia. Polypodium Kze. Linn. **25**. 748. 1853. HB. 305; Phegopteris Mett. n. 34. 1858; Nephrodium alleopterum Diels 1899. NPfl. 168.

alsophilacea O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. tenuifolia.

(P) **alsophiloides** (Fourn.) C. Chr. Ind. 251. 1905 — N. Caledonia. — Phegopteris Fourn. Ann. sc. nat. V. **18**. 289. 1873; Polypodium Balansae Bak. 1891.

amabilis O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(P) **amaurophylla** (Christ) C. Chr. Ind. 251. 1905 — Yunnan. Phegopteris Christ, Bull. Boiss. **7**. 14. 1899.

amboinensis O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. parasitica*.

(D) **ameristoneura** (Fée) C. Chr. Ind. 251. 1905 — India occ. Aspidium Fée, 8 mém. 104. 1857; Lastrea Moore 1858. Nephrodium Grisebachii Bak. 1867. HB. 285; Dryopteris O. Ktze. 1891; Nephrodium amplum Hk. 1862.

(D) **amphioxypteris** (Sod.) C. Chr. Ind. 251. 1905 — Ecuador. Nephrodium Sod. Rec. 51. 1883; Cr. vasc. quit. 230. 1893.

(D) **ampla** (H.B.Willd.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Amer. trop. Polypodium H.B.Willd. sp. **5**. 207. 1810; Phegopteris Fée 1850-52; Aspidium Mett. n. 176. 1858. Chr. 264; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 285. NPfl. 177 (non Hk. 1862) — Polypodium Sloanei Pöppig; Spr. 1827. — vide Polypodium pulverulentum Poir.

(D) **amplificata** (Fée) C. Chr. Ind. 251. 1905 — Mexico. Phegopteris Fée, 8 mém. 89. 1857; Aspidium Fée; Fourn. 1872.

(D) **amplissima** (Pr.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Guiana. Brasilia. Polystichum Pr. 1836 (nomen); Epim. 58. 1849; Aspidium Mett. n. 160. 1858. Chr. 265; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 502

— Aspidium fallax Fisch. 1850, 1855; Acrophorus Regel 1857; Lastrea Moore 1858 — Aspidium biforme, latissimum, macrum, phaeochlamys, ramosum Fée 1869. — (an potius Polystichum?).

(D) **anateinophlebia** (Bak.) C. Chr. Ind. 252. 1905 — Madagascar.
Nephrodium Bak. Journ. Linn. Soc. **16**. 202. 1877; Aspidium Kuhn 1879.

(M) **Andreana** (Sod.) C. Chr. Ind. 252. 1905 — Ecuador. Costa Rica. — Meniscium Sod. Rec. 71. 1883; Cr. vasc. quit. 392. 1893; Phegopteris Christ 1901.

(P) **aneitensis** (Hk.) C. Chr. Ind. 252. 1905 — N. Hebridae.
Polypodium Hk. sp. **4**. 268. 1862. (HB. 312); Phegopteris Kuhn 1869.

angulata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(M) **angustifolia** (Willd.) Urban, Symb. Ant. **4**. 21. 1903 — Amer. trop.
Meniscium Willd. sp. **5**. 133. 1810. HB. 391; Phegopteris Mett. 1859.

(D) **angustifrons** (Moore) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India bor.
Lastrea Moore msc. (non Ind. 85. 1858); Bedd. 1867; Nephrodium splendens var. angustifrons Hk. sp. **4**. 126. 1862; N. angustifrons Bak. 1867. HB. 283. NPfl. 174.

anomala O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum aculeatum var. 37.

(C) **anoptera** (Kze.) C. Chr. Ind. 252. 1905 — Brasilia. — Aspidium Kze.; Kuhn, Linn. **36**. 113. 1869; A. nitidulum Kze.; Ettingsh. 1865; Nephrodium Bak. 1870. HB. 502 (vix Pr. 1849); Dryopteris O. Ktze. 1891. Aspidium catacolobum Kze.; Ettingsh. 1865 (nomen).

(G) **aoristisora** (Harr.) C. Chr. Ind. 252. 1905 — Ins. Philipp.
Polypodium Harr. Journ. Linn. Soc. **16**. 30. 1877.

(D) **aperta** (Fée) C. Chr. Ind. 252. 1905 — Mexico.
Aspidium Fée, 8 mém. 106. 1857; Lastrea Moore 1858.

(D) **apicalis** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ins. Lord Howe.
Nephrodium Bak. Syn. 499. 1874; Aspidium Benth. 1878.

(P) **apicidens** (Bak.) C. Chr. Ind. 252. 1905 — Yunnan.
Polypodium Bak. Kew Bull. **1895**. 54.

(D) **apiciflora** (Wall.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India bor.
Aspidium Wall. 1828 (nomen); Mett. Pheg. u. Asp. 54. n. 128. 1858; Lastrea Pr. 1836 (nomen); Nephrodium Hk. 1862. HB. 271. NPfl. 173.

apiifolia O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = Polystichum.

(C) **appendiculata** (Bl.) C. Chr. Ind. 252. 1905 — Java. — Gymnogramma Bl. Enum. add. 1828; Goniopteris Pr. 1836; Phegopteris Mett.; Salom. 1883; Nephrodium Racib. 1898; Aspidium pilosiusculum Mett. 1864. (HB. 293).

(D) **aquilina** (Thouars) C. Chr. Ind. 252. 1905 — Tristan d'Acunha. Ins. Amsterdam.
Polypodium Thouars, Fl. Trist. d'Ac. 32. 1804. HB. 311; Phegopteris Mett.; Kuhn 1868; Nephrodium Hemsl. 1885.

(D) **aquilinoides** (Desv.) C. Chr. Ind. 252. 1905 — Réunion.
Nephrodium Desv. Prod. 261. 1827; Aspidium Mett.; Kuhn 1868.

(D) **aquilonaris** Maxon, Bull. Torr. Cl. **27**. 638. 1900 — Alaska.

arborescens O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. ludens.

(C) **arbuscula** (Willd.) O. Ktze. Rev. Gen. Pl. **2** 812. 1891 — India-Malesia-Polynesia. Ins. Mascarenae. — Aspidium Willd. sp. **5**. 233. 1810. Mett. n. 253. Chr. 249; Nephrodium Desv. 1827. HB. 292. NPfl. 179. Aspidium Hookeri Wall. 1829. Hk. 1854; Nephrodium Moore et Houlst. —; J. Sm. 1857.

(D) **Arechavaletae** (Hier.) C. Chr. Ind. 252 1905 — Uruguay.
Aspidium Hier. Engl. Jahrb. **22**. 370. 1896.

(G) **arfakiana** (Bak.) C. Chr. Ind. 253. 1905 — N. Guinea.
Polypodium Bak. Malesia **3**. 45. 1886.

(D) **argentina** (Hier.) C. Chr. Ind. 253. 1905 — Argentina.
Aspidium Hier. Engl. Jahrb. **22**. 367. 1896.

(C) **arida** (Don) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India bor.-Malesia-Ins. Philipp. — Aspidium Don, Prod. Fl. Nepal. 4. 1825. Chr. 247; Nephrodium J. Sm. 1841. HB. 291. NPfl. 178 — Aspidium obscurum Bl. 1828. Mett. n. 260; Nephrodium Moore 1858 — ? N. plumiferum Desv. 1827; Aspidium Kuhn 1889.

aristata O. Ktze. Rev. Gen. Pl. **2** 812. 1891 = Polystichum.

(G) **arthrothrix** (Hk.) C. Chr. Ind. 253. 1905 — Madagascar. Ins. Mascarenae.
Polypodium Hk. sp. **5**. 14. 1863. HB. ed. I. 315; P. tomentosum Bory, Bél. Voy. Bot. **2**. 31. 1833. HB. ed. II. 315 (non Thouars 1804); Phegopteris Mett.; Kuhn 1868.

(D) **ascensionis** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ascension.
Nephrodium Hk. sp. **4**. 124 t. 257. 1862. HB. 279. NPfl. 174; Aspidium Kuhn 1868; Lastrea J. Sm. 1875.

(P) **asperula** (J. Sm.) C. Chr. Ind. 253. 1905 — Ins. Philipp.
Polypodium J. Sm. 1841 (nomen); Hk. sp. **4**. 257. 1862. HB. 311; Dicksonia Fée 1850-52 (nomen).

(L) **aspidioides** (Willd.) C. Chr. Ind. 253. 1905 — Amer. austr. trop.
Ceterach Willd. sp. **5**. 137. 1810; Gymnogramma Klf. 1824; Leptogramme Kl. 1847; Phegopteris Mett. 1856. n. 24; Grammitis Moore 1857. Gymnogramma asplenioides Sw. 1817. HB. 376; Grammitis Pr. 1822; Leptogramma J. Sm. 1841; Phegopteris Mett. 1856. Chr. 273; Nephrodium Diels 1899. NPfl. 172 — Polypodium saxicola Sw. 1817.

(C) **asplenioides** (Sw.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India occ.
Polypodium Sw. Schrad. Journ. **1800²**. 26. 1801; Goniopteris Pr. 1836; Aspidium Gris. 1864. Chr. 243; Nephrodium Bak. 1867. HB. 293; Aspidium scolopendrioides var. 3 Mett. n. 235. 1858; A. sclerophyllum Kze. 1827; Nephrodium Pr. 1836.

(D) **athamantica** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Africa austr.
Aspidium Kze. Linn. **18**. 123. 1844. Mett. n. 154. Chr. 258; Lastrea Moore 1853; Nephrodium Hk. 1862. HB. 277.

(D) **athyrioides** (Mart. et Gal.) O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 — Mexico.
Aspidium Mart. et Gal. Mém. Ac. Brux. **15**. 67 t. 18. 1842; Lastrea Liebm. 1849; Athyrium sphaerocarpon Fée 1850-52; Nephrodium Hk. 1862. HB. 277. NPfl. 174; Aspidium Christ 1905.

(D) **atrorubens** (Mett.) C. Chr. Ind. 253. 1905 — Peru.
Aspidium Mett.; Kuhn, Linn. **36**. 112. 1869.

attenuata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. stenobasis.

auriculata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(L) **aurita** (Hk.) C. Chr. Ind. 253. 1905 — India bor.
Gymnogramme Hk. Ic. pl. t. 974, 989. 1854. HB. 377. (Chr. 274); Grammitis Moore 1856; Phegopteris J. Sm. 1857. Mett n. 22; Polypodium Lowe 1858; Leptogramme Bedd. 1883; Aspidium Christ 1904.

(D) **Balbisii** (Spr.) Urban, Symb. Ant. **4**. 14. 1903 — Amer. trop.
Polypodium Spr. Nova Acta **10**. 228. 1821; Aspidium Kuhn 1877; A. Sprengelii Klf. 1823. Mett. n. 195. Chr. 252; Lastrea Pr. 1836; Nephrodium Hk. 1862. HB. 268. NPfl. 172; Dryopteris O. Ktze. 1891.

(D) **barbigera** (Moore) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Afghanistan-India bor.-Yunnan. — Lastrea Moore hb.; Nephrodium Hk. sp. **4**. 113. 1862. HB. 274; NPfl. 173; Lastrea Bedd. 1867; Aspidium Chr. 260. 1897 — Nephrodium Falconeri Hk. 1862; HB. 277; Lastrea Bedd. 1865; Dryopteris O. Ktze. 1891.

Barclayi O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(P) **Baroni** (Bak.) C. Chr. Ind. 254. 1905 — Madagascar. Polypodium Bak. Journ. Linn. Soc. **22**. 535. 1887.

Barteri O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(P) **Barteriana** (Hk.) C. Chr. Ind. 254. 1905 — Africa occ. trop. — Polypodium Hk. sp. **4**. 253. 1862. (HB. 309); Phegopteris Mett.; Kuhn 1868.

(D) **basiattenuata** (Jenm.) C. Chr. Ind. 254. 1905 — Jamaica. Nephrodium Jenm. Gard. Chr. III. **15**. 330. 1894; Aspidium Jenm. 1895.

(C) **basilaris** (Pr.) C. Chr. Ind. 254. 1905 — Ins. Philipp. Nephrodium Pr. 1849 (nomen); N. caudiculatum J. Sm. 1841 (nomen, non Aspidium Sieb.; Kze. 1851); N. philippinense Bak. Ann. Bot. **5**. 327. 1891 (non Phegopteris (Fée) Mett. 1858); Aspidium Christ 1898.

(D) **Beccariana** (Cesati) C. Chr. Ind. 254. 1905 — Borneo. Nephrodium Cesati, Atti Ac Napoli 7^{8}. 23. 1876.

(D) **Beddomei** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — China austr. India austr.-Ceylon. Ins. Philipp. — Nephrodium Bak. Syn. 267. 1867; NPfl. 172; Lastrea Bedd. 1876; Aspidium Prantl 1882; Lastrea gracilescens Bedd. 1863-65 (non Aspidium Bl. 1828).

(D) **Bedieri** (Cordem.) C. Chr. Ind. 254. 1905 — Réunion. — Nephrodium Cordem. Bull. soc. sci. et arts Réunion **1890-91**. —; Fl. Réun. 77. 1895.

(D) **Bergiana** (Schlecht.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Africa austr. Polypodium Schlecht. Adumbr. 20 t. 9. 1825; Aspidium Mett. n. 188. 1858; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 269 (pt.). NPfl. 172.

(C?) **bermudiana** (Bak.) Gilb. Bull. Torr. Cl. **25**. 600. 1898 — Bermuda. Nephrodium Bak. ap. Hemsl. Chall. Exp. Bot. $\mathbf{1}^{1}$. 86 t. 13. 1885.

(C) **biaurita** (Bedd.) C. Chr. Ind. 254. 1905 — Assam. Nephrodium Bedd. Handb. Suppl. 68. 1892.

(C) **bibrachiata** (Jenm.) C. Chr. Ind. 254. 1905 — Jamaica. Nephrodium Jenm. Gard. Chr. III. **15**. 230. 1894.

(G?) **biformis** (Boiv.) C. Chr. Ind. 254. 1905 — Ins. afric. orient. Lastrea Boiv. msc.; Phegopteris Mett.; Kuhn, Fil. Afr. 121. 1868; Polypodium Bak. 1874. HB. 505; Nephrodium Diels 1899. NPfl. 178. Polypodium polyxiphion Bak. 1885.

(P) **biserialis** (Bak.) C. Chr. Ind. 254. 1905 — Peru. Ecuador. Polypodium Bak. Syn. 309. 1867; Nephrodium Diels 1899. NPfl. 170.

(P) **Bissetiana** (Bak.) C. Chr. Ind. 254. 1905 — Japonia. Nephrodium Bak. JoB. **1877**. 366.

(P) **bivestita** (Tausch) C. Chr. Ind. 254. 1905 — Mauritius. — Polypodium Tausch; Pr. 1836 (nomen); Phegopteris Mett. Pheg. u. Asp. 27. n. 57. 1858.

(P) **blanda** (Fée) C. Chr. Ind. 254. 1905 — Mexico. Phegopteris Fée, 8 mém. 91. 1857; Polypodium Bak. 1891 (non Fée 1857).

(D) **Blanfordii** (Hope) C. Chr. Ind. 254. 1905 — India bor. Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 624 t. 11. 1899.

(D) **Bodinieri** (Christ) C. Chr. Ind. 254. 1905 — China. Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 248 c. fig.

(D) **Boivini** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Madagascar.
Nephrodium Bak. Syn. 501. 1874; Aspidium Kuhn 1879.

Bojeri O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. filix mas *elongata var.

Boottii Und. Our nat. Ferns ed. IV. 117. 1893 = ? D. cristata × spinulosa.

(D) **borbonica** (Bak.) C. Chr. Ind. 255. 1905 — Réunion.
Nephrodium Bak. Ann. Bot. **5**. 317. 1891.

(D?)?**borneensis** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Borneo.
Lastrea Hk. Ic. pl. t. 993. 1854; Nephrodium Hk. 1862. HB. 273; Leucostegia J. Sm. 1866; Davallia J. Sm.; Kuhn 1869.

Borreri Newm. Hist. Br. Ferns ed. III. 189. 1854 = D. filix mas.

(D) **Boryana** (Willd.) C. Chr. Ind. 255. 1905 — Japonia. India bor. China austr.-Malesia. Réunion. — Aspidium Willd. sp. **5**. 285. 1810; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 284; NPfl. 177. Aspidium divisum Wall. 1828; Lastrea Moore 1858; Nephrodium Hk. 1862; Dryopteris O. Ktze. 1891.

(D) **Bourgaei** (Fourn.) C. Chr. Ind. 255. 1905 — Mexico.
Aspidium Fourn. Mex. pl. **1**. 98. 1872.

(C) **brachyodus** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ind. occ. Costa Rica-Peru. Galapagos. Malesia. — Polypodium Kze. Linn. **9**. 48. 1834; Phegopteris Mett. n. 38. 1858; Nephrodium Hk. 1862. HB. 295. NPfl. 181; Aspidium Gris. 1864. Chr. 248. Phegopteris Seemanni J. Sm. 1854 — ? Phegopteris abbreviata Mett. n. 45. 1858; (Polypodium Sieb. exsicc.; Goniopteris Pr. 1836).

(D) **brachypoda** (Bak.) C. Chr. Ind 255. 1905 — Guiana.
Nephrodium Bak. Tr. Linn. Soc. II. Bot. **2**. 290. 1887.

(D) **Brackenridgei** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Polynesia.
Aspidium Mett. 1861; Lastrea Carr. 1873; Nephrodium Bak. 1874 HB. 494; Lastrea attenuata Brack. Expl. Exp. **16**. 193 t. 26 f. 2. 1854 (non J. Sm. 1841). Nephrodium inaequilaterale Bak. 1868; Aspidium Luerss. 1871.

(P) **braineoides** (Bak.) C. Chr. Ind. 255. 1905 — China austr. occ.
Polypodium Bak. JoB. **1888**. 229; Nephrodium Diels 1899. NPfl. 171.

(D) **Brauniana** (Karst.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Columbia.
Aspidium Karst. Fl. Col. **1**. 63 t. 31. 1859; Nephrodium Hk. 1862. HB. 260.

Braunii Und. ap. Britt. and Brown, Ill. Fl. North. U. S. **1**. 15. 1896 = [Polystichum.

(M) **brevipinna** C. Chr. Ind. 255. 1905 — Borneo.
Meniscium stenophyllum Bak. JoB. **1891**. 108 (non Nephrodium Sod. 1883 nec Bak. 1884 et 1886).

(P) **brunnea** (Wall.) C. Chr. Ind. 255. 1905 — China austr. Formosa. India bor.-Malesia. Japonia. — Polypodium Wall. 1828 (nomen); Phegopteris J. Sm. 1875. Polypodium distans Don, Prod. Fl. Nepal. 2. 1825. HB. 308 (non Klf. 1824); Phegopteris Mett. n. 23. 1858. Chr. 274; Nephrodium Diels 1899. NPfl. 170; Aspidium Christ 1902 — A. paludosum Bl. 1828 (non Raddi 1825); Polypodium Bl. 1829; Lastrea Moore 1858; Phegopteris J. Sm. 1875 — Nephrodium microstegium Hk. 1862; Lastrea Bedd. 1865 — Polypodium Griffithii Hk. 1862 — P. pyrrhorachis Kze. 1851.

(P) **brunneo-villosa** C. Chr. Ind. 255. 1905 — N. Guinea.
Cheilanthes gigantea Ces. Rend. Ac. sci. Nap. **16**. 25, 29. 1877 (non Dryopteris infra); Polypodium cheilanthoides Bak. Malesia **3**. 45. 1886 (non Aspidium Kze. 1851).

(D) **Brunoniana** (Wall.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India bor. Aspidium Wall. 1828 (nomen): Mett. Pheg. u. Asp. 54 n. 127. 1858. Chr. 259; Lastrea Pr. 1836 (nomen); Nephrodium Hk. sp. **4**. 113 t. 251. 1862. HB. 274. NPfl. 173.

(D) **Buchanani** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Sikkim. Natal. Madagascar. Réunion. — Nephrodium Bak. Syn 498. 1874; Ic. pl. t. 1662; NPfl 174; Lastrea Bedd. 1883.

Obs. Plant. ind. f. Clarke ad D. pulvinuliferam, plant. afr. f. Kuhn ad D. squamisetam pertinentes.

(D) **Buchholzii** (Kuhn) C. Chr. Ind. 256 1905 — Kamerun. — Aspidium Kuhn, v. Deck. Reisen **3**[3]. Bot. 47. 1879; Nephrodium Bak. 1891.

caduca O. Otze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum Hookerianum.

(P) **caespitosa** (Fourn.) C. Chr. Ind. 256. 1905 — Mexico. Phegopteris Fourn. Mex. pl. **1**. 89. 1872; Polypodium Bak. 1874. HB. ed. II. 305 (non Link 1833); Nephrodium Diels 1899. NPfl. 172.

(D) **calcarata** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — China austr.-India-Malesia-Samoa. — Aspidium Bl. Enum. 159. 1828; Chr. 255; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 264, 494. NPfl. 171 — Aspidium Reinwardtianum Kze. 1848. Mett. n. 206; Lastrea Moore 1858 — L. falciloba Hk. 1857; Aspidium Benth. 1861; Nephrodium Hk. 1862 — Aspidium ciliatum Wall. 1828; Nephrodium Clarke 1880; Lastrea sericea Scott; [Bedd. 1869.

calcarea O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

californica O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(D) **callopsis** Franch. et Sav.) C. Chr. Ind. 256. 1905 — Japonia. Aspidium Franch. et Sav. En. pl. Jap. **2**. 240, 634. 1879.

(D) **callosa** (Bl.) C. Chr. Ind. 256. 1905 — Java. Sumatra. Aspidium Bl. Enum. 152. 1828. Mett. n. 259; Nephrodium Keys. 1873.

(P) **camporum** (Lindm.) C. Chr. Ind. 256. 1905 — Brasilia austr. Polypodium Lindm. Ark. f. Bot. **1**. 228 t. 10 f. 13. 1903.

(P) **camptocaulis** (Fée) C. Chr. Ind. 256. 1905 — Brasilia. — Phegopteris Fée, Cr. vasc. Br. **2**. 60. t. 98 f. 1. 1872-73; Polypodium Bak. 1891.

(D) **cana** (J. Sm.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Sikkim. Lastrea J. Sm. 1857; Nephrodium Bak. Syn. 267. 1867 (non Aspidium Wall. 1828, nomen). Lastrea Bergiana Bedd. 1876 (non Moore 1858).

(D) **Cañadasii** (Sod.) C. Chr. Ind. 256. 1905 — Ecuador. Nephrodium Sod. Rec. 48. 1833; Cr. vasc. quit. 236. 1893.

(C) **canescens** (Bl.) C. Chr. Ind. 256. 1905 — Java. Celebes. Ins. Philipp. Fiji. Polypodium Bl. Enum. 133. 1828; Gymnogramma Bl. 1828; Goniopteris Pr. 1836; Phegopteris Mett. 1864; Nephrodium Christ 1895. NPfl. 178; Aspidium Chr. 244. 1897; Nephrodium Blumei J. Sm. 1841 (non Hk. 1862); Aspidium Kze.; Mett. 1856. n. 236; Pleocnemia Moore 1857.

—* Nephrodium simplicifolium J. Sm. 1841. NPfl. 178; Abacopteris Fée 1850-52; Aspidium Hk. 1854; Polypodium Hk. 1863. HB. 313; Goniopteris Carr. 1873 (cum var. Fiji).

—* Haplodictyum heterophyllum Pr. 1849; Aspidium Hk. 1854; Nephrodium Hk. 1862. HB 295; Dryopteris O. Ktze. 1891.

—* Goniopteris asymetrica Fée 1850-52.

—* ? Nephrodium acrostichoides J. Sm. 1841; Cyclodium J. Sm. 1842; Pronephrium Pr. 1849.

—* Nephrodium Bakeri Harr. 1877; Bak. Ic. pl. t. 1664. NPfl. 178. Conf. Christ, Ann. Jard. Buit. **15**. 130.

capensis O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum adiantiforme.

(D) **caribaea** (Jenm.) C. Chr. Ind. 257. 1905 — Jamaica.
Nephrodium Jenm. **1886**. 270.

caripensis O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. submarginalis.

(P) **Carrii** (Bak.) C. Chr. Ind. 257. 1905 — Brasilia. — Polypodium Bak. Fl. bras. 1^2. 597. 1870. HB. ed. II. 305; Nephrodium Diels 1899. NPfl. 170.

caryotidea Und.; Heller, Minn. bot. St. **1**. 779. 1897 = Polystichum falcatum*.

(D) **catocarpa** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Venezuela.
Aspidium Linn. **9**. 95. 1834. Mett. n. 174; Nephrodium Hk. 1862. HB. 285.

catoptera O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. lanuginosa.

(D) **caucaensis** (Hier.) C. Chr. Ind. 257. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. **34**. 444. 1904.

(P) **caudata** (Raddi) C. Chr. Ind. 257. 1905 — Cuba-Brasilia.
Polypodium Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 25 t. 39. 1825; Klf. Enum. **113**. 1824. HB. 309; Phegopteris Fée 1850-52. Mett. n. 56. Chr. 276; Nephrodium Diels 1899. NPfl. 170. Polypodium pedicularifolium Pr. 1822 (non Hoffm. 1795).

(D) **Cavalerii** (Christ) C. Chr. Ind. 257. 1905 — China.
Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 116.

(M) **Cesatiana** C. Chr. Ind. 257. 1905 — N. Guinea. Fiji. — Meniscium Beccarianum Ces. Rend. Ac. Napoli **16**. 27, 30. 1877 (non Nephrodium Ces. 1876).

(D) **chaerophyllifolia** (Zippel) C. Chr. Ind. 257. 1905 — Java. Aneiteum.
Aspidium Zippel; Mett. Ann. Lugd. Bat. **1**. 228. 1864.

(D) **chartacea** C. Chr. Ind. 257. 1905 — Jamaica. — Nephrodium paucijugum Jenm. JoB. **1886**. 270 (non Aspidium Kl. 1847).

(D) **cheilanthoides** (Kze.) C. Chr. Ind. 257. 1905 — Brasilia.
Aspidium Kze. Linn. **22**. 578. 1849. Mett. n. 203; Lastrea Moore 1858.

(D) **cheiloplotia** (Fée) C. Chr. Ind. 257. 1905 — Mexico.
Aspidium Fée, 8 mém. 103. 1857; Lastrea Moore 1858.

(P) **chiapasensis** (Christ) C. Chr. Ind. 257. 1905 — Mexico.
Aspidium Christ, Bull. Boiss. II. **5**. 727. 1905.

(D) **chlorophylla** C. Chr. Ind. 257. 1905 — Mexico. — Aspidium pallidum Fourn. Mex. pl. **1**. 96. 1872 (non Bl. 1828 nec Link 1841).

(D) **chontalensis** (Fourn.) C. Chr. Ind. 257. 1905 — Nicaragua.
Aspidium Fourn. Bull. Soc. Fr. **19**. 254. 1872; Nephrodium Bak. 1891.

(M) **Christii** C. Chr. Ind. 257. 1905 — Ecuador. — Meniscium opacum Bak. JoB. **1877**. 166; Phegopteris Christ 270. 1897 (non Dryopteris infra).

(D) **chrysocoma** (Christ) C. Chr. Ind. 257. 1905 — China merid.
Aspidium filix mas var. Chrysocoma Christ, Bull. Boiss. **6**. 966. 1898: A. Chrysocoma Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 253.

(D) **chrysoloba** (Klf.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Brasilia. Columbia.
Aspidium Klf.; Link, Hort. Berol. **2**. 117. 1833. Mett. n. 219; Lastrea Pr. 1836; Nephrodium Fée 1850-52. HB. 262.

(D) **chrysotricha** (Bak.) C. Chr. Ind. 257. 1905 — Samoa.
Nephrodium Bak. Ann. Bot. **5**. 328. 1891; Aspidium Christ 1896.

cicutaria O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(D) **cirrhosa** (Schum.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Africa trop.
Aspidium Schum. Vid. Selsk. Skr. **4**. 231 (seors. 457). 1827; Nephrodium Bak. 1874. HB. 263. NPfl. 172. N. crinibulbon Hk. 1862; Lastrea J. Sm. 1875.

(D) **Clarkei** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Sikkim.
Nephrodium Bak. Syn. 497. 1874 — (an D. filix mas var.?).

(G) **coalescens** (Bak.) C. Chr. Ind. 258. 1905 — Ecuador.
Polypodium Bak. 1877. 164.

(D) **coarctata** (Kze.) C. Chr. Ind. 258. 1905 — Caracas. Columbia.
Aspidium Kze. Bot. Zeit. 1845. 287. Mett. n. 182; Polypodium Kl. 1847. Lastrea Moore 1858; Nephrodium Hier. 1904.

(D) **cochleata** (Don) C. Chr. Ind. 258. 1905 — India bor. et austr. Malacca.
Nephrodium Don, Prod. Fl. Nepal. 6. 1825; Aspidium Spr. 1827; Lastrea Moore 1857; Nephrodium Filix-mas δ HB. 272; Aspidium Filix mas var. Chr. 257. Arthrobotrys avara Wall. 1828 — A. macrocarpa Wall. 1828; Lastrea Pr. 1836.

(D) **cognata** (Pr.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — St. Helena.
Lastrea Pr. Epim. 40. 1849; Aspidium Fée 1850-52. Mett. n. 145; Nephrodium Hk. 1862. HB. 279. NPfl. 174.

(D) **concinna** (Willd.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Mexico-Chile.
Polypodium Willd. sp. 5. 201. 1810; Phegopteris Fée 1850-52; Aspidium (Link 1833?) Mett. 1856. n. 183. (Chr. 252); Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 268. NPfl. 172.

confluens O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Aspidium Mülleri.

(D) **conformis** (Sod.) C. Chr. Ind. 258. 1905 — Ecuador.
Nephrodium Sod. Rec. 45. 1883; Cr. vasc. quit. 240. 1893.

coniifolia Und.; Heller, Minn. bot. St. 1. 778. 1897 = Polystichum carvifolium.

(P) **connexa** (Klf.) C. Chr. Ind. 258. 1905 — Brasilia. — Polypodium Klf. Enum. 120. 1824. HB. 312; Phegopteris Fée 1850-52. Mett. n. 61.

connexa O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = D. Martini.

contermina O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = D. opposita.

(D) **conterminoides** C. Chr. Ind. 258. 1905 — N. Guinea.
Nephrodium simulans Bak. JoB. 1890. 106 (non 1874 nec 1888).

(D) **corazonensis** (Bak.) C. Chr. Ind. 258. 1905 — Ecuador.
Nephrodium (carazunense) Bak. JoB. 1877. 163; Aspidium Chr. 254. 1897.

(D? P?) **cordata** (Fée) Urban, Symb. Ant. 4. 18. 1903 — Cuba. Puerto Rico. Mexico.
Phegopteris Fée, Gen. 244. 1850-52; 6 mém. 13 t. 6 f. 3; Polypodium Moore 1857 (non Desv. 1827 nec Kze. 1834); Nephrodium Diels 1899. NPfl. 168; Polypodium cubanum Bak. 1867. HB. 304; Aspidium reptans var. 1 cordata Mett. n. 237. 1858.

costalis O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = D. densisora.

(C) **costularis** (Bak.) C. Chr. Ind. 258. 1905 — Madagascar.
Nephrodium Bak. Journ. Linn. Soc. 16. 203. 1877; Aspidium Kuhn 1879.

craspedosora O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Polystichum.

(D) **crassifolia** (Bl.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Birma. Malesia.
Aspidium crassifolium Bl. Enum. 158. 1828; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 266. NPfl. 171. Lastrea lata J. Sm. 1841; Aspidium Kze.; Mett. n. 231. 1858. Lastrea nephrodioides Bedd. 1866 (non Moore 1858).

(D) **crassipes** (Sod.) C. Chr. Ind. 258. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 234. 1893.

(D) **Creaghii** (Bak.) C. Chr. Ind. 258. 1905 — Borneo.
Nephrodium Bak. Kew Bull. 1898. 230.

(D) **crenata** (Forsk.) O. Ktze. Rev. Gen. Pl. 2. 811. 1891 — China austr. India bor. Malacca. Réunion. Arabia. Abyssinia. Ins. Caboverd. — Polypodium Forsk.

Fl. Aeg.-Arab. 185. 1775; Aspidium Kuhn 1868. Chr. 262; Lastrea Bedd. 1876; Nephrodium Bak. 1877. NPfl. 175; Hypodematium Kuhn 1879. Aspidium odoratum Bory; Willd. 1810; Cystopteris Desv. 1827; Lastrea Wawra 1863; Nephrodium Bak. 1867. HB. 280. Aspidium eriocarpum Wall. 1828. Mett. n. 142; Lastrea Pr. 1836; Nephrodium Dcne. 1841. N. hirsutum Don 1825; Lastrea Moore 1857 — Hypodematium onustum Kze. 1833 — H. Rüppellianum Kze. 1840 — Aspidium chrysolepis Fée 1857 — Nephrodium Fordii Bak. 1889.

(D) **crenulaea** (Jenm.) C. Chr. Ind. 259. 1905 — Jamaica. Nephrodium Jenm. Bull. Dep. Jam. **3**. 68. 1896.

(P) **crenulato-serrulata** (Mak.) C. Chr. Ind. 259. 1905 — Japonia. Athyrium Mak. Bot. Mag. Tokyo **13**. 26. 1899; Phegopteris Mak. 1903.

(P) **crinalis** (Hk. et Arn.) C. Chr. Ind. 259. 1905 — Ins. Hawaii. — Polypodium Hk. et Arn. Bot. Beech. Voy. 105. 1832. HB. 311; Phegopteris Mann 1868.

(C) **crinipes** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India bor. Malacca. Nephrodium Hk. sp. **4**. 71. 1862. HB. 294.

(D) **crinita** (Poir.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ins. afric. orient. Polypodium Poir. Enc. **5**. 528. 1804; Nephrodium Desv. 1827. HB. 265; Aspidium Wall. 1828. Chr. 254; Phegopteris Fée 1850-52; Lastrea Moore 1857 — Aspidium nitidum Bory; Willd. 1810; Nephrodium Desv. 1827; Lastrea Pr. 1836 — Aspidium pauciflorum Klf.; Spr. 1827. Mett. n. 215; Lastrea Pr. 1836; Nephrodium Kiærsk. 1874 — Aspidium sulcatum Klf.; Bory 1833. Mett. n. 216; Lastrea Pr. 1836; Nephrodium Kiærsk. 1874 — Aspidium strigosum Mett. n. 217. 1858 (non Willd. 1810). Alsophila erubescens Kze. 1844.

(D) **cristata** (L) Gray, Man. ed. I. 631. 1848 — Europa bor. et centr. Caucasus. Sibiria occ. Amer. bor. orient. — Polypodium L. sp. **2**. 1090. 1753; Polystichum Roth 1799; Aspidium Sw. 1801. Mett. n. 132. Chr. 258; Nephrodium Michx. 1803. HB. 273 (excl. β). NPfl. 174; Lastrea Pr. 1836; Polypodium Callipteris Ehrh. 1784; Polystichum Bernh. 1799 (pt.); Lastrea Newm. 1844; Lophodium Newm. 1851.

(D) **cristata × marginalis** Dav. Bot. Gaz. **19**. 497. 1894 (syn.) — New England. New Jersey. — Aspidium Dav. Bot. Gaz. **19**. 494. 1894.

(D) **cristata × spinulosa** (Milde) C. Chr. Ind. 259. 1905 — Europa centr. Sibiria. ? Alaska. ? Amer. bor. orient. (D. Boottii). — Aspidium Milde, Nova Acta **26**. 533. 1858; A. spinulosum × cristatum Lasch, A Br. et Milde, 33 Jahresber. Schles. Ges. **1855**. 94; ? Polystichum Sanio 1884; Lastrea uliginosa Newm. 1849; Lophodium Newm. 1851; Aspidium Nyman 1884; Nephrodium cristatum β HB. 273 — ? Aspidium Boottii Tuck. 1843. (Chr. 261); Polystichum Gandog. 1891; Dryopteris Und. 1893; Nephrodium spinulosum δ Boottii Bak. 1867. HB. 275.

(P) **Crossii** (Bak.) C. Chr. Ind. 259. 1905 — Andes loxenses. Polypodium Bak. Ann. Bot. **5**. 455. 1891.

(P) **cruciata** (Willd.) C. Chr. Ind. 259. 1905 — Ins. Mascarenae. Aspidium Willd. sp. **5**. 278. 1810; Lastrea Pr. 1836; Phegopteris Mett.; Kuhn 1868; Polypodium Bak. 1877. P. sessilifolium Hk. 1862; P. Bojeri Hk. 1862. HB. 311. Cystopteris villosa Desv. 1827.

(D) **crystallina** (Kze.) Maxon in C. Chr. Ind. 259. 1905 — India occ. Polypodium Kze. Farnkr. **2**. 85 t. 135. 1851; (Aspidium Mett. 1856); Lastrea Moore 1858.

(D) **ctenitis** (Link) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Brasilia.
Aspidium Link, Hort. Berol. 2. 122. 1833. Mett. n. 222; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 265. NPfl. 171.

(P) **ctenoides** (Fée) C. Chr. Ind. 260. 1905 — Jamaica. Guadeloupe.
Phegopteris Fée, 11 mém. 54 t. 14 f. 2. 1866; Polypodium Jenm. 1897.

(D) **cubensis** (Kuhn) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Cuba.
Aspidium Kuhn, Linn. 36. 108. 1869; Nephrodium Bak. 1874. HB. 501.

(C) **Cumingiana** (Kze.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Panama.
Aspidium Kze. Farnkr. 1. 17 t. 9 f. 2. 1840. Mett. n. 234. (Chr. 244); Nephrodium J. Sm. 1854. HB. 287.

(D) **Currori** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Africa occ. trop.
Aspidium Mett.; Kuhn, Fil. Afr. 130. 1868; Nephrodium Bak. 1874. HB. 500.

(P) **cyatheifolia** (Desv.) C. Chr. Ind. 260. 1905 — Ins. Mascarenae.
Polypodium Desv. Prod. 239. 1827. HB. ed. II. 306; Phegopteris Mett.; Kuhn 1868; Polypodium Sieberianum Klf.; Spr. 1827. HB. ed. I. 306; Phegopteris Fée 1850-52. Mett. n. 39.

(C) **cyatheoides** (Klf.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Ins. Hawaii. N. Guinea. Sumatra. — Aspidium Klf. Enum. 234. 1824. Mett. n. 263; Nephrodium Pr. 1836. HB. 293. NPfl. 179. Aspidium Dubrueillianum Gaud. 1827; Polystichum Gaud. 1827; Nephrodium Hk. et Arn. 1832 — Aspidium Boydiae Eat. 1879.

(D) **cycadina** (Franch. et Sav.) C. Chr. Ind. 260. 1905 — Japonia.
Aspidium Franch. et Sav. En. pl. Jap. 2. 236, 630. 1879.

(P) **cyclocolpa** (Christ) C. Chr. Ind. 260. 1905 — Costa Rica.
Polypodium Christ, Bull. Boiss. 4. 659. 1896.

(D) **cystolepidota** (Miq.) C. Chr. Ind. 260. 1905 — Japonia.
Aspidium Miq. Ann. Lugd. Bat. 3. 177. 1867.

cystostegia O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Polystichum.

(L) **dasyphylla** C. Chr. Ind. 260. 1905 — Brasilia. — Gymnogramme villosa Link, Hort. Berol. 2. 51. 1833. HB. 376 (non Dryopteris infra); Grammitis Pr. 1836; Leptogramma J. Sm. 1841; Phegopteris Mett. 1856. n. 32.

(D) **davallioides** (Brack.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Polynesia.
Lastrea Brack. Expl. Exp. 16. 202. 1854; Nephrodium Bak. 1867. HB. 287; Aspidium Kuhn 1869 (non Sw. 1801); Polystichum Diels 1899. NPfl. 193. — (f. Kirk = D. glabella).

(D) **Davenportii** C. Chr. Ind. 260. 1905 — Mexico. — Aspidium scabriusculum Dav. Bot. Gaz. 21. 255. 1896 (non Mett.; Salom. 1883); Dryopteris Dav. 1896; Lastrea Dav. 1896 (non Pr. 1849); Nephrodium Dav. 1896.

(D) **Dayi** (Bedd.) C. Chr. Ind. 260. 1905 — Perak. — Nephrodium Bedd. JoB. 1887. 323; Aspidium Bedd. 1888; Lastrea Bedd. 1892.

(G) **debilis** (Mett.) C. Chr. Ind. 260. 1905 — Amboina.
Phegopteris Mett. Ann. Lugd. Bat. 1. 223 t. 6 f. 1. 1864; Polypodium Bak. 1867. HB. 313; Nephrodium Diels 1899. NPfl. 178.

(C) **decadens** (Bak.) C. Chr. Ind. 260. 1905 — Fiji.
Nephrodium Bak. JoB. 1886. 183.

(D) **decipiens** (Hk.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — China. Liu Kiu.
Nephrodium Hk. sp. 4. 86 t. 243. 1862. HB. 260. NPfl. 168; Aspidium Luerss. 1883.

(D) **decomposita** (R. Br.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891. — Australia. Tasmania. Ins. Norfolk. N. Zealand. ? Africa austr.
Nephrodium R. Br. Prod. Fl. N. Holl. 149. 1810. HB. 281. NPfl. 175;

Aspidium Spr. 1827. Mett. n. 167. Chr. 263; Lastrea J. Sm. 1846; Polystichum Keys. 1873. Nephrodium pentangularum Col. 1844 — ? Aspidium quinquangulare Kze. 1850; Lastrea J. Sm. 1857 (Africa austr.).
— ? Deparia nephrodioides Bak. 1872. HB. 463; Davallia F. Muell. 1877; Dicksonia F. Muell. 1882 (Ins. Lord Howe).
— Conf. Kirk: Tr. N. Zeal. Inst. **10**. 398. — conf. D. glabella.

(D) **decrescens** (Kze.) O. Otze. Rev. Gen. Pl. **2**. 812. 1891 — Columbia. Venezuela. Aspidium Kze.; Mett. Pheg. u. Asp. 84 n. 202. 1858; Lastrea Moore 1858; Nephrodium Bak. 1874. HB. 497.

(D) **decumbens** C. Chr. Ind. 261. 1905 — Guadeloupe. — Aspidium asperulum Fée, 11 mém. 80 t. 23 f. 1. 1866. (HB. 266) (non Dryopteris supra).

decurrens O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(L) **decurrenti-alata** (Hk.) C. Chr. Ind. 261. 1905 — Japonia. — Gymnogramme Hk. sp. **5**. 142 t. 294. 1864. HB. 377; Leptogramma J. Sm. 1875; Phegopteris Christ 274. 1897; Nephrodium Diels 1899. NPfl. 171.

(D) **decursive-pinnata** (van Hall) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Asia orient. Polypodium van Hall, Nieuwe Verhdl. v. d. Nederl. Inst. **5**. 204 c. tab. 1836; Aspidium Kze. 1844. Mett. n. 177; Phegopteris Fée 1850-52. Chr. 275; Lastrea J. Sm. 1866; Nephrodium Bak. 1867. HB. 259. NPfl. 170.

(P) **decussata** (L.) Urban, Symb. Ant. **4**. 19. 1903 — Amer. trop. Polypodium L. sp. **2**. 1093. 1753. HB. 307; Glaphyropteris Pr. 1848; Phegopteris Mett. 1856. n. 29. Chr. 273; Gymnogramme Fée 1857; Nephrodium Diels 1899. NPfl. 171 — Polypodium velutinum Sod. 1883.

(D) **deflexa** (Klf.) C. Chr. Ind. 261. 1905 — Brasilia austr. — Polypodium Klf. Enum. 114. 1824. P. vestitum Raddi 1819 (non Forst. 1786); Lastrea J. Sm. 1846; Nephrodium Bak. 1867. HB. 265. NPfl. 171; Aspidium Fée 1869; Dryopteris O. Ktze. 1891; Aspidium Raddianum Mett. n. 221. 1858; Nephrodium Hk. 1862. ? Polypodium lepigerum Schrad. 1824.

(C) **dejecta** (Jenm.) C. Chr. Ind. 261. 1905 — Demerara. Nephrodium Jenm. Gard. Chr. III. **18**. 640. 1895.

(D) **deltoidea** (Sw.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India occ. Polypodium Sw. Prod. 133. 1788; Aspidium Sw. 1801. Mett. n. 225. Chr. 244; Nephrodium Desv. 1827. HB. 270. NPfl. 172; Lastrea Moore 1858.

(P) **demerarana** (Bak.) C. Chr. Ind. 261. 1905 — Demerara. Polypodium Bak. Tr. Linn. Soc. II. Bot. **2**. 290. 1887.

(D) **densiloba** C. Chr. Ind. 261. 1905 — Brasilia. Aspidium Gardnerianum Kze. msc. (non Mett. 1858); Nephrodium Bak. Fl. bras. 1^2. 474. 1870. HB. 496; Dryopteris O. Ktze. 1891.

(D) **densisora** C. Chr. Ind. 261. 1905 — Venezuela. Aspidium costale Mett.; Kuhn, Linn. **36**. 111. 1869 (non Bl. 1828); Nephrodium Bak. 1874. HB. 495; Dryopteris O. Ktze. 1891.

denticulata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(D) **deparioides** (Moore) O. Ktze. Rev. Gen. Pl **2**. 812. 1891 — India austr. Ceylon. Diclisodon Moore, Ind. XCVI. 1857; Aspidium Hk. 1858. Chr. 262; Nephrodium Hk. 1859, 1862. HB. 277. NPfl. 174; Lastrea Bedd. 1863-65. —* Aspidium concinnum Thwait. 1864 (non Link 1833); Lastrea Bedd. 1863-65; Nephrodium Thwaitesii Bak. 1867. HB. 277; Lastrea Bedd. 1876; Dryopteris O. Ktze. 1891.

devexa O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = Aspidium.

(C) **devolvens** (Bak.) C. Chr. Ind. 261. 1905 — Brasilia. Nephrodium Bak. JoB. **1885**. 217.

(P) **Dianae** (Hk.) C. Chr. Ind. 262. 1905 — St. Helena.
Polypodium Hk. sp. 4. 234. 1862; P. mollis Roxb. 1816. HB. 308 (non Jacq. 1789); Phegopteris Kuhn 1868.

(C) **dicarpa** (Fée) C. Chr. Ind. 262. 1905 — Réunion.
Nephrodium Fée, Gen. 305. 1850-52.

(D) **Dickinsii** (Franch. et Sav.) C. Chr. Ind. 262. 1905 — Japonia. China.
Aspidium Franch. et Sav. Enum. pl. Jap. 2. 236, 629. 1879; Nephrodium Bak. 1886. NPfl. 168.

(C) **didymosora** (Parish) C. Chr. Ind. 262 1905 — Tenasserim. Perak. Singapore.
Nephrodium Parish; Bedd. Ferns br. Ind. t. 200. 1866; Aspidium tectum Wall. 1828 (nomen); Nephrodium Bedd. Handb. Suppl. 79. 1892.

(P) **Dielsii** C. Chr. Ind. 262. 1905 — China.
Polypodium stenopteron Bak. JoB. 1888. 229 (non 1886); Phegopteris Christ 1898; Nephrodium Diels 1899. NPfl. 170 (non Bak. 1890).

(D) **diffracta** (Bak.) C. Chr. Ind. 262. 1905 — Yunnan.
Nephrodium Bak. Kew Bull. 1898. 230; Aspidium Christ 1899.

dilatata Gray, Man. ed. I. 631. 1848 = D. spinulosa*.

(D) **dimidiata** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Africa occ. trop.
Aspidium Mett.; Kuhn, Fil. Afr. 131. 1868; Nephrodium Bak. 1874. HB. 500.

(L) **diplazioides** (Desv.) Urban, Symb. Ant. 4. 21. 1903 — Amer. trop.
Gymnogramma Desv. Prod. 214. 1827. HB. 377; Phegopteris Mett. 1864; Leptogramma Und. 1902; Nephrodium Hier. 1904 (non Hk. 1862) — Gymnogramme polypodioides Link 1833 (non Spr. 1827); Grammitis Linkiana Pr. 1836; Leptogramma J. Sm. 1841; Gymnogramme Kze. 1844; Phegopteris Mett. 1856. n. 26. Chr. 273; Nephrodium Diels 1899. NPfl. 172.

diplazioides O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = D. Moritziana.

(D) **dissecta** (Forst.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Ind. bor.-Malesia-Polynesia-Austr. trop. Madagascar. — Polypodium Forst. Prod. 81. 1786 (non Sw. 1788); Polystichum Bernh. 1802; Nephrodium Desv. 1827. HB. 282. NPfl. 175; Lastrea Carr. 1873; Aspidium Christ 1896 (non Mett. 1864); Polypodium laciniatum Gmel. 1791; Aspidium attenuatum Sw. 1801; Nephrodium membranifolium Hk. 1862 (pt.). HB. ed I. 282 (non Pr. 1825) — N. ingens Atk.; Clarke 1880 — ? Aspidium Gardnerianum Mett. n. 170. 1858; Lastrea Moore 1858.

dissidens O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Aspidium.

(P) **dissitifolia** (Bak.) C. Chr. Ind. 262. 1905 — Yunnan.
Polypodium Bak. Kew Bull. 1895. 54.

(C) **distans** (Hk.) O. Ktze. Rev. Gen. Pl. 2 812. 1891 — Ins. Comorae. Madagascar.
Nephrodium Hk. sp. 4. 76 in adnota 1862. HB. 288; Aspidium Kuhn 1868.

divisa O. Ktze. Rev. Gen. Pl. 2. 811. 1891 = D. Boryana.

(P) **Drakei** C. Chr. Ind. 262. 1905 — Tahiti. — Polypodium Vescoi Drake. Fl. Polyn. franç. 307. 1893 (non Nephrodium Drake l. c. 301).

(D) **Eatoni** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Liu Kiu. Kakeah.
Nephrodium Bak. Syn. 276. 1867.

(D) **echinata** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Borneo. Celebes. N. Guinea. — Aspidium Mett. Ann. Lugd. Bat. 1. 230. 1864; Nephrodium Bak. 1867. HB. 266. NPfl. 171.

(D) **edentula** (Kze.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Java.
Aspidium Kze. Bot. Zeit. 1846. 474. Mett. n. 146; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 279. NPfl. 174.

(D) **effusa** (Sw.) Urban. Symb. Antill. **4**. 16. 1903 — Amer. trop.
Polypodium Sw. Prod. 134. 1788; Phegopteris Fée 1850-52; Aspidium Gris. 1864. Chr. 263; Nephrodium Bak. 1867. HB. 287; Polystichum Keys. 1873. NPfl. 193. Polypodium multifidum Jacq. 1789; Dryopteris O. Ktze. 1891. Polypodium divergens Willd.; Schkuhr 1806; Phegopteris Fée 1850-52. Mett. n. 16 — Nephrodium xanthotrichium Sod. 1883. —* Aspidium excultum Mett. n. 162. 1858; Lastrea Moore 1858; Nephrodium Hk. 1862.

(P) **efulensis** (Bak.) C. Chr. Ind. 263. 1905 — Kamerun.
Polypodium Bak. Kew Bull. **1897**. 299.

(D) **Eggersii** (Hier) C. Chr. Ind. 263. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. **34**. 441. 1904.

(C) **elata** (Mett.) C. Chr. Ind. 263. 1905 — Guinea cum insulis. — Aspidium Mett.; Kuhn 1868 (non Boj. 1837, nomen); Nephrodium venulosum Hk. sp. **4**. 71. 1862. HB. 291 (non Desv. 1827); Dryopteris O. Ktze. 1891.

elata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium chimborazense.

(D) **elegantula** (Sod.) C. Chr. Ind. 263. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 243. 1893.

elongata O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = D. khasiana.

(D) **Elwesii** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Sikkim.
Nephrodium Bak. Syn. 497. 1874; Lastrea Bedd. 1876.

(D) **eminens** (Bak.) C. Chr. Ind. 263. 1905 — Sumatra.
Nephrodium Bak. JoB. **1880**. 213.

(D) **enneaphylla** (Bak.) C. Chr. Ind. 263. 1905 — China occ.
Nephrodium Bak. JoB. **1887**. 170. NPfl. 168.

(D) **eriocaulis** (Fée) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Brasilia.
Aspidium Fée, Cr. vasc. Br. **1**. 136 t. 44 f. 1. 1869; Nephrodium Bak. 1874. HB. 495; N. ramentaceum Bak. 1870.

(D) **eriosora** (Fée) C. Chr. Ind. 263. 1905 — Brasilia.
Aspidium Fée, Cr. vasc. Br. **2**. 73 t. 101. 1872-73.

(P) **erubescens** (Wall.) C. Chr. Ind. 263. 1905 — China austr. India bor. Malesia.
Polypodium Wall. 1828 (nomen); Hk. sp. **4**. 236. 1862. HB. 306; Glaphyropteris Fée 1872-73; Phegopteris J. Sm. 1875. Chr. 272; Nephrodium Diels 1899. NPfl. 171; Aspidium Christ 1902.

(D) **erythrosora** (Eat.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — China. Japonia. Luzon. — Aspidium Eat. in Perry, Narr. Exp. to China **2**. 230. 1856. (Chr. 257); Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 273. NPfl. 173 — Aspidium prolificum Maxim. 1879; Lastrea Moore 1882; Nephrodium Diels 1899. NPfl. 173. (f. monstr.).

(D) **erythrostemma** (Christ) C. Chr. Ind. 263. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. II. **4**. 961. 1904.

(P) **euchlora** (Sod.) C. Chr. Ind. 263. 1905 — Ecuador.
Polypodium Sod. Rec. 58. 1883; Cr. vasc. quit. 290. 1893.

(C) **eurostotricha** (Bak.) C. Chr. Ind. 263. 1905 — Madagascar.
Nephrodium Bak. JoB. **1880**. 329.

(C) **evoluta** (Clarke) C. Chr. Ind. 263. 1905 — India bor.
Nephrodium amboinense var. evolutum Clarke, Journ. Linn. Soc. **24**. 417. 1888; N. evolutum Bedd. Handb. Suppl. 76. 1892 (excl. β).

excellens O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium.

(D) **excelsa** (Desv.) C. Chr. Ind. 264. 1905 — Mexico. Columbia. ? India occ. Polypodium Desv. Prod. 243. 1827 (non Bak. 1874). Aspidium equestre Kze. 1844. Mett. n. 175 (t. Fourn.); Lastrea Liebm. 1849 — L. ciliata Liebm. 1849; Nephrodium Kiærsk. 1874 (non Desv. 1827); Aspidium Mett.; Salom. 1883.

exigua O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. philippina.

(C) **extensa** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India austr. Ceylon. Birma-Malesia-Ins. Philipp. — Aspidium Bl. Enum. 156. 1828. Chr. 247; Nephrodium Moore 1858. HB. 289. NPfl. 178. N. conionevron Fée 1850-52; Aspidium Mett. n. 243. 1858.

(D) **Faberi** (Bak.) C. Chr. Ind. 264. 1905 — China. Nephrodium Bak. Ann. Bot. **5**. 316. 1891. NPfl. 168.

falcata O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(D) **falciculata** (Raddi) O. Ktze. Rev. Gen. Pl. **3**². 378. 1898 — Brasilia. Aspidium Raddi, Opusc. sci. Bol. **3**. 289. 1819; Pl. Bras. **1**. 31 t. 47. 1825. Chr. 252; Nephrodium Desv. 1827. HB. 264. NPfl. 171; Polystichum Gaud. 1827; Lastrea Pr. 1836 — ? Polypodium falciculatum Raddi 1819; Phegopteris Fée 1869.

falcinella O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

Falconeri O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. barbigera.

(D) **Fargesii** (Christ) C. Chr. Ind. 264. 1905 — China merid. Aspidium Christ, Bull. Soc. Fr. **52** Mém. I. 42. 1905.

(D) **fasciculata** (Fourn.) C. Chr. Ind. 264. 1905 — N. Caledonia. Aspidium Fourn. Ann. sc. nat. V. **18**. 295. 1873; Nephrodium Bak. 1891.

(D) **Feei** C. Chr. Ind. 264. 1905 — Mexico. — Aspidium puberulum Fée 10 mém. 40. 1865 (non Desv. 1827 nec Gaud. 1827); Nephrodium Bak. 1874. HB. 495; Dryopteris O. Ktze. 1891.

(C) **Fendleri** (Eat.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Venezuela. Aspidium Eat. Mem. Amer. Ac. n. s. **8**. 210. 1860; Nephrodium Hk. 1862. HB. 295. — (f. Jenm. = D. venusta).

(C) **ferox** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — ? Kumaon. Perak. Malesia-Ins. Philipp. — Aspidium Bl. Enum. 153. 1828. Chr. 247; Polypodium Hassk. 1844; Nephrodium Moore 1858. HB. 294. NPfl. 180; Phegopteris Mett. 1864; Goniopteris barbata Fée 1850-52; Phegopteris Mett. n. 48. 1858; Polypodium Hk. 1863 (non Kze. 1827).

ferruginea O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. peranemiformis.

(P) **fibrillosa** (Bak.) C. Chr. Ind. 264. 1905 — Peru. Polypodium Bak. Syn. 307. 1867.

(D) **fijiensis** (Hk.) C. Chr. Ind. 264. 1905 — Fiji. Nephrodium Hk. 2 Cent. t. 67. 1861. (HB. 280); Lastrea Carr. 1873.

(D) **filix mas** (L.) Schott, Gen. Fil. ad t. 9. 1834 — Europa: Island-Algérie et Madeira. Asia minor. Asia septent. et temp.-Himalaya-Java. Ins. Mascarenae. Madagascar. Ins. Amsterdam. Amer. bor.-Mexico-Peru. Brasilia. Jamaica. Ins. Hawaii. — Polypodium L. sp. **2**. 1090. 1753; Polystichum Roth 1799; Aspidium Sw. 1801. Mett. n. 129. Chr. 256; Nephrodium Rich. 1801. HB. 272, 498. NPfl. 173; Tectaria Cav. 1802; Lastrea Pr. 1836; Lophodium Newm. 1851.

—* Polypodium elongatum Ait. 1789; Aspidium Sw. 1801. Mett. n. 148; Tectaria Cav. 1802; Nephrodium Hk. et Grev. 1831; Lastrea Pr. 1836 — Nephrodium Boryanum Hk. 1862 (non Aspidium Willd. 1810);

N. Bojeri Bak. 1867. HB. 280. NPfl. 175; Dryopteris O. Ktze. 1891 — Ins. Atlant. Ins. Mascar.

—* Aspidium oligodonton Desv. Berl. Mag. **5**. 322. 1811; Nephrodium Desv. 1827; Aspidium elongatum Willd. 1810 (non Sw. 1801). A. canariense A. Br. 1841. Mett. n. 150; Lastrea Moore 1857; Polystichum Keys. 1873; Nephrodium Kiærsk. 1874. Aspidium aquilinoides Bolle 1866 (non Dryopteris supra) — Ins. Atlant. Ins. Mascar. ? India. — (conf. Kuhn, v. Deck. Reis. **3**[3]. Bot. 45).

—* Aspidium paleaceum Sw. Syn. 52. 1806; Lastrea Moore 1858 — Aspidium parallelogrammum Kze. 1839; Lastrea Liebm. 1849; Dichasium Fée 1850-52. Lastrea truncata Brack. 1854 — Mexico-Peru. Ins. Hawaii. (? Asia trop.).

—* Aspidium Ludovicianum Kze. Amer. Journ. Sci. II. **6**. 93. 1848 — U. S. A. (an eadem ac *oligodonta supra?).

—* Aspidium Hawaiiense Hill. Fl. Haw. 575. 1888 — Hawaii.

—* Nephrodium antarcticum Bak. Journ. Linn. Soc. **14**. 479. 1875; Aspidium Fourn. 1875 — Ins. Amsterdam.

Subspecies Asiae centralis.

—* Aspidium patentissimum Wall. 1828; Lastrea Pr. 1836; Dichasium Fée 1850-52; Nephrodium Clarke 1876. Aspidium paleaceum Don, Prod. Fl. Nepal. 4. 1825 (non Sw. 1806); A. Donianum Spr. 1827 — Nephrodium filix mas var. fibrillosa Clarke, Tr. Linn. Soc. II. Bot. **1**. 520 t. 70. 1880 — Himalaya.

—* Nephrodium filix mas var. khasiana Clarke, Tr. Linn. Soc. II. Bot. **1**. 519 t. 69 f. 1. 1880. Aspidium patentissimum Franch. 1887 (non Wall. 1828) — Himalaya. Thibet. China merid.

—* Nephrodium Kingii Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 621 t. 9. 1899 (non Dryopteris infra) — India bor.

—* Nephrodium filix mas var. nidus Clarke; Bak. Syn. 498. 1874; (Bedd. Ferns br. Ind. Suppl. 17 t. 372. 1876); N. nidus Clarke 1876 — Sikkim.

—* Nephrodium filix mas var. panda Clarke, Tr. Linn. Soc. II. Bot. **1**. 519 t. 68 f. 1. 1880; N. pandum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 623. 1899 — Himalaya. China merid.

—* Lastrea filix mas var. serrato-dentata Bedd. Handb. Suppl. 55. 1892; Nephrodium serrato-dentatum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 622 t. 10. 1899; N. filix mas var. odontoloma Bak. Syn. 498. 1874; N. odontoloma Clarke, Tr. Linn. Soc. II. Bot. **1**. 521. 1880; Lastrea Bedd. 1883 (non Moore 1858; Bedd. 1863-65) — India bor.

—* Lastrea odontoloma Moore 1858; Bedd. Ferns S. Ind. 39 t. 114. 1863-65; Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 736 t. 31. 1903; N. filix mas var. 2 normalis Clarke, Tr. Linn. Soc. II. Bot. **1**. 521 t. 71. 1880 — Himalaya. India austr.

—* Nephrodium filix mas var. Schimperiana Clarke, Tr. Linn. Soc. II. Bot. **1**. 520 t. 69 f. 2. 1880; Nephrodium Schimperianum Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 733. 1903 (vix eadem ac Dryopteris Schimperiana infra); Lastrea intermedia Bedd. 1863-65 (non Pr. 1836) — India bor. et austr.

—* Aspidium marginatum Wall. 1828; Clarke, Tr. Linn. Soc. II. Bot. **1**. 521 t. 71. 1880 (var.); Nephrodium Hope, Journ. Bomb. Nat. Hist.

Soc. **14**. 740 t. 33. 1903; Nephrodium vel Lastrea filix mas var. elongata auctt. quoad pl. ind. — India bor. China merid.

—* Nephrodium assamensis Hope, JoB. **1890**. 326 — Assam.

—* Aspidium oxyodon Franch. Nouv. Arch. Mus. II. **7**. 163. 1883 — China.

—* Aspidium adnatum Bl. Enum. 162. 1828 — Java.

— *Obs.* conf. D. apiciflora, chrysocoma, Clarkei, erythrosora, lacera, lunanensis, subarborea, subtriangularis, quae t. auctt. plur. subspecies vel varietates D. filicis maris sunt.

(D) **filix mas × spinulosa** (A.Br.) C. Chr. Ind. 266. 1905 — Europa centr. England. Aspidium A.Br. in Döll, Fl. Bad. 30. 1857; A. rigidum var. remotum A.Br. 1843. Chr. 261; A. remotum A.Br. 1850. Mett. n. 135; Lastrea Moore 1858; Nephrodium Moore 1861; N. spinulosum γ Bak. 1867. HB. 275.

(D) **firma** (Bak.) C. Chr. Ind. 266. 1905 — Jamaica. Nephrodium Bak.; Jenm. JoB. **1879**. 260.

(D) **firmifolia** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Peru. Nephrodium Bak. Syn. 501. 1874; N. acutum Hk. 1862 (pt. t. 271).

(G) **firmula** (Bak.) C. Chr. Ind. 266. 1905 — Borneo. Polypodium Bak. Kew Bull. **1893**. 211.

(D) **Fischeri** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Brasilia. Aspidium Mett. Pheg. u. Asp. 80 n. 192. 1858; Nephrodium Bak. 1870. HB. 476; Aspidium obscurum Fisch. et Mey. 1833 (v. pag. 85 supra; non Bl. 1828); Lastrea Moore 1858.

(D) **flaccida** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — China austr. India bor.-Ceylon. Java. — Aspidium Bl. Enum. 161. 1828; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 274. NPfl. 175.

flexa O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum Berterianum.

(D) **flexilis** (Christ) C. Chr. Ind. 266. 1905 — China. Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 252.

(D) **flexuosa** (Fée) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Brasilia. Aspidium Fée, Cr. vasc. Br. **1**. 138 t. 46 f. 2. 1869; Nephrodium Bak. Fl. bras. **1**[2]. 483. 1870. HB. 501.

(D) **floridana** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Florida. Alabama. Texas. — Nephrodium Hk. Fil. exot. t. 99. 1859. HB. 273. NPfl. 174; Aspidium Eat. 1860; Lastrea J. Sm. 1877; Aspidium cristatum var. Floridanum Christ 258. 1897.

foeniculacea O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Polystichum.

(D) **formosana** (Christ) C. Chr. Ind. 266. 1905 — Formosa. Aspidium Christ, Bull. Boiss. II. **4**. 615. 1904.

(D) **Fournieri** (Bak.) C. Chr. Ind. 266. 1905 — Mexico. Nephrodium Bak. 1891 (Ann. Bot. **5**. 317, non 328); Aspidium flaccidum Fourn. Bull. Soc. Fr. **27**. 328. 1880 (non Bl. 1828).

(D) **fragilis** (Bak.) C. Chr. Ind. 266. 1905 — Madagascar. Polypodium Bak. Journ. Linn. Soc. **16**. 203. 1877; Phegopteris Kuhn 1879; Nephrodium Bak. 1891.

(D) **fragrans** (L.) Schott, Gen. Fil. ad t. 9. 1834 — Asia bor. Japonia. Caucasus. Amer. arct.-Minnesota-New York. — Polypodium L. sp. **2**. 1089. 1753; Aspidium Sw. 1801. Mett. n. 133. Chr. 260; Nephrodium Richards. 1823. HB. 275. NPfl. 173; Lastrea Pr. 1836; Polystichum Ledeb. 1853.

(D) **Fraseri** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ecuador. — Aspidium Mett.; Kuhn, Linn. **36**. 109. 1869; Nephrodium Bak. 1874. HB. 495.

frondosa O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Polystichum Webbianum.

(D) **fructuosa** (Christ) C. Chr. Ind. 267. 1905 — Yunnan.
Aspidium Christ, Bull. Soc. Fr. 52. Mém. I. 38. 1905.

(D) **Funckii** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Columbia.
Aspidium Mett. Ann. sc. nat. V. 2. 246. 1864; Nephrodium (Funkii) Bak. 1874. HB. 496.

(D) **furcata** (Kl.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Columbia-Peru. Galapagos.
Aspidium Kl. Linn. 20. 371. 1847; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 285 — ? Polypodium paleaceum Hk.fil. 1847 (non Borck. 1798); Nephrodium Diels 1899. NPfl. 177. (Galapagos).

(D) **fusca** C. Chr. Ind. 267. 1905 — Ecuador. — Nephrodium polylepis Sod. Cr. vasc. quit. 260. 1893 (non Dryopteris infra).

(D) **Galanderi** (Hier.) C. Chr. Ind. 267. 1905 — Argentina.
Aspidium Hier. Engl. Jahrb. 22. 363. 1896.

(P) **Galeottii** (Mart.) C. Chr. Ind. 267. 1905 — Mexico. — Polypodium Martens, Mém. Ac. Brux. 15. 43 t. 7 f. 3. 1842; Phegopteris Fée 1850-52. Ph. stenolepis Fée 1857. Polypodium Sloanei Liebm. 1849 (non Spr. 1827).

(D) **Gamblei** (Hope) C. Chr. Ind. 267. 1905 — India bor.
Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. 12. 533 t. 7. 1899.

Gardneriana O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = D. densiloba.

(D) **geropogon** (Fée) C. Chr. Ind. 267. 1905 — Mexico. Cuba.
Aspidium Fée, 10 mém. app. 1865; Nephrodium Kiærsk. 1874.

(P) **Germaniana** (Fée) C. Chr. Ind. 267. 1905 — Guadeloupe. — Phegopteris Fée, 11 mém. 55 t. 13 f. 2. 1866; Polypodium Bak. 1867. HB. 306.

(G) **Ghiesbreghtii** (Linden) C. Chr. Ind. 267. 1905 — Mexico. Guatemala.
Polypodium Linden, Cat. 1858; Bak. Syn. 315. 1867 (Gheisbreghtii) Goniopteris (Gheisbeghtii) J. Sm. 1866; Fourn. 1872; Nephrodium Diels 1899. NPfl. 179; Aspidium (Geisbreghtii) Christ 1904. Goniopteris mollis Fée 1850-52 (non Aspidium Sw. 1801) (§ Microdictyon Fée 1866). Polypodium crenatum Hk. Fil. exot. t. 84. 1859 (non Forsk. 1775).

gigantea O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Aspidium.

(M) **gigantea** (Mett.) C. Chr. Ind. 267. 1905 — Columbia-Peru.
Meniscium Mett. Fil. Lechl. 1. 19. 1856. HB. 391; Phegopteris Mett. 1859. Chr. 269; Nephrodium Diels 1899. NPfl. 177 (non Bak. 1867. 1874).

(?) **Gilberti** Clute, Fern Bull. 8. 67. 1900 — Jamaica.

(D) **glabella** (A. Cunn.) C. Chr. Ind. 267. 1905 — N. Zealand. Australia. ? Polynesia.
Nephrodium A. Cunn. in Hk. Comp. Bot. Mag. 2. 367. 1836. NPfl. 175; Lastrea Moore et Houlst. (1851?); Brack. 1854; Aspidium Lowe 1857. Mett. n. 164; Nephrodium decompositum β HB. 281. — Conf. Kirk. Tr. N. Zeal. Inst. 10. 398. Vide D. davallioides.

(D) **glabra** (Brack.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Ins. Hawaii. Tahiti.
Lastrea Brack. Expl. Exp. 16. 200. 1854; Aspidium Mett. n. 140. 1858; Nephrodium Bak. 1867. HB. 278. NPfl. 174 — Phegopteris dicksonioides Mett. 1869.

(D) **glabrata** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 812. 1891 — Fernando Po.
Aspidium Mett.; Kuhn, Fil. Afr. 133. 1868; Nephrodium Bak. 1874. HB. 500.

(P) **glandulifera** (Liebm.) C. Chr. Ind. 267. 1905 — Mexico. — Polypodium Liebm. Vid. Selsk. Afh. V. 1. 206 (seors. 54). 1849. HB. 310; Phegopteris (glanduligera) Fourn. 1872; Nephrodium Kiærsk. 1874. NPfl. 175.

(C) **glandulosum** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Perak. Malesia. Ins. Philipp. — Aspidium Bl. Enum. 144. 1828. Mett. n. 265. Chr. 245; Cyclodium Pr. 1836; Abacopteris Fée 1850-52; Nephrodium Moore 1857. HB. 291 (pt.). NPfl. 179 (vix J. Sm. 1841). N. meniscioides Clarke 1880. — Conf. Mett. Ann. Lugd. Bat. **1**. 235; Clarke, Tr. Linn. Soc. II. Bot. **1**. 531; Bedd. Handb. Suppl. 70.

(C) **glaucostipes** (Bedd.) C. Chr. Ind. 268. 1905 — Perak.
Nephrodium Bedd. Handb. Suppl. 80. 1892.

(D) **Glaziovii** (Christ) C. Chr. Ind. 268. 1905 — Brasilia.
Aspidium Christ, Bull. Boiss. II. **2**. 633. 1902.

(D) **gleichenioides** (Christ) C. Chr. Ind. 268. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. II. **4**. 960. 1904.

(D) **globulifera** (Brack.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Ins. Hawaii.
Lastrea Brack. Expl. Exp. **16**. 194. 1854; Nephrodium Hk. 1862. HB. 269. NPfl. 172; Aspidium Mann 1868.

(D) **Goldiana** (Hk.) Gray, Man. ed. I. 631. 1848 — U. S. A. atlant.
Aspidium Hk.; Goldie, Edinb. Phil. Journ. **6**. 333. 1822. Mett. n. 131; Nephrodium Hk. et Grev. 1829. HB. 272. NPfl. 173; Lastrea Pr. 1836; Polystichum Keys. 1873; Aspidium Filix mas var. Goldieanum Chr. 258. A. americanum Torrey 1818 (nomen).

(C) **gongylodes** (Schkuhr) O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 — China austr. India bor.-Malesia-Polynesia-Australia-N. Zealand-Hawaii. Algeria. Africa trop. et austr. cum ins. Florida-Brasilia. — Aspidium (goggilodus) Schkuhr, Kr. Gew. **1**. 193 t. 33 c. 1809. Mett. n. 241; Polystichum Gaud. 1827, Cyclosorus Link 1833; Nephrodium Schott 1834. N. unitum R.Br. 1810. HB. 289. NPfl. 178; Aspidium Mett. 1864. Chr. 246; Dryopteris Maxon 1901 — Aspidium Ecklonii Kze. 1836. Mett. n. 240; Nephrodium Pr. 1849. —* Nephrodium propinquum R.Br. 1810; Polystichum Gaud. 1827; Aspidium Sw. 1829. Mett. n. 242; Dryopteris Gilb. 1896. Aspidium unitum Sw. 1801 (non Polypodium L. 1759); A. unitum β hirsuta Mett. 1864.

(P) **Gordoni** (Bak.) C. Chr. Ind. 268. 1905 — Fiji.
Polypodium Bak. JoB. **1879**. 297.

(P) **gracilenta** (Jenm.) C. Chr. Ind. 268. 1905 — Jamaica. Cuba.
Polypodium Jenm. Bull. Dep. Jam. **4**. 129. 1897.

(D) **gracilescens** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India austr. Perak. Malesia. — *Japonia. China. India bor. Java.
Aspidium Bl. Enum. 155. 1828; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 262 — ? Lastrea hirsutipes Bedd. 1892.
—* Aspidium glanduligerum Kze. 1837. Mett. n. 207; Lastrea Moore 1858. — ? Aspidium angustifrons Miq. 1867.

(P) **gracilifrons** C. Chr. Ind. 268. 1905 — Japonia. — Polypodium leptophyllum Bak. Ann. Bot. **5**. 458. 1891 (non L. 1753 nec Moritz; Mett. 1864).

(D) **grammitoides** (Christ) C. Chr. Ind. 268. 1905 — Luzon.
Aspidium Christ, Bull. Boiss. **6**. 193. 1898.

(P) **grandis** (Pr.) C. Chr. Ind. 268. 1905 — Brasilia.
Polypodium Pr. Del. Prag. **1**. 171. 1822; Phegopteris Mett. 1870. Polypodium auriculatum Raddi 1819 (non L. 1753). P. splendidum Klf. 1824. HB. 312; Phegopteris Fée 1850-52. Mett. n. 60. Chr. 277. Polypodium formosum Raddi 1825; Phegopteris Fée 1850-52.

—* Polypodium macropterum Klf. 1824; Phegopteris Fée 1850-52. Alsophila Fischeriana Regel 1855; Polypodium Bak. 1868.

(G) **granulosa** (Pr.) C. Chr. Ind. 269. 1905 — Luzon. Timor.
Polypodium Pr. Rel. Haenk. **1**. 24 t. 4 f. 2. 1825; Goniopteris Pr. 1836; Nephrodium J. Sm. 1857.

(D) **grenadensis** (Jenm.) C. Chr. Ind. 269. 1905 — Grenada.
Nephrodium Jenm. Gard. Chr. III. **15**. 198. 1894.

Griffithii O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = Aspidium multicaudatum.

(D) **grisea** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — India austr.
Nephrodium Bak. Syn. 271. 1867; Lastrea Bedd. 1870.

Grisebachii O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 = D. ameristoneura.

(P) **grossa** (Christ) C. Chr. Ind. 269. 1905 — China austr.
Phegopteris Christ, Bull. Boiss. **7**. 13. 1899.

(C) **Grunowii** (Bolle) C. Chr. Ind. 269. 1905 — Ins. Caboverdicae.
Aspidium Bolle, Bonpl. **3**. 123. 1855; Nephrodium Moore 1858.

(C) **guadalupensis** (Fée) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Guadeloupe. Jamaica. Costa Rica. — Nephrodium Fée, 11 mém. 89 t. 24 f. 3. 1866. HB. 503; Aspidium Christ 1896.

(D) **guatemalensis** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Mexico. Guatemala.
Nephrodium Bak. Syn. 498. 1874.

(D) **Gueintziana** (Mett.) C. Chr. Ind. 269. 1905 — Africa trop. et austr. Réunion. St. Helena. — Aspidium Mett. Pheg. u. Asp. 83. n. 201. 1858. (HB. 269): Lastrea Moore 1858; Nephrodium Hier. 1900.

(C) **Gustavi** (Bedd.) C. Chr. Ind. 269. 1905 — India.
Nephrodium Bedd. JoB. **1893**. 227.

(P) **gymnogrammoides** (Bak.) C. Chr. Ind. 269. 1905 — China centr.
Polypodium Bak. JoB. **1888**. 229; Nephrodium Diels 1899. NPfl. 170.

(D) **gymnophylla** (Bak.) C. Chr. Ind. 269. 1905 — China austr.
Nephrodium Bak. JoB. **1887**. 170. NPfl. 174.

(D) **gymnopoda** (Bak.) C. Chr. Ind. 269. 1905 — Borneo.
Nephrodium Bak. Tr. Linn. Soc. II. Bot. **4**. 252. 1894.

(D) **gymnosora** (Mak.) C. Chr. Ind. 269. 1905 — Japonia.
Nephrodium Mak. Bot. Mag. Tokyo **13**. 64. 1899.

(C) **Haenkeana** (Pr.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Malesia-Polynesia.
Nephrodium Pr. Epim. 46. 1849. HB. 290. NPfl. 178; Aspidium Fourn. 1873; Nephrodium serratum Pr. 1825 (non Aspidium Sw. 1806 nec Meniscium Cav. 1802); N. multilineatum Pr. 1836; ? Diels 1901; Aspidium Mett. n. 258. 1858.

(D) **Hallieri** (Christ) C. Chr. Ind. 269. 1905 — Borneo.
Aspidium Christ, Ann. Jard. Buit. II. **5**. 106. 1905.

(C) **Harrisoni** (Bak.) C. Chr. Ind. 269. 1905 — Costa Rica.
Nephrodium Bak. 1891; Aspidium Christ 1896. (Chr. 244); Nephrodium stenophyllum Bak. JoB. **1884**. 363 (non Sod. 1883).

(D) **Harveyi** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 812. 1891 — Polynesia.
Aspidium Mett.; Kuhn, Linn. **36**. 115. 1869; Lastrea Carr. 1873; Nephrodium Bak. 1874. HB. 497. NPfl. 172.

(P) **Hasseltii** (Bl.) C. Chr. Ind. 269. 1905 — Java. — Polypodium Bl. Fl. Jav. Fil. 195 t. 92. 1829. HB. 309; Phegopteris Mett. n. 14. 1858.

(C) **hastata** (Fée) Urban, Symb. Ant. **4**. 21. 1903 — India occ.
Goniopteris Fée, 11 mém. 65 t. 18 f. 1. 1866; Nephrodium Jenm. 1879; Aspidium Kuhn 1897.

(D) **Helliana** (Fée) C. Chr. Ind. 270. 1905 — Réunion. — Phegopteris Fée. Gen. 247. 1850-52. Aspidium subglandulosum Mett. msc. (Kuhn 1868); Nephrodium Bak. 1867. HB. 285. NPfl. 177; Dryopteris O. Ktze. 1891.

(P) **Hemsleyana** (Bak.) C. Chr. Ind. 270. 1905 — Guatemala.
Polypodium Bak. in Hemsl. Biol. Centr. Amer. **3**. 660 t. 108. 1885.

(D) **Hendersoni** (Bedd.) C. Chr. Ind. 270. 1905 — India bor.
Lastrea Bedd. Ferns br. Ind. Suppl. 17 t. 377. 1876; Aspidium spectabile Wall. 1828 (nomen, non Bl. 1828); Nephrodium Clarke 1880; Lastrea Bedd. 1883; L. Wallichii Moore 1858 (nomen).

(P) **Henriquesii** (Bak.) C. Chr. Ind. 270. 1905 — Ins. S. Thomé (Guinea).
Polypodium Bak. Bol. Soc. Broter. **4**. 154 t. 1. 1886.

(C) **heterocarpa** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Malesia. Hongkong.
Aspidium Bl. Enum. 155. 1828; Nephrodium Moore 1858. HB. 293 — Aspidium dimorphum Kze. 1848; Nephrodium Moore 1858.

(L) **heteroclita** (Desv.) C. Chr. Ind. 270. 1905 — India occ.
Polypodium Desv. Berl. Mag. **5**. 318. 1811; Phegopteris Kuhn 1897. Gymnogramma gracilis Hew. 1838. HB. 377; Leptogramma J. Sm. 1842; Phegopteris Mett. n. 27. 1858; Grammitis Hewardii Moore 1856; Polypodium Gris. 1864 — Gymnogramme consimilis Fée (1868); Jenm. 1897.

(G) **heterophlebia** (Bak.) C. Chr. Ind. 270. 1905 — Costa Rica.
Polypodium Bak. JoB. **1884**. 363.

heterophylla O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. canescens*.

(D) **heteroptera** (Desv.) C. Chr. Ind. 270. 1905 — Réunion. Madagascar.
Nephrodium Desv. Prod. 256. 1827; Aspidium Mett.; Kuhn 1868. A. stipulaceum Mett. n. 198. 1858. (HB. 269); Lastrea Moore 1858. Phegopteris scalpturata Fée 1850-52. — Conf. Kuhn, v. Deck. Reis. **3**[3]. Bot. 40.

heterosora O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(P) **hexagonoptera** (Michx.) C. Chr. Ind. 270. 1905 — U. S. A. orient.-Minnesota-Louisiana. Simla. — Polypodium Michx. Fl. Bor. Am. **2**. 271. 1803. HB. 309; Phegopteris Fée 1850-52. Mett. n. 20. Chr. 275; Nephrodium Diels 1899. [NPfl. 170.

(G) **Hillii** (Bak.) C. Chr. Ind. 270. 1905 — Queensland.
Polypodium Bak. Syn. 505. 1874; Phegopteris Luerss.; Salom. 1883; Goniopteris Ghiesbrechtii Bail. 1874 (non Dryopteris supra).

hirsuta O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. adenophora.

(D) **hirsutifrons** C. Chr. Ind. 270. 1905 — Fernando Po. — Nephrodium Hendersoni Bak. Ann. Bot. **5**. 323. 1891 (non Dryopteris supra).

(D) **hirta** (Sw.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India occ. Amer. centr. Africa occ. trop. — Polypodium Sw. Prod. 133. 1788; Aspidium Sw. 1801. Mett. n. 271. Chr. 265; Lastrea Pr. 1836; Cystopteris Kl. 1847; Nephrodium Hk. 1862. HB. 278. NPfl. 174 — ? Cystopteris rufescens Fée 1850-52.

(D) **hirtipes** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — China austr. India bor.-Malesia-Polynesia. — Aspidium Bl. Enum. 148. 1828. Chr. 259; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 261. NPfl. 168; Aspidium atratum Wall. 1828. Mett. n. 126. 1858; Lastrea Pr. 1836 — ? Polypodium Scottii Bedd. 1870; Phegopteris Bedd. 1876.

(P) **hirtula** (Kze.) C. Chr. Ind. 270. 1905 — Brasilia.
Polypodium Kze.; Steud. 1824; P. hirsutum Sw. Vet. Ak. Hdl. **1817**. 61. HB. 505 (non Desv. 1811); Phegopteris Fée 1872-73. Chr. 272;

Nephrodium Diels 1899. NPfl. 177; Polypodium lasiernos Spr. 1827; Phegopteris Mett. n. 58. 1858 — Polypodium myriotrichum Bak. 1885!

hispida O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(C) **hispidula** (Dcne.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Malesia orient. Ins. Philipp. N. Guinea. Samoa. — Aspidium Dcne. Nouv. Ann. Mus. **3**. 346. 1834; Nephrodium Bak. 1867. HB. 293. NPfl. 179; N. angustifolium Pr. 1849; Aspidium Mett.; Salom. 1883 — Nephrodium Smithianum [Pr. 1849.

(D) **Holmei** (Bak.) C. Chr. Ind. 271. 1905 — Montserrat. Nephrodium Bak. Ann. Bot. **5**. 317. 1891.

(G) **holophylla** (Bak.) C. Chr. Ind. 271. 1905 — Borneo. Polypodium Bak. JoB. **1888**. 325 (non 1879).

(P) **honesta** (Kze.) C. Chr. Ind. 271. 1905 — Peru. Polypodium Kze. Linn. **9**. 49. 1834; Phegopteris Mett. n. 59. 1858.

(P) **honolulensis** (Hk.) C. Chr. Ind. 271. 1905 — Ins. Hawaii. Polypodium Hk. 1862; Phegopteris Mann 1868; Polypodium Hillebrandii Hk. sp. **4**. 254. 1862 (non 228). HB. 311; Phegopteris Luerss. 1875.

(D) **Hopeana** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891. — Fiji. Nephrodium Bak. Syn. 494. 1874; Lastrea Moore 1882.

(D) **Hornei** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Sechellae. Nephrodium Bak. Syn. 500. 1874; Aspidium Kuhn 1879.

(M) **Hosei** (Bak.) C. Chr. Ind. 271. 1905 — Borneo. Meniscium Bak. Journ. Linn. Soc. **22**. 230. 1886.

(D) **huatuscensis** C. Chr. Ind. 271. 1905 — Mexico. — Aspidium obtusilobum Fée, 8 mém. 105. 1857; Lastrea Moore 1858 (non Dryopteris infra).

(D) **Humblotii** (Bak.) C. Chr. Ind. 271. 1905 — Ins. Comorae. Nephrodium Bak. Bull. Linn. Soc. Paris **1**. 534. 1885.

(D) **humida** (Cord.) C. Chr. Ind. 271. 1905 — Ins Mascarenae. Nephrodium Cord. Fl. Réun. 74. 1895. N. crinitum var. nudatum Bak. Fl. Maur. 495. 1877. (an Aspidium cyclochlamys Fée 1850-52?).

(P) **hydrophila** (Fée) C. Chr. Ind. 271. 1905 — Guadeloupe. Phegopteris Fée, 11 mém. 56 t. 13 f. 3. 1866; Polypodium Bak. 1891.

(P) **ichtiosma** (Sod.) C. Chr. Ind. 271. 1905 — Ecuador. Polypodium Sod. Rec. 59. 1883; Cr. vasc. quit. 294. 1893.

ilicifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(C) **imbricata** (Liebm.) C. Chr. Ind. 271. 1905 — Mexico. Polypodium Liebm. Vid. Selsk. Skr. V. **1**. 210 (seors. 58). 1849; Aspidium Fourn. 1872. A. varians Mett.; Kuhn, Linn. **36**. 114. 1869 (f. Fourn., non Nephrodium Fée 1866).

(D) **immersa** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Malacca. Malesia-Ins. Philipp. — Aspidium Bl. Enum. 156. 1828. Mett. n. 214; Lastrea Moore 1857; Nephrodium Hk. 1862. HB. 263. NPfl. 171.

(G) **imponens** (Ces.) C. Chr. Ind. 271. 1905 — N. Guinea. Polypodium Cesati, Rend. Ac. Napoli **16**. 27, 29. 1877.

(D) **inaequalis** (Schlecht.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Africa austr. Aspidium Schlecht. Adumbr. 23 t. 12. 1825. Mett. n. 151. (Chr. 261) Lastrea Pr. 1836; Nephrodium Hk. 1862. HB. 277. NPfl. 174; Polystichum Keys. 1873.

Obs. f. Christ D. *spinulosa var.; f. Kuhn a D. filice mare *oligodonta non diversa. — Vide Kuhn, v. Deck. Reisen **3**[3]. Bot. 45.

(D) **incana** (Christ) C. Chr. Ind. 272. 1905 — Amazonas.
Aspidium Christ, Hedwigia **44**. 367. 1905.

(C) **incisa** (Sw.) O. Ktze. Rev. Gen Pl. **2**. 813. 1891 — India occ. Columbia.
Polypodium Sw. Prod. 131. 1788; Goniopteris Pr. 1836; Aspidium Gris. 1864 (non Sw. 1806). Chr. 243; Nephrodium Bak. 1867. HB. 288. NPfl. 178; Aspidium scolopendrioides var. incisa Mett. n. 235. 1858. A. stenopteris Kze. 1849; Polystichum Moore 1857; Pleocnemia Moore 1858; Nephrodium Eat. 1859; Hk. 1862.

(P) **incrassata** (Christ) C. Chr. Ind. 272. 1905 — China austr.
Phegopteris Christ, Bull. Boiss. **6**. 963. 1898.

(D) **indecora** (Liebm.) C. Ghr. Ind. 272. 1905 — Mexico. Columbia.
Lastrea Liebm. Vid. Selsk. Skr. V. **1**. 272 (seors. 120). 1849; Aspidium Fourn. 1872; Nephrodium Kiærsk. 1874. Aspidium inquinans Fée 1857; Lastrea Moore 1858.

(D) **insignis** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Columbia.
Aspidium Mett. Ann. sc. nat. V. **2**. 247. 1864; Nephrodium Bak. 1867. [HB. 262.

intermedia Gray, Man. ed. I. 630. 1848 = D. spinulosa*.

intermedia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. rhodolepis.

(C) **invisa** (Forst.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Polynesia.
Polypodium Forst. Prod. 81. 1786 (non Sw. 1788); Nephrodium Carr. 1873. HB. 290. NPfl. 179; Aspidium Christ 1896; Polypodium exosum Gmel. 1791; Aspidium dissectum Mett. 1864 (non Polypodium Forst. 1786); Nephrodium Bak. 1867. HB. ed. I. 290.

irrigua O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(D) **isabellina** (Fée) C. Chr. Ind. 272. 1905 — Brasilia.
Aspidium Fée, Cr. vasc. Br. **1**. 137 t. 45 f. 2. 1869.

(D) **itatiaiensis** C. Chr. Ind. 272. 1905 — Brasilia. — Aspidium tijucense Fée, Cr. vasc. Br. **2**. 72 t. 102. 1872-73 (non D. tijuccana infra).

(C) **jaculosa** (Christ) C. Chr. Ind. 272. 1905 — Formosa.
Aspidium Christ, Bull. Boiss. II. **4**. 615. 1904.

(C) **jamaicense** (Bak.) C. Chr. Ind. 272. 1905 — Jamaica.
Nephrodium Bak.; Jenm. JoB. **1877**. 264.

(D) **japonica** (Bak.) C. Chr. Ind. 272. 1905 — Japonia.
Nephrodium Bak. Ann. Bot. **5**. 318. 1891.

(D) **Jenmani** (Bak.) C. Chr. Ind. 272. 1905 — Jamaica. St. Vincent.
Nephrodium Bak.; Jenm. JoB. **1877**. 263; Lastrea Bull 1887.

juglandifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Phanerophlebia.

(D) **Karwinskyana** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891. — Mexico. Guatemala. — Aspidium Mett. Pheg. u. Asp. 59 n. 141. 1858; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 279.

Kaulfussii O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. oligocarpa*.

(P) **Keraudreniana** (Gaud.) C. Chr. Ind. 272. 1905 — Ins. Hawaii. ? Java.
Polypodium Gaud. Freyc. Voy. Bot. 362 t. 7. 1827. HB. 313; Phegopteris Mann 1868; Nephrodium Diels 1899. NPfl. 177 — Polypodium procerum Brack. 1854 (non Willd. 1810); Phegopteris Mann 1868.

(D) **khasiana** C. Chr. Ind. 272. 1905 — China austr. India bor.
Polypodium elongatum Wall. 1828 (nomen); Hk. sp. **4**. 234. 1862 (non Ait. 1789); Phegopteris J. Sm. 1875; Dryopteris O. Ktze. 1891; Aspidium cuspidatum Mett. Pheg. u. Asp. 92 n. 223. 1858 (non Meniscium Bl. 1828): Nephrodium Bak. 1867. HB. 260. NPfl. 168; Lastrea Bedd. 1870.

(D) **kiboschensis** (Hier.) C. Chr. Ind. 273. 1905 — Africa orient.
Aspidium Hier. in Engl. Pflw. Ostafr. C. 85. 1895.

(D) **kilemensis** (Kuhn) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 (kilmensis) — Kilimanjaro. — Aspidium Kuhn, Fil. Deck. 24. 1867 (kilmense ex err.); Nephrodium Bak. 1874. HB. 498.

(P) **Kingii** (Bedd.) C. Chr. Ind. 273. 1905 — Perak.
Phegopteris Bedd. Handb. Suppl. 84. 1892.

(D) **Koordersii** (Christ) C. Chr. Ind. 273. 1905 — Celebes.
Aspidium Christ, Ann. Jard. Buit. 15. 128 t. 15 f. 17. 1897.

(D) **Kuhnii** (Hier.) C. Chr. Ind. 273. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. 34. 440. 1904.

(D) **Kunzeana** (Hk.) C. Chr. Ind. 273. 1905 — Peru. — Nephrodium Hk. 1862; Aspidium abruptum Kze. Linn. 9. 93. 1834. Mett. n. 227 (non Bl. 1828); Lastrea Pr. 1836; Nephrodium Bak. 1867. HB. 263; Dryopteris O. Ktze. 1891.

(D) **Labordei** (Christ) C. Chr. Ind. 273. 1905 — China.
Aspidium Christ, Bull. Soc. Fr. 52 Mém. I. 40. 1905.

(G) **labuanensis** C. Chr. Ind. 273. 1905 — Borneo.
Polypodium Borneense Hk. sp. 5. 11. 1863. HB. 314 (non Dryopteris supra).

(D) **lacera** (Thbg.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Japonia. Korea. China. Polypodium Thbg. Fl. Jap. 337. 1784; Aspidium Sw. 1801; Polystichum Pr. 1849; Lastrea Eat. 1858; Nephrodium Bak. 1867. HB. 273. NPfl. 173; Aspidium Filix mas var. lacerum Chr. 257.

(D) **laeta** (Komarov) C. Chr. Ind. 267. 1905 — Manshuria.
Nephrodium Komarov, Fl. Mansh. 124. 1901.

(P) **laevigata** (Mett.) C. Chr. Ind. 273. 1905 — Peru.
Phegopteris Mett.; Kuhn, Linn. 36. 112. 1869; Polypodium Bak. 1874. HB. 505 (non 348); P. Lechleri Bak. 1891.

(C) **laevis** (Mett.) C. Chr. Ind. 273. 1905 — Luzon.
Aspidium Mett. Pheg. u. Asp. 104 n. 249. 1858; Nephrodium Moore 1858.

(D) **Lagerheimii** (Sod.) C. Chr. Ind. 273. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 252. 1893.

(D) **lanceolata** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Guatemala.
Nephrodium Bak. Syn. 498. 1874.

(D) **lanciloba** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Queensland.
Nephrodium Bak. Syn. 499. 1874; Lastrea J. Sm. 1877. (t. Benth. = D. decomposita var.).

(D) **lanigera** (Kuhn) C. Chr. Ind. 273. 1905 — Congo.
Aspidium Kuhn, Fil. Afr. 135. 1868. (an D. protensa var.?).

(D) **lanuginosa** (Willd.) C. Chr. Ind. 273. 1905 — Africa austr. et occ. Ins. Mascarenae. — Aspidium Willd.; Klf. Enum. 244. 1824; Nephrodium Desv. 1827; Lastrea Moore 1858; Polystichum Keys. 1873. Aspidium odoratum Sieber; Mett. n. 273. 1858. Chr. 264 (non Bory 1810); Lastrea Pr. 1836. Aspidium catopteron Kze. 1836; Lastrea Pappe et Raws. 1858; Nephrodium Hk. 1862. HB. 284. NPfl. 177; Dryopteris O. Ktze. 1891.

(C) **laruteusis** (Bedd.) C. Chr. Ind. 273. 1905 — Perak. Borneo.
Nephrodium Bedd. Handb. Suppl. 73. 1892.

laserpitiifolia O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Polystichum Standishii.

(P) **laserpitiifolia** (Scort.) C. Chr. Ind. 273. 1905 — Perak.
Polypodium Scort.; Bedd. JoB. 1887. 324; Phegopteris Bedd. 1892.

(D) **lasiopteris** (Sod.) C. Chr. Ind. 274. 1905 — Ecuador.
Nephrodium Sod. Rec. 45. 1883; Cr. vasc. quit. 242. 1893.

(D) **Lastii** (Bak.) C. Chr. Ind. 274. 1905 — Africa orient. trop.
Nephrodium Bak. Ann. Bot. **5**. 324. 1891; Aspidium Hier. 1895.

(P) **latealata** (Christ) C. Chr. Ind. 274. 1905 — Paraguay.
Phegopteris Christ, Ann. Cons. Jard. bot. Genève. **3**. 36. 1899.

(C) **latebrosa** (Kze.) C. Chr. Ind. 274. 1905 — Java. Sumatra.
Aspidium Kze. 1848 (nomen); Mett. Pheg. u. Asp. 104. n. 248. 1858; Nephrodium Moore 1858. (HB. 204).

(P) **laterepens** (Trotter) C. Chr. Ind. 274. 1905 — India bor. — Polypodium Trotter; Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 628 t. 14. 1899.

latifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(D) **latifrons** (Brack.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Hawaii.
Lastrea Brack. Expl. Exp. **16**. 196. 1854; Nephrodium Hk. 1862. HB. 282. NPfl. 177; Aspidium Mann 1868.

latipinna O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. parasitica var.

(D) **laxa** (Franch. et Sav.) C. Chr. Ind. 274. 1905 — Japonia. China.
Aspidium Franch. et Sav. Enum. pl. Jap. **2**. 237. 1876, 631. 1879; Nephrodium Diels 1900.

(D) **Layardi** (Bak.) C. Chr. Ind. 274. 1905 — »Polynesia« (Fiji?).
Nephrodium Bak. Ann. Bot. **5**. 323. 1891.

(P) **Lebeufii** (Bak.) C. Chr. Ind. 274. 1905 — Cambodia.
Polypodium Bak. Ann. Bot. **5**. 461. 1891.

(M) **Lechleri** (Hier.) C. Chr. Ind. 274. 1905 — Peru.
Nephrodium Hier. Engl. Jahrb. **34**. 448. 1904.

(C) **Lenormandi** C. Chr. Ind. 274. 1905 — N. Caledonia. N. Hebridae.
Aspidium resiniferum Lenorm. hb. (non Klf. 1824); A. aneitense Fourn. Ann. sc. nat. V. **18**. 297. 1873; Nephrodium Bak. 1891 (non Dryopteris supra).

(D) **lepida** (Moore) C. Chr. Ind. 274. 1905 — »Polynesia«.
Lastrea Moore, Gard. Chr. n. s. **26**. 681. 1886; Nephrodium Bak. 1891.

lepidocaulis O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(P) **lepidorachis** C. Chr. Ind. 274. 1905 — China.
Polypodium rheosorum Bak. Ann. Bot. **5**. 457. 1891 (non 1884).

(D) **lepigera** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Bonin.
Nephrodium Bak. Syn. 284. 1867.

(D) **Lepinei** (Kuhn) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Tahiti.
Aspidium Kuhn, Linn. **36**. 118. 1869; Nephrodium Bak. 1874. HB. 501.

(D) **Leprieurii** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Guiana. Peru. Costa Rica. — Nephrodium Hk. sp. **4**. 106. 1862. HB. 266. NPfl. 171. Aspidium Tonduzii Christ 1901.

(D) **leptosora** C. Chr. Ind. 274. 1905 — Ecuador. — Nephrodium microsorum Hk. sp. **4**. 106. 1862. HB. 271 (non Endl. 1833); Dryopteris O. Ktze. 1891.

(G) **leucophlebia** (Christ) C. Chr. Ind. 274. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. II. **4**. 961. 1904.

(D) **leucorhachis** (Cheesem.) C. Chr. Ind. 274. 1905 — Raratonga (Cook Ins.).
Nephrodium Cheesem. Tr. Linn. Soc. II. Bot. **6**. 309. 1903.

(D) **leucostipes** (Bak.) C. Chr. Ind. 274. 1905 — Formosa.
Nephrodium Bak. JoB. **1885**. 105; Aspidium Christ 1904.

Leuzeana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(L) **Levingii** (Clarke) C. Chr. Ind. 275. 1905 — Himalaya. — Gymnogramme aurita var. Levingei Clarke, Tr. Linn. Soc. II. Bot. **1**. 568. 1880; G. Levingei Bak. Ann. Bot. **5**. 483. 1891; Leptogramme Bedd. 1892.

(C) **Levyi** (Fourn.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Nicaragua. — Aspidium Fourn. Bull. Soc. Fr. **19**. 255. 1872; Nephrodium Bak. 1874. HB. 502.

(D) **L'Herminieri** (Kze.) C. Chr. Ind. 275. 1905 — Guadeloupe. Jamaica. Costa Rica. Aspidium Kze.; Mett. Pheg. u. Asp. 85. n. 205. 1858; Lastrea Moore 1858.

(D) **libanotica** (Rosenstock) C. Chr. Ind. 275. 1905 — Libanon. Aspidium Rosenstock, Mém. l'Herb. Boiss. n. 9. 1900.

(D) **ligulata** (J. Sm.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Philipp. Lastrea J. Sm. 1841 (nomen); Pr. Epim. 35. 1849; Aspidium Kze.; Mett. n. 213. 1858; Nephrodium Hk. 1862. HB. 264 — Lastrea philippina Pr. 1849; Aspidium Fée 1850-52.

(D) **limbata** (Sw.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India occ. Aspidium Sw. Schrad. Journn. **1800**[2]. 35. 1801. Chr. 251; Nephrodium Desv. 1827. HB. 269; Lastrea Moore 1858. Amauropelta Breutelii Kze. 1843; Aspidium Mett. 1856. n. 204; Lastrea Moore 1857.

(D) **Lindeni** (Kuhn) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Mexico. Guatemala. Aspidium Kuhn, Linn. **36**. 116. 1869 (non Fourn. 1872); Nephrodium Bak. 1874. HB. 493.

(P) **Lindigii** C. Chr. Ind. 275. 1905 — Columbia-Peru. Nephrodium deflexum Pr. Rel. Haenk. **1**. 36 t. 5 f. 2. 1825. NPfl. 172. (non Dryopteris supra); Lastrea Pr. 1836; Phegopteris Mett. Ann. sc. nat. V. **2**. 241. 1864; Polypodium Bak. 1867. HB. 305 (non alior.).

(C) **lineata** (Bl.) C. Chr. Ind. 275. 1905 — Malesia. — Aspidium Bl. Enum. 144. 1828. Mett. n. 264; Meniscium Kze. 1848; Nephrodium Pr. 1849. NPfl. 178; Pronephrium Pr. 1849; Nephrodium glandulosum pt. HB. 291.

(P) **Linnaeana** C. Chr. Ind. 275. 1905 — Europa-Asia sept. Japonia. China. Himalaya occ. Amer. bor. arct. et temp. — Polypodium Dryopteris L. sp. **2**. 1093. 1753. HB. 309; Polystichum Roth 1799; Bernh. 1799; Nephrodium Michx. 1803. NPfl. 175; Lastrea Bory 1826; Aspidium Baumg. 1846; Phegopteris Fée 1850-52. Mett. n. 1. Chr. 275; Gymnocarpium Newm. 1851.

Lobbii O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(C) **Loheriana** (Christ) C. Chr. Ind. 275. 1905 — Luzon. Aspidium Christ, Bull. Boiss. **6**. 191. 1898.

(D) **lomatopelta** (Kze.) C. Chr. Ind. 275. 1905 — Cuba. Aspidium Kze.; Mett. Pheg. u. Asp. 72 n. 172. 1858; Lastrea Moore 1858.

Lonchitis O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(D) **lonchodes** (Eat.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Cuba. Aspidium Eat. Mem. Amer. Ac. n. s. **8**. 210. 1860; Nephrodium Hk. 1862. HB. 270. NPfl. 172; Phegopteris Fée 1866.

(D) **longicaulis** (Bak.) C. Chr. Ind. 275. 1905 — Columbia. Nephrodium Bak. JoB. **1881**. 204. NPfl. 168.

(D) **longicuspis** (Bak.) C. Chr. Ind. 275. 1905 — Madagascar. Nephrodium Bak. Journ. Linn. Soc. **16**. 202. 1877; Aspidium Kuhn 1879.

(C) **longipes** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Java. Celebes. Aspidium Bl. Enum. 155. 1828; Nephrodium Moore 1858. HB. 290.

(P) **longipetiolata** C. Chr. Ind. 275. 1905 — Ecuador. Polypodium dentatum Bak. Ann. Bot. **5**. 456. 1891 (non Forsk. 1775). ? Nephrodium crinitum Sod. 1893 (non Desv. 1827).

(C) **longissima** (Brack.) C. Chr. Ind. 276. 1905 — Tahiti.
Goniopteris Brack. Expl. Exp. **16**. 29 t. 5. 1854; Polypodium Hk. 1863. (HB. 316, non Bl. 1828); Nephrodium Diels 1899. NPfl. 179.

(D) **Lorentzii** (Hier.) C. Chr. Ind. 276. 1905 — Argentina.
Aspidium Hier. Engl. Jahrb. **22**. 368. 1896.

(P) **loxoscaphoides** (Bak.) C. Chr. Ind. 276. 1905 — N. Guinea.
Polypodium Bak. JoB. **1890**. 107.

(C) **lucida** (Bak.) C. Chr. Ind. 276. 1905. — Madagascar.
Nephrodium Bak. Gard. Chr. n. s. **8**. 456. 1877.

(D) **ludens** (Bak.) C. Chr. Ind. 276. 1905 — Samoa.
Nephrodium Bak. Ann. Bot. **5**. 325. 1891. NPfl. 177; Aspidium Christ 1896; Nephrodium arborescens Bak. 1867 (pt.?). HB. 286; Aspidium Luerss. 1871; Dryopteris O. Ktze. 1891.

(D) **Luerssenii** (Harr.) C. Chr. Ind. 276. 1905 — Ins. Philipp.
Nephrodium Harr. Journ. Linn. Soc. **16**. 29. 1877; Aspidium Christ 1898.

(D) **lugubris** (Kze.) C. Chr. Ind. 276. 1905 — Mexico. Brasilia.
Polypodium Kze. hb.; Aspidium Mett. Pheg. u. Asp. 94. n. **230**. 1858. (HB. 266); Lastrea Moore 1858; Nephrodium Rosenst. 1904; Hier. 1904 — ? Aspidium conspersoides Fée 1857; Lastrea Moore 1858.

(D) **lunanensis** (Christ) C. Chr. Ind. 276. 1905 — China austr.
Aspidium Christ, Bull. Boiss. **6**. 966. 1898.

(P) **lustrata** (Hier.) C. Chr. Ind. 276. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. **34**. 443. 1904.

(D) **Macarthyi** (Bak.) C. Chr. Ind. 276. 1905 — China.
Nephrodium Bak. 1891; N. puberulum Bak. JoB. **1875**. 201 (non 1874).

(D) **macradenia** (Sod.) C. Chr. Ind. 276. 1905 — Ecuador.
Nephrodium Sod. Rec. 47. 1883; Cr. vasc. quit. 242. 1893.

(D) **macrostegia** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Guiana. Amazonas.
Nephrodium Hk. sp. **4**. 148. 1862. HB. 501; Aspidium Fée 1872-73.

(D) **macrotis** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Peru. — Nephrodium Hk. sp. **4**. 86 t. 242 B. 1862. HB. 261. NPfl. 169; Lastrea J. Sm. 1875.

macroura O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. patens*.

(C) **madagascariensis** (Fée) C. Chr. Ind. 276. 1905 — Madagascar.
Goniopteris Fée, Gen. 251. 1850-52; Aspidium malgassicum Kuhn 1868.

(D) **magna** (Bak.) C. Chr. Ind. 276. 1905 — Madagascar.
Nephrodium Bak. JoB. **1884**. 142; Ic. pl. t. 1663. NPfl. 177.

(P) **manipurensis** (Bedd.) C. Chr. Ind. 276. 1905 — Manipur. India bor.
Polypodium Bedd. JoB. **1888**. 235; Phegopteris Bedd. 1892.

(P) **Manniana** (Hk.) C. Chr. Ind. 276. 1905 — Fernando Po.
Polypodium Hk. sp. **4**. 253. 1862. HB. 310; Phegopteris Kuhn 1868.

(D) **maranguensis** (Hier.) C. Chr. Ind. 276. 1905 — Africa orient. trop.
Aspidium Hier. in Engl. Pflw. Ostafr. C. 85. 1895.

(D) **marginalis** (L.) Gray, Man. ed. I. 632. 1848 — U. S. A. orient.
Polypodium L. sp. **2**. 1091. 1753; Aspidium Sw. 1801. Mett. n. 130; Nephrodium Michx. 1803. HB. 273. NPfl. 173; Lastrea Pr. 1836; Polystichum Keys. 1873; Aspidium Filix mas var. marginale Chr. 257.

(marginalis × spinulosa vide D. pittsfordensis).

(C) **Martini** C. Chr. Ind. 276. 1905 — Cayenne.
Aspidium connexum Mett. msc. (Salom. 1883); Nephrodium Kuhn; Bak. Fl. bras. **1**². 489. 1870. HB. 502; Dryopteris O. Ktze. 1891 (non supra).

martinicensis O. Ktze. Rev. Gen. Pl. 2. 812. 1891 = Aspidium.

(C) **mascarensis** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Madagascar. Polypodium Bak. Syn. ed. I. 455. 1868; Nephrodium (mascaranense) Bak. 1874. HB. 503; Aspidium Kuhn 1879 (non Klf.; Spr. 1827).

(D) **mascarenarum** Urban, Symb. Ant. 4. 14. 1903 — Madagascar. Ins. Mascarenae. Aspidium oppositum Klf.; Spr. Syst. 4. 108. 1827; Lastrea Pr. 1836; Nephrodium Hk. 1862. HB. 283; NPfl. 177; Dryopteris O. Ktze. 1891 (non infra).

(D) **Matsumurae** (Mak.) C. Chr. Ind. 277. 1905 — Japonia. Aspidium Mak. 1892 (nomen); Nephrodium Mak. Bot. Mag. Tokyo 13. 48. 1899.

(G) **mauiensis** C. Chr. Ind. 277. 1905 — Ins. Hawaii. (Maui). Polypodium Clarkei Bak. Ann. Bot. 5. 457. 1891 (non Dryopteris supra).

(C) **mauritiana** (Fée) C. Chr. Ind. 277. 1905 — Madagascar. Ins. Mascarenae. Natal. — Nephrodium Fée, Gen. 308. 1850-52; Aspidium Kuhn 1868 — A. procerum Bojer; Kuhn 1868 (nomen); v. Deck. Reis. 3[3]. Bot. 41. 1879 (non Spr. 1827); Nephrodium Bak. 1874. HB. 502; Dryopteris O. Ktze. 1891 — Aspidium elatum Boj. 1837 (nomen); Nephrodium Bak. Syn. 502. 1874 (non 298. 1867).

(D) **maxima** (Bak.) C. Chr. Ind. 277. 1905 — Fiji. — Nephrodium Bak. JoB. 1886. 182.

(D) **Maximowiczii** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Japonia. Nephrodium Bak. Syn. 499. 1874; Lastrea Moore 1881; Aspidium commutatum Franch. et Sav. 1876.

(P) **Mazei** (Fourn.) C. Chr. Ind. 277. 1905 — Guadeloupe. Phegopteris Fourn. hb.; Polypodium Bak. Ann. Bot. 5. 457. 1891.

(G) **megacuspis** (Bak.) C. Chr. Ind. 277. 1905 — Tonkin. Polypodium Bak. JoB. 1890. 266.

(G) **megalodus** (Schkuhr) Urban, Symb. Ant. 4. 21. 1903 — Amer. trop. Polypodium Schkuhr, Kr. Gew. 1. 24 t. 19 b. 1806; Goniopteris Pr. 1836; Phegopteris Mett. n. 47. 1858; Aspidium Mett. 1859; Polypodium tetragonum β Bak. 1867. HB. 317. — (an P. pennatum Poir. 1804?).

(C) **megaphylla** (Mett.) C. Chr. Ind. 277. 1905 — India bor. (Bengal)-Ceylon-Malesia. Ins. Comorae. Africa occ. — Aspidium Mett. Ann. Lugd. Bat. 1. 233. 1864; A. pennigerum Bl. Enum. 153. 1828. Chr. 248 (non Sw. 1801); Nephrodium Moore 1858. HB. 292. NPfl. 179 (non Pr. 1825); Aspidium multilineatum Wall. 1828 (nomen; part.); Dryopteris O. Ktze. 1891. — (quod est Nephrodium pennigerum Bedd. Handb. Suppl. 73 1892?).

melanocaulis O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Aspidium.

melanochlamys O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Polystichum.

melanosticta O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Polystichum.

(G) **meniscioides** (Liebm.) C. Chr. Ind. 277. 1905 — Mexico. Panama. Guadeloupe. Polypodium Liebm. Vid. Selsk. Skr. V. 1. 211 (seors. 59). 1849. HB. ed. II. 314; Goniopteris Fourn. 1872; Phegopteris Mett.; Salom. 1883 — Goniopteris rostrata Fée 1866.

meniscioides O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Cyclodium.

(G) **Metcalfei** (Bak.) C. Chr. Ind. 277. 1905 — N. Hebridae. Polypodium Bak. Ann. Bot. 5. 461. 1891.

Metteniana O. Ktze. Rev. Gen. Pl. 2. 811. 1891 = Polystichum multifidum.

mexicana O. Ktze. Rev. Gen. Pl. 2. 811. 1891 = D. patula.

Meyanthidis O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Aspidium menyanthidis.

(P) **Michaelis** (Bak.) C. Chr. Ind. 277. 1905 — Ecuador. Polypodium Bak. JoB. 1877. 164.

(D) **microbasis** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Africa occ.
Nephrodium Bak. Syn. 496. 1874.

(D) **microchlaena** (Fée) C. Chr. Ind. 278. 1905 — Mexico. Columbia.
Aspidium Fée, 8 mém. 102. 1857; Lastrea Moore 1858. Aspidium microcarpon Fée 1857 (non Bl. 1828); Lastrea Moore 1858.

microsora O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. leptosora.

(D) **Millettii** C. Chr. Ind. 278. 1905 — Java. — Aspidium setosum Bl. hb. (non Sw. 1801 nec Schkuhr 1809); Nephrodium Bak. Syn. 274. 1867; Lastrea J. Sm. 1877 (non Pr. 1849); Dryopteris O. Ktze. 1891.

(D) **Milnei** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Fiji.
Nephrodium Hk. 2 Cent. t. 62. 1861. HB. 282.

(D) **Miqueliana** (Maxim.) C. Chr. Ind. 278. 1905 — Japonia.
Aspidium Maxim.; Franch. et Sav. Enum. pl. Jap. **2**. 240. 1876, 634. 1879; Nephrodium Yabe 1903.

mohrioides O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(D) **mollicula** (Kze.) C. Chr. Ind. 278. 1905 — Amer. trop.
Polypodium Kze.; Link, Fil. sp. 130. 1841; Aspidium Mett. 1856 (an Fourn. 1872?); Nephrodium Kiærsk. 1874 — ? Aspidium pilosulum Kl. et Karst.; Kze. 1850. Mett. n. 185; Lastrea Moore 1858; Nephrodium Hk. 1862. (HB. 262) — Aspidium lasiesthes Kze. 1850.

(D) **mollissima** (Christ) C. Chr. Ind. 278. 1905 — Yunnan.
Aspidium Christ, Bull. Boiss. **6**. 968. 1898.

(C) **molliuscula** (Wall.) C. Chr. Ind. 278. 1905 — India bor.
Polypodium Wall. 1828 (nomen); Aspidium Kuhn, Bot. Zeit. **1868**. 41. (HB. 293); Nephrodium Bedd. 1892; N. extensum var. microsorum Clarke 1880; N. microsorum Bedd. 1883 (non Endl. 1833); N. Hopei Bak. 1891.

(D) **mongolica** (Franch.) C. Chr. Ind. 278. 1905 — Mongolia.
Aspidium Franch. hb.; Nephrodium Bak. Ann. Bot. **5**. 317. 1891 (nomen).

(D) **monocarpa** (Cordem.) C. Chr. Ind. 278. 1905 — Réunion.
Nephrodium Cordem. Fl. Réun. 74. 1895.

(D) **monodonta** C. Chr. Ind. 278. 1905 — Perak.
Lastrea unidentata Bedd. Handb. Suppl. 53. 1892 (non Dryopteris infra).

montana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. oreopteris.

(D) **monticola** (Mak.) C. Chr. Ind. 278. 1905 — Japonia. Korea. China.
Nephrodium Mak. Bot. Mag. Tokyo **13**. 80. 1899; Aspidium Goldieanum Christ 1902 (non Hk. 1822).

(D) **Moreletii** C. Chr. Ind. 278. 1905 — Mexico.
Aspidium Lindeni Fourn. Mex. pl. **1**. 97. 1872 (non Kuhn 1869).

(D) **Moritziana** Urban, Symb. Ant. **4**. 21. 1903 — Columbia.
Aspidium diplazioides Moritz; Mett. Pheg. u. Asp. 83. n. 200. 1858; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 270. NPfl. 172; Dryopteris O. Ktze. 1891 (non supra).

(P) **Morlae** (Sod.) C. Chr. Ind. 278. 1905 — Ecuador.
Polypodium Sod. Cr. vasc. quit 641. 1893.

(D) **Motleyana** (Hk.) C. Chr. Ind. 278. 1905 — Borneo. — Nephrodium Hk. Syn. 266. 1867 (sub N. crassifolio); Aspidium Christ, Ann. Jard. Buit. II. **5**. 105. 1905.

(C) **moulmeinensis** (Bedd.) C. Chr. Ind. 278. 1905 — China-India. Celebes. Fiji.
Nephrodium Bedd. Ferns br. Ind. Correct. 1870 (nomen). (HB. 503). NPfl. 178; Aspidium Christ 1902. Polypodium multilineatum Wall.;

Hk. sp. 5. 11. 1863. HB. 316; Goniopteris Bedd. 1867; Phegopteris Luerss. 1871; Aspidium Christ 1897. 249 (non Wall. 1828 nec Mett. 1858).
mucronata O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum echinatum.
multifida O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = D. effusa.
multijuga O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. ochthodes.
multilineata O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = D. megaphylla.
(D) **multiseta** (Bak.) C. Chr. Ind. 279. 1905 — Borneo.
Nephrodium Bak. Journ. Linn. Soc. **22**. 227. 1886.
munita O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.
(D) **mutica** (Franch. et Sav.) C. Chr. Ind. 279. 1905 — Japonia.
Aspidium Franch. et Sav. En. pl. Jap. **2**. 240, 635. 1879.
(D) **myriolepis** (Bak.) C. Chr. Ind. 279. 1905 — San Domingo.
Nephrodium Bak. JoB. **1888**. 34.
(C) **namburensis** (Bedd.) C. Chr. Ind. 279. 1905 — Assam.
Nephrodium Bedd. Handb. Suppl. 69. 1892.
(D) **Napoleonis** (Bory) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — St. Helena.
Nephrodium Bory, Bél. Voy. Bot. **2**. 61 t. 6. 1833. HB. 279. NPfl. 174; Lastrea Pr. 1836; Aspidium Fée 1850-52. Mett. n. 144; Polystichum Keys. 1873.
(D) **negligens** (Jenm.) C. Chr. Ind. 279. 1905 — Jamaica.
Nephrodium Jenm. Bull. Dep. Jam. **3**. 21. 1896.
(C) **nemoralis** (Sod.) C. Chr. Ind. 279. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 267. 1893.
(D) **nemorosa** (Willd.) Urban, Symb. Ant. **4**. 15. 1903 — India occ. — Aspidium Willd. sp. **5**. 255. 1810; Nephrodium Desv. 1827; Lastrea Moore 1858.
nephrodioides O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = D. orbicularis.
(P) **nervosa** (Kl.) C. Chr. Ind. 279. 1905 — Guiana.
Polypodium Kl. Linn. **20**. 386. 1847; Nephrodium Hier. 1904.
nevadensis Und. Our. Nat. Ferns ed. IV. 113. 1893 = D. oregana.
(P) **nicaraguensis** (Fourn.) C. Chr. Ind. 279. 1905 — Nicaragua.
Phegopteris Fourn. Bull. Soc. Fr. **19**. 252. 1872.
nicaraguensis O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
nigrescens O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
(G) **nigrescentia** (Jenm.) C. Chr. Ind. 279. 1905 — Jamaica.
Polypodium Jenm. Gard. Chr. III. **17**. 100. 1895.
(P) **nigritiana** (Bak.) C. Chr. Ind. 279. 1905 — Fernando Po. Africa occ. trop.
Polypodium Bak. Syn. 313. 1867.
nigritiana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. tomentella.
(D) **nigrovenia** (Christ) C. Chr. Ind. 279. 1905 — Honduras.
Nephrodium Christ ap. Donn. Smith, Bot. Gaz. **20**. 545. 1895.
(D) **nimbata** (Jenm.) C. Chr. Ind. 279. 1905 — Jamaica.
Nephrodium Jenm. Gard. Chr. III. **15**. 264. 1894.
(C) **nipponica** (Franch. et Sav.) C. Chr. Ind. 279. 1905 — Japonia. China. Tibet orient. — Aspidium Franch. et Sav. Enum. pl. Jap. **2**. 242, 636. 1879.
nitidula O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. anoptera.
(D) **Nockiana** (Jenm.) C. Chr. Ind. 279. 1905 — Jamaica.
Nephrodium Jenm. JoB. **1886**. 270.
(D) **noveboracense** (L.) Gray, Man. ed. I. 630. 1848 — Amer. septentr. atlant.
Polypodium L. sp. **2**. 1091. 1753; Aspidium Sw. 1801. Mett. n. 187. Chr. 253; Nephrodium Desv. 1827. HB. 267. NPfl. 172; Lastrea Pr. 1836.
(D) **nuda** Und. ap. Heller, Minn. bot. St. **1**. 780. 1897 — Ins. Hawaii.

(D) **obliquata** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — N. Caledonia. Aspidium Mett. Ann. sc. nat. IV. **15**. 75. 1861; Nephrodium Bak. 1867. HB. 264. NPfl. 172 — Aspidium Germinyi Linden 1880.

(G) **obliterata** (Sw.) C. Chr. Ind. 280. 1905 — India occ. Mexico. — Polypodium Sw. Prod. 132. 1788; Goniopteris Pr. 1836; Phegopteris Mett. n. 46. 1858; Nephrodium Keys. 1873. NPfl. 179 — Polypodium faucium Liebm. 1849. HB. 315; Goniopteris Fourn. 1872; Phegopteris Salomon 1883.

(D) **obovata** (Bak.) C. Chr. Ind. 280. 1905 — Tonkin. Assam. Nephrodium Bak. JoB. **1890**. 265; Lastrea Bedd. 1892.

obtusa O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(P) **obtusiloba** (Desv.) C. Chr. Ind. 280. 1905 — Africa trop. Madagascar. Ins. Mascarenae. — Polypodium Desv. Berl. Mag. **5**. 317. 1811. HB. 305; Phegopteris Christ 1897. 273; Aspidium Desvauxii Mett.; Kuhn 1868.

obtusiloba O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. peranemiformis var.

(D) **ochropteroides** (Bak.) C. Chr. Ind. 280. 1905 — Jamaica. Nephrodium Bak. Ann. Bot. **5**. 325. 1891.

(D) **ochrorachis** (Bak.) C. Chr. Ind. 280. 1905 — Madagascar. Nephrodium Bak. JoB. **1884**. 142.

(D) **ochthodes** (Kze.) C. Chr. Ind. 280. 1905 — Japonia. China austr. India. Malacca. — Aspidium Kze. Linn. **24**. 282. 1851. Mett. n. 197; Lastrea Moore 1858; Nephrodium Hk. 1862; N. prolixum β HB. 268 — Aspidium xylodes Kze. 1851 (tylodes auctt.); Lastrea Moore 1858; Nephrodium Hope 1903 — Nephrodium multijugum Bak. 1867. HB. 291; Dryopteris O. Ktze. 1891 — Nephrodium punctatum Bak. 1874. HB. 503 (non Parish 1866); Dryopteris O. Ktze. 1891.

(C) **occulta** (Hope) C. Chr. Ind. 280. 1905 — India bor. Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 627 t. 13. 1899.

oculata O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum Richardi var.

(P) **Oldhami** (Bak.) C. Chr. Ind. 280. 1905 — Formosa. Polypodium Bak. Syn. 311. 1867.

(D) **oligocarpa** (H.B.Willd.) O. Ktze. Rev. Gen. Pl. **3**^{2}. 378. 1898 — Amer. trop. Polypodium H.B.Willd. sp. **5**. 201. 1810; Aspidium HBK. 1815. Mett. n. 184; Nephrodium Desv. 1827; Lastrea Moore 1858 — Polypodium pubescens Raddi 1825 (non L. 1759); Phegopteris Fée 1869; Aspidium Fée 1872-73. Nephrodium conterminum part. HB. 268 — Aspidium Kaulfussii Link 1833. Mett. n. 189; Lastrea Pr. 1836; Nephrodium Hk. 1862. HB. 268. NPfl. 172; Dryopteris O. Ktze. 1891.

(D) **oligophlebia** (Bak.) C. Chr. Ind. 280. 1905 — China. Formosa. Japonia. Nephrodium Bak. JoB. **1875**. 291; Aspidium Christ 1904.

(P) **omeiensis** (Bak.) C. Chr. Ind. 280. 1905 — China occ. Polypodium Bak. JoB. **1888**. 229; Nephrodium Diels 1899. NPfl. 171.

(C) **oochlamys** C. Chr. Ind. 280. 1905 — Mexico. — Aspidium Kunzei Fée, 10 mém. 37 t. 41 f. A, B. 1865; Fourn. Mex. pl. **1**. 96 (non D. Kunzeana supra).

(C) **oosora** (Bak.) C. Chr. Ind. 280. 1905 — Borneo. Nephrodium Bak. Kew Bull. **1896**. 41.

(L) **opaca** (Don) C. Chr. Ind. 280. 1905 — India bor. Java. Hemionitis Don, Prod. Fl. Nepal. 13. 1825; Gymnogramme Spr. 1827. HB. 378; Phegopteris Mett. n. 21. 1858; Leptogramme Bedd. 1883. Gymnogramma obtusata Bl. 1828; Grammitis Pr. 1836; Leptogramma J. Sm. 1875; Phegopteris Christ 274. 1897; Nephrodium Diels 1899. NPfl. 171.

(D) **opposita** (Vahl) Urban, Symb. Ant. **4**. 14. 1903 — Amer. trop.
Polypodium Vahl, Ecl. Amer. **3**. 53. 1807; Aspidium Sw. 1829. Chr. 252 (non Klf. 1827); Nephrodium Diels 1899. NPfl. 172 (non 177). Aspidium conterminum Willd. sp. **5**. 249. 1810. Mett. n. 186; Nephrodium Desv. 1827. HB. 268 (pt.); Polystichum Gaud. 1827; Lastrea Pr. 1836; Dryopteris O. Ktze. 1891. (Oochlamys Fée 1850-52) — Polypodium rivulorum Raddi 1825; Aspidium Link 1833. Mett. n. 193; Lastrea Pr. 1836.

opposita O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. mascarenarum.

(G) **oppositifolia** (Hk.) C. Chr. Ind. 281. 1905 — Ins. S. Thomé (Guinea).
Polypodium Hk. sp. **5**. 8. 1863. HB. 316; Phegopteris Kuhn 1868; Goniopteris J. Sm. 1875.

(C?) **orbicularis** C. Chr. Ind. 281. 1905 — »Malesia«. — Aspidium nephrodioides Hk. sp. **4**. 42 t. 235. 1862 (non Kl. 1847); Dryopteris O. Ktze. 1891; Aspidium Hookeri Bak. 1867. HB. 257 (non Wall. 1829 nec Kl. 1847).

(D) **oregana** C. Chr. Ind. 281. 1905 — California. Oregon.
Aspidium nevadense Eat. Ferns N. Am. **1**. 73 t. 10. 1878 (non Boiss. 1838); Nephrodium Bak. 1891; Dryopteris Und. 1893.

(D) **oreopteris** (Ehrh.) Maxon, Proc. U. S. Nat. Mus. **23**. 638. 1901 — Europa. Madeira. Japonia. Amer. bor. pacif. (Alaska-Washington).
Polypodium Ehrh.; Willd. Prod. 292. 1787; Ehrh. Beitr. **4**. 44. 1789; Aspidium Sw. 1801. Mett. n. 266. Chr. 253; Polystichum Bernh. 1799; Lastrea Bory 1826; Nephrodium Desv. 1827; Phegopteris Fée 1850-52; Hemestheum Parm. 1899; Polypodium montanum Vogler, Dissert. 1781 (non Lam. 1778); Polystichum Roth 1799; Hemestheum Newm. 1851; Lastrea Moore 1857; Aspidium Aschers. 1859; Nephrodium Bak. 1867. HB. 271. NPfl. 173; Dryopteris O. Ktze. 1891.

(D) **orientalis** (Gmel.) C. Chr. Ind. 281. 1905 — Africa trop. Ins. Mascarenae. Amboina. Fiji. — Polypodium Gmel. Syst. Nat. 2^2. 1312. 1791; P. pectinatum Forsk. Fl. Aeg.-Arab. 185. 1775 (non L. 1753); Arthropteris Kuhn 1889; Aspidium Christ 1897. 250; Nephrodium Hier. 1899. Aspidium albopunctatum Bory; Willd. 1810. Mett. n. 123; Nephrodium Desv. 1827. HB. 264; Lastrea Pr. 1836; Arthropteris J. Sm. 1857; Dryopteris O. Ktze. 1891. Aspidium leucosticton Kze. 1850.
— Nephrodium subbiauritum Hk. 1862. HB. 261; Arthropteris J. Sm. 1875; Dryopteris O. Ktze. 1891.
— Nephrodium punctulatum Bak. 1867. HB. 261; Dryopteris O. Ktze. 1891.
— Lastrea articulata Brack. 1854. — Fiji.

(D) **orizabae** (Fée) C. Chr. Ind. 281. 1905 — Mexico.
Aspidium Fée, 8 mém. 104. 1857; Lastrea Moore 1858.

(P) **ornata** (Wall.) C. Chr. Ind. 281. 1905 — China austr. India bor.-Malesia-Australia bor.-Polynesia. — Polypodium Wall. 1828 (nomen); Bedd. Ferns S. Ind. t. 171. 1863-65; Phegopteris Fée 1850-52. Chr. 277; Nephrodium Christ 1895. NPfl. 177; Aspidium Christ 1897; Nephrodium tenericaule Hk. 1862 (sp. **4**. t. 269 non pag. 142); N. setigerum pt. HB. 284 — Polypodium pallidum Brack. 1854.

(C) **oshimensis** (Christ) C. Chr. Ind. 281. 1905 — Liu Kiu.
Aspidium Christ, Bull. Boiss. II. **1**. 1018. 1901.

(C) **otaria** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Assam. India austr. Ceylon. Ins. Philipp. — Aspidium Kze.; Mett. Pheg. u. Asp. 34. n. 73. 1858. Chr. 249; Nephrodium Bak. 1867. HB. 288. NPfl. 178. Aniso-

campium Cumingianum Pr. Epim. 59. 1849 (non Dryopteris supra); Cyclodium Moore 1857. Goniopteris aristata Fée 1850-52 (non Aspidium Sw. 1801); Nephrodium Hk. 1862; Pleocnemia Bedd. 1863-65. Polypodium subpectinatum Wall. 1828 (nomen); Goniopteris J. Sm. 1875; Nephrodium Bedd. 1892; Aspidium Christ 1902. Oligocampia microcarpa Trevis. 1851 (nomen).

(D) **otarioides** (Christ) C. Chr. Ind. 282. 1905 — China.
Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 247.

(P) **oxyodus** (Bak.) C. Chr. Ind. 282. 1905 — Sulu Archip.
Polypodium Bak. JoB. **1879**. 66.

(P) **oyamensis** (Bak.) C. Chr. Ind. 282. 1905 — Japonia. China.
Polypodium Bak. JoB. **1877**. 366; P. Krameri Franch. et Sav. 1876 (nomen); 1879; Phegopteris Mak. 1899; Nephrodium Diels 1899. NPfl. 170; Aspidium Christ 1900 (Crameri), 1905.

pachyphylla O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(D) **pachyrachis** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Venezuela.
Aspidium Kze.; Mett. Pheg. u. Asp. 83. n. 199. 1858; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 270. NPfl. 172.

(D) **padangensis** (Bedd.) C. Chr. Ind. 282. 1905 — Perak.
Lastrea Bedd. Handb. Suppl. 60. 1892.

palladivenia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. striata.

(D) **palustris** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Brasilia. — Aspidium Mett. msc.; Fée 1872-73; Nephrodium Bak. Syn. 270. 1867. NPfl. 172.

(C) **papilio** (Hope) C. Chr. Ind. 282. 1905 — India bor. Ceylon.
Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 625 t. 12. 1899.

(D) **pandiformis** (Christ) C. Chr. Ind. 282. 1905 — China.
Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 255.

(D) **papuana** C. Chr. Ind. 282. 1905 — N. Guinea. — Nephrodium dissitifolium Bak. Kew Bull. **1899**. 120. (non Dryopteris supra).

(C) **papyracea** (Bedd.) C. Chr. Ind. 282. 1905 — India bor.
Nephrodium Bedd. Handb. Suppl. 69. 1892.

(P) **paraguayensis** (Christ) C. Chr. Ind. 282. 1905 — Paraguay.
Phegopteris Christ in litt.; Ph. subsimilis Christ, Ann. Cons. Jard. bot. Genève **3**. 36. 1899 (non Mett. 1868).

(D) **parallela** (Bak.) C. Chr. Ind. 282. 1905 — Madagascar.
Nephrodium Bak. Journ. Linn. Soc. **15**. 417. 1876; Aspidium Kuhn 1879.

(C) **parasitica** (L.) O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 — Trop. et Subtrop. Ins. Atlantidis. N. Zealand. — Polypodium L. sp. **2**. 1090. 1753; Aspidium Sw. 1801. Chr. 243; Nephrodium Desv. 1827. NPfl. 181. Polypodium molle Jacq. Coll. **3**. 188. 1789 (non Schreb. 1771 nec All. 1785); ? Aspidium Sw. 1801 (certe auct. plur.). Mett. n. 246; Nephrodium R. Br. 1810. HB. 293; Polystichum Gaud. 1827; Hemestheum Gandog. 1891 — Polypodium nymphale Forst. 1786; Aspidium Schkuhr 1806; Nephrodium Desv. 1827; Polystichum Gaud. 1827 — Polypodium dentatum Forsk. 1775; Aspidium violascens Link 1833. Mett. n. 247; Nephrodium Fée 1850-52 — N. latipinna Hk. 1867. HB. 292; Aspidium Hance 1873; Dryopteris O. Ktze. 1891.

—* Aspidium amboinense Willd. 1810. Mett. n. 251; Nephrodium Pr. 1836. HB. 292; Dryopteris O. Ktze. 1891.

—* Nephrodium Jamesoni Hk. 1862 — Ecuador. Peru.

(D) **parathelypteris** (Christ) C. Chr. Ind. 283. 1905 — China.
Aspidium Christ, Bull. Soc. Fr. **52** Mém. I. 36. 1905.

(D) **Parishii** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Birma. — Nephrodium Hk. sp. **4**. 131 t. 260. 1862. HB. 280. NPfl. 175; Lastrea Bedd. 1865.

(D) **patens** (Sw.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Amer. trop. et subtrop. (Alabama-Chile). Africa trop. Japonia. China. Polynesia.
Polypodium Sw. Prod. 133. 1788; Polystichum (Bernh. 1799); Gaud. 1827; Aspidium Sw. 1801. Mett. n. 208. Chr. 251; Nephrodium Desv. 1827. HB. 262. NPfl. 170; Lastrea Pr. 1836 — Aspidium deversum Kze. 1850 (pt.). Mett n. 209; Lastrea Moore 1858; Nephrodium Rosenst. 1904. —* Aspidium stipulare Willd. 1810; Nephrodium Desv. 1827. ? Aspidium macrourum Klf. 1823. Mett. n. 210; Lastrea Pr. 1836; Nephrodium Schott 1834. HB. 262; Dryopteris O. Ktze. 1891. Nephrodium schizotis Hk. 1862.
—* Polypodium invisum Sw. 1788 (non Forst. 1786); Aspidium Sw. 1801. Mett. n. 211; Nephrodium Desv. 1827; Lastrea Pr. 1836; Nephrodium Sloanei Bak. 1874. HB. 263 (non Pr. 1825); Dryopteris O. Ktze. 1891. Nephrodium Serra Hk. 1862 (non Desv. 1827).

(D) **patula** (Sw.) Und. Our nat. Ferns ed. IV. 117. 1893 — Amer. trop. Arizona.
Aspidium Sw. Vet. Ak. Hdl. **1817**. 64. Chr. 263; Lastrea Moore 1858; Nephrodium Bak. 1870. HB. 276. NPfl. 174. N. mexicanum Pr. 1825; Lastrea Pr. 1836; Aspidium Kze. 1839. Mett. n. 152; Dryopteris O. Ktze. [1891.

(M) **pauciflora** (Hk.) C. Chr. Ind. 283. 1905 — Africa occ. trop.
Meniscium Hk. sp. **5**. 164. 1864. HB. 391; Phegopteris Kuhn 1868.

(G) **paucinervata** C. Chr. Ind. 283. 1905 — Peru. — Polypodium oligophlebium Bak. Syn. 506. 1874 (non Kze. 1850); Phegopteris Salom. 1883.

(G) **paucivenia** C. Chr. Ind. 283. 1905 — Madagascar.
Polypodium oligophlebium Bak. JoB. **1891**. 6 (non 1874 nec Kze. 1850).

(P) **Pavoniana** (Kl.) C. Chr. Ind. 283. 1905 — Peru. — Polypodium Kl. Linn. **20**. 386. 1847; Phegopteris Mett. 1869; Nephrodium Hier. 1904.

(D) **pedata** (Desv.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India occ.
Aspidium Desv. Prod. 244. 1827; Mett. n. 277; Camptodium Fée 1850-52; Lastrea Moore 1857; Nephrodium Hk. 1862. HB. 259. NPfl. 170.

(D) **pellucida** (Franch.) C. Chr. Ind. 283. 1905 — China austr. — Aspidium Franch. Nouv. Arch. Mus. II. **10**. 119. 1887; Christ, Bull. Soc. Fr. **52** Mém. I. 41. 1905; Nephrodium Diels 1900. (an D. sparsa *nitidula?).

(Ph) **pellucido-punctata** C. Chr. Ind. 283. 1905 — Peru.
Polypodium macrophyllum Hk. sp. **4**. 241. 1862. HB. 308 (non Mett. 1856); Phegopteris J. Sm. 1875.

(C) **penangiana** (Hk.) C. Chr. Ind. 283. 1905 — China. India bor. Penang.
Polypodium (Panangianum) Hk. sp. **5**. 13. 1863; Goniopteris Bedd. 1867. Polypodium lineatum Colebr.; Wall. 1828; Hk. sp. **5**. 12. 1863. HB. 316, 503 (non Dryopteris supra); Goniopteris Pr. 1836; Phegopteris Mett.; Salom. 1883; Polypodium costatum Wall. hb.; Goniopteris J. Sm. 1841 (nomen); Nephrodium Bedd. 1867.

(G) **pennigera** (Forst.) C. Chr. Ind. 283. 1905 — N. Zealand. Polynesia. ? Ins. Philipp. — Polypodium Forst. Prod. 82. 1786. HB. 317; Aspidium Sw. 1801; Nephrodium Pr. 1825. NPfl. 179; Polystichum Gaud. 1827; Lastrea Pr. 1836; ? Goniopteris J. Sm. 1841; G. Forsteri Moore 1858.

— Polypodium Adenopteris Kze. hb.; Aspidium Mett. n. 255. 1858; Nephrodium Moore 1858.

—* Goniopteris costata Brack. 1854; Polypodium Hk. 1863. HB. 316; Nephrodium Diels 1899. NPfl. 179 (non Bedd. 1867).

(D) **Pentheri** (Krasser) C. Chr. Ind. 284. 1905 — Africa austr.
Nephrodium Krasser, Ann. Hofmus. Wien **15**. 5 t. 2 f. 1-5. 1900.

(C) **perakensis** (Bedd.) C. Chr. Ind. 284. 1905 — Perak.
Aspidium Bedd. JoB. **1888**. 4; Nephrodium Bak. 1891.

(D) **peranemiformis** C. Chr. Ind. 284. 1905 — India austr. Ceylon. Celebes.
Lastrea ferruginea Bedd. Ferns S. Ind. t. 100. 1863-65 (non Aspidium Fée 1850-52); Nephrodium Bak. 1867. HB. 283. NPfl. 177; Dryopteris O. Ktze. 1891.

—* Nephrodium obtusilobum Bak. 1867. HB. 284; Lastrea Bedd. 1868; Aspidium Prantl 1882; Dryopteris O. Ktze. 1891 (non supra).

(P) **Percivalii** (Jenm.) C. Chr. Ind. 284. 1905 — Guiana.
Polypodium Jenm.; Bak. Ann. Bot. **5**. 456. 1891.

(D) **peregrina** C. Chr. Ind. 284. 1905 — China.
Nephrodium regulare Bak. JoB. **1875**. 200 (non Desv. 1827).

(C) **Peripae** (Sod.) C. Chr. Ind. 284. 1905 — Ecuador.
Nephrodium Sod. Rec. 52. 1883; Cr. vasc. quit. 265. 1893.

(P) **persimilis** (Bak.) C. Chr. Ind. 284. 1905 — Samoa.
Polypodium Bak. JoB. **1876**. 344.

(D) **phaeostigma** (Ces.) C. Chr. Ind. 284. 1905 — N. Guinea.
Aspidium Ces. Rend. Acad. Napoli **16**. 26, 29. 1877.

(D) **phanerophlebia** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Salomonis.
Nephrodium Bak. Syn. 494. 1874.

(P) **phegopteris** (L.) C. Chr. Ind. 284. 1905 — Europa. Asia minor. Asia septentr.-Himalaya-Japonia. Amer. bor. arct. et temp. — Polypodium L. sp. **2**. 1089. 1753. HB. 308; Polystichum Roth 1799; Bernh. 1799; Lastrea Bory 1826; Aspidium Baumg. 1846; Gymnocarpium Newm. 1851; Phegopteris Keys. 1873; Nephrodium Prantl 1884. NPfl. 170; Phegopteris polypodioides Fée 1850-52. Chr. 275; Ph. vulgaris Mett. n. 19. 1856.

(D) **phegopteroidea** (Kuhn) C. Chr. Ind. 284. 1905 — Timor.
Hypodematium Kuhn, Forschr. Gazelle **4** Farne 8. 1889.

(D) **philippina** (Pr.) C. Chr. Ind. 284. 1905 — Ins. Philipp.
Physematium Pr. Epim. 34. 1849. Lastrea exigua J. Sm. 1841 (nomen); Haplodictyum? Fée 1850-52 (pt.); Aspidium Kze.; Mett. n. 180. 1858. Chr. 245; Nephrodium Hk. 1862. HB. 267; Dryopteris O. Ktze. 1891 — ? Phegopteris nervosa Fée 1850-52.

(D) **physematioides** (Kuhn et Christ) C. Chr. Ind. 284. 1905 — San Domingo.
Aspidium Kuhn et Christ, Engl. Jahrb. **24**. 115. 1897.

Pica O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(L) **pilosa** (Mart. et Gal.) C. Chr. Ind. 284. 1905 — Mexico. — Gymnogramme Mart. et Gal. Mém. Ac. Brux. **15**. 27 t. 4 f. 1. 1842. HB. 376; Phegopteris Mett. n. 30. 1858; Leptogramma Und. 1902 — Gymnogramme procurrens Fée 1857. (HB. 515); Phegopteris Mett.; Salomon 1883.

(C) **pilosiuscula** (Zippel) C. Chr. Ind. 284. 1905 — Java.
Nephrodium Zippel; Racib. Pterid. Buit. 189. 1898.

(D) **pittsfordensis** Slosson, Rhodora **6**. 75. 1904 — Vermont.
Nephrodium Woolson 1904. (? D. marginalis × spinulosa).

plantaginea O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
Plaschnickiana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.
platanifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
(P) **platyloba** (Bak.) C. Chr. Ind. 285. 1905 — Peru. — Polypodium Bak. Syn. 306. 1867; P. rotundatum Hk. sp. **4**. 238. 1862 (non Dryopteris infra).
platypus O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum aristatum var.
(D) **platyrachis** (Fée) C. Chr. Ind. 285. 1905 — Brasilia.
Aspidium Fée, Cr. vasc. Br. **2**. 71 t. 100 f. 2. 1872-73.
pleiotoma O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
(D) **podophylla** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — China austr.
Aspidium Hk. JoB. **5**. 236 t. 1. 1853. Mett. n. 125. Chr. 250; Lastrea Hk. 1857; Nephrodium Hk. 1862. HB. 260. NPfl. 168.
(G) **poecilophlebia** (Hk.) C. Chr. Ind. 285. 1905 — Australia.
Polypodium Hk. sp. **5**. 14. 1863. HB. 314; Phegopteris Luerss. 1873; Goniopteris Bail. 1874.
(G) **Poiteana** (Bory) Urban, Symb. Ant. **4**. 20. 1903 — Amer. trop.
Lastrea Bory, Dict. class. **9**. 233. 1826. Polypodium crenatum Sw. Prod. 132. 1788. HB. 315 (non Forsk. 1775); Goniopteris Pr. 1836; Phegopteris Mett. 1856. Chr. 268; Nephrodium Keys. 1873 — Goniopteris Rivoirei Fée 1850-52; Polypodium Bak. 1891; Nephrodium Diels 1899. NPfl. 179.
(D) **polylepis** (Franch. et Sav.) C. Chr. Ind. 285. 1905 — Japonia. China.
Aspidium Franch. et Sav. En. pl. Jap. **2**. 236. 1876; 631. 1879; Nephrodium Bak. 1891.
polymorpha O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
(D) **polypodiformis** (Mak.) C. Chr. Ind. 285. 1905 — Japonia. — Aspidium Mak. 1892 (nomen); Nephrodium Mak. Bot. Mag. Tokyo **13**. 58. 1899.
(L) **polypodioides** (Raddi) C. Chr. Ind. 285. 1905 — Brasilia.
Ceterach Raddi, Opusc. sci. Bol. **3**. 284. 1819; Pl. Bras. **1**. 10 t. 22. 1825; Gymnogramme Spr. 1827. HB. 377; Grammitis Pr. 1836; Leptogramma J. Sm. 1841; Phegopteris Mett. 1856. n. 33 (non Fée 1850-52); Gymnogramme Raddiana Link 1833. Polystichum gymnogrammoides Gaud. 1827.
(D) **Poolii** C. Chr. Ind. 285. 1905 — Madagascar.
Nephrodium fibrillosum Bak. Journ. Linn. Soc. **15**. 418. 1876 (non Dryopteris supra); Aspidium Kuhn 1879.
(D) **popayanensis** (Hier.) C. Chr. Ind. 285. 1905 — Columbia.
Nephrodium Hier. Engl. Jahrb. **34**. 447. 1904.
(C) **porphyrophlebia** (Christ) C. Chr. Ind. 285. 1905 — China.
Aspidium Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 117.
(D) **Powellii** (Bak.) C. Chr. Ind. 285. 1905 — Samoa. — Nephrodium Bak. Ann. Bot. **5**. 325. 1891. NPfl. 177; Aspidium Christ 1896.
(D) **praetervisa** (Kuhn) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Venezuela.
Aspidium Kuhn, Linn. **36**. 111. 1869; Nephrodium Bak. 1874. HB. 495.
(P) **prasina** (Bak.) C. Chr. Ind. 285. 1905 — Costa Rica-Peru.
Polypodium Bak. Syn. 312. 1867; P. punctatum Spruce; Hk. sp. **4**. 262. 1862 (non Thbg. 1784).
(D) **Prenticei** (Carr.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Fiji. Samoa.
Lastrea Carr. in Seem. Fl. Vit. 359. 1873; Nephrodium Bak. 1874. HB. 494; Aspidium Christ 1896.
Prescottiana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(D) **Preslii** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Philipp.
Nephrodium Bak. Syn. 272. 1867; Lastrea propinqua Pr. Epim. 38. 1849 (J. Sm. 1841 part., non Nephrodium R. Br. 1810); Aspidium Fée 1850-52.

(P) **prionites** (Kze.) C. Chr. Ind. 286. 1905 — Brasilia.
Polypodium Kze. Flora **1839**[1]. Beibl. 29; Phegopteris (Prionitis) Fée 1850-52. Mett. n. 37; Nephrodium Diels 1899. NPfl. 169; Polypodium flavopunctatum *β* HB. 304.

procera O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = D. mauritiana.

(C) **procurrens** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Java. Celebes. India bor. — Aspidium Mett. Ann. Lugd. Bat. **1**. 231. 1864; Nephrodium Bak. 1867. HB. 290.

(C) **producta** (Klf.) C. Chr. Ind. 286. 1905 — Luzon.
Aspidium Klf. Enum. 237. 1824. Mett. n. 261; Nephrodium Pr. 1836.

(P) **prolifera** (Retz.) C. Chr. Ind. 286. 1905 - China austr.-India bor.-Malesia-Polynesia-Australia. Africa trop. Ins. Mascarenae.
Hemionitis Retz. Obs. **6**. 38. 1791; Meniscium Sw. 1806; Polypodium Roxb.; Wall. 1828. HB. 315; Goniopteris Pr. 1836; Nephrodium Keys. 1873; Phegopteris Kuhn 1879. Chr. 270; Aspidium Hier. 1895 — Ampelopteris elegans Kze. 1848. A. firma Kze. 1851 — Polypodium luxurians Kze. 1850; Phegopteris Mett. n. 51. 1858.

(D) **prolixa** (Willd.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India. Africa trop. Ins. Mascar. — Aspidium Willd. sp. **5**. 251. 1810. Chr. 254; Nephrodium Desv. 1827. HB. 268 (pt.). NPfl. 172; Lastrea Pr. 1836 — Aspidium appendiculatum Wall. 1828 (pt.); Mett. n. 196 (non Bl. 1828); Nephrodium Schott 1834; Hk. 1862; ? Lastrea Pr. 1836.

(D) **prominula** (Christ) C. Chr. Ind. 286. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. **4**. 659. 1896.

propinqua Gilb. Bull. Torr. Cl. **4**. 659. 1896 = D. gongylodes*.

(D) **protensa** (Afz.) C. Chr. Ind. 286. 1905 — Amer. trop. Africa occ. trop. Samoa.
Aspidium Afz.; Sw. Schrad. Journ. **1800**[2]. 36. 1801; Polypodium Poir. 1804; Lastrea Moore 1858; Nephrodium Bak. 1870. NPfl. 175. Aspidium subquinquefidum Pal. Beauv. 1804. Chr. 264; Lastrea Pr. 1836; Nephrodium Hk. 1862. HB. 281 (excl. *β*); Polystichum Keys. 1873; Dryopteris O. Ktze. 1891.

—* Aspidium funestum Kze. 1834. Mett. n. 169; Lastrea Moore 1858; Nephrodium Hk. 1862. NPfl. 175. — Conf. D. lanigera.

— (t. Kze. Flora **1839**[1]. Beibl. 33 Aspidium cicutarium Sw. 1801 et certe Willd. 1810, non Polypodium L. 1759, est eadem sp. ac D. protensa).

(D) **pseudomontana** (Hier.) C. Chr. Ind. 286. 1905 — Argentina. Brasilia austr.
Aspidium Hier. Engl. Jahrb. **22**. 373. 1896; Nephrodium Rosenst. 1904.

(C) **pseudoreptans** C. Chr. Ind. 286. 1905 — Sumatra.
Nephrodium debile Bak. JoB. **1880**. 212 (non Dryopteris supra).

(D) **pseudotetragona** Urban, Symb. Ant. **4**. 20. 1903 — Columbia-Peru. Brasilia.
Nephrodium Hier. 1904; N. tetragonum Pr. Rel. Haenk. **1**. 35. 1825. HB. 266. NPfl. 169; Lastrea Pr. 1836; Aspidium Mett. n. 232. 1858; Dryopteris O. Ktze. 1891 (non infra) — Aspidium setosum Kl. 1847 (non Sw. 1801); Nephrodium Keys. 1873 — Aspidium quadrangulare Fée 1869 (non Nephrodium Fée 1850-52).

(D) **pseudothelypteris** (Rosenst.) C. Chr. Ind. 286. 1905 — Brasilia austr.
Nephrodium Rostenst. Hedwigia **43**. 225. 1904.

(D) **pseudovaria** (Christ) C. Chr. Ind. 287. 1905 — China merid.
Aspidium Christ, Bull. Soc. Fr. **52** Mém. I. 42. 1905.

(D) **ptarmica** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Brasilia austr.
Aspidium Kze.; Mett. Pheg. u. Asp. 80. n. 191. 1858; Lastrea Moore 1858; Nephrodium Bak. 1870. HB. 496.

(D) **pterifolia** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Columbia. Bolivia.
Aspidium Mett.; Kuhn, Linn. **36**. 110. 1869; Nephrodium Bak. 1874. HB. 497.

(P) **pteroidea** (Kl.) C. Chr. Ind. 287. 1905 — Columbia-Ecuador. Brasilia.
Polypodium Kl. Linn. **20**. 389. 1847. HB. 308; Phegopteris Mett. n. 3. 1858; Nephrodium Diels 1899. NPfl. 171 — Gymnogramme patula Fée 1869.

(C) **pteroides** (Retz.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — China-Birma-Malesia-Polynesia-Queensland. — Polypodium Retz. Obs. **6**. 39. 1791; Aspidium Sw. 1801. Chr. 247; Nephrodium J. Sm. 1857. HB. 289. NPfl. 178 — Aspidium obtusatum Sw. 1801. Mett. n. 244; Nephrodium Desv. 1827. Aspidium terminans Wall. 1828; Nephrodium J. Sm. 1846; Hk. 1862 (pt.).

puberula O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. Feei.

(D) **pubescens** (L.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India occ.
Polypodium L. Syst. ed. X. **2**. 1327. 1759; Aspidium Sw. 1801. Mett. n. 166; Nephrodium Desv. 1827. HB. 280. NPfl. 175; Lastrea Pr. 1836 — Polypodium chaerophylloides Poir. 1804; ? Nephrodium Desv. 1827; Polystichum Pr. 1836. Polypodium muscosum Vahl 1807; Aspidium Sw. 1829. Polypodium portoricense Spr. 1821; Phegopteris Fée 1850-52. Mett. n. 17 — Phegopteris sericea Mett. 1860; Polypodium Hk. 1862.

(D) **pubirachis** (Bak.) C. Chr. Ind. 287. 1905 — Samoa.
Nephrodium Bak. JoB. **1876**. 344; Aspidium Christ 1896.

(D) **pulvinulifera** (Bedd.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Sikkim.
Lastrea Bedd. Ferns br. Ind. t. 333. 1870; Nephrodium Bak. 1874. HB. 500. NPfl. 174.

(P) **punctata** (Thbg.) C. Chr. Ind. 287. 1905 — Japonia. Asia trop. Australia. Polynesia. N. Zealand. Hawaii. Ins. afric. occ. et orient. Amer. trop.-Chile.
Polypodium Thbg. Fl. Jap. 337. 1784. HB. 312 (non Sw. 1801); Phegopteris Mett. 1864. Chr. 277; Hypolepis Mett.; Kuhn 1868; ? Bedd. 1892; Nephrodium Diels 1899. NPfl. 177 — Polypodium Poeppigii Kze. 1834. Phegopteris Fée; Gay 1853. Mett. n. 13. (Chile) — Polypodium amplum Col. 1892 (non H.B.Willd. 1810). (N. Zealand) — adhuc species plur. Cheilanthidis et Hypolepidis auctt.
—* Polypodium rugosulum Lab. 1806; Hypolepis (rugulosa) J. Sm. 1846; Phegopteris Fée 1850-52. Mett. n. 12.
— (an potius Hypolepis?).

punctata O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. ochthodes.

punctulata O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — D. orientalis var.

pungens O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(D) **pusilla** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Columbia.
Aspidium Mett. Ann. sc. nat. V. **2**. 245. 1864; Nephrodium Bak. 1867. HB. 260. NPfl. 168; Hypodematium Kuhn 1889.

(D) **ramosum** (Hope) C. Chr. Ind. 287. 1905 — India bor. — Nephrodium Hope, JoB. **1896**. 126; Journ. Bomb. Nat. Hist. Soc. **14**. 739 t. 32.

(C) **rampans** (Bak.) C. Chr. Ind. 287. 1905 — China merid.
Nephrodium Bak. JoB. **1889**. 177; Aspidium Christ 1898.

(D) **recedens** (J. Sm.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India austr. Ceylon. Ins. Philipp. — Polypodium J. Sm. 1841 (nomen); Lastrea J. Sm.; Moore, Gard. Chr. **1855**. 708 c. fig.; Aspidium Lowe 1859; Nephrodium Hk. 1862. HB. 281. NPfl. 175.

(C) **refracta** (Fisch. et Mey.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Brasilia. Paraguay. Argentina. — Polypodium Fisch. et Mey.; Kze. Linn. **23**. 283, 321. 1850; Aspidium A. Br. 1856. Mett. n. 238. Chr. 245; Goniopteris J. Sm. 1857; Nephrodium Hk. 1862. HB. 292. NPfl. 179; Phegopteris Fourn. 1872. — an Polypodium retroflexum L. 1753; Aspidium Willd. 1810; Nephrodium Desv. 1827 (Trinidad)?

(P) **refulgens** (Kl.) C. Chr. Ind. 288. 1905 — Panama-Guiana. Polypodium Kl. hb. HB. 307. 1867; Phegopteris Mett. Ann. sc. nat. V. **2**. 240. 1864; Nephrodium Diels 1899. NPfl. 170.

repanda O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(D) **repens** (Hope) C. Chr. Ind. 288. 1905 — India bor. Sierra Leone. Nephrodium Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 535 t. 8. 1899.

(C) **reptans** (Gmel.) C. Chr. Ind. 288. 1905 — Florida-Brasilia. Polypodium Gmel. Syst. Nat. **2**[2]. 1309. 1791; Sw. Schrad. Journn. **1800**[2]. 25. 1801. HB. 316; Goniopteris Pr. 1836; Aspidium Mett. n. 237. var. 3. 1858. Chr. 244; Phegopteris Eat. 1883; Nephrodium Diels 1899. NPfl. 168; Polypodium repens Sw. Prod. 132 (non 130). 1788 (non Aubl. 1775); P. radicans Poir. 1804 — ? Goniopteris gracilis Moore et Houlst. 1856; Polypodium Lowe 1858.

—* Goniopteris asterothrix Fée 1850-52; Phegopteris Mett. n. 40. 1858; Polypodium Hk. 1863.

(D) **resinofoetida** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Costa Rica-Bolivia. Nephrodium Hk. sp. **4**. 105. 1862 (pt.). HB. 269. NPfl. 172; Aspidium Christ 1896. Chr. 253.

(M) **reticulata** (L.) Urban, Symb. Ant. **4**. 22. 1903 — Amer. trop. Polypodium L. Syst. Nat. ed. X. **2**. 1325. 1759; Meniscium Sw. 1803. HB. 392; Phegopteris Mett. 1859. Chr. 269; Nephrodium Keys. 1873. NPfl. 179.

—* Asplenium sorbifolium Jacq. 1788; Meniscium Desrouss. 1797; Phegopteris Mett. 1859; Nephrodium Hier. 1904.

—* Meniscium macrophyllum Kze. 1839; Phegopteris Mett. 1859; Nephrodium Keys. 1873.

—* Meniscium falcatum Liebm. 1849; Phegopteris Mett. 1859.

—* Phegopteris membranacea Mett. 1859.

—* Phegopteris mollis Mett. 1864; Meniscium Bak. 1868. HB. 392 (in obs.). — Conf. Mett. Fil. Lechl. **2**. 23-25.

(D) **retrorsa** (Sod.) C. Chr. Ind. 288. 1905 — Ecuador. Nephrodium Sod. Rec. 51. 1883; Cr. vasc. quit. 244. 1893.

(P) **retusa** (Sw.) C. Chr. Ind. 288. 1905 — Brasilia. — Polypodium Sw. Vet. Ak. Hdl. **1817**. 61; Lindm. Ark. för Bot. **1**. 227 t. 10 f. 11; Aspidium Mett. 1870.

rhizophylla O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.

(D) **rhodolepis** (Clarke) C. Chr. Ind. 288. 1905 — Japonia. China. Asia trop. Nephrodium Clarke, Tr. Linn. Soc. II. Bot. **1**. 526 t. 72. 1880. Aspidium intermedium Bl. Enum. 161. 1828. Chr. 263 (non Willd. 1810); Nephrodium Bak. 1867. HB. 283; Lastrea Bedd. 1870; Polystichum Keys. 1873; Dryopteris O. Ktze. 1891; Lastrea Blumei Moore 1858; Nephrodium Hk. 1862 (pt.). NPfl. 177; Aspidium Thwait. 1864 (non

Kze.; Mett. 1856) — Nephrodium sarawakensis Bak. 1886 — N. Mannii Hope 1890.

Richardi O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Polystichum.

(D) **Richardsi** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — N. Caledonia.
Nephrodium Bak. Syn. 494. 1874; Lastrea Moore 1881.

(D) **rigescens** (Sod.) C. Chr. Ind. 289. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 239. 1893.

rigida A. Gray, Man. ed. I. 631. 1848 = D. cristata × spinulosa (Boottii).

(D) **rigida** (Hoffm.) Und. Our nat. Ferns ed. IV. 116. 1893 — Europa. Reg. Mediterraneae-Afghanistan. Alaska-California.
Polypodium Hoffm. Deutschl. Fl. 2. 6. 1795; Aspidium Sw. 1801. Mett. n. 134. Chr. 260; Polystichum Lam. et DC. 1805; Nephrodium Desv. 1827. HB. 275. NPfl. 173; Lastrea Pr. 1836; Lophodium Newm. 1851. —* Aspidium rigidum var. australe Tenore, Atti Ist. Incor. Napoli 5. 144 t. 2. f. 4 B. 1830; Nephrodium pallidum Bory 1832. NPfl. 173; Aspidium Link 1841 (non Bl. 1828); Polystichum Todaro 1866. (Mediterr.). — vide Sagorski, Österr. Bot. Zeit. 53. 76. 1903.
—* Aspidium argutum Klf. Enum. 212. 1824. Chr. 260; Lastrea Brack. 1854; Nephrodium Diels 1899. NPfl. 173. (Amer. bor. pacif.).

rigidula O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = D. scalpturoides.

(G) **riograndensis** (Lindm.) C. Chr. Ind. 289. 1905 — Brasilia austr.
Polypodium Lindm. Ark. för Bot. 1. 230 t. 9 f. 6. 1903.

(P) **Robertiana** (Hoffm.) C. Chr. Ind. 289. 1905 — Europa. Afghanistan. Amer. bor. temp. — Polypodium Hoffm. Deutschl. Fl. 2. 20. 1795; Gymnocarpium Newm. 1851; Phegopteris A.Br.; Aschers. 1859. Chr. 276; Nephrodium Prantl 1884. NPfl. 175; Aspidium Luerss.; Aschers. 1896. Polypodium calcareum Sm. 1804; Lastrea Bory 1826; Aspidium Baumg. 1846; Phegopteris Fée 1850-52. Mett n. 2; Polypodium Dryopteris β Robertianum [HB. 309.

(C) **Rodigasiana** (Linden) C. Chr. Ind. 289. 1905 — Samoa.
Aspidium Linden msc.; Nephrodium Moore, L'Ill. Hort. 29. 27 t. 442. 1882.

(P) **roraimensis** (Bak.) C. Chr. Ind. 289. 1905 — Guiana.
Polypodium Bak. Tr. Linn. Soc. II. Bot. 2. 291. 1887.

(D) **Rosthornii** (Diels) C. Chr. Ind. 289. 1905 — China.
Nephrodium Diels, Engl. Jahrb. 29. 190. 1900.

(P) **rotundata** (Willd.) C. Chr. Ind. 289. 1905 — Amer. trop. — Aspidium Willd. sp. 5. 247. 1810; Nephrodium Desv. 1827. NPfl. 168; Lastrea Pr. 1836; Phegopteris Fée 1866; Polypodium Bak. 1870. P. flavopunctatum Klf. 1824. HB. 304 (excl. β); Phegopteris Fée 1850-52. Mett. n. 35. Chr. 271 — Nephrodium Imrayanum Hk. 1862; Aspidium? Fée 1866; Phegopteris J. Sm. 1875.

(G) **rubida** (J. Sm.) C. Chr. Ind. 289. 1905 — Ins. Philipp. ? Borneo.
Goniopteris J. Sm. 1841 (nomen); Polypodium Hk. sp. 5. 12. 1863. HB. 314; Nephrodium Copeland 1905.

(D) **rubiginosa** (Brack.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Hawaii. ? Fiji.
Lastrea Brack. Expl. Exp. 16. 201. 1854; Nephrodium Hk. 1862. HB. 280. NPfl. 174; Aspidium Mann 1868.

(G) **rubrinervis** (Mett.) C. Chr. Ind. 289. 1905 — Polynesia.
Phegopteris Mett.; Kuhn, Linn. 36. 116. 1869; Goniopteris Carr. 1873; Polypodium Bak. 1874. HB. ed. II. 315; Nephrodium Diels 1899. NPfl. 178.

(P) **rudis** (Kze.) C. Chr. Ind. 289. 1905 — Mexico-Peru.
Polypodium Kze. Linn. 13. 133. 1839. HB. 307; Glaphyropteris Pr. 1848; Phegopteris Mett. 1856. n. 28; Nephrodium Diels 1899. NPfl. 171.

(D) **rufa** (Poir.) C. Chr. Ind. 290. 1905 — Chile. — Polypodium (ruffum) Poir. Enc. **5**. 532. 1804; Nephrodium Desv. 1827; Aspidium Mett. 1870.

(P) **rufescens** (Bl.) C. Chr. Ind. 290. 1905 — Ceylon. Java. Queensland. N. Caledonia. Aspidium Bl. Enum. 168. 1828; Polypodium Bl. Fl. Jav. Fil. 194 t. 91. 1829. HB. 309; Polystichum? Moore 1858; Phegopteris Mett. 1864. Chr. 276; Nephrodium Diels 1899. NPfl. 175. Phegopteris munita Mett. n. 18. [1858.

(D) **rufostraminea** (Christ) C. Chr. Ind. 290. 1905 — China. Aspidium Christ, Bull. Soc. Fr. **52** Mém. I. 36. 1905.

(L) **rupestris** (Kl.) C. Chr. Ind. 290. 1905 — Columbia. Venezuela. Leptogramme Kl. Linn. **20**. 415. 1847; Gymnogramme Kze. 1850; Phegopteris Mett. 1856. Pheg. n. 25. Gymnogramme diplazioides pt. HB. 377.

(P) **rustica** (Fée) C. Chr. Ind. 290. 1905 — Guadeloupe. Costa Rica. Phegopteris Fée, 11 mém. 55 t. 13 f. 1. 1866; Polypodium Bak. 1867. HB. 306.

(D) **Sabaei** (Franch. et Sav.) C. Chr. Ind. 290. 1905 — Japonia. Aspidium Franch. et Sav. Enum. pl. Jap. **2**. 259, 632. 1879.

(D) **sagenioides** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Birma. Malesia. Ins. Philipp. — Aspidium Mett. Pheg. u. Asp. 113 n. 269. 1858. Chr. 255; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 271. NPfl. 173. N. melanopus Hk. 1862; Lastrea Bedd. 1865. Aspidium fuscipes pt. Wall. 1828 (nomen); Lastrea Moore 1858; Bedd. 1883 — ? Aspidium nigricaule Fée 1850-52; Lastrea Moore 1858.

—* Phegopteris obscura Fée 1850-52 (non Aspidium Bl. 1828); Polypodium Hk. 1862. HB. 308; Nephrodium Diels 1899. NPfl. 173. Phegopteris philippinensis var. 2 Mett. n. 55. 1858.

(P) **sagittata** (Sw.) C. Chr. Ind. 290. 1905 — India occ. — Polypodium Sw. Prod. 132. 1788; P. hastaefolium Sw. 1801. HB. 304; Phegopteris J. Sm. 1857; Nephrodium Diels 1899. NPfl. 168; Polypodium hastatum Sw. 1806; Aspidium reptans var. 2. hastaefolia Mett. n. 237. 1858.

(C) **sagittifolia** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Java. Perak. Aspidium Bl. Enum. 153. 1828; Nephrodium Moore 1858. HB. 294.

(C) **sakayensis** (Zeiller) C. Chr. Ind. 290. 1905 — Perak. Nephrodium Zeiller, Bull. Soc. Fr. **32**. 75. 1885.

(M) **salicifolia** (Wall.) C. Chr. Ind. 290. 1905 — Penang. Singapore. Meniscium Wall. 1828 (nomen); Hk. Ic. pl. t. 990. 1854. HB. 391; Phegopteris Mett. 1859; Nephrodium Diels 1899. NPfl. 178.

(D) **Salvini** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Guatemala. Nephrodium Bak. Syn. 274. 1867.

(P) **samoensis** C. Chr. Ind. 290. 1905 — Samoa. Polypodium paleaceum Powell; Bak. Syn. 505. 1874 (non Hk. fil. 1847).

(D) **sancta** (L.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India occ. Guatemala. Quito. — Acrostichum L. Syst. Nat. ed. X. **2**. 1320. 1759; Polypodium Sw. 1788; Phegopteris Fée 1850-52; Aspidium Mett. n. 181. 1858. Chr. 252; Lastrea J. Sm. 1866; Nephrodium Bak. 1867. HB. 267. NPfl. 172.

(D) **sancti-gabrieli** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Amazonas. Polypodium Hk. sp. **4**. 233. 1862; Nephrodium Bak. 1870. HB. 493; NPfl. 168; Aspidium Fée 1872-73.

(P) **sandwicense** (Hk. et Arn.) C. Chr. Ind. 290. 1905 — Ins. Hawaii. Fiji. Pitcairn. — Polypodium Hk. et Arn. Bot. Beech. Voy. 105. 1832. HB. 312; Phegopteris Mann 1868.

(D) **Sangnellii** (Bak.) C. Chr. Ind. 291. 1905 — N. Caledonia.
Nephrodium Bak. Ann. Bot. 5. 318. 1891.

(P) **Sauvallei** C. Chr. Ind. 291. 1905 — Cuba. — Polypodium Wrightii Bak. Syn. 304. 1867; Nephrodium Diels 1899. NPfl. 168 (non Dryopteris infra).

(D) **savaiiensis** (Bak.) C. Chr. Ind. 291. 1905 — Samoa.
Nephrodium Bak. Ann. Bot. 5. 318. 1891; Aspidium Christ 1896.

scabriuscula Dav. Bot. Gaz. 21. 256. 1896 (syn.) = D. Davenportii.

(D) **scabrosa** (Kze.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — India austr.
Aspidium Kze. Linn. 24. 286. 1851. Mett. n. 139; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 283. NPfl. 177; Polypodium nigrocarpum Bedd. 1863-65; Gymnothalamium aspidioides Zenker msc.

(L) **Scallanii** (Christ) C. Chr. Ind. 291. 1905 — China.
Aspidium Christ, Bull. Soc. Bot. It. 1901. 296.

(D) **scalpturoides** (Fée) C. Chr. Ind. 291. 1905 — Cuba. Jamaica.
Phegopteris Fée, 11 mém. 51. 1866. Aspidium rigidulum Mett.; Kuhn 1869; Nephrodium Bak. 1874. HB. 496; Dryopteris O. Ktze. 1891.

(C) **Schaffneri** (Fée) C. Chr. Ind. 291. 1905 — Mexico. — Nephrodium Fée, 8 mém. 108. 1857; 10 mém. 44 t. 43. NPfl. 179; Aspidium Kuhn 1869.

(D) **Schimperianum** (Hochst.) C. Chr. Ind. 291. 1905 — Abyssinia.
Aspidium Hochst.; A.Br. Flora 1841. 708; Mett. Pheg. u. Asp. 63. n. 149. 1858; Polystichum Keys. 1873. (Conf. D. filix mas *Schimperiana).

(C) **scolopendrioides** (L.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — India occ.
Polypodium L. sp. 2. 1085. 1753; Goniopteris Pr. 1836; Aspidium Mett. n. 235. var. 2 subpinnata 1858. Chr. 243; Nephrodium Hk. 1862. HB. 288. NPfl. 178 — Goniopteris strigosa Fée 1866; Nephrodium Jenm. 1896 (non Desv. 1827).

(D) **securidiformis** (Hk.) C. Chr. Ind. 291. 1905 — Africa trop.
Nephrodium subquinquefidum γ securidiforme Hk. sp. 4. 130. 1862; Aspidium securidiforme Mett.; Kuhn, Fil. Afr. 141. 1868; Nephrodium Diels 1899. NPfl. 175; Nephrodium subquinquefidum pt. HB. 281.

Seemanni O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Polystichum.

(D) **Seemanni** (Bak.) C. Chr. Ind. 291. 1905 — Ins. Hawaii. (Oahu).
Nephrodium Bak. Ann. Bot. 5. 320. 1891.

semibipinnata O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Aspidium.

semicordata O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Cyclopeltis.

(D) **semihastata** (Kze.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Peru.
Aspidium Kze. Linn. 9. 91. 1834. Mett. n. 179; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 260. NPfl. 168.

(D) **semilunata** (Sod.) C. Chr. Ind. 291. 1905 — Ecuador.
Nephrodium Sod. Rec. 46. 1883; Cr. vasc. quit. 245. 1893.

(D) **senanensis** (Franch. et Sav.) C. Chr. Ind. 291. 1905 — Japonia.
Aspidium Franch. et Sav. En. pl. Jap. 2. 241, 636. 1879.

(C) **serra** (Sw.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — India occ.
Polypodium Sw. Prod. 132. 1788; Aspidium Sw. 1801. Mett. n. 226. Chr. 246; Nephrodium Desv. 1827. HB. 289. NPfl. 178 (non Hk. 1862); Lastrea Pr. 1836; ? Cyclodium A. Brongn. 1846 — Aspidium augescens Link 1841; Nephrodium J. Sm. 1846; ? Lastrea J. Sm. 1856 — Tectaria incisa Cav. 1802; Aspidium Sw. 1806.

(M) **serrata** (Cav.) C. Chr. Ind. 291. 1905 — Amer. trop.
Meniscium Cav. Descripc. 548. 1803. HB. 392; Phegopteris Mett. 1864;

Nephrodium Keys. 1873. NPfl. 179 — Meniscium palustre Raddi 1819; Polypodium Lowe 1858; Phegopteris Mett. 1859.

(C) **serrulata** (Sw.) C. Chr. Ind. 292. 1905 — Jamaica.
Polypodium Sw. Schrad. Journ. **1800**[2]. 25. 1801; Goniopteris J. Sm. 1857; Aspidium Mett. n. 252. 1858; Nephrodium Jenm. 1896. Polypodium Lunanianum Hew. 1838.

(D) **setigera** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Japonia. China. India bor.-Malesia-Polynesia-Austr. trop. — Cheilanthes Bl. Enum. 138. 1828; Hypolepis Hk. 1858; Aspidium Kuhn 1869. Chr. 265; Lastrea Bedd. 1870; Nephrodium Bak. 1874. HB. 284 (pt.). NPfl. 177. Aspidium uliginosum Kze. 1847. Mett. n. 171. Polypodium tenericaule Wall. 1828; Lastrea Moore 1858; Nephrodium Hk. 1862 (excl. t. 259); Aspidium Thwait. 1864.

setosa O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. Millettii.

(P) **setosa** (Pr.) C. Chr. Ind. 292. 1905 — Ins. Philipp.
Lastrea Pr. 1849; Polypodium hirtum Pr. Rel. Haenk. **1**. 27 (non Sw. 1788); Phegopteris Christ 1898; Nephrodium Copeland 1905 (non Hk. 1862).

(D) **setulosa** (Bak.) C. Chr. Ind. 292. 1905 — Tonkin.
Nephrodium Bak. JoB. **1890**. 265.

(D) **Sewellii** (Bak.) C. Chr. Ind. 292. 1905 — Madagascar.
Nephrodium Bak. Journ. Linn. Soc. **15**. 418. 1876; Aspidium Kuhn 1879.

(P) **Sharpiana** (Bak.) C. Chr. Ind. 292. 1905 — Madagascar.
Polypodium Bak. JoB. **1880**. 369.

(D) **Sheareri** (Bak.) C. Chr. Ind. 292. 1905 — China.
Nephrodium Bak. JoB. **1875**. 200. N. isolatum Bak. 1880.

(D) **shikokiana** (Mak.) C. Chr. Ind. 292. 1905 — Japonia. — Aspidium Mak. 1892 (nomen); Nephrodium Mak. Bot. Mag. Tokyo **13**. 62. 1899.

(D) **siambonensis** (Hier.) C. Chr. Ind. 292. 1905 — Argentina.
Aspidium Hier. Engl. Jahrb. **22**. 372. 1896.

(D) **Sieboldii** (van Houtte) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Japonia. China.
Aspidium van Houtte; Mett. Fil. Lips. 87 t. 20 f. 1-4. 1856; Asp. n. 124. Chr. 250; Pycnopteris Moore 1855; Lastrea Moore 1857; Nephrodium Hk. 1859. HB. 261. NPfl. 168; Polystichum Keys. 1873; Lastrea podophylla J. Sm. 1857; Aspidium Lowe 1857 (non Hk. 1853).

siifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(D) **sikkimensis** (Bedd.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Sikkim.
Polystichum Bedd. Ferns br. Ind. t. 127. 1866; Aspidium Bak. 1867. HB. 256; Nephrodium Clarke 1880; Lastrea Bedd. 1883.

(G) **silvatica** (Pappe et Raws.) C. Chr. Ind. 292. 1905 — Natal. Kamerun.
Goniopteris Pappe et Raws. Syn. Fil. Afr. austr. 39. 1858. Gymnogramme unita Kze. Linn. **18**. 115. 1844 (non Dryopteris infra); Phegopteris Mett. n. 42. 1858; Polypodium Hk. 1863. HB. 317; Goniopteris J. Sm. 1875 — G. patens Fée 1850-52 (non Dryopteris supra); Nephrodium J. Sm. 1875.

(C) **simillima** C. Chr. Ind. 292. 1905 — Borneo.
Nephrodium simulans Bak. JoB. **1888**. 325 (non 1874).

Simonsii O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

(M) **simplex** (Hk.) C. Chr. Ind. 292. 1905 — China austr. Formosa.
Meniscium Hk. Lond. JoB. **1**. 294 t. 11. 1842. HB. 390; Polypodium Lowe 1858; Phegopteris Mett. 1859; Aspidium Hance 1873; Nephrodium Diels 1899. NPfl. 177.

(D) **simplicissima** (Christ) C. Chr. Ind. 293. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. II. **4**. 959. 1904.
simulans O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium Thwaitesii.
(D) **simulata** Dav. Bot. Gaz. **19**. 497. 1894 (syn.) — Amer. bor. atlant. — Aspidium Dav. l. c. 495. Nephrodium Dav. 1894. NPfl. 172; Lastrea Dav. 1894.
(D) **singalanensis** (Bak.) C. Chr. Ind. 293. 1905 — Sumatra. Perak.
Nephrodium Bak. JoB. **1880**. 212; Lastrea Bedd. 1892.
singaporiana O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.
(C) **Sintenisii** (Kuhn) Urban, Symb. Ant. **4**. 19. 1903 — Puerto Rico.
Aspidium Kuhn (Bak. Ann. Bot. **5**. 327. 1891, nomen); Christ, Engl. Jahrb. **24**. 119. 1897; Nephrodium Bak. 1891 (nomen).
(C) **Skinneri** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Guatemala. Ecuador.
Aspidium Hk. Ic. pl. t. 924. 1854. Mett. n. 224; Nephrodium Moore 1858. HB. 287. (Chr. 244) — Aspidium Francoanum Fourn. 1872.
Sloanei O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. patens* invisa.
(C) **sophoroides** (Thbg.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Japonia. China. Formosa. — Polypodium Thbg. 1794; Aspidium Sw. 1801. Chr. 246; Nephrodium Desv. 1827. HB. 289. NPfl. 178. Polypodium unitum Thbg. Fl. Jap. 336. 1784 (non L. 1753).
(D) **sparsa** (Ham.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — China austr. India bor.-Ceylon. Malesia. Mauritius. — Polypodium Ham. msc.; Nephrodium Don, Prod. Fl. Nepal. 6. 1825. HB. 276. 498 (excl. var. 3 et 4). NPfl. 174; Aspidium Spr. 1827; Chr. 262; Lastrea Moore 1858; Polystichum Keys. 1873. Aspidium densum Wall. 1828. Mett. n. 153; Lastrea Pr. 1836.
—* Aspidium purpurascens Bl. 1828; Nephrodium Hk. 1862.
—* Aspidium nitidulum Wall. 1828; Kuhn, Linn. **36**. 117. 1869 (non Kze. Mett. 1858); Lastrea Bedd. 1876.
—* Aspidium obtusissimum Mett.; Kuhn 1869.
—* Aspidium undulatum Thwaites 1864 (non Afz.; Sw. 1801); Lastrea Bedd. 1863-65; Nephrodium Bak. 1867. HB. 276. NPfl. 174; Dryopteris O. Ktze. 1891.
(D) **Spekei** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Afr. occ. Ins. Comorae.
Nephrodium Bak. Syn. 263. 1867. NPfl. 172; Aspidium Kuhn 1868.
(P) **sphaeropteroides** (Bak.) C. Chr. Ind. 293. 1905 — Yunnan.
Polypodium Bak. Kew Bull. **1895**. 55; Phegopteris Christ 1899.
(D) **spinulosa** (Müll.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Europa. Asia septentr.-Himalaya. Amer. bor. — Polypodium Müller, (Fl. Fridrichsd. 193 f. 2. 1767) Fl. dan. t. 707. 1777; Aspidium Sw. 1801. Mett. n. 136. Chr. 261; Polystichum Lam. et DC. 1805; Nephrodium Strempel 1822. HB. 275 (excl. γ et δ). NPfl. 174; Lastrea Pr. 1836. Polystichum spinosum Roth 1799; Lastrea Newm. 1844; Lophodium Newm. 1851 — Polystichum spinulosum × dilatatum Sanio 1884.
—* Aspidium intermedium Mühl.; Willd. 1810; Nephrodium Pr. 1825; Lastrea Pr. 1836; Dryopteris A. Gray 1848. (Amer. bor.).
—* Polypodium dilatatum Hoffm. 1795; Polystichum Schum. 1803; Aspidium Sm. 1804. Mett. n. 137; Nephrodium Desv. 1827; Lastrea Pr. 1836; Dryopteris A. Gray 1848. Polypodium aristatum Vill. 1789 (non Forst. 1786) — var. patuloides Christ, Bull. Soc. Fr. **52** Mém. I. 41. 1905. (China merid.).

(D) **splendens** (Hk.) O.Ktze. Rev. Gen. Pl. **2**. 813. 1891 — China austr. India bor. Malacca. — Nephrodium Hk. sp. **4**. 126. 1862 (excl. β). HB. 282. NPfl. 177; Lastrea Bedd. 1865; Aspidium Christ 1897. 259.

Sprengelii O.Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. Balbisii.

(D) **Sprucei** (Bak.) O.Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ecuador. Nephrodium Bak. Syn. 269. 1867. NPfl. 172.

(P) **squamaestipes** (Clarke) C. Chr. Ind. 294. 1905 — India bor. China. Polypodium appendiculatum var. squamaestipes Clarke, Tr. Linn. Soc. II. Bot. **1**. 543 t. 79 f. 2. 1880. Polypodium auriculatum Wall. 1828 (nomen); Hk. sp. **4**. 237. 1862. HB. 306 (non L. 1753); Phegopteris J. Sm. 1875. Chr. 271; Nephrodium Diels 1899. NPfl. 171 — Polypodium appendiculatum Bedd. 1868. (an HB. 306. 1867?); Phegopteris J. Sm. 1875; Aspidium Christ 1902 (non Dryopteris supra).

(D) **squamigera** (Hk. et Arn.) O.Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Hawaii. Nephrodium Hk. et Arn. Bot. Beech. Voy. 106. 1832. HB. 280. NPfl. 174; Lastrea Brack. 1854; Aspidium Mann 1868.

(D) **squamiseta** (Hk.) O.Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Africa trop. Réunion. Nephrodium Hk. sp. **4**. 140 t. 268. HB. 278. NPfl. 174; Aspidium Kuhn 1867.

(D) **squamosissima** (Sod.) C. Chr. Ind. 294. 1905 — Ecuador. Nephrodium Sod. Cr. vasc. quit. 256. 1893.

(S) **stegnogramme** (Bl.) C. Chr. Ind. 294. 1905 — India bor. Ceylon. Java. Gymnogramme Bl. Fl. Javae Fil. 98 t. 44. 1828; Phegopteris Mett. 1856. n. 54. Chr. 268; Nephrodium Diels 1899. NPfl. 181; Stegnogramma aspidioides Bl. 1828 (non Dryopteris supra); Gymnogramme Hk. 1854. HB. 378; Syneuron J. Sm. 1854; Grammitis Hk. 1864.

(S) **stegnogrammoides** (Bak.) C. Chr. Ind. 294. 1905 — Ins. Hawaii. Polypodium Bak. 1867. HB. 317; Phegopteris Luerss. 1875; Goniopteris J. Sm. 1877; Polypodium polycarpon Hk. et Arn. Bot. Beech. Voy. 104. 1832 (non Sw. 1801); Phegopteris Hill. 1888; Nephrodium Diels 1899. NPfl. 179; Stegnogramme Sandwicense Brack. 1854; Polypodium Hk. 1863 (sp. **5**. 5, non **4**. 267 = Hk. et Arn. 1832); Goniopteris J. Sm. 1875; Phegopteris microdendron Eat. 1868.

(D) **stenobasis** C. Chr. Ind. 294. 1905 — Ins. Philipp. Celebes. Lastrea attenuata J. Sm. 1841 (nomen, non Brack. 1854); Nephrodium Fée 1850-52 (nomen). HB. 263. NPfl. 171; Aspidium Kze.; Mett. Pheg. u. Asp. 96 n. 233. 1858 (non Sw. 1801); Dryopteris O.Ktze. 1891.

(C) **stenolepis** (Bak.) C. Chr. Ind. 294. 1905 — China austr. — Polypodium Bak. Kew Bull. **1898**. 231; Aspidium Christ 1899; A. yunnanense Christ 1898.

(D) **stenophylla** (Sod.) C. Chr. Ind. 294. 1905 — Ecuador. — Nephrodium Sod. Rec. 44. 1883; Cr. vasc. quit. 229. 1893 (non Bak. 1884 nec 1886).

(C) **stipellata** (Bl.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Java. Aspidium Bl. Enum. 152. 1828; Nephrodium Moore 1858. HB. 294.

(D) **stipitata** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Ins. Ladrones. Aspidium Mett. Pheg. u. Asp. 60 n. 143. 1858; Lastrea Moore 1858; Nephrodium Bak. 1867. HB. 278.

(G) **straminea** (Bak.) C. Chr. Ind. 294. 1905 — Venezuela. Polypodium Bak. Syn. 316. 1867; Phegopteris Salom. 1883; Polypodium salicifolium Hk. sp. **4**. 242. 1862 (non Vahl 1807 nec Willd. 1810).

(C) **striata** (Schum.) C. Chr. Ind. 294. 1905 — Africa trop. Aspidium Schum. Vid. Selsk. Afh. **4**. 230 (seors. 456). 1827! Polypodium

pallidivenium Hk. sp. 5. 8. 1863; Nephrodium Bak. 1867. HB. 290 NPfl. 179; Dryopteris O. Ktze. 1891.

(D) **strigilosa** Dav. Bot. Gaz. **21**. 257. 1896 (syn.) — Mexico.
Aspidium Dav. l. c. 256; Nephrodium Dav. 1896; Lastrea Dav. 1896.

(D) **strigosa** (Willd.) C. Chr. Ind. 295. 1905 — Madagascar. Ins. Mascarenae.
Aspidium Willd. sp. 5. 249. 1810; Nephrodium Desv. 1827; Lastrea Pr. 1836 — Aspidium pulchrum Bory; Willd. 1810; Nephrodium Desv. 1827; Lastrea Pr. 1836. — Conf. Kuhn, v. Deck. Reis. 3^3. Bot. 37-38.

(P) **Sturmii** (Phil.) C. Chr. Ind. 295. 1905 — Chile.
Phegopteris Phil. Anal. Univ. Chil. **94**. 358. 1896.

(D) **subarborea** (Bak.) C. Chr. Ind. 295. 1905 — Malesia-Ins. Philipp. N. Guinea.
Polypodium Bak. Journ. Linn. Soc. **24**. 259. 1887 — Nephrodium megaphyllum Bak. Journ. Linn. Soc. **22**. 227. 1886; Lastrea Bedd. 1892; Aspidium Christ 1898 (non Mett. 1864); A. Filix mas var. Chr. 257 — Alsophila Warburgii Christ 1897. NPfl. 138.

subbiaurita O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. orientalis var.

(P) **subconnexa** (Christ) C. Chr. Ind. 295. 1905 — Batjan.
Phegopteris Christ in Warb. Monsunia **1**. 83. 1900.

(P) **subcoriacea** C. Chr. Ind. 295. 1905 — Africa occ. trop.
Polypodium Vogelii Hk. sp. **4**. 271. 1862. HB. 313; Phegopteris Kuhn 1868 (non Dryopteris infra).

(D) **subcrenulata** (Bak.) C. Chr. Ind. 295. 1905 — Madagascar. — Nephrodium Bak. Journ. Linn. Soc. **16**. 202. 1877; Ic. pl. t. 1660; Aspidium Kuhn 1879.

(C) **subcuneata** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Cayenne.
Nephrodium Bak. Fl. bras. 1^2. 487. 1870. HB. 503.

(P) **subdecussata** (Christ) C. Chr. Ind. 295. 1905 — Costa Rica.
Aspidium Christ, Bull. Boiss. II. **4**. 960. 1904.

(P) **subdryopteris** (Christ) C. Chr. Ind. 295. 1905 — Costa Rica.
Phegopteris Christ, Prim. Fl. costar. **3**. 36. 1901.

(D) **subexaltata** (Christ) C. Chr. Ind. 295. 1905 — Formosa.
Aspidium Christ, Bull. Boiss. II. **4**. 616. 1904.

(D) **subfusca** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Cayenne.
Nephrodium Bak. Syn. 267. 1897. NPfl. 271.

(D) **subglabra** (Sod.) C. Chr. Ind. 295. 1905 — Ecuador.
Nephrodium Sod. Cr. vasc. quit. 259. 1893.

subglandulosa O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. Helliana.

(D) **subgranulosa** C. Chr. Ind. 295. 1905 — Madagascar.
Nephrodium granulosum Bak. JoB. **1891**. 4 (non Dryopteris supra).

(P) **subincisa** (Willd.) Urban, Symb. Ant. **4**. 19. 1903 — Amer. trop. Chile.
Polypodium Willd. sp. **5**. 202. 1810; Lastrea Pr. 1836; Phegopteris Fée 1850-52. Mett. n. 65. Chr. 276; Nephrodium Christ 1903; N. villosum pt. HB. 286 — Polypodium spectabile Klf. 1824; Phegopteris Fée 1850-52. Mett. n. 62; Nephrodium Hier. 1904 (non Hk. 1862).
—* Alsophila martinicensis Spr. 1822; Phegopteris Fourn. 1872.
—* Polypodium vastum Kze. 1834; Phegopteris Mett. 1859; Nephrodium Hier. 1904 (non Bak. 1867).
—* Polypodium Karstenianum Kl. 1847; Phegopteris Mett. n. 63. 1858; Polypodium barbatum Desv. 1827 (non Spr. 1827).
—* Phegopteris Lechleri Mett. 1859.

—* Phegopteris canescens Mett. n. 64. 1858; Polypodium Kze.; Hk. 1862 (non Bl. 1828); Phegopteris cana Mett. 1864.

—* Phegopteris vestita Phil. 1857; Polypodium Hk. 1862 (non Forst. 1786).

—* Polypodium Berteroanum Hk. 1862 (non Spr. 1827).

Obs. Subspecies omnes f. auctt. nonnullis species distinctae sunt, f. auctt. aliis = formae exindusiatae D. villosae.

(C) **subintegra** (Bak.) C. Chr. Ind. 296. 1905 — Ecuador.
Polypodium Bak. JoB. 1877. 164; Nephrodium Sod. 1883.

(D) **subjuncta** (Bak.) C. Chr. Ind. 296. 1905 — Samoa.
Nephrodium Bak. Ann. Bot. 5. 321. 1891; Aspidium Christ 1896.

(D) **submarginalis** (Langsd. et Fisch.) C. Chr. Ind. 296. 1905 — Mexico-Brasilia-Uruguay-Peru. — Polypodium Langsd. et Fisch. Ic. Fil. 12 t. 13. 1810; Willd. sp. 5. 202. (n. 137). 1810; Phegopteris J. Sm. 1857. Polypodium caripense H.B.Willd. sp. 5. 202. (n. 138). 1810; Aspidium Mett. 1856. n. 220. Chr. 254; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 265. NPfl. 171; Polystichum Keys. 1873; Dryopteris O. Ktze. 1891 — Nephrodium Tarapotense Hk. 1862.

(D) **subobliquata** (Bak.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — Brasilia. Guiana. Columbia. — Polypodium Hk. sp. 4. 240. 1862; Nephrodium Bak. 1867. HB. 261. NPfl 168; Aspidium Fée 1872-73. — (Conf. Aspidium emarginatum).

(P) **subobscura** (Christ) C. Chr. Ind. 296. 1905 — Yunnan.
Phegopteris Christ, Bull. Boiss. 6. 836. 1898.

subquinquefida O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = D. protensa.

(D) **subsericea** (Mett.) O. Ktze. Rev. Gen. Pl. 2. 813. 1891 — N. Caledonia.
Aspidium Mett. Ann. sc. nat. IV. 15. 74. 1861; Nephrodium Bak. 1867. HB. 281. NPfl. 175; Leucostegia Fourn. 1873.

(L) **subsimilis** (Hk.) C. Chr. Ind. 296. 1905 — Fernando Po.
Gymnogramme Hk. sp. 5. 142 t. 293. 1864. (HB. 313); Phegopteris Mett. 1868; Leptogramma J. Sm. 1875.

(P) **subspinosa** C. Chr. Ind. 296. 1905 — Samoa.
Polypodium muricatum Powell; Bak. Syn. 506. 1874 (non L. 1753). Nephrodium Diels 1899. NPfl. 179 — Goniopteris glandulifera Brack. 1854 (non Dryopteris supra); Aspidium Mett. 1864; Polypodium Brackenridgei Hk. 1863 (non Dryopteris supra).

(D) **subspinulosa** (Christ) C. Chr. Ind. 296. 1905 — Korea.
Aspidium Christ, Bull. Boiss. II. 2. 829. 1902.

(D) **subthelypteris** (Christ) C. Chr. Ind. 296. 1905 — China.
Aspidium Christ in litt.; A. melanorhizum Christ, Bull. Soc. bot. It. 1901. 295 (non Desv. 1827).

(D) **subtriangularis** (Hope) C. Chr. Ind. 296. 1905 — Assam.
Nephrodium Hope, JoB. 1890. 327. (f. Bedd. = D. filix mas var.).

subtriphylla O. Ktze. Rev. Gen. Pl. 2. 813. 1891 = Aspidium.

(D) **subtripinnata** (Miq.) O. Ktze. Rev. Gen. Pl. 2. 811. 1891 — China. Korea. Japonia. — Aspidium Miq. Ann. Lugd. Bat. 3. 179. 1867; Nephrodium Bak. 1868. NPfl. 174; N. chinense Bak. 1867. HB. 278.

(P) **sulcinervia** (Hier.) C. Chr. Ind. 296. 1905 — Africa orient.
Aspidium Hier. in Engl. Pflw. Ostafr. C. 85. 1895.

(D) **supina** (Sod.) C. Chr. Ind. 296. 1905 — Ecuador. Columbia.
Nephrodium Sod. Cr. vasc. quit. 241. 1893.

(P) **sylvicola** (Bak.) C. Chr. Ind. 297. 1905 — Columbia.
Polypodium Bak. JoB. **1881**. 205.

(D) **syrmatica** (Willd.) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — India bor.-Malesia. Ins. Philipp. — Aspidium Willd. sp. **5**. 237. 1810 (patria erronea). Chr. 255; Nephrodium Desv. 1827. HB. 272. NPfl. 173; Lastrea Moore 1858. Aspidium spectabile Bl. 1828. Mett. n. 268; Lastrea J. Sm. 1841; Nephrodium Hk. 1862.

(D) **tablana** (Christ) C. Chr. Ind. 297. 1905 — Mexico.
Aspidium Christ, Bull. Boiss. II. **5**. 727. 1905.

(C) **taiwanensis** C. Chr. Ind. 297. 1905 — Formosa.
Aspidium lobulatum Christ, Bull. Boiss. II. **4**. 614. 1904 (non Bl. 1828).

(G) **tannensis** C. Chr. Ind. 297. 1905 — N. Hebridae.
Polypodium excelsum Bak. Syn. 505. 1874 (non Desv. 1827). ? Aspidium glanduliferum Kuhn 1869 (non Dryopteris supra).

(P) **tarapotensis** (Bak.) C. Chr. Ind. 297. 1905 — Peru.
Polypodium Bak. Syn. 505. 1874.

(C) **tenebrica** (Jenm.) C. Chr. Ind. 297. 1905 — Jamaica.
Nephrodium Jenm. JoB. **1882**. 326.

(D) **tenera** (R.Br.) C. Chr. Ind. 297. 1905 — Australia. — Nephrodium R.Br. Prod. Fl. N. Holl. 149. 1810; Aspidium Spr. 1827; Lastrea Moore 1858.

tenuifolia O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. tenuifrons.

(D) **tenuifolia** (Pr.) C. Chr. Ind. 297. 1905 — Brasilia.
Lastrea Pr. Epim. 37. 1849. ? Polypodium aspidioides Pr. Del. Prag. **1**. 170. 1822 (f. Bak. = Polystichum aculeatum); ? Lastrea Pr. 1849; Moore 1858; ? Phegopteris Fée 1850 52 (non Dryopteris supra). Aspidium alsophileum Kze. 1837 (nomen); A. alsophilaceum Kze.; Mett. Pheg. u. Asp. 90. n. 218. 1858; Nephrodium Bak. 1870. HB. 495; Dryopteris O.Ktze. 1891.

(D) **tenuifrons** C. Chr. Ind. 297. 1905 — Fiji. — Lastrea tenuifolia Brack. Expl. Exp. **16**. 199. 1854 (non Pr. 1849); Nephrodium Hk. 1862. HB. 278. NPfl. 174; Aspidium Luerss. 1871; Dryopteris O. Ktze. 1891.

ternata O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Aspidium.

tetragona O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. pseudotetragona.

(D) **tetragona** (Sw.) Urban, Symb. Ant. **4**. 20. 1903 — Florida-Brasilia-Peru. Polypodium Sw. Prod. 132. 1788. HB. 317; Aspidium Sw. 1801; Goniopteris Pr. 1836; Phegopteris Mett. 1856. n. 44. Chr. 268; Nephrodium Keys. 1873. Polypodium androgynum Poir. 1804; Nephrodium Diels 1899. NPfl. 179 — Goniopteris sp. plur. Fée 1850-52, 1866 — Monogonia palustris Pr. 1836 (f. Milde, Fil. Eur. 45).

(D) **thelypteris** (L.) A. Gray, Man. ed. I. 630. 1848 — Europa. Algeria. Asia temp. Himalaya. China merid. India austr. Amer. septent. atlant. Africa trop. et austr. N. Zealand. — Acrostichum L. sp. **2** 1071. 1753; Polypodium Weis 1770; Polystichum Roth 1799; Bernh. 1799; Aspidium Sw. 1801. Mett. n. 267. Chr. 253; Athyrium Spr. 1804; Nephrodium Strempel 1822. HB. 271. NPfl. 172; Lastrea Bory 1826; Hemestheum Newm. 1851; Parment. 1899; Thelypteris palustris Schott 1834; Lastrea J. Sm 1857. —* Aspidium Thelypteris var. squamigerum Schlecht. Adumbr. 23 t. 11. 1825; Lastrea squamulosa Pr. 1836; Nephrodium Hk.fil. 1854; Aspidium squamigerum Fée 1857; Lastrea invisa Armstr. 1882 — L. Fairbankii Bedd. 1867 (India austr.).

(D) **thibetica** (Franch.) C. Chr. Ind. 298. 1905 — Tibet orient.
Aspidium Franch. Nouv. Arch. Mus. II. **10**. 118. 1887; Nephrodium (tibeticum) Bak. 1891; Lastrea Bedd. 1892.
Thomsonii O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = Polystichum.
(P) **Thomsonii** (Jenm.) C. Chr. Ind. 298. 1905 — Jamaica.
Polypodium Jenm. JoB. **1886**. 272.
Thwaitesii O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 = D. deparioides var.
(M) **Thwaitesii** (Hk.) C. Chr. Ind. 298. 1905 — India austr. Ceylon.
Meniscium Hk. Fil. exot. t. 83. 1859. HB. 391; Phegopteris Mett. 1859.
(P) **tijuccana** (Raddi) C. Chr. Ind. 298. 1905 — Brasilia. Guiana.
Polypodium (tyucanum); Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. bras. **1**. 25 t. 37. 1825. HB. 305; Phegopteris Fée 1850-52; Mett. n. 36; Nephrodium Diels 1899. NPfl. 170.
(D) **tokyoensis** (Matsum.) C. Chr. Ind. 298. 1905 — Japonia. Korea.
Aspidium Matsum. hb.; Nephrodium Mak. Bot. Mag. Tokyo **12**. 87. 1898. Aspidium transitorium Christ 1899.
(D) **tomentella** C. Chr. Ind. 298. 1905 — Ins. Principis (Guinea).
Aspidium Nigritianum Mett.; Kuhn, Fil. Afr. 138. 1868; Nephrodium Bak. 1874. HB. 499; Dryopteris O. Ktze. 1891 (non supra).
(D) **tomentosa** (Thouars) O. Ktze. Rev. Gen. Pl. **2**. 813. 1891 — Tristan d'Acunha. Madagascar. Ins. Mascarenae. — Polypodium Thouars, Fl. Trist. d'Ac. 32 t. 3. 1804; Nephrodium Desv. 1827. HB. 269. NPfl. 172; Lastrea Moore 1858; Aspidium Kuhn 1879. A. riparium Bory; Willd. 1810; Nephrodium Desv. 1827; Lastrea Pr. 1836. Polypodium thelypteroides Sieb.; Spr. 1827; Phegopteris Fée 1850-52; Aspidium Mett. n. 194. 1858; Nephrodium Hk. 1862. — Conf. Kuhn, v. Deck. Reis. **3**[3]. Bot. 38.
(D) **Trianae** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Columbia. Mexico.
Aspidium Mett. Ann. sc. nat. V. **2**. 243. 1864; Nephrodium Bak. 1867. HB. 286. Aspidium extensum Fée 1850-52 (non Bl. 1828).
triangula O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Polystichum.
(D) **tricholepis** (Bak.) C. Chr. Ind. 298. 1905 — Guatemala. Costa Rica.
Nephrodium Bak.; Hemsl. Biol. Centr. Amer. **3**. 651. 1885; Aspidium Christ 1905.
(D) **trichophlebia** (Bak.) C. Chr. Ind. 298. 1905 — Madagascar.
Nephrodium Bak. Journ. Linn. Soc. **16**. 201. 1877; Aspidium Kuhn 1879.
(D) **trichophora** (Fée) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Guadeloupe.
Aspidium Fée, 11 mém. 81 t. 23 f. 2. 1866; Nephrodium Bak. 1867. HB. 265 — Aspidium Capitainei Fée et L'Herm. 1866.
(D) **trichopoda** C. Chr. Ind. 298. 1905 — Borneo.
Nephrodium polytrichum Bak. JoB. **1891**. 107 (non Schrad. 1824).
tridens O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Polystichum.
trifoliata O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Aspidium.
(P) **trinidadensis** (Jenm.) C. Chr. Ind. 298. 1905 — Trinidad.
Polypodium Jenm. Gard. Chr. III. **18**. 235. 1895.
(M) **triphylla** (Sw.) C. Chr. Ind. 298. 1905 — China austr. India bor.-Malesia. Ins. Philipp. Queensland. — Meniscium Sw. Schrad. Journ. **1800**[2]. 16. 1801. HB. 391; Phegopteris Mett. 1859. Chr. 269; Nephrodium Diels 1899. NPfl. 178 — Meniscium Parishii Bedd. 1866.
triptera O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Polystichum.
(D) **tristis** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Columbia-Peru. Brasilia.
Polypodium Kze. Linn. **9**. 47. 1834; Aspidium Mett. n. 229. 1858. Chr.

255 (non Bl. 1828); Nephrodium Hk. 1862. HB. 266. NPfl. 169; Lastrea flebilis Moore 1858 — Polypodium monosorum Pr. 1836 (nomen, non Desv. 1811); Lastrea Pr. 1849. Aspidium monostichum Kze.; Mett. n. 228. 1858; Lastrea Moore 1858.

(C) **truncata** (Poir.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — India bor.-Malesia-Australia trop.-Polynesia. Madagascar. Ins. Mascarenae.
Polypodium Poir. Enc. **5**. 534. 1804; Polystichum Gaud. 1827; Aspidium Gaud. 1827. Mett. n. 254. Chr. 248; Nephrodium Pr. 1836. HB. 294. NPfl. 181 — Aspidium caudiculatum Sieb.; Kze. 1851. Mett. n. 250; Nephrodium Pr. 1836.

(P) **tuberculata** (Ces.) C. Chr. Ind. 299. 1905 — N. Guinea. — Nephrodium Ces. Rend. Ac. Napoli **16**. 26, 29. 1877; Polypodium Bak. 1891.

(D) **Tuerckheimii** (Donn. Smith) C. Chr. Ind. 299. 1905 — Guatemala.
Nephrodium Donn. Smith, Bot. Gaz. **12**. 133 t. 11. 1887.

(P) **Ulei** (Christ) C. Chr. Ind. 299. 1905 — Brasilia austr. — Phegopteris Christ in Schwacke, Pl. Nov. Mineiras **2**. 29. 1900; Bull. Boiss. II. **2**. 634. 1902.

undulata O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = D. sparsa*.

(P) **unidentata** (Hk. et Arn.) C. Chr. Ind. 299. 1905 — Ins. Hawaii.
Polypodium Hk. et Arn. Bot. Beech. Voy. 105. 1832. HB. 310; Phegopteris J. Sm. 1866; Nephrodium Diels 1899. NPfl. 174.

(C) **unita** (L.) O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 — India austr. Ceylon-Malesia-Polynesia. Ins. Mascarenae et Sechellae. — Polypodium L. Syst. Nat. ed. X. **2**. 1326. 1759; Aspidium Sieb.; Mett. n. 257. 1858 (non Sw. 1801 et auctt. pl.); Polystichum Gaud. 1827; Nephrodium Bory 1833. Aspidium cucullatum Bl. 1828. Chr. 247; Nephrodium Bak. 1867. HB. 290. NPfl. 178.

unita Maxon, Proc. U. S. Nat. Herb. **23**. 639. 1901 = D. gongylodes.

(D) **Urbani** (Sod.) C. Chr. Ind. 299. 1905 — Ecuador.
Polypodium Sod. Cr. vasc. quit. 301. 1893; Nephrodium Sod. l. c. 638.

(C) **urophylla** (Wall.) C. Chr. Ind. 299. 1905 — India bor. China austr. Malesia-Polynesia-Australia trop. — Polypodium Wall. 1828 (nomen); Hk. sp. **5**. 9. 1863. HB. 315; Goniopteris Pr. 1836; Phegopteris Mett. n. 53. 1858; Nephrodium Keys. 1873. NPfl. 178; Aspidium Christ 1897. 248. Polypodium asperum Pr. 1825 (non L. 1753); Goniopteris Pr. 1836; Aspidium Mett. 1864. A. repandum Bl. 1828 (non Willd. 1810); Polypodium cuspidatum Roxb. 1844 (non Don 1825 nec Pr. 1825).
— Meniscium Kennedyi F. Muell. 1864; Polypodium F. Muell. 1864. HB. ed. I. 314; Phegopteris Luerss. 1873; Goniopteris Bail. 1874 — Polypodium Pinwillii Bak. 1891.
— ? Meniscium cuspidatum Bl. 1828. HB. 392. (NPfl. 178); Phegopteris Mett. n. 52. 1858.

(C) **usitata** (Jenm.) C. Chr. Ind. 299. 1905 — Jamaica.
Nephrodium Jenm. JoB. **1879**. 261.

(D) **valdepilosa** (Bak.) C. Chr. Ind. 299. 1905 — Columbia.
Nephrodium Bak. JoB. **1881**. 204.

varia O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Polystichum.

(C) **varians** (Fée) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Trinidad. Amazonas.
Nephrodium Fée, 11 mém. 88 t. 24 f. 2. 1866. HB. 288.

variolosa O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Aspidium.

vasta O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Aspidium.

(D) **velata** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Cuba.
Aspidium Kze.; Mett. Pheg. u. Asp. 79. n. 190. 1858; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 270.

(D) **vellea** (Willd.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — India occ. — Aspidium Willd. sp. **5**. 255. 1810; Nephrodium Desv. 1827. HB. 265; Lastrea Moore 1858. Nephrodium aureovestitum Hk. 1862; Phegopteris Fée 1866.

(D) **velutina** (A. Rich.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — N. Zealand. ? Fiji. Aspidium A. Rich. Fl. N. Zél. 70. 1832. Mett. n. 275; Nephrodium Hk. fil. 1854. HB. 281. NPfl. 175; Lastrea Brack. 1854.

venulosa O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = D. elata.

(C) **venusta** (Hew.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Jamaica.
Aspidium Hew. Mag. Nat. Hist. II. **2**. 464. 1838; Lastrea J. Sm. 1841; Nephrodium Hew.; Moore 1855. HB. 294.

(D) **Vescoi** (Drake) C. Chr. Ind. 300. 1905 — Tahiti.
Nephrodium Drake, Fl. Polyn. franç. 301. 1893.

vestita O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = D. deflexa.

(D) **Vieillardii** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — N. Caledonia.
Aspidium Mett. Ann. sc. nat. IV. **15**. 75. 1861; Nephrodium Bak. 1867. HB. 282. NPfl. 175.

(D) **vilis** (Kze.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Java.
Aspidium Kze. Bot. Zeit. **1846**. 474. Mett. n. 272; Lastrea Moore 1858; Nephrodium Bak. 1874. HB. 499.

(D) **villosa** (L.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Amer. trop.
Polypodium L. sp. **2**. 1093. 1753; Aspidium Sw. 1801. Mett. n. 274. Chr. 264. Nephrodium Pr. 1825. HB. 286 (pt.). NPfl. 177; Lastrea Pr. 1836. — (Conf. D. subincisa).

(D) **viridescens** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Japonia Liu Kiu. China. — Nephrodium Bak. Syn. 275. 1867. NPfl. 173; Aspidium Miq. 1870.

viscidula O. Ktze. Rev. Gen. Pl. **2**. 811. 1891 = Adenoderris.

(D) **viscosa** (J. Sm.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Perak. Borneo. Ins. Philipp. — Lastrea J. Sm. 1841 (nomen); Nephrodium Bak. Syn. 264. 1867; Aspidium Bedd. 1888.

vivipara O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 = Polystichum.

(G) **vivipara** (Raddi) C. Chr. Ind. 300. 1905 — Brasilia. Ecuador. Peru.
Polypodium Raddi, Pl. bras. **1**. 22 t. 32. 1825; Goniopteris Brack. 1854. Polypodium diversifolium Sw. Vet. Ak. Handl. **1817**. 60. HB. 314 (non Willd. 1810 nec R. Br. 1810); Phegopteris Kuhn 1879. Chr. 267; Nephrodium Diels 1899. NPfl. 179; Polypodium Swartzii Wikstr. 1826 (non Bak. 1868). P. proliferum Klf. 1824; Phegopteris Mett. 1856. Pheg. n. 49; Goniopteris Fée 1869 (non Dryopteris supra) — Aspidium macropus Mett. 1859.

(D) **Vogelii** (Hk.) C. Chr. Ind. 300. 1905 — Africa trop. Madagascar.
Aspidium Hk. Ic. pl. t. 921. 1854. Mett. n. 178; Lastrea Moore 1857; Nephrodium Hk. 1862. NPfl. 175 — N. variabile Hk. 1862; Lastrea J. Sm. 1875. Aspidium speciosum Mett.; Ettingsh. 1865. A. fraternum Mett; Kuhn 1868. Nephrodium subquinquefidum β HB. 281.

(D) **vulcanica** (Bak.) C. Chr. Ind. 300. 1905 — Java.
Nephrodium Bak. Ann. Bot. **8**. 127. 1894.

(P) **waiwaiensis** C. Chr. Ind. 300. 1905 — Fiji. — Polypodium alsophiloides Bak. JoB. **1879**. 297 (non Liebm. 1849 nec Dryopteris supra).

(C) **Wakefieldii** (Bak.) C. Chr. Ind. 301. 1905 — Africa orient. trop.
Nephrodium Bak. Ann. Bot. **5**. 326. 1891; Aspidium Hier. 1895. (f. Bedd. = D. extensa).
(D) **Walkerae** (Hk.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Ceylon.
Polypodium Hk. sp. **4**. 233. 1862. HB. ed. I. 305; Lastrea Bedd. 1870; Nephrodium Bak. 1874. HB. 493; Phegopteris J. Sm. 1875.
(D) **Wardii** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Ins. Sechellae.
Nephrodium Bak. Syn. 500. 1874; Aspidium Kuhn 1879.
(D) **Welwitschii** (Bak.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Africa occ.
Nephrodium Bak. Syn. 274. 1867; Aspidium Kuhn 1868; A. pinnatifido-serratum Mett.; Kuhn, Fil. Afr. 139. 1868.
(D) **Wigmanii** (Racib.) C. Chr. Ind. 301. 1905 — Aru Archip.
Aspidium Racib. Bull. int. Ac. Cracovie **1902**. 61.
(P) **Willsii** (Bak.) C. Chr. Ind. 301. 1905 — Brasilia.
Polypodium Bak. Ann. Bot. **5**. 458. 1891.
(D) **Wilsoni** (Bak.) C. Chr. Ind. 301. 1905 — Uganda.
Nephrodium Bak. Ann. Bot. **5**. 321. 1891; Aspidium Hier. 1895.
(C) **Wrightii** (Mett.) O. Ktze. Rev. Gen. Pl. **2**. 814. 1891 — Cuba. Jamaica.
Aspidium Mett.; Eat. Mem. Amer. Ac. n. s. **8**. 210. 1860; Nephrodium Hk. 1862. HB. 288.
(?) **yaeyamensis** (Mak.) C. Chr. Ind. 301. 1905 — Liu Kiu.
Aspidium Mak. Bot. Mag. Tokyo **11**. 18. 1897.
(D) **zambesiaca** (Bak.) C. Chr. Ind. 301. 1905 — Africa orient.
Nephrodium Bak. Ann. Bot. **5**. 318. 1891; Aspidium Hier. 1895.
Zollingeriana O. Ktze. Rev. Gen. Pl. **2** 814. 1891 = Aspidium brachiatum.

DRYOSTACHYUM J. Smith, Journ. Bot. **3**. 399. 1841 et in Hk. Gen. Fil. t. 95. 1841 — (Polypodiaceae Gen. **104**).
caudatum J. Sm. JoB. **4**. 62. 1841 = Polypodium Feei var.
drynarioides (Hk.) Kuhn, Ann. Lugd. Bat. **4**. 296. 1869. NPfl. 328 — Malacca-Malesia-Ins. Salomonis. — Acrostichum Hk. sp. **5**. 282. 1864. HB. 425; Photinopteris Bedd. 1869; Polypodium Christ 117. 1897 (non Gris. 1857).
pilosum J. Sm. JoB. **3**. 399. 1841 = D. splendens var.
speciosum Kuhn, Ann. Lugd. Bat. **4**. 296. 1869 = Photinopteris.
splendens J. Sm. JoB. **3** 399. 1841 et in Hk. Gen. t. 95. 1841. NPfl. 328 — Malesia-Ins. Philipp. N. Guinea. — Polypodium Hk. 1863. HB. 367. Chr. 121; Drynaria Bedd. 1869 — Dryostachyum pilosum J. Sm. 1841; Kze. Farnkr. **1**. 139 t. 61. 1844; Polypodium Salom. 1883.
Thomsoni (Bak.) Diels, Nat. Pfl. **1**⁴. 328. 1899 — Ins. Admiralitatis.
Acrostichum Bak. Journ. Linn. Soc. **15**. 111. 1876; Ic. pl. t. 1694.

DYCTIOGRAMME Presl, Epim. Bot. 263. 1849 = **Coniogramme.**
japonica Pr. Epim. 263. 1849. (conf. Notogramme).

EATONIOPTERIS Bommer, Bull. Soc. France **20**. XIX. 1873 = **Cyathea.**

Ectoneura Fée, Hist. des Acrost. 75. 1845 *(Polybotrya §)* = Polybotrya.

EGENOLFIA Schott, Gen. Fil. tab. 16. 1834 = **Polybotrya.**
appendiculata J. Sm. Ferns br. and for. 111. 1866.
asplenifolia, intermedia, neglecta, nodiflora, rhizophylla, Schottii, serrulata Fée, Gen. 48. 1850-52; Gaudichaudiana, marginata, montana Fée, Gen. 358. 1850-52 = P. appendiculata varr.

bipinnatifida J. Sm. Hist. Fil. 132. 1875 } = P. appendiculata varr.
Hamiltoniana Schott, Gen. t 16. 1834 }
?nana Fée, Gen. 48. 1850-52 = ? Blechnum lanceolatum.

ELAPHOGLOSSUM Schott, Gen. Fil. ad tab. 14. 1834 — (Polypodiaceae [Gen. **107**).

accedens (Mett.) Christ, Mon. 91. 1899 — Bolivia.
Acrostichum Mett.; Kuhn, Linn. **36**. 49. 1869. HB. 523.

achroalepis (Bak.) C. Chr. Ind. 302. 1905 -- Madagascar.
Acrostichum Bak. JoB. **1880**. 371.

acrocarpum (Mart.) Moore, Ind. 4. 1857; Chr. Mon. 107 — Brasilia.
Acrostichum Mart. Ic. Cr. Bras. 85 t. 23. 1834. HB. 522; Olfersia Pr. 1836.

actinolepis (Sod.) C. Chr. Ind. 302. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 11. 1905.

actinotrichum (Mart.) Moore, Ind. 4. 1857; Chr. Mon. 68 — Brasilia.
Acrostichum Mart. Ic. Cr. Bras. 86. 1834.

acutissimum Christ, Bull. Boiss. II. **5**. 730. 1905 — Mexico.

adenolepis (Kze.) Moore, Ind. 4. 1857 — Peru.
Acrostichum Kze. Linn. **9**. 27. 1834.

aemulum (Klf.) Brack. Expl. Exp. **16**. 71. 1854 — Ins. Hawaii.
Acrostichum Klf. Enum. 63. 1824. (HB. 401. Chr. Mon. 48); Olfersia Pr. 1836. Acrostichum helleri Und. 1897.

affine (Mart. et Gal.) Moore, Ind. 4. 1857 — Amer. trop. — Acrostichum Mart. et Gal. Mém. Ac. Brux. 24 t. 3 f. 1. 1842. A. unitum Bory; Fée 1845. (HB. 401).

alatum Gaud. Voy. Bonite Bot. t. 135. 1846 = E. sessile.

alatum Moore, Ind. XVI. 1857 = E. pteropus.

albescens (Sod.) Christ, Mon. 123 f. 68. 1899. NPfl. 333 — Columbia-Peru.
Acrostichum Sod. Rec. 75. 1883; Cr. vasc. quit. 455. 1893.

alcicorne Keys. Pol. Cyath. Hb. Bung. 37. 1873 = Platycerium.

alismaefolium Moore, Ind. **5**. 1857 = E. latifolium var.

alpestre (Gardn.) Moore, Ind. 5. 1857 — Columbia. Brasilia.
Acrostichum Gardn. in Field. et Gardn. Sert. pl. t. 25. 1844. HB. 405.

amplissimum (Fée) Christ, Bull. Boiss. II. **2**. 328. 1902. — Brasilia.
Acrostichum Fée, Cr. vasc. Br. **1**. 5 t. 6. 1869; Chr. Mon. 40.

amygdalifolium (Mett.) Christ, Mon. 101 f. 53. 1899 — Amer. centr. Cuba.
Acrostichum Mett.; Kuhn, Linn. **36**. 41. 1869. HB. 520 — A. Tatei Bak. [1874. HB. 518.

andicola Moore, Ind. 5. 1857 = E. latifolium var.

Andreanum Christ, Mon. 100 f. 50. 1899 — Columbia.

angamarcanum (Sod.) C. Chr. Ind. 302. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 5. 1905.

angulatum (Bl.) Moore, Ind. 5. 1857 — Java. — Acrostichum Bl. Enum. 101. 1828. (HB. 401; Chr. Mon. 46, 49); Olfersia Pr. 1836.

angustum (Fée) Christ, Bull. Boiss. II. **2**. 361. 1902 — Brasilia. — Acrostichum Fée, Cr. vasc. Br. **2**. 12 t. 86 f. 3. 1872-73. A. Galvinii Glaz.; Bak. 1891!

antioquianum Hier. Engl. Jahrb. **34**. 543. 1904 — Columbia.

aphlebium Moore, Ind. 5. 1857 = E. conforme.

apodum (Klf.) Schott, Gen. ad t. 14. 1834. Chr. 38; Mon. 111. NPfl. 333 — Amer. trop.-Chile. — Acrostichum Klf. Enum. 59. 1824. HB. 407; Olfersia Pr. 1836. Acrostichum platynevron Fée 1845 (non L. 1753); Elaphoglossum Moore 1857 — Acrostichum Webbii Bory; Fée 1845; Elaphoglossum Moore 1857. (Chile).

araneosum (Eat.) C. Chr. Ind. 303. 1905 — Mexico. — Acrostichum Eat. Proc. Amer. Ac. **22**. 461. 1887. A. venustum Fée, 8 mém. 68. 1857 (non Liebm. 1849); Elaphoglossum Moore 1857; Chr. Mon. 95.

argyrophyllum Christ, Farnkr. 37. 1897 = E. Lindigii.

Aschersonii *Hier.* Engl. Jahrb. **34**. 555. 1904 — *Columbia.*

aspidiolepis (Bak.) C. Chr. Ind. 303. 1905 — Madagascar.
Acrostichum Bak. JoB. **1880**. 371.

attenuatum Moore, Ind. 5. 1857 = E. hybridum var. melanopus.

attenuatum Christ, Mon. 36. 1899 = E. firmum.

Aubertii (Desv.) Moore, Ind. 5. 1857; Chr. Mon. 105, 148. NPfl. 333 — Fernando Po. Africa austr. Amer. trop. — Acrostichum Desv. Berl. Mag. **5**. 309. 1811. HB. 406 — A. Boivini Mett.; Kuhn 1868 (Mauritius) — A. hirtipes Fée 1869 — A. stramineum Fée 1869 (non Mett. 1864) — A. Beaurepairii Fée 1872-73 — A. Macahense Fée 1872-73 — A. setigerum Sod. 1893; Elaphoglossum Diels 1899. NPfl. 333.

aureo-nitens Diels, Nat. Pfl. 1^4. 334. 1899 = Trachypteris pinnata.

auricomum (Kze.) Moore, Ind. 7. 1857 — Amer. trop.
Acrostichum Kze. Linn. **9**. 28. 1834. HB. 410: Olfersia Pr. 1836 (adhuc f. Christ E. laminarioides) — Acrostichum Tanajense Kuhn 1869.

auripilum Christ, Bull. Boiss. *II*. **5**. 8. 1905 — Costa Rica.

Backhousianum Moore, Gard. Chr. n. s. **17**. 672 f. 103-105. 1882 — Mexico.
Acrostichum Bak. 1891.

Bakeri (Sod.) Christ, Mon. 132. 1899. NPfl. 333 — Ecuador.
Acrostichum Sod. Rec. 77. 1883; Cr. vasc. quit. 443. 1893.

Balansae C. Chr. Ind. 303. 1905 — Paraguay.
Acrostichum tenerum Bak. JoB. **1878**. 302 (non Fée; Kuhn 1869); Elaphoglossum Christ, Mon. 62. 1899.

Bangii Christ, Mon. 99 f. 51. 1899 = E. Moorei.

Banksianum (Fée) Moore, Ind. 7. 1857 — ? — Acrostichum Fée, Acrost. 64. 1845.

barbatum Hier. Engl. Jahrb. **34**. 553. 1904 = E. scolopendrifolium.

Barteri (Bak.) C. Chr. Ind. 303. 1905 — Africa occ. trop.
Acrostichum Bak. Syn. 519. 1874.

Beccarianum (Bak.) C. Chr. Ind. 303. 1905 — Borneo.
Acrostichum Bak. Malesia **3**. 27. 1886; A. Norrisii Ces. 1876 (non Hk. 1864).

Bellermannianum (Kl.) Moore, Ind. 7. 1857; Chr. Mon. 80. NPfl. 334 — Mexico-Ecuador. — Acrostichum Kl. Linn. **20**. 426. 1847. HB. 410.

Blanchetii (Mett.) C. Chr. Ind. 303. 1905 — Brasilia.
Acrostichum Mett.; Kuhn, Linn. **36**. 55. 1869.

blepharodes Moore, Ind. 7. 1857 = E. hybridum var.

Blumeanum J. Sm. JoB. **3**. 400. 1841 — Ins. Philipp.
Acrostichum Fée Acrost. 62. 1845. — (verisim. = E. petiolatum var).

boragineum (Sod.) Christ, Mon. 122 f. 67. 1899 — Ecuador. Columbia.
Acrostichum Sod. Rec. 80. 1883; Cr. vasc. quit. 456. 1893.

borbonicum (Bak.) C. Chr. Ind. 303. 1905 — Réunion.
Acrostichum Bak. Ann. Bot. **5**. 491. 1891.

borneense (Burck) C. Chr. Ind. 303. 1905 — Borneo.
Acrostichum Burck, Ann. Jard. Buit. **4**. 99. 1884.

Boryanum (Fée) Moore, Ind. 7. 1857; Chr. Mon. 130. NPfl. 333 — India occ. Costa Rica-Quito. — Acrostichum Fée, Acrost. 40 t. 1. 1845. HB. 407.

brachyneuron (Fée) J. Sm. Cat. cult. Ferns 26. 1857 — Amer. trop.
Acrostichum Fée, Acrost. 49 t. 22 f. 1. 1845. A. Stenopteris Kl. 1847. HB. 402; Elaphoglossum Moore 1857; Chr. Mon. 61.

brevipes (Kze.) Moore, Ind. 7. 1857 — Amer. austr. trop. Puerto Rico. Guadeloupe.
Acrostichum Kze. 1845; Fée, Acrost. 29. 1845. (Chr. Mon. 45).

Burchellii (Bak.) C. Chr. Ind. 304. 1905 — Brasilia.
Acrostichum Bak. Syn. 401. 1868; Chr. Mon. 62.

caespitosum (Sod.) Christ, Mon. 100 f. 52. 1899. NPfl. 332 — Ecuador.
Acrostichum Sod. Rec. 74. 1883; Cr. vasc. quit. 427. 1893.

Calaguala Moore, Ind. 7. 1857 = E. huacsaro.

callaefolium J. Sm. Bot. Mag. **72**. Comp. 17. 1846. = E. brevipes.

callifolium (Bl.) Moore, Ind. 7. 1857; Chr. Mon. 34 — Java.
Acrostichum Bl. Enum. 100. 1828; Olfersia Pr. 1836.

callolepis (Fée) Moore, Ind. 7. 1857 — Mexico.
Acrostichum Fée, 8 mém. 69. 1857.

calophyllum (Kze.) Moore, Ind. 7. 1857 — Peru. Guiana.
Acrostichum Kze. Linn. **9**. 27. 1834.

cardiophyllum (Hk.) Moore, Ind. 7. 1857. NPfl. 333 — Ecuador.
Acrostichum Hk. Ic. pl. t. 715. 1845. HB. 404; A. cochleariæfolium Fée 1850-52; Elaphoglossum Moore 1857; Chr. Mon. 57.

carnosum Keys. Pol. Cyath. Hb. Bung. 36. 1873 = Drymoglossum.

cartilagineum (Mett.) C. Chr. Ind. 304. 1905 — Brasilia.
Acrostichum Mett.; Kuhn, Linn. **36**. 58. 1869.

castaneum (Bak.) Diels, Nat. Pfl. 1^4. 333. 1899 — Ecuador. Columbia. Costa Rica.
Acrostichum Bak. JoB. **1877**. 166.

caudatum Moore, Ind. 7. 1857 = E. petiolosum.

caulolepia (Karst.) Hier. Engl. Jahrb. **34**. 548. 1904 — Columbia.
Achrostichum Karst. Fl. Col. **1**. 121. t. 60. 1860.

chartaceum (Bak.) C. Chr. Ind. 304. 1905 — Jamaica.
Acrostichum Bak.; Jenm. JoB. **1882**. 325.

Christii (Sod.) C. Chr. Ind. 304. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 4 t. 1. 1905.

ciliatum (Pr.) Moore, Ind. 8. 1857; Chr. Mon. 93. NPfl. 333 — Amer. austr. trop.
Acrostichum Pr. Rel. Haenk. **1**. 15. 1825. HB. 406 (non Desv. 1811); Olfersia Pr. 1836. — (nomen malum; an = E. Preslianum?).

cinnamorneum (Bak.) Diels, Nat. Pfl. 1^4. 333. 1899. — Africa occ. trop.
Acrostichum Bak. Syn. 409. 1868; A. Mannianum Mett. 1868 (nomen); Elaphoglossum Christ, Mon. 148. 1899.

cladotrichium (Sod.) C. Chr. Ind. 304. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 9. 1905.

clarenceanum (Bak.) C. Chr. Ind. 304. 1905 — Fernando Po.
Acrostichum Bak. Kew Bull. **1896**. 42.

cochleariaefolium Moore, Ind. 8. 1857 = E. cardiophyllum.

cognatum Moore, Ind. 10. 1857 = E. elongatum.

conforme (Sw.) Schott, Gen. ad t. 14. 1834. Chr. 33; Mon. 43. NPfl. 332 — Trop. (Typ. St. Helena). — Acrostichum Sw. Syn. 10, 192 t. 1 f. 1. 1806 HB. 401 (pt.); Olfersia Pr. 1836.
— Acrostichum angustatum Schrad. 1818; Olfersia Pr. 1836. Vittaria acrostichoides Hk. et Grev. 1830; Drymoglossum Moore 1857. (Afr. austr.).
— Acrostichum glandulosum Carm.; Hk. et Grev. 1827 (St. Helena).

— Acrostichum didynamum Fée 1845; Elaphoglossum Moore 1857. (Réunion).
— Acrostichum Lepervanchii Fée 1845; Elaphoglossum Moore 1857 (Réunion).
— Acrostichum marginatum Wall. 1828; Fée 1845 (non Schkuhr 1809); Olfersia Pr. 1836; Elaphoglossum Moore 1857. (India bor.-Malesia).
— Acrostichum aphlebium Kze.; Kl. 1847; Elaphoglossum Moore 1857 (Merida).
— Acrostichum glaucum Fée 1857 (non Cav. 1799); Elaphoglossum Moore [1857. (Mexico).
— Acrostichum spissum Fée 1869. (Brasilia).
— Acrostichum ovatum Fée 1872-73 (non Hk. et Grev. 1829). (Brasilia).
— Acrostichum compactum Mett.; Kuhn 1869. (Columbia).
— Acrostichum Fendleri Mett.; Chr. Mon. 44. 1899. (Venezuela).
— Conf. E. gorgoneum.

confusum Christ, Mon. 124. 1899 — Ecuador. Columbia.
Acrostichum hirtipes Sod. Cr. vasc. quit. 442. 1893 (non Fée 1869).

conopodium Moore, Ind. 354. 1862 = E. conforme.

consobrinum Moore, Ind. 8. 1857 = E. latifolium var.

corazonense Christ, Mon. 86. 1899 — Ecuador.

Corderoanum (Sod.) Christ, Mon. 81. 1899 — Ecuador.
Acrostichum Sod. Cr. vasc. quit. 470. 1893.

crassinerve Moore, Ind. 8. 1857 = E. latifolium var.

crassipes (Hier.) Diels, Nat. Pfl. 1[4]. 334. 1899 — Argentina.
Acrostichum Hier. Engl. Jahrb. **22**. 409. 1896; Chr. Mon. 77.

crenatum Moore, Ind. 354. 1862 = E. Feei.

crinitum (L.) Christ, Mon. 112 f. 53. 1899. NPfl. 334 — India occ. Mexico. Costa Rica. — Acrostichum L. sp. **2**. 1068. 1753. HB. 421; Olfersia Pr. 1836; Hymenodium Fée 1845; Dictyoglossum J. Sm. 1846; Anetium Pr. 1849; Chrysodium Mett. 1856. Chr. 48.

crispatulum Moore, Ind. 8. 1857 = E. ovatum.

Cruegerianum (Kze.) C. Chr. Ind. 305. 1905 — Trinidad.
Acrostichum Kze.; Kuhn, Engl. Jahrb. **24**. 139. 1897.

cubense (Mett.) C. Chr. Ind. 305. 1905 — Cuba. Jamaica.
Acrostichum Mett.; Kuhn, Linn. **36**. 46. 1869. HB. 520; Chr. Mon. 108.

Cumingii Moore, Ind. 8. 1857 = E. decurrens var.

Curtisii (Bak.) C. Chr. Ind. 305. 1905 — Madagascar.
Acrostichum Bak. 1891; A. schizolepis Bak. JoB. **1881**. 368 (non 1876).

curvans (Kze.) Moore, Ind. 8. 1857; Chr. Mon. 90 — Peru.
Acrostichum Kze. Linn. **9**. 30. 1834.

cuspidatum (Willd.) Moore, Ind. XVI. 1857; Chr. Mon. 77. NPfl. 334 — Amer. trop.
Acrostichum Willd. sp. **5**. 106. 1810. HB. 411; Olfersia Pr. 1836.

decipiens Hier. Engl. Jahrb. **34**. 554. 1904 — Columbia.

Deckenii (Kuhn) C. Chr. Ind. 305. 1905 — Africa. orient. trop.
Acrostichum Kuhn. Fil. Deck. 16. 1867. HB. 523; Chr. Mon. 147.

decoratum (Kze.) Moore. Ind. 8. 1857. Chr. 35; Mon. 40. NPfl. 333 — Amer. trop.
Acrostichum Kze. Linn. **9**. 25. 1834. HB. 404; Olfersia Pr. 1836.

decurrens (Desv.) Moore, Ind. 8. 1857 — Java. Ins. Philipp.
Acrostichum Desv. Berl. Mag. **5**. 310. 1811. HB. 403. (Chr. Mon. 51); Olfersia Pr. 1836 — Acrostichum obtusifolium Spr. 1827; Bl. 1828 (an Willd. 1810?); Elaphoglossum J. Sm. 1841; Olfersia Blumeana Pr. 1836 — Acrostichum Cumingii Fée 1845. HB. 407; Chr. Mon. 52; Elaphoglossum Moore 1857.

deltoideum (Sod.) Christ, Mon. 56 f. 19. 1899. NPfl. 334 — Columbia-Ecuador.
Acrostichum Sod. Cr. vasc. quit. 466. 1893.

dendricolum (Bak.) Christ, Mon. 94. 1899. NPfl. 333 — Columbia-Ecuador.
Acrostichum Bak. Syn. 521. 1874.

didymoglossoides C. Chr. Bot. Tidsskr. **26**. 299. 1904 — Brasilia.

didynamum Moore, Ind. 8. 1857 = E. conforme.

dimorphum (Hk. et Grev.) Moore, Ind. 8. 1857. Chr. 35; Mon. 98. NPfl. 333 — St. Helena. — Acrostichum Hk. et Grev. Ic. Fil. t. 145. 1829. HB. 406; Olfersia Pr. 1836; Microstaphyla J. Sm. 1875. Anogramme? paradoxa part. Fée 1850-52; Microstaphyla furcata part. Fée 1857.

discolor (Kuhn) C. Chr. Ind. 306. 1905 — Amazonas.
Acrostichum Kuhn, Linn. **36**. 53. 1869. HB. 521; Elaphoglossum sancti-gabrielis Christ, Mon. 37. 1899.

dissimile (Kze.) Moore, Ind. 8. 1857 — Mexico-Peru. Brasilia.
Acrostichum Kze. Linn. **9**. 28. 1834; Olfersia Pr. 1836 — Acrostichum Schiedei Kze. 1837; Elaphoglossum Moore 1857; Chr. Mon. 94 — Acrostichum Schmitzii Mett. 1869. HB. 523.

Dombeyanum Moore et Houlst. Gard. Mag. Bot. **3**. 95. 1851 = E. perelegans.

drabifolium Christ, Mon. 111 f. 59. 1899 — Cuba.

durum Moore, Ind. 9. 1857 = E. simplex var. martinicense.

Eatonianum (E. Britt.) C. Chr. Ind. 306. 1905 — Bolivia.
Acrostichum E. Britt. Bull. Torr. Cl. **15**. 252. 1888.

ecuadorense C. Chr. Ind. 306. 1905 — Ecuador.
Acrostichum fimbriatum Sod. Rec. 84. 1883; Cr. vasc. quit. 474. 1893 (vix Cav. 1799); Elaphoglossum Christ, Mon. 85. 1899. NPfl. 334 (non Moore [1862).

Eggersii (Bak.) Christ, Mon. 71. 1899 — San Domingo.
Acrostichum Bak. JoB. **1888**. 34.

elegans Hier. Engl. Jahrb. **34**. 549. 1904 = ? E. plumosum.

ellipticum (Fée) Moore, Ind. 9. 1857 — Mauritius.
Acrostichum Fée, Acrost. 30 t. 4. 1845; Chr. Mon. 34. — (an E. Sieberi var?).

elongatum (Kze.) Moore, Ind. 9. 1857 — Mexico-Peru. Brasilia.
Acrostichum Kze. Linn. **9**. 31. 1834; Chr. Mon. 68 — A. intermedium Fée 1857; Elaphoglossum cognatum Moore 1857.

Engelii (Karst.) Christ, Mon. 81. 1899 — Columbia-Bolivia.
Acrostichum Karst. Fl. Col. **1**. 118. t. 59. 1860.

erinaceum Moore, Ind. 9. 1857 = E. scolopendrifolium.

erythrolepis (Fée) Moore, Ind. 9. 1857 — Peru.
Acrostichum Fée, Acrost. 60. 1845.

erythrotrichum Kl. msc. — Caracas. — Acrostichum Salom. Nom. 5. 1883 [(nomen).

eximium (Mett.) Christ, Mon. 107. 1899 — Columbia.
Acrostichum Mett. Ann. sc. nat. V. **2**. 199. 1864.

falcatum Moore, Ind. 9. 1857 = E. obductum.

Feei (Bory) Moore. Ind. XVI. 1857. Chr. 34; Mon. 99. NPfl. 333 — India occ.-Ecuador. — Acrostichum Bory; Fée. Acrost. 48. t. 8 f. 2. 1845. HB. 404.

feejeense Brack. Expl. Exp. **16**. 72. 1854 — Fiji. Samoa.
Acrostichum Hk. 1864. HB. 401, 519.

ferrugineum (Linden) Moore, Ind. 9. 1857; Chr. Mon. 34 — Columbia-Ecuador.
Acrostichum Linden 1856; Bak. Syn. 522. 1874. — (an E. sporadolepis var.?)

fimbriatum Moore, Ind. 356. 1862 = E. hybridum.

fimbriatum Christ, Mon. 85. 1899 = E. ecuadorense.

firmum (Mett.) Urban, Symb. Ant. **4**. 59. 1903 — Cuba. Puerto Rico. Acrostichum Mett.: Kuhn, Linn. **36**. 55. 1869 — A. attenuatum Kze.; Kuhn 1869 (non Fée 1850-52); Elaphoglossum Christ, Mon. 36. 1899.

flabellatum (H. B. Willd.) Christ, Mon. 58. 1899 — Costa Rica-Peru. Acrostichum H. B. Willd. sp. **5**. 110. 1810. HB. 415; Olfersia Pr. 1836; Rhipidopteris Fée 1845. NPfl. 331; Polybotrya Mett. 1864; Rhipidopteris Humboldtiana Schott 1834 — Acrostichum tripartitum Hk. et Grev. 1829; Rhipidopteris Schott 1834; Olfersia Pr. 1836; Polybotrya J. Sm. 1841.

flaccidum (Fée) Moore. Ind. 356. 1862. Chr. 34; Mon. 60. NPfl. 332 — Amer. trop. Acrostichum Fée, Acrost. 35 t. 7 f. 2. 1845. HB. 401 — A. lonchophyllum Fée 1857. HB. 519; Elaphoglossum Moore 1857 — Acrostichum Schaffneri Fée; Fourn. 1872 — A. papyraceum Fée 1869 — A. ambiguum Mett.; Chr. Mon. 60. 1899.

foeniculaceum (Hk. et Grev.) C. Chr. Ind. 307. 1905 — Ecuador. Acrostichum Hk. et Grev. Ic. Fil. t. 119. 1829. HB. 416; Chr. Mon. 59; Rhipidopteris Schott 1834. NPfl. 331; Olfersia Pr. 1836; Polybotrya J. Sm. 1841.

Fonki Moore, Ind. 356. 1862 = E. Mathewsii.

Forsythii-Majoris Christ, Bull. Boiss. II. **3**. 32. 1903 — Madagascar.

Fraseri (Mett.) Christ, Mon. 113. 1899 — Columbia-Ecuador. Acrostichum Mett; Kuhn, Linn. **36**. 43. 1869. HB. 520.

frigidum J. Sm. Cat. Kew Ferns 4. 1856 = E. perelegans.

Funckii (Fée) Moore. Ind. 9. 1857 — Columbia-Venezuela. Acrostichum Fée, Acrost. 36 t. 6 f. 1. 1845.

funiculum Christ, Mon. 129. 1899 — Merida.

furcatum (L.fil.) Christ. Mon. 98. f. 50. 1899 — St. Helena. Adiantum L.fil. Suppl. 447. 1781; Microstaphyla Fée 1857. Chr. 46. NPfl. 265; Polybotrya Mett.; Kuhn 1868; Osmunda bifurcata Jacq. 1789; Acrostichum Sw. 1801. HB. 413; Gymnogramma Klf. 1827; Olfersia Pr. 1836; Polybotrya J. Sm. 1841; Microstaphyla Pr. 1849; Darea furcans Bory 1828; Anogramme ? paradoxa part. Fée 1850-52.

furcatum Keys. Pol. Cyath. Hb. Bung. 37. 1873 = Eschatogramme.

furfuraceum (Mett.) Christ, Mon. 126. 1899 — Costa Rica. Acrostichum Mett.; Kuhn, Linn. **36**. 51. 1869. HB. 523.

furfuraceum Diels, Nat. Pfl. 1^4. 334. 1899 = E. vulcanicum.

fuscopunctatum Christ, Bull. Boiss. **6**. 867. 1898; Mon. 51 — Yunnan.

Gardnerianum (Kze.) Moore, Ind. XVI. 1857; Chr. Mon. 121 — Columbia-Ecuador. Brasilia. — Acrostichum Kze.; Fée. Acrost. 55 t. 15 f. 3. 1845. HB. 405.

Gayanum (Fée) Moore, Ind. 10. 1857; Chr. Mon. 46 — Amer. austr. subtrop. Acrostichum Fée, Acrost. 37 t. 19 f. 2. 1845. (HB. 401).

Gilleanum Diels, Nat.Pfl. 1^4. 334. 1899 = Trachypteris pinnata.

glabellum J. Sm. Lond. JoB. **1**. 197. 1842 — Amer. trop. Acrostichum Kl. 1847. A. martinicense Fée 1845 (non Desv. 1811).

glabratum (Mett.) Moore, Ind. 357. 1862 — N. Caledonia. Acrostichum Mett. Ann. sc. nat. IV. **15**. 55. 1861. HB. 518; Chr. Mon. 47.

glaucum Moore, Ind. 10. 1857 = E. conforme.

glossoides (André) Christ, Mon. 97 t. 4. 1899 — Columbia. Acrostichum André msc.

glossophyllum Hier. Hedwigia **44**. 180. 1905 — Columbia. E. linguiforme Hier. Engl. Jahrb. **34**. 542. 1904 (non Moore 1857).

glutinosum (Spruce) Christ, Mon. 120. 1899 — Columbia-Ecuador.
Acrostichum Spruce hb. n. 5614.

gorgoneum (Klf.) Brack. Expl. Exp. **16**. 74. 1854. NPfl. 332 — Malesia-Polynesia. Acrostichum Klf. Enum. 63. 1824. HB. 416; Chr. Mon. 33; Olfersia Pr. 1836; Aconiopteris J. Sm. 1875. — (f. Christ = forma E. latifolii). — ? Aconiopteris obtusa Fée 1845; Olfersia Moore 1857. Acrostichum pellucido-marginatum Christ 1895. — (f. Christ = formae E. conformis).

gracile Christ in Schwacke, Pl. Nov. Mineiras **2**. 10. 1900 = E. lineare.

gracilipes (Fée) C. Chr. Ind. 308. 1905 — Ecuador.
Acrostichum Fée; Kuhn, Linn. **36**. 47. 1869.

gramineum (Jenm.) Urban, Symb. Ant. **4**. 60. 1903 — Jamaica. Puerto Rico.
Acrostichum Jenm. JoB. **1879**. 263.

grandifolium (Mett.) C. Chr. Ind. 308. 1905 — Venezuela.
Acrostichum Mett.; Kuhn, Linn. **36**. 62. 1869.

gratum (Fée) Moore, Ind. 10. 1857; Chr. Mon. 118 — Mexico.
Acrostichum Fée, 8 mém. 69. 1857; 10 mém. 5 t. 30 f. 1. HB. 521.

guatemalense (Kl.) Moore. Parkers Cat. 1858: Ind. 357. 1862 — Guatemala.
Acrostichum Kl. Allgem. Gartenzeit. **1855**. 66.

Hartwegii Moore, Ind. XVI. 1857 = E. Mathewsii.

heliconiifolium (Sod.) Christ, Farnkr. 35. 1897; Mon. 134. NPfl. 334 — Ecuador.
Acrostichum Sod. Rec. 78. 1883; Cr. vasc. quit. 464. 1893.

Herminieri (Bory et Fée) Moore, Ind. XVI. 1857; Chr. Mon. 62. NPfl. 332 — Amer. trop. — Acrostichum Bory et Fée; Fée, Acrost. 43 t. 11. 1845. HB. 402.

heterolepis Moore, Ind. 10. 1857 — E. tomentosum.

heteromorphum (Kl.) Moore, Ind. 10. 1857; Chr. Mon. 117. NPfl. 333 — Columbia-Ecuador. — Acrostichum Kl. Linn. **20**. 424. 1847. HB. 405.

Hieronymi (Sod.) C. Chr. Ind. 308. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 11. 1905.

hirtum (Sw.) C. Chr. Ind. 308. 1905 — Amer. trop. Ins. Azor. Madeira. Ins. Afr. orient. India austr. Ceylon. Ins. Hawaii. — Acrostichum Sw. Schrad. Journ. **1800**[2]. 10. 1801 — A. squamosum Sw. 1801. HB. 411 (non Cav. 1799); Elaphoglossum J. Sm. 1841. Chr. 36; Mon. 63. NPfl. 334. A. vestitum (paleaceum in tab.) Hk. et Grev. 1831; Elaphoglossum J. Sm. 1866 (non Schott 1834); Olfersia paleacea Pr. 1836 — Acrostichum micans Mett. 1869. (Ins. Hawaii.) — A. splendens Willd. 1810; Olfersia Pr. 1836; Elaphoglossum Brack. 1854.

Hoffmanni (Mett.) Christ, Mon. 36. 1899 — Costa Rica. Peru.
Acrostichum Mett.; Kuhn **36**. 56. 1869 — A. polystichum Mett. 1869.

horridulum J. Sm. Bot. Voy. Herald 232. 1854 = E. spathulatum var.

huacsaro (Ruiz) Christ, Mon. 96. 1899 (Huacssaro) — Columbia-Peru. ? Jamaica. Acrostichum Ruiz 1799; Bertol. Opusc. sci. Bol. **1**. 241 t. 8. 1817 (Cuacsaro HB. 405; Elaphoglossum Ruizianum Moore 1857 — Acrostichum Calaguala Kl. 1847; Elaphoglossum Moore 1857.

hyalinum Christ, Farnkr. 37. 1897; Mon. 70 — Costa Rica.

hybridum (Bory) Moore. Ind. 10. 1857. Chr. 38; Mon. 103, 148. NPfl. 333 — Africa austr. cum. ins. Amer. trop. — Acrostichum Bory, Voy. **3**. 95. 1804. HB. 403; Olfersia Pr. 1836 — Acrostichum melanopus Kze. 1850. HB. 403; Elaphoglossum Moore 1857 — Acrostichum crinitum Mart. et Gal. 1842 (non L. 1753); A. blepharodes Fée 1845; Elaphoglossum Moore 1857 — Acrostichum nitidum Liebm. 1849; Elaphoglossum fimbriatum Moore

1857; Acrostichum mexicanum Fourn. 1872. (an sp. propr.?) — A. Lindbergii Mett.; Kuhn 1869.

hybridum Brack. Expl. Exp. **16**. 69. 1854 = E. scolopendrifolium.

hystrix (Kze.) Moore, Ind. 10. 1857 — Peru. Mexico.
Acrostichum Kze. Linn. **9**. 26. 1834. — (an E. scolopendrifolium var. ?).

impressum Moore, Ind. 10. 1857 = E. latifolium.

inaequalifolium (Jenm.) C. Chr. Ind. 309. 1905 — Jamaica.
Acrostichum Jenm. JoB. **1886**. 273.

intermedium Brack. Expl. Exp. **16**. 69. 1854 = E. Gardnerianum.

inversum (Cord.) Christ, Mon. 146. 1899 — Réunion. — Acrostichum Cord. Bull. Soc. sci. et arts Réunion **1890-91**. 84. 1891; Fl. Réun. 95. 1895.

isophyllum (Sod.) Christ, Mon. 131. 1899. NPfl. 333 — Ecuador.
Acrostichum Sod. Cr. vasc. quit. 441. 1893.

Jamesoni (Hk. et Grev.) Moore, Ind. 10. 1857 — Mexico-Bolivia. — Acrostichum Hk. et Grev. Ic. Fil. t. 86. 1829; Chr. Mon. 113. A. spathulatum var. auctt.

Junghuhnianum Moore, Ind. 10. 1857 = E. callifolium.

Karstenianum Moore, Ind. 10. 1857 = A. petiolatum.

laminarioides (Bory) Moore, Ind. 10. 1857; Chr. Mon. 72 — Amer. trop.
Acrostichum Bory; Fée, Acrost. 57 t. 12. 1845. HB. 523. (f. Christ = E. auricomum) — A. Orbignyanum Fée 1845; Elaphoglossum Moore 1857.

lanceolatum Keys. Pol. Cyath. Hb. Bung. 37. 1873 = Paltonium.

Langsdorffii Moore, Ind. 11. 1857 = E. muscosum.

Lastii (Bak.) C. Chr. Ind. 309. 1905 — Zanzibar.
Acrostichum Bak. Ann. Bot. **5**. 491. 1891.

latifolium (Sw.) J. Sm. Lond. JoB. **1**. 197. 1842. Chr. 34; Mon. 31. NPfl. 332 — Amer. trop. — Acrostichum Sw. Prod. 128. 1788. HB. 403 — A. scalpellum Mart. 1834; Elaphoglossum Moore 1857 — Acrostichum andicola Fée 1845; Elaphoglossum Moore 1857 — Acrostichum alismaefolium Fée 1845; Elaphoglossum Moore 1857 — Acrostichum crassinerve Kze. 1845; Elaphoglossum Moore 1857 — Acrostichum hymenodiastrum Fée 1869 — A. Tovarense Moritz; Eat. 1860; Kuhn 1869; Elaphoglossum Moore 1862 — Acrostichum membranaceum Fourn. & Mazé msc.; Elaphoglossum Hier. 1904. — Conf. E. gorgoneum et E. pachyphyllum.

latifolium Bedd. Handb. 416. 1883 = E. laurifolium.

laurifolium (Thouars) Moore, Ind. XVI. 1857; Chr. Mon. 36 — Tristan d'Acunha. Ins. Mascarenae. India austr. Ceylon. Java. Celebes. — Acrostichum Thouars, Fl. Trist. d'Ac. 31. 1804. (HB. 401 non 412); Olfersia Pr 1836 — Acrostichum conforme Bl. 1828 (non Sw. 1806); A. commutatum Mett.; Kuhn 1869.

Lechlerianum (Mett.) Moore, Ind. 11. 1857; Chr. Mon. 61 (Lechleri) — Columbia-Peru. — Acrostichum Mett. Fil. Lechl. **1**. 3. 1856. (HB. 401).

Lehmannianum Christ. Mon. 81 f. 36. 1899 — Columbia.

Lepervanchii Moore, Ind. 11. 1857 = E. conforme.

lepidoglossum (Bak.) C. Chr. Ind. 309. 1905 — Columbia.
Acrostichum Bak. Ann. Bot. **5**. 492. 1891.

lepidotum J. Sm. Cat. Kew Ferns 1856; Cat. cult. Ferns 26. 1857 = E. plicatum.

leprosum (Mett.) Christ, Mon. 119. 1899 — Peru.
Acrostichum Mett.; Kuhn, Linn. **36**. 48. 1869.

leptophlebium (Bak.) C. Chr. Ind. 309. 1905 — Guiana.
Acrostichum Bak. Tr. Linn. Soc. II. Bot. **2**. 295. 1886.

leptophyllum (Fée) Moore, Ind. 11. 1857. NPfl. 332 — Amer. trop.
Acrostichum Fée, Acrost. 45 t. 17 f. 1. 1845. HB. 400. (Chr. Mon. 48).

Lessonii (Mett.) C. Chr. Ind. 310. 1905 — Vanikoro. Sumatra.
Acrostichum Mett.; Kuhn. Linn. **36**. 60. 1869. (Chr. Mon. 49).

Liebmanni Moore, Ind. 360. 1862 = E. Lindenii.

Lindenii (Bory) Moore, Ind. XVI. 1857; Chr. Mon. 110 — Amer. trop.
Acrostichum Bory; Fée, Acrost. 48 t. 18 f. 3. 1845. HB. 404 — A. venustum Liebm. 1849; Elaphoglossum Liebmanni Moore 1862.

Lindigii (Karst.) Moore, Ind. 360. 1862; Chr. Mon. 84 — Columbia-Ecuador.
Acrostichum Karst. Fl. Col. **1**. 5 t. 3. 1858 — A. argyrophyllum Sod. 1883; Elaphoglossum Christ 1897. 37.

lineare (Fée) Moore, Ind. 11. 1857 — Amer. trop. ? Réunion.
Acrostichum Fée, Acrost. 47 t. 15. f. 2. 1845. HB. 406 — A. gracile Fée 1872-73; Elaphoglossum Christ 1900.

lingua (Raddi) Brack. Expl. Exp. **16**. 74. 1854; Chr. Mon. 45. NPfl. 332 — Amer. trop. — Acrostichum Raddi, Opusc. sci. Bol. **3**. 283. 1819: Pl. Bras. **1**. 5 t. 15 f. 4. 1825. HB. 402 (non Thbg. 1784) — A. scandens Bory; Fée 1845 (non Raddi 1819); Elaphoglossum Moore 1857.

linguiforme (Cav.) Moore, Ind. 11. 1857 — Ecuador-Peru.
Acrostichum Cav. Anal. Hist. Nat. **1**. 103. 1799; Descr. 238. 1802.

linguiforme Hier. Engl. Jahrb. **34**. 542. 1904 = E. glossophyllum.

litanum (Sod.) C. Chr. Ind. 310. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 10. 1905.

lloense (Hk.) Moore, Ind. 11. 1857; Chr. Mon. 54. NPfl. 333 — Mexico-Peru.
Acrostichum Hk. Ic. pl. t. 657. 1844 — A. proximum Bomm. 1896; Elaphoglossum Christ 1905.

longifolium (Jacq.) J. Sm. Bot. Mag. **72** Comp. 17. 1846 — Amer. trop.
Acrostichum Jacq. Coll. **2**. 105. 1788; Olfersia Pr. 1836; Aconiopteris Fée 1845; Chr. Mon. 33.

longipes Brack. Expl. Exp. **16**. 70 t. 9 f. 2. 1854 — Peru.

longissimum (Sod.) C. Chr. Ind. 310. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 3. 1905.

Lorentzii (Hier.) Christ, Mon. 121. 1899 — Argentina.
Acrostichum Hier. Engl. Jahrb. **22**. 408. 1896.

luridum Christ, Mon. 33. 1899 = E. Schomburgkii.

macrophyllum (Kl.) Christ, Bull. Boiss. II. **5**. 10. 1905 — Brasilia. Costa Rica.
Olfersia Kl. hb.; Acrostichum Mett.; Kuhn, Linn. **36**. 61. 1869.

macropodium Moore, Ind. 11. 1857 = E. Sieberi.

macrorhizum (Bak.) C. Chr. Ind. 310. 1905 — Brasilia.
Acrostichum Bak. Ann. Bot. **5**. 490. 1891.

madagascariense (Kuhn) C. Chr. Ind. 310. 1905 — Madagascar.
Acrostichum Kuhn; Bak. 1891 (nomen); Chr. Mon. 145. 1899.

malgassicum C. Chr. Ind. 310. 1905 — Madagascar.
Acrostichum subsessile Bak. Kew Bull. **1897**. 300 (non 1884).

magnum (Bak.) C. Chr. Ind. 310. 1905 — Demerara.
Acrostichum Bak. Gard. Chr. n. s. **20**. 135. 1883.

Mandoni (Mett.) Christ, Mon. 128. 1899 — Bolivia.
Acrostichum Mett.; Kuhn, Linn. **36**. 47. 1869. HB. 522.

Mannianum Christ, Mon. 148. 1899 = E. cinnamomeum.

marginale (Bak.) Christ, Mon. 131. 1899 — Ecuador.
Acrostichum Bak. Syn. 519. 1874.

marginatum Moore, Ind. 11. 1857 = E. conforme.

martinicense Moore, Ind. 11. 1857 = E. simplex var.

Mathewsii (Fée) Moore, Ind. 12. 1857; Chr. Mon. 87. NPfl 334 — Mexico-Chile.
Acrostichum Fée, Acrost. 54 t. 2 f. 2. 1845. HB. 408 — A. Hartwegii Fée 1845; Elaphoglossum Moore 1857 — Acrostichum assurgens Bak. 1874. HB. ed. II. 408 — A. Fonki Phil. 1857; Elaphoglossum Moore 1862 (Chile).

maximum (Bak.) C. Chr. Ind. 311. 1905 — Columbia.
Acrostichum Bak. Ann. Bot. **5**. 490. 1891.

melanolepis (Fée) Moore, Ind. 12. 1857 —?
Acrostichum Fée, Gen. 43. 1850-52 (nomen).

melanopus Moore, Ind. 12. 1857 = E. hybridum var.

melanostictum (Bl.) Moore, Ind. 361. 1862; Chr. Mon. 63 — Penang. Java.
Acrostichum Bl. Fl. Javae Fil. 26 t. 7. 1828. A. Norrisii Hk. 1864. HB. 401; Elaphoglossum Bedd. 1870. NPfl. 332.

membranaceum Hier. Engl. Jahrb. **34**. 540. 1904 = E. latifolium var.

meridense (Kl.) Moore, Ind. 12. 1857 — Amer. austr. trop.
Acrostichum Kl. Linn. **20**. 427. 1847. (HB. 410); Chr. Mon. 73.

Mettenii (Kuhn) Christ, Mon. 125. 1899 — Ecuador-Columbia.
Acrostichum Kuhn, Linn. **36**. 48. 1869. HB. 520.

micradenium (Fée) Moore, Ind. 12. 1857; Chr. Mon. 53. NPfl. 332 — Ins. Hawaii.
Acrostichum Fée, Acrost. 43 t. 8 f. 1. 1845. HB. 400. Elaphoglossum nitidum Brack. 1854.

microlepis (Kze.) J. Sm. Cat. Kew Ferns 1856; Cat. cult. Ferns 26. 1857 — Venezuela.
Acrostichum Kze. Linn. **23**. 214, 299. 1850.

microlepis Diels, Nat. Pfl. 1^4. 332. 1899 = E. subnudum.

Miersii (Bak.) C. Chr. Ind. 311. 1905 — Brasilia.
Acrostichum Bak. Ann. Bot. **5**. 490. 1891.

miniatum Christ, Mon. 72 f. 30. 1899 — Celebes.
Acrostichum Christ, Verh. Nat. Ges. Basel **11**. 254 t. 3 f. 25-27. 1895.

minutum (Pohl) Moore, Ind. 12. 1857 — Brasilia. Costa Rica.
Acrostichum Pohl; Fée, Acrost. 39 t. 10 f. 3. 1845. (HB. 401); Chr. Mon. 47.

Moorei (E. Britt.) Christ, Bull. Boiss. II. **3**. 148. 1903 — Bolivia. Columbia.
Acrostichum E. Britt. Mem. Torr. Club. **4**. 271. 1895; Microstaphyla Und. 1905; Rhipidopteris Rusbyi Christ 1897. 46. NPfl. 331; Elaphoglossum Bangii Christ, Mon. 99. 1899; Bull. Boiss. II. **1**. 588. 1901; Microstaphyla [Hier. 1904.

Moritzianum Moore, Ind. 12. 1857 = E. villosum var.

Mülleri (Fourn.) C. Chr. Ind. 311. 1905 — Mexico.
Acrostichum Fourn. Mex. pl. **1**. 68 t. 1 f. 2. 1872. HB. 522.

muscosum (Sw.) Moore, Ind. 12. 1857. Chr. 36; Mon. 74. NPfl. 334 — Amer. trop.
Acrostichum Sw. Prod. 128. 1788. HB. 410; Olfersia Pr. 1836 — Acrostichum Langsdorffii Hk. et Grev. 1829; Olfersia Pr. 1836; Elaphoglossum Moore 1857 — Acrostichum Dombeyanum Fée 1845.

Neintneri Moore, Ind. 362. 1862 (nomen) — Ceylon.

nervosum (Bory) Christ, Mon. 50 f. 13. 1899. NPfl. 332 — St. Helena.
Acrostichum Bory, Dup. Voy. Bot. **1**. 252. 1828; Olfersia Moore 1857. Aconiopteris J. Sm. 1866; Acrostichum subdiaphanum Hk. et Grev. 1831. HB. 416; Aconiopteris Pr. 1836; Olfersia Moore 1857.

nigrescens (Hk.) Diels, Nat. Pfl. 1^4. 332. 1899 — Guiana.
Acrostichum Hk. sp. **5**. 214. 1864. HB. 400.
nitidum Brack. Expl. Exp. **16**. 70 t. 9 f. 3. 1854 = E. micradenium.
Norrisii Bedd. Ferns br. Ind. ad finem 1870 = E. melanostictum.
notatum (Fée) Moore, Ind. 12. 1857; Chr. Mon. 119 — Bolivia.
Acrostichum Fée, Acrost. 38 t. 10 f. 1. 1845.
obductum (Klf.) Moore, Ind. 12. 1857; Chr. Mon. 88 — Ins. Mascarenae.
Acrostichum Klf.; Spr. Syst. **4**. 34. 1827. HB. 410; Olfersia Pr. 1836 — Acrostichum falcatum Fée 1845; Elaphoglossum Moore 1857.
oblanceolatum C. Chr. Ind. 312. 1905 — Ecuador.
Acrostichum subsessile Sod. Sert. Fl. Ecuad. 6. 1905 (non Bak. 1884).
obliquatum Christ, Mon. 56. 1899 = E. Sellowianum.
obscurum (Fourn.) C. Chr. Ind. 312. 1905 — Mexico.
Acrostichum Fourn. Mex. pl. **1**. 68. 1872.
obtusifolium J. Sm. JoB. **3**. 401. 1841 = E. decurrens.
oleandropsis (Sod.) C. Chr. Ind. 312. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 7. 1905. (Aconiopteris).
opacum Hier. Engl. Jahrb. **34**. 541. 1904 — Columbia.
Orbignyanum Moore, Ind. XVI. 1857 = E. laminarioides.
ornatum (Mett.) Christ, Mon. 38. 1899 — Amer. austr. trop.
Acrostichum Mett.; Kuhn, Linn. **36**. 44. 1869. HB. 522.
ovalifolium (Fée) Christ, Bull. Boiss. II. **2**. 361. 1902 — Brasilia.
Acrostichum Fée, Cr. vasc. Br. **1**. 3 t. 4 f. 1. 1869.
ovatum (Hk. et Grev.) Moore, Ind. XVI. 1857. Chr. 37; Mon. 127. NPfl. 334 Amer. trop. (excl. Ind. occ.) — Acrostichum Hk. et Grev. Ic. Fil. t. 146. 1829. HB. 408 — A. rabdolepis Fée 1850-52; Elaphoglossum Moore 1857 — Acrostichum ramosissimum Fée 1845; Elaphoglossum Moore 1857 — Acrostichum crispatulum Fée 1850-52; Elaphoglossum Moore 1857.
oxyphyllum Moore, Ind. 362. 1862 = E. flaccidum.
pachycraspedon Christ, Hedwigia **44**. 362. 1905 — Peru.
pachydermum (Fée) Moore, Ind. 12. 1857 — Brasilia.
Acrostichum Fée, Acrost. 47. 1845.
pachyphyllum (Kze.) C. Chr. Ind. 312. 1905 — Costa Rica-Peru.
Acrostichum Kze. Linn. **9**. 26. 1834. HB. 421; Chr. Mon. 33. Olfersia Pr. 1836; Anetium Pr. 1849; Hymenodium Moore 1857; H. Kunzeanum Fée 1845 — Acrostichum corpulentum Mett.; Kuhn 1869. — (f. Christ = E. latifolium var.).
pala André; Christ, Mon. 35 t. 1. 1899 — Ecuador.
pallidum (Bak.) C. Chr. Ind. 312. 1905 — Jamaica.
Acrostichum Bak. JoB. **1879**. 263.
palustre (Hk.) J. Sm. Ferns br. and for. ed. II. 298. 1877; Chr. Mon. 147. 1899 — Africa occ. trop. — Acrostichum Hk. sp. **5**. 214. 1864. HB. 402; Lomariopsis Mett.; Kuhn 1868.
pangoanum (Sod.) C. Chr. Ind. 312. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 6. 1905.
papillosum (Bak.) Christ, Mon. 130. 1899. NPfl. 333 — Ecuador.
Acrostichum Bak. JoB. **1877**. 167.
Patini (Bak.) Christ, Mon. 58. 1899 — Columbia.
Acrostichum Bak. Syn. 519. 1874.

pellucidum Gaud. Voy. Bonite Bot. t. 50 f. 5-7. 1846 (sine descript.) = E. [micradenium.
pellucidum (Sod.) C. Chr. Ind. 313. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 7. 1905.

peltatum (Sw.) Urban, Symb. Ant. **4**. 60. 1903 — Amer. trop.
Osmunda Sw. Prod. 127. 1788; Acrostichum Sw. 1801. HB. 416; Chr. Mon. 59; Rhipidopteris Schott 1834. Chr. 45. NPfl. 331; Olfersia Pr. 1836; Peltapteris Link 1841; Polybotrya Link 1841.

perelegans (Fée) Moore, Ind. XVI. 1857. Chr. 36; Mon. 73 — Amer. trop.
Acrostichum Fée, Acrost. 55 t. 23. 1845. HB. 411.

petiolatum (Sw.) Urban, Symb. Ant. **4**. 61. 1903 — Amer., Africa, Asia trop.
Acrostichum Sw. Prod. 128. 1788; A. viscosum Sw. 1806. HB. 406; Elaphoglossum Schott 1834. Chr. 39; Mon. 91, 148. NPfl. 333; Olfersia Pr. 1836 — Acrostichum salicifolium Willd.; Klf. 1824; Olfersia Pr. 1836 — Acrostichum neriifolium Wall. 1828; Olfersia Pr. 1836 — Acrostichum stelligerum Wall. 1829. HB. 521; Elaphoglossum Moore 1857 — Acrostichum Karstenianum Kze. 1850; Elaphoglossum Moore 1857 — Acrostichum angustissimum Fée 1865 — A. tenuiculum Fée 1865.

petiolosum (Desv.) Moore, Ind. 12. 1857. Chr. 34; Mon. 115. NPfl. 332 — Ecuador-Peru. — Acrostichum Desv. Berl. Mag. **5**. 309. 1811. HB. 400; Olfersia Pr. 1836; Acrostichum acuminatum Juss.; Poir. 1810 (non Willd. 1810) — A. caudatum Hk. 1840; Elaphoglossum Moore 1857.

pichinchae Christ, Mon. 68. 1899 — Columbia-Ecuador.

piloselloides Moore, Ind. 13. 1857 = E. spathulatum var.

piloselloides Keys. Pol. Cyath. Hb. Bung. 36. 1873 = Drymoglossum heterophyllum.

pilosum (H.B.Willd.) Moore, Ind. 13. 1857. Chr. 120. NPfl. 334 — Mexico-Columbia. — Acrostichum H.B.Willd. sp. **5**. 103. 1810. HB. 406; Olfersia [Pr. 1836.

platyneuron Moore, Ind. 13. 1857 = E. apodum.

plicatum (Cav.) C. Chr. Ind. 313. 1905 — Mexico-Peru.
Acrostichum Cav. Anal. Hist. Nat. **1**. 102. 1799; Descr. 238. 1802; A. lepidotum Willd. sp. **5**. 102. 1810. HB. 409; Olfersia Pr. 1836; Elaphoglossum J. Sm. 1856; Chr. Mon. 77. NPfl. 334.

Plumieri Moore, Ind. 13. 1857 = E. villosum var.

plumosum (Fée) Moore, Ind. 364. 1862; Chr. Mon. 65 — Amer. trop.
Acrostichum Fée, Acrost. 54 t. 20 f. 1. 1845 — A. Wageneri Kze. 1853; Elaphoglossum Moore 1857 — Acrostichum acuminans Fée 1869 — A. chrysolepis Fée 1869 — A. elegans Fée 1872-73.

Poeppigianum Moore, Ind. 13. 1857 = E. villosum.

polytrichum Christ, Mon. 113. 1899 — Ecuador.

Poolii (Bak.) Christ, Bull. Boiss. II. **3**. 32. 1903 — Madagascar.
Acrostichum Bak. Ann. Bot. **5**. 493. 1891.

potosianum Christ, Mon. 119. 1899 — Mexico.

praecelsum (André) Christ, Mon. 121 t. 3. 1899 — Columbia.
Acrostichum André msc.

Preslianum (Fée) Christ, Mon. 95. 1899 — Venezuela-Peru. Brasilia.
Acrostichum Fée, Acrost. 46 (t. 24 f. 1?). 1845. (Conf. E. ciliatum) — A. praelongum Fée 1869.

Prestoni J. Sm. Ferns br. and for. ed. II. 298. 1877 = A. scolopendrifolium var.

Pringlei (Dav.) C. Chr. Ind. 313. 1905 — Mexico.
Acrostichum Dav. Bot. Gaz. **21**. 253. 1894.

procurrens (Mett.) Moore, Ind. 365. 1862; Chr. Mon. 110 — Cuba. Montserrat.
Acrostichum Mett.; Eat. Mem. Amer. Ac. n. s. 8. 194. 1860. HB. 408.
propinquum (Mett.) Christ, Mon. 115. 1899 — Peru.
Acrostichum Mett.; Kuhn, Linn. **36**. 45. 1869.
proximum Christ, Bull. Boiss. II. **5**. 9. 1905 = E. lloense var.
pseudodidynamum Hier. Engl. Jahrb. **34**. 544. 1904 — Columbia.
pteropodum (Sod.) C. Chr. Ind. 314. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 8. 1905.
pteropus C. Chr. Ind. 314. 1905 — Amer. trop.
Acrostichum alatum Fée, Acrost. 35 t. 5 f. 2. 1845. (HB. 401): Chr. Mon. 46 (non Roxb. 1844); Elaphoglossum Moore 1857 (non Gaud. 1846).
pusillum (Mett.) C. Chr. Ind. 314. 1905 — Cuba.
Acrostichum Mett.; Kuhn, Linn. **36**. 43. 1869.
pygmaeum (Mett.) Christ, Mon. 111 f. 6. 1899. NPfl. 333 — Ecuador-Columbia.
Acrostichum Mett; Kuhn, Linn. **36**. 42. 1869. HB. 520.
quitense (Bak.) C. Chr. Ind. 314. 1905 — Ecuador.
Acrostichum Bak. Ann. Bot. **5**. 493. 1891.
rabdolepis Moore, Ind. 13. 1857 = E. ovatum.
Raddianum Brack. Expl. Exp. **16**. 67. 1854 = E. spathulatum var.
ramosissimum Moore, Ind. XVI. 1857 = E. ovatum.
rampans (Bak.) Christ, Mon. 56. 1899 — Cuba. Amer. centr.
Acrostichum Bak. Syn. 518. 1874.
Reichenbachii (Moritz) Christ, Mon. 106. 1899 — Venezuela.
Acrostichum Moritz; Kuhn, Linn. **36**. 42. 1869.
repians (Cav.) Moore, Ind. 365. 1862 — Ecuador.
Acrostichum Cav. Anal. Hist. Nat. **1**. 104. 1799; Descr. 239. 1802.
reticulatum (Klf.) Gaud. Voy. Bonite, Bot. t. 79 f. 1-4. 1846; Chr. Mon. 50. NPfl. 332 — Ins. Hawaii. — Acrostichum Klf. Enum. 64. 1824. HB. 421; Anetium Pr. 1849; Hymenodium Moore 1857; Acrostichum crassifolium Gaud. 1827; Hymenodium Fée 1845.
revolutum (Liebm.) Moore, Ind. 365. 1862 — Mexico. Cuba. Columbia.
Acrostichum Liebm. Vid. Selsk. Skr. V. **1**. 163 (seors. 11). 1849 — A. humile Mett. 1864; Chr. Mon. 47.
revolvens (Fée) C. Chr. Ind. 314. 1905 — Columbia.
Acrostichum Fée; Mett. 1864 (nomen); Kuhn, Linn. **36**. 57. 1869.
rhynchophyllum Christ, Mon. 57 f. 20. 1899 — Venezuela. Columbia.
Richardi (Bory) Christ, Mon. 35. 1899 — Réunion. — Aconiopteris Bory; Fée, Acrost. 80 t. 16 f. 4. 1845; Olfersia Moore 1857; Acrostichum Hk. 1864.
Rimbachii (Sod.) Christ, Mon. 126 f. 70. 1899 — Ecuador.
Acrostichum Sod. Cr. vasc. quit. 478. 1893.
Roezlii (Schaffn.) Moore, Ind. 14. 1857 (Roeslii) — Mexico.
Acrostichum Schaffn.; Fée, 8 mém. 69. 1857; 10 mém. 7 t. 30 f. 2.
rubescens (Kuhn) Christ, Mon. 93 f. 47-48. 1899 — Guatemala.
Acrostichum Kuhn msc.
rubiginosum J. Sm. Cat. cult. Ferns 26. 1857 = E. perelegans.
rufescens (Liebm.) Moore, Ind. 365. 1862; Chr. Mon. 119 — Mexico.
Acrostichum Liebm. Vid. Selsk. Skr. V. **1**. 171 (seors. 18). 1849.
rufidulum (Willd.) C. Chr. Ind. 314. 1905 — Madagascar.
Acrostichum Willd.; Kuhn, Fil. Afr. 47. 1868. HB. 522.
Ruizianum Moore, Ind. 10. 1857 = E. huacsaro.

rupestre (Karst.) Christ, Farnkr. 37. 1897; Mon. 83 f. 40. NPfl. 334 — Costa Rica-Ecuador. — Acrostichum Karst. Fl. Col. **1**. 118 t. 59. 1860.

Salvini (Bak.) C. Chr. Ind. 315. 1905 — Guatemala.
Acrostichum Bak.: Hemsl. Biol. Centr. Amer. **3**. 688. 1885.

samoense Brack. Expl. Exp. **16**. 68 t 9 f. 1. 1854. NPfl. 333 — Samoa.
Acrostichum Bak. 1868. HB. 407; Chr. Mon. 109.

sancti-gabrielis Christ, Mon. 37. 1899 = E. discolor.

savaiense (Bak.) Diels, Nat. Pfl. 1^4. 332. 1899 — Samoa.
Acrostichum Bak. Ann. Bot. **5**. 494. 1891.

scalpellum Moore, Ind. 14. 1857 = E. latifolium.

scandens Moore, Ind. 14. 1857 = E. lingua.

scapellum (Kze.) Moore, Ind. 365. 1862 — Mexico.
Acrostichum Kze.; Fée, 9 mém. 1. 1857. (= scalpellum?).

Schiedei Moore, Ind. 14. 1857 = E. dissimile.

schizolepis (Bak.) Christ, Bull. Boiss. II. **3**. 32. 1903 — Madagascar.
Acrostichum Bak. Journ. Linn. Soc. **15**. 421. 1876.

Schlimense (Fée) Moore, Ind. 366. 1862; Chr. Mon. 35 — Columbia-Ecuador.
Acrostichum Fée, 8 mém. 68. 1857. HB. 402.

Schomburgkii (Fée) Moore, Ind. 14. 1857 — Trinidad. Guiana. Costa Rica.
Acrostichum Fée. Acrost. 32 t. 8 f. 2. 1845 — A. luridum Fée, 1845; Elaphoglossum Christ, Mon. 37. 1899.

scolopendrifolium (Raddi) J. Sm. Bot. Mag. **72** Comp. 17. 1846. Chr. 38; Mon. 37. NPfl. 333 — Amer. trop. — Acrostichum Raddi, Pl. Bras. **1**. 4 t. 16. 1825. HB. 407; Olfersia Pr. 1836 — Acrostichum erinaceum Fée 1845; Elaphoglossum Moore 1857 — Acrostichum barbatum Karst. 1869; Elaphoglossum Hier. 1904 — Acrostichum Glaziovii Fée 1869 — A. insigne Fée 1872-73 — A. Prestoni Bak. 1872. HB. 519; Elaphoglossum J. Sm. 1877 — Acrostichum versatile Sod. 1883. — (an Acrostichum fimbriatum Cav. 1799?).

Sellowianum (Pr.) Moore, Ind. 366. 1862 (nomen) — Amer. austr. trop.
Olfersia Pr. 1836 (nomen); Acrostichum Kl.: Kuhn, Linn. **36**. 52. 1869; Chr. Mon. 37 — A. obliquatum Fée 1869; Elaphoglossum Christ, Mon. 56. 1899.

sessile (Fée) Moore, Ind. 14. 1857 — Ins. Hawaii. — Acrostichum Fée 1850-52; Elaphoglossum alatum Gaud. Voy. Bonite Bot. t. 135. 1845 (non Moore 1857).

setigerum Diels, Nat. Pfl. 1^4. 333. 1899 = E. Aubertii.

setosum Moore, Ind. 14. 1857 = E. villosum.

Sherringii (Bak.) C. Chr. Ind. 315. 1905 — Grenada.
Acrostichum Bak. Ann. Bot. **6**. 100. 1892; Chr. Mon. 61.

Sieberi (Hk. et Grev.) Moore, Ind. 14. 1857; Chr. Mon. 35. 145 — Ins. afric. orient. Fernando Po. — Acrostichum Hk. et Grev. Ic. Fil. t. 237. 1831. HB. 403; Olfersia Pr. 1836 — Acrostichum macropodum Fée 1845; Elaphoglossum Moore 1857.

siliquoides (Jenm.) C. Chr. Ind. 315. 1905 — Jamaica.
Acrostichum Jenm. JoB. **1881**. 53.

simplex (Sw.) Schott, Gen. ad t. 14. 1834; Chr. Mon. 47, 146. NPfl. 332 — Amer. trop. Africa trop. Madagascar. — Acrostichum Sw. Prod. 128. 1788. HB. 400; Olfersia Pr. 1836 — Acrostichum martinicense Desv. 1811 (non Fée 1845); Elaphoglossum Moore 1857 — Acrostichum lineatum Kuhn; Christ. Mon. 146. 1899 (non Cav. 1799). (Madagascar).

Smithii (Bak.) Christ. Mon. 106. 1899 — St. Vincent.
Acrostichum Bak. Ann. Bot. **5**. 169. 1891.

Sodiroi (Bak.) Christ, Mon. 122. 1899. NPfl. 333 — Ecuador. Columbia.
Acrostichum Bak. JoB. **1877**. 167; Ic. pl. t. 1688.

spathulatum (Bory) Moore, Ind. 14. 1857. Chr. 39; Mon. 112. NPfl. 333 — Africa austr. cum insulis. Ceylon. Amer. trop.
Acrostichum Bory, Voy. **1**. 363 t. 20 f. 1. 1804. HB. 408; Olfersia Pr. 1836 — Acrostichum ciliare Thouars 1804; A. obtusatum Carm. 1818 (Tristan d'Acunha). — A. piloselloides Pr. 1825; Olfersia Pr. 1836; Elaphoglossum Moore 1857 — Acrostichum horridulum Klf. 1824; Olfersia Pr. 1836; Elaphoglossum J. Sm. 1854. Acrostichum spathulinum Raddi 1825; A. Raddii Desv. 1827; A. Raddianum Hk. et Grev. 1827; Elaphoglossum Brack. 1854. (an sp. propr.?) — (Conf. E. Jamesoni).
— Acrostichum Hayesii Mett. 1869 — A. acutum Mett. 1869. (Ceylon).

splendens Brack. Expl. Exp. **16**. 68. 1854 = E. hirtum.

sporadolepis (Kze.) Moore, Ind. 367. 1862 — Venezuela. Ecuador.
Acrostichum Kze.; Eat. 1860; Kuhn, Linn. **36**. 59. 1869. — (Conf. E. [ferrugineum).

Sprucei (Bak.) Diels, Nat. Pfl. 1^4. 334. 1899 — Ecuador.
Acrostichum Bak. Syn. 411. 1868.

squamatum Moore, Ind. 15. 1857 = Acrostichum squamosum Cav.

squamipes (Hk.) Moore, Ind. 15. 1857; Chr. Mon. 53. NPfl. 333 — Amer. trop.
Acrostichum Hk. Ic. pl. t. 197. 1837. HB. 404 — A. craspedariaeforme Fée 1869 — ? A. heterolepis Bak. 1874. HB. 521 (non Fée 1845) — A. ovatum Liebm. 1849 (non Hk. et Grev. 1829); Elaphoglossum subaequale [Moore 1862.

squamosum J. Sm. JoB. **4**. 148. 1841 = E. hirtum.

squarrosum (Kl.) Moore, Ind. 15. 1857; Chr. Mon. 116 — Mexico-Ecuador.
Acrostichum Kl. Linn. **20**. 424. 1847. HB. 405.

stelligerum Moore, Ind. 15. 1857 = E. petiolatum.

stenophyllum (Sod.) Diels, Nat. Pfl. 1^4. 333. 1899 — Ecuador. Columbia.
Acrostichum Sod. Cr. vasc. quit. 468. 1893; Chr. Mon. 90.

stenopteris Moore, Ind. 15. 1857 = E. brachyneuron.

stigmatolepis (Fée) Moore, Ind. XVI. 1857; Chr. Mon. 52 — India austr.
Acrostichum Fée, Acrost. 62 t. 24 f. 2. 1845. HB. 521.

stipitatum (Bory) Moore. Ind. 15. 1857; Chr. Mon. 88 — Réunion.
Acrostichum Bory; Fée, Acrost. 38 t. 4 f. 3. 1845.

stramineum (Mett.) Christ, Mon. 117 f. 66. 1899 — Columbia.
Acrostichum Mett. Ann. sc. nat. V. **2**. 200. 1864. HB. 403.

strictum (Raddi) Moore, Ind. 15. 1857. Chr. 37; Mon. 70 — Brasilia. Costa Rica-Columbia. — Acrostichum Raddi, Pl. Bras. **1**. 3 t. 15. f. 3. 1825. HB. 409; Olfersia Pr. 1836 — Acrostichum Liaisianum Glaz.; Fée 1872-73.

subaequale Moore, Ind. 368. 1862 = E. squamipes.

subnudum C. Chr. Ind. 316. 1905 — Ecuador.
Acrostichum microlepis Sod. Cr. vasc. quit. 437. 1893; Chr. Mon. 55 (non Kze. 1850); Elaphoglossum Diels 1899. NPfl. 332 (non J. Sm. 1856).

subsessile (Bak.) C. Chr. Ind. 316. 1905 — Madagascar.
Acrostichum Bak. JoB. **1884**. 143 (non 1897).

succisifolium (Thouars) Moore. Ind. 15. 1857. Chr. Mon. 52 — Tristan d'Acunha. Ins. Amsterdam. Ins. Mascarenae.
Acrostichum Thouars, Fl. Trist. d'Ac. 31. 1804 (an succisum?); Juss.; Poir. Enc. Suppl. **1**. 120. 1810. HB. 408; Olfersia Pr. 1836.

tabanense Christ, Mon. 125 t. 2. 1899 — Columbia.
tahitense Brack. Expl. Exp. **16**. 73. 1854; Chr. Mon. 52 — Tahiti. Fiji.
Acrostichum Carr. 1873. HB. 522.
tambillense (Hk.) Moore. Ind. 15. 1857; Chr. Mon. 54. NPfl. 332 — Guatemala. Columbia-Ecuador. ? Mexico. — Acrostichum Hk. Ic. pl. t. 636. 1844. HB. 400 — ? A. Bourgaeanum Fourn. 1872.
tectum (H.B.Willd.) Moore, Ind. 15. 1857; Chr. Mon. 89. NPfl. 334 — Amer. trop. — Acrostichum H.B.Willd. sp. **5**. 102. 1810. HB. 410; Olfersia Pr. 1836 — Acrostichum rubiginosum Fée 1845.
tenerum (Fée) Hier. Engl. Jahrb. **34**. 557. 1904 (descr.) — Ecuador. Columbia.
Acrostichum Fée; Kuhn, Linn. **36**. 52. 1869.
tenerum Christ, Mon. 62. 1899 = E. Balansae.
tenuiculum (Fée) Moore ap. Bak. Ann. Bot. **5**. 491. 1891 (syn.) — Venezuela-Bolivia. — Acrostichum Fée, 10 mém. 6 t. 29 f. 2. 1865.
tenuifolium (Liebm.) Moore, Ind. 368. 1852; Chr. Mon. 56 — Mexico. Jamaica. Guiana. — Acrostichum Liebm. Vid. Selsk. Skr. V. **1**. 163 (seors. 11). [1849.
tolimense (Kuhn) Christ, Mon. 124 f. 69. 1899 — Ecuador.
Acrostichum Kuhn msc.
tomentosum (Bory) Christ, Farnkr. 37. 1897; Mon. 79 — Réunion.
Acrostichum Bory; Willd. sp. **5**. 101. 1810. HB. 411; Olfersia Pr. 1836
Acrostichum heterolepis Fée 1845; Elaphoglossum Moore 1857 — Olfersia glabrescens Pr. 1836; Aconiopteris Pr. 1849; Acrostichum Kuhn 1868.
Tonduzii Christ, Prim. Fl. costaric. **3**. 8. 1901 — Costa Rica.
Acrostichum Christ 1899 (Mon. 45 sub E. conforme).
tosaense (Yatabe) Makino, Phan. et Pter. Jap. Ic. **1**. t. 53-54. 1901 — Japonia.
Acrostichum Yatabe, Bot. Mag. Tokyo **5**. 149 t. 24. 1891.
tovarense Moore, Ind. 369. 1862 = E. latifolium var.
tragiaefolium Moore, Ind. 15, 369. 1857-62 = E. Feei.
Trianae Christ, Mon. 91. 1899 — Columbia. Ecuador.
tricholepis (Bak.) C. Chr. Ind. 317. 1905 — Madagascar.
Acrostichum Bak. JoB. **1891**. 5.
trichophorum (Sod.) C. Chr. Ind. 317. 1905 — Ecuador.
Acrostichum Sod. Sert. Fl. Ecuad. 10. 1905.
trivittatum (Sod.) Christ, Mon. 85 f. 38-39. 1899. NPfl. 334 — Ecuador.
Acrostichum Sod. Rec. 82. 1883; Cr. vasc. quit. 479. 1893.
truncicola (Karst.) Hier. Engl. Jahrb. **34**. 548. 1904 — Columbia.
Acrostichum Karst. Fl. Col. **1**. 121 t. 60. 1860.
Ulei Christ, Mon. 114 f. 62. 1899 — Brasilia austr.
undulatum (Willd.) Moore, Ind. 16. 1857; Chr. Mon. 129 — India occ. Mexico-Ecuador. — Acrostichum Willd. sp. **5**. 105. 1810; Olfersia Pr. 1836: Acrostichum podotrichum Desv. 1811.
vagans (Mett.) Hier. Engl. Jahrb. **34**. 543. 1904 — Brasilia.
Acrostichum Mett.; Kuhn, Linn. **36**. 58. 1869. (HB. 401).
venustum Moore, Ind. 16. 1857 = E. araneosum.
vestitum (Schlecht. et Cham.) Schott, Gen. ad t. 14. 1834 — Mexico.
Acrostichum Schlecht. et Cham. Linn. **5**. 605. 1830; Olfersia Pr. 1836 — Acrostichum fulvum Mart. et Gal. 1842.
vestitum J. Sm. Ferns br. and for. 106. 1866 = E. hirtum.
Vieillardii (Mett.) Moore, Ind. 369. 1862 — N. Caledonia. Fiji.
Acrostichum Mett. Ann. sc. nat. IV. **15**. 55. 1861; Olfersia Carr. 1873.

villosum (Sw.) J. Sm. JoB. **4**. 148. 1841. Chr. 38; Mon. 108. NPfl. 333 — Amer. trop. — Acrostichum Sw. Prod. 128. 1788. HB. 409; Olfersia Pr. 1836 — Acrostichum Poeppigianum Fée (1845) 1850-52; Elaphoglossum Moore 1857 — Acrostichum Plumieri Fée 1845; Elaphoglossum Moore 1857 — Acrostichum Moritzianum Kl. 1847; Elaphoglossum Moore 1857 — Acrostichum setosum Liebm. 1849; Elaphoglossum Moore 1857 — Acrostichum recognitum Kze.; Fourn. 1872. HB. 520 — A. Haynaldii Sod. 1883 — A. mollissimum Fée 1869 — A. acutum Fée 1872-73 (non Kuhn 1869).

viride (Fourn.) C. Chr. Ind. 318. 1905 — Mexico.
Acrostichum Fourn. Mex. pl. **1**. 66. 1872.

viridifolium (Jenm.) C. Chr. Ind. 318. 1905 — Jamaica.
Acrostichum Jenm. JoB. **1886**. 273.

viscidum (Fée) Christ in Schwacke, Pl. Nov. Mineiras **2**. 17. 1900 — Brasilia.
Acrostichum Fée, Cr. vasc. Br. **2**. 12 t. 86 f. 1. 1872-73; Chr. Mon. 70.

viscosum Schott, Gen. ad t. 14. 1834 = E. petiolatum.

Volkensii (Hier.) C. Chr. Ind. 318. 1905 — Africa orient. trop.
Acrostichum Hier. in Engl. Pflw. Ostafr. C. 81. 1895.

vulcanicum Christ, Mon. 131. 1899 — Ecuador.
Acrostichum furfuraceum Bak. JoB. **1877**. 166 (non Mett. 1869); Elaphoglossum Diels 1899. NPfl. 334.

Wageneri Moore, Ind. 16. 1857 = E. plumosum.

Wawrae (Luerss.) C. Chr. Ind. 318. 1905 — Ins. Hawaii.
Acrostichum Luerss. Flora **1875**. 420.

Webbii Moore, Ind. 16. 1857 = E. apodum.

Welwitschii (Bak.) C. Chr. Ind. 318. 1905 — Africa occ. trop.
Acrostichum Bak. Syn. 521. 1874.

Wercklei Christ, Bull. Boiss. II. **5**. 9. 1905 — Costa Rica.

Wrightii (Mett.) Moore, Ind. 370. 1862; Chr. Mon. 51 — Cuba. Costa Rica.
Acrostichum Mett.; Eat. Mem. Amer. Ac. n. s. **8**. 194. 1860. HB. 401.

xanthoneuron (Kze.) Moore, Ind. 370. 1862; Chr. Mon. 89 — Venezuela.
Acrostichum Kze.; Eat. Mem. Amer. Ac. n. s. **8**. 194. 1860 (nomen).

yarumalense Hier. Engl. Jahrb. **34**. 556. 1904 — Columbia.

Yatesii (Sod.) Christ, Mon. 83. f. 37. 1899 — Ecuador.
Acrostichum Sod. Anal. Univ. quit. **4**. 174. 1890; Cr. vasc. quit. 469. 1893.

Yoshinagae (Yatabe) Mak. Phan. et Pter. Jap. Ic. **1**. t. 51-52. 1901 — Japonia.
Acrostichum Yatabe, Bot. Mag. Tokyo **5**. 109 t. 23. 1891.

Eleutheria Kunze, Bot. Zeit. **1844**. 296 (*Hemitelia §*) = Hemitelia.

Eleutherophlebia Gris. Abh. Gött. Ges. Wiss. **7**. 282, 285. 1857 (*Polypodium* et *Aspidium §*) = Polypodium et Dryopteris.

ELLOBOCARPUS Kaulfuss, Enum. Fil. 117. 1824 = **Ceratopteris**.
cornutus Klf. Enum. 148. 1824. } = C. thalictroides.
oleraceus Klf. Enum. 148. 1824. }

ENTEROSORA Baker, Trans. Linn. Soc. II. Bot. **2**. 294. 1886 — (Polypodiaceae. Gen. **101**).

Campbellii Bak. Tr. Linn. Soc. II. Bot. **2**. 294. t. 55. 1886. NPfl. 322 — Guiana. Jamaica. Costa Rica. — E. Fawcettii Jenm. 1895.

Fawzettii Jenm. Gard. Chr. III. **18**. 62. 1895 = E. Campbellii.

EREMOPODIUM Trevisan, Atti dell' Ist. Veneto V. **3**. 589. 1877 = **Asplenium**.
sundense Trevis. Atti. Ist. Veneto V. **3**. 589. 1877 = ? A. vittaeforme.
vittaeforme Trevis. Atti. Ist. Veneto V. **3**. 589. 1877.

Eriochosma J. Smith. Journ. Bot. 4. 50. 1841 *(Nothochlaena §)* = Notholaena.
ERIOSORUS Fée, Gen. Fil. 152. 1850-52 = **Gymnogramma.**
?Ruizianus Fée. Gen. 152 t. 13 f. 2. 1850-52 = Gymnopteris ferruginea.
scandens Fée, Gen. 152. 1850-52 = Gymnogramma aureo-nitens.
ESCHATOGRAMME Trevisan, Atti dell' Ist. Veneto II. **2**. 168. 1851 — Polypodiaceae Gen. **95**).
furcata (L.) C. Chr. Bot. Tidsskr. **26**. 285. 1904 — Amer. trop.
Pteris L. sp. **2**. 1073. 1753; Taenitis Willd. 1810. HB. 397; Pteropsis Desv. 1827; Cuspidaria Fée 1851-52. Chr. 132; Dicranoglossum J. Sm. 1854. NPfl. 303; Taeniopsis Moore 1857; Vittaria Bak. 1870; Elaphoglossum Keys. 1873; Oetosis O. Ktze. 1891 — ? Pteris tricuspidata L. 1753; Taenitis Spr. 1827; Pteropsis Desv. 1827; Taeniopsis Moore 1857; Cuspidaria tricuspis Fée 1851-52 — Taenitis Desvauxii Kl. 1847 (Hk. et Grev. t. 7). Cuspidaria subpinnatifida Fée 1851-52; Dicranoglossum Moore 1861.
Euphorophyllum van den Bosch, Plant. Jungh. **1**. (19). 1856. *(Hymenophyllum §)* = Hymenophyllum.
EUPODIUM J. Smith in Hook. Gen. Fil. t. 118. 1842 = **Marattia.**
Kaulfussii J. Sm. in Hk. Gen. t. 118. 1842.
EUPTERIS Newman, The Phytologist **2**. 278. 1845 = **Pteridium.**
aquilina Newm. Phytol. **2**. 278. 1845.
Euryostichum Presl. Epim. Bot. 188. 1849 *(Anapausia §)* = Leptochilus.
FADYENIA Hooker, Gen. Fil. t. 53. 1840 — (Polypodiaceae Gen. **11**).
Fadyenii (Mett.) C. Chr. Ind. 319. 1905 — India occ.
Aspidium Mett. 1856. Chr. 234; Asplenium proliferum Sw. Prod. 129. 1788 (non Lam. 1786); Aspidium Hk. et Grev. 1829; Fadyenia Hk. 1840. HB. 303. NPfl. 181.
prolifera Hk. Gen. t. 53. 1840 = F. Fadyenii.
FEEA Bory, Dict. class. d'Hist. Nat. **6**. 446. 1824 = **Trichomanes.**
Boryi v.d.B. Ned. Kr. Arch. **4**. 347. 1859 = T. diversifrons.
botryoides v.d.B. Ned. Kr. Arch. **4**. 346. 1859.
Humboldtii v.d.B. Ned. Kr. Arch. **4**. 347. 1859 = T. heterophyllum.
nana Bory, Dict. class. **6**. 446. 1824; **17** t. 42 f. 1 = T. botryoides.
polypodina Bory. Dict. class. **6**. 446. 1824; **17** t. 43 = T. osmundoides.
spicata Pr. Abh. böhm. Ges. V. **5**. 330. 1848 = T. osmundoides.
Spruceana v.d.B. Ned. Kr. Arch. **4**. 347. 1859 = T. heterophyllum var.
FILIX Ludwig. Instit. Hist. Phys. Regni Veg. ed. II. 142. 1757 = ? **Cystopteris.**
bipinnata Gilib. Exerc. Phyt. **2**. 557. 1792 = Athyrium filix femina.
bulbifera Und. Our nat. ferns ed. VI. 119. 1900.
fragilis Und. Our nat. ferns ed. VI. 119. 1900.
montana Und. Our nat. ferns ed. VI. 119. 1900.
FOURNIERA Bommer, Bull. Soc. France **20**. XIX. 1873.
funebris Fourn. L'Ill. Hort. **23**. 99. 1876 = Cyathea.
lepidotricha Fourn. Ann. sc. nat. V. **18**. 349. 1873 = Alsophila.
Novae Caledoniae Bomm.; Fourn. Ann. sc. nat. V. **18**. 349. 1873 = Alsophila.
FURCARIA Desvaux. Prodrome 292. 1827 = **Ceratopteris.**
cornuta Desv. Prod. 292. 1827 } = C. thalictroides.
thalictroides Desv. Prod. 292. 1827 } = C. thalictroides.
GALEOGLOSSA Presl. Epim. Bot. 132. 1849 = **Cyclophorus.**
nummularifolia Pr. Epim. 133. 1849 } = C. nummularifolius.
obovata Pr. Epim. 133. 1849 } = C. nummularifolius.
rotundifolia Pr. Epim. 133. 1849 } = C. nummularifolius.

GISOPTERIS Bernhardi, Schrad. Journ. **1800**[2]. 129. 1801 = **Lygodium.**
palmata Bernh. Schrad. Journ. **1800**[2]. 129 t. 2 f. 1. 1801.
GLAPHYROPTERIS Presl, Abhandl. böhm. Ges. Wiss. V. **5**. 344. 1848 = **Dryo-**
decussata Pr. Abh. böhm. Ges. V. **5**. 344. 1848. [**pteris.**
erubescens Fée, Cr. vasc. Br. **2**. 41. 1872-73.
rudis Pr. Abh. böhm. Ges. V. **5**. 344. 1848.
GLEICHENIA Necker, Element. bot. **3**. 314. 1790 = **Dryopteris.**
GLEICHENIA Smith. Mém. Acad. Turin **5**. 419. 1793 — (Gleicheniaceae Gen. **2**).
acutifolia Hk. sp. **1**. 7 t. 8 A. 1844 = G. quadripartita.
affinis Mett.; Kuhn, Linn. **36**. 167. 1869 — Peru.
alpina R. Br. Prod. Fl. N. Holl. 161. 1810 — Tasmania. N. Zealand. Samoa.
Mertensia Poir. 1813; Platyzoma Desv. 1827; Calymella Pr. 1836; Gleichenia dicarpa β HB. 12 — G. hecistophylla A. Cunn. 1836.
arachnoidea A. Cunn.; Col. Tasm. Jour. **2**. 162. 1844 = G. Cunninghamii.
arachnoides Mett. Ann. Lugd. Bat. **1**. 47. 1863 = G. glauca var.
argentea Klf. Enum. 36. 1824 = G. polypodioides var.
attenuata Wall. List n. 156. 1828 = G. flagellaris.
axialis Christ, Bull. Boiss. II. **5**. 14. 1905 — Costa Rica.
Bancroftii Hk. sp. **1**. 5 t. 4 A. 1844 — Amer. trop. — Mertensia Kze. 1844: Gleichenia glauca vel G. longissima auctt. quoad pl. americ.
Bibreae Mett.; Kuhn, Linn. **36**. 166. 1869. HB. 449 — Chile.
bifida (Willd.) Spr. Syst. **4**. 27. 1827. Chr. 341 — Amer. trop.
Mertensia Willd. Vet. Ak. Nya Hdl. **1804**. 168 t. 5 f. B. M. immersa Klf. 1824; Gleichenia Spr. 1827. G. pubescens HB. 14 (pt.). NPfl. 353 (pt.).
bifurcata Bl. Enum. 250. 1828 = G. laevigata.
blepharolepis Sod. Rec. 5. 1883 — Ecuador.
G. pubescens var. Sod. Cr. vasc. quit. 34. 1893.
Boryi Kze. Farnkr. **1**. 162 t. 70 f. 1. 1844. HB. 12. Chr. 339. NPfl. 355 — Réunion. — Gleicheniastrum Pr. 1848.
Brackenridgei Fourn. Ann. sc. nat. V. **18**. 269. 1873 — Fiji. N. Caledonia.
Mertensia flagellaris Brack. 1854 (non Bory 1810).
brasiliana Spr. Syst. **4**. 27. 1827 = G. pectinata.
Brunei Christ, Bull. Boiss. II. **5**. 13. 1905 — Costa Rica.
bullata Moore, Ind. 374. 1862 = G. glauca.
ciliata Col. Tr. N. Zeal. Inst. **29**. 414. 1897 — N. Zealand.
circinnata Sw. Schrad. Journ. **1800**[2]. 107. 1801. HB. 11. Chr. 338. NPfl. 355 — Australia. Tasmania. N. Zealand. N. Caledonia. Borneo. Malacca.
Mertensia Poir. 1813. Gleichenia microphylla R. Br. 1810; Mertensia Klf. 1824; Calymella Pr. 1836; Gleicheniastrum Pr. 1848 — Gleichenia speluncae R. Br. 1810; Mertensia Poir. 1813; Platyzoma Desv. 1827; Gleicheniastrum Pr. 1848 — Gleichenia semivestita Lab. 1824; Gleicheniastrum Pr. 1848.
Obs. f. Mett. Ann. Lugd. Bat. **1**. 47 est G. circinnata Sw. = G. dicarpa R.Br. G. circinnata supra verisimiliter G. microphylla appellanda est et G. dicarpa infra G. circinnata.
circinnata Mett. Ann. Lugd. Bat. **1**. 47. 1863 = G. dicarpa.
compacta Christ, Bull. Boiss. II. **5**. 254. 1905 — Costa Rica.
cryptocarpa Hk. sp. **1**. 7 t. 6 A. 1844. HB. 13. NPfl. 353 — Chile austr. Chiloë. Ins. Falkland. — Mertensia Gay 1853.

Cunninghamii Hew.; Hk. sp. **1**. 6 t. 6 B. 1844. HB. 13. Chr. 340. NPfl. 353 — N. Zealand. — Mertensia J. Sm. 1843. (nomen).

dicarpa R Br. Prod. Fl. N. Holl. 161. 1810. HB. 12 (excl. var.). Chr. 338. NPfl. 355 — Austr. Tasmania. N. Caledonia. — Mertensia Poir. 1813; Platyzoma Desv. 1827; Calymella Pr. 1848. — Conf. *Obs.* sub G. circinnata.

dichotoma Hk. sp. **1**. 12. 1844 = G. linearis.

elongata Bak. Kew Bull. **1901**. 137 — Uganda.

emarginata Moore, Ind. 377. 1862 = G. linearis var.

excelsa J. Sm. JoB. **3**. 420. 1841; Hk. sp. **1**. 5 t. 4 B. 1844 = G. glauca.

farinosa Hk. sp. **1**. 9. 1844 = G. bifida vel G. furcata.

ferruginea Bl. Enum. 249. 1828 = G. linearis var.

flabellata R.Br. Prod. Fl. N. Holl. 161. 1810. HB. 12. Chr. 340. NPfl. 353 — Austr. Tasmania. N. Caledonia. N. Zealand. — Mertensia Poir. 1813.

flagellaris (Bory) Spr. Syst. **4**. 25. 1827. HB. 14. Chr. 347. NPfl. 353 — Ins. Mascar. Ins. Sechellae. — Mertensia Bory; Willd. sp. **5**. 74. 1810.

flagellaris auctt. quoad pl. males. = G. laevigata.

flexuosa (Schrad.) Mett. Ann. Lugd. Bat. **1**. 50. 1863 — Amer. trop. Mertensia Schrad. Gött. gel. Anz. **1824**. 863; Mart. Ic. Cr. Bras. 108 t. 60 f. 1. M. rigida Kze. 1834; Gleichenia Bomm. et Christ 1896.

furcata (L.) Spr. Syst. **4**. 26. 1827 — Amer. trop. Acrostichum L. Syst. Nat. ed. X. **2**. 1321. 1759; Polypodium Sw. 1801; Mertensia Willd. 1804 — M. tomentosa Cav.; Sw. 1804; Gleichenia Spr. 1827 — Mertensia pubescens H.B.Willd. 1810; Gleichenia HBK. 1815. HB. 13 (pt.). NPfl. 353 (pt.) — ? Pteris dichotoma L. 1753.

Gaenslyi Rosenst. Festschr. Alb. v. Bamberg 56. 1905 — Brasilia.

gigantea Wall. List n. 157. 1828; Hk. sp. **1**. 5 t. 3 A. 1844 = G. glauca.

glabra Mann, Proc. Amer. Ac. **7**. 211. 1868 = G. glauca.

glauca (Thbg.) Hk. sp. **1**. 4 t. 3 B. Chr. 340. NPfl. 353 — Japonia. China. Asia, Austr., Polynesia trop. — Polypodium Thbg.; Houtt. Nat. Hist. **14**. 177. 1783; Thbg. Fl. Jap. 338. 1784; Mertensia Sw. 1804; Gleichenia japonica Spr. 1827 — G. longissima Bl. 1828. HB. 12; Mertensia Kze. 1846; Mesosorus Hassk. 1856 — Gleichenia gigantea Wall. 1828; Mertensia Pr. 1836; Mesosorus Hassk. 1856. Hicriopteris speciosa Pr. 1849 — Gleichenia excelsa J. Sm. 1841; Mertensia J. Sm. 1843; Mesosorus Hassk. 1856 — Mertensia pinnata Kze. 1837; Gleichenia Moore 1862. Mertensia glabra Brack. 1854; Gleichenia Mann 1868 (Hawaii) — Mertensia arachnoides Hassk. 1855; Mesosorus Hassk. 1856; Gleichenia Mett. 1863. G. bullata Moore 1862 (an sp. propr.? Java).

glauca Sw. Syn. 165, 393. 1806 = G. polypodioides var.

glauca auctt. quoad pl. americ. = G. Bancroftii.

glaucescens HBK. Nov. Gen. et Sp. **1**. 29. 1815 = G. pectinata.

gnidioides Mett. Ann. sc. nat. V. **2**. 266. 1864 — Columbia.

gracilis (Mart.) Moore, Ind. 378. 1862 — Brasilia. Mertensia Mart. Ic. Cr. Bras. 107 t. 59 f. 2. 1834.

grandis Moore, Ind. 378. 1862 = M. Bancroftii.

Hallieri Christ, Ann. Jard. Buit. II. **5**. 138. 1905 — Borneo.

? *Hamiltoniana* Zoll. Syst. Verz. 47. 1854 (nomen) — Malacca.

hawaiiensis Hill. Fl. Haw. 544. 1888 = G. owhyhensis.

hecistophylla A. Cunn. in Hk. Comp. Bot. Mag. **2**. 361. 1836 = G. alpina.

Hermanni R.Br. Prod. Fl. N. Holl. 161. 1810 = G. linearis.

Hermanni Hk. et Grev. Ic. Fil. t. 14. 1827 = G. pectinata.
hirta Bl. Enum. 250. 1828. HB. 14 — Malesia.
hispida Mett.; Kuhn, Verh. zool. bot. Ges. **25**. 600. 1875 — Java. Celebes. Ternate.
hypoleuca Sod. Rec. 8. 1883; Cr. vasc. quit. 36. 1893 — Ecuador.
immersa Spr. Syst. **4**. 27. 1827 = G. bifida.
intermedia Bak. JoB. **1887**. 24 (excl. var.?) — Costa Rica.
intermedia Bak. l. c. { var. dissitifolia = ? G. orthoclada. var. flexuosa = G. retroflexa.
japonica Spr. Syst. **4**. 25. 1827 = G. glauca.
javanica Spr. Syst. **4**. 25. 1827 = G. laevigata.
Klotzschii Hk. sp. **1**. 13 t. 5 B. 1844 = G. linearis var.
Koordersii Christ, Ann. Jard. Buit. **15**. 76 t. 13 f. 1. 1897 — Celebes. Luzon.
laevigata (Willd.) Hk. sp. **1**. 10. 1844; Mett. Ann. Lugd. Bat. **1**. 49. 1863 — Malesia. — Mertensia Willd. sp. **5**. 75. 1810; Sticherus Pr. 1836. Gleichenia bifurcata Bl. 1828; Mertensia Kze. 1846; Mesosorus Hassk. 1856 — Mertensia plumaeformis Pr. 1848 — ? M. truncata Willd. 1804; Gleichenia Spr. 1827 — G. flagellaris auctt. quoad pl. moles.
laevissima Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 268 — China.
lanigera Don, Prod. Fl. Nepal. 17. 1825 = G. linearis.
lanosa Christ in Schwacke, Pl. Nov. Mineiras **2**. 35. 1900. Bull. Boiss. II. **2**. 691. 1902 — Brasilia.
Lechleri Mett.; Kuhn, Linn. **36**. 167. 1869 — Peru.
Lehmannii Hier. Engl. Jahrb. **34**. 562. 1905 — Columbia.
leucocarpa Sod. Rec. 8. 1883; Cr. vasc. quit. 36. 1893 — Ecuador.
Liebmanni Moore, Ind. 379. 1862 — Mexico.
Mertensia gleichenioides Liebm. Vid. Selsk. Skr. V. **1**. 296 (seors. 144). 1849.
linearis (Burm.) Clarke, Tr. Linn. Soc. II. Bot. **1**. 428. 1880. Chr. 343. NPfl. 355 — Trop. et subtrop. N. Zealand. — Polypodium Burm. Fl. Ind. 235 t. 67 f. 2. 1768 (non Thbg. 1784); Mertensia Fritsch 1901; Polypodium pedatum Houtt. 1783; P. dichotomum Thbg. 1784; Mertensia Willd. 1804; Dicranopteris Bernh. 1806; Gleichenia Hk. 1844. HB. 15; Mesosorus Hassk. 1856 — Gleichenia Hermanni R.Br. 1810; Mertensia Poir. 1813 — Gleichenia lanigera Don 1825; Sticherus Pr. 1836 — Gleichenia ferruginea Bl. 1828 (non Mertensia Desv. 1811) — Mertensia rufinervis Mart. 1834; Gleichenia Hk. 1844. G. Klotzschii Hk. 1844; Mertensia Brack. 1854 — M. emarginata Brack. 1854 (non Raddi 1825); Gleichenia Moore 1862 — Mertensia Lessonii A. Rich. 1834. Gleichenia rigida J. Sm. 1841; Mertensia J. Sm. 1843 (non Kze. 1834); M. crassifolia Pr. 1848 — M. pteridifolia Pr. 1848; Gleichenia Ces. 1876.
littoralis (Phil.) C. Chr. Ind. 322. 1905 — Chile.
Mertensia Phil. Anal. Univ. Chil. **1873**. 104.
littoralis Col. Tr. N. Zeal. Inst. **16**. 334. 1884 = G. flabellata.
longipes (Fée) Christ in Schwacke, Pl. Nov. Mineiras **2**. 35. 1900; Bull. Boiss. II. **2**. 690. 1902 — Brasilia.
Mertensia Fée, Cr. vasc. Br. **2**. 87 t. 105 f. 2. 1872-73.
longipinnata Hk. sp. **1**. 9. 1844 — Amer. trop.
Mertensia Kl. 1844. Gleichenia pubescens pt. HB. 14.
longissima Bl. Enum. 250. 1828 = G. glauca.
maritima Hier. Engl. Jahrb. **34**. 562. 1905 — Columbia.

Mathewsii Hk. sp. **1**. 9 t. 7 B. 1844 — Mexico-Peru. Jamaica.
Mertensia Liebm. 1849.

microphylla R.Br. Prod. Fl. N. Holl. 161. 1810 = G. circinnata.

microphylla Christ, Farnkr. 339. 1897 = G. platyzoma.

Milnei Bak. Syn. 449. 1874; Ic. pl. t. 1602. NPfl. 353 — Aneiteum. Vanecolla.

moniliformis Moore, Ind. 379. 1862 = Stromatopteris.

mucronata Reinw.; J. Sm. JoB. **3**. 430. 1841 = G. linearis.

muricata Klf.; Spr. Syst. **4**. 27. 1827 = G. flagellaris.

nervosa (Klf.) Spr. Syst. **4**. 25. 1827. Chr. 343 — Brasilia austr. Paraguay. Uruguay. Argentina. — Mertensia Klf. Enum. 37. 1824; Gleichenia dichotoma var. HB. 449 — G. neglecta Sturm 1859 — Mertensia latissima Fée 1869.

nitida Pr. Rel. Haenk. **1**. 70. 1825 — Mexico. — Mertensia Pr. 1836.

Norrisii Mett.; Kuhn, Linn. **36**. 165. 1869. HB. 449 — Malacca.

nuda (Moritz) Moore, Ind. 380. 1862 — Caracas.
Mertensia Moritz; Reichardt, Denkschr. Ak. Wien 17^2. 28 t. 1 f. 1-4. 1859.

oceanica Kuhn, Verh. zool. bot. Ges. **19**. 583. 1869. HB. 449. Chr. 342. NPfl. 353 — Celebes. Polynesia.

oligocarpa (Phil.) C. Chr. Ind. 323. 1905 — Chile.
Mertensia Phil. Anal. Univ. Chil. **94**. 361. 1896.

orthoclada Christ, Bull. Boiss. II. **5**. 16. 1905 — Costa Rica.
? G. intermedia var. dissitifolia Bak. JoB. **1887**. 24.

owhyhensis Hk. sp. **1**. 9. 1844. HB. 14. NPfl. 353 — Ins. Hawaii.
Mertensia hawaiiensis Brack. 1854; Gleichenia Hill. 1888.

pallescens Mett. Ann. sc. nat. V. **2**. 267. 1864 — Columbia.

palmata (Schaffn.) Moore, Ind. 380. 1862 — Mexico. Puerto Rico.
Mertensia Schaffn.; Fée, 9 mém. 32. 1857; Fourn. Mex. pl. **1**. 137. 1872.

patens Col. Tr. N. Zeal. Inst. **20**. 212. 1888 = G. circinnata.

pectinata (Willd.) Pr. Rel. Haenk. **1**. 71. 1825. HB. 15. Chr. 342. NPfl. 354 — Amer. trop. — Mertensia Willd. Vet. Ak. Nya Handl. **1804**. 168 t. 4. M. glaucescens H.B.Willd. 1810; Gleichenia HBK. 1815. Mertensia brasiliana Desv. 1811; Gleichenia Spr. 1827. G. Hermanni Hk. et Grev. 1827 (non R.Br. 1810). Mertensia emarginata Raddi 1825.

pedalis (Klf.) Spr. Syst. **4**. 26. 1827. HB. 13. NPfl. 353 — Chile. Juan Fernandez. — Mertensia Klf. Enum. 39. 1824.

pennigera (Mart.) Moore, Ind. 381. 1862 — Brasilia. Costa Rica.
Mertensia Mart. Ic. Cr. Bras. 130 t. 59 f. 1. 1834; M. gracilis Mart. l. c. 106 (non 107).

pinnata Moore, Ind. 381. 1862 = G. glauca var.

platyzoma F. Muell. Veg. of the Chatham Islands 63. 1864 — Australia.
Platyzoma microphylla R.Br. Prod. Fl. N. Holl. 160. 1810. HB. 11; Gleichenia Christ 1897. 339. NPfl. 355 (non R.Br. 1810).

polypodioides (L.) Sm. Mém. Ac. Turin **5**. 419 t. 9 f. 10. 1793. HB. 11. Chr. 338. NPfl. 355 — Africa austr. Ins. Amsterdam.
Onoclea L. Mant. 306. 1771 — Gleichenia glauca Sw. 1806. G. argentea Klf. 1824. Mertensia coeruleo-glauca Poir. 1813.

pruinosa (Mart.) Mett. Ann. Lugd. Bat. **1**. 49. 1863 — Brasilia.
Mertensia Mart. Ic. Cr. Bras. 109. 1834. M. subflabellata Brack. 1854.

pteridifolia Ces. Atti Ac. Napoli 7^8. 2. 1876 = G. linearis.

pubescens HBK. Nov. Gen. et Sp. **1**. 29. 1815 = G. furcata.

pubescens Hk. Syn. 13. 1865 et auctt. = G. bifida, furcata, longipinnata, Mathewsii.

quadripartita (Poir.) Moore, Ind. 382. 1862. HB. 13. Chr. 340. NPfl. 353 — Amer. antarct. — Polypodium Poir. Enc. **5**. 543. 1804; Mertensia Poir. 1813. Gleichenia acutifolia Hk. 1844; Mertensia Gay 1853.

remota (Klf.) Spr. Syst. **4**. 27. 1827 — Brasilia. Columbia.
Mertensia Klf. Enum. 39. 1824.

retroflexa Bomm.; Christ, Bull. Boiss. **4**. 657. 1896; II. **5**. 15 c. fig. — Costa Rica. ? G. intermedia var. flexuosa Bak. JoB. **1887**. 24 (non G. flexuosa Mett. 1863).

revoluta HBK. Nov. Gen. et Sp. **1**. 29. 1815. HB. 13. Chr. 341. NPfl. 353 — Jamaica. Costa Rica-Peru. Brasilia. — Mertensia Desv. 1827.

rigida J. Sm. JoB. **3**. 420. 1841 = G. linearis.

rigida Bomm. et Christ, Bull. Soc. bot. Belg. **35**. 174. 1896 = G. flexuosa.

rubiginosa Mett. Ann. sc. nat. V. **2**. 267. 1864 — Columbia. Peru.

rufinervis Hk. sp. **1**. 11. 1844 = G. linearis var.

rupestris R.Br. Prod. Fl. N. Holl. 160. 1810. HB. 12 — Australia. N. Caledonia.
Mertensia Poir. 1813; Platyzoma Desv. 1827; Gleicheniastrum Pr. 1848.

Sartorii (Fée) Moore, Ind. 383. 1862 — Mexico. — Mertensia Fée, 9 mém. 32. 1857.

seminuda (Kl.) Moore, Ind. 383. 1862 — Brasilia. Caracas.
Mertensia Kl. Linn. **18**. 538. 1844.

semivestita Lab. Sert. Austr. Cal. 8 t. 11. 1824 = G. circinnata.

simplex (Desv.) Hk. Ic. pl. t. 92. 1837. HB. 13. Chr. 343. NPfl. 353 — Peru-Ecuador. — Mertensia Desv. Prod. 200. 1827.

speluncae R.Br. Prod. Fl. N. Holl. 160. 1810 = G. circinnata var.

squamulosa (Desv.) Moore, Ind. 383. 1862 — Amer. austr.
Mertensia Desv. Journ. de bot. appl. **1**. 268. 1813.

strictissima Christ, Bull. Boiss. II. **5**. 13. 1905 — Costa Rica.

subandina Sod. Rec. 5. 1883; Cr. vasc. quit. 34. 1893 — Ecuador.

subpectinata Christ, Bot. Tidsskr. **24**. 111. 1901 — Siam (Koh Chang).

subtrisperma (Fée) Krug, Engl. Jahrb. **24**. 78. 1897 — Guadeloupe.
Mertensia Fée, 11 mém. 122 t. 32 f. 2. 1866.

tenera R.Br. Prod. Fl. N. Holl. 161. 1810. HB. 12 — Tasmania.
Mertensia Poir. 1813.

tenuis Pr. Rel. Haenk. **1**. 70. 1825 — Mexico. — Mertensia Pr. 1836.

tomentosa Spr. Syst. **4**. 27. 1827 = M. furcata.

trifurcans (Fée) Christ in Schwacke, Pl. Nov. Mineiras 2. 35. 1900; Bull. Boiss. II. **2**. 691. 1902 — Brasilia. — Mertensia Fée, Cr. vasc. Br. **1**. 201 t. 74 f. 1. 1869. (an M. decurrens Raddi 1825?).

truncata Spr. Syst. **4**. 25. 1827 = M. laevigata.

tuberculata Mett.; Kuhn, Linn. **36**. 166. 1869 — Peru.

umbraculifera (Kze.) Moore, Ind. 384. 1862. HB. 13. Chr. 340. NPfl. 353 — Africa austr. — Mertensia Kze. Linn. **18**. 114. 1844.

velata (Kze.) Mett. Fil. Lips. 113. 1856; Sod. Cr. vasc. quit. 37. 1893 — Andes.
Mertensia Kze. Linn. **9**. 15. 1834.

vestita Bl. Enum. 249. 1828. HB. 14. NPfl. 353 — Malesia.
Mertensia Kze. 1848; Mesosorus Hassk. 1856.

volubilis Jungh. Reisen durch Java 452. 1845; Flora **1847**. 525 = G. glauca var. arachnoides.

vulcanica Bl. Enum. 251. 1828; Mett. Ann. Lugd. Bat. **1**. 46. 1863 — Malesia.
Calymella Pr. 1836. Gleichenia dicarpa part. HB. 12.

Warburgii Christ, Ann. Jard. Buit. **15**. 78. 1897. NPfl. 353 — Celebes. Batjan. Borneo.

GLEICHENIASTRUM Presl, Abh. böhm. Ges. V. **5**. 338. 1848 = **Gleichenia.**
Boryi Pr. l. c. 338.
microphyllum Pr. l. c. 338 = G. circinnata.
rupestre Pr. l. c. 338.
semivestitum Pr. l. c. 338 = G. circinnata.
speluncae Pr. l. c. 338 = G. circinnata var.
GLOSSOPTERIS Rafinesque, Journ. Phys. **89**. 262. 1819 = **Phyllitis.**
GLYPHOTAENIUM J. Smith, Hist. Fil. 187. 1875 = **Polypodium.**
Ctenopteris § Glyphotaenium J. Sm. Bot. Voy. Herald 227. 1854.
crispatum J. Sm. Hist. Fil. 188. 1875 = P. goniopteroides.
Goniolepicystis Diels, Nat. Pflanzenfam. **1**[4]. 323. 1899 (*Lepicystis* §) = Polypodium.
GONIOPHLEBIUM (Bl.) Presl, Tent. Pterid. 185. 1836 = **Polypodium.**
Polypodium § Goniophlebium Blume, Fl. Javae Fil. 132. 1828.
acuminatum Fée, 11 mém. 68 t. 19 f. 1. 1866 = P. brasiliense var.
adnatum Moore, Ind. 385. 1862.
albo-punctatum J. Sm. Bot. Mag. **72** Comp. 12. 1846 = P. brasiliense.
amoenum J. Sm. in Hk. Gen. ad t. 51. 1840.
angustifolium Brack. Expl. Exp. **16**. 33. 1854.
anisomeron Fée, 8 mém. 92. 1857 = P. plesiosorum var.
appendiculatum Moore, Gard. Chr. **1856**. 820 = ? P. plesiosorum var.
arcuatum Fée, Gen. 255. 1850-52 — Martinique. Brasilia.
an P. loriceum vel P. falcaria var?
areolatum Pr. Tent. 186. 1836 = P. aureum*.
argutum J. Sm. in Hk. Gen. ad t. 51. 1840.
articulatum Pr. Tent. 186. 1836 = P. fraxinifolium.
assimile Lodd. Cat. 1849 = P. chnoodes.
attenuatum Pr. Tent. 186. 1836.
aurisetum Brack. Expl. Exp. **16**. 31. 1854 = P. tectum.
biauriculatum Moore, Ind. 386. 1862.
blepharodes Moore, Ind. 386. 1862 = Marginaria.
Calaguala Fée, 8 mém. 93. 1857 = P. falcaria.
Californicum Fée, Gen. 255. 1850-52 = P. synammia.
californicum Moore, Ind. 386. 1862.
cardiophyllum Moore, Ind. 387. 1862 = ? P. vacciniifolium.
Catharinae J. Sm. in Hk. Gen. ad t. 51. 1840.
caudiceps Moore, Gard. Chr. n. s. **25**. 234. 1886.
ceteraccinum Fée, Cr. vasc. Br. **1**. 107. 1869 = P. polypodioides.
chnoodes Fée, Gen. 255. 1850-52.
chondrocheilon Fée, 8 mém. 92. 1857.
ciliatum J. Sm. in Hk. Gen. ad t. 51. 1840.
colpodes J. Sm. Cat. Kew Ferns 1. 1856 = P. plesiosorum.
confluens Moore, Ind. 388. 1862 = P. conterminans.
cordatum Moore, Ind. 388. 1862 = P. Kunzeanum.
coriaceum Fée, 10 mém. 33 t. 37 f. 2. 1865 = P. coriaceum Kuhn.
crassimargo Moore, Ind. 388. 1862 = P. lepidopteris.
crenatum Moore, Ind. 388. 1862 (nomen) — Ecuador.
? cuspidatum Pr. Tent. 186. 1836 = P. persicifolium.
dasypleuron Moore, Ind. LXXIV. 1857.
deflexum Moore et Houlst. Gard. Mag. Bot. **3**. 61. 1851 = P. fraxinifolium.
deltoideum Moore, Ind. 388. 1862 = P. catharinae.

demissum Fée, Cr. vasc. Br. **2**. 63. 1872-73.
dissimile J. Sm. JoB. **4**. 57. 1841 = ? P. dissimile vel potius P. chnoodes.
distans J. Sm. Lond. JoB. **1**. 195. 1842 = P. fraxinifolium.
dulce Moore, Ind. 388. 1862 = ? P. aureum.
elatius Moore, Ind. 389. 1862 = P. brasiliense.
elatum Fée, Gen. 256. 1850-52 = ? P. attenuatum an P. sororium?
eleutherophlebium Fée, Gen. 255. 1850-52.
ensifolium Brack. Expl. Exp. **16**. 33. 1854 = P. angustifolium var.
ensiforme Fée, Gen. 255. 1850-52.
erythrocarpum Bedd. Ferns br. Ind. Suppl. 21 t. 382. 1876.
excelsior Fée, Cr. vasc. Br. **1**. 244 t. 77 f. 4. 1869 = P. brasiliense.
falcarium Fée, Gen. 255. 1850-52 = P. falcaria.
Fieldingianum Moore, Ind. 389. 1862 = P. lachnopus.
fraxinifolium Moore, Ind. LXXIV. 1857.
furfuraceum Moore, Ind. LXXIV. 1857.
Gauthieri Fée, Cr. vasc. Br. **1**. 112 t. 34 f. 3. 1869.
giganteum Moore, Ind. 390. 1862.
gladiatum Fée, Gen. 256. 1850-52 = P. attenuatum.
glaucophyllum Fée, Gen. 255. 1850-52.
glaucum J. Sm. Cat. cult. ferns 3. 1857 = P. catharinae.
grammatoides Fée, Cr. vasc. Br. **1**. 110 t. 34 f. 2. 1869 = P. latipes.
grandidens Fée, Gen. 255. 1850-52 = P. persicifolium var.
guatemalense Moore, Ind. 390. 1862.
Haenkei Pr. Tent. 186 t. 7 f. 14. 1836 = P. attenuatum.
harpeodes J. Sm. Bot. Mag. **72** Comp. 11. 1846 = P. latipes.
Hasskarlii Kl.; van Houtte, Cat. 1858; Moore, Ind. 390. 1862 — ? Java.
Hendersoni Bedd. Ferns br. Ind. Suppl. 21 t. 383. 1876 = P. Atkinsoni.
heteroneuron Fée, 9 mém. 24. 1857 — Mexico.
hirsutissimum Brack. Expl. Exp. **16**. 33. 1854 = P. lepidopteris.
hirsutulum Moore, Ind. 390. 1862.
inaequale J. Sm. Cat. cult. ferns 3. 1857 = P. Lowei.
incanum J. Sm. JoB. **4**. 56. 1841 = P. polypodioides.
intermedium Fée, 11 mém. 69 t. 18 f. 3. 1866 = P. brasiliense.
invertens Fée, 9 mém. 25. 1857; 10 mém. 33 t. 37 f. 3. 1865 = P. falcaria.
javanense Moore, Ind. 390. 1862 = P. accedens.
Korthalsii Bedd. Handb. Suppl. 91. 1892.
lachnopus J. Sm. in Hk. Gen. ad t. 51. 1840.
laetum J. Sm. Bot. Voy. Herald 231. 1854 = P. loriceum var.
lanuginosum Moore, Ind. 391. 1862.
lasiopus Moore, Ind. 391. 1862.
latipes J. Sm. Bot. Mag. **72** Comp. 12. 1846.
lepidopteris Moore, Ind. LXXIV. 1857.
lepidotrichum Fée, 8 mém. 93. 1857 = P. squamatum.
ligustrifolium Moore, Ind. 392. 1862 = P. lycopodioides.
longicaule Fée, 8 mém. 95. 1857.
loriceum J. Sm. in Hk. Gen. ad t. 51. 1840.
macrocarpum J. Sm. JoB. **4**. 56. 1841 = P. pycnocarpum.
menisciifolium J. Sm. in Hk. Gen. ad t. 51. 1840 = P. brasiliense var.
microlepis Fée, Cr. vasc. Br. **1**. 107. 1869 = P. polypodioides.
microrhizoma Bedd. Ferns br. Ind. Suppl. 21 t. 384. 1876.

molestum Fée, 8 mém. 92. 1857 = P. plesiosorum.
molle Bedd. Ferns br. Ind. t. 206. 1867 = P. Beddomei.
moupinense Bedd. Handb. Suppl. 91. 1892.
myrtillifolium Moore, Ind. LXXIV. 1857 = P. lycopodioides.
neriifolium Hk. Gen. t. 70 B. 1841 = P. brasiliense.
nummularium Moore, Ind. LXXIV. 1857 = P. pyrolifolium.
oleandrifolium Moore, Ind. 394. 1862 = P. attenuatum.
ornatum Fée, Gen. 255. 1850-52 = P. fraxinifolium.
owariense Lodd. Cat. 1849 = P. lycopodioides.
pachynevron Fée, Gen. 255. 1850-52 (nomen)?
pachyphyllum Moore, Ind. 394. 1862 = P. Scouleri.
?pallens Pr. Tent. 186. 1836 = P. subauriculatum var.
patens J. Sm. Bot. Voy. Herald 230. 1854.
pectinans Fée, Cr. vasc. Br. 1. 109 t. 34 f. 1. 1869 = P. latipes.
pectinatum J. Sm. JoB. **4**. 57. 1841.
pectinatum J. Sm. Bot. Voy. Herald 230. 1854 = P. loriceum var.
pellitum Moore, Ind. 395. 1862.
persicaefolium Bedd. Ferns br. Ind. Correct. 1870.
pictum Fée, Cr. vasc. Br. 1. 244 t. 77 f. 3. 1869 = P. catharinae.
piloselloides J. Sm. in Hk. Gen. ad t. 51. 1840.
plectolepis Fée, 8 mém. 95. 1857.
Pleopeltis Fée, Gen. 256. 1850-52 = P. subauriculatum.
plesiosorum Fée, 9 mém. 24. 1857.
Prainii Bedd. JoB. **1893**. 226.
pringlei Maxon, Proc. U. S. Nat. Herb. **27**. 953 t. 48. 1904.
pubescens Moore, Ind. 395. 1862 = P. Gilliesii.
punctulatum Moore, Ind. 395. 1862 = P. falcaria var.
pyrrholepis Fée, 8 mém. 94. 1857 = P. lepidopteris.
ramosum Fée, Gen. 255. 1850-52 = P. loriceum.
Reinwardtianum Fée, Gen. 255. 1850-52 } = P. subauriculatum.
Reinwardtii De Vriese, Ned. kr. Arch. 1. 257. 1846 } = P. subauriculatum.
retrofractum Moore, Ind. 396. 1862.
revertens Fée, 8 mém. 94. 1857.
rachipterygium Moore, Ind. 396. 1862 = P. synammia var.
rhagadiolepis Fée, 7 mém. 69 t. 19 f. 3. 1857 = P. thyssanolepis.
rhizocaulon Pr. Tent. 186. 1836 = P. fraxinifolium var.
rhodopleuron Fée, 9 mém. 24. 1857 = P. plesiosorum var.
rigidulum Moore, Ind. 396. 1862 = Drynaria r.
Schaffnerianum Moore, Ind. 396. 1862 = P. squamatum.
Scouleri J. Sm. Hist. Fil. 91. 1875.
semipinnatifidum Fée, Gen. 256. 1850-52 = P. glaucophyllum var.
sepultum J. Sm. in Hk. Gen. ad t. 51. 1840 = P. lepidopteris.
?serpens Fée, Gen. 255. 1850-52 = P. Swartzii.
serratifolium Brack. Expl. Exp. **16**. 35. 1854 = P. subauriculatum var.
serratum Fée, 8 mém. 95. 1857 = P. sessilifolium.
solutum Fée, Gen. 255. 1850-52 = P. angustifolium.
squamatum Moore, Ind. 391. 1862.
subamoenum Bedd. Handb. 317. 1883.
?subauriculatum Pr. Tent. 186. 1836.
surucuchense Moore, Ind. LXXIV. 1857.

Synammia Fée; Gay, Fl. chil. **6**. 510. 1853.
tectum J. Sm. in Hk. Gen. ad t. 51. 1840.
thyssanolepis Moore, Ind. 396. 1862.
translucens Fée, Gen. 255. 1850-52.
trilobum Moore, Ind. LXXIV. 1857 = P. synammia.
Tweedianum J. Sm. JoB. **4**. 56. 1841.
vacciniifolium J. Sm. in Hk. Gen. ad t. 51. 1840.
vacillans Fée, Gen. 255. 1850-52 = P. latipes.
verrucosum J. Sm. in Hk. Gen. ad t. 51. 1840.
Villeminianum Fée, 7 mém. 63 t. 27 f. 3. 1857 = P. Mathewsii.
xiphophorum Fée, Cr. vasc. Br. **1**. 108. 1869 = P. attenuatum.
yunanense Bedd. Handb. Suppl. 89. 1892.

GONIOPTERIS Presl, Tent. Pterid. 181. 1836 = **Dryopteris.**
abbreviata Pr. Tent. 183. 1836 = ? D. brachyodus.
affinis Fée, Gen. 250. 1850-52 = D. scolopendrioides.
amplexicaulis Pr. Tent. 183. 1836 (nomen) — ?
appendiculata Pr. Tent. 183 t. 7 f. 11. 1836.
aristata Fée, Gen. 253. 1850-52 = D. otaria.
aspera Pr. Tent. 183. 1836 = D. urophylla.
aspera J. Sm. JoB. **3**. 396. 1841 = D. ferox.
asplenioides Pr. Tent. 182. 1836.
asterothrix Fée, Gen. 253. 1850-52 = D. reptans var.
asymetrica Fée, Gen. 253. 1850-52 = D. canescens*.
Bahiensis Fée, Cr. vasc. Br. **2**. 61. 1872-73 = D. tetragona.
barbata Fée, Gen. 252. 1850-52 = D. ferox.
Berteriana Pr. Tent. 183. 1836 (nomen) — ?
Besckeana Pr. Tent. 182. 1836 (nomen) — ?
canescens Pr. Tent. 183. 1836.
cheilocarpa Fée, Gen. 251. 1850-52 = D. gongylodes.
costata J. Sm. JoB. **4**. 54. 1841 = D. penangiana.
costata Brack. Expl. Exp. **16**. 28. 1854 = D. pennigera.
crenata Pr. Tent. 183 t. 7 f. 10. 1836 = D. Poiteana.
crenato-dentata Fée, Gen. 252. 1850-52 = D. otaria.
crispata Moore, Ind. LXXII. 1857 = Polypodium goniopteroides.
Dalhousiana Fée, 8 mém. 92. 1857 — India.
(poss. = D. urophylla).
domingensis Fée, 11 mém. 59. 1866 = D. scolopendrioides.
faucia Fourn. Mex. pl. **1**. 88. 1872 = D. obliterata.
Féei L'Herm.; Fée, 11 mém. 61 t. 17 f. 1. 1866 = D. tetragona var.
ferax Fée, Gen. 250. 1850-52 = D. scolopendrioides.
Forsteri Moore, Ind. 99. 1858 = D. pennigera.
fraxinifolia Pr. Tent. 182 t. 7 f. 9. 1836 = D. vivipara.
Ghiesbrechtii Bail. Hdb. Ferns Queensl. 40. 1874 = D. Hillii.
Ghiesbreghtii J. Sm. Ferns br. and for. 138. 1866 (Gheisbeghtii); Fourn. Mex. pl. **1**. 89. 1872.
glandulifera Brack. Expl. Exp. **16**. 29. 1854 = D. subspinosa.
gracilis Moore et Houlst. Gard. Chr. **1856**. 301. c. fig. = D. reptans var.
granulosa Pr. Tent. 183. 1836.
guadalupensis Fée, 11 mém. 64 t. 17 f. 2. 1866 = D. tetragona var.
hastata Fée, 11 mém. 65 t. 18 f. 1. 1866.

hastata Fée, Cr. vasc. Br. 1. 107 t. 33 f. 2. 1869 = D. tetragona var. bahiensis.
incisa Pr. Tent. 182. 1836.
Kennedyi Bail. Hdb. Ferns Queensl. 41. 1874 = D. urophylla.
leptocladia Fée, 11 mém. 60 t. 16 f. 1. 1866 = D. tetragona var.
lineata Pr. Tent. 183. 1836 = D. penangiana.
lineata Bedd. Ferns br. Ind. t. 3. 1865 = D. moulmeinensis.
Lobbiana Fée, Gen. 251. 1850-52 = D. ferox.
longissima Brack. Expl. Exp. 16. 29 t. 5. 1854.
lucida Fée, Gen. 253. 1850-52 — Guadeloupe.
luzonensis Pr. Tent. 182. 1836 — Luzon.
Polypodium Pr. Rel. Haenk. 1. 26. 1825.
macrocladia Fée, Cr. vasc. Br. 1. 106 t. 33 f. 3. 1869 = D. tetragona.
Madagascariensis Fée, Gen. 251. 1850-52.
megalodus Pr. Tent. 182. 1836.
meniscioides Fée, Gen. 253. 1850-52 = D. prolifera.
meniscioides Fourn. Mex. pl. 1. 88. 1872.
mollis Fée, Gen. 252. 1850-52 = P. Ghiesbreghtii.
multilineata Bedd. Ferns br. Ind. t. 231. 1867 = D. moulmeinensis.
obliterata Pr. Tent. 183. 1836.
oppositifolia J. Sm. Hist. Fil. 192. 1875.
patens Fée, Gen. 253. 1850-52 = D. silvatica.
Penangiana Bedd. Ferns br. Ind. t. 232. 1867.
pennigera Pr. Tent. 183. 1836 = D. gongylodes.
pennigera J. Sm. JoB. 4. 54. 1841.
pilosa Pr. Tent. 182. 1850-52 = Polypodium p. (Schkuhr).
platypes Fée, Cr. vasc. Br. 1. 106 t. 33 f. 1. 1869 = D. tetragona.
poecilophlebia Bail. Handb. Ferns Queensl. 40. 1874.
portoricensis Pr. Tent. 182. 1836 = D. megalodus.
prolifera Pr. Tent. 183. 1836.
prolifera Fée, Cr. vasc. Br. 1. 105. 1869 = D. vivipara.
pyramidata Fée, 11 mém. 61 t. 16 f. 2. 1866 = D. tetragona.
quadrangularis Fée, 11 mém. 63 t. 16 f. 3. 1866 = D. tetragona.
refracta J. Sm. Cat. cult. ferns 20. 1857.
repanda Fée, Gen. 251. 1850-52 — India. (an = D. urophylla?).
reptans Pr. Tent. 182. 1836.
Rivoirei Fée, Gen. 253. 1850-52 = D. Poiteana.
rostrata Fée, 11 mém. 64 t. 17 f. 3. 1866 = D. meniscioides.
rubida J. Sm. JoB. 3. 395. 1841.
rubrinervis Carr. in Seem. Fl. Vit. 366. 1873.
sandvicense J. Sm. Hist. Fil. 192. 1875 = D. stegnogrammoides.
scolopendrioides Pr. Tent. 182. 1836.
serrulata J. Sm. Cat. cult. ferns 20. 1857.
silvatica Pappe et Raws Syn. Fil. Afr. austr. 39. 1858.
simplicifolia Carr. in Seem Fl. Vit. 366. 1873 = D. canescens*.
spectabilis (Willd.) Pr. Tent. 183. 1836 (nomen) — ? — Polypodium Willd. hb.
stegnogrammoides J. Sm. Ferns br. and for. ed. II. 303. 1877.
strigosa Fée, 11 mém. 59 t. 15 f. 1. 1866 = D. scolopendrioides.
submarginalis Pr. Tent. 182. 1836 = D. tetragona.
subpectinata J. Sm. Hist. Fil. 192. 1875 = D. otaria.
subtetragona Pr. Tent. 182. 1836 = D. tetragona.

tenera Fée, 11 mém. 60 t. 15 f. 3. 1866 = D. asplenioides.
tetragona Pr. Tent. 183. 1836.
trifurcata Moore, Ind. LXXII. 1857 = Polypodium t.
unita J. Sm. Hist. Fil. 192. 1875 = D. silvatica.
urophylla Pr. Tent. 183. 1836.
vivipara Brack. Expl. Exp. **16**. 27. 1854.

GONOCORMUS Van den Bosch, Hymenoph. Jav. 7. 1861 = **Trichomanes.**
dichotomus Prantl, Hym. 51. 1875.
diffusus v.d.B. Hym. Jav. 9 t. 4. 1861.
digitatus Prantl, Hym. 51. 1875.
flabellatus Prantl, Hym. 51. 1875 = T. digitatum var.
minutus v.d.B. Hym. Jav. 7 t. 3. 1861.
nitidulus Prantl, Hym. 51. 1875.
palmatifidus Prantl, Hym. 51. 1875.
palmatus v.d.B. Hym. Jav. 11 t. 6. 1861 = T. proliferum.
prolifer Prantl, Hym. 51. 1875.
saxifragoides v.d.B. Hym. Jav. 9. 1861.
Teysmani v.d.B. Hym. Jav. 10 t. 5. 1861.

GRAMMATOSORUS Regel, Ind. Sem. ht. Petr. 75. 1866; Gartenflora **15**. 335. 1866 = ? **Aspidium.**
Blumeanus Regel, l. c.; Gartenfl. **15**. 335 c. figg. 1866 — Hort.
Aspidium Grammatosorus Salomon 1883. — Conf. Milde, Bot. Zeit. **1868**. 614.

Grammitastrum Fournier, Ann. sc. nat. V. **18**. 282. 1873 (*Polypodium §*) = Polypodium.

GRAMMITIS Swartz, Schrad. Journ. **1800**[2]. 17. 1801 = **Polypodium.**
acuminata Wall. List n. 7. 1828 = P. scolopendrinum.
adiantoides Griff. Notulae **2**. 609; Ic. **2**. t. 134 f. 2. 1849 = Anogramma leptophylla.
adspersa Bl. Fl. Jav. Fil. 115 t. 48 f. 2. 1828 = P. hirtellum var.
affinis Wall. List n. 11. 1828 = Coniogramme fraxinea.
ampla F. Muell. Fragm. **5**. 188. 1866 (nomen); Benth. Fl. austr. **7**. 777. 1878 [= P. ellipticum.
angustata Desv. Prod. 221. 1827 = ? P. Billardieri.
angustifolia Hew. Mag. Nat. Hist. II. **2**. 458. 1838.
araucana Phil. Linn. **33**. 304. 1865 = P. Billardieri.
Ascensionis Hk. Ic. pl. t. 967. 1854 = Anogramma a.
aspidioides Moore, Ind. LIII. 1857 = Dryopteris a.
aspidioides Hk. in Thwait. Enum. pl. Zeyl. 383. 1864 = Dryopteris stegno- [gramme.
asplenioides Pr. Del. Prag. **1**. 163. 1822 = Dryopteris apidioides.
athroosperma Fourn. Bull. Soc. Fr. **16**. 423. 1869 = P. Deplanchei.
attenuata Kze. Linn. **24**. 251. 1851 = P. mediale var.
aurea Sw. Syn. 23, 215. 1806 = Ceterach a.
aurita Moore, Gard. Chr. **1856**. 389 c. fig. = Dryopteris a.
australis R.Br. Prod. Fl. N. Holl. 146. 1810 = P. Billardieri.
avenia Bl. Enum. 117. 1828 = P. Blumeanum.
Billardieri Willd. sp. **5**. 139. 1810.
bisulcata Hk. Ic. pl. t. 998. 1854.
blechnoides Grev. Ann. Mag. Nat. Hist. II. **1**. 320 t. 17. 1848.
Blumeana Pr. Tent. 209. 1836 = Dryopteris africana.
caespitosa Bl. Enum. 115. 1828.
capensis Moore, Ind. LIII. 1857 = Ceterach cordatum.

caudata Wall. List n. 4. 1828 = Coniogramme fraxinea.
caudiformis Hk. Bot. Mag. t. 5328. 1862 = P. Feei var.
Ceterach Sw. Syn. 23. 1806 = Ceterach officinarum.
cheilanthoides Sw. Syn. 23. 219, 419. 1806 = Gymnogramma c.
ciliata Col. Tasm. Journ. 2. 166. 1844 = P. Billardieri.
clavifer Hk. 2 Cent. t. 5. 1860.
conformis J. Sm. Hist. Fil. 181. 1875 = P. Hookeri var.
congener Bl. Enum. 115. 1828.
cordata Sw. Syn. 23, 217. 1806 = Ceterach c.
coriacea Klf.; Spr. Syst. 4. 41. 1827 = P. loxogramme.
coriacea Bl. Enum. 117. 1828 = P. Blumeanum.
crassa Fée, Gen. 234. 1850-52 = P. Billardieri.
cucullata Bl. Fl. Javae, Fil. 119 t. 50 f. 3. 1828.
cuspidata Zenker, Pl. Ind. 1. t. 2. 1835 = P. scolopendrinum.
decurrens Wall.; Hk. et Grev. Ic. Fil. t. 6. 1827 = P. ellipticum.
decurrens Bl. Enum. 119. 1828 = P. revolutum.
denticulata Bl. Fl. Javae, Fil. 121 t. 50 f. 4. 1828.
diversifolia Wall. List n. 8. 1828 = P. heterocarpum.
elongata Sw. Syn. 22, 213. 1806 = P. lanceolatum*.
fasciata Bl. Enum. 116. 1828.
fasciculata Bl. Fl. Javae, Fil. 112 t. 47 f. 2. 1828 = P. setigerum.
Finlaysoniana Wall. List pag. 248 (n. 776) 1829 = P. digitatum.
flavescens Wall. List n. 6. 1828 = ? P. scolopendrinum.
flexuosa Humb. et Bonpl. (? Pl. aequin. 2. 167 t. 138. 1809); HBK. Nov. Gen. et Sp. 1. 5. 1815 = Gymnogramma f.
fluminensis Fée, Cr. vasc. Br. 1. 85 t. 19 f. 3. 1869 = P. marginellum.
furcata Hk. et Grev. Ic. Fil. t. 62. 1828 = P. dicranophyllum.
graminoides Sw. Schrad. Journ. 1800². 18. 1801 = Monogramma g.
grammitidis Keys. Pol. Cyath. Hb. Bung. 34. 1873.
Hamiltoniana Wall. List n. 9. 1828 = P. pedunculatum.
heterocarpa Bl. Enum. 118. 1828.
heterophylla Lab. N. Holl. pl. spec. 2. 90 t. 239. 1806 = P. grammitidis.
Hewardii Moore, Gard. Chr. 1856. 261 c. fig. = Dryopteris heteroclita.
hirta HBK. Nov. Gen. et Sp. 1. 4. 1815 = Gymnogramma h.
hirta Bl. Enum. 115. 1828 = P. diplosorum.
hispanica Cosson, Notes sur quelq. pl. du midi de l'Espagne 48. —. = Pleurosorus Pozoi.
humilis Homb. et Jacq. Voy. Pôle sud t. 2 f. H. 1852 = P. Billardieri.
involuta Don, Prod. Fl. Nepal. 14. 1825 = P. scolopendrinum.
Iserti Sw. Adnot. 65. 1829 = Monogramma seminuda.
lanceolata Sw. Schrad. Journ. 1800². 18. 1801 = P. loxogramme.
lanceolata Schkuhr, Kr. Gew. 1. 9 t. 7. 1806 = P. lanceolatum*.
leptophylla Sw. Syn. 23. 218 t. 1. f. 6. 1806 = Anogramma l.
limbata Fée, Gen. 233. 1850-52; 6 mém. 6 t. 5 f. 1 = P. marginellum
linearifolia Steud. Nom. 2. 187. 1824 = Monogramma l.
linearis Sw. Schrad. Journ. 1800². 17. 1801 = P. gramineum.
Linkiana Pr. Tent. 209. 1836 = Dryopteris diplazioides.
longa Fée, Gen. 233. 1850-52; 6 mém. 6 t. 4 f. 1 = P. fasciatum (vel P. congenerum).
longifolia Bl. Enum. 119. 1828 = P. revolutum.
macrophylla Bl. Enum. 119. 1828.

macrophylla Wall. List n. 10. 1828 = P. scolopendrinum.
magellanica Desv. Berl. Mag. **5**. 313. 1811 = P. Billardieri.
marginella Sw. Schrad. Journ. **1800**^{2}. 17. 1801.
membranacea Bl. Enum. 118. 1828 = P. selliguea.
microphylla Bedd. Ferns br. Ind. t. 148. 1866 = Anogramma m.
Muelleri F. Muell. Fragm. **5**. 138. 1866 = Gymnopteris M.
muscosa Fée, Cr. vasc. Br. **2**. 51 t. 95 f. 2. 1872-73 = P. Wittigianum.
myosuroides Sw. Schrad. Journ. **1800**^{2}. 18. 1801.
muosuroides Schkuhr, Kr. Gew. **1**. 9 t. 7. 1804 = P. delitescens.
nana Fée, 6 mém. 7 t. 6 f. 1. 1853 = ? P. hirtellum.
nana Brack. Expl. Exp. **16**. 1. 1854 = P. Billardieri.
obliquata Hassk. Cat. pl. Bogor. 4. 1844.
obscura Bl. Enum. 115. 1828 = P. Reinwardtii var.
obtusa Willd.; Klf. Enum. 84. 1824 = ? P. furcatum.
obtusata Pr. Tent. 209. 1836 = Dryopteris opaca.
organensis Gardn. in Hk. Ic. pl. t. 509. 1843; Fée, Cr. vasc. Br. **1**. 264 t. 78 [f. 1. 1869.
parietina Fée, Gen. 233. 1850-52.
paucinervata Fée, Cr. vasc. Br. **2**. 51 t. 96 f. 1. 1872-73.
pedunculata Bl. Enum. 118. 1828.
pilosiuscula Bl. Enum. 115. 1828.
pinnata F. Muell. Fragm. **6**. 124. 1868 = Syngramma p.
polypodioides Pr. Tent. 209. 1836 = Dryopteris p.
procera Wall. List n. 3. 1828 = Coniogramme fraxinea.
pseudaustralis Fourn. Bull. Soc. Fr. **16**. 423. 1869.
pubinervis Bl. Enum. 116. 1828.
pumila Sw. Syn. 214, 419. 1806 = Monogramma graminea.
pumila Armstr. Tr. N. Zeal. Inst. **13**. 341. 1882 = P. Billardieri.
punctata Raddi, Pl. Bras. **1**. 11 t. 22 bis f. 1. 1825 = Monogramma seminuda.
pusilla Bl. Fl. Javae, Fil. 109 t. 46 f. 4-6. 1828 = P. hirtellum.
quaerenda Bolle, Zeitschr. f. Erdk. n. Folge **14**. 324. 1863 = P. marginellum.
Reinwardtii Bl. Enum. addend. 1828.
repanda Kze.; Mett. Fil. Lechl. 2. 1856 = P. jungermannioides.
reticulata Griff. Calc. Journ. **4**. 466. 1844 = P. scolopendrinum.
revoluta Spr.; Willd. sp. **5**. 139. 1810 = P. lanceolatum*.
Reynoldsii Benth. Fl. austr. **7**. 775. 1878 = Gymnopteris R.
rigida Hombr. et Jacq. Voy. Pôle sud t. 2 f. F. 1852 = P. Billardieri.
robusta Phil. Anal. Univ. Chil. **94**. 358. 1896 = P. lanceolatum*.
rutaefolia R.Br. Prod. Fl. N. Holl. 146. 1810 = Pleurosorus r.
Salvinii Hk. 2 Cent. t. 71. 1861 = P. mexicanum.
Sayeri F. Muell. Sec. Cens. **1**. 234. 1889 = P. selliguea var.
scolopendrina Bory, Dup. Voy. Bot. **2**. 257 t. 30 f. 1. 1828.
scolopendrioides Gaud. Freyc. Voy. Bot. 310. 1827 = ? P. scolopendrinum.
seminuda Willd. sp. **5**. 140. 1810 = Monogramma s.
serrulata Sw. Schrad. Journ. **1800**^{2}. 18. 1801.
serrulata Thwait. Enum. pl. Zeyl. 382. 1864 = Coniogramme fraxinea.
sessilifolia J. Sm. Hist. Fil. 181. 1875 = P. subevenosum var.
setigera J. Sm. Hist. Fil. 181. 1875 = P. Hookeri.
setosa Bl. Enum. 116. 1828 = P. diplosorum var.
setosa Pr. Tent. 208. 1836 = P. micropteris.
severa Pr. Tent. 209. 1836 = Dryopteris polypodioides.

Sprucei J. Sm. Hist. Fil. 181. 1875.
squamulosa Splitgb. Tijds. Nat. Gesch. **7**. 398. 1840 = P. lanceolatum*.
subpinnatifida Bl. Fl. Javae, Fil. 118 t. 49 f. 2. 1828.
tenella Klf. Enum. 84. 1824 = P. pseudogrammitis.
totta Pr. Tent. 209 t. 9 f. 4. 1836 = Dryopteris africana.
triloba Desv. Prod. 222. 1827 = P. synammia.
?vestita Wall. List n. 12. 1828 = Gymnopteris v.
villosa Pr. Tent. 209. 1836 = Dryopteris dasyphylla.
vulcanica Bl. Enum. 118. 1828 = P. Feei.
Wightiana Wall. List n. 2161. 1829 — India austr.
Wittigiana Fée, Cr. vasc. Br. **2**. 50 t. 95 f. 1. 1872-73.
zeylanica Fée, Gen. 234. 1850-52.

Granulina Bory; Fée, Gen. Fil. 48. 1850-52 (*Egenolfia §*) = Polybotrya.

GUERINIA J. Smith, Hist. Fil. 272. 1875 = **Schizoloma.**
articulata J. Sm. Hist. Fil. 272. 1875 = S. Guerinianum.

GYMNIA Hamilton msc. (Don, Prod. Fl. Nepal. 16. 1825) = **Cheilanthes.**

GYMNOCARPIUM Newman, The Phytologist **4**. 371. 1851 = **Dryopteris.**
Dryopteris Newm. Phytol. **4**. app. XXIV. 1851.
Phegopteris Newm. Phytol. **4**. app. XXIII. 1851.
Robertianum Newm. Phytol. **4**. app. XXIV. 1851.

Gymnoglossa Christ, Mon. Elaph. 22. 1899 (*Elaphoglossum §*) = Elaphoglossum.

GYMNOGRAMMA Desvaux, Berl. Mag. **5**. 304. 1811 — (Polypodiaceae Gen. **60**).
abyssinica Bak. Syn. 517. 1874 = Polypodium loxogramme.
?acrostichoides Desv. Berl. Mag. **5**. 304. 1811 = Leptochilus.
acrostichoides Pr. Tent. 219. 1836 = Cryptogramma.
acuminatum Klf. Enum. 70. 1824 = Gymnopteris rufa.
acuminata Bak. JoB. **1888**. 326 = Polypodium interruptum.
adiantoides Karst.; Mett. Ann. sc. nat. V. **2**. 212. 1864 = Ceropteris tartarea [var.
affinis Pr. Tent. 218. 1836 = Coniogramme fraxinea.
alismaefolia Hk. sp. **5**. 154. 1864 = Syngramma.
alpina Potts, Tr. N. Zeal. Inst. **10**. 361. 1879 = Pleurosorus rutifolius var.
Alstoni ht. Birkenh.; Gard. Chr. III. **19**. 652. 1896 — Hort.
amaurophylla Christ, Bull. Boiss. II. **4**. 1097. 1904 = Paesia anfractuosa.
ambigua Hk. sp. **5**. 150. 1864 = Aspidium.
ampla Bak. Ann. Bot. **5**. 486. 1891 = Polypodium ellipticum.
Andersoni Bedd. Ferns br. Ind. t. 190. 1866 = Woodsia lanosa.
anfractuosa Christ, Bull. Soc. bot. Belg. **35**. 235. 1896 = Paesia.
angustifrons Bak. Syn. 380. 1868 = G. elongata.
appendiculata Bl. Enum. addend. 1828; Fl Javae Fil. 92 t. 39. 1828 = Dryopteris.
argentea (Willd.) Mett.; Kuhn, Fil. Afr. 59. 1868. HB. ed. II. 385. Chr. 69. NPfl. 260 — Africa austr. Ins. Mascar. — Hemionitis Willd. sp. **5**. 132. 1810; Ceropteris Kuhn 1879 — Gymnogramma rosea Desv. 1811. HB. ed. I. 385; Anogramme Fée 1850-52 — Gymnogramme conspersa Kze. 1844; Anogramme Fée 1850-52 — Hemionitis aurea Willd. 1810: Gymnogramma Desv. 1811.
arifolia Kuhn, Ann. Lugd. Bat. **4**. 283. 1869 = Hemionitis.
Ascensionis Hk. sp. **5**. 137. 1864 = Anogramma.
aspidioides Klf. Enum. 81. 1824 = Dryopteris.
aspidioides Desv. Prod. 214. 1827 = Dryopteris polypodioides.
aspidioides Bl. Enum. 112. 1828 = Dryopteris africana.

aspidioides Hk. Ic. pl. t. 950. 1854 = Dryopteris stegnogramme.
Asplenioides Sw. Vet. Akad. Handl. **1817**. 56 t. 3 f. 4 = Dryopteris aspidioides.
attenuata Fée, Cr. vasc. Br. **1**. 59 t. 14 f. 2. 1869 = Diplazium striatum var.
aurea Desv. Berl. Mag. **5**. 305. 1811 = G. argentea*.
aureo-nitens Hk. Ic. pl. t. 820. 1852. HB. 382. Chr. 74. NPfl. 259 — Peru.
Psilogramme Kuhn 1882. Eriosorus scandens Fée 1850-52; Gymnogramme Mett. 1856.
auriculatum Klf. Enum. 79. 1824 = Polybotrya appendiculata.
aurita Hk. Ic. pl. t. 974. 1854 = Dryopteris.
aurita var. Levingii Clarke, Tr. Linn. Soc. II. Bot. **1**. 568. 1880 = Dryopteris [Levingii.
avenia Bak. Syn. 388. 1868 = Polypodium Blumeanum.
axillaris Ces. Atti Acad. Napoli 7^8. 29. 1876 = Leptochilus.
Baileyi Bak. Kew Bull. **1892**. 86 = Polypodium.
Biardii (Fée) Bak. Syn. 516. 1874 — Brasilia.
Anogramme Fée, Cr. vasc. Br. **1**. 241 t. 77 f. 1. 1869; Psilogramme Kuhn 1882; Gymnogramme extensa Bak. 1870.
bidentata Pr. Rel. Haenk. **1**. 18 t. 2 f. 3. 1825 = Ceropteris calomelanos.
bifurcatum Klf. Wes. d. Farrnkr. 81. 1827 = Elaphoglossum furcatum.
Billardierii Klf. Wes. d. Farrnkr. 81. 1827 = Polypodium grammitidis.
Blumei Franch. et Sav. Enum. pl. Jap. 248. 1876 = Polypodium Blumeanum.
bogotensis Hier. Engl. Jahrb. **34**. 478. 1904 = Jamesonia rotundifolia var.
?**Bommeri** Christ, Bull. Soc. Bot. Belg. **35**. 237. 1896 — Costa Rica.
(Gymnopteris?; Ceropteris?).
Borneensis Hk. sp. **5**. 154 t. 301. 1864 = Syngramma.
Boucheana A.Br. App. Ind. sem. ht. Berol. 18. 1854 = Ceropteris tartarea var. [hort.
Brackenridgei Carr. in Seem. Fl. Vit. 370. 1873 = Ceropteris tartarea.
Brownii Kuhn, Bot. Zeit. **1869**. 458 = Notholaena.
Brunoniana Pr. Tent. 219. 1836 = Cryptogramma.
Caeciliae Mett. Fil. Lips. 40. 1856 = Syngramma alismifolia.
calomelanos Klf. Enum. 76. 1824 } = Ceropteris calomelanos.
calomelas Link, Hort. Berol. **2**. 52. 1833 } = Ceropteris calomelanos.
campyloneuroides Bak. Journ. Linn. Soc. **24**. 261. 1887 = Polypodium Hosei.
candida Mett. Cheil. 6. 1859 = Pellaea nivea var.
canescens Bl. Enum. addend. 1828; Fl. Javae Fil. 93 t. 40. 1828 = Dryopteris.
canescens Kl. Linn. **20**. 407. 1847 = Jamesonia.
cantoniense Bak. in Hk. Ic. plant. t. 1685. 1887 = Polypodium.
capensis Spr.; Klf. Linn. **6**. 183. 1831 = Ceterach cordatum.
caracasana Bak. Syn. 516. 1874 (non Kl. msc.) — Venezuela. Columbia.
G. glandulosa Karst. Fl. Col. **1**. 196 t. 97. 1861. NPfl. 259 (non infra); Psilogramme Kuhn 1882.
cartilagidens Bak. Syn. 386. 1868 = Syngramma.
caudata Pr. Tent. 218. 1836 = Coniogramme fraxinea.
caudiformis Hk. sp. **5**. 158. 1864 = Polypodium Feei var.
ceratolepis var. Atirrensis Donn, Smith, Bot. Gaz. **23**. 251. 1897 = Diplazium.
Ceterach Spr. Syst. **4**. 38. 1827 = Ceterach officinarum.
chaerophylla Desv. Berl. Mag. **5**. 305. 1811 = Anogramma.
cheilanthoides Sw. Vet. Akad. Handl. **1817**. 59. HB. 382. NPfl. 259 — Tristan d'Acunha. — Grammitis Sw. Syn. 23, 219, 419. 1806 (patria erron.); Psilogramme Kuhn 1882. Asplenium filipendulaefolium Thouars 1804 (?1809); Gymnogramma Desv. 1811.

cheilanthoides Mett. Fil. Lechl. 1. 10. 1856 = G. elongata.
cheilosora Fée, 8 mém. 79. 1857 = Dryopteris diplazioides.
chilensis Brack. Expl. Exp. **16**. 22. 1854 = Pleurosorus papaverifolius.
chrysoconia Moore, Ind. 7. 1857 = Acrostichum.
chrysophyllum Klf. Enum. 74. 1824 = Ceropteris calomelanos var.
?**chrysosora** Bak. Journ. Linn. Soc. **24**. 260. 1887 — Borneo.
(§ Cerosora Bak.) Gen. dub.
ciliata Hier. Engl. Jahrb. **34**. 477. 1904 = Jamesonia.
cinnamomea Kl. Linn. **20**. 407. 1847 = Jamesonia.
Cominsii Bak. Journ. Linn. Soc. **19**. 296. 1882 = Diplazium.
congesta Christ, Bull. Boiss. II. **4**. 1098. 1904 — Costa Rica.
consanguinea A.Br. App. Ind. Sem. ht. Berol. 18. 1854 — Hort.
consimilis Fée; (Bak. Syn. 377. 1868) Jenm. Bull. Dep. Jam. **4**. 203. 1897 = Dryopteris heteroclita var.
conspersa Kze. Linn. **18**. 116. 1844 = G. argentea.
cordata Schlecht. Adumbr. 16. 1825 = Ceterach.
cruciata Moore, Ind. 8. 1857 = Acrostichum.
cyclophylla Bak. Tr. Linn. Soc. II. Bot. **2**. 293 t. 53. 1887 = Pterozonium.
Dayi Bedd. JoB. **1888**. 5 t. 279 B. = Syngramma.
dealbata Pr. Rel. Haenk. **1**. 18 t. 3 f. 1. 1825 = Ceropteris tartarea.
dealbata Nutt.; Mett. Cheil. 6. 1859 = Pellaea.
decipiens Mett. Ann. sc. nat. IV. **15**. 60. 1861 = Aspleniopsis.
decomposita Bak. Gard. Chr. **1872**. 1587. HB. 517 — Hort. (Andes?).
decurrens Hk. sp. **5**. 161. 1864 = Polypodium ellipticum.
decurrenti-alata Hk. sp. **5**. 142 t. 294. 1864 = Dryopteris.
decussata Fée, 7 mém. 44. 1857 = Dryopteris.
Delavayi Bak. Ann. Bot. **5**. 484. 1891 = Gymnopteris.
digitata Bak. JoB. **1890**. 267 = Polypodium.
diplazioides Desv. Prod. 214. 1827 = Dryopteris.
distans Link, Hort. Berol. **2**. 53. 1833 = Ceropteris calomelanos.
Dobraydense ht. Birkenhead; Gard. Chr. III. **23**. 332. 1898 — Hort.
domingensis Bak. Ann. Bot. **5**. 485. 1891 — San Domingo.
Psilogramme Und. 1902.
dubia (Kuhn) Bak. Ann. Bot. **5**. 485. 1891 — Brasilia.
Psilogramme Kuhn, Chaetopt. 337. 1882.
edulis Ces. Atti Ac. Napoli 7^8. 28. 1876 = Diplazium esculentum.
Eggersii Christ, Bull. Soc. Bot. Belg. $\mathbf{33^2}$. 92. 1894 = Anogramma.
Ehrenbergiana Kl. Linn. **20**. 411. 1847 = Gymnopteris.
elaphoglossoides Bak. Tr. Linn. Soc. II. Bot **2**. 293 t. 54. 1887 = Syngramma.
elegantissima Bull, Cat. **1889**. 6, 8 c. fig. — Hort.
elliptica Bak. Syn. 389. 1868 = Polypodium.
elongata Hk. et Grev. JoB. **1**. 61 t. 119. 1834. NPfl. 259 — Columbia-Peru.
Jamesonia J. Sm. 1875; Psilogramme Kuhn1882; Gymnogramme angustifrons Bak. 1868. HB. 380. Chr. 73.
elongata Hk. sp. **5**. 157. 1864 = Polypodium lanceolatum*.
expansa Fée, Cr. vasc. Br. **1**. 60 t. 14 f. 4. 1869 = Diplazium ambiguum.
extensa Bak. Fl. bras. $\mathbf{1^2}$. 599. 1870 = G. Biardii.
falcata J. Sm. JoB. **4**. 51. 1841 = Coniogramme fraxinea.
farinifera Lind. et Rod. L'Ill. Hort. **33**. 113 t. 604. 1886 — Hort.
Fauriei Christ, Bull. Boiss. II. **2**. 561. 1902 = Anogramma.

Feei Hk. sp. **5**. 158. 1864 = Polypodium.
Fendleri Mett. Cheil. 7. 1859 = Pellaea.
ferruginea Kze. Linn. **9**. 34. 1834 = Gymnopteris.
filipendulaefolia Desv. Berl. Mag. **5**. 304. 1811 = G. cheilanthoides.
Finlaysoniana Bak. Ann. Bot. **5**. 486. 1891 = Polypodium digitatum.
flabellata Hk. et Grev. JoB. **1**. 61 t. 120. 1834. HB. 382. NPfl. 259 — Peru. Anogramme Fée 1850-52; Psilogramme Kuhn 1882.
flavens Klf. Enum. 77. 1824 = Pellaea.
flexicaulis Kl.; Salom. Nom. 184. 1883 — Mexico. — (an seq.?).
flexilis Kl. Linn. **20**. 414. 1847 — Columbia. — (Ceropteris sp.).
flexuosa (H.B.) Desv. Berl. Mag. **5**. 306. 1811. HB. 384. Chr. 72 — Costa Rica-Peru. — Grammitis Humb. et Bonpl. (? Pl. aequin. **2**. 167 t. 138. 1809); HBK. Nov. Gen. et. Sp. **1**. 5. 1815; Psilogramme Kuhn 1882. Gymnogramme refracta Kze.; Kl. 1847; Anogramme Fée 1850-52. Cryptogramma retrofracta Grev. et Hk. 1833; Gymnogramme Kze.; Mett. 1856 — G. Ruiziana Kl. 1847 — G. recurvata Karst. 1869.
fraxinea Bedd. Ferns br. Ind. Suppl. 24. 1876 = Coniogramme.
Gaerdtiana Koch. Wochenschr. Gärtn. Pflzk. **1858**. 192 — Hort. (Ceropteris hybr.).
Gardneri Bak. Syn. 515. 1874 = Gymnopteris.
gigantea Bak. JoB. **1889**. 177 = Diplazium maximum var.
?**glandulosa** (Sw.) Christ in Schwacke, Pl. Nov. Mineiras **2**. 18. 1900; Bull. Boiss. II. **2**. 364. 1902 — Brasilia. — Cheilanthes Sw. Vet. Ak. Handl. **1817**. 77. HB. 475; Pteris Mett. 1861; Paesia Kuhn 1882. — (an potius Paesia sp.?).
glandulosa Karst. Fl. Col. **1**. 196 t. 97. 1861 = G. caracasana.
glutinosa Mett. Ann. sc. nat. V. **2**. 209. 1864 = Jamesonia.
Goudotii Hier. Engl. Jahrb. **34**. 476. 1904 = Jamesonia.
gracilis Hew. Mag. Nat. Hist. II. **2**. 457. 1838 = Dryopteris heteroclita.
grammitoides Bak. JoB. **1889**. 178 = Polypodium.
grandis Bak. Syn. 377. 1868 = Diplazium striatum var.
grandis Racib. Pterid. Buit. 72. 1898 = Polypodium Raciborskii.
Griffithii Hance, JoB. **1887**. 14 = Aspidium.
guianensis Kl. Linn. **20**. 413. 1847 = Ceropteris tartarea.
haematodes Christ, Bull. Boiss. II. **4**. 1097. 1904 — Costa Rica.
Hamiltoniana Hk. sp. **5**. 161. 1864 = Polypodium pedunculatum.
Haughtoni Hk.; Bak. Syn. 381. 1868 = Ceterach cordatum var.
helenensis Fée, 8 mém. 79. 1857 = Dryopteris punctata.
helveola Fée, 8 mém. 79. 1857 — Columbia. — (Dryopteris sp.).
Henryi Bak. JoB. **1887**. 171 = Polypodium.
heterocarpa Hk. sp. **5**. 160. 1864 = Polypodium.
heterophlebia Gilbert, Bull. Torr. Cl. **23**. 449. 1896 = Polypodium persicariifolium.
Heyderi Lauche, Monatsschr. Ver. Beförd. Gartenb. Preuss. St. **1877**. 421 t. 4; Gartenfl. **27**. 216. 1878 — Hort.
hirsutula Mett. Ann. sc. nat. V. **2**. 209. 1864. HB. 380. NPfl. 259 — Columbia. Psilogramme Kuhn 1882.
hirta (HBK.) Klf. Enum. 72. 1824. HB. 384. Chr. 72 — Columbia. Costa Rica. Grammitis HBK. Nov. Gen. et Sp. **1**. 4. 1815; Psilogramme Kuhn 1882. Gymnogramme laserpitiifolia Kze. 1845; Anogramme Fée 1850-52 — G. petroselinifolia Kl. 1847; Anogramme Fée 1850-52.

hispida Mett.; Kuhn, Linn. **36**. 72. 1869 = Gymnopteris.

hispidula (Kze.) Kl. Linn. **20**. 407. 1847 — Columbia. — Jamesonia Kze. Bot. Zeit. **1844**. 739; Anogramme? Fée 1850-52; Psilogramme Kuhn 1882.

Hookeri J. Sm.; Bak. Syn. 381. 1868 — Columbia. Bolivia. (an potius Ceropteris sp.?).

hybrida Martens, Bull. sc. Brux. **4**. 50. 1837 = Ceropteris calomelanos.

imbricata Kl. Linn. **20**. 407. 1847 = Jamesonia.

immersa Moore, Ind. 232. 1861 = Pleurosorus papaverifolius.

incisa Mart. et Linden; Kze. Farnkr. **2**. 78 t. 132. 1851. HB. 380 — Columbia. Psilogramme Kuhn 1882.

insignis Mett.; Kuhn, Linn. **36**. 70. 1869. HB. 516. NPfl. 259 — Brasilia. Psilogramme Kuhn 1882.

involuta Hk. sp. **5**. 155. 1864 = Polypodium scolopendrinum.

iridifolia Christ, Verh. Nat. Ges. Basel **11**. 248. 1895 = Polypodium.

Jamesoni Bak. Syn. 516. 1874. Chr. 71 — Ecuador. Columbia.

japonica Desv. Berl. Mag. **5**. 305. 1811 = Coniogramme.

javanica Bl. Enum. 112. 1828 = Coniogramme fraxinea.

Karstenii Mett. Ann. sc. nat. V. **2**. 210. 1864. HB. 380. NPfl. 259 — Columbia. Psilogramme Kuhn 1882.

Kunzei Moritz; Ettingsh. Farnkr. 56 t. 31 f. 5. 1865 = G. flexuosa.

lanata Kl.; A.Br. App. Ind. Sem. ht. Berol. 17. 1854 = Gymnopteris ferruginea [var.

lanceolata Hk. sp. **5**. 156. 1864 = Polypodium loxogramme.

lanceolata Luerss. Fil. Graeff. 113. 1871 = Syngramma Hookeri.

lanuginosa A.Br. Ind. Sem. ht. Berol. 17. 1854 = Notholaena vellea.

laserpitiifolia Kze. Bot. Zeit. **1845**. 285 = G. hirta.

Lathamiae Moore, Gard. Chr. n. s. **22**. 360. 1884 — Hort.

Laucheana Koch, Allgm. Gartenzeit **1858**. 53 — Hort. (Ceropteris hybr.).

laxa Mett.; Kuhn, Linn. **36**. 69. 1869 = Jamesonia.

Lechleri Mett.; Kuhn, Linn. **36**. 71. 1869. HB. 516 — Peru. Psilogramme Kuhn 1882.

Lehmannii Hier. Engl. Jahrb. **34**. 474. 1904 — Columbia. (§ Isgnogramme Hier.).

leptophylla Desv. Berl. Mag. **5**. 305. 1811 = Anogramma.

Levingei Bak. Ann. Bot. **5**. 483. 1891 = Dryopteris.

L'Herminieri Bory; Kze. Linn. **23**. 255. 310. 1850 = Ceropteris calomelanos.

Lindigii Mett. Ann. sc. nat. V. **2**. 210. 1864. HB. 381. Chr. 73. NPfl. 259 — Columbia. — Psilogramme Kuhn 1882.

Linkiana Kze. Linn. **18**. 310. 1844 = Dryopteris diplazioides.

Lobbiana Hk. sp. **5**. 153 t. 300. 1864 = Syngramma.

longifolia Bak. Ann. Bot. **5**. 484. 1891 — Brasilia.

longipes Bak. JoB. **1878**. 301 = Trismeria.

longipetiolata Hier. Engl. Jahrb. **34**. 479. 1904 — Columbia. (an potius Jamesonia sp.?).

Lorentzii Hier. Engl. Jahrb. **22**. 397. 1896 = Anogramma.

Lowei Hk. et Grev. Ic. Fil. t. 89. 1829 = Dryopteris africana.

macrophylla Hk. sp. **5**. 159. 1864 = Polypodium.

macrotis Kze. Bot. Zeit. **1848**. 114 = Dryopteris lineata.

Maingayi Bak. Syn. 517. 1874 = Polypodium.

Makinoi Maxim.; Mak. Bot. Mag. Tokyo **8**. 481 c. tab. 1894 = Anogramma.

Marantae Mett. Fil. Lips. 43. 1856 = Notholaena.

marginata Mett. Ann. sc. nat. IV. **15**. 59. 1861 = Syngramma.
maritima Bory; Salom Nom. 185. 1883 — Columbia.
Martensii Bory; Kze. Linn. **23**. 255. 1850 } = Ceropteris calomelanos varr. hort.
Massoni Loud.; Kze. Linn. **23**. 255. 1850 }
Mathewsii Hk. sp. **5**. 128 t. 290. 1864. HB. 382 — Peru.
Psilogramme Kuhn 1882.
Mayeriana A.Br. Allgm. Gartenzeit. **1857**. 279 — Hort. (Hybr.)
membranacea Hk. sp. **5**. 159. 1864 = Polypodium selliguea.
membranacea Bail. Syn. Queensl. Fl. Suppl. III. 94. 1890 = Polypodium Baileyi.
mexicana Bak. Syn. 387. 1868 = Polypodium.
microcarpon Fée, 7 mém. 43 t. 20 f. 5. 1857 = Dryopteris decussata.
microphylla Hk. Ic. pl. t. 916. 1854 = Anogramma.
microphylloides Ces. Rend. Ac. Napoli **16**. 27, 30. 1877 = Cheilanthes tenuifolia.
mohriaeformis Kze. Farnkr. **2**. 79. 1851 (nomen); Mett. Fil. Lechl. **1**. 9. 1856 [= G. rufescens.
mollis Fée, 8 mém. 80. 1857 = Dryopteris sp. (Hort.).
mollissima Fisch.; Kze. Linn. **23**. 255. 310. 1850 = Dryopteris africana.
Muelleri Hk. sp. **5**. 143 t. 295. 1864 = Gymnopteris.
myriophylla Sw. Vet. Ak. Handl. **1817**. 58. HB. 383. Chr. 72. NPfl. 259 — Brasilia. — Psilogramme Kuhn 1882. Cheilanthes glandulifera Fée 1872-73 (non Liebm. 1849).
namaquensis Pappe et Raws. Syn. Fil. Afr. austr. 42. 1858 = Ceterach cornivea [datum.
nivea Mett. Cheil. 7. 1859 = Pellaea.
Novae-Zealandiae Col. Tasm. Journ. **2**. 165. 1844 = Anogramma leptophylla.
obtusa Moore, Ind. 231. 1861 = Ceropteris.
obtusata Bl. Enum. 113. 1828 = Dryopteris opaca.
obtusifolia Hk. sp. **5**. 153 t. 299. 1864 = Syngramma.
ochracea Pr. Rel. Haenk. **1**. 17. 1825 = Ceropteris tartarea var.
opaca Spr. Syst. **4**. 39. 1827 = Dryopteris.
opaca Kze. Linn. **24**. 268. 1851 = Diplazium polypodioides.
oppositans Fée, Cr. vasc. Br. **1**. 58 t. 14 f. 1. 1869 = Dryopteris diplazioides.
Orbignyana Mett.; Kuhn, Linn. **36**. 70. 1869. HB. 517 — Bolivia.
Psilogramme Kuhn 1882.
Ornithopteris Kl. Linn. **20**. 413. 1847 = Ceropteris tartarea var.
Ottonis Kl. Linn. **20**. 408. 1847. HB. 381. NPfl. 259 — Venezuela.
Anogramme Fée 1850-52; Psilogramme Kuhn 1882.
pallescens Moore, Ind. 231. 1861 = Ceropteris tartarea.
palmata Link, Hort. Berol. **2**. 48. 1833 = Hemionitis.
palmata Bak. Syn. 389. 1868 = Polypodium christovalense.
papaverifolia Kze. Anal. 12 t. 8 f. 2. 1837 = Pleurosorus.
patula Fée, Cr. vasc. Br. **1**. 59 t. 14 f. 3. 1869 = Dryopteris pteroidea.
Pearcei Moore, Gard. Chr. **1864**. 340. HB. 384 — Peru. Chile.
pedatum Klf. Enum. 69. 1824 = Gymnopteris.
pentaphylla Bak. Kew Bull. **1898**. 233 = Polypodium ellipticum var.
peruviana Desv. Berl. Mag. **5**. 329. 1811 = Ceropteris.
petroselinifolia Kl. Linn. **20**. 409. 1847 = G. hirta.
Philippinensis Fée, Gen. 181. 1850-52 = Aspidium ambiguum.
pilosa Mart. et Gal. Mém. Ac. Brux. **15**. 27 t. 4 f. 1. 1842 = Dryopteris.
pilosa Brack. Expl. Exp. **16**. 22 t. 4 f. 1. 1854 = Coniogramme fraxinea.
pinnata Hk. sp. **5**. 151. 1864 = Syngramma.
pinnatifida Salom. Nom. 186. 1883 = Hemionitis.

plantaginea Spr. Syst. **4**. 39. 1827 = Hemionitis (Don).
plicata Moore, Ind. 231. 1861 = Ceropteris peruviana.
podophylla Hk. sp. **5**. 152 t. 296. 1864 = Gymnopteris Ehrenbergiana.
polypodioides Spr. Syst. **4**. 40. 1827 = Dryopteris.
polypodioides Link, Hort. Berol. **2**. 50. 1833 = Dryopteris diplazioides var.
pothifolia Spr. Syst. **4**. 39. 1827 = Polypodium ellipticum.
Pozoi Desv. Prod. 216. 1827 = Pleurosorus.
prehensibilis Bak. Syn. 517. 1874; Ic. pl. t. 1683. Chr. 72. NPfl. 259 — Columbia-Ecuador.
procera Moore, Ind. LXII. 1857 = Coniogramme fraxinea.
procurrens Fée, 8 mém. 78. 1857 = Dryopteris pilosa var.
prolifera Fée, Gen. 182. 1850-52 } = Dryopteris diplazioides v. Linkiana.
pteroides Fée, Gen. 182. 1850-52 } = Dryopteris diplazioides v. Linkiana.
pteridiformis Ces. Rend. Ac. Napoli **16**. 27. 30. 1877 = Notholaena.
?**pulchella** Linden; Moore, Gard. Chr. **1856**. 597 c. fig.; Hk. Fil. exot. t. 74. 1859. HB. 385 — Venezuela. — prob. Ceropteris sp.
pumila Spr. Tent. Suppl. Syst. **4**. 31. 1828 = Hecistopteris.
quinata Hk. sp. **5**. 152 t. 297. 1864 = Syngramma.
Raddiana Link, Hort. Berol. **2**. 50. 1833 = Dryopteris polypodioides.
recurvata Karst. Fl. Col. **2**. 165 t. 187 f. 1. 1869 = G. flexuosa.
refracta Kze.; Kl. Linn. **20**. 410. 1847 = G. flexuosa.
Regnelliana Christ in Schwacke, Pl. Nov. Mineiras **2**. 18. 1900; Bull. Boiss. II. **2**. 365. 1902 = Cheilanthes.
Regnelliana Lindm. Ark. för Bot. **1**. 240 t. 8 f. 6. 1903 = Anogramma Lorentzii var.
regularis Bak. Syn. 388. 1868 = Polypodium.
remota Moore; Salom. Nom. 187. 1883 — India occ.
reniformis Mart. Ic. Cr. Bras. 88 t. 26. 1834 = Pterozonium.
retrofracta Kze.; Mett. Fil. Lechl. **1**. 10. 1856 = G. flexuosa.
rhizophylla Klf. Enum. 78. 1824 = Polybotrya appendiculata.
robusta Hier. Engl. Jahrb. **34**. 476. 1904 = Jamesonia
rosea Desv. Berl. Mag. **5**. 306. 1811 = G. argentea.
rotundifolia Mett. Ann. sc. nat. V. **2**. 209. 1864 = Jamesonia.
rufa Desv. Berl. Mag. **5**. 304. 1811 = Gymnopteris.
rufescens Fée, Gen. 181 t. 19 C f. 3. 1850-52. NPfl. 259 — Peru.
Psilogramme Kuhn 1882; Gymnogramme mohriaeformis Kze. 1851 (nomen); Mett. 1856. HB. 380.
Ruiziana Kl. Linn. **20**. 410. 1847 = G. flexuosa.
Ruiziana Mett. Fil. Lips. 42. 1856 = Gymnopteris ferruginea.
rupestris Kze. Linn. **23**. 256. 1850 = Dryopteris.
rutaefolia Desv. Prod. 213. 1827 = Pleurosorus.
rutaefolia var. hispanica Hk. Ic. pl. t. 935. 1854 = Pleurosorus Pozoi.
sadlerioides Und. ap. Heller, Minn. bot. St. **1**. 781 t. 43. 1897 = Sadleria polystichoides.
sagittata Ettingsh. Denkschr. Ak. Wien **22**. 102 t. 11 f. 10. 1864 = Hemionitis arifolia.
salicifolia Mak. Phan. et Pterid. Jap. Ic. t. 34. 1899 = Polypodium Makinoi.
Salvinii Hk. sp. **5**. 157. 1864 = Polypodium mexicanum.
Sayeri F. Muell. et Bak. JoB. **1887**. 163 = Polypodium selliguea var.
scalaris Kl. Linn. **20**. 407. 1847 = Jamesonia.
scandens Mett. Fil. Lips. 42. 1856 = G. aureo-nitens.

scandens (Fée) Bak. Journ. Linn. Soc. **14**. 25. 1873. HB. 516. Chr. 73 — Brasilia. — Neurogramme Fée. Cr. vasc. Br. **1**. 263. 1869; **2**. 39 t. 92.
Schaffneri Moore, Ind. 231. 1861 = Ceropteris peruviana.
Schaffneri Bak. Ann. Bot. **5**. 484. 1891 = Gymnopteris hispida.
schizophylla Bak.; Jenm. JoB. **1877**. 266 = Anogramma.
Schomburgkiana Kze.; Kl. Linn. **20**. 408. 1847. HB. 382. NPfl. 259 — Guiana. Anogramme Fée 1850-52; Psilogramme Kuhn 1882.
Schwackeana Christ in Schwacke, Pl. Nov. Mineiras **2**. 18. 1900; Bull. Boiss. II. **2**. 365. 1902 — Brasilia.
?**sciatraphis** Donn. Smith, Bot. Gaz. **19**. 266 t. 26. 1894. Chr. 65 — Costa Rica. (an Asplenium sp. aff. A. ferulaceo?).
scolopendrioides Bak. JoB. **1879**. 299 = Syngramma.
Sellowiana (Pr.) Mett.; Kuhn, Linn. **36**. 69. 1869. HB. 515. NPfl. 259 — Brasilia. — Cheilanthes Pr. 1836 (nomen); Psilogramme Kuhn 1882.
serrata Moore, Ind. 232. 1861 = Ceropteris calomelanos.
serrulata Bl. Enum. 113. 1828 = Coniogramme fraxinea.
serrulata Fée, 8 mém. 79. 1857 — Mexico. — (Dryopteris sp.).
setulosa Hier. Engl. Jahrb. **34**. 479. 1904 — Columbia.
sinuata Pr. Tent. 219. 1836 = Notholaena.
?**sinuata** Moore; Bak. Ann. Bot. **5**. 483. 1891 — Ecuador. — (Gen. dub.)
Sprengeriana ht.; Wiener Ill. Gartenzeit **1897**. 257 — Hort.
spuria Bak. Syn. 388. 1868 = Polypodium.
Stegnogramme Bl. Fl. Javae Fil. 98 t. 44. 1828 = Dryopteris.
Stelzneri Koch, Wochenschr. Gärtn. Pflzk. **1859**. 183 = Ceropteris tartarea (hybr.).
Stierii Rosenst. Festschr. Alb. v. Bamberg 64. 1905 = Dryopteris.
subcordata Eat. et Dav. Contr. U. S. Nat. Herb. **5**. 138 t. 16. 1897 = Gymnopteris.
subdigitata Keys Pol. Cyath. Hb. Bung. 31. 1873 = Monachosorum.
subglandulosa Hk. et Grev. Ic. Fil. t. 9. 1827 = Pleurosorus rutifolius.
subscandens Sod. Cr. vasc. quit. 401. 1893 — Ecuador.
subsimilis Hk. sp. **5**. 142 t. 293. 1864 = Dryopteris.
subtrifoliata Hk. sp. **5**. 152 t. 298. 1864 = Stenochlaena sorbifolia var. 17.
sulphurea (Sw.) Desv. Berl. Mag. **5**. 305. 1811. HB. 385. NPfl. 260 — Ind. occ. Acrostichum Sw. Prod. 129. 1788; Ceropteris Fée 1850-52.
tartarea Desv. Berl. Mag. **5**. 305. 1811 = Ceropteris.
tenera Mett. Cheil. 7. 1859 = Pellaea.
Thiebautii Desv. Prod. 215. 1827 = G. argentea.
tolimensis Hier. Engl. Jahrb. **34**. 475. 1904 = Jamesonia.
tomentosa Desv. Berl. Mag. **5**. 304. 1811 = Gymnopteris.
tortuosa Sod. Cr. vasc. quit. 401. 1893 — Ecuador.
totta Schlecht. Adumbr. 15 t. 6. 1825 = Dryopteris africana.
totta Mart. et Gal. Mém. Ac. Brux. **15**. 26. 1842 = Dryopteris diplazioides.
triangulare Klf. Enum. 73. 1824 = Ceropteris.
triangularis var. viscosa Eat. Ferns N. Am. **2**. 16 t. 48 f. 5. 1881 = Ceropteris viscosa.
triangulata Jenm. Bull. Dep. Jam. **4**. 206. 1897 = Ceropteris.
trichomanoides J. Sm. Hist. Fil. 159. 1875 — Ceylon. — Pleurosorus sp.?
trifoliata Desv. Berl. Mag. **5**. 305. 1811 = Trismeria.
unita Kze. Linn. **18**. 115. 1844 = Dryopteris silvatica.
valleculata Bak. JoB. **1888**. 325 = Syngramma.
Veitchii ht.; Gard. Chr. III. **16**. 446. 1894 — Hort.

vellea Kuhn, Fil. Afr. 61. 1868 = Notholaena.
vellea Bak. JoB. **1881**. 206; Ic. pl. t. 1681 — Columbia.
verticalis Kl. Linn. **20**. 410. 1847 = Jamesonia.
vestita Pr. Tent. 218. 1836 = Gymnopteris.
villosa Link, Hort. Berol. **2**. 51. 1833 = Dryopteris dasyphylla.
villosa Moore, Ind. 78. 1858 = Anogramma chaerophylla.
vittaeformis Mett. Fil. Lips. 40. 1856 = Syngramma.
Wallichii Hk. sp. **5**. 155 t. 302. 1864 = Syngramma alismifolia var.
Warscewiczii Mett. Ann. sc. nat. V. **2**. 211. 1864. HB. 382. NPfl. 259 — Costa Rica. Columbia. — Psilogramme Kuhn 1882.
Wetenhalliana Moore, Gard. Chr. **1860**. 934 = G. pulchella.
Wrightii Hk. sp. **5**. 160 t. 303. 1864 = Polypodium.
xerophila Bak. JoB. **1881**. 206. Chr. 71. NPfl. 260 — Columbia.
Neurogramme Diels 1899. NPfl. 263.
Zollingeri Salom. Nom. 188. 1883 = Syngramma.

GYMNOGRAMMITIS Griffith, Icon. Pl. Asiat. **2**. t. 129 f. 1. 1849 = **Davallia** Clarkei.

GYMNOPTERIS Bernhardi, Schrad. Journ. **1799**[1]. 297 — (Polypodiaceae Gen. **64**).
Ceterach Bernh. Schrad. neu. Journ. **1**[2]. 22. 1806 = Ceterach officinarum.
Delavayi (Bak.) Und. Bull. Torr. Cl. **29**. 627. 1902 — Yunnan.
Gymnogramme Bak. Ann. Bot. **5**. 484. 1891. Chr. 67; Neurogramme Diels 1899. NPfl. 262.
discolor Bernh. Schrad. neu. Journ. **1**[2]. 20. 1806 = Blechnum.
Ehrenbergiana (Kl.) C. Chr. Ind. 341. 1905 — Mexico-Guatemala.
Gymnogramme Kl. Linn. **20**. 411. 1847; Stegnogramme Fourn. 1872; Bommeria Und. 1902. Gymnogramme podophylla Hk. 1864. HB. 390; Hemionitis J. Sm. 1875; Dictyogramme Trev. 1877; Bommeria Fourn. 1880.
ferruginea (Kze.) Und. Bull. Torr. Cl. **29**. 628. 1902 — Guatemala-Peru.
Gymnogramme Kze. Linn. **9**. 34. 1834. HB. 381. Chr. 74; Notholaena Eat. 1860 (non Desv. 1811); Neurogramme Diels 1899. NPfl. 263 — Gymnogramme lanata Kl. 1854.
Gardneri (Bak.) Und. Bull. Torr. Cl. **29**. 627. 1902 — Brasilia.
Gymnogramme Bak. Syn. 515. 1874.
hispida (Mett.) Und. Our nat. Ferns ed. VI. 84. 1900 — Texas-New Mexico. Mexico. — Gymnogramme Mett.; Kuhn, Linn. **36**. 72. 1869. HB. 516 Chr. 68; Neurogramme Diels 1899. NPfl. 264; Bommeria Und. 1902. B. Schaffneri Fourn. 1880; Gymnogramme Bak. 1891.
Muelleri (Hk.) Und. Bull. Torr. Cl. **29**. 627. 1902 — Austr. trop.
Gymnogramme Hk. sp. **5**. 143 t. 295. 1864. HB. 379. Chr. 67; Grammitis F. Muell. 1866; Hemionitis J. Sm. 1875; Neurogramme Diels 1899. NPfl. 262.
pedata (Sw.) C. Chr. Ind. 341. 1905 — Mexico-Guatemala.
Hemionitis Sw. Syn. 20, 209 t. 1 f. 3. 1806; Gymnogramma Klf. 1824. HB. 383. Chr. 67; Neurogramme Link 1841. NPfl. 264; Bommeria Fourn. 1880.
quercifolia Bernh. Schrad. neu. Journ. **1**[2]. 20. 1806 = Leptochilus zeylanicus.
Reynoldsii (F. Muell.) C. Chr. Ind. 341. 1905 — Australia.
Nothochlaena F. Muell. Fragm. **8**. 175. 1874; Grammitis Benth. 1878; Neurogramme Diels 1899. NPfl. 262.
rufa (L.) Bernh. (Schrad. Journ. **1799**[1]. 297): Und. Bull. Torr. Cl. **29**. 627. 1902 — Amer. trop. — Pteris (ruffa) L. sp. **2**. 1074. 1753; Acrostichum L. 1759; Hemionitis Sw. 1801; Gymnogramma Desv. 1811. HB. 379; Neurogramme Link 1841. Chr. 62. NPfl. 262.

serrulata Bernh. Schrad. neu. Journ. 1^2. 48 t. 2 f. 4 a. 1806 = Polypodium.
subcordata (Eat. et Dav.) Und. Bull. Torr. Cl. **29**. 628. 1902 — Mexico.
Gymnogramme Eat. et Dav. Contr. U. S. Nat. Herb. **5**. 138 t. 16. 1897.
tomentosa (Lam.) Und. Bull. Torr. Cl. **29**. 627. 1902 — Brasilia-Peru.
Asplenium Lam. Enc. **2**. 308. 1786; Gymnogramma Desv. 1811. HB. 380; Hemionitis Raddi 1819; Neurogramme Link 1833. Chr. 62. NPfl. 263. Ceterach lobatum Pr. 1822.
triangularis Und. Our nat. Ferns ed. VI. 84. 1900 = Ceropteris.
vestita (Wall.) Und. Bull. Torr. Cl. **29**. 627. 1902 — China. India bor.
Grammitis? Wall. 1828 (nomen); Gymnogramma Pr. 1836 (nomen); Hk. Ic. pl. t. 115. 1837. HB. 379. Chr. 66; Syngramma Moore 1857; Hemionitis J. Sm. 1875; Neurogramme Diels 1899. NPfl. 262.

GYMNOPTERIS Presl, Tent. Pterid. 244. 1836 = **Leptochilus.**
acuminata Pr. Tent. 244 t. 11 f. 6. 1836 = L. nicotianifolius.
aliena Pr. Tent. 244. 1836.
aurita Keys. Pol. Cyath. Hb. Bung. 33. 1873 = Stenosemia a.
axillaris Pr. Tent. 244 t. 11 f. 4-5. 1836.
Bernoullii Kuhn; Diels, Nat. Pfl. 1^4. 201. 1899.
Blumeana Bedd. Ferns br. Ind. Suppl. 27. 1876 = L. lomarioides.
Bonii Christ, Bull. Boiss. II. **4**. 610. 1904.
Cadieri Christ, Journ. de Bot. **19**. 126. 1905.
contaminans Bedd. Ferns br. Ind. Suppl. 27. 1876 = L. virens.
contaminoides Christ. Ann. Cons. Jard. bot. Genève **3**. 32. 1899.
costaricensis Christ, Bull. Boiss. II. **4**. 964. 1904 = L. Lindigii var.
costata Bedd. Ferns br. Ind. Suppl. 27. 1876 = L. scalpturatus.
decurrens Moore, Ind. XXIX. 1857.
decurrens Hk. JoB. **9**. 359. 1857 = L. Harlandii.
dentata Fée, Acrost. 85. 1845 = L. alienus.
Feei Moore, Ind. XXIX. 1857 = L. lanceolatus.
flagellifera Bedd. Ferns br. Ind. Suppl. 27. 1876 = L. heteroclitus.
Gaboonensis J. Sm. Ferns br. and for. 118. 1866.
Guyanensis Christ. Bull. Boiss. II. **1**. 70. 1901 = L. guianensis.
Harlandii J. Sm. Hist. Fil. 138. 1875.
hastata Fourn. Mex. pl. **1**. 70. 1872 = L. alienus.
Heudelotii Bory; Fée, Acrost. 84 t. 45. 1845.
Hollrungii Kuhn in Schum. Fl. Kais. Wilh. Land. 8. 1889.
inconstans Copeland in Perkins, Fragm. 177. 1905.
irregularis Fourn. Mex. pl. **1**. 70. 1872 = L. alienus.
lanceolata Moore, Ind. 10. 1857 = L. Linnaeanus.
lanceolata Bedd. Ferns br. Ind. Suppl. 26. 1876.
latifolia Pr. Tent. 244. 1836.
Linnaeana Christ, Journ. de Bot. **19**. 125. 1905.
lomarioides Christ, Bull. Boiss. II. **1**. 456. 1901.
metallica Bedd. Ferns br. Ind. Suppl. 26 t. 390. 1876.
minor Hk. 2 Cent. t. 78. 1861.
Naumannii Diels in Schum. et Laut. Fl. deut. Südsee 117. 1901.
neglecta Diels, Nat. Pfl. 1^4. 200. 1899.
nicotianaefolia Pr. Tent. 244. 1836.
normalis J. Sm. JoB. **3**. 403. 1841 = L. lanceolatus var.
nummularifolia Pr. Tent. 244. 1836 = Cyclophorus n.

obtusifolia Pr. Tent. 244. 1836 = ? L. decurrens.
oligarchica J. Sm. Hist. Fil. 138. 1875.
pandurifolia Hk. 2 Cent. t. 87. 1861.
Platyrhynchos J. Sm. JoB. **3**. 403. 1841 = Hymenolepis p.
portoricensis Fée, Acrost. 85. 1845 = L. alienus.
Presliana J. Sm. Hist. Fil. 138. 1875.
punctata Christ, Farnkr. 50. 1897 = L. auriculatus.
repanda Christ, Farnkr. 50. 1897 = L. cuspidatus.
salicina J. Sm. Hist. Fil. 138. 1875.
scandens Christ, Farnkr. 51. 1897 = L. guianensis.
semicordata J. Sm. Hist. Fil. 138. 1875 = L. virens.
semipinnatifida Fée, Acrost. 83 t. 44. 1845 = L. alienus.
serratifolia Diels, Nat. Pfl. **1**⁴. 201. 1899.
spicata Pr. Tent. 244 t. 11 f. 7. 1836 = Hymenolepis s.
subcrenata Bedd. Handb. 437. 1883.
subquinquefida Pr. Epim. 151. 1849 = L. latifolius.
subrepanda J. Sm. JoB. **3**. 403. 1841 = L. Zollingeri.
subsimplex Fée, Acrost. 83 t. 40. 1845 = ? L. Zollingeri.
taccaefolia J. Sm. JoB. **3**. 403. 1841 = L. latifolius var.
terminans Bedd. Ferns br. Ind. Suppl. 27. 1876 = L. subcrenatus.
tricuspis Bedd. Ferns br. Ind. t. 53. 1866.
trilobata J. Sm. JoB. **3**. 403. 1841; Hk. Ic. pl. t. 907. 1854 = L. latifolius var.
variabilis Bedd. Ferns br. Ind. t. 272. 1868 = L. decurrens.
Vespertilio Hk. Lond. JoB. **5**. 193 t. 7-8. 1846 = Cheiropleuria bicuspis.
virens Keys. Pol. Cyath. Hb. Bung. 33. 1873.
Wallichiana Pr. Tent. 244. 1836 = L. decurrens.
Wallii Bedd. Ferns br. Ind. Suppl. 26 t. 389. 1876.

GYMNOSPHAERA Blume, Enum pl. Javae 242. 1828 = **Alsophila.**
aculeata J. Sm. Lond. JoB. **1**. 667. 1842 = A. aspera.
gigantea J. Sm. Lond. JoB. **1**. 667. 1842 = A. glabra.
glabra Bl. Enum. 242. 1828.
squamulata Bl. Enum. 243. 1828.
squamulata J. Sm. JoB. **3**. 419. 1841 et in Hk. Gen. t. 100. 1842 = A. commutata.

GYMNOTHALAMIUM Zenker msc. (Kunze, Linn. **24**. 287. 1851) = **Dryopteris** (scabrosa).

GYMNOTHECA Presl. Suppl. Tent. Pterid. 12. 1845 = **Marattia.**
cicutaefolia Pr. Suppl. 13. 1845.
Douglasii Moore, Ind. CXXI. 1857.
laxa Pr. Suppl. 15. 1845.
Loddigesiana de Vriese, Mon. 11. 1853 — Hort. (America).
Mertensiana Pr. Abh. böhm. Ges. V. **5**. 321. 1848; Epim. 9 t. 1. 1849.
obtusidens Pr. Suppl. 15. 1845 = M. cicutifolia.
podolepis de Vriese, Mon. 10. 1853.
polyodon Pr. Suppl. 14. 1845.
Raddiana Pr. Suppl. 13. 1845 = M. Raddii.
Verschaffeltiana de Vriese, Mon. 10. 1853.
weinmanniaefolia de Vriese. Mon. 11. 1853.

GYROSORIUM Presl. Epim. Bot. 139. 1849 = **Cyclophorus.**
africanum Pr. Epim. 140. 1849.
angustatum Pr. Epim. 141. 1849 — Luzon.
fissum Pr. Epim. 141. 1849 = ? C. acrostichoides.
samarense Pr. Epim. 140. 1849.

HABRODICTYON Van den Bosch, Hym. Jav. 17. 1861 = **Trichomanes.**
(= Abrodictyum Pr.).
Cumingii v.d.B. Hym. Jav. 17 t. 12. 1861 } = T. Smithii.
Smithii v.d.B. Versl. Akad. Wet. Amsterd. **11**. 321. 1861 } = T. Smithii.
HAPLODICTYUM Presl, Epim. bot. 50. 1849; Fée, Gen. 309. 1850-52 = **Dryopteris.**
?exiguum Fée. Gen. 309. 1850-52 = D. philippina vel D. canescens.
heterophyllum Pr. Epim. 51. 1849 = D. canescens*.
Haplophlebia Martius, Icon. Sel. Cr. Bras. 62. 1834 *(Alsophila §)* = Alsophila.
Haplophlebia Gris. Abh. Ges. Wiss. Gött. 7. 286. 1857 *(Aspidium §)* = Oleandra.
HAPLOPTERIS Presl, Tent. Pterid. 141. 1836 = **Vittaria.**
lineata J. Sm. Ferns br. and for. 132. 1866.
scolopendrina Pr. Tent. 141 t. 5 f. 21. 1836.
HECISTOPTERIS J. Smith, Lond. Journ. Bot. **1**. 193. 1842 — (Polypodiaceae Gen. **90**.
pumila (Spr.) J. Sm. Lond. JoB. **1**. 193. 1842. Chr. 55. NPfl. 300 — Amer. trop. Gymnogramme Spr. Tent. Suppl. Syst. 31. 1828. HB. 379; Vittaria Goebel 1897. —* ceratophylla, fimbriata, lineata Christ, Hedw. **44**. 366. 1905.
HELMINTHOSTACHYS Kaulfuss, Flora **1822**[1]. 103; Enum. 28. 1824 — [(Ophioglossaceae Gen. **3**).
crenata Pr. Suppl. 60. 1845 = H. zeylanica.
dulcis Klf. Flora **1822**[1]. 103; Enum. 28 t. 1 f. 1. 1824 = H. zeylanica.
integrifolia Pr. Suppl. 60. 1845 = H. zeylanica.
mexicana Spr. Syst. **4**. 23. 1827 = H. zeylanica.
zeylanica (L.) Hk. Gen. t. 47. 1840. HB. 447. Chr. 365. NPfl. 472 — Asia, Australia trop. N. Caledonia. — Osmunda L. sp. **2**. 1063. 1753; Botrychium Sw. 1801; Ophiala Desv. 1827; Helminthostachys dulcis Klf. 1822. Botryopteris mexicana Pr. 1825 (patria err.); Helminthostachys Spr. 1827.
HEMESTHEUM Newman, Phytol. **4**. app. XXII. 1851 = **Dryopteris.**
fragrans Gandog. Fl. Eur. **27**. 176. 1891 = D. oreopteris.
montanum Newm. Phytol. **4**. app. XXII. 1851 = D. oreopteris.
molle Gandog. Fl. Eur. **27**. 178. 1891 = D. parasitica.
oreopteris Parment. Ann. sc. nat. VIII. **9**. 300. 1899.
Thelypteris Newm. Phytol. **4**. app. XXII. 1851.
Hemianeimia Prantl, Schiz. 86. 1881 *(Aneimia §)* = Aneimia.
HEMICARDION Fée, Gen. Fil. 282. 1850-52 = **Cyclopeltis.**
Cochinchinae Fée, Gen. 282. 1850-52 = C. Presliana.
crenatum Fée, Gen. 283 t. 22 A. f. 1. 1850-52 = C. Presliana.
Cumingianum Fée, Gen. 283 t. 22 A. f. 2. 1850-52 = C. Presliana.
macrosorum Fée, 8 mém. 101. 1857 = C. semicordata.
Nephrolepis Fée, Gen. 282. 1850-52 = C. semicordata.
subhastatum Fée, Gen. 282. 1850-52 (nomen) — Amboyna.
HEMIDICTYUM Presl, Tent. Pterid. 110. 1836.
?Brunonis Pr. Tent. 111 t. 3 f. 25, 26. 1836 = Diplaziopsis javanica.
Ceterach Bedd. Ferns br. Ind. Suppl. 13. 1876 = Ceterach officinarum.
Douglasii Pr. Tent. 111. 1836 = Phyllitis plantaginea.
Drummondii Fourn. Ann. sc. nat. V. **18**. 311. 1873 = ? Diplaziopsis javanica.
Finlaysonianum Moore, Ind. L. 1857 = Asplenium.
Hookerianum Moore, Ind. 130. 1859 = Asplenium Finlaysonianum.
limbatum Pr. Epim. 74. 1849 = Diplazium marginatum.
marginatum Pr. Tent. 111. 1836 = Diplazium.
peruvianum Pr. Epim. 74. 1849 = Diplazium marginatum.
Purdieanum Moore, Ind. L. 1857 = Asplenium.

HEMIGONUM J. Smith »1841«. (Hist. Fil. 204. 1875) = **Polystichum.**

Hemionites J. Smith, Hist. Fil. 318. 1875 *(Asplenium §)* = Asplenium.

Hemionitidastrum Fée, Gen. Fil. 190. 1850-52 *(Asplenium §)* = Asplenium.

HEMIONITIS Linné (Gen. 944); Spec. plant. **2**. 1077. 1753 — (Polypodiaceae Gen. **63**).

acrostichoides Afz.; Sw. Schrad. Journ. **1800**². 17. 1801 = Leptochilus.

alismaefolia Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Syngramma.

argentea Willd. sp. **5**. 132. 1810 = Gymnogramma.

arifolia (Burm.) Moore, Ind. 114. 1859. Chr. 60. NPfl. 262 — Asia trop. Asplenium Burm. Fl. Ind. 231. 1768; Gymnogramme Kuhn 1869. Hemionitis cordata Hk. et Grev. 1828. HB. 398; H. cordifolia Roxb.; Wall. 1828 — H. sagittata Fée 1850-52; Gymnogramme Ettingsh. 1864.

asplenioides Vell. Fl. flum. **11**. t. 93. 1827; Arch. Mus. Nac Rio Jan. **5**. 454. 1881 = Gymnopteris tomentosa.

aurea Willd. sp. **5**. 131. 1810 = Gymnogramma argentea var.

aureo-nitens J. Sm. Hist. Fil. 151. 1875 = Trachypteris pinnata.

Blumeana Pr. Tent. 221. 1836 = Antrophyum latifolium.

Boryana Willd. sp. **5**. 128. 1810 = Antrophyum.

Boryana Bl. Enum. 111. 1828 = Antrophyum latifolium.

brasiliana Desv. Prod. 216. 1827 = Antrophyum.

callaefolia Pr. Tent. 221. 1836 = Antrophyum.

cajenensis Desv. Berl. Mag. **5**. 311. 1811 = Antrophyum cayennense.

chaerophylla Poir. Enc. Suppl. **3**. 38. 1813 = Anogramma.

citrifolia Hk. sp. **5**. 193. 1864 = Anetium.

concava Pr. Tent. 221 t. 9 f. 20. 1836 = Antrophyum coriaceum.

cordata Hk. et Grev. Ic. Fil. t. 64. 1828 } = H. arifolia.
cordifolia Roxb.; Wall. List n. 44. 1828 }

coriacea Don, Prod. Fl. Nepal. 13. 1825 = Antrophyum.

Cumingiana Fée, 10 mém. 22. 1865 = H. arifolia.

dealbata Willd. Enum. 1067. 1809 = Ceropteris tartarea var.

discolor Schkuhr, Kr. Gew. **1**. 7 t. 6. 1806 = Blechnum.

elegans Dav. Gard. and For. **4**. 484 f. 75. 1891. Chr. 60. NPfl. 262 — Mexico.

elongata Cav. Descr. 261. 1802 — Puerto Rico.

elongata Brack. Expl. Exp. **16**. 66 t. 8. 1854 = Syngramma pinnata.

esculenta Retz. Obs. **6**. 38. 1791 = Diplazium.

falcata Willd. sp. **5**. 126. 1810 = Antrophyum zosteraefolium.

falcata De Vriese, Ned. kr. Arch. **1**. 12. 1846 = Antrophyum callifolium.

gigantea Desv. Prod. 216. 1827 = Antrophyum Desvauxii.

grandifolia Sw. Fl. Ind. occ. **3**. 1605. 1806 = Diplazium.

Griffithii Hk.fil. et Thoms.; Hk. sp. **5**. 193. 1864 = Aspidium.

gymnopteroidea Copeland in Perkins. Fragm. 183. 1905 — Ins. Philipp.

hastata R.Br.; Wall. List n. 2170. 1829 = H. arifolia.

hederifolia J. Sm. Bot. Voy. Herald 339. 1854. HB. 398. NPfl. 262 — Mexico. Dictyogramme Trev. 1877.

Hookeriana Pr. Tent. 221. 1836 = Antrophyum immersum.

Hosei Bak. JoB. **1891**. 108 = Syngramma.

humilis Vell. Fl. flum. **11**. t. 94. 1827; Arch. Mus. Nac. Rio Jan. **5**. 454. 1881 = Gymnopteris tomentosa.

immersa Bory; Willd. sp. **5**. 127. 1810 = Antrophyum.

incisa Blanco, Fl. Filip. 829. 1837 = Diplazium esculentum.

?intermedia Fée, Gen. 172. 1850-52 = H. arifolia.
japonica Thbg. Fl. Jap. 333. 1784 = Coniogramme.
lanceolata L. sp. **2**. 1077. 1753 = Antrophyum.
lanceolata Hk. 2 Cent. t. 55. 1861 = Syngramma Hookeri.
latifolia Keys. Pol. Cyath. Hb. Bung. 32. 1873 = Antrophyum.
leptophylla Lag. Gen. et Sp. 33. 1816 = Anogramma.
Lessonii Pr. Tent. 221. 1836 = Antrophyum plantagineum var.
Levyi Fourn. Bull. Soc. Fr **17**. 237. 1870 — Nicaragua.
lineata Sw. Prod. 129. 1788 = Antrophyum.
Muelleri J. Sm. Hist. Fil. 151. 1875 = Gymnopteris.
obtusa Bory; Willd. sp. **5**. 127. 1810 = Antrophyum.
opaca Don, Prod. Fl. Nepal. 13. 1825 = Dryopteris.
palmata L. sp. **2**. 1077. 1753. HB. 398. Chr. 60. NPfl. 262 — Amer. trop.
Gymnogramme Link 1833.
parasitica L. Syst. Nat. ed. X. **2**. 1322. 1759 = Anetium citrifolium.
parvula Pr. Tent. 221. 1836 = Antrophyum.
pedata Sw. Syn. 20, 209 t. 1 f. 3. 1806 = Gymnopteris.
pedatifida Sm. in Rees, Cycl. **17**. —. 1819 = Doryopteris pilosa.
pinnata J. Sm. JoB. **4**. 69. 1841 (nomen) = H. Smithii.
pinnata Hk.fil. Tr. Linn. Soc. **20**. 167. 1847 = Trachypteris.
pinnatifida Bak. Syn. 399. 1868; Kuhn, Bot. Zeit **1869**. 144. NPfl. 262 — Amer. centr. — Dictyogramme Trev. 1877; Gymnogramme Salom. 1883.
plantaginea Sm. Mém. Ac. Turin **5**. 421 t. 9 f. 1. 1793 = Diplazium plantaginifolia.
plantaginea Cav. Descr 260. 1802 = Antrophyum.
plantaginea Don, Prod. Fl. Nepal. 13. 1825 — Nepal.
Gymnogramme Spr. 1827.
podophylla J. Sm. Hist. Fil. 150. 1875 = Gymnopteris Ehrenbergiana.
pothifolia Don. Prod. Fl. Nepal. 13. 1825 = Polypodium ellipticum.
Pozoi Lag. Gen. et Sp. 33. 1816 = Pleurosorus.
prolifera Retz. Obs. **6**. 38. 1791 = Dryopteris.
Reinwardtiana Pr. Tent. 221. 1836 = Antrophyum plantagineum.
reticulata Forst. Prod. 79. 1786 = Antrophyum.
reticulata Bory. Voy. **1**. 214. 1804 = Antrophyum Boryanum.
reticulata Raddi. Opusc. sci. Bol. **3**. 284. 1819 = Antrophyum cayennense.
rigida Cav. Descr. 260. 1802 = ?Antrophyum semicostatum.
rufa Sw. Schrad. Journ. **1800²**. 16. 1801 = Gymnopteris.
sagittata Fée, Gen. 172 t. 14 D. 1850-52 = H. arifolia.
semicostata Pr Tent. 221. 1836 = Antrophyum.
sessilifolia Cav. Descr. 261. 1802 — Ins. Mauban.
Smithii (Trev.) C. Chr. Ind. 346. 1905 — Jamaica. Costa Rica.
Dictyogramme Trev. 1877. Hemionitis pinnata J. Sm. 1841 (nomen. non Hk. fil. 1847); HB. 399. 1868. NPfl. 262.
spatulata Pr. Tent. 221. 1836 = Anetium citrifolium.
tomentosa Raddi. Opusc. sci. Bol. **3**. 284. 1819 = Gymnopteris.
toxotis Trev. Atti Ist. Veneto II. **2**. 168. 1851 = H. arifolia.
trifoliata HBK. Nov. Gen. et Sp. **1**. 4. 1815 = Trismeria.
vestita J. Sm. Hist. Fil. 151. 1875 = Gymnopteris.
vittaeformis Trev. Atti Ist. Veneto V. **3**. 591. 1877 = Syngramma.
Wilfordii Hk. Fil. Exot. t. 93. 1859 = Aspidium Griffithii var.
Zollingeri Kurz. Journ. As. soc. Bengal **30²**. 90 t. 5. 1870 = Syngramma.

HEMIPHLEBIUM Presl, Hymenoph. 25. 1843 = **Trichomanes.**
apodum Prantl, Hym. 46. 1875 = T. hymenoides.
bimarginatum Luerss. Bot. Centralbl. **9**. 439. 1882.
cuspidatum, Hookeri, Kapplerianum, Kraussii, membranaceum, Motleyi, Petersii, sphenoides, sublimbatum Prantl. Hym. 46. 1875 = T. sp. homonym.
Hookeri Pr. Abh. böhm. Ges. V. **5**. 336. 1848 = T. sphenoides.
lineolatum v.d.B.; Salom. Nom. 191. 1883.
muscoides Prantl, Hym. 46. 1875 = T. hymenoides.
peltatum Luerss. Bot. Centralbl. **11**. 28. 1882 = P. omphalodes.
punctatum v.d.B. Hym. Jav. 6. 1861.
pusillum Pr. Hym. 25 t. 9. 1843.
reptans Pr. Abh. böhm. Ges. V. **5**. 336. 1848.

HEMISTEGIA Presl, Abhandl. böhm. Ges. Wiss. V. **5**. 354. 1848 = **Hemitelia.**
ameristonevra Fée, 8 mém. 110. 1857.
contracta Fée, 8 mém. 110. 1857 — India.
decurrens Fourn. Mex. pl. **1**. 135. 1872.
elegantissima Fée, 8 mém. 110. 1857 = H. decurrens.
grandifolia Pr. Abh. böhm. Ges. V. **5**. 355. 1848.
horrida Fée, Gen. 351. 1850-52.
insignis Fée, 11 mém. 98 t. 26. 1866.
Kohautiana Pr. Abh. böhm. Ges. V. **5**. 355. 1848 = H. grandifolia.
lucida Fée, Gen. 351. 1850-52 = H. decurrens.
marginalis Pr. Abh. böhm. Ges. V. **5**. 355. 1848 = H. petiolata.
mexicana Fourn. Mex. pl. **1**. 135. 1872 = H. decurrens.
?monilifera Pr. Abh. böhm. Ges. V. **5**. 355. 1848.
?munita Pr. Abh. böhm. Ges. V. **5**. 355. 1848.
obtusa Pr. Abh. böhm. Ges. V. **5**. 355. 1848 = H. grandifolia.
repanda Fée, Gen. 351. 1850-52.; 11 mém. 98. 1866 — Cuba.
speciosa Fée, Gen. 351. 1850-52.
spectabilis Fée, Gen. 351. 1850-52 = H. subincisa.
?Willdenowii Fée, Gen. 351. 1850-52 = H. grandifolia.

HEMITELIA R. Brown. Prod. Fl. N. Holl. 158. 1810 — (Cyatheaceae Gen. 6). (Hemithelia Fée).
aculeata Fée, Gen. 350. 1850-52 = Alsophila armata.
acuminata Karst.: Schlecht. Bot. Zeit. **1856**. 474 = H. horrida.
alternans Hk. Ic. pl. t. 622 1844 = Cyathea.
ameristoneura (Fée) C. Chr. Ind. 347. 1905 — Guiana.
Hemistegia Fée, 8 mém. 110. 1857.
andina Karst. Linn. **28**. 452. 1856; Fl. Col. **2**. 183. t. 197. f. 1. 1869. HB. 455. NPfl. 131 — Columbia.
apiculata Hk. Syn. 29. 1865. Chr. 330. NPfl. 131 — Mexico. Brasilia austr.
arborea Fée, Gen. 350. 1850-51 = Cyathea.
aspera Fée, Gen. 350. 1850-51 = Alsophila.
atrovirens Trevis. Atti Ist. Veneto II. **2**. 164. 1851 = H. nigricans.
australis Pr. Epim. 33. 1849 — Australia trop. — Amphicosmia Moore 1857.
Beddomei Clarke, Tr. Linn. Soc. II. Bot. **1**. 429. 1880 = Cyathea spinulosa.
bella Reichb. fil.: Mett. Fil. Lips. 110. 1856. HB. 29. NPfl. 131 — Caracas.
Beyrichiana Pr. Abh. böhm. Ges. V. **5**. 353. 1848 = H. setosa. Brasilia.
boninsimensis Christ; Diels. Nat. Pfl. **1**[4]. 132. 1899 — Ins. Bonin.
Alsophila Christ in Warb. Monsunia **1**. 90. 1900.

Boryana Mett.: Kuhn, Linn. **36**. 161. 1869 — Guiana.
Brunoniana Clarke. Tr. Linn. Soc. II. Bot. **1**. 430. 1880 = Cyathea.
bullata Christ. Engl. Jahrb. **24**. 81. 1897 — Grenada.
calolepis Hk. Syn. 29. 1865. NPfl. 131 — Cuba.
capensis (L.fil.) (R.Br. Prod. Fl. N. Holl. 158. 1810) Klf. Enum. 253. 1824. HB. 29. NPfl. 132 — Africa austr. Brasilia. ? Java.
Polypodium L.fil. Suppl. 445. 1781; Cyathea Sm. 1793. Chr. 322; Aspidium Sw. 1801; Alsophila J. Sm. 1842; Cormophyllum Newm. 1854; Amphicosmia Kl. 1856. Cyathea riparia Willd. 1810. Hemitelia Desv. 1827; Amphicosmia Gardn. 1842 — Cyathea polypodioides Sw. 1817. — Folia advent.: Trichomanes incisum Thbg. 1800; Tr. cormophyllum Klf. 1824.
caudata Mett. Fil. Lechl. **2**. 30. 1859 = Alsophila.
commutata Schlecht. Bot. Zeit. **1856**. 474 = H. horrida.
cordata Desv. Prod. 321. 1827 = Cyathea marattioides.
costaricensis (Kl.) Mett. Ann. sc. nat. V. **2**. 265. 1864 (nomen); Kuhn, Linn. **36**. 159. 1869. HB. 455. NPfl. 131 — Costa Rica. Panama.
Cyathea Kl. msc.
crenata Sod. Rec. 18. 1883; Cr. vasc. quit. 524. 1893. NPfl. 131 — Ecuador.
Cyathea Christ 1897. 323.
crenulata Mett. Ann. Lugd. Bat. **1**. 55. 1863. — Java.
cruciata Desv. Prod. 320. 1827 = H. grandifolia.
Cumingii Trevis. Atti Ist. Veneto II. **2**. 164. 1851 = Alsophila extensa.
cyathoides Desv. Prod. 321. 1827 — Guiana.
cystolepis (Sod.) Bak. Ann. Bot. **5**. 187. 1891; Sod. Cr. vasc. quit. 523. 1893. NPfl. 131 — Ecuador. — Cyathea Sod. Rec. 15. 1883.
decipiens Scott, Tr. Linn. Soc. **30**. 33 t. 14. 1874 — Cyathea.
decurrens Liebm. Vid. Selsk. Skr. V. **1**. 286 (seors. 134). 1849 — Mexico.
Hemistegia Fourn. 1872. Hemitelia mexicana Liebm. 1849. NPfl. 131. Hemistegia Fourn. 1872. H. lucida Fée 1850-52. H. elegantissima Fée 1850-52.
denticulata Hk. Syn. 31. 1865. NPfl. 132 — Ins. Elisabethæ (mare pacif. austr.)
Amphicosmia Cumingii Moore 1857 (nomen).
D'Urvillei Mett.; Kuhn, Linn. **36**. 160. 1869 = H. tahitensis.
escuquensis Karst. Fl. Col. **2**. 181 t. 196. 1869. NPfl. 131 — Columbia. Amazonas. Puerto Rico. — H. multiflora β Sprucei Bak. 1870.
extensa Pr. Abh. böhm. Ges. V. **5**. 351. 1848 = Alsophila. — (f. Mett. = A. caudata.
falciloba Col. Tr. N. Zeal. Inst. **24**. 394. 1892 — N. Zealand.
firma Bak. JoB. **1877**. 161. NPfl. 131 — Ecuador.
Gardneriana Pr. Abh. böhm. Ges. V. **5**. 320. 1848 = H. capensis.
glandulosa Kuhn; Bak. Ann. Bot. **5**. 188. 1891. NPfl. 132 — Madagascar.
grandifolia (Willd.) Spr. Syst. **4**. 125. 1827. HB. 28. NPfl. 131 — Amer. trop.
Cyathea Willd. sp. **5**. 490. 1810; Hemistegia Pr. 1848; Microstegnus Pr. 1848; Cnemidaria Kuhn msc. (1897); Hemistegia Willdenowii Fée 1850-52. Hemitelia cruciata Desv. 1827. Chr. 329. H. spectabilis Kze. 1848. Cnemidaria Kohautiana Pr. 1836; Hemitelia Kze. 1844; Hemistegia Pr. 1848 — Hemitelia obtusa Klf. 1824; Cnemidaria Pr. 1836; Hemistegia Pr. 1848 — Hemitelia Klotzschiana Karst.; Kl. 1852.
?**Godeffroy** Luerss. Journ. Mus. Godeffr. **3**. 4. 1873 — Queensland
?guianensis Hk. Ic. pl. t. 648. 1844 = H. multiflora.
Hartii Bak. JoB. **1886**. 243. NPfl. 131 — Costa Rica. Columbia.

Hookeri Pr. Abh. böhm. Ges. V. **5.** 350. 1848 = H. horrida var.
Hookeriana Schlecht. Bot. Zeit. **1856.** 474 = H. horrida var. Hookeri.
horrida (L.) (R.Br. Prod. Fl. N. Holl. 158. 1810) Spr. Syst. 125. 1827 (excl. β). Chr. 330. NPfl. 131 — Ind. occ. Guatemala-Ecuador.
Polypodium L. sp. **2.** 1092. 1753; Cyathea Sm. 1793; Cnemidaria Pr. 1836; Actinophlebia Pr. 1848; Hemistegia Fée 1850-52; Cormophyllum Newm. 1854 — Cyathea commutata Spr. 1804; Hemitelia Schlecht. 1856 — H. Hookeri Pr. 1848 = H. horrida Hk. sp. **1.** 30 t. 15. Conf. Schlecht. Bot. Zeit. **1856.** 449, 465.
Hostmanni Hk. Ic. pl. t. 646. 1844 = H. multiflora var.
Imrayana Hk. Ic. pl. t. 669. 1844 — Ind. occ. Peru. Ecuador.
H. horrida β Imrayana HB. 28. H. serrata J. Sm. 1842 (nomen).
insignis (Fée) C. Chr. Ind. 349. 1905 — Guadeloupe. Brasilia.
Hemistegia Fée, 11 mém. 98 t. 26. 1866.
integrifolia Kl. Linn. **18.** 539. 1844 — H. speciosa.
javanica Pr. Epim. 31. 1849 = H. Junghuhniana.
Joadii Bak. Ann. Bot. **5.** 187. 1891. NPfl. 131 — Santa Marta.
Junghuhniana (Kze.) Mett. Fil. Lechl. **2.** 31. 1859. HB. 31. NPfl. 132 — Java. Sumatra. — Alsophila Kze. Bot. Zeit. **1848.** 284. A. lunulata Bl. 1828 (non R.Br. 1810). Hemitelia javanica Pr. 1849; Amphicosmia Moore 1857.
Karsteniana Kl. Allg. Gartenzeit. **20.** 42. 1852; Bot. Zeit. **1854.** 439. HB. 27. [NPfl. 131 — Caracas.
Kegelii Kze. Linn. **21.** 235. 1848 — H. multiflora.
Klotzschiana Karst.; Kl. Allg. Gartenzeit. **20.** 42. 1852; Bot. Zeit. **1854.** 440 = H. grandifolia var.
Kohautiana Kze. Bot. Zeit. **1844.** 298 — H. grandifolia.
laciniata Spr. Syst. **4.** 126. 1827 = ? Aspidium viridans.
latebrosa Mett. Fil. Lips. 111. 1856 = Alsophila.
Lechleri Mett. Fil. Lechl. **2.** 28. 1859. NPfl. 131 (Lechleriana) — Peru.
Leprieurii Kze. Bot. Zeit **1844.** 296 — Guiana.
(an Jenm. W. Ind. and Guiana Ferns 47. 1898?).
Lindeni Hk. Ic. pl. t. 706. 1844. HB. 29. NPfl. 131 — Caracas.
Lindeni L. Lind. Gard. Chr. III. **15.** 663. 1894 — Hort.
Lindigii Bak. Syn. 454. 1874. NPfl. 131 — Columbia.
lingulata Pr. Epim. 32. 1849 = H. multiflora.
Macarthurii F. Muell. Fragm. **8.** 176. 1874 = Cyathea.
macrocarpa Pr. Abh. böhm. Ges. V. **5.** 352. 1848. HB. ed. II. 30. NPfl. 131 — Brasilia. Guiana. — Amphicosmia Moore 1857. Hemitelia Moricandiana Kze.; Hk. 1865. HB. ed. I. 30.
macrosora Jenm. W. Ind. and Guiana Ferns 45. 1898 = Alsophila.
manilensis Pr. Abh. böhm. Ges. V. **5.** 351. 1848 — Luzon.
Alsophila Pr. 1836 (nomen); Amphicosmia Moore 1857.
marginalis J. Sm. Lond. JoB. **1.** 622. 1842 = H. petiolata.
marginalis Jenm. W. Ind. and Guiana Ferns 43. 1898 = Alsophila.
megalosora Trevis. Atti Ist. Veneto II. **2.** 164. 1851 = H. macrocarpa.
Melleri Bak. Syn. 456. 1874. NPfl. 132 — Madagascar.
Mertensiana Pr. Epim. 34. 1849 = Alsophila.
mexicana Liebm. Vid. Selsk. Skr. V. **1.** 287 (seors. 135). 1849 = H. decurrens
microphylla Col. Tr. N. Zeal. Inst. **27.** 399. 1895 — N. Zealand.
microsepala Jenm. JoB. **1886.** 266 = H. parvula.
monilifera J. Sm. Lond. JoB. **1.** 622. 1842 (nomen) — ? — Hemistegia Pr. 1848.

Moorei Bak. Gard. Chr. **1872**. 252. HB. 455 — Ins. Lord Howe.
Alsophila J. Sm. 1875 (non 1866); Cyathea F. Muell. 1882 (non Bak. 1874).
Moricandiana Kze.; Hk. Syn. ed. I. 30. 1865 = H. macrocarpa.
multiflora (Sm.) (R.Br. Prod. Fl. N. Holl. 158. 1810.) Spr. Syst. 126. 1827. HB. 30. NPfl. 131 — Amer. austr. trop. — Cyathea Sm. Mém. Ac. Turin **5**. 416. 1793; Sw. Syn. 140 (vide Obs. sub Alsophila phalerata). Chr. 323; Alsophila Pr. 1836; Amphicosmia Gardn. 1842. Hemitelia guianensis Hk. 1844 — H. lingulata Pr. 1849; Amphicosmia Moore 1857 — Hemitelia Kegelii Kze. 1848; Amphicosmia Moore 1857 — Polypodium pauciflorum Hk. 1862 — Hemitelia servitensis Karst. 1856.
—* Hemitelia Hostmanni Hk. 1844; Alsophila J. Sm. 1846; Amphicosmia Moore 1857 — Hemitelia urolepis Kze. 1848; Amphicosmia Moore 1857 — Hemitelia surinamensis Miq. 1843; Alsophila J. Sm. 1866.
—* Hemitelia Parkeri Hk. 1844; Amphicosmia Moore 1857; Alsophila J. Sm. 1875. A. Weigeltii Roem.; Pr. 1836 (nomen).
multiflora β Sprucei Bak. Fl. bras. 1^2. 313. 1870 = H. escuquensis.
munita (Willd.) Hk. sp. **1**. 32. 1844; Kuhn. Linn. **36**. 162. 1869 — »America«. Cyathea Willd.; Klf. 1824 (nomen); Cnemidaria Pr. 1836 (nomen); Hemistegia Pr. 1848 (nomen).
muricata Fée, Gen. 350. 1850-52 = Cyathea.
nigricans Pr. Epim. 31. 1849. HB. 454. NPfl. 131 — Guatemala. Peru. H. obscura Mett. 1864.
obscura Mett. Ann. sc. nat. V. **2**. 264. 1864; Karst. Fl. Col. **2**. 183 t. 197 f. 2. [= H. nigricans.
obtusa Klf. Enum. 252. 1824 = H. grandifolia.
obtusa Hk. sp. **1**. 29 t. 14 B. 1844 = H. subincisa.
obtusiloba Mett. msc.; Salomon, Nomenkl. 193. 1883 = Alsophila.
Parkeri Hk. Ic. pl. t. 643. 1844 = H. multiflora*.
parvula (Jenm.) Bak. Ann. Bot. **5**. 188. 1891. NPfl. 131 — Jamaica. Alsophila Jenm. JoB. **1879**. 258 — Hemitelia microsepala Jenm. 1886.
petiolata Hk. sp. **1**. 31 t. 16. 1844. HB. 28. NPfl. 131 — Panama. H. marginalis J. Sm. 1842 (nomen); Hemistegia Pr. 1848.
platylepis Hk. 2 Cent. t. 100. 1861. HB. 29. NPfl. 131 — Amazonas.
riparia Desv. Prod. 322. 1827 = H. capensis.
rufescens Mett.; Kuhn, Linn. **36**. 159. 1869. HB. 454 — Peru.
sagittifolia Jenm. W. Ind. and Guiana Ferns 43. 1898 = Alsophila.
samoensis Mett. Fil. Lechl. **2**. 31. 1859 = Alsophila.
Sellowiana Pr. Abh. böhm. Ges. V. **5**. 353. 1848 = ? Alsophila paleolata.
serra Desv. Prod. 221. 1827 = Cyathea.
serrata J. Sm. Lond. JoB. **1**. 662. 1842 = H. Imrayana.
servitensis Karst. Linn. **28**. 451. 1856 = H. multiflora.
sessilifolia Jenm. W. Ind. and Guiana Ferns 44. 1898 = Alsophila.
setosa (Klf.) Mett. Fil. Lechl. **2**. 30. 1859. HB. 455. NPfl. 131 — Brasilia. Alsophila Klf. Enum. 249. 1824. Cyathea Beyrichiana Pr. 1836; Hk. Ic. pl. t. 623. 1844; Amphicosmia Moore 1857; Alsophila J. Sm. 1875. (f. Rosenstock Cyathea sp.)
Sherringii Jenm. JoB. **1886**. 266. NPfl. 131 — Jamaica.
Smithii (Hk.fil.) Hk. Syn. 31. 1865. NPfl. 132 — N. Zealand. Cyathea Hk.fil. Fl. N. Zeal. **2**. 8 t. 22. 1854 — H. stellulata Col. 1886.
speciosa Klf. Enum. 252. 1824 = H. subincisa.

speciosa (H.B.Willd.) Desv. Prod. 320. 1827. HB. 28. Chr. 330. NPfl. 131 — Columbia. — Cyathea H.B.Willd. sp. **5**. 490. 1810; Hemistegia Fée 1850-52 — Hemitelia integrifolia Kl. 1844 — (§ Eleutheria Kze. 1844).
spectabilis Kze. Linn. **21**. 233. 1848 = H. grandifolia.
stellulata Col. Tr. N. Zeal. Inst. **18**. 222. 1886 = H. Smithii.
stigmosa Desv. Prod. 321. 1827 — »America calidior«.
subcaesia Sod. Cr. vasc. quit. 522. 1893. NPfl. 131 — Ecuador.
subincisa Kze. Bot. Zeit. **1844**. 296. HB. 27. NPfl. 131 — Guatemala-Peru. Guiana. — H. speciosa Klf. Enum. 252. 1824 (non Desv. 1827); Cnemidaria Pr. 1836; Hemitelia obtusa Hk. 1844 (non Klf. 1824); Actinophlebia Pr. 1848; Hemistegia spectabilis Fée 1850-52.
suprastrigosa Christ, Prim. Fl. costar. **3**. 44. 1901 — Costa Rica.
surinamensis Miq. »Diar. Inst. Reg. Bat. **1843**. 7« = H. multiflora *Hostmanni.
tahitensis (Brack.) Mett. Fil. Lechl. **2**. 31. 1859. HB. 455. NPfl. 132 — Tahiti. Alsophila Brack. Expl. Exp. **16**. 288 t. 40 f. 2. 1854; Amphicosmia Moore 1857 — Hemitelia D'Urvillei Mett. 1869.
Traillii Bak. Ann. Bot. **5**. 188. 1891. NPfl. 131 — Amazonas.
trinitensis Jenm. W. Ind. and Guiana Ferns 46. 1898 — Trinidad.
truncata Christ, Ann. Jard. Buit. **15**. 81 t. 13. f. 4. 1797 = Alsophila.
urolepis Kze. Linn. **21**. 235. 1848; **23**. 311. 1850 = H. multiflora *Parkeri.
?**velaminosa** Diels, Nat. Pfl. **1**[4]. 132. 1899 — Kamerun.
venosa Reichb.; Kuhn, Linn. **36**. 161. 1869. HB. 454 — Caracas.
Vrieseana Trevis. Atti Ist Veneto II. **2**. 164. 1851 = H. Junghuhniana.
Walkerae (Hk.) Pr. Abh. böhm. Ges. V. **5**. 351. 1848. HB. 30. NPfl. 132 — Ceylon. Cyathea Hk. Ic. pl. t. 647. 1844; Amphicosmia Moore 1857; Alsophila J. Sm. 1875; Hemitelia latebrosa v. Walkerae Mett. Fil. Lechl. **2**. 31. 1859.
Wilsoni Hk. Syn. 30. 1865. NPfl. 131 — Jamaica.

Hemiteliella Diels. Nat. Pfl. **1**[4]. 132. 1899 (*Hemitelia §*) = Hemitelia (capensis).

HETERODANAEA Presl, Suppl. Tent. Pter. 38. 1845 = **Danaea**.
stenophylla Pr. Suppl. 38. 1845.

Heteroglossum Diels, Nat. Pflanzenfam. **1**[4]. 334. 1899 (*Elaphoglossum §*) — Trachypteris.

HETEROGONIUM Presl, Epim. Bot. 142. 1849 = **Aspidium**.
aspidioides Pr. Epim. 143. 1849 = A. ambiguum.

HETERONEVRON Fée, Hist. Acrost. 20. 91. 1845 = **Leptochilus**.
argutum Fée, Acrost. 96 t. 25 f. 2. 1845 = L. cuspidatus.
bipinnatifidum Kuhn, Forschr. Gazelle **4**. Farne 6. 1889 = L. cuspidatus var.
cuspidatum Pr. Epim. 169. 1849.
diversifolium Fée, Acrost. 91. 1845.
heteroclitum Fée, Acrost. 92. 1845.
lobulosum Fée, Gen. 60. 1850-52 = Poecilopteris l.
lonchophorum Fée, Acrost. 94. 1845 = L. cuspidatus.
meniscioides Fée, Acrost. 93 t. 55. 1845 = L. nicotianifolius.
Naumanni Kuhn, Forschr. Gazelle **4**. Farne 5 t. 1. 1889.
?paradoxum Fée. 6 mém. 3 t. 1 f. 4. 1853 = Polypodium thyssanolepis (f. Fourn.).
Preslianum Fée, Acrost. 92 t. 39 f. 1. 1845.
proliferum Fée, Acrost. 95 t. 55. 1845.
punctulatum Fée, Acrost. 91 t. 54. 1845 = L. auriculatus.
Quoyanum Fée, Acrost. 96. 1845 = L. cuspidatus var.
Raddianum Fée, Acrost. 94. 1845 = L. guianensis.
repandum Fée, Acrost. 96 t. 57. 1845 = L. cuspidatus.

scalpturatum Fée, Acrost. 95 t. 56. 1845.
serratifolium Fée, Acrost. 94 t. 55. 1845.
sinuosum Fée, Acrost. 95 t. 55 f. 2. 1845 = L. cuspidatus.
subcrenatum Fée, Acrost. 93. 1845.
virens Fée, Acrost. 93. 1845.
HETEROPHLEBIUM Fée, Genera Filicum 139. 1850-52 = **Pteris.**
grandifolium Fée, Gen. 140 t. 11 A. f. 9-12. 1850-52.
HETEROPTERIS Fée. Dix. session du Congr. Sci. France. 1. 178. 1843 = **Paltonium.**
formosa Fée l. c. = P. lanceolatum.
lanceolata Diels, Nat. Pfl. 1^4. 305. 1899.
HETEROPTERIS Fée, Crypt. vasc. Brésil 1. 123. 1869 = **Doryopteris.**
Doryopteris Fée, Cr. vasc. Br. 1. 123 t. 10 f. 2. 1859 = D. lonchophora
HETEROPTERIS Presl; (Ettingsh. Farnkr. 96. 1865) = **Pteris** (lineata).
Heteropterygium Diels, Nat. Pflanzenf. 355. 1900 (*Gleichenia §*) = Gleichenia.
HEWARDIA J. Smith. Journ. Bot. 3. 432. 1841 = **Adiantum.**
adiantoides J. Sm. JoB. 3. 432 t. 16, 17. 1841.
diphylla Fée, Cr. vasc. Br. 1. 39 t. 9 f. 3. 1869 = A. adiantoides.
dolosa Fée, Gen. 122. 1850-52.
Leprieurii Fée, Gen. 122. 1850-52.
serrata Fée, Gen. 122. 1850-52 = A. dolosum.
Wilsoni Fée, Gen. 122. 1850-52 = A. dolosum.
HIATEA Menzies msc. (Hk. sp. 1. 82. 1844) = **Cibotium.**
HICRIOPTERIS Presl, Epim. Bot. 26. 1849 = **Gleichenia.**
speciosa Pr. Epim. 27. 1849 = G. glauca.
HIPPODIUM Gaudichaud, (Freyc. Voy. 340. 1827) = **Didymochlaena.**
HISTIOPTERIS (Agardh) J. Smith, Hist. Fil. 295. 1875 — (Polypodiaceae Pteris § Histiopteris Agardh, Rec. Gen. Pter. 76. 1839. [Gen. 84).
aurita J. Sm. Hist. Fil. 295. 1875 = H. incisa var.
incisa (Thbg.) J. Sm. Hist. Fil. 295. 1875. NPfl. 294 — Trop. et subtrop.-Ins. antarct. — Pteris Thbg. Prod. Fl. Cap. 171. 1800. HB. 172. Chr. 163; Litobrochia Pr. 1836; Phegopteris Keys. 1873. Pteris Vespertilionis Lab. 1806; Litobrochia Pr. 1836; Histiopteris J. Sm. 1875 — Pteris elegans Sw. 1817; Litobrochia Brack. 1854. Pteris pallida Raddi 1819; Litobrochia Pr. 1836 — L. sinuata Brack. 1854; Histiopteris J. Sm. 1875 — Pteris aurita Bl. 1828; Litobrochia Pr. 1836; Histiopteris J. Sm. 1875.
montana Copeland in Perkins, Fragm. 193. 1905 — Ins. Philipp.
sinuata J. Sm. Hist. Fil. 295. 1875 = H. incisa.
Vespertilionis J. Sm. Hist. Fil. 295. 1875 = H. incisa.
Holcochlaena Baker. Syn. Fil. 153. 1867 (*Pellaea §*) = Pellaea.
HOLCOSORUS Moore, Index Fil. XXIX. 1857 = **Polypodium.**
pentagonus Moore, Ind. XXX. 1857 = P. bisulcatum.
Holodanaea Presl, Suppl. Tent. Pterid. 37. 1845 (*Danaea §*) = Danaea.
Holophlebium Christ, Farnkr. 27. 1897 (*Trichomanes §*) = Trichomanes.
Holopterygium Diels, Nat. Pflanzenfam. 1^4. 353. 1900 (*Gleichenia §*) = Gleichenia.
Homaloneuron Klotzsch, Linnaea 20. 354. 1847 (*Asplenium §*) = Asplenium.
HOMOEOTES Presl. Abh. böhm. Ges. Wiss. V. 5. 331. 1848 = **Trichomanes.**
heterophylla Pr. l. c. 332. 1848.
Homopteris Ruprecht. Distr. Cr. vasc. Ross. 48. 1845 (*Allosorus §*) = Cryptogramma (Stelleri).
HUMATA Cavanilles, Descripc. plant. 272. 1802 — (Polypodiaceae Gen 28).

aemula (Mett.) Carr. in Seem. Fl. Vit. 335. 1873 — Aneiteum.
Davallia Mett.; Kuhn, Linn. **36**. 144. 1869. HB. 467.
affinis Mett. Fil. Lips. 102 t. 27 f. 5, 6. 1856 = Davallia hymenophylloides.
alpina Moore, Ind. XCII. 1857 = H. repens var.
Anderssonii (Mett.) C. Chr. Ind. 353. 1905 — Tahiti.
Davallia Mett.; Kuhn, Linn. **36**. 143. 1869. HB. 467.
angustata J. Sm. JoB. **3**. 416. 1841 = H. heterophylla var.
asplenioides Desv. Prod. 324. 1827 — India orient. — Davallia Moore 1861.
bipinnatifida Moore, Ind. 290. 1861 = H. vestita.
botrychioides Brack. Expl. Exp. **16**. 231 t. 32 f. 1. 1854. NPfl. 209 — Celebes. Ins. Philipp. Polynesia. — Davallia Bak. 1867. HB. 90. 467 — Humata multifida Carr. 1873 — H. rigida (Sol.) Carr. 1873; Davallia Sol.; Salom. 1883.
chaerophylla Mett. Fil. Lips. 102. 1856 = Davallia pulchra.
chaerophylloides Desv. Prod. 325. 1827 = Davallia.
cicutarioides Diels, Nat. Pfl. 1⁴. 210. 1899 = Davallia.
contigua Desv. Prod. 323. 1827 = Davallia.
Cumingii Brack. Expl. Exp. **16**. 230. 1854 = H. lepida.
dareoidea Mett. Aspl. 65 t. 6 f. 21, 22. 1859 = Asplenium davallioides.
elata Desv. Prod. 325. 1827 = Davallia denticulata.
elegans Desv. Prod. 324. 1827 = Davallia denticulata.
epiphylla Desv. Prod. 325. 1827 = Davallia denticulata var.
falcata Cav. Hort. Matrit. —; Desv. Prod. 324. 1827 = Nephrolepis cordifolia.
falcinella Copeland, Polyp. Philipp. 51. 1905 = Davallia.
ferulacea Diels, Nat. Pfl. 1⁴. 210. 1899 = Davallia.
Gaimardiana (Gaud.) J. Sm. Lond. JoB. **1**. 425. 1842. NPfl. 209 — Birma-Malesia. Polynesia. — Nephrodium Gaud. Freyc. Voy. Bot 335 t. 12 f. 1. 1827; Nephrolepis Pr. 1836; Davallia? Pr. 1836. D. parallela Wall. 1828. HB. 89. Chr. 304; Pachypleuria Pr. 1849; Pteronevron Fée 1850-52; Humata Brack. 1854; Oleandra Keys. 1873 — Davallia intermarginalis Bl. 1828; Pachypleuria Pr. 1849; Humata Moore 1861.
heterophylla (Sm.) Desv. Prod. 323. 1825. NPfl. 209. — Malesia. Polynesia. Davallia Sm. Mém. Ac. Turin **5**. 415. 1793; Sw. Schrad. Journ. 1800². 86. 1801. HB. 88. Chr. 303; Trichomanes Poir. 1808. Humata ophioglossa Cav. 1802 — H. pinnatifida Cav. 1802; Davallia Sw. 1806.
—* D. angustata Wall. 1828; Hk. et Grev. Ic. Fil. t. 231. 1831. HB. 88; Humata J. Sm. 1841; Pachypleuria Pr. 1849.
hirsuta Desv. Prod. 324. 1827 = Microlepia pilosella.
Hookeri Diels, Nat. Pfl. 1⁴. 209. 1899 = Davallia Clarkei.
hymenophylloides Copeland, Polyp. Philipp. 51. 1905 = Davallia.
immersa Mett. Fil. Lips. 102. 1856 = Davallia.
?Imrayana Moore, Ind. XCII. 1857 = Saccoloma.
intermarginalis Moore, Ind. 296. 1861 = H. Gaimardiana.
intermedia C. Chr. Ind. 353. 1905 — Borneo. Perak.
Davallia pinnatifida Bak. Journ. Linn. Soc. **24**. 257. 1887; Humata Bedd. 1892 (non Cav. 1802.)
Kingii Diels, Nat. Pfl. 1⁴. 210. 1899 = Davallia.
lepida (Pr.) Moore, Ind. XCII. 1857 — Ins. Philipp.
Davallia Pr. Tent. 128. 1836 (nomen); Goldm. Nova Acta **19** Suppl. I. 464. 1843; Pachypleuria Pr. 1849. Davallia Cumingii Hk. 1846. HB. 90; Humata Brack. 1854 — Davallia longula Kze. 1848; Humata Moore 1861.

longula Moore, Ind. 296. 1861 = H. lepida.
?Mannii Diels, Nat. Pfl. 1⁴. 210. 1899 = Davallia.
membranulosa Diels, Nat. Pfl. 1⁴. 209. 1899 = Davallia.
micans Diels, Nat. Pfl. 1⁴. 209. 1899 = Davallia assamica.
Moritziana (Kl.) Moore, Ind. 297. 1861 (nomen) — Venezuela.
Davallia Kl. msc.; Kze. Bot. Zeit. **1850**. 54 (nomen).
multidentata Diels, Nat. Pfl. 1⁴. 209. 1899 = Davallia.
multifida Carr. in. Seem. Fl. Vit. 335. 1873 = H. botrychioides var.
neoguineensis C. Chr. Ind. 354. 1995 — N. Guinea.
Davallia bipinnatifida Bak. Kew Bull. **1899**. 119 (non Bl. 1828).
ophioglossa Cav. Descr. 273. 1802 = H. heterophylla.
parallela Brack. Expl. Exp. **16**. 229. 1854 = H. Gaimardiana.
parvula Mett. Fil. Lips. 102 t. 27 f. 7, 8. 1856 = Davallia.
patens Desv. Prod. 325. 1827 = Davallia denticulata.
pectinata (Sm.) Desv. Prod. 323. 1827 — Polynesia.
Davallia Sm. Mém. Ac. Turin **5**. 415. 1793. HB. 89; Trichomanes Poir. 1808; Pachypleuria Pr. 1849.
pedata J. Sm. JoB. **3**. 416. 1841 = H. repens.
pinnata Desv. Prod. 324 t. 8 f. 1. 1827 = Davallia pentaphylla.
pinnatifida Cav. Descr. 273. 1802 = H. heterophylla.
pinnatifida Bedd. Handb. Suppl. 12. 1892 = H. intermedia.
polypodioides Desv. Prod. 325. 1827 = Microlepia speluncae.
polypodioides Brack. Expl. Exp. **16**. 228 t. 32 f. 1. 1854 = H. sessilifolia.
pulcherrima Diels, Nat. Pfl. 1⁴. 210. 1899 ≐ Davallia.
pulchra Diels, Nat. Pfl. 1⁴. 209. 1899 = Davallia.
pusilla (Mett.) Carr. in Seem. Fl. Vit. 335. 1873. NPfl. 209 — Melanesia.
Davallia Mett. Ann. sc. nat. IV. **15**. 79. 1861. HB. 89.
pyxidata Desv. Prod. 325. 1827 = Davallia.
repens (L.fil.) Diels, Nat. Pfl. 1⁴. 209. 1899 — Japonia. China-Asia trop. Austr. trop. Ins. Mascarenae. Ins. Sechellae. — Adiantum L.fil. Suppl. 446. 1781; Davallia Kuhn 1867. D. pedata Sm. 1793. HB. 89. Chr. 304; Humata J. Sm. 1841; Pachypleuria Pr. 1849. Humata trifoliata Cav. 1802; Pachypleuria Pr. 1849. Davallia serrata Willd. 1810; Humata Desv. 1827 — Davallia alpina Bl. 1828. HB. 89. Chr. 304; Humata Moore 1857. NPfl. 209.
rigida Carr. in Seem. Fl. Vit. 335. 1873 = H. botrychioides var.
scandens Desv. Prod. 324. 1827 = Odontosoria.
serrata Desv. Prod. 323. 1827 = H. repens.
serrata Brack. Expl. Exp. **16**. 230. 1854 = H. repens var. alpina.
sessilifolia (Bl.) Mett. Fil. Lips. 102. 1856. NPfl. 209 — Java. Celebes. Fiji.
Davallia Bl. Enum. 231. 1828. HB. 89; Pachypleuria Pr. 1849 — Humata polypodioides Brack. 1854.
solida Desv. Prod. 324. 1827 = Davallia.
trifoliata Cav. Descr. 273. 1802 = H. repens.
Tyermanni Moore. Gard. Chr. **1871**. 870 f. 178. NPfl. 209 — China.
Davallia Moore 1871. HB. 467.
vestita (Bl.) Moore. Ind. XCII. 1857. NPfl. 209 — Malesia.
Davallia Bl. Enum. 233. 1828. HB. 90; Pachypleuria Pr. 1849.

HYALOLEPIS Kunze, Linnaea **23**. 258. 1850 = **Hymenolepis.**
ophioglossoides Kze. Linn. **23**. 258. 1850 = H. spicata.
revoluta Kze. Linn. **23**. 258. 1850 = H. spicata.

HYDROGLOSSUM Willdenow, Schr. Akad. Wiss. Erfurt **1802**. 20 = **Lygodium.**
auriculatum Willd. sp. **5**. 84. 1810 = L. flexuosum.
circinnatum Willd. Schr. Akad. Erfurt **1802**. 24.
cubense Steud. Nomencl. **2**. 205. 1824.
dichotomum Willd. sp. **5**. 82. 1810 = L. circinnatum.
dissectum Steud. Nomencl. **2**. 205. 1824 = L. japonicum.
elegans Steud. Nomencl. **2**. 205. 1824 = L. flexuosum.
expansum Poir. Enc. Suppl. **3**. 77. 1813 = L. volubile var.
flexuosum Willd. Schr. Akad. Erfurt **1802**. 23 t. 1 f. 3.
hastatum Willd. sp. **5**. 79. 1810 = L. polymorphum.
heptaphyllum Schrad. Gött. gel. Anz. **1824**. 863 = L. volubile.
heterodoxum Moore, Ind. CXIV. 1857.
hirsutum Willd. sp. **5**. 80. 1810 = L. polymorphum.
japonicum Willd. Schr. Akad. Erfurt **1802**. 26.
lanceolatum Steud. Nomencl. **2**. 205. 1824.
longifolium Willd. Schr. Akad. Erfurt **1802**. 22 t. 2 = L. circinnatum.
madagascariense Poir. Enc. Suppl. **3**. 78. 1813 = L. lanceolatum.
mexicanum Fée, 9 mém. 42. 1857 = L. heterodoxum.
microstachyum Steud. Nomencl. **2**. 206. 1824 = L. flexuosum.
oligostachyum Willd. sp. **5**. 81. 1810.
palmatum Willd. Schr. Akad. Erfurt **1802**. 25 t. 1 f. 2.
pedatum Willd. Schr. Akad. Erfurt **1802**. 25 = L. circinnatum.
pinnatifidum Willd. Schr. Akad. Erfurt **1802**. 21 = L. flexuosum.
polycarpum Willd. sp. **5**. 79. 1810 = L. reticulatum.
scandens Willd. Schr. Akad. Erfurt **1802**. 20 = { L. scandens. / L. polymorphum (β).
scandens Pr. Suppl. 113. 1845 = L. reticulatum.
semipinnatum Poir. Enc. Suppl. **3**. 78. 1813 = L. flexuosum.
spectabile Liebm. Vid. Selsk. Skr. V. **1**. 299 (seors. 147). 1849 = L. heterodoxum.
volubile Willd. sp. **5**. 78. 1810.

HYMENOCYSTIS C. A. Meyer, Verz. Pl. Cauc. 229. 1831 = **Woodsia.**
caucasica Meyer, Verz. Pl. Cauc. 229. 1831 = W. fragilis.

HYMENODIUM Fée. Hist. Acrost. 20, 90. 1845 = **Elaphoglossum.**
crassifolium Fée, Acrost. 91 t. 63 f. 1. 1845 = E. reticulatum.
crinitum Fée, Acrost. 90. 1845.
Kunzeanum Fée, Acrost. 90 t. 58. 1845 = E. pachyphyllum.
pachyphyllum Moore, Ind. XIX. 1857.
reticulatum Moore, Ind. XIX. 1857.

HYMENOGLOSSUM Presl, Hymenoph. 35. 1843 = **Hymenophyllum.**
cruentum Pr. Hym. 35. 1843.

HYMENOLAENA Mey. (Fée. Gen. 358. 1850-52) = **Woodsia.**

HYMENOLEPIS Kaulfuss, Enum. Fil. 146. 1824 — (Polypodiaceae Gen. **99**).
abbreviata Fée, 10 mém. 12. 1865 = Drymoglossum.
brachystachys (Hk.) J. Sm. Ferns br. and for. 92. 1866. NPfl. 305 — Malesia.
Acrostichum spicatum β brachystachys Hk. Gard. Ferns t. 3. 1862. HB. 424; A. brachystachyum Racib. 1898.
callifolia Christ, Ann. Jard. Buit. II. **5**. 128. 1905 — Borneo.
mucronata Fée, Gen. 82. 1850-52 = H. spicata var.
ophioglossoides Klf. Enum. 146 t. 1 f. 9. 1824 = H. spicata.

platyrhynchos (J. Sm.) Kze. Farnkr. **1**. 101. 1842. Chr. 129. NPfl. 305 — Celebes. Ins. Philipp. — Gymnopteris J. Sm. 1841 (nomen); Macroplethus Pr. Epim. 141. 1849; Acrostichum Hk. 1864. HB. 424; Taenitis Mett. 1869.

revoluta Bl. Enum. 201. 1828 = H. spicata.

spicata (L.fil.) Pr. Epim. 159. 1849. Chr. 129. NPfl. 305 — Madagascar. Ins. Mascar. Asia. Austr., Polynesia trop. — Acrostichum L.fil. Suppl. 444. 1781. HB. 424; Schizaea Sm. 1793; Onoclea Sw. 1801; Belvisia Mirb. 1803; Lomaria Willd. 1810; Gymnopteris Pr. 1836; Taenitis Mett. 1868. Hymenolepis ophioglossoides Klf. 1824; Hyalolepis Kze. 1850; Taenitis Mett. 1856; Oetosis O.Ktze. 1891 — Hymenolepis revoluta Bl. 1828; Hyalolepis Kze. 1850; Taenitis Mett. 1856 — Hymenolepis validinervis Kze. 1848; Taenitis Mett. 1868-69.

validinervis Kze. Bot. Zeit. **1848**. 122 = H. spicata var.

Hymenoloma Davenport, Bot. Gazette **21**. 262. 1896 (*Pellaea §*) = Pellaea (membranacea)

HYMENOPHYLLUM Smith, Mém. Acad. Turin **5**. 418. 1793 — (Hymenophyllaceae Gen. **2**).

abietinum Hk. et Grev. Ic. Fil. t. 127. 1829 = H. polyanthos var.

abruptum Hk. sp. **1**. 88 t. 31 B. 1844. HB. 57. Chr. 16 — Amer. trop. Patagonia. Leptocionium Pr. 1848 — Hymenophyllum brevifrons Kze. 1847. NPfl. 108.

acrosorum v.d.B. Pl. Jungh. **1**. 564. 1856 = H. Blumeanum.

aculeatum (J. Sm.) Racib. Pterid. Buit. 21. 1898 — Ceylon. Malesia. Trichomanes J. Sm. 1841 (nomen, non Sw. 1788); Didymoglossum v.d.B 1856; Leptocionium v.d.B. 1859; Hym. Jav. 41 t. 31. 1861. Hymenophyllum Neesii HB. 71 (pt.). Chr. 22 (pt.?). NPfl. 112 — Didymoglossum acanthoides v.d.B. 1856; Leptocionium v.d.B. 1859: Hym. Jav. 42 t. 32. 1861.

aculeolatum v.d.B. Ned. kr. Arch. **5**[3]. 179. 1863 — Brasilia.

acutum Mett. (Sadeb. NPfl. 108. 1899) = H. polyanthos var.

adiantoides v.d.B. Ned. kr. Arch. **5**[3]. 184. 1863 — Peru.

aequabile Kze. Bot. Zeit. **1847**. 200 = H. interruptum.

aeruginosum (Poir.) Carm. Tr. Linn. Soc. **12**. 513. 1818. HB. 64. NPfl. 110 — Tristan d'Acunha. — Trichomanes Poir. Enc. **8**. 76. 1808; Thouars. Fl. Trist. d'Ac. 34. 1804(?).

aeruginosum β Franklinianum Hk. sp. **1**. 93 t. 34 A. 1844 = H. ferrugineum

affine Brack. Expl. Exp. **16**. 265 t. 37 f. 1. 1854 — Fiji.

affine Racib. Pterid. Buit. 20. 1898 = H. holochilum.

alatum Schkuhr, Kr. Gew. **1**. 133 t. 135 b. 1809 = Trichomanes bipunctatum.

alatum Willd. sp. **5**. 526. 1810 = H. tunbridgense.

alatum Sm. Engl. Bot. t. 1417. 1805 = Trichomanes radicans var.

alpinum Col. Tr. N. Zeal. Inst. **31**. 263. 1899 — N. Zealand.

alveolatum C. Chr. Ind. 356. 1905 — Ecuador.
H. divaricatum Sod. Cr. vasc. quit. 29. 1893 (non v.d.B. 1863).

amoenum Sturm, Fl. bras. **1**. 291. 1859 = H. crispum.

andinum v.d.B. Ned. kr. Arch. **4**. 397. 1859. HB. 60 — Ecuador.

angustatum Thouars; Kze. Bot. Zeit. **1847**. 186 = H. ciliatum var.

angustifrons Christ, Bull. Boiss. II. **4**. 940. 1904 — Costa Rica-Columbia.

angustum v.d.B. Ned. kr. Arch. **5**[3]. 183. 1863 = H. hirsutum.

antarcticum Pr. Hym. 31. 50 t. 12 A. 1843 = H. tunbridgense.

antillense Jenm. Bull. Dep. Jam. n. 18. 6. 1890 — Jamaica.
H. lineare var. antillense Jenm. JoB. **1879**. 258.

apicale v.d.B. Ned. kr. Arch. **4**. 397. 1859 = H. axillare.
apiculatum Mett.; Kuhn, Linn. **35**. 391. 1868. (NPfl. 108) — Venezuela.
apterum v.d.B. Ned. kr. Arch. **5**[3]. 194. 1863 = H. ciliatum var.
arbuscula Desv. Prod. 332. 1827 = H. ciliatum.
Armstrongii (Bak.) Kirk, Tr. N. Zeal. Inst. **10**. XLIII. t. 21 f. A. 1878 — N. Zealand. — Trichomanes Bak. Syn. ed. I. 452. 1868; ed. II. 465. 1874. NPfl. 105. Hymenophyllum melanocheilos Col. 1885.
asperulum Kze. Linn. **9**. 109. 1834 = H. tunbridgense.
asplenioides Sw. Schrad. Journ. **1800**[2]. 98. 1801. HB. 57. Chr. 15. NPfl. 108 — Amer. trop. — Trichomanes Sw. Prod. 136. 1788.
asterothrix Kze. Farnkr. **1**. 160. 1844 (nomen) = H. fusugasugense.
atrosanguineum v.d.B. Ned. kr. Arch. **5**[3]. 165. 1859 — Columbia.
atrovirens Col. Tasm. Journ. **2**. 186. 1844 = H. australe var.
atrovirens Fée et L'Herm.; Fée, 11 mém. 120 t. 30 f. 4. 1866 = H. fucoides.
atrovirens Christ, Bull. Boiss. II. **4**. 941. 1904 = H. subrigidum.
attenuatum Hk. sp. **1**. 99 t. 36 B. 1844 = H. magellanicum.
Aucklandicum v.d.B. Ned. kr. Arch. **4**. 393. 1859 = H. australe var.
australe Willd. sp. **5**. 527. 1810. Chr. 17. NPfl. 108 — India bor-Malesia-Australia-N. Zealand. Ins. antarct. — Sphærocionium Pr. 1843; Hymenophyllum javanicum Spr. 1827; v.d.B. Hym. Jav. 50 t. 40. HB. 60 — H. crispatum Wall. 1828; Hk. et Grev. t. 77; Sphærocionium Pr. 1843 — H. flexuosum A. Cunn. 1836 — H. fimbriatum J. Sm. 1841 — H. atrovirens Col. 1844 — H. dædaleum Bl. 1828 — H. erosum Bl. 1828; v.d.B. Hym. Jav. 54 t. 43 — H. micranthum v.d.B. 1856; Hym. Jav. 52 t. 41.
axillare Sw. Schrad. Journ. **1800**[2]. 101. 1801. HB. 58. NPfl. 108 — Ind. occ. Venezuela. — Trichomanes Poir. 1808; Sphærocionium Pr. 1843 — Hymenophyllum apicale v.d.B. 1859.
axillare Raddi, Opusc. sci. Bol. **3**. 297. 1819 = Trichomanes rupestre.
badium Hk. et Grev. Ic. Fil. t. 76. 1828. HB. 60. NPfl. 108 — Himalaya. [Malacca.
badium Wall. List n. 172. 1828 = H. Blumeanum.
Balansæ Fourn. Ann. sc. nat. V. **18**. 265. 1873 — N. Caledonia.
Baldwinii Eat. Bull. Torr. Cl. **6**. 293. 1879. NPfl. 108 — Ins. Hawaii.
Balfourii Bak. Ann. Bot. **5**. 192. 1891 — Réunion.
barbatum (v.d.B.) Bak. Syn. 68. 1874. NPfl. 110 — Korea. Tsus Sima. China. Leptocionium v.d.B. Ned. kr. Arch. **5**[3]. 146. 1863.
Berteroi Hk. sp. **1**. 93 t. 33 C. 1844 = H. ferrugineum.
Beyrichianum Kze. Linn. **9**. 108. 1834 = H. microcarpum.
Bibraianum Sturm. Flora **1853**. 361 = H. magellanicum.
bifarium Mett.; Sadeb. Nat. Pfl. **1**[4]. 108. 1899 = H. mnioides.
bismarckianum Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 34. 1905 — N. Guinea.
bivalve (Forst.) Sw. Schrad. Journ. **1800**[2]. 99. 1801. HB. 69. NPfl. 110 — N. S. Wales. N. Zealand. — Trichomanes Forst. Prod. 84. 1786; Sphaerocionium Pr. 1843 — Hymenophyllum pyriforme v.d.B. 1863.
bivalve J. Sm. JoB. **3**. 418. 1841 = H. serrulatum.
blandum Racib. Pterid. Buit. 20. 1898. NPfl. 110 — Java.
blepharodes Pr. Hym. 32, 51. 1843 = H. fucoides.
Blumeanum Spr. Syst. **4**. 131. 1827; v.d.B. Hym. Jav. 46 t. 36. NPfl 108 — Asia trop. — Meringium? Pr. 1843; Hymenophyllum pectinatum Nees et Bl. Nova Acta **11**. 124 t. 12 f. 5. 1823 (non Cav. 1802); H. acrosorum

v.d.B. 1856 — H. integrum v.d.B. 1856; Hym. Jav. 49 t. 38 — H. pycnocarpum v.d.B. Hym. Jav. 48 t. 37 — H. Himalaianum. Osmundoides Sphaerocarpum v.d.B. 1863 — H. polyanthos auctt. quoad pl. asiat.

borneense Hk. Syn. 62. 1866 — Borneo.

Boryanum Willd. sp. **5**. 518. 1810 = H. ciliatum var.

botryoides v.d.B. Ned. kr. Arch. **5**[3]. 160. 1863 — Mexico.

Boutonii Bak. Fl. Maur. 462. 1877 — Mauritius.

brachyglossum A.Br.; Kze. Bot. Zeit. **1847**. 227. (NPfl. 112) — Java. Borneo. Didymoglossum Braunii v.d.B. 1856; Leptocionium v.d.B. 1859: Hym. Jav. 43 t. 33. 1861.

brachypus Sod. Cr. vasc. quit. 28. 1893 — Ecuador.

Breutelii v.d.B. Ned. kr. Arch. **5**[3]. 154. 1863 = H. polyanthos var.

brevifrons Kze. Bot. Zeit. **1847**. 185 — H. abruptum.

brevistipes Liebm. Vid. Selsk Skr. V. **1**. 290 (seors. 138). 1849 — Mexico. Caracas. Cuba.

Bridgesii Hk. sp. **1**. 97 t. 35 C 1844 = H. dentatum.

caespitosum Gaud. Ann. sc. nat. **5**. 99. 1825 = Trichomanes.

caespitosum Fée. 11 mém. 119 t. 30 f. 3. 1866 — Guadeloupe.

caespitosum Christ apud Dusén, Svensk Exp. till Magelhl. **3**. 242. 1899 = [H. glebarium.

calodictyon v.d.B. Ned. kr. Arch. **5**[3]. 172. 1863 — Peru.

capense Schrad. Gött. gel. Anz. **1818**. 919 = H. fumarioides.

capillaceum Roxb. in Beats. St. Helena 311. 1816 (nomen?); Hk. sp. **1**. 109 t. 38 B. 1844. HB. 58. NPfl. 108 — St. Helena.
H. infortunatum Bory. Dup. Boy. Bot. **2**. 284 t. 38 f. 3. 1828 (an nomen opt.?); Sphaerocionium Pr. 1843.

capillare Desv. Prod. 333. 1827 — Tristan d'Acunha. Réunion.
H. pendulum Bory 1833. (NPfl. 110); Sphaerocionium Pr. 1843 — Hymenophyllum fulvum v.d.B. 1863 — H. lineare pt. HB. 66.

carnosum Christ, Bull. Boiss. II. **4**. 938. 1904 — Costa Rica.

Catherinae Hk. Syn. 67. 1867 = H. lineare.

caudatellum Christ, Bull. Boiss. II. **4**. 939. 1904 — Costa Rica.

caudatum v.d.B. Ned. kr. Arch. **4**. 394. 1859 = H. caudiculatum.

caudiculatum Mart. Ic. Cr. Bras. 102 t. 67. 1834. HB. 61. Chr. 17. NPfl. 108 — Brasilia-Patagonia. Peru-Juan Fernandez — Sphaerocionium Pr. 1843 — S. productum Pr. 1843; Hymenophyllum Sturm 1858 (non Kze. 1848); H. caudatum v.d.B. 1859 — H. Martii Sturm 1859.

caulopteron Fée, Cr. vasc. Br. **1**. 197 t. 70 f. 3. 1869 = H. ciliatum.

ceratophylloides Christ, Bull. Boiss. II. **4**. 942. 1904 — Costa Rica. Quito.

Cheesemanni Bak. in Hk. Ic. pl. t. 1132. 1873. HB. 464 — N. Zealand.

Chiloense Hk. sp. **1**. 90 t. 32 A. 1844 = H. dicranotrichum.

chrysothrix Sturm, Fl. bras. **1**[2]. 298. 1859; Bot. Zeit. **1859**. 299 — Venezuela. Brasilia.

ciliatum Sw. Schrad. Joun. **1800**[2]. 100. 1801. HB. 63. Chr. 19. NPfl. 110 — Amer. trop. Africa trop. et austr. cum ins. Sikkim. N. Zealand.
Trichomanes Sw. Prod. 136. 1788; Sphaerocionium Pr. 1843. Hymenophyllum Boryanum Willd. 1810; Sphaerocionium Pr. 1843. Hymenophyllum Arbuscula Desv. 1827 — H. Plumieri Hk. et Grev. 1829; Sphaerocionium Pr. 1843 — S. vestitum, commutatum. Schiedeanum. Sieberi, Grevilleanum Pr. 1843; Hymenophyllum commutatum, Sieberi v.d.B. 1859. H. vestitum v.d.B. 1863 — H. remotum. Surinamense v.d.B. 1859

— H. apterum v.d.B. 1863 — H. splendidum v.d.B. 1863 — H. trapezoidale Liebm. 1849 — H. Gardnerianum Sturm 1859 — H. microcarpon Fée 1869 (non Desv. 1827) — H. Fraseri Mett.; Fourn. 1872 — H. elatius Christ 1900.

clavatum Sw. Schrad. Journ. **1800**². 101. 1801 = H. polyanthos var.

coloratum A. Br.; v.d.B. Pl. Jungh. **1**. 565. 1856 = H. paniculiflorum.

commutatum v.d.B. Ned. Kr. Arch. **4**. 414. 1859 = H. ciliatum.

constrictum Christ, Bull. Boiss. II. **4**. 939. 1904 — Costa Rica.

contortum v.d.B. Ned. kr. Arch. **5**³. 170. 1863 — Costa Rica. Columbia. Bolivia.

contractile Sod. Cr. vasc. quit. 22. 1893 — Ecuador.

convolutum v.d.B. Ned. kr. Arch. **4**. 400. 1859 = H. reniforme.

corrugatum Christ, Bull. Boiss. II. **3**. 508. 1903 — China.

corticola Hk. in Thwait. Enum. pl. zeyl. 397. 1864 = Trichomanes digitatum.

costaricanum v.d.B. Ned. kr. Arch. **5**³. 161. 1863 — Costa Rica. Columbia. [— Bolivia.

crispatulum v.d.B. Ned. kr. Arch. **4**. 412. 1859 — Peru.

crispatum Wall. List n. 169. 1828 — H. australe.

crispum HBK. Nov. Gen. et Sp. **1**. 26. 1815. HB. 59. Chr. 16. NPfl. 108 — Amer. trop. — Sphaerocionium Kl. 1844 — Hymenophyllum amoenum Sturm 1859 — H. polycarpum Mett.; Kuhn 1868.

crispum Nees et Bl. Nova Acta **11**. 128 t. 14 f. 1. 1823 = H. australe.

crispum var brasiliana Fée, Cr. vasc. Br. **1**. 195 t. 71 f. 2. 1869 = H. ciliatum.

cristatum Hk. et Grev. Ic. Fil. t. 148. 1829 = H. fucoides var.

cruentum Cav. Descr. 275. 1802. HB. 56. Chr. 15. NPfl. 108 — Chile. Juan Fernandez. — Hymenoglossum Pr. 1843.

Crügeri Müll. Bot. Zeit. **1854**. 722 = H. hirsutum vel H. lineare.

cubense Sturm, Bot. Zeit. **1859**. 298. NPfl. 110 — Cuba.

Cumingii Pr. Hym. 32, 56 t. 11 A. 1843 = H. cuneatum.

Cumingii v.d.B. Ned. kr. Arch. **4**. 395. 1859 = H. dilatatum *emarginatum.

cuneatum Kze. Anal. 50. 1837. — Chile. Juan Fernandez.
H. Cumingii Pr. 1843.

cupressiforme Lab. Nov. Holl. pl. spec. **2**. 102 t. 250 f. 2. 1806 = H. tunbridgense.

daedaleum Bl. Enum. 222. 1828 = H. australe.

Darwinii Hk. fil.; v.d.B. Ned. Kr. Arch. **5**³. 157. 1863 — Amer. antarct.

decurrens (Jacq.) Sw. Schrad. Journ. **1800**². 99. 1801 — Ind. occ.
Adiantum Jacq. Coll. **2**. 103 t. 2 f. 1, 2. 1788; Trichomanes Poir. 1808.

decurrens β Sieberianum Pr. Hym. 32. 1843 = H. Kohautianum.

dejectum Bak. in Hk. Ic. pl. t. 1610. 1886 — Guiana.

Delavayi Christ, Bull. Soc. Fr. **52** Mém. I. 11. 1905 — Yunnan.

delicatissimum Fée, Cr. vasc. Br. **2**. 83 t. 105 f. 1. 1872-73 = H. lineare.

demissum (Forst.) Sw. Schrad. Journ. **1800**². 100. 1801. HB. 61. Chr. 16. NPfl 108 — Polynesia. N. Zealand. Malesia. — Trichomanes Forst. Prod. 85. 1786; Sphaerocionium Pr. 1843 — Hymenophyllum productum Kze. 1848 (Malesia) — H. megalocarpum Col. 1883.

densifolium Phil. Linn. **29**. 108. 1857 = Trichomanes caespitosum.

densum Wall. List n. 171. 1828 = H. exsertum et Trichomanes bipunctatum.

dentatum Cav. Descr. 276. 1802. NPfl. 112 — Chile. Chiloe.
Leptocionium v.d.B. 1859. Hymenophyllum Bridgesii Hk. 1844. HB. 70; Sphaerocionium Kl. 1844.

denticulatum Sw. Schrad. Journ. **1800**². 100. 1801. HB. 71. NPfl. 112 — Asia trop.
Trichomanes Poir. 1808; Didymoglossum Hassk. 1857-58; Leptocionium

v.d.B. 1859; Hym. Jav. 39 t. 29. Hymenophyllum humile Nees et Bl. 1823. — H. dichotomum Nees et Bl. 1823 (non Cav. 1802); Trichomanes Neesii Bl. 1828; Didymoglossum Pr. 1843; Hymenophyllum Hk. 1844. HB. 71 (pt.). Chr. 22 (pt.). NPfl. 112; Leptocionium v.d.B. 1859; Hym. Jav. 40 t. 30.

Deplanchei Mett.; Kuhn, Linn. **35**. 393. 1868. NPfl. 110 — N. Caledonia.

dichotomum Cav. Descr. 276. 1802. HB. 70. NPfl. 112 — Amer. antarct.

Myrmécostylum? Pr. 1843; Ptychophyllum Pr 1848; Leptocionium v.d.B. 1859. Hymenophyllum plicatum Klf. 1824; Ptychophyllum Pr. 1843 — ?Hymenophyllum nigricans Colla 1836.

dichotomum Nees et Bl. Nova Acta **11**. 127 t. 13 f. 4. 1823; Bl. Enum. 222. 1828 = H. denticulatum v. Neesii.

dicranotrichum (Pr.) Sadeb. Nat. Pfl. 1^4. 110. 1899 — Chile. Chiloe.

Leptocionium Pr. Hym. 27 t. 11 f. D. 1843; Hymenophyllum Chiloense Hk. 1844. HB. 63.

dilatatum (Forst.) Sw. Schrad. Journ. **1800^2**. 100. 1801. HB. 62, 463 Chr. 17. NPfl. 108 — Malesia. Austr. trop. Polynesia. N. Zealand.

Trichomanes Forst. Prod. 85. 1786; Sphaerocionium Pr. 1843; Diploöphyllum v.d.B. 1861 — Leptocionium sororium Pr. 1848; Hymenophyllum v.d.B. 1859 — H. eximium Kze. 1846; v.d.B. Hym. Jav. 57 t. 46. —* H. emarginatum Sw. 1801; Trichomanes Poir. 1808 — Hymenophyllum formosum Brack. 1854; v.d.B. Hym. Jav. 59 t. 47, 48 — Sphaerocionium macrocarpum Pr. 1843; Hymenophyllum v.d.B. 1859 — Sphaerocionium badium Pr. 1843 (non Hymenophyllum Hk. et Grev. 1828); Hymenophyllum Cumingii v.d.B. 1859 (non Pr. 1843).

dilatatum Bl. Enum. 221. 1828 = H. Junghuhnii.

dimidiatum Mett.; Kuhn, Linn. **35**. 393. 1868. (NPfl. 110) — N. Caledonia.

dimorphum Christ, Bull. Boiss. II. **4**. 941. 1904 — Costa Rica.

dipteroneuron A.Br.; Kze. Bot. Zeit. **1847**. 225 = H. fuscum.

discosum Christ, Bull. Boiss. **6**. 140. 1898 — Luzon.

divaricatum v.d.B. Ned. kr. Arch. 5^3. 190. 1863. (NPfl. 110) — Venezuela.

divaricatum Sod. Cr. vasc. quit. 29. 1893 = H. alveolatum.

diversilobum (Pr.) Fée, 11 mém. 118. 1866 — Ind. occ.

Sphaerocionium Pr. Hym. 34, 59. 1843.

Dregeanum Pr. Hym. 32, 52. 1843 = H. tunbridgense.

Durandii Christ, Bull. Boiss. **4**. 657. 1896 — Costa Rica.

Dusenii Christ apud Dusén, Svensk Exp. till Magelhl. **3**. 242. 1899 — Tierra del Fuego.

ectocarpon Fée, 11 mém. 115 t. 31 f. 1. 1866 — Ind. occ.

edentulum (v.d.B.) C. Chr. Ind. 360. 1905 — Assam. Borneo.

Leptocionium v.d.B. Ned. kr. Arch. 5^3. 148. 1863.

elasticum Bory; Willd. sp. **5**. 520. 1810 = H hygrometricum.

elatius Christ in Schwacke, Pl. Nov. Mineiras **2**. 13. 1900; Bull. Boiss. II. **2**. 319. 1902 = H. ciliatum var.

elegans Spr. Syst. **4**. 133. 1827 — Amer. trop. — H. lineare pt. HB. 66.

elegantissimum Fée, 11 mém. 118 t. 29 f. 2. 1866 — Guadeloupe. Jamaica.

elegantulum v.d.B. Ned. kr. Arch. **4**. 408. 1859. HB. 66. NPfl. 110 — Columbia. Ecuador.

emarginatum Sw. Schrad. Journ. **1800^2**. 101. 1801 = H. dilatatum*.

emersum Bak. Syn. ed. I. 457. 1868 = H. tenellum.

endiviaefolium Desv. Prod. 334. 1827 — Peru.

erecto-alatum Col. Tr. N. Zeal. Inst. **11**. 431. 1879 = H. demissum.
eriophorum v.d.B. Ned. kr. Arch. **5**^{3}. 180. 1863 = H trichophyllum.
erosum Bl. Enum: 221. 1828 = H. australe.
exiguum Bedd. Ferns br. Ind. t. 275. 1868 = Trichomanes.
eximium Kze. Bot. Zeit. **1846**. 478 = H. dilatatum.
exsertum Wall. List n. 170. 1828; Hk. sp. **1**. 109 t. 38 A. 1844. HB. 58. Chr. 18. NPfl. 109 — Himalaya-Ceylon. — H. macroglossum v.d.B. 1863.
falklandicum Bak. Syn. 68. 1867 — Falkland Ins.
farallonense Hier. Engl. Jahrb. **34**. 430. 1904 — Columbia. Ecuador.
fastigiosum Christ, Bull. Boiss. **7**. 3 1899 — Yunnan.
fecundum v.d.B. Ned. kr. Arch. **5**^{3}. 153. 1863 — Costa Rica.
feejeense Brack. Expl. Exp. **16**. 266 t. 37 f. 2. 1854 = H. multifidum.
Fendlerianum Sturm, Fl. bras. **1**^{2}. 291. 1859 — Venezuela-Bolivia Brasilia.
ferax v.d.B. Ned. kr. Arch. **4**. 392. 1859 — Venezuela-Ecuador.
ferrugineum Colla, Mem. Ac. Torino **39**. 30. 1836 — Chile. Juan Fernandez. N. Zealand. — H. subtilissimum Kze. 1837. HB. 64. NPfl. 110. H. Berteroi Hk. 1844 — H. Frankliniae Col. 1841; H. Franklinianum Col. 1844; H. aeruginosum *β* Hk. 1844.
filicula Bory; Willd. sp. **5**. 528. 1810 = Trichomanes bipunctatum.
fimbriatum J. Sm. JoB. **3**. 418. 1841; Hk. sp. **1**. 102 t. 36 C. 1844 = H. australe.
flabellatum Lab. Nov. Holl. pl. spec. **2**. 101 t. 250 f. 1. 1806. HB. 61. Chr. 17. NPfl. 109 — Austr. N. Zealand. Auckland Ins. Samoa.
H. nitens R.Br. 1810 — H. Hookeri Bory 1833.
flaccidum v.d.B. Ned. kr. Arch. **4** 400. 1859 = H. undulatum var.
flaccidum Bak. Syn. ed. I. 451. 1868 = H. khasianum.
flavo-aureum Bory, Bél. Voy. Bot. **2**. 84. 1833 = H. hygrometricum.
flexile Mak. Bot. Mag. Tokyo **13**. 45. 1899; Phan. et Pter. Jap. Ic. **1**. t. 18. 1899 — Japonia.
flexuosum A. Cunn. in Hk. Comp. Bot. Mag. **2**. 369. 1836 = H. australe var.
floribundum HBK. Nov. Gen. et Sp. **1**. 27. 1815 — Cumana.
formosum Brack. Expl. Exp. **16**. 268 t. 32 f. 3. 1854 = H. dilatatum*.
Francavillei v.d.B.Ned. kr. Arch. **4**. 411. 1859 — India occ.
Frankliniae Col. Tasm. Journ. **1**. 378. 1841 } = H. ferrugineum.
Franklinianum Col. Tasm. Journ. **2**. 183. 1844. } = H. ferrugineum.
fraternum Pr. Hym. 32, 54 t. 12 B. 1843 = H. polyanthos*.
fraternum Harr. Journ. Linn. Soc. **16**. 26. 1877 = H. Steerei.
Fraseri Mett.; Fourn. Mex. pl. **1**. 63. 1872 = H. ciliatum.
fuciforme Sw. Syn. 148. 1806. HB. 62. Chr. 18. NPfl. 109 — China austr. Juan Fernandez. — Leptocionium Pr. 1848; Hymenophyllum fucoides Cav. Descr. 275. 1802 (non Sw. 1801). H. semiteres Colla 1836.
fucoides Sw. Schrad. Journ. **1800**^{2}. 99. 1801. HB. 69. Chr. 22. NPfl. 110 — Amer. trop. Samoa. — Trichomanes Sw. Prod. 136. 1788; Leptocionium Pr. 1843 — Hymenophyllum spinulosum HBK. 1815 — H. pedicellatum Kze. 1847; Leptocionium Fourn. 1872 — Hymenophyllum cristatum Hk. et Grev. 1829; Sphaerocionium Pr. 1843; Dermatophlebium peruvianum Hk. et Grev. 1831 — H. blepharodes Pr. 1843 — H. podocarpum Fée 1869.
fucoides Cav. Descr. 275. 1802 = H. fuciforme.
fulvum v.d.B. Ned. kr. Arch. **5**^{3}. 196. 1863 = H. capillare.
fumarioides Willd. sp. **5**. 526. 1810 — Afr. austr. cum ins.
H. rarum auctt. quoad pl. Afr. (HB. 58. Chr. 16). H. capense Schrad. 1818.

H. Thunbergii Pr. 1843. H. Natalense, Zeyheri. tabulare v.d.B. 1859 — H. Limminghii v.d.B. 1863.

Funckii v.d.B. Ned. kr. Arch. 5³. 166. 1863 — Venezuela. Jamaica.

fuscum (Bl.) v.d.B. Hym. Jav. 62 t. 51-52. 1861. (NPfl. 109) — Malesia. Trichomanes Bl. Enum. 225. 1828; Amphipterum Pr. 1849; Didymoglossum Hassk. 1857-58 — Hymenophyllum dipteroneuron A.Br.; Kze. 1847.

fusugasugense Karst.; Sturm, Bot. Zeit. **1859**. 297. NPfl. 110 — Columbia. H. asterothrix Kze. 1844 (nomen).

Gardneri v.d.B. Ned. kr. Arch. **4**. 417. 1859 — Ceylon.

Gardnerianum Sturm, Fl. bras. 1². 297. 1859 = H. ciliatum var.

Glaziovii Bak. in Hk. Ic. pl. t. 1612. 1886 — Brasilia.

glebarium Christ in C. Chr. Ind. 362. 1905 — Tierra del Fuego. — H. caespitosum Christ apud Dusén, Svensk Exp. till Magelhl. **3**. 242. 1899 (non Gaud. 1825 nec Fée 1866).

Gollmeri v.d.B. Ned. kr. Arch. 5³. 151. 1863 — Caracas.

gracile Bory; Willd. sp. **5**. 527. 1810 = H. inaequale.

gratum Fée, 11 mém. 118 t. 30 f. 1, 2. 1866 — Jamaica. Guadelou (an = H. lineare?).

Grevilleanum Pr. Hym. 32. 1843 = H. polyanthos.

Gunnii v.d.B.; Bak. Syn. 463. 1874 = H. rarum var.

haematochroum Mett. (Sad. NPfl. 1⁴. 110. 1899) = H. Simonsianum.

helicoideum Sod. Cr. vasc. quit. 20. 1893 — Ecuador.

Henryi Bak. JoB. **1889**. 176 — China.

Himalaianum v.d.B. Ned. kr. Arch. 5³. 156. 1863 = H. Blumeanum.

hirsutum (L.) Sw. Schrad. Journ. **1800²**. 99. 1801. HB. 63. Chr. 19. NPfl. 110 — Amer. trop. Afr. austr. cum ins. — Trichomanes L. sp. **2**. 1098. 1753; Sphaerocionium Pr. 1843 — Hymenophyllum venustum Desv. 1827. H. Raddianum Müll. 1854 — H. angustum v.d.B. 1863 — H. latifrons v.d.B. 1859.

hirtellum Sw. Schrad. Journ. **1800²**. 101. 1801. HB. 64. NPfl. 110 — India occ. Amer. centr. Mexico. — Trichomanes Poir. 1808; Sphaerocionium Pr. 1843.

holochilum (v.d.B.) C. Chr. Ind. 362. 1905 — Java. Didymoglossum v.d.B. Pl. Jungh. **1**. 561. 1856; Leptocionium v.d.B. 1859; Hym. Jav. 45 t. 34 — Didymoglossum affine v.d.B. 1856; Leptocionium v.d.B. 1859; Hym. Jav. 45 t. 35; Hymenophyllum Racib. 1898 (non Brack. 1854). H. Smithii pt.. HB. 69.

Hookeri Bory, Bél. Voy. Bot. **2**. 84. 1833 = H. flabellatum var.

Houstonii Jenm. JoB. **1886**. 42 — Jamaica.

Humboldtianum Fourn. Ann. sc. nat. V. **18**. 265. 1873 — N. Caledonia.

humile Nees et Bl. Nova Acta **11**. 125 t. 13 f. 3. 1823 = H. denticulatum.

hygrometricum (Poir.) Desv. Prod. 333. 1827. Chr. 19. NPfl. 110 — Madagascar. Ins. Mascar. Ins. Sechellae. — Trichomanes Poir. Enc. **8**. 79. 1808; Hymenophyllum elasticum Bory; Willd. 1810. HB. 64; Sphaerocionium Pr. 1843 — Hymenophyllum flavo-aureum Bory 1833.

imbricatum Bl. Enum. 220. 1828 — Java.

imbricatum Col. Tasm. Journ. **2**. 187. 1844 = H. rarum.

imbricatum Kze.; Sturm, Fl. bras. 1². 287. 1859 = H. polyanthos.

inaequale (Poir.) Desv. Prod. 335. 1827 — Natal Madagascar. Ins. Mascarenae. Trichomanes Poir. Enc. **8**. 74. 1808; Hymenophyllum gracile Bory; Willd 1810. HB. 58. NPfl. 108; Sphaerocionium Pr. 1843.

inclinatum v.d.B. Pl. Jungh. **1**. 570. 1856 = H. dilatatum var. eximium.

infortunatum Bory. Dup. Voy. Bot. **2**. 284 t. 38 f. 3 1828 = H. capillaceum

integrum v.d.B. Pl. Jungh. **1**. 563. 1856 = H. Blumeanum.
intercalatum Christ, Bull. Boiss. II. **4**. 942. 1904 — Costa Rica.
intermedium Mett. (Sad. NPfl. 110 1899) = H. hirsutum.
interruptum Kze. Linn. **9**. 107. 1834; Anal. 48 t. 30. HB. 65. Chr. 20. NPfl. 110 — India occ. Mexico-Peru. Galapagos. — Sphaerocionium Pr. 1843 — Hymenophyllum aequabile Kze. 1847.
intricatum v d.B. Ned. kr. Arch. **5**³ 168. 1863 — Tasmania.
jalapense Schlecht et Cham. Linn. **5**. 619. 1830 = H polyanthos var.
Jamesoni Hk. sp. **1**. 96 t. 35 A. 1844. HB. 68. NPfl. 110 — Ecuador. Dermatophlebium Pr. 1849.
japonicum Miq. Ann. Lugd. Bat. **3**. 183. 1867. HB. ed. I. 451 — Japonia.
javanicum Spr. Syst. **4**. 132. 1827 = A. australe.
Junghuhnii v.d.B. Pl. Jungh. **1**. 570. 1856; Hym. Jav. 60 t. 49. 1861 — Java. H. dilatatum Bl. Enum. 221. 1828 (non Sw. 1801). HB. 62 (pt.)
kaieteurum Jenm. W. Ind. and Guiana Ferns 15. 1898 — Guiana.
Karstenianum Sturm, Bot. Zeit. **1859**. 298. (NPfl. 110) — Venezuela.
khasianum Bak. Syn. 464. 1874 — Khasia. Birma. Leptocionium flaccidum v.d.B. Ned. kr. Arch. **5**³. 149. 1863. Hymenophyllum Bak. 1868 (non v.d.B. 1859).
klabatense Christ, Verh. Nat. Ges. Basel **11**. 4. 1894 = H. multifidum.
Kohautianum Pr. Hym. 32, 56. 1843 — Ind. occ. Guiana. — H. decurrens β Sieberianum Pr. 1843; H. Martinicense v.d.B. 1859.
Krauseanum Phil. Linn. **30**. 208. 1860 = H. magellanicum var.
Kuhnii C. Chr. Ind. 363. 1905 — Kilimanjaro. H. Meyeri Kuhn in Engler, Hochgbfl. trop. Afr. 94. 1892 (non Pr. 1843).
Kurzii Prantl, Hym. 54. 1875 — Java.
laciniosum Christ, Bull. Boiss. II. **4**. 938. 1904 = H. nigrescens.
laeve Ham.: Wall. List n. 2198. 1829 (nomen) — Mauritius.
lanatum Fée, 11 mém. 116 t. 31 f. 3. 1866 — Guadeloupe. Jamaica.
lanceolatum Hk. et Arn. Bot. Beech. Voy. 109. 1832. HB. 64. NPfl. 110 — Ins. Hawaii.
latifrons v.d.B. Ned. kr. Arch. **4**. 407. 1859 = H. hirsutum var.
Lehmanni Hier.; Sadeb. Nat. Pfl. **1**⁴. 108. 1899 (nomen); Hier. Engl. Jahrb. **34**. 428. 1904 — Columbia.
leptodictyon K. Müll. Bot. Zeit. **1854**. 734 = H. dilatatum.
Levingei Clarke, Tr. Linn. Soc. II. Bot. **1**. 439 t. 49 f. 3. 1880 — Sikkim.
L'Herminieri Mett.: Kuhn, Linn. **35**. 392. 1868. HB. 463 — Guadeloupe.
Limminghii v.d.B. Ned. kr. Arch. **5**³. 151. 1863 = H. fumarioides.
Lindeni Hk. sp. **1**. 94 t. **34** C. 1844. HB. 65. NPfl. 110 — Venezuela. Ecuador. — H. terminale v.d.B. 1863.
Lindigii Mett. Ann. sc. nat. V. **2**. 197. 1864 — Columbia.
lindsaeoides Bak. Ann. Bot. **8**. 123. 1894 — Madagascar.
lineare Sw. Schrad. Journ. **1800**². 100. 1801. HB. 66 (pt.). Chr. 20. NPfl. 110 — Ind. occ. — Trichomanes Sw. Prod. 137. 1788; Didymoglossum Desv. 1827; Sphaerocionium Pr. 1843 — H. trifidum Hk. et Grev. 1830; Sphaerocionium Pr. 1843 — Hymenophyllum Moritzianum Sturm 1859 — H. Catherinae Hk. 1867. HB. 67.
lineare var. antillense Jenm. JoB. **1879**. 258 = H. antillense.
Liukiuense Christ, Bull. Boiss. II. **1**. 1021. 1901 = H. riukiuense.
lobato-alatum Kl. Linn. **20**. 438. 1847 = H. pyramidatum var.
lobato papillosum Sadeb. Nat Pfl. **1**⁴. 110. 1899 = T. lobato-alatum.

Lobbii Moore; v.d.B. Ned. Kr. Arch. 5^3. 176. 1863 — »Malesia«. Assam.
lophocarpum Col. Tr. N. Zeal. Inst. **17**. 255. 1885 — N. Zealand.
Lyallii Hk. fil. Fl. N. Zeal. **2**. 16. 1854 = Trichomanes.
macrocarpum v.d.B. Ned. kr. Arch. **4**. 395. 1859 = H. dilatatum*.
macroglossum v.d.B. Ned. kr. Arch. 5^3. 156. 1863 = H. exsertum.
macrothecum Fée, 11 mém. 115 t. 31 f. 2. 1866 — Guadeloupe.
magellanicum Willd.; Kze. Bot. Zeit. **1847**. 226. HB. 70. Chr. 22. NPfl. 112 — Brasilia austr. Amer. antarct. — Didymoglossum Desv. Prod. 331. 1827; Ptychophyllum Kl. 1844; Leptocionium v.d.B. 1859 — Hymenophyllum attenuatum Hk. 1844; Leptocionium v.d.B 1859 — Hymenophyllum Bibraianum Sturm 1853 — H. Krauseanum Phil. 1860.
Malingii (Hk.) Mett. Abh. Sachs. Ges. Wiss. **7**. 489. 1864; Bak. Syn. 66. 1867. Chr. 20. NPfl. 110 — N. Zealand. Tasmania. — Trichomanes Hk. Gard. [Ferns t. 64. 1862.
Mannianum Mett.; Kuhn, Fil. Afr. 40. 1868 = H. triangulare.
marginatum Hk. et Grev. Ic. Fil. t. 34. 1828. HB. 57. Chr. 15. NPfl. 108 — N. S. Wales. Tasmania. — Pachyloma v.d.B. 1861.
Martii Sturm, Fl. bras. 1^2. 287. 1859 = H. caudiculatum.
Martinicense v.d.B. Ned. kr. Arch. **4**. 389 1859 = H. Kohautianum.
Mathewsii v.d.B. Ned. kr. Arch. 5^3. 162. 1863 = H. polyanthos.
Mazei Fourn.; Christ, Engl. Jahrb. **24**. 85. 1897 — Guadeloupe.
megachilum Pr. Abh. böhm. Ges. V. 5. 337. 1848; Epim. 22 t. 8 B = H. peltatum.
megalocarpum Col. Tr. N. Zeal. Inst. **15**. 308. 1883 = H. demissum var.
melanocheilos Col. Tr. N. Zeal. Inst. **17**. 255. 1885 = H. Armstrongii.
Menziesii Pr. Hym. 31, 51. 1843 = H. peltatum.
Mettenii v.d.B. Ned. kr. Arch. 5^3. 171. 1863 = H. peltatum.
Meyeri Pr. Hym. 31, 50. 1843 = H. peltatum.
Meyeri Kuhn in Engler, Hochgbfl. trop. Afr. 94. 1892 = H. Kuhnii.
micans Christ. Bull. Boiss. II. **5**. 260. 1905 — Costa Rica. Brasilia.
H. nitens Werckle; Christ. Bull. Boiss. II. **4**. 940. 1904 (non R. Br. 1810).
micranthum v.d.B. Pl. Jungh. **1**. 566. 1856; Ned. kr. Arch. **4**. 399. 1859 = H. australe.
microcarpon Fée, Cr. vasc. Br. **1**. 245 t. 69 f. 3. 1869 = H. ciliatum.
microcarpum Desv. Prod. 333. 1827. HB. 65. Chr. 18. NPfl 110 — Amer. trop. — H. Beyrichianum Kze. 1834 — H. Organense Hk. 1844 — H. pteropodum v.d.B. 1863.
microphyllum Mett. Ann. sc. nat. V. **2**. 195. 1864. HB. 463 — Columbia.
microsorum v.d.B. Ned. kr. Arch. 5^3. 155. 1863. HB. 59. NPfl. 108 — China. [Sikkim.
millefolium Schlecht. et Cham. Linn. **5**. 620. 1830 — Mexico.
minimum A. Rich. Fl. N. Zél. 91 t. 14 f. 2. 1832. HB. 464. NPfl. 110 — N. Zealand. Ins. Auckland.
mnioides Bak. Syn. 57. 1867; Mett.; Kuhn, Linn. **35**. 390. 1868 — N. Caledonia H. bifarium Mett.; Sadeb. 1899. NPfl. 108.
modestum v.d.B. Ned. kr. Arch. 5^3. 163. 1863 — Ins. Philipp.
montanum Kirk. Tr. N. Zeal. Inst. **10**. 394 t. 21 B. 1878 — N. Zealand.
Moorei Bak. Syn. 464. 1874 — Ins. Lord Howe.
Moritzianum Sturm, Fl. bras. 1^2. 295. 1859 — H. lineare.
multifidum (Forst.) Sw. Schrad. Journ. 1800^2. 102. 1801. HB. 69. Chr. 22. NPfl. 110 — N. Zealand N. S. Wales Polynesia. Celebes.
Trichomanes Forst. Prod. 85. 1786; Davallia Spr. 1799. Hymenophyllum Feejeense Brack. 1854.

myriocarpum Hk. sp. **1**. 106 t. 37 D. 1844. HB. 59. NPfl. 108 — Mexico-Peru. — Sphaerocionium Kl. 1844.
nanum Sod. Cr. vasc. quit. 28. 1893 — Ecuador.
Natalense v.d.B. Ned. kr. Arch. **4**. 386. 1859 = H. fumarioides.
Neesii Hk. sp. **1**. 99. 1844. HB. 71 — H. denticulatum et aculeatum.
nigrescens Liebm. Vid. Selsk. Skr. V. **1**. 292 (seors. 140). 1849 — Mexico-Peru. Sphaerocionium nigricans Pr.; Kl. Linn. **18**. 536. 1844; Hymenophyllum Kze. 1847 (non Colla 1836). H. laciniosum Christ 1904.
nigricans Colla, Mem. Ac. Torino **39**. 32 t. 62. 1836 = H. dichotomum.
nigricans Kze. Bot. Zeit. **1847**. 244 = H. nigrescens.
nitens R.Br. Prod. Fl. N. Holl. 159. 1810 = H. flabellatum.
nitens Werckle; Christ, Bull. Boiss. II. **4**. 940. 1904 = H. micans.
notabile Fée, Cr. vasc. Br. **1**.193 t. 69 f. 2. 1869 — Brasilia.
nudum (Poir.) Desv. Prod. 332. 1827 — Guadeloupe. Trichomanes Poir. Enc. **8**. 73. 1808.
obtusum Hk. et Arn. Bot. Beech. Voy. 109. 1832. HB. 63. NPfl. 110 — Ins. Hawaii. N. Guinea.
oligocarpum Col. Tr. N. Zeal. Inst. **31**. 264. 1899 — N. Zealand.
oligosorum Mak. Bot. Mag. Tokyo **13**. 44. 1899; Phan. et Pter. Jap. Ic. **1**. t. 19. 1899 — Korea. Japonia.
ooides F. Muell. et Bak. JoB. **1890**. 105 — N. Guinea.
Orbignianum v.d.B. Ned. kr. Arch. **5**^{3}. 191. 1863 — Bolivia.
Organense Hk. sp. **1**. 90 t. 32 B. 1844 = H. microcarpum.
Osmundoides v.d.B. Ned. kr. Arch. **5**^{3}. 164. 1863 = H. Blumeanum.
oxyodon Bak. JoB. **1890**. 262. NPfl. 110 — Tonkin.
pachydermicum Ces. Atti Ac. Napoli **7**^{8}. 8. 1876 — Borneo.
palmatum v.d.B. Ned. kr. Arch. **4**. 385. 1859 = Trichomanes sibthorpioides.
paniculiflorum Pr. Hym. 32, 55. 1843; v.d.B. Hym. Jav. 49 t. 39 — Malesia. [Japonia.
pannosum Christ, Bull. Boiss. II. **5**. 250. 1905 — Costa Rica.
parvifolium Bak. Journ. Linn. Soc. **9**. 348 t. 8 f. E. 1866. HB. 56 — Moulmein. — Trichomanes microphyllum Kuhn 1868.
parvulum C. Chr. Ind. 365. 1905 — Chile. H. terminale Phil. Linn. **33**. 306. 1865 (non v.d.B. 1863).
pastoense Hk. Syn. 67. 1867 = H. Ruizianum.
paucicarpum Jenm. Bull. Dep. Jam. **1**. n. 18. 4. 1890 — Jamaica.
pectinatum Cav. Descr. 275. 1802. HB. 68. Chr. 21. NPfl. 110 — Chile austr. Juan Fernandez.
pectinatum Nees et Bl. Nova Acta **11**. 124 t. 12 f. 5. 1823 = H. Blumeanum.
pedicellatum Kze.; Kl. Linn. **20**. 439. 1847 = H. fucoides.
pedicularifolium Ces. Rend. Ac. Napoli **16**. 24, 28. 1877 — N. Guinea.
peltatum (Poir.) Desv. Prod. 333. 1827 — Eur. atlant. et. merid. Ins. Atlant. Afr. austr. c. ins. Amer. austr. temp. et antarct. Ins. antarct. Tasmania. N. Zealand. — Trichomanes Poir. Enc. **8**. 76. 1808; Hymenophyllum tunbridgense β HB. 57. NPfl. 110. H. unilaterale Bory; Willd. 1810. (Chr. 21.) (Ins. Mascar.) — H. Wilsoni Hk. 1830 (Eur. atlant.) — H. Meyeri Pr. 1843 (Afr. austr.) — H. Menziesii Pr. 1843 (Amer. antarct. Ins. antarct.) — H. Mettenii v.d.B. 1863 (Chile) — H. megachilum Pr. 1848 (Brasilia) — H. zeelandicum v.d.B. 1863 (N. Zealand.)
pendulum Bory, Bél. Voy. Bot. **2**. 81 t. 8 f. 2. 1833 = H. capillare.
pendulum Sod. Cr. vasc. quit. 31. 1893 = H. Sodiroi.

peruvianum Hk. et Grev. Ic. Fil. t. 208. 1831 = H. fucoides.

physocarpum Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 35 t. 1 A. 1905 — N. Guinea.

platylobum v.d.B. Ned. kr. Arch. **5**^{3}. 189. 1863 = H. valvatum var.

plicatum Klf. Enum. 268. 1824 = H. dichotomum.

Plumieri Hk. et Grev. Ic. Fil. t. 123. 1829 = H. ciliatum.

plumosum Klf. Enum. 267. 1824 — Amer. trop.
Sphaerocionium aureum Pr. 1843.

podocarpon Fée, Cr. vasc. Br. **1**. 196 t. 71 f. 3 1869 = H. fucoides.

Poeppigianum Pr. Hym. 32, 54. 1843 = H. polyanthos var.

polyanthos Sw. Schrad. Journ. **1800**^{2}. 102. 1801. HB. 60 (pt.). Chr. 16 (pt.). NPfl. 108 (pt.) — Amer. trop.-antarct. Afr. occ. trop. Madagascar. Mauritius. ? Japonia. ? Samoa. — Trichomanes Sw. Prod. 137. 1788 — Hymenophyllum abietinum Hk. et Grev. 1829; Sphaerocionium Pr. 1843 — Hymenophyllum protrusum Hk. 1844; Leptocionium Pr. 1848 — Hymenophyllum Poeppigianum Kl. 1843 — H. imbricatum Sturm 1859 (non Bl. 1828) — H. Schomburgkii Pr. 1843 — Sphaerocionium Breutelii Pr. 1848; Hymenophyllum v.d.B. 1863 — H. Sturmii v.d.B. 1863 — H. Matthewsii v.d.B. 1863 — H. pusillum Schott; Sturm 1859 — H. jalapense Schlecht. et Cham. 1830 — H. viridissimum Fée 1869.
—* Trichomanes clavatum Sw. 1788; Hymenophyllum Sw. 1801; Myrmecostylum Pr. 1843 — Hymenophyllum fraternum Pr. 1843.

polyanthos auctt. quoad pl. asiat. trop. = H. Blumeanum.

polyanthos auctt. quoad pl. Novae Zealandiae = H. sanguinolentum.

polyanthos Hk. in Night. Ocean. Sketch. App. 132. 1835 = Trichomanes Bauerianum var.

polyanthos Bedd. Ferns S Ind. t. 267. 1863-65 = H. tenellum.

polycarpum Mett.; Kuhn, Linn. **35**. 391. 1868 = H. crispum var.

polychilum Col. Tr. N. Zeal. Inst. **24**. 395. 1892 — N. Zealand.

polyodon Bak. Journ. Linn. Soc. **15**. 105. 1876 — Ins. Admiralitatis.

Pooli Bak. Journ. Lin. Soc. **15**. 413. 1876; Ic. pl. t. 1609 — Madagascar.

portoricense Kuhn, Engl. Jahrb. **24**. 86. 1897 = H. lineare.

praetervisum Christ, Engl. Jahrb. **23**. 338. 1896; Farnkr. 21 — Borneo. Samoa. — Trichomanes denticulatum Bak. Syn. 82. 1867. NPfl. 107 (non Hymenophyllum Sw. 1801).

prionema Kze.; Sturm, Fl. bras. **1**^{2}. 284. 1859 — Brasilia.

procerum v.d.B. Ned. kr. Arch. **4**. 409. 1859 = H. trichophyllum.

producens Fée, Cr. vasc. Br. **1**. 196 t. 71 f. 4. 1869 = H. valvatum.

productum Kze. Bot. Zeit. **1848**. 305 = H. demissum var.

productum Sturm, Enum. pl. vasc. cr. chil. 43. 1858 = H. caudiculatum var.

protrusum Hk. sp. **1**. 104 t. 37 B. 1844 = H. polyanthos var.

pteropodum v.d.B. Ned. kr. Arch. **5**^{3}. 187. 1863 = H. microcarpum.

puellum Ces. Rend. Ac. Napoli **16**. 24, 28. 1877 = ? Trichomanes Bauerianum.

pulchellum Schlecht. et Cham. Linn. **5**. 618. 1830 = H. sericeum.

pulcherrimum Col. Tasm. Journ. **2**. 185. 1844; Hk. sp **1**. 103 t. 37 A. 1844. HB. 62. Chr. 18. NPfl. 109 — N. Zealand.

pumilum C. Moore; Bak. Syn. 464. 1874 = N. S. Wales.

pusillum Schott; Sturm, Fl. bras. **1**^{2}. 289. 1859 = H. polyanthos.

pusillum Col. Tr. N. Zeal. Inst. **12**. 365. 1880 = H tunbridgense.

pycnocarpum v.d.B. Pl. Jungh. **1**. 564. 1856 = H. Blumeanum.

pygmaeum Col Tr. N. Zeal. Inst. **13**. 276. 1881 = H. tunbridgense.
pyramidatum Desv. Prod. 332. 1827. NPfl. 110 — Andes.
— H. lobato-alatum Kl. 1847 (lobato-papillosum Sad. NPfl. 110).
pyriforme v.d.B. Ned. kr. Arch. **5**[3]. 173. 1863 = H. bivalve.
quadrifidum Phil. Linn. **30**. 208. 1860 – Chile.
Raddianum K. Müll. Bot. Zeit. **1854**. 723 = H. hirsutum.
ramosissimum Ham.; Don, Prod. Fl. Nepal. 12. 1825 = Odontosoria chinensis.
rarum R. Br. Prod. Fl. N. Holl. 159. 1810. HB. 58 (pt.). Chr. 16 (pt.). NPfl. 108 (pt.) — Tasmania. N. Zealand. Auckland. ? Perak — H. semibivalve Hk. et Grev. 1829 — H. imbricatum Col. 1844 (non Bl. 1828).
rarum auctt. quoad pl. afric. = H. fumarioides.
recurvum Gaud. Freyc. Voy. Bot. 376. 1827. HB. 61. NPfl. 108 — Ins. Hawaii.
refrondescens Sod. Rec. 2. 1883 = H. sericeum var.
Reinwardtii v.d.B. Ned. kr. Arch. **4**. 399. 1859; Hym. Jav. 52 t. 42 — Java. [Sumatra.
remotum v.d.B. Ned. kr. Arch. **4**. 413. 1859 = H. ciliatum.
reniforme Hk. sp. **1**. 110 t. 38 C. 1844. HB. 59. NPfl. 109 — Ecuador-Peru.
revolutum Col. Tasm. Journ. **2**. 186. 1844 = H tunbridgense.
ricciaefolium Bory; Willd. sp. **5**. 531. 1810 = H. tenellum.
Rimbachii Sod. Cr. vasc. quit. 633. 1893 — Ecuador.
ringens Christ, Ann. Jard. Buit. II. **4**. 34. 1904 — Celebes.
riukiuense Christ, Ann. Cons. Jard. bot. Genève **4**. 208. 1900 — Liu Kiu.
rufescens Kirk, Tr. N. Zeal. Inst. **11**. 457 t. 19 A. 1879 — N. Zealand.
rufum Fée, Cr. vasc. Br. **1**. 198 t. 70 f. 4. 1869 — Brasilia. Costa Rica.
Ruizianum (Kl.) Kze. Bot. Zeit. **1847**. 199. NPfl. 110 — Venezuela-Peru.
Sphaerocionium Kl. Linn. **18**. 535. 1844. Hymenophyllum pastoense Hk. [1867. HB. 67.
rupestre Raddi, Pl. Bras. **1**. 67 t. 80. 1825 = Trichomanes.
sabinifolium Bak. Syn. 71. 1867. Chr. 32. NPfl. 112 — Java. Sumatra.
salakense Racib. Pterid. Buit. 18. 1898. NPfl. 109 — Java.
samoense Bak. JoB. **1876**. 10 — Samoa.
sanguinolentum (Forst.) Sw. Schrad. Journ. **1800**[2]. 101. 1801 — N. Zealand. Trichomanes Forst. Prod. 84. 1786; Sphaerocionium Pr. 1843; Mecodium Pr. 1849.
scabrum A. Rich. Fl. N. Zél. 90 t. 14 f. 1. 1832. HB. 61. NPfl. 109 — N. Zealand. — Sphaerocionium Pr. 1843; Diploöphyllum v.d.B. 1861 — Sphaerocionium glanduliferum Pr. 1849.
Schiedeanum K. Müll. Bot. Zeit. **1854**. 720 = H. ciliatum vel H. lineare.
Schomburgkii Pr. Hym. 32. 1843 = H. polyanthos.
secundum Hk. et Grev. Ic. Fil. t. 133. 1829. HB. 68. Chr. 22. NPfl. 110 — Amer. antarct. — H. serra Pr. 1843.
semibivaloe Hk. et Grev. Ic. Fil. t. 83. 1829 = H. rarum.
semiteres Colla. Mem. Ac. Torino **39**. 32 t. 61. 1836 = H. fuciforme.
sericeum Sw. Schrad. Journ. **1800**[2]. 99. 1801. HB. 66 (pt.). Chr. 19. NPfl. 110 (pt.) — Amer. trop. — Trichomanes Sw. Prod. 136. 1788; Sphaerocionium Pr. 1843 — Hymenophyllum pulchellum Schlecht. et Cham. 1830; Sphaerocionium Pr. 1843 — Hymenophyllum speciosum v.d.B. 1863 — H. refrondescens Sod. 1883.
serpens Wall. List n. 173. 1828 = H. australe.
serra Pr. Hym. 32, 53 t. 11 B. 1843 = H. secundum.
serrulatum (Pr.) C. Chr. Ind. 367. 1905 — Ins. Philipp. N. Guinea.
Didymoglossum Pr. Hym. 23, 48. 1843; Leptocionium v.d.B. 1859. Hymen-

ophyllum Smithii Hk. 1844. HB. 69 (pt.). (NPfl. 110 pt.) — Leptocionium Preslii v.d.B. 1859.
seselifolium Pr. Hym. 38, 52. 1843 — Chile.
Leptocionium v.d.B. 1859.
sibthorpioides Mett.; Kuhn, Fil. Afr. 41. 1868 = Trichomanes.
Sieberi v.d.B. Ned. kr. Arch. **4**. 414. 1859 = H. ciliatum.
siliquosum Christ, Bull. Boiss. II. **4**. 939. 1904 — Costa Rica.
Silveirae Christ in Schwacke, Pl. Nov. Mineiras **2**. 14. 1900; Bull. Boiss. II. **2**. 320. 1902 — Brasilia.
Simonsianum Hk. 2 Cent. t. 13. 1860. HB. 68. NPfl. 110 — India bor.
Didymoglossum v.d.B. 1863.
Smithii Hk. sp. **1**. 97 t. 35 B. 1844 = H. serrulatum.
Smithii Hk. & Bak. Syn. 69. 1867 = H. serrulatum et holochilum.
Sodiroi C. Chr. Ind. 368. 1905 — Ecuador.
H. pendulum Sod. Cr. vasc. quit. 31. 1893 (non Bory 1833).
sororium v.d B. Ned. kr. Arch. **4**. 395. 1859 = H. dilatatum.
spathulatum Col. Tasm. Journ. **2**. 184. 1844 = H. bivalve.
speciosum v.d.B. Ned. kr. Arch. **5**^{3}. 181. 1859 = H. sericeum var.
spectabile Mett.; Kuhn, Linn. **35**. 392. 1868 — Bolivia.
spectabile Moritz; (Hk. Bak. Syn. 65. 1867) = H. Lindeni.
Sphaerocarpum v.d.B. Ned. kr. Arch. **5**^{3}. 167. 1863 = H. Blumeanum.
spinulosum HBK. Nov. Gen. et Sp. **1**. 26. 1815 = H. fucoides.
splendidum v.d.B. Ned. kr. Arch. **5**^{3}. 192. 1863 = H. ciliatum var.
Sprucei Bak. Syn. 65. 1867 = Peru.
Steerei C. Chr. Ind. 368. 1905 — Ins. Philipp.
H. fraternum Harr. Journ. Linn. Soc. **16**. 26. 1876 (non Pr. 1843).
streptophyllum Fourn. Ann. sc. nat. V. **18**. 266. 1873 — N. Caledonia.
Sturmii v.d.B. Ned. kr. Arch. **5**^{3}. 152. 1863 = H. polyanthos.
Sturmii Mett. (Sadeb. NPfl. 110. 1899) = H. sericeum an H. Karstenianum?
subdeltoideum Christ, Bull. Boiss. II. **2**. 322. 1902 = H. polyanthos.
subdemissum Christ. Bull. Boiss. **6**. 140. 1898 — Luzon.
subflabellatum Ces. Atti Ac. Napoli **7**^{8}. 8. 1876 — Borneo.
subrigidum Christ, Bull. Boiss. II. **5**. 260. 1905 — Costa Rica
H. atrovirens Christ, Bull. Boiss. II. **4**. 941 1904 (non Col. 1844 nec Fée [1866.
subtilissimum Kze. Anal. 49. 1837 = H. ferrugineum.
Surinamense v.d.B. Ned. kr. Arch. **4**. 414. 1859 = H. ciliatum.
tabulare v.d.B. Ned. kr. Arch. **4**. 397. 1859 = H. fumarioides.
Tasmannicum v.d.B. Ned. kr. Arch. **4**. 399. 1859 = H. australe var. atrovirens.
Telfairianum Wall. List n. 168. 1828 = H. hygrometricum.
tenellum Don, Prod. Fl. Nepal. 12. 1825 — Nepal.
tenellum (Jacq.) Kuhn, Fil. Afr. 42. 1868. HB. 57 — Madagascar. Ins. Mascarenae. Ceylon. — Adiantum Jacq. Coll. **3**. 287 t. 21 f. 3. 1789; Hymenophyllum Bory; Willd. 1810. NPfl. 108. Sphaerocionium Pr. 1843 — Hymenophyllum emersum Bak. 1868. NPfl. 108.
tenerrimum v.d.B. Ned. kr. Arch. **5**^{3}. 185. 1863. HB. 68 — Peru.
tenerum v.d.B. Ned. kr. Arch. **5**^{3}. 167. 1863 — Ecuador.
terminale v.d.B. Ned. kr. Arch. **5**^{3}. 186. 1863 = H. Lindeni.
terminale Phil. Linn. **33**. 306. 1865 = H. parvulum.
thuidium Harr. Journ. Linn. Soc. **16**. 25. 1877 — Ins. Philipp.
Thunbergii Eckl.; Pr. Hym. 32. 1843 = H. fumarioides.

tomentosum Kze. Linn. 9. 107. 1834. NPfl. 110 — Andes.
Sphaerocionium Pr. 1843; Dermatophlebium Pr. 1849.
torquescens v.d.B Ned. kr. Arch. 5³. 178. 1863 — Martinique.
tortuosum Hk. et Grev. Ic. Fil. t. 129. 1829. HB. 70. NPfl. 112 — Amer. antarct. Juan Fernandez. — Myrmecostylum Pr. 1843; Leptocionium v.d.B. 1859.
trapezoidale Liebm. Vid. Selsk. Skr. V. 1. 293 (seors. 141). 1849 — Mexico. Columbia.
Treubii Racib. Pterid. Buit. 15. 1898; Nat. Tijds. Ned. Ind. **59** t. 3. NPfl. 109 [— Java.
Trianae Hier. Engl. Jahrb. **34**. 429. 1904 — Columbia.
triangulare Bak. Syn. 69. 1867; Ic. pl. t. 1613 — Fernando Po.
H. Mannianum Mett.; Kuhn 1868. NPfl. 110.
trichocaulon Phil. Anal. Univ. Chil. **94**. 360. 1896 — Chile.
trichomanoides v.d.B. Ned. kr. Arch. 5³. 158. 1863 — Columbia.
trichomanoides Bail. Rep. Govt. Sci. Exp. Bellenden-Ker 74. 1889; Queensl. Fl. Suppl. III. 90. 1890 — Queensland.
trichophyllum HBK. Nov. Gen. et Sp. **1**. 27. 1815 — Andes.
H. procerum v.d.B. 1859 — H. eriophorum v.d.B. 1863.
trifidum Hk. et Grev. Ic. Fil. t. 196. 1830 = H. lineare.
truncatum Col. Tr. N. Zeal. Inst. **23**. 390 1891 — N. Zealand.
tunbridgense (L.) Sm. in Sowerby, Engl. Bot. t. 162. 1794. HB. 67 (excl. 3). Chr. 20 (pt.). NPfl. 110 (pt.) — Europa atlant. et medit. Ins. Atlant. Africa austr. Austr. temp. Tasmania. N. Zealand. Jamaica. Venezuela. Chile.
Trichomanes tunbrigense L. sp. **2**. 1098. 1753. Hymenophyllum alatum Willd. 1810 — H. cupressiforme Lab. 1806 (Austr. Tasmania) — H. Dregeanum Pr. 1843 (Afr. austr.) — H. antarcticum Pr. 1843 (Austr. Tasmania) — H. revolutum Col. 1844 (N. Zealand) — H. pusillum Col. 1880 (N Zealand) — H. pygmaeum Col. 1881 (N. Zealand) — H. asperulum Kze. 1834 (Chile).
Ulei Christ et Giesenh. Flora **1899**. 85 f. 6-7. NPfl. 110 — Brasilia.
undulatum Sw. Schrad. Journ. **1800²**. 101. 1801. HB. 59. NPfl. 108 — Amer. trop.
Trichomanes Sw. Prod. 137. 1788; Sphaerocionium Pr. 1843 — Hymenophyllum flaccidum v.d.B. 1859.
uniforme Phil. Bot. Zeit. **1856**. 631 (err. typ.) = H. reniforme.
unilaterale Bory; Willd. sp. **5**. 521. 1810 = H. peltatum.
vacillans Christ in Schwacke, Pl. Nov. Mineiras **2**. 14. 1900; Bull. Boiss. II. **2**. 322. 1902 — Brasilia.
valvatum Hk. et Grev. Ic. Fil. t. 219. 1831. HB. 65. Chr. 19. NPfl. 110 — Amer. trop. — H. platylobum v.d.B. 1863 — H. producens Fée 1869.
venustum Desv. Prod. 332. 1827 = H. hirsutum.
vestitum v.d.B. Ned. kr. Arch. 5³. 193. 1863 = H. ciliatum.
villosum Col. Tasm. Journ. **2**. 185. 1844; Kirk. Tr. N. Zeal. Inst. **10**. 395. 1878 — N. Zealand.
vincentinum Bak. Ann. Bot. **5**. 164 t. 10. 1891 — St. Vincent. Puerto Rico.
violaceum Meyen. Ned. kr. Arch. 5³. 147. 1863 (syn.) — Ins. Philipp.
Leptocionium v.d.B. Ned. kr. Arch. 5³. 147. 1863.
viridissimum Fée, Cr. vasc. Br. **1**. 194 t. 49 f. 3. 1869 = H. polyanthos.
Wercklei Christ, Bull. Boiss. II. **4**. 940. 1904 — Amer. trop.
Wilsoni Hk. Brit. Fl. *I*. 446. 1830 = H. peltatum.
Wilsoni Fée, Cr. vasc. Br. **1**. 193. 1869 = H. lineare.
Wrightii v.d.B. Ned. kr. Arch. **4**. 391. 1859. (NPfl. 58) — Japonia.

Zeelandicum v.d.B. Ned. kr. Arch. 5[3]. 175. 1863 = H. peltatum.
Zeyheri v.d.B. Ned. kr. Arch. 4. 388. 1859 = H. fumarioides.
Zollingerianum Kze. Bot. Zeit. **1848**. 305; v.d.B. Hym. Jav. 61 t. 50. HB. 62. Chr. 18. NPfl. 109 — Java. — Didymoglossum Zollingeri Hassk. 1857-58.
HYMENOPTERIS Klf.; Link, Fil. sp. 48, 73. 1841 (nomen) — ?
HYMENOSTACHYS Bory, Dict. class. d'hist. nat. **6**. 588; **8**. 462. 1825 = **Trichomanes.**
diversifrons Bory, Dict. class. **8**. 462. 1825; **17** t. 42 f. 2.
elegans Pr. Hym. 11. 1843 = T. diversifrons.
osmundoides Pr. Hym. 11. 1843.
Hymenostegia J. Smith, Lond. Journ. Bot. **1**. 666. 1842 *(Alsophila §)* = Alsophila.
HYMENOTOMIA Gaudichaud. Freyc. Voy. Bot. 379. 1827 = **Lindsaya.**
HYPOCHLAMYS Fée, Gen. Fil. 200. 1850-52 = **Athyrium.**
pectinata Fée, Gen. 200 t. 17 C. f. 2. 1850-52 = A. costale.
Sorgonensis Fée, Gen. 200. 1850-52 = Diplazium sorzogonense.
squamulosa Fée, Gen. 201. 1850-52 = A. costale.
Tussaci Fée, Gen. 201. 1850-52 = A. costale.
HYPODEMATIUM Kunze, Flora **1833**[2]. 690 = **Dryopteris.**
crenatum Kuhn, v. Deck. Reis. **3**[3]. Bot. 37. 1879.
cystopteroides Kuhn, Forschr. Gazelle **4**. Farne 8. 1889 = Athyrium c.
nivale Fée, Gen. 298. 1850-52 = D. rigida *australis.
onustum Kze. Flora **1833**[2]. 690 = D. crenata.
phegopteroideum Kuhn, Forschr. Gazelle **4**. Farne 8. 1889.
pusillum Kuhn, Forschr. Gazelle **4**. Farne 8. 1889.
Rüppellianum Kze. Farnkr. **1**. 41 t. 21. 1840 = D. crenata.
HYPODERRIS R. Brown in Wallich, Pl. Asiat. Rar. **1**. 16. 1830; J. Smith in Hook. Gen. Fil. t. 1. 1838 — (Polypodiaceae Gen. **4**).
adnata Fourn. Bull. Soc. Fr. **19**. 260. 1872 — Nicaragua.
Brownii J. Sm. in Hk. Gen. t. 1. 1838. HB. 46. Chr. 283. NPfl. 163 — Ind. occ. Haïti. Puerto Rico. Grenada. Trinidad. — Woodsia Mett. 1856.
marginalis Fourn. Bull. Soc. Fr. **19**. 260. 1872 = H. Seemanni.
Seemanni Prentice, JoB. **1869**. 240. HB. 460. Chr. 284. NPfl. 163 — Nicaragua. H. marginalis Fourn. 1872.
HYPOLEPIS Bernhardi, Schrad. neu. Journ. **1**[2]. 34. 1806 — (Polypodiaceae Gen. **73**)
aculeata J. Sm. JoB. **4**. 157. 1841 = H. repens.
alpina (Bl.) Hk. sp. **2**. 63. 1852 — Java. — Cheilanthes Bl. Enum. 138. 1828.
amaurorachis Hk. sp. **2**. 62. 1852 = Dryopteris punctata.
anthriscifolia Pr. Tent. 162. 1836 = H. sparsisora.
aquilinaris (Fée) Christ, Bull. Boiss. II. **2**. 636. 1902 — Brasilia. Cheilanthes Fée, Cr. vasc. Br. **2**. 37 t. 91 f. 1. 1869.
aspera Pr. Tent. 162. 1836 = H. sparsisora var.
aspidioides Christ, Engl. Jahrb. **23**. 343. 1896 — Samoa.
Bergiana (Schlecht.) Hk. sp. **2**. 67. 1852. HB. 130. Chr. 279 — Afr. austr. Cheilanthes Schlecht. (Adumbr. 51 t. 31 ined.); Kze. Linn. **10**. 541. 1836.
bogotensis Karst.; Mett. Ann. sc. nat. V. **2**. 238. 1864 (nomen); Karst. Fl. Col. **2**. 91 t. 147. 1864-69 — Columbia.
Boryana Pr. Tent. 162. 1836 = H. tenuifolia.
brasiliana (Pr.) Kuhn, Chaetopt. 347. 1882 — Brasilia. Aspidium Pr. Del. Prag. **1**. 176. 1822; Cystopteris? Pr. 1836. (an adhuc Cheilanthes rigescens Kze. 1834?).

californica (Nutt.) Hk. sp. 2. 71 t. 88 A. 1852. HB. 131. NPfl. 279 — California. Aspidotis Nutt. msc.; Cheilanthes Mett. 1859. Chr. 146 — C. amoena A. A. Eat. 1877.

Capensis Hk. sp. 2. 71 t. 77 C. 1852 = Adiantopsis.

chilensis Fée, 8 mém. 76. 1857 = Dryopteris punctata.

commutata Fée, Gen. 146. 1850-52 = H. anthriscifolia.

coniifolia Pr. Tent. 162. 1836 = Adiantopsis chlorophylla.

delicatula Fée, 11 mém. 28. 1866 = H. repens.

dicksonioides (Bory) Hk. sp. 2. 61. 1852 — Polynesia. N. Zealand. Australia. Adiantum Bory hb.; Cheilanthes Endl. Prod. Fl. Norf. 15. 1833. Hypolepis Endlicheriana Pr. 1836.

dicksonioides Fée. 11 mém. t. 21. f. 1. 1866 (delicatula in text.) = H. repens.

dissecta Brack. Expl. Exp. 16. 89. 1854 = H. tenuifolia.

distans (Col.) Hk. sp. 2. 70 t. 95 C. 1852. HB. 129. NPfl. 277 — N. Zealand. Cheilanthes Col. msc.

Eckloniana Pr. Tent. 162. 1836 = H. sparsisora.

elata Hk. sp. 2. 67. 1852 = H. Bergiana.

elegans Carr. in Seem. Fl. Vit. 347. 1873 — Fiji. Aneiteum. Ins. Lord Howe.

Endlicheriana Pr. Tent. 162. 1836 = H. dicksonioides.

flexuosa Sod. Cr. vasc. quit. 634. 1893 — Ecuador. Columbia.

Gardneri Hk. sp. 2. 74 t. 92 B. 1852 = Adiantopsis monticola.

guianensis Kl. Linn. 20. 339. 1847 — Guiana.

Helenensis Fée, Gen. 147. 1850-52 = Dryopteris punctata.

hostilis (Kze.) Pr. Tent. 162. 1836. HB. 130. NPfl. 278 — Jamaica. Columbia-Peru. Brasilia. — Cheilanthes Kze. Linn. 9. 86. 1834.

incisa (Kze.) C. Chr. Ind. 371. 1905 — Brasilia. — Cheilanthes Kze.; Mett. Cheil. 44 n. 65 t. 3 f. 28-31. 1859. HB. 133. Chr. 147. Cystopteris leptophylla Pr. 1836 (nomen); Hypolepis Pr. 1849 (nomen); H. serrata Fée 1869.

inermis Christ, Bull. Boiss. II. 4. 1095. 1904 = H. repens var.

javanica Moore, Ind. 253. 1861 = Dryopteris setigera.

leptophylla Pr. Epim. 66. 1849 = H. incisa.

meifolia (Eat.) Bak. Ann. Bot. 5. 209. 1891. NPfl. 279 — Mexico. Cheilanthes Eat. Proc. Amer. Ac. 18. 185. 1883.

mexicana Liebm. Vid. Selsk. Skr. V. 1. 260 (seors. 108). 1849 — Mexico.

millefolium Hk. sp. 2. 68 t. 95 B. 1852. HB. 130. NPfl. 278 — N. Zealand.

mitis Kze.; Kuhn, Linn. 36. 105. 1869 — Brasilia.

monticola Hk. sp. 2. 114. 1852 = Adiantopsis.

nigrescens Hk. sp. 2. 66 t. 90 C. 1852. HB. 129. NPfl. 278 — India occ. Venezuela. Amazonas.

nuda Mett.; Kuhn, Linn. 36. 105. 1869 — Venezuela.

obtusata (»Pr.«) Kuhn, Chaetopt. 347. 1882 (nomen). — an Cheilanthes ob. Pr.?

pallida Hk. sp. 2. 64. 1852 = H. tenuifolia.

parallelogramma (Kze.) Pr. Tent. 162. 1836. HB. 130. NPfl. 278 — Guatemala-Peru-Brasilia. — Cheilanthes Kze. Linn. 9. 85. 1834.

parviloba Fée. Cr. vasc. Br. 1. 53 t. 20 f. 1. 1869 = H. hostilis.

paupercula Hk. sp. 2. 73 t. 88 C. 1852 = Adiantopsis.

pedata Hk. sp. 2. 73 t. 92 A. 1852 = Adiantopsis.

Poeppigiana Mett. Fil. Lechl. 1. 18. 1856 = Dryopteris punctata.

polypodioides Hk. sp. 2. 63. 1852 = Dryopteris punctata.

pteridioides Hk. 2 Cent. t. 59. 1861 = Pteris brevisora.

pteroides Mett. Fil. Lechl. **1**. 17 t. 3 f. 7-13. 1856 — Peru.
punctata Mett.; Kuhn, Fil. Afr. 120. 1868 = Dryopteris.
punctata Bedd. Handb. Suppl. 19. 1892 = ? Dryopteris.
Purdieana Hk. sp. **2**. 69 t. 91 B. 1852. HB. 130 — Columbia. Jamaica.
radiata Hk. sp. **2**. 72 t. 91 A. 1852 = Adiantopsis.
repens (L.) Pr. Tent. 162. 1836. HB. 129. Chr. 278. NPfl. 278 — Amer. trop. Lonchitis L. sp. **2**. 1078. 1753; Cheilanthes Klf. 1824. Dicksonia aculeata Spr. 1821; Cheilanthes Klf. 1823; Hypolepis J. Sm. 1842.
repens J. Sm. JoB. **3**. 404. 1841 = Saccoloma moluccanum.
resinifera Hk. sp. **2**. 63. 1852 = Dennstaedtia.
resistens Hk. sp. **2**. 64. 1852 = Dryopteris punctata.
rigescens (Kze.?) Moore, Ind. 252. 1861? — Brasilia. Columbia. ? Cheilanthes Kze. Flora **1839**[1]. Beibl. 51; Polypodium punctatum var. Bak. 1870. (Cheil. rigescens Kze. est. fortasse = H. brasiliana).
rugulosa J. Sm. Bot. Mag. **72** Comp. 8. 1846 = Dryopteris punctata*.
rugulosa Hk. sp. **2**. 68. 1852 = ? Paesia rugosula.
Schimperi (Kze.) Hk. sp. **2**. 70. 1852. NPfl. 279 — Afr. orient. Cheilanthes Kze. Farnkr. **1**. 52 t. 26. 1840. HB. 133. Chr. 146; Adiantopsis [Moore 1857.
Sellowiana Kl. Linn. **14**. 286. 1840 (nomen) = ? Gymnogramma.
serrata Fée, Cr. vasc. Br. **1**. 53 t. 13 f. 3. 1869 = H. incisa.
setigera Hk. sp. **2**. 62. 1852 = Dryopteris.
Smithii Moug.: Fée, Gen. 147. 1850-52 = Dennstaedtia scandens.
sparsisora (Schrad.) Kuhn, Fil. Afr. 120. 1868 — Afr. austr. Ins. Marscarenae. Fernando Po. — Cheilanthes Schrad. Gött. gel. Anz. **1818**. 918; Phegopteris Keys. 1873; Cheilanthes anthriscifolia Schlecht. (1832) (non Willd. 1810); Hypolepis Pr. 1836. HB. 129. Chr. 279. NPfl. 278 — Cheilanthes aspera Klf. 1831; Hypolepis Pr. 1836. (conf. Kuhn, Bot. Zeit. **1869**. 392).
spectabilis Link, Fil. sp. 67. 1841 = Adiantopsis chlorophylla.
stenophylla Moore, Ind. XXXIX. 1857 = Dryopteris setigera.
stolonifera Fée, Cr. vasc. Br. **2**. 35 t. 91 f. 2. 1872-73 = H. repens.
tenuifolia (Forst.) Bernh. Schrad. neu. Journ. **1**[2]. 34. 1806. HB. 129. Chr. 278. NPfl. 278 — China-Malesia-Australia-Polynesia-N.Zealand. — Lonchitis Forst. Prod. 80. 1786; Phegopteris Keys. 1873. Cheilanthes arborescens Sw. 1806; Adiantum Poir. 1810 — Cheilanthes pallida Bl. 1828; Hypolepis [Hk. 1852.
trichodes Fée, Gen. 146. 1850-52 = Dryopteris setigera.
viscosa (Karst.) Mett. Ann. sc. nat. V. **2**. 238. 1864 (nomen): Karst. Fl. Col. **2**. 89 t. 145-46. 1869 — Columbia. — Cheilanthes Karst. Bot. Zeit. **1854**. 855 (nomen; non Carm. 1818 nec Link 1833).

HYPOPELTIS Michaux, Fl. Bor. Am. **2**. 226. 1803 (in obs.); Bory, Bél. Voy. Bot. **2**. 63. 1833 = **Polystichum.**

aculeata Todaro, Syn. pl. acot. vasc. Sicil. 33. 1866 = α P. lobatum. β P. aculeatum.
amygdalina Bory, Bél. Voy. Bot. **2**. 64. 1833 — Java. (an Nephrolepis sp.?)
articulata Bory. Bél. Voy. Bot. **2**. 64. 1833 = Oleandra a.
biserrata Bory, Bél. Voy. Bot. **2**. 65. 1833 = Nephrolepis b.
coriacea Bory. Bél. Voy. Bot. **2**. 70. 1833 = P. adiantiforme.
exaltata Bory, Bél. Voy. Bot. **2**. 66. 1833 = Nephrolepis e.
Hamiltonii Bory, Bél. Voy. Bot. **2**. 69. 1833 = P. aristatum var.
hastulata Todaro. Syn. pl. acot. vasc. Sicil. 32. 1866 = P. aculeatum var. 42.
lobulata Bory, Exp. scientif. de Morée **3**[2]. 286. 1832 = P. aculeatum.

Lonchitis Todaro, Syn. pl. acot. vasc. Sicil. 32. 1866.
marginifera Bory, Bél. Voy. Bot. **2**. 69. 1833 — Java.
Aspidium Mett.; Salom. 1883. — (Dryopteris sp. dub.)
mauritiana Bory, Bél. Voy. Bot. **2**. 67. 1833 = P. aculeatum var. 1.
nitida Bory, Bél. Voy. Bot. **2**. 68. 1833 — India.
palmoides Bory, Bél. Voy. Bot. **2**. 65. 1833 = Nephrolepis biserrata.
prolifera Bory, Bél. Voy. Bot. **2**. 67. 1833 = P. aculeatum var. 29.
propinquoides Bory, Bél. Voy. Bot. **2**. 69. 1833 — Java.
Aspidium Mett.; Salom. 1883. (= Dryopteris gongylodes?).
setosa Bory, Bél. Voy. Bot. **2**. 68. 1833 = P. varium.
squarrosa Bory, Bél. Voy. Bot. **2**. 67. 1833 = P. lobatum var. 4.
straminea Bory, Bél. Voy. Bot. **2**. 67. 1833 = P. aculeatum var. 1.

HYSTEROCARPUS Langsd. (Fée; Gen. Fil. 216. 1850-52) = **Didymochlaena**.

Isgnogramme Hieronymus, Engl. Jahrb. **34**. 475. 1904 (*Gymnogramme §*) = Gymnogramma.

ISOLOMA J. Smith, Journ. Bot. **3**. 414. 1841 et in Hk. Gen. Fil. t. 102. 1842 = [**Lindsaya**.
divergens J. Sm. JoB. **3**. 414. 1841.
Guerinianum Fée, Gen. 108. 1850-52.
lanuginosum J. Sm. in Hk. Gen. t. 102. 1842 = Nephrolepis acutifolia.
ovatum Pr. Epim. 101. 1849 = L. Gueriniana var.
Walkerae Pr. Epim. 101. 1849.

Isotes Moore, Ind. Fil. XXXVII. 1857 (*Hewardia §*) = Adiantum (Leprieurii).

JAMESONIA Hooker et Greville, Icon. Fil. t. 178. 1830 — (Polypodiaceae Gen. **61**).

adnata Kze. Farnkr. **2**. 80 t. 133 f. 1. 1851 = Polypodium moniliforme.
bipinnata Fée, Gen. 161. 1850-52 = Gymnogramma elongata.
Bogotensis Karst. Fl. Col. **2** 29 t. 115. 1862 = J. rotundifolia.
brasiliensis Christ, Farnkr. 75. 1897. NPfl. 261 — Brasilia.
canescens (Kl.) Kze. Farnkr. **1**. 195. 1846; **2**. 81. t. 133 f. 2. 1851. Chr. 74. NPfl. 260 — Venezuela-Ecuador. — Gymnogramme Kl. Linn. **20**. 407. 1847 (non Bl. 1828); Psilogramme Kuhn 1882; Jamesonia imbricata β HB. 369 — J. robusta Karst. 1862; Gymnogramme Hier. 1904.
ciliata Karst. Fl. Col. **2**. 85 t. 143 f. 13. 1864-69 (nomen) — Columbia.
Gymnogramme Hier. Engl. Jahrb. **34**. 477. 1904.
cinnamomea Kze. Bot Zeit. **1844**. 738. Chr. 75 — Costa Rica-Venezuela.
Gymnogramme Kl. 1847; Psilogramme Kuhn 1882.
elongata J. Sm. Hist. Fil. 168. 1875 = Gymnogramma.
glutinosa Karst. Fl. Col. **2**. 85 t. 143. 1864-69. NPfl. 260 — Columbia.
Gymnogramme Mett. 1864; Psilogramme Kuhn 1882.
Goudotii (Hier.) C. Chr. Ind. 373. 1905 — Columbia.
Gymnogramme Hier. Engl. Jahrb. **34**. 476. 1904.
hispidula Kze. Bot. Zeit. **1844**. 739 = Gymnogramma.
imbricata (Cav.) Hk. et Grev. Ic. Fil. addend. 1831. HB. 369 (excl. varr.). NPfl. 261 — Columbia-Peru. — Pteris Cav. hort. Matrit. **1**. —; Sw. Syn. 102. 1806; Cheilanthes Desv. 1827; Allosorus Pr. 1836; Gymnogramme Kl. 1847; Psilogramme Kuhn 1882. Pteris orbiculata Poir. 1804; Jamesonia pulchra Hk. et Grev. 1830.
imbricata b. laxa Mett. Fil. Lechl. **1**. 10. 1856 (nomen) = J. scalaris.
laxa Kze.; Kuhn, Linn. **36**. 69. 1869 (syn.). NPfl. 260 — Venezuela.
Gymnogramme Mett.; Kuhn, Linn. **36**. 69. 1869; Psilogramme Kuhn 1882.

nivea Karst. Fl. Col. **2**. 29 t. 115. 1862. Chr. 75. NPfl. 260 — Venezuela-Ecuador. — Psilogramme Kuhn 1882.
paleacea Kze. Bot. Zeit. **1844**. 739 = Polystichum aculeatum var. 27.
Pearcei Bak. Syn. 514. 1874 (imbricata var.) — Ecuador.
pulchra Hk. et Grev. Ic. Fil. t. 178. 1830 = J. imbricata.
robusta Karst. Fl. Col. **2**. 29 t. 115. 1862 = J. canescens.
rotundifolia Fée, 7 mém. 41 t. 10 f. 3. 1857. Chr. 75 — Costa Rica-Venezuela. Gymnogramme Mett. 1864; Psilogramme Kuhn 1882 — Jamesonia Bogotensis Karst. 1862; Gymnogramme Hier. 1904.
scalaris Kze. Bot. Zeit. **1844**. 738. Chr. 74. NPfl. 261 — Costa Rica-Peru. Gymnogramme Kl. 1847; Psilogramme Kuhn 1882.
scalaris Fée, Cr. vasc. Br. **2**. 38. 1872-73 = J. brasiliensis.
tolimensis (Hier.) C. Chr. Ind. 374. 1905 — Columbia. Gymnogramme Hier. Engl. Jahrb. **34**. 475. 1904.
verticalis Kze. Bot. Zeit. **1844**. 739. Chr. 75. NPfl. 260 — Columbia. Gymnogramme Kl. 1847; Psilogramme Kuhn 1882; Jamesonia imbricata [γ HB. 369.
JENKINSIA Hooker. Genera Fil. t. 75 B. 1840 = **Leptochilus.**
undulata Hk. Gen. t. 75 B. 1840 = L. scalpturatus var.
KAULFUSSIA Blume, Enum. pl. Jav. 260. 1828 = **Christensenia.**
aesculifolia Bl. Enum. 260. 1828.
assamica Griff. Asiat. Research. **19**. 108 t. 18. 1836 } = C. aesculifolia varr.
Korthalsii de Vriese, Ned. kr. Arch. **3**. 186. 1852; Mon. 14 t. 5 f. 1. } = C. aesculifolia varr.
Lobbiana de Vriese. Mon. Maratt. 14. 1853 } = C. aesculifolia varr.
LACAUSSADEA Gaudichaud, Voy. Bonite Bot. t. 118-120. 1846 = **Polybotrya.**
appendiculata Gaud. l. c. t. 119. 1846 } = P. appendiculata.
montana Gaud. l. c. t. 118. 1846 } = P. appendiculata.
rhizophylla Gaud. l. c. t. 120. 1846 } = P. appendiculata.
LACOSTEA van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 320. 1861 = [**Trichomanes.**
Ankersii, auriculata, guianensis, javanica, tanaica Prantl, Hym. 50. 1875 = T. sp. homonym.
Boryana, rhomboidea, Singaporiana, Zollingeri Prantl, Hym. 50. 1875 = T. javanicum varr.
brachypus Prantl, Hym. 50. 1875 = T. pedicellatum.
Lacosteopsis Prantl, Die Hymenoph. 53. 1875 (*Trichomanes §*) = Trichomanes.
LASTREA Bory. Dict. Class. d'hist. nat. **6**. 588. 1824; **9**. 232. 1826 = **Dry-** [**opteris.**
abbreviata Wollaston, Phytol. n. s. **1**. 172. 1855-56 = D. filixmas.
abrupta Pr. Tent. 75. 1836 = D. Kunzeana.
abrupta Liebm. Vid Selsk. Skr V. **1**. 274 (seors. 122). 1849 = D. patens.
acuminata Moore, Ind. 84. 1858 = D. decomposita.
aemula Brack. Expl. Exp. **16** 200. 1854.
affinis Bedd. Hdb. 230. 1883 = Polystichum speciosum.
agatholepis Moore, Ind. 85. 1858 = D. patula.
albicaulis Moore, Ind. 85. 1858 = D. patens.
albopunctata Pr. Tent. 77. 1836 = D. orientalis.
amabilis Moore, Ind. 85. 1858 = Polystichum a.
ameristoneura Moore, Ind. 85. 1858.
ampla Moore, Ind. 85. 1858.
amplissima Moore, Ind. 85. 1858.
angustata Pr. Tent. 75. 1836 = D. patens *stipularis.

angustifrons Moore, Ind. 85. 1858 = Polypodium a.
angustifrons Moore msc. (Hk. sp. **4**. 126) 1862; Bedd. Ferns br. Ind. t. 226. 1867.
anisoptera (Kze.) Moore, Ind. 85. 1858 —? — Aspidium Kze. hb.
aperta Moore, Ind. 85. 1858.
apiciflora Pr. Tent. 76. 1836.
appendiculata Pr. Tent. 75. 1836 = ? D. prolixa.
arguta Brack. Expl. Exp. **16**. 196. 1854 = D. rigida var.
aristata Moore, Ind. 85. 1858 = Polystichum a.
aristata var. Hamiltonii Bedd Ferns br. Ind. Suppl. t. 369. 1876 = Polystichum speciosum.
articulata Brack. Expl. Exp. **16**. 191 t. 26 f. 1. 1854 = D. orientalis.
ascendens J. Sm. JoB. **4**. 193. 1841 = Polystichum apiifolium.
ascensionis J. Sm. Hist. Fil. 215. 1875.
aspidioides Pr. Epim. 41. 1849 = ? Polystichum aculeatum var.
aspidioides Moore, Ind. 85. 1858 = D. tenuifolia.
athamantica Moore, JoB. **5**. 311. 1853.
athyrioides Liebm. Vid. Selsk. Skr. V. **1**. 274 (seors. 122). 1849.
atrata Pr. Tent. 77. 1836 = D. hirtipes.
atrovirens J. Sm. Cat. cult. Ferns 59. 1857 = D. decomposita.
attenuata J. Sm. JoB. **3**. 412. 1841 = D. stenobasis.
attenuata Brack. Expl. Exp. **16**. 193 t. 26 f. 2. 1854 = D. Brackenridgei.
augescens Houlst.; J. Sm. Cat. Kew Ferns 6. 1856 = D. serra.
Balbisiana Bory, Dict. class. **9**. 234. 1826 — India occ.
barbigera Moore; Bedd. Ferns br. Ind. t. 227. 1867.
Barclayi Carr. in Seem. Fl. Vit. 359. 1873 = Aspidium B.
Beddomei Bedd. Ferns. br. Ind. Corr. II. 1870 et Suppl. 16. 1876.
Bergiana Moore, Ind. 86. 1858.
Bergiana Bedd. Ferns br. Ind. Suppl. 16 t. 370. 1876 = D. cana.
bidentata Moore, Ind. 86. 1858 = ? Nephrolepis b.
Blumei Moore, Ind. 94. 1858 = D. rhodolepis.
borneensis Hk. Ic. pl. t. 993. 1854.
Boryana Moore, Ind. 86. 1858.
Brackenridgei Carr. in Seem. Fl. Vit. 359. 1873.
Breutelii Moore, Ind. 59. 1857 = D. limbata.
Brunoniana Pr. Tent. 76. 1836.
Buchanani Bedd. Hdb. 255. 1883.
bulbosa Pr. Tent. 75. 1836 = Woodsia mollis.
caduca Pr. Tent. 77. 1836 = *Cyclopeltis semicordata*.
calcarata Moore, Ind. 87. 1858.
calcarea Bory, Dict. class. **9**. 233. 1826 = D. Robertiana.
Callipteris Newm. Hist. Br. Ferns ed. II. 12. 1844 = D. cristata.
cana J. Sm. Cat. cult. Ferns 57. 1857.
canariensis Moore, Ind. 87. 1858 = D. filix mas *oligodonta.
caripensis Moore, Ind. 87. 1858 = D. submarginalis.
catoptera Pappe et Raws. Syn. Fil. Afr. austr. 12. 1858 = D. lanuginosa.
caudiculata Pr. Epim. 36. 1849 = D. immersa.
cheilanthoides Moore, Ind. 88. 1858.
cheiloplotia Moore, Ind. 88. 1858.
chrysocarpa Moore, Ind. 88. 1858 = D. filix mas.
chrysolepis Moore, Ind. 88. 1858 = D. crenata.
chrysoloba Pr. Tent. 75. 1836.

ciliata Liebm. Vid. Selsk. Skr. V. 1. 273 (seors. 121). 1849 = D. excelsa.
ciliata Hk. JoB. 9. 338. 1857 = D. calcarata var.
ciliata Pr. Epim. 41. 1849 = Polypodium c.
coarctata Moore, Ind. 88. 1858.
cochleata Moore, Ind. LXXXVIII. 1857.
cognata Pr. Epim. 40. 1849.
collina Newm. Hist. br. Ferns ed. II. 224. 1844 = D. spinulosa *dilatata.
concinna Moore, Ind. 86. 1858.
concinna Bedd. Ferns S. Ind. t. 246. 1863-65 = D. deparioides var.
confluens Moore, Ind. 88. 1858 = Aspidium c. (Fée).
coniifolia Moore, Ind. LXXXVIII. 1857 = Polystichum carvifolium.
consobrina Moore, Ind. 88. 1858 = Aspidium c. (Bory).
conspersoides Moore, Ind. 88. 1858 = D. lugubris.
contermina Pr. Tent. 75. 1836 = D. opposita.
corusca Bull. Cat. **1890**. 7 — Hort. (Japonia).
crassifolia Moore, Ind. 89. 1858.
crenata Bedd. Ferns br. Ind. Suppl. 18. 1876.
crinibulbon J. Sm. Hist. Fil. 214. 1875 = D. cirrhosa.
crinita Moore, Ind. LXXXIX. 1857.
cristata Pr. Tent. 77. 1836.
cruciata Pr. Tent. 77. 1836.
Ctenitis Moore, Ind. 89. 1858.
ctenochlaena Moore, Ind. 89. 1858 = ? Aspidium Linkii.
crystallina Moore, Ind. 89. 1858.
Cumingiana Pr. Tent. 75. 1836; Epim. 37. 1849 = D. concinna.
cuspidata Bedd. Ferns br. Ind. Correct. 1870 = D. khasiana.
cyclochlamys Moore, Ind. 89. 1858 = Aspidium c.
davallioides Brack. Expl. Exp. **16**. 202. 1854.
Dayi Bedd. Handb. Suppl. 54. 1892.
decomposita J. Sm. Bot. Mag. **72**. Comp. 34. 1846.
decrescens Moore, Ind. 89. 1858.
decurrens J. Sm. Bot. Mag. **72**. Comp. 32. 1846 } = D. decursive-pinnata.
decursivo-pinnata J. Sm. Ferns br. a. for. 154. 1866 } = D. decursive-pinnata.
decussata (Sieb.) Pr. Tent. 75. 1836. — ? — Aspidium Sieb. Syn. fil. exsic. n. 189.
deflexa Pr. Tent. 76. 1836 = D. Lindigii.
deltoidea Moore, Ind. 89. 1858.
deltoidea Bedd. Ferns S. Ind. t. 248. 1863-65 = D. sparsa var.
densa Pr. Tent. 76. 1836 = D. sparsa.
denticulata Moore, Ind. 90. 1858 = Polystichum d.
deparioides Bedd. Ferns S. Ind. t. 104. 1863-65.
deversa Moore, Ind. 90. 1858 = D. patens.
diastematocarpa Moore, Ind. 90. 1858 = D. patula.
dilatata Pr. Tent. 77. 1836 = D. spinulosa*.
diplazioides Moore, Ind. 90. 1858 = D. Moritziana.
dissecta Carr. in Seem. Fl. Vit. 360. 1873.
distans Brack. Expl. Exp. **16**. 192. 1854 — Brasilia.
divisa Moore, Ind. 90. 1858 = D. Boryana.
Dryopteris Bory, Dict. class. 9. 233. 1826 = D. Linnaeana.
eburnea J. Sm. Bot. Mag. **72**. Comp. 34. 1846 = Athyrium drepanopterum.
edentula Moore, Ind. 90. 1858.

elegans Moore et Houlst.; J. Sm. Cat. cult. Ferns 57. 1857 = Aspidium e.
elongata Pr. Tent. 76. 1836 = D. filix mas*.
Elwesii Bedd. Ferns br. Ind. Suppl. 18 t. 376. 1876.
equestris Liebm. Vid. Selsk. Skr. V. 1. 273 (seors. 121). 1849 = D. excelsa.
eriocarpa Pr. Tent. 77. 1836 = D. crenata.
erosa Deakin, Fl. brit. 4. 101. 1848 = D. filix mas.
erythrosora Moore, Ind. 91. 1858.
exculta Moore, Ind. 91. 1858 = D. effusa*.
exigua J. Sm. JoB. 3. 412. 1841 = D. philippina.
expansa Pr. Tent. 76. 1836 = D. spinulosa *dilatata.
expansa Moore, Ind. 91. 1858 = Aspidium e. (Mart.).
Fairbankii Bedd. Ferns br. Ind. t. 254. 1867 = D. thelypteris*.
falciculata Pr. Tent. 75. 1836.
falcifera Moore. Ind. 92. 1858 = Aspidium Finlaysonianum.
falciloba Hk. JoB. 9. 337. 1857 = D. calcarata var.
Falconeri Bedd. Ferns br. Ind. t. 41. 1865 = D. barbigera.
fallax Moore, Ind. 91. 1858 = Dryopteris amplissima.
ferruginea Moore, Ind. 91. 1858 = Asyndium f. (Fée).
ferruginea Bedd. Ferns S. Ind. t. 100. 1863-65 = D. peranemiformis.
Fijiensis Carr. in Seem. Fl. Vit. 360. 1873.
Filix foemina Colomb, Compt. rendus 107. 1013. 1888 = Athyrium f.
filix mas Pr. Tent. 76. 1836.
flaccida Moore, Ind. 92. 1858.
flaccida Bedd. Ferns S. Ind. t. 99. 1863-65 = D. setigera.
flebilis Moore, Ind. 107. 1858 = D. tristis.
Floridanum J. Sm. Ferns br. and for. ed. II. 306. 1877.
Foenisecii Watson, Phytol. 2. 568. 1846 = D. aemula.
foeniculacea Bedd. Ferns br. Ind. t. 36. 1865 = Polystichum f.
formosa Moore, Ind. 92. 1858 = Polysticum denticulatum.
fragrans Pr. Tent. 76. 1836.
frondosa Moore, Ind. 85. 1858 = Polystichum Webbianum.
funesta Moore, Ind. 88, 92. 1858 = D. protensa var.
furcata Moore, Ind. 92. 1858.
fuscipes Moore, Ind. 92. 1858; Bedd. Hdb. 243. 1883 = D. sagenioides.
Gardneriana Moore, Ind. 92. 1858 = D. dissecta.
glabella Moore et Houlst. Gard. Mag. Bot. 3. —. 1851; Brack. Expl. Exp. 16.
glabra Brack. Expl. Exp. 16. 200. 1854. [199. 1854.
glandulifera Moore, Ind. 93. 1858 = D. prolixa.
glanduligera Moore, Ind. 93. 1858 = D. gracilescens*.
glandulosa Newm. Phytol. 4. 258. 1851 = D. spinulosa *dilatata.
globulifera Brack. Expl. Exp. 16. 194. 1854.
Goeringiana Moore, Ind. 93. 1858 = Athyrium G.
Goldiana Pr. Tent. 76. 1836.
gracilescens Hk. JoB. 9. 338. 1857 = D. gracilescens *glanduligera.
gracilescens Moore, Ind. 93. 1858.
gracilescens Bedd. Ferns S. Ind. t. 110. 1863-65 = D. Beddomei.
gracilis Moore; Bedd. Ferns br. Ind. t. 198. 1866 = ? Polystichum Thomsoni.
grandifolia Pr. Epim. 39. 1849 — Brasilia.
grandis Moore, Ind. 93. 1858 = D. filix mas *elongata.
grisea Bedd, Ferns br. Ind. t. 335. 1870.

grossa Pr. Epim. 41. 1849 = D. patens.
Gueinziana Moore, Ind. 93. 1858.
Harveyi Carr. in Seem. Fl. Vit. 359. 1873.
Hendersoni Bedd. Ferns br. Ind. Suppl. 17 t. 377. 1876.
hirsuta Moore, Ind. LXXXVIII. 1857 = D. crenata.
hirsutipes Bedd. Handb. Suppl. 52. 1892 = D. gracilescens var.
hirta Pr. Tent. 76. 1836.
hirtipes Moore, Ind. 85. 1858.
hispida Houlst.; Moore (et Houlst. Gard. Mag. Bot. **3**. —. 1851) Ind. LXXXVIII. [1857 = Polystichum.
Hopeana Moore, Gard. Chr. n. s. **18**. 744. 1882.
Hookeriana Pr. Tent. 77. 1836 = Polystichum.
immersa Moore, Ind. LXXXIX. 1857.
impressa Moore, Ind. 94. 1858 = Aspidium i. (Fée).
inaequalis Pr. Tent. 77. 1836.
indecora Liebm. Vid. Selsk. Skr. V. **1**. 272 (seors. 120). 1849.
inquinans Moore, Ind. 94. 1858 = D. indecora.
intermedia Pr. Tent. 77. 1836 = D. spinulosa*.
intermedia Bedd. Ferns S. Ind. t. 113. 1863-65 = D. filix mas *Schimperiana.
intermedia Bedd. Ferns br. Ind. correct. II. 1870 = D. rhodolepis.
invisa Pr. Tent. 75. 1836 = D. patens*.
invisa Armstr. Tr. N. Zeal. Inst. **13**. 363. 1882 = D. thelypteris *squamu-[ligera.
Jenmani Bull, Cat. **1887**. 11.
Karwinskiana Moore, Ind. 94. 1858.
Kaulfussii Pr. Tent. 75. 1836 = D. oligocarpa*.
Kohautiana Pr. Tent. 76. 1836 = D. patens.
Kunthii Moore, Ind. 99. 1858 = D. patens.
lacera Eat. Pr. Amer. Ac. **4**. 110. 1858.
laeta Moore, Ind. LXXXVIII. 1857 = Polystichum denticulatum*.
lancastriensis J. Sm. Cat. Kew Ferns 6. 1856 = D. cristata.
lanciloba J. Sm. Ferns br. and for. ed. II. 305. 1877.
lanuginosa Moore, Ind. 87. 1858.
lata J. Sm. JoB. **3**. 412. 1841 = D. crassifolia.
latifrons Brack. Expl. Exp. **16**. 196. 1854.
Leiboldiana Pr. Epim. 41. 1849 = D. patens.
lepida Moore, Card. Chr. n. s. **26**. 681. 1886.
lepidota Moore; Handb. brit. Ferns ed. III. 136. 1857 = D. spinulosa *dilatata.
leptorachis Liebm. Vid. Selsk. Skr. V. **1**. 272 (seors. 120). 1849 = D. patula.
leucolepis Pr. Epim. 39. 1849 = D. setigera.
L'Herminieri Moore, Ind. 95. 1858.
ligulata J. Sm. JoB. **3**. 412. 1841; Pr. Epim. 35. 1849.
limbata Moore, Ind. 95. 1858.
lomatopelta Moore, Ind. 95. 1858.
lugubris Moore, Ind. 95. 1858.
?lutescens Moore, Ind. 95. 1858 = Polypodium pulverulentum.
macrocarpa Pr. Tent. 77. 1836 = D. cochleata.
?macrocarpa Moore, Ind. 95. 1858 = Athyrium m.
macrotis J. Sm. Hist. Fil. 214. 1875.
macroura Pr. Tent. 75. 1836 = D. patens*.
maculata Deakin, Florigr. brit. **4**. 110. 1848 = D. spinulosa *dilatata.
madagascariensis Moore, Ind. 96. 1858 = Aspidium m.

malaccensis Pr. Epim. 35. 1849 = D. extensa.
manilensis Pr. Tent. 76. 1836; Epim. 39. 1849 — Luzon.
Aspidium Mett.; Salom. 1883.
marginalis Pr. Tent. 77. 1836.
marginata Moore, Ind. 96. 1858 = Aspidium m. (Wall.).
mascarenensis Moore, Ind. 96. 1858 = D. crinita.
Maximowiczii Moore, Gard. Chr. n. s. **15**. 626. 1881.
megaphylla Bedd. Handb. Suppl. 63. 1892 = D. subarborea.
melanochlamys Moore, Ind. 96. 1858 = Polystichum m.
melanopus Bedd. Ferns br. Ind. t. 38. 1865 = D. sagenioides.
melanosticta Moore, Ind. 91. 1858 = Polystichum m.
membranifolia Pr. Tent. 76. 1836 = Aspidium m.
membranifolia Bedd. Ferns S. Ind. t. 102. 1863-65 = D. dissecta.
meridionalis Pr. Tent. 76. 1836 = Polypodium m.
mexicana Pr. Tent. 76. 1836 = D. patula.
microcarpa Moore, Ind. 96. 1858 = D. microchlaena.
microchlaena Moore, 97. 1858.
microchlamys de Vriese, Tijdschr. Wis. en Nat. Wet. Amsterd. **1**. 155. 1848 = [Mesochlaena polycarpa.
microsora Pr. Tent. 76. 1836 = D. decomposita.
microstegia Bedd. Ferns br. Ind. t. 39. 1865 = D. brunnea.
moesta Moore, Ind. 107. 1858 = Aspidium triste Kze.; Fée.
monosora Pr. Epim. 36. 1849 = D. tristis var.
monosticha Moore, Ind. 97. 1858 = D. tristis var.
montana Moore, Hdb. br. Ferns ed. II. 100 —. = D. oreopteris.
mucronata Pr. Tent. 75. 1836; Epim. 37. 1849 = D. falciculata.
multiflora Newm. Hist. br. Ferns ed. II. 215. 1844 = D. spinulosa *dilatata.
multijuga J. Sm. JoB. **4**. 193. 1841 = ? D. extensa.
Napoleonis Pr. Tent. 77. 1836.
nemophila Moore, Ind. 87. 1858 = D. catocarpa.
nemorosa Moore, Ind. 95. 1858.
nephrodioides Moore, Ind. 98. 1858.
nephrodioides Bedd. Ferns br. Ind. t. 199. 1866 = D. crassifolia.
nigricaulis Moore, Ind. 98. 1858 = ? D. sagenioides.
nitida Pr. Tent. 75. 1836 = D. crinita.
nivalis Moore, Ind. 98. 1858 = D. rigida *australis.
noveboracensis Pr Tent. 75. 1836.
obovata Bedd. Handb. Suppl. 60. 1892.
obscura Moore, Ind. 92. 1858 = D. Fischeri.
obtusiloba Moore, Ind. 98. 1858 = D. huatuscensis.
obtusiloba Bedd. Ferns. br. Ind. t. 296. 1868 = D. peranemiformis var.
ochthodes Moore, Ind. 98. 1858.
odontoloma Moore, Ind. 90. 1858; Bedd. Ferns S. Ind. t. 114. 1863-65 = D. filix mas.*
odontoloma Bedd. Handb. 248. 1883 = D. filix mas *serrato-dentata.
odorata Pr. Tent. 77. 1836 = D. lanuginosa.
odorata Bedd. Ferns br. Ind. Correct. II. 1870 = D. crenata.
oligocarpa Moore, Ind. 86. 1858.
opaca Hk. JoB. **9**. 339. 1857 = Polystichum varium.
opposita Pr. Tent. 77. 1836 = D. mascarenarum.
Oreopteris Bory, Dict. class. **9**. 233. 1826.
Orizabae Moore, Ind. 99. 1858.

pachyrachis Moore, Ind. 99. 1858.
padangensis Bedd. Handb. Suppl. 60. 1892.
paleacea Moore, Ind. 99. 1858 = D. filix mas*.
pallens Brack. Expl. Exp. **16**. 197. 1854 — Luzon. — Aspidium Mett.: Salom. 1883.
pallida Moore, Ind. 99. 1858 = Aspidium p. Bl.
paludosa Moore, Ind. 99. 1858 = D. brunnea.
palustris J. Sm. Cat. cult. Ferns 56. 1857 = D. thelypteris.
panamensis Pr. Tent. 76. 1836 = D. opposita
parallelogramma Liebm. Vid. Selsk. Skr. V. **1**. 271 (seors. 119). 1849 = D. filix mas *parallelogramma.
paradoxa Moore, Ind. 99. 1858 = Aspidium p.
Parishii Bedd. Ferns br. Ind. t. 43. 1865.
patens Pr. Tent. 75. 1836.
patentissima Pr. Tent. 76. 1836 = D. filix mas.*
patula Moore, Ind. 99. 1858.
pauciflora Pr. Tent. 75. 1836 = D. crinita.
paucijuga Moore, Ind. 99. 1858 = Aspidium p.
pauperis Moore, Ind. 99. 1858 = D. patens.
paupertina Liebm. Vid. Selsk. Skr. V. **1**. 272 (seors. 120). 1849 — D. patula.
pedata Moore, Ind. LXXXVIII. 1857.
pellita Pr. Epim. 37. 1849 = ? D. parasitica.
pennigera Pr. Tent. 76. 1836.
pentagona Moore, JoB. **5**. 227. 1853 = D. inaequalis.
Phegopteris Bory, Dict. class. **9**. 233. 1826.
philippina Pr. Epim. 36. 1849 = D. ligulata.
pilosissima J. Sm.; Moore, Gard. Chr. **1855**. 677 c. fig. = D. Vogelii.
pilosula Moore, Ind. 100. 1858 = ? D. mollicula.
Plantii Moore, JoB. **5**. 227. 1853 = D. athamantica.
platypus Moore, Ind. 100. 1858 = Polystichum aristatum var.
podophylla Hk. JoB. **9**. 339. 1857.
podophylla J. Sm. Cat. cult. Ferns 57. 1857 = D. Sieboldii.
Poeppigiana Pr. Epim. 40. 1849 — Cuba.
Poiteana Bory, Dict. class. **9**. 233. 1826.
polystichoides Pr. Epim. 38. 1849 — Brasilia. — Aspidium Mett.; Salom. 1883.
polyphylla Pr. Tent. 75. 1836 = D. opposita.
Prenticei Carr. in Seem. Fl. Vit. 359. 1873.
Presliana J. Sm. JoB. **3**. 412. 1841 = Cyclopeltis.
prionitis Moore, Ind. 101. 1858 = ? D. prionites.
prolifica Moore, Gard. Chr. n. s. **18**. 744. 1882 = D. erythrosora f. monstr.
prolixa Pr. Tent. 75. 1836.
prolonga Moore, Ind. 101. 1858 = Aspidium p.
propinqua J. Sm. JoB. **3**. 412. 1841 = D. Blumei et D. Preslii.
propinqua Pr. Epim. 38. 1849 = D. Preslii.
protensa Moore, Ind. 101. 1858.
pseudomas Wollast. Phytol. n. s. **1**. 172. 1855-56 = D. filix mas.
Ptarmica Moore, Ind. 101. 1858.
pubescens Pr. Tent. 76. 1836.
pulchra Pr. Tent. 75. 1836 = D. strigosa.
pulvinulifera Bedd. Ferns. br. Ind. t. 333. 1870.
pumila J. Sm. Hist. Fil. 214. 1875 — Hort. (Yorkshire).
quinquangularis J. Sm. Cat. cult. Ferns 57. 1857 = ? D. decomposita.

radicans Moore, Ind. 102. 1858 = Aspidium r. Fée.
recedens J. Sm., Moore Gard. Chr. **1855**. 708.
recurva Newm. Nat. Alm. 23. 1844 = D. aemula.
Reinwardtiana Moore, Ind. 102. 1858 = D. calcarata.
remota Moore, Ind. 102. 1858 = D. filix mas × spinulosa.
Richardsii Moore, Gard. Chr. n. s. **15**. 104. 1881.
rigida Pr. Tent. 77. 1836.
riparia Pr. Tent. 75. 1836 = D. tomentosa.
Rivoirei Moore, Ind. 102. 1858 = D. oligocarpa.
rivulorum Pr. Tent. 75. 1836 = D. opposita.
Robertiana Newn. Natur. Alm. 17. 1844; Hist. Br. Ferns ed. II. 17. 1844.
rotundata Pr. Tent. 77. 1836.
rubiginosa Brack. Expl. Exp. **16**. 201. 1854.
rufidula Pr. Tent. 76. 1836 = Woodsia ilvensis.
sagenioides Moore, Ind. 96. 1858.
sancta Moore, Ind. 103. 1858 = D. opposita.
sancta J. Sm. Ferns br. and for. 159. 1866.
scabra Pr. Epim. 41. 1849 = Polypodium s.
scabriuscula Pr. Tent. 75. 1836; Epim. 35. 1849 = D. patens.
scabriuscula Dav. Bot. Gaz. **21**. 256 (syn.). 1896 = D. Davenportii.
scabrosa Moore, Ind. 103. 1858.
Schiedeana Pr. Tent. 76. 1836 (nomen) — Mexico.
Schomburgkii Moore, Ind. 103. 1858 — Aspidium S.
semicordata Pr. Tent. 77. 1836 = Cyclopeltis s.
semihastata Moore, Ind. 103. 1858.
sericea Schott; Bedd. Ferns br. Ind. t. 308. 1869 = D. calcarata.
serra Pr. Tent. 75. 1836.
setifera Pr. Tent. 75. 1836 (nomen) —?
setigera Moore, Ind. 103. 1858 = Aspidium s. (Sw.).
setigera Bedd. Ferns br. Ind. Correct. II. 1870.
setosa Pr. Epim. 40. 1849.
setosa Moore, Ind. 103. 1858 = Polystichum varium.
setosa Bedd. Ferns br. Ind. t. 262. 1868 = Cystopteris s.
setosa J. Sm. Ferns br. and for. ed. II. 306. 1877 = D. Millettii.
Shepherdi Moore, Ind. 90. 1858 = D. decomposita var.
Sieberiana Pr. Tent. 75. 1836 = D. crinita.
Sieboldii Moore, Ind. LXXXVIII. 1857.
sikkimensis Bedd. Hdb. 259. 1883.
similis J. Sm. JoB. **3**. 412. 1841 = D. crassifolia.
simulata Dav. Bot. Gaz. **19**. 497 (syn.). 1894.
singalanensis Bedd. Handb. Suppl. 54. 1892.
sparsa Moore, Ind. 87, 104. 1858.
spectabilis J. Sm. JoB. **3**. 412. 1841 = D. syrmatica.
spectabilis Bedd. Hdb. 257. 1883 = D. Hendersoni.
spinescens Moore et Houlst. (? Gard. Mag. Bot. **3**. —. 1851) = Aspidium s.
spinosa Newm. Nat. Alm. 21. 1844 = D. spinulosa.
spinulosa Pr. Tent. 76. 1836.
splendens Bedd. Ferns br. Ind. t. 42. 1865.
Sprengelii Pr. Tent. 75. 1836 = D. Balbisii.
squamigera Brack. Expl. Exp. **16**. 198. 1854.

squamulosa Pr. Tent. 76. 1836 = D. thelypteris var.
Standishii Moore, Gard. Chr. **1863**. 292 = Polystichum.
stipitata Moore, Ind. 105. 1858.
stipulacea Moore, Ind. 105. 1858 = D. heteroptera.
strigilosa Dav. Bot. Gaz. **21**. 257. 1896 (syn.).
strigosa Pr. Tent. 75. 1836.
subincisa Pr. Tent. 76. 1836.
subquinquefida Pr. Tent. 76. 1836 = D. protensa.
sulcata Pr. Tent. 75. 1836 = D. crinita.
?syrmatica Moore, Ind. 105. 1858.
tenera Moore, Ind. 105. 1858.
tenericaulis Moore, Ind. 99. 1858 = D. setigera.
tenuicula Moore, Ind. 105. 1858 = Aspidium t.
tenuifolia Pr. Epim. 37. 1849.
tenuifolia Brack. Expl. Exp. **16**. 199. 1854 = D. tenuifrons.
tetragona Pr. Tent. 76. 1836 = D. pseudotetragona.
Thelypteris Bory, Dict. class. **9**. 233. 1826.
thelypteroides Moore, Ind. 106. 1858 = D. tomentosa.
thibetica Bedd. Handb. Suppl. 52. 1892
Thwaitesii Bedd. Ferns br. Ind. Suppl. 18. 1876.
tomentosa Moore, Ind. 86. 1858.
Torresiana Moore, Ind. 86. 1858 = Polystichum T.
trichotoma Moore, Ind. 106. 1858 = Aspidium t.
tristis Moore, Ind. 107. 1858 = Alsophila t.
truncata Brack. Expl. Exp. **16**. 195 t. 27. 1854 = D. filix mas.
Tussacii Moore, Ind. 107. 1858 = Aspidium T.
tylodes Bedd. Ferns br. Ind. Correct. II. 16. 1870 = D. ochthodes.
uliginosa Newm. Phytol. **3**. 679. 1849 = D. cristata × spinulosa.
undulata Bedd. Ferns S. Ind. t. 271. 1863-65 = D. sparsa*.
unidentata Bedd. Handb. Suppl. 53. 1892 = D. monodonta.
varia Moore, Ind. 107. 1858 = Polystichum v.
variabilis J. Sm. Hist. Fil. 217. 1875 = D. Vogelii.
velata Moore, Ind. 107. 1858.
?vellea Moore, Ind. 107. 1858.
velutina Brack. Expl. Exp. **16**. 198. 1854.
venusta J. Sm. JoB. **4**. 193. 1841.
verrucosa J. Sm. JoB. **3**. 412. 1841; Pr. Epim. 36. 1849 = D. immersa.
vestita J. Sm. Bot. Mag. **72**. Comp. 33. 1846 = D. deflexa.
vilis Moore, Ind. 107. 1858.
villosa Pr. Tent. 76. 1836.
viscosa J. Sm. JoB. **3**. 412. 1841.
Vogelii Moore, Ind. 108. 1858.
Walkerae Bedd. Ferns br. Ind. Correct. II. 1870; Suppl. 16. 1876.
Wallichii Moore, Ind. 104. 1858 = D. Hendersoni.
Wattii Bedd. Handb. Suppl. 49. 1892 = Polystichum.
xylodes Moore, Ind. 107. 1858 = D. ochthodes.
zeylanica Moore, Ind. 108. 1858 = Aspidium z.

Lastreastrum Presl, Epim. Bot. 37. 1849 (*Lastrea §*) = Dryopteris.
LECANIUM Reinwardt, Flora **1825**[2]. 3 Beibl. 48 = **Lecanopteris.**
LECANIUM Presl, Hymenoph. 11. 1843 = **Trichomanes.**
membranaceum Pr. Hym. 12 t. 1. 1843.

LECANOPTERIS Reinwardt, Flora 1825[2]. 3 Beibl. 48; Blume, Enum. 120. 1828 — (Polypodiaceae Gen. **103**).

carnosa (Reinw.) Bl. Enum. 120. 1828. NPfl. 328 — Malesia. Perak.
Onychium Reinw. Syll. Pl. **2**. 3. 1824; Polypodium Christ 1894; Farnkr. 115; Yapp, Ann. Bot. **16**. 188. 1902; P. Lecanopteris Mett. 1866. P. patelliferum Burck 1884.

Curtisii Bak. JoB. **1881**. 366; Ic. pl. t. 1607. NPfl. 328 — Sumatra.

deparioides (Ces.) Bak. JoB. **1881**. 366. NPfl. 328 — Borneo.
Davallia Ces. Atti Ac. Napoli 7[8]. 13 t. 4 f. 8. 1876; Polypodium Christ 1897. 116 (non Bak. 1879).

incurvata Bak. Ann. Bot. **8**. 123. 1894 — Sumatra.

Macleayii Bak. Malesia **2**. 244. 1886. NPfl. 328 — Java.

Nieuwenhuisii Christ, Ann. Jard. Buit. II. **5**. 127 t. 3 f. 1. 1905 — Borneo.

pumila Bl. Fl. Javae, Fil. t. 94 B. 1829 (sine descr.); Copeland, Polyp. Philipp. 134. 1905 — »Malesia«. Ins. Philipp.

LEMAPTERIS Rafinesque, Journ. Phys. **89**. 262. 1819 = **Pteris**.

LEMMA Jussieu, »Ht. Trian. 1759«; Adanson, Fam. d. pl. **2**. 21. 1763; Aublet, Hist. Guian. **2**. 960. 1775 = **Marsilea**.

minuta Desrouss.; Lam. Enc **3**. 721. 1789.

quadrifolia Desrouss.; Lam. Enc. **3**. 720. 1789.

LEMMAPHYLLUM Presl, Epim. Bot. 157. 1849 = **Drymoglossum**.

carnosum Pr. Epim. 158. 1849.

microphyllum Pr. Epim. 263. 1849.

piloselloides Luerss. Bot. Centralbl. **11**. 78. 1882. = D. heterophyllum.

spatulatum Pr. Epim. 158. 1849 = D. heterophyllum.

Lepichosma J. Smith, Journ. Bot. **4**. 50. 1841 (*Nothochlaena* §) = Notholaena.

Lepichroa Moore, Index LXIII. 1857 (*Grammitis* §) = Ceterach.

LEPICYSTIS J. Smith, Lond. Journ. Bot. **1**. 195. 1842 = **Polypodium**.
Goniophlebium § Lepicystis J. Sm. JoB. **4**. 56. 1841.

angusta, fallax, Friederichsthaliana, furfuracea, lanceolata, murorum, percussa, piloselloides, plebeja, thyssanolepis, vacciniifolia Diels, Nat. Pfl. **1**[4]. 323. 1899 — chrysolepis Diels, l. c. 322 — leucospora Diels, l. c. 324 = Polypodum sp. homonym.

incana J. Sm. Lond. JoB. **1**. 195. 1842 = P. polypodioides.

lepidopteris J. Sm. Hist. Fil. 112. 1875.

macrocarpa Diels, Nat. Pfl. **1**[4]. 322. 1899 = P. pycnocarpum.

onusta Diels, Nat. Pfl. **1**[4]. 323. 1899 = P. monosorum.

rhagadiolepis J. Sm. Ferns br. and for. 80. 1866 = P. thyssanolepis.

sepulta J. Sm. Lond. JoB. **1**. 195. 1842 = P. lepidopteris.

squamata J. Sm. Cat. Kew Ferns 1. 1856.

Lepidoglossa Christ, Mon. Elaph. 21. 1899 (*Elaphoglossum* §) = Elaphoglossum.

LEPIDONEVRON Fée, Gen. Fil. 301. 1850-52 = **Nephrolepis**.

acuminatum Fée, Gen. 301. 1850-52 = N. biserrata var.

biauritum Fée, Gen. 301. 1850-52.

bidentatum Fée, Gen. 301. 1850-52.

biserratum Fée, Gen. 301. 1850-52.

distans Fée, Gen. 302. 1850-52 — San Domingo.

hirsutulum Fée, Gen. 301. 1850-52.

longifolium Fée, Gen. 301. 1850-52 = N. hirsutula.

obtusatum Fée, Gen. 301. 1850-52 — Brasilia.

punctulatum Fée, Gen. 301 t. 23 C f. 1. 1850-52 = N. biserrata*.
rufescens Fée, Gen. 301. 1850-52 = N. hirsutula.
sesquipedale Fée, Cr. vasc. Br. 1. 148. 1869 = N. rivularis.
trichomanoides Fée, Gen. 301. 1850-52 = Arthropteris obliterata.
volubile Fée, Gen. 301. 1850-52 = N. radicans.

Lepisorus J. Smith, Bot. Mag. **72**. Comp. 13. 1846 (*Drynaria* §) = Polypodium.

LEPTOCHILUS Kaulfuss, Enum. Fil. 147. 1824 — (Polypodiaceae Gen. **23**).

acrostichoides (Afz.) C. Chr. Ind. 384. 1905 — Afr. trop. centr. et occ.
Hemionitis Afz.; Sw. Schrad. Journ. **1800²**. 17. 1801; Gymnogramma? Desv. 1811; Polybotrya Mett. 1868; Acrostichum Afzelii Carr. 1901.

alienus (Sw.) C. Chr. Bot. Tidsskr. **26**. 285. 1904 — Amer. trop.
Acrostichum Sw. Schrad. Journ. **1800²**. 13. 1801. HB. 419; Gymnopteris Pr. 1836. Chr. 51. NPfl. 201; Anapausia Pr. 1849; Chrysodium Mett. 1856 — Acrostichum brunneum Willd. 1810; Poecilopteris Pr. 1836; Acrostichum cladorrhizans Spr. 1821 — A. portoricense Spr. 1821; Gymnopteris Fée 1845; Anapausia Pr. 1849 — Gymnopteris semipinnatifida Fée 1845; Anapausia Pr. 1849; Acrostichum Hk. 1864 — ? A. hastatum Liebm. 1849 (non Thbg. 1784); Gymnopteris Fourn. 1872 — Acrostichum irregulare Liebm. 1849; Gymnopteris Fourn. 1872 — Chrysodium maracaybense Karst. 1860.

antrophyoides (Bak.) C. Chr. Ind. 384. 1905 — Borneo.
Acrostichum Bak. Journ. Linn. Soc. **22**. 231. 1886.

auriculatus (Lam.) C. Chr. Ind. 384. 1905 — Africa trop.
Acrostichum Lam. Enc. **1**. 36. 1783; A. punctatum L.fil. Suppl. 444. 1781 (non L. 1863); Chrysodium Mett; Kuhn 1868; Gymnopteris Christ 1897. 50. NPfl. 201; Leptochilus C. Chr. 1904; Acrostichum punctulatum Sw. 1801. HB. 419; Poecilopteris Pr. 1836; Cyrtogonium J. Sm. 1841; Heteronevron Fée, 1845; Chrysodium Mett. 1856.

axillaris (Cav.) Klf. Enum. 147 t. 1 f. 10. 1824 — Asia trop.
Acrostichum Cav. Anal. Hist. Nat. **1**. 101. 1799; Descr. 239. 1802. HB. 320; Gymnopteris Pr. 1836. NPfl. 199; Gymnogramme Ces. 1876.

Bernoullii (Kuhn) C. Chr. Bot. Tidsskr. **26**. 296 f. 8. 1904 — Mexico-Costa Rica. — Acrostichum Kuhn; Christ, Bull. Soc. bot. Belg. **35**. 244. 1896; Gymnopteris Kuhn; Diels 1899. NPfl. 201.

Boivini (Mett.) C. Chr. Ind. 384. 1905 — Ins. Comorae. Angola.
Chrysodium Mett.; Kuhn, Fil. Afr. 51. 1868; Acrostichum Welw.; Carr. 1901.

Bonii (Christ) C. Chr. Ind. 384. 1905 — Formosa. Tonkin.
Gymnopteris Christ, Bull. Boiss. II. **4**. 610. 1904.

Cadieri (Christ) C. Chr. Ind. 384. 1905 — Annam.
Gymnopteris Christ, Journ. de Bot. **19**. 126. 1905.

celebicus (Bak.) C. Chr. Ind. 384. 1905 — Celebes.
Acrostichum Bak. Kew Bull. **1901**. 145.

contaminoides (Christ) C. Chr. Bot. Tidsskr. **26**. 294 f. 6. 1904 — Brasilia austr. Paraguay. — Gymnopteris Christ, Ann. Cons. Jard. bot. Genève **3**. 32. 1899; Acrostichum Lindm. 1904.

curupirae (Lindm.) C. Chr. Bot. Tidsskr. **26**. 296 f. 7. 1904 — Brasilia.
Acrostichum Lindm. Hedwigia **43**. 311. 1904; A. scalpturatum Lindm. Ark. för Bot. **1**. 253 t. 8 f. 9. 1903 (non Kze. 1848).

cuspidatus (Pr.) C. Chr. Ind. 384. 1905 — Asia-Austr. trop. Polynesia. Ins. Sechellae. — Nephrodium? Pr. Rel. Haenk. **1**. 31. 1825; Polystichum

Pr. 1836; Heteroneuron Pr. 1849; Chrysodium Kuhn 1869. Acrostichum repandum Bl. 1828. HB. 419; Bolbitis Schott 1834; Campium Pr. 1836; Cyrtogonium J. Sm. 1841; Heteronevron Fée 1845; Poecilopteris Pr. 1849; Chrysodium Mett. 1870; Gymnopteris Christ 1897. 50. NPfl. 201 — Acrostichum proliferum Bl. 1828 — A. Quoyanum Gaud. 1827; Heteronevron Fée 1845; Poecilopteris Pr. 1849; Chrysodium Ettingsh. 1864 — C. bipinnatifidum Mett.; Kuhn 1868; Heteroneuron Kuhn 1889. Ins. [Sechellae).

decurrens Bl. Enum. 206. 1828 — Asia trop.
Anapausia Pr. 1849; Gymnopteris Moore 1857; Acrostichum Thwait. 1864; A. variabile Hk. 1864. HB. 417; Gymnopteris Bedd. 1868. Chr. 48. NPfl. 199.

diversifolius (Bl.) C. Chr. Ind. 9. 1905; 385. 1906 — Malesia.
Acrostichum Bl. Enum. 103. 1828; Bolbitis Schott 1834; Poecilopteris Pr. 1826; Cyrtogonium J. Sm. 1841; Heteronevron Fée 1845; Chrysodium Ettingsh. 1864.

exculptus (Bak.) C. Chr. Ind. 9. 1905; 385. 1906 — Borneo.
Acrostichum Bak. JoB. **1888**. 326.

Fendleri (Eat.) C. Chr. Bot. Tidsskr. **26**. 285. 1904 — Trinidad.
Meniscium Eat. Bot. Gaz. **3**. 89. 1878; Acrostichum Bak. 1887.

fluviatilis (Hk.) C. Chr. Ind. 10. 1905; 385. 1906 — Africa occ. trop.
Acrostichum Hk. sp. **5**. 274. 1864. HB. 422; Chrysodium Kuhn 1868; Poecilopteris J. Sm. 1875. — (f. Carr. = L. auriculatus var.)

gaboonensis (Hk.) C. Chr. Ind. 10. 1905; 385. 1906 — Africa occ. trop.
Acrostichum Hk. sp. **5**. 270. 1864. HB. 417; Gymnopteris J. Sm. 1866. NPfl. 199; Chrysodium Kuhn 1868.

guianensis (Aublet) C. Chr. Bot. Tidsskr. **26**. 288 f. 1. 1904 — Guiana-Brasilia. Puerto Rico. — Polypodium Aublet, Hist. Guian. 962. 1775; Acrostichum Bak. 1870; Gymnopteris Christ 1901; Acrostichum scandens Raddi 1819; Cyrtogonium J. Sm. 1841; Poecilopteris Pr. 1849; Neurocallis Moore 1857; Gymnopteris Christ 1897. 51. NPfl. 201; Heteronevron Raddianum Fée 1845; Acrostichum Kze.; Hk. 1864. HB. 423; Lomariopsis Mett. 1856.

Harlandii (Hk.) C. Chr. Ind. 11. 1905; 385. 1906 — Hongkong. Formosa.
Acrostichum Hk. sp. **5**. 274. 1864. HB. 418; Chrysodium Hance 1873; Gymnopteris J. Sm. 1875. NPfl. 199; G. decurrens Hk. JoB. **9**. 359. 1857 (non Leptochilus supra).

heteroclitus (Pr.) C. Chr. Ind. 11. 1905; 385. 1906 — Asia trop. et subtrop. Melanesia. — Acrostichum Pr. Rel. Haenk. **1**. 15 t. 2 f. 2. 1825; Poecilopteris Pr. 1836; Heteronevron Fée 1845; Chrysodium Kuhn 1869. Acrostichum flagelliferum Wall.; Hk. et Gr. 1827. HB. 418; Bolbitis Schott 1834; Cyrtogonium J. Sm. 1841; Chrysodium Mett. 1856; Poecilopteris J. Sm. 1857; Gymnopteris Bedd. 1876. Chr. 49. NPfl. 201.

Heudelotii (Bory) C. Chr. Ind. 11. 1905; 385. 1906 — Africa trop.
Gymnopteris Bory; Fée, Acrost. 84 t. 45. 1845. Chr. 51. NPfl. 201; Anapausia Pr. 1849; Acrostichum Hk. 1864. HB. 422; Chrysodium Kuhn 1868.

hilocarpus Fée, Acrost. 87 t. 48 f. 1. 1845 = L. latifolius.

Hollrungii (Kuhn) C. Chr. Ind. 11. 1905; 385. 1906 — N. Guinea.
Gymnopteris Kuhn in Schum. Fl. Kais. Wilh. Land 8. 1889. NPfl. 200; Acrostichum Bak. 1891.

Humblotii (Bak.) C. Chr. Ind. 11. 1905; 385. 1906 — Madagascar.
Acrostichum Bak. JoB. **1884**. 114.

inconstans (Copeland) C. Chr. Ind. 386. 1906 — Luzon.
Gymnopteris Copeland in Perkins, Fragm. 178. 1905.

labrusca (Christ) C. Chr. Ind. 386. 1906 — Congo.
Acrostichum Christ, Ann. Mus. Congo Bot. 5. **1**. 10. 1903.

lanceolatus Fée, Acrost. 87 t. 47 f. 1. 1845 — Asia trop.
Acrostichum Hk. 1864. HB. 420 (non L. 1753); Gymnopteris Bedd. 1876. NPfl. 199; G. Feei Moore 1857 — G. normalis J. Sm. 1841 (nomen); Dendroglossa Pr. 1849.

lanceolatus Zoll. Syst. Verz. 33. 1854 = L. Linnaeanus.

latifolius (Meyen) C. Chr. Ind. 19. 1905; 386. 1906 — Ins. Philipp.
Polybotrya Meyen hb.; Gymnopteris Pr. 1836 (nomen); Goldm. Nova Acta **19**. Suppl. I. 460. 1843; Pr. Epim. 150. 1849; Dendroglossa Fée 1850-52 — Gymnopteris taccaefolia J. Sm. 1841 (nomen). NPfl. 199; Leptochilus Fée, Acrost. 89 t. 50. 1845; Dendroglossa Fée 1850-52; Polybotrya Mett. 1859; Acrostichum Hk. 1864. HB. 418 — Gymnopteris trilobata J. Sm. 1841 (nomen): Polybotrya Mett. 1856. Leptochilus subquinquefidus Fée 1845; Gymnopteris Pr. 1849; Dendroglossa Fée 1850-52; Polybotrya [Mett. 1859.

Laurentii (Christ) C. Chr. Ind. 12. 1905; 386. 1906 — Congo.
Acrostichum Christ, Bull. Soc. bot. Belg. **36**^{2}. 94. 1897.

Lindigii (Mett.) C. Chr. Bot. Tidsskr. **26**. 290 f. 4. 1904 — Costa-Rica-Ecuador.
Chrysodium Mett. Ann. sc. nat. V. **2**. 204. 1864; Acrostichum Bak. 1868. HB. 423 (non Karst. 1858) — Gymnopteris costaricensis Christ 1904.

Linnaeanus Fée, Acrost. 87 t. 47 f. 2. 1845 — Malesia. Annam.
Acrostichum Hk. 1860. HB. 417; Chrysodium Kuhn 1869; Gymnopteris Christ 1905; Dendroglossa lanceolata Pr. 1849 (vix Acrostichum L. 1753); Leptochilus Zoll. 1854 (non Fée 1845); Gymnopteris Moore 1857.

Listeri (Bak.) C. Chr. Ind. 12. 1905; 386. 1906 — Christmas Ins. (Mare ind.).
Acrostichum Bak. Journ. Linn. Soc. **25**. 361. 1889.

lomarioides Bl. Enum. 206. 1828 — Assam. Malesia-Polynesia.
Neurocallis Pr. 1849; Polybotrya Kuhn 1869; Lomagramma J. Sm. 1875; Acrostichum Christ 1895; Gymnopteris Christ 1901; Cheilolepton Blumeanum Fée 1845; Polybotrya Mett. 1856; Acrostichum Hk. 1864. HB. 423; Poecilopteris Bedd. 1870; Gymnopteris Bedd. 1876. NPfl. 201 — Lomagramma pteroides J. Sm. 1841; Polybotrya Kuhn 1869 — Acrostichum cultratum Bak. 1874. HB. 524 — Chorizopteris bipinnata Moore 1855; Neurocallis Moore 1857. Chorizopteris pinnata Moore 1855; Neurocallis Moore 1857; Cyrtogonium J. Sm. 1875.

metallicus (Bedd.) C. Chr. Ind. 386. 1906 — Ceylon.
Gymnopteris Bedd. Ferns br. Ind. Suppl. 26 t. 390. 1876. Chr. 49. NPfl. 199.

minor Fée, Acrost. 87 t. 25 f. 3. 1845 — Ins. Philippinae. Ind. bor.
Acrostichum Mett. 1856. HB. 420; Gymnopteris Hk. 1861. Chr. 48. NPfl. 199.

minutulus Fée, 10 mém. 8 t. 31 f. 2. 1865 — Ceylon.

modestus (Bak.) C. Chr. Ind. 14. 1905; 386. 1906 — Borneo.
Acrostichum Bak. Journ. Linn. Soc. **22**. 231. 1886.

Naumanni (Kuhn) C. Chr. Ind. 386. 1906 — N. Hannover.
Heteroneuron Kuhn, Forschr. Gazelle **4**. Farne. 5 t. 1. 1889; Gymnopteris Diels 1901.

neglectus (Bailey) C. Chr. Ind. 14. 1905: 386. 1906 — Queensland.
Acrostichum Bailey, Proc. Linn. Soc. N. S. Wales **5**. 32. 1881; Bak. Ic. pl. t. 1689. 1887; Gymnopteris Diels 1899. NPfl. 200.

nicotianifolius (Sw.) C. Chr. Bot. Tidsskr. **26**. 285. 1904 — Amer. trop. Acrostichum Sw. Syn. 13, 199. 1806. HB. 419; Gymnopteris Pr. 1836; Chr. 51. NPfl. 201; Anapausia Pr. 1849; Chrysodium Mett. 1856 — Acrostichum acuminatum Willd. 1810; Gymnopteris Pr. 1836; Anapausia Pr. 1849; Chrysodium Mett. 1856 — Heteroneuron meniscioides Fée 1845; Poecilopteris Pr. 1849.

obtusifolius Pr. Epim. 185. 1849 = L. decurrens.

oligarchicus (Bak.) C. Chr. Bot. Tidsskr. **26**. 285. 1904 — Peru. Ecuador. Acrostichum Bak. Syn. 418. 1868; Gymnopteris J. Sm. 1875; Chrysodium Salom. 1883.

oligodictyus (Bak.) C. Chr. Ind. 14. 1905; 387. 1906 — Borneo. Acrostichum Bak. Journ. Linn. Soc. **24**. 261. 1887.

opacus (Mett.) C. Chr. Bot. Tidsskr. **26**. 292 f. 5. 1904 — Columbia. Chrysodium Mett. Ann. sc. nat. V. **2**. 204. 1864.

pandurifolius (Hk.) C. Chr. Bot. Tidsskr. **26**. 286. 1904 — Peru. Ecuador. Gymnopteris Hk. 2 Cent. t. 87. 1861; Acrostichum Hk., 1864. HB. 422.

perakensis (Bedd.) C. Chr. Ind. 387. 1906 — Perak. Lomagramma Bedd. Handb. Suppl. 107. 1892.

phanerodictyus (Bak.) C. Chr. Ind. 15. 1905; 387. 1906 — Ins. S. Thomé (Afr. occ.). Acrostichum Bak. Bol. Soc. Brot. **4**. 156 t. 2 f. C. 1886.

Preslianus (Fée) C. Chr. Ind. 16. 1905; 387. 1906 — Ins. Philipp. India austr. Madagascar. — Heteronevron Fée, Acrost. 92 t. 39 f. 1. 1845; Poecilopteris Moore 1857; Acrostichum Hk. 1864. HB. 420; Gymnopteris J. Sm. 1875. Acrostichum punctulatum Pr. 1825 (non Sw. 1801); Campium Pr. 1836.

prolifer (Bory) C. Chr. Ind. 387. 1906 — Asia trop. N. Guinea. Polybotrya Bory, Bél. Voy. Bot. **2**. 18. 1833; Heteronevron Fée 1845; Poecilopteris Pr. 1849.

punctatus C. Chr. Bot. Tidsskr. **26**. 292 in obs. 1904 = L. auriculatus.

quercifolius Fée, Acrost. 88. 1845 = L. zeylanicus.

Rawsoni (Bak.) C. Chr. Ind. 16. 1905; 387. 1906 — Mauritius. Acrostichum Bak. Ann. Bot. **5**. 496. 1891.

rivularis (Brack.) C. Chr. Ind. 17. 1905; 387. 1906 — Fiji. Cyrtogonium Brack. Expl. Exp. **16**. 85 t. 11 f. 3. 1854; Neurocallis Moore 1867; Acrostichum Hk. 1864. HB. 524; Poecilopteris Carr. 1873.

sagenioides (Kuhn) C. Chr. Ind. 182. 1905; 387. 1906 — Aneiteum. Chrysodium Kuhn, Linn. **36**. 63. 1869.

salicinus (Hk.) C. Chr. Ind. 17. 1905; 387. 1906 — Afr. occ. trop. Acrostichum Hk. sp. **5**. 265. 1864. HB. 422; Chrysodium Kuhn 1868; Gymnopteris J. Sm. 1875.

scalpturatus (Fée) C. Chr. Ind. 17. 1905; 387. 1906 — India bor. Heteronevron Fée. Acrost. 95 t. 56. 1845; ? Acrostichum Kze. 1848; Chrysodium Mett. 1856. Acrostichum costatum Wall. 1828 (nomen); Campium Pr. 1836 (nomen); Cyrtogonium J. Sm. 1841 (nomen); Poecilopteris Moore 1857; Gymnopteris Bedd. 1876. Chr. 50. NPfl. 201; Acrostichum virens δ HB. 420 — Nothochlaena undulata Wall. 1828; Jenkinsia Hk. 1842; Campium Pr. 1849; Lomariopsis Mett. 1856 — Meniscium deltigerum Wall. 1828.

serratifolius (Mertens) C. Chr. Bot. Tidsskr. **26**. 289 f. 3. 1904 — Brasilia. Guiana. — Acrostichum Mertens; Klf. Enum. 66. 1824. HB. 423; Bolbitis Schott 1834; Cyrtogonium J. Sm. 1841; Heteronevron Fée 1845; Chry-

sodium Ettingsh. 1864; Gymnopteris Diels 1899. NPfl. 201; Acrostichum fraxinifolium Pr. 1822 (non R.Br. 1810); Poecilopteris Pr. 1836; Chrysodium Mett. 1856; Polybotrya Mett. 1859.

serratus (Kuhn) C. Chr. Bot. Tidsskr. **26**. 289 f. 2. 1904 — Peru.
Chrysodium Kuhn, Linn. **36**. 63. 1869; Acrostichum Bak. 1874. HB. 524.

subcrenatus (Hk. et Grev.) C. Chr. Ind. 19. 1905; 388. 1906 — India austr. Ceylon. — Acrostichum Hk. et Grev. Ic. Fil. t. 110. 1829; Campium Pr. 1836; Cyrtogonium J. Sm. 1841; Heteronevron Fée 1845; Poecilopteris Moore 1857; Gymnopteris Bedd. 1883. NPfl. 201 — Acrostichum proliferum Hk. 1844 (non Bl. 1828); Campium Pr. 1849; Polybotrya Mett. 1856 (non Bory 1833); Acrostichum virens γ HB. 420. Poecilopteris Hookeriana Moore 1857; Acrostichum Thwait. 1864.

subquinquefidus Fée, Acrost. 88 t. 49. 1845 = L. latifolius.

taccifolius Fée, Acrost. 89 t. 50. 1845 = L. latifolius.

Taylori (Bailey) C. Chr. Ind. 19. 1905; 388. 1906 — Queensland.
Acrostichum Bailey, Proc. Roy. Soc. Queensl. **1**. 89. 1884; Syn. Queensl. Fl. Suppl. I. 65. 1886.

Thwaitesianus Fée, 10 mém. 7. 1865 = L. lanceolatus.

tricuspis (Hk.) C. Chr. Ind. 20. 1905; 388. 1906 — Sikkim.
Acrostichum Hk. sp. 5. 272 t. 304. 1864. HB. 422; Gymnopteris Bedd. 1866. Chr. 49. NPfl. 199; Cheiropleuria J. Sm. 1875.

umbrosus (Liebm.) C. Chr. Ind. 20. 1905; 388. 1906 — Mexico.
Acrostichum Liebm. Vid. Selsk. Skr. V. **1**. 174 (seors. 22). 1849.

varians (Mett.) Fourn. Bull. Soc. Fr. **15**. 394. 1868 — N. Caledonia.
Acrostichum Mett. Ann. sc. nat. IV. **15**. 56. 1861. HB. 524.

virens (Wall.) C. Chr. Ind. 20. 1905; 388. 1906 — India austr. Ceylon. Birma. Ins. Philipp. — Acrostichum Wall. (1828 dub.); Hk. et Grev. Ic. Fil. t. 221. 1831. HB. 420 part.; Bolbitis Schott 1834; Campium Pr. 1836; Cyrtogonium J. Sm. 1841; Heteronevron Fée 1845; Poecilopteris Moore 1857; Gymnopteris Keys. 1873; Acrostichum contaminans Wall. 1828 (nomen); Poecilopteris Moore 1857; Gymnopteris Bedd. 1876. Chr. 50. NPfl. 201 — Acrostichum crispatulum Wall. 1828 (nomen); Cyrtogonium J. Sm. 1841; Campium Pr. 1849; Poecilopteris J. Sm. 1857 — Acrostichum terminans Wall. 1829 (nomen); Bolbitis Schott 1834; Poecilopteris Moore 1857 — Poecilopteris semicordata Moore; Bedd. 1868; Acrostichum Bak. 1868. HB. 422; Gymnopteris J. Sm. 1875. (Hk. 2 Cent. t. 88).

Wallii (Bak.) C. Chr. Ind. 20. 1905; 388. 1906 — Ceylon. — Acrostichum Bak. JoB. **1872**. 146. HB. 523; Gymnopteris Bedd. 1876. NPfl. 199.

zeylanicus (Houtt.) C. Chr. Ind. 16. 1905; 388. 1906 — Asia trop.
Ophioglossum Houtt. Nat. Hist. **14**. 43. 1783; Pfl. Syst. (ed. germ.) **13**[1]. 47 t. 94 f. 1. 1786. Osmunda trifida Jacq. 1789. Acrostichum quercifolium Retz. 1791. HB. 418; Onoclea Willd. 1802; Gymnopteris Bernh. 1806. Chr. 49. NPfl. 200; Leptochilus Fée 1845; Dendroglossa Fée 1850-52; Polybotrya Mett. 1859.

zeylanicus Fée, 10 mém. 8 t. 31 f. 1. 1865 = L. decurrens.

Zollingeri (Kze.) Fée. Gen. 55. 1850-52 — Malesia. ? Fiji.
Acrostichum Kze. Bot. Zeit. **1846**. 419; Anapausia Pr. 1849; Chrysodium Kuhn 1869. Gymnopteris subrepanda J. Sm. 1841 (nomen); Poecilopteris Pr. 1849; Acrostichum Hk. 1864. HB. 419 — ? Gymnopteris subsimplex Fée 1845.

LEPTOCIONIUM Presl. Hymenoph. 26. 1843 = **Hymenophyllum.**
- abruptum Pr. Abh. böhm. Ges. V. **5**. 336. 1848.
acanthoides v.d.B. Ned. kr. Arch. **4**. 383. 1859 } = H. aculeatum.
aculeatum v.d.B. Ned. kr. Arch. **4**. 383. 1859 }
affine v.d.B. Ned. kr. Arch. **4**. 383. 1859 = H. holochilum.
attenuatum v.d.B. Ned. kr. Arch. **4**. 384. 1859 = H. magellanicum.
barbatum v.d.B. Ned. kr. Arch. **5**³. 416. 1863.
Braunii v.d.B. Ned. kr. Arch. **4**. 383. 1859 = H. brachyglossum.
caespitosum Pr. Abh. böhm. Ges. V. **5**. 336. 1848 = Trichomanes c.
dentatum v.d.B. Ned. kr. Arch. **4**. 384. 1859.
denticulatum v.d.B. Ned. kr. Arch. **4**. 382. 1859.
dichotomum v.d.B. Ned. kr. Arch. **4**. 384. 1859.
dicranotrichum Pr. Hym. 27 t. 11 f. D. 1843.
edentulum v.d.B. Ned. kr. Arch. **5**³. 148. 1863.
flaccidum v.d.B. Ned. kr. Arch. **5**³. 149. 1863 = H. khasianum.
fuciforme Pr. Abh. böhm. Ges. V. **5**. 336. 1848.
?fucoides Pr. Hym. 27. 1843.
holochilum v.d.B. Ned. kr. Arch. **4**. 383. 1859.
Magellanicum v.d.B. Ned. kr. Arch. **4**. 384. 1859.
Neesii v.d.B. Ned. kr. Arch. **4**. 383. 1859 = H. denticulatum.
pedicellatum Fourn. Bull. Soc. Fr. **19**. 249. 1872 = H. fucoides var.
Preslii v.d.B. Ned. kr. Arch. **4**. 384. 1859 = H. serrulatum.
protrusum Pr. Abh. böhm. Ges. V. **5**. 336. 1848 = H. polyanthos var.
serrulatum v.d.B. Ned. kr. Arch. **4**. 383. 1859.
Seselifolium v.d.B. Ned. kr. Arch. **4**. 384. 1859.
sororium Pr. Abh. böhm. Ges. V. **5**. 336. 1848; Epim. 21 t. 11. 1849 = H. dila-
tortuosum v.d.B. Ned. kr. Arch. **4**. 384. 1859. [tatum.
violaceum v.d.B. Ned. kr. Arch. **5**³. 147. 1863.

LEPTOGRAMMA J. Smith. Journ. Bot. **4**. 51. 1841 = **Dryopteris.**
aspidioides Kl. Linn. **20**. 415. 1847.
asplenioides J. Sm. JoB. **4**. 52. 1841 = D. aspidioides.
attenuata J. Sm. Cat. Kew Ferns 1856 = D. heteroclita.
aurita Bedd. Handb. 377. 1883.
decurrenti-alata J. Sm. Hist. Fil. 232. 1875.
diplazioides Und. Bull. Torr. Cl. **29**. 626. 1902.
gracilis J. Sm. JoB. **4**. 52. 1841 = D. heteroclita.
Levingii Bedd. Handb. Suppl. 99. 1892.
Linkiana J. Sm. JoB. **4**. 52. 1841 = D. diplazioides.
Lovei J. Sm. JoB. **4**. 52. 1841 = D. africana.
obtusata J. Sm. Hist. Fil. 232. 1875 = D. opaca.
opaca Bedd. Handb. 379. 1883.
pilosa Und. Bull. Torr. Cl. **29**. 626. 1902.
polypodioides J. Sm. JoB. **4**. 52. 1841.
rupestris Kl. Linn. **20**. 415. 1847.
subsimilis J. Sm. Hist. Fil. 232. 1875.
totta J. Sm. JoB. **4**. 52. 1841 = D. africana.
villosa J. Sm. JoB. **4**. 52. 1841 = D. dasyphylla.

LEPTOLEPIA Mettenius; Kuhn, Chaetopterides 348. 1882 — (Polypodiaceae Gen. **81**).

Andersoni Mett.: Kuhn, Chaetopt. 348. 1882 (nomen) — Ins. Hawaii.
Davallia Bak. 1891 (nomen). ?Microlepia jamaicensis Hill. 1888 (non Fée 1866).

aspidioides Mett.; Kuhn, Chaetopt. 348 t. 2. f. 1-3. 1882 (nomen) — N. Caledonia. Davallia Bak. 1891 (nomen).

novae-zelandiae (Col.) Kuhn, Chaetopt. 348. 1882. NPfl. 212 — N. Zealand. Austr. bor. — Davallia Col. Tasm. Journ. **2**. 182. 1844; Hk. sp. **1**. 158 t. 51 B. 1846. HB. 91; Microlepia J. Sm. 1856. Chr. 308; Dennstaedtia Keys. 1873. Acrophorus hispidus Moore 1854.

tripinnata (F. Muell.) Kuhn, Chaetopt. 348. 1882 — Queensland. Davallia F. Muell.; Benth. Fl. Austr. **7**. 717. 1878.

Leptomanes Prantl, Die Hymenoph. 52. 1875 *(Trichomanes §)* = Trichomanes.

Leptophyllum Blume, Fl. Javae, Fil. Suppl. t. 95. 96 —. *(Olfersia §)* = Stenochlaena.

LEPTOPLEURIA Presl, Tent. Pterid. 136. 1836 = **Nephrolepis.**

abrupta Pr. Tent. 137 t. 5 f. 9-11. 1836.

LEPTOPTERIS Presl, Suppl. Tent. Pterid. 70. 1845 — (Osmundaceae Gen. **2**).

alpina (Bak.) C. Chr. Ind. 390. 1906 — N. Guinea. Todea Bak. Kew Bull. **1899**. 121.

Fraseri (Hk. et Grev.) Pr. Suppl. 71. 1845. Chr. 334. NPfl. 378 — N. S. Wales. N. Caledonia. — Todea Hk. et Grev. Ic. Fil. t. 101. 1829. HB. 427 (excl. β); Osmunda F. Muell. 1882.

hymenophylloides (A. Rich.) Pr. Suppl. 71. 1845. Chr. 335. NPfl. 378 — N. Zealand. Todea A. Rich. Fl. Nouv. Zél. 97 t. 16. 1832. HB. 427; Osmunda Armstr. 1881. Todea pellucida Carm. 1833 — T. grandipinnula Moore 1886.

marginata (Col.) C. Chr. Ind. 390. 1906 — N. Zealand. Todea Col. Tr. N. Zeal. Inst. **29**. 419. 1897.

Moorei (Bak.) Christ, Farnkr. 335. 1897. NPfl. 378 — Ins. Lord Howe. Todea Bak. JoB. **1873**. 16. HB. 524; Osmunda F. Muell. 1882.

superba (Col.) Pr. Abh. böhm. Ges. V. **5**. 326. 1848; Hk. Ic. pl. t. 910. 1854 Chr. 335. NPfl. 378 — N. Zealand. — Todea Col. Tasm. Journ. **2**. 188. 1844. HB. 428; Osmunda Armstr. 1881.

Wilkesiana (Brack.) Christ, Farnkr. 334. 1897. NPfl. 378. — Polynesia. N. Guinea. Todea Brack. Expl. Exp. **16**. 309 t. 43. 1854; T. Fraseri β HB. 427.

LEPTOSTEGIA Don, Prod. Fl. Nepal. 14. 1825 = **Onychium.**

lucida Don, Prod. 14. 1825 = O. japonicum var.

LEPTOSTEGIA Zippel msc. (Mett. Pol. 50. 1857) = **Polypodium.**

LEUCOMANES Presl, Epim. Bot. 258. 1849 = **Trichomanes.**

album Pr. Epim. 258. 1849 = T. pallidum.

LEUCOSTEGIA Presl, Tent. Pterid. 94. 1836 = **Davallia.**

affinis J. Sm. JoB. **3**. 416. 1841 = D. hymenophylloides.

assamica J. Sm. Hist. Fil. 84. 1875.

borneensis J. Sm. Ferns br. and for. 77. 1866 = Dryopteris b.

chaerophylla J. Sm. Lond. JoB. **1**. 426. 1842 = D. pulchra.

dareaeformis Bedd. Ferns br. Ind. Suppl. 4. 1876 = Polypodium d. et Davallia Clarkei.

falcinella J. Sm. JoB. **3**. 416. 1841.

Griffithiana J. Sm. Hist. Fil. 84. 1875.

hirsuta J. Sm. JoB. **3**. 416. 1841 = Microlepia h.

Hookeri Bedd. Handb. 52. 1883 = D. Clarkei.

hymenophylla Bedd. Ferns br. Ind. Suppl. 4. 1876.

hymenophylloides Bedd. Ferns br. Ind. Suppl. 4. 1876.

immersa Pr. Tent. 95 t. 4 f. 11. 1836.

ligulata J. Sm. Lond. JoB. **1**. 426. 1842 = Acrophorus stipellatus.

Mac Gillivrayi Fourn. Ann. sc. nat. V. **18**. 344. 1873.

maxima Fourn. Ann. sc. nat. V. **18**. 344. 1873.
membranulosa J. Sm. Hist. Fil. 84. 1875.
multidentata Bedd. Ferns br. Ind. Suppl. 4. 1876.
nodosa Bedd. Ferns br. Ind. Suppl. 4. 1876 = Acrophorus stipellatus.
parvula J. Sm. Lond. JoB. **1**. 426. 1842.
pseudo-cystopteris Bedd. Ferns br. Ind. Suppl. 4. 1876.
pulchra J. Sm. Lond. JoB. **1**. 426. 1842.
subsericea Fourn. Ann. sc nat. V. **18**. 344. 1872 = Dryopteris s.
yaklaensis Bedd. Handb. Suppl. 13. 1892.

LINDSAYA Dryander; J. Smith, Mém. Ac. Turin **5**. 413. 1793: Dry. Tr. Linn. Soc. **3**. 39. 1797 (Lindsaea); Klf. Enum. 218. 1824 — (Polypodiaceae Gen. **40**).
aculeata Mett. Ann. sc. nat. IV. **15**. 65. 1861 = Odontosoria.
acutifolia Desv. Prod. 312. 1827 = Nephrolepis.
adiantoides (Bl.) Kuhn, Ann. Lugd. Bat. **4**. 278. 1869 — Malesia. Ins. Philipp. Fiji. Samoa. — Aspidium Bl. Enum. 145. 1828; Saccoloma Pr. 1836; Odontoloma Pr. 1849; Acrophorus Moore 1857; Davallia adiantifolia Hk. 1846. Odontoloma pulchellum J. Sm. 1841; Davallia Hk. 1846. HB. 93; Lindsaya Mett. 1869 — Saccoloma cuneifolium Pr. 1836; Davallia? Hk. 1844; Odontoloma Pr. 1849; Acrophorus Moore 1857.
adiantoides J. Sm. JoB. **3**. 415. 1841 = L. humilis.
Alexandri Hill. Fl. Haw. 622. 1888 = Diellia.
alutacea Mett. Ann. sc. nat. IV. **15**. 63. 1861. HB. 110 — N. Caledonia. Schizoloma Kuhn 1882.
ambigens Ces. Rend. Ac. Napoli **16**. 25. 29. 1877 = L. decomposita.
amboynensis Mett.; Kuhn, Ann. Lugd. Bat. **4**. 279. 1869 = Tapeinidium.
angustata Wall. List pag. 62. 1829 = Schizoloma ensifolia.
angustifolia Bernh. Schrift. Ak. Erfurt **1802**. 14 f. 6 = Odontosoria.
apiculata Kze. Farnkr. **1**. 206 t. 85 f. 2. 1846 = L. cultrata.
apoensis Copeland in Perkins, Fragm. 182 t. 4 f. A. 1905 — Mindanao.
aquatica Jenm. W. Ind. and Guiana Ferns 78. 1899 — Guiana.
arcuata Kze. Linn. **9**. 86. 1834 = L. lancea var.
attenuata Wall. List n. 151. 1828 = L. cultrata.
attenuata Wall. List n. 2192. 1829 = Schizoloma ensifolium.
azurea Christ, Verh. Nat. Ges. Basel **11**. 429. 1896; Ann. Jard. Buit **15**. 101 t. 14. f. 12. 1897 — Celebes.
Balansae Fourn. Ann. sc. nat. V. **18**. 335. 1873 — N. Caledonia.
bantamensis Bl. Enum. 218. 1828 = L. hymenophylloides.
bifida Mett.; Kuhn, Chaetopt. 346. 1882 = Odontosoria.
Billardieri Carr. in Seem. Fl. Vit. 337. 1873 = Schizoloma ensifolium var.
bilobata Pr. Tent. 132. 1836 = L. decomposita.
bipinnata Roxb. Calc. Journ. **4**. 511. 1844 = ? L. lancea.
blanda Mett.; Kuhn, Linn. **36**. 80. 1869 — Java.
Blumeana (Hk.) Kuhn, Ann. Lugd. Bat. **4**. 277. 1869. NPfl. 222 — Celebes. Ins. Philipp. — Davallia Hk. sp. **1**. 117 t. 54 A. 1846. HB. 93. (Chr. 294); Odontoloma Mett. 1856. O. tenuifolium J. Sm. 1841 (nomen, non Lindsaya Bl. 1828); Stenoloma Fée 1850-52.
Boivini Mett.; Kuhn, Fil. Afr. 67, 206. 1868 = Schizoloma.
borneensis Hk. Syn. 107. 1867 — Borneo. Perak.
botrychioides St. Hil. Voy. Distr. Diam. **1**. 379. 1833. HB. 105. NPfl. 221 — Brasilia.
brachypoda Salom. Nom. 212. 1883 = L. cultrata var.

brasiliensis Desv. Berl. Mag. **5**. 330. 1811 = L. lancea.
brevifolia Reinw.; Pr. Tent. 131. 1836 = L. gracilis.
Calomelanos Kze. Bot. Zeit. **1848**. 214 = L. cultrata.
campylophylla Fourn. Ann. sc. nat. V. **18**. 335. 1873 — N. Caledonia.
capillacea Christ, Bull. Boiss. **6**. 144 t. 4 f. 7. 1898 — Luzon.
Catharinae Hk. sp. **1**. 212 t. 65 B. 1846 = L. virescens var.
caudata Hk. sp. **1**. 215. 1846 = L. lancea var.
caudifera Fée, Gen. 106. 1850-52 — Guadeloupe.
centifolia Hill. Fl. Haw. 621. 1888 = Diellia.
cheiroides Fourn. Ann. sc. nat. V. **18**. 334. 1873 — N. Caledonia.
chinensis Mett; Kuhn, Fil. Afr. 67. 1868 = Odontosoria.
clavata Mett. Ann. sc. nat. IV. **15**. 64. 1861 = Odontosoria.
colobodes Kze. Bot. Zeit. **1846**. 446 = L. gracilis.
concinna J. Sm. JoB. **3**. 415. 1841; Hk. sp. **1**. 205 t. 61 B. 1846 = L. gracilis.
consanguinea Fée, 11 mém. 16 t. 6 f. 3. 1866 = L. lancea.
Copelandi C. Chr. Ind. 392. 1906 — Ins. Philipp.
L. montana Copeland in Perkins, Fragm. 182. 1905 (non Fée 1866).
cordata Gaud.; Desv. Prod. 312. 1827 = Schizoloma.
coriifolia Lindm. Ark. för Bot. **1**. 201 t. 8 f. 1. 1903 — Venezuela.
crenata Kl. Linn. **18**. 546. 1844 — Guiana.
crenulata Fée, Gen. 105 t. 28 f. 2. 1850-52 = L. gracilis.
crispa Bak. JoB. **1879**. 39; Ic. pl. t. 1627. NPfl. 221 — Borneo.
cultrata (Willd.) Sw. Syn. 119. 1806. HB. 105 (pt.). Chr. 292. NPfl. 221 — Asiă trop.-Queensland. Madagascar. Réunion.
Adiantum Willd. Phytogr. 14 t. 10 f. 2. 1794 — Lindsaya apiculata Kze. 1846 — L. Calomelanos Kze. 1848 — Davallia brachypoda Bak. 1874. HB. 468; Lindsaya Salom. 1883.
cultrata β japonica Bak. Syn. 105. 1867 = L. japonica.
cuneata (Forst.) C. Chr. Ind. 25. 1905; 392. 1906 — Austr. Tasmania. N. Zealand. Fiji. — Adiantum Forst. Prod. 84. 1786; Davallia Spr. 1799. Lindsaea trichomanoides Dry. Tr. Linn. Soc. **3**. 43 t. 11. 1797. HB. 110. Chr. 295; Adiantum Poir. 1810; Schizoloma Kuhn 1882 — Lindsaea Lessonii Bory 1828. — (an potius Schizoloma sp.?).
cuneata Willd. sp. **5**. 423. 1810 = Schizoloma.
cuneifolia Pr. Rel. Haenk. **1**. 60. 1825 = Odontosoria retusa.
curvans Fée, Gen. 106. 1850-52 = L. quadrangularis.
davallioides Bl. Enum. 218. 1828. HB. 111. Chr. 293. NPfl. 221 — Malesia. Synaphlebium J. Sm. 1842; Schizoloma Moore 1857 — Davallia Kunzeana Hk. 1846.
decomposita Willd. sp. **5**. 425. 1810 — Asia trop. Austr. trop. Polynesia. L. lobata Poir. 1813. HB. 111, 471. Chr. 293. NPfl. 221; Davallia? Desv. 1827; Schizoloma Bedd. 1876. Lindsaya nitens Bl. 1828; Schizoloma Bedd. 1870; Synaphlebium J. Sm. 1875 — Lindsaya recurvata Wall. 1828; Synaphlebium J. Sm. 1841; Schizoloma Moore 1857 — Synaphlebium obtusum J. Sm. 1841; Lindsaya Hk. 1846 — L. propinqua Hk. 1835; Synaphlebium Fée 1850-52 — Lindsaya intermedia Hk. 1846 — Schizoloma malabaricum Bedd. 1868. (HB. 471) — Lindsaya Seemanni Carr. 1873 (non J. Sm. 1854). Synaphlebium pulchrum Brack. 1854; Lindsaea Carr. 1873. (an = L. stolonifera?).
decomposita J. Sm. JoB. **3**. 415. 1841 = L. scandens.

delicatula Christ, Verh. Nat. Ges. Basel **11**. 258. 1895 — Celebes.
Davallia Christ, l. c.; Ann. Jard. Buit. **15**. 97 t. 14. f. 11. 1897.
deltoidea C. Chr. Ind. 393. 1906 — N. Caledonia. N. Hebridae. Ins. Pinorum.
L. elongata Lab. Sert. Austr. Cal. 6 t. 9. 1824. HB. 110 (non Cav. 1802); Schizoloma Kuhn 1882.
Denhami Mett.; Kuhn, Verh. zool. bot. Ges. **19**. 573. 1869 = Tapeinidium.
dentata Wall. List n. 146. 1828 (nomen) — Singapore.
dentata Fée, Cr. vasc. Br. **2**. 20 t. 87 f. 2. 1872-73 — Brasilia.
dimorpha Bailey, Handb. Queensl. Ferns 19. 1874 — Queensland.
L. heterophylla Prentice, JoB. **1873**. 295 (non Dry. 1797).
divaricata Kl. Linn. **18**. 547. 1844 = L. lancea var.
divaricata Mett.; Kuhn, JoB. **1877**. 231 = Odontosoria Schlechtendalii.
divergens Wall.; Hk. et Grev. Ic. Fil. t. 226. 1831 = Schizoloma.
dolabra Kze. Farnkr. **2**. 47. 1849 — Venezuela.
dubia Spr. Syst. **4**. 79. 1827. HB. 105. NPfl. 220 — Guiana. Venezuela. Amazonas. — L. tenera Klf. Enum. 219. 1824 (non Dry. 1797); Wibelia pectinalis Fée 1850-52.
elata Desv. Berl. Mag. **5**. 326. 1811 = L. stricta.
elegans Hk. Ic. pl. t. 98. 1837 = L. stricta var.
elegans Fée, Gen. 106. 1850-52 = L. Feei.
elongata Cav. Descr. 265. 1802 = L. stricta.
elongata Lab. Sert. Austr. Cal. 6 t. 9. 1824 = L. deltoidea.
ensifolia Sw. Schrad. Journ. **1800**². 77. 1801 = Schizoloma.
erecta Mett. Ann. sc. nat. IV. **15**. 65 in obs. 1861 = Diellia.
exilis Fourn. Ann. sc. nat. V. **18**. 335. 1873 — N. Caledonia.
falcata Dry. Tr. Linn. Soc. **3**. 41 t. 7 f. 2. 1797 — Amer. trop. — L. nitidissima Willd. 1810 — L. Leprieurii Hk. 1846. L. trapeziformis ; HB. 107.
falcata Hk. Syn. 113. 1867 = Diellia.
falciformis Hk. sp. **1**. 208 t. 64 B. 1846. HB. 104 — Guiana.
Feei C. Chr. Ind. 393. 1906 — Venezuela.
L. elegans Fée, Gen. 106. 1850-52 (non Hk. 1837).
ferruginea Kuhn, Fil. Afr. 68. 1867 = Odontosoria Melleri.
filiformis Hk. sp. **1**. 212 t. 63 D. 1846 = L. tenuis.
Finlaysoniana Wall. List n. 2197. 1829 = Schizoloma heterophyllum.
flabellifolia Kuhn, v. Deck. Reis. **3**³. Bot. 62. 1879 = Odontosoria.
flabellulata Dry. Tr. Linn. Soc. **3**. 41 t. 8 f. 2. 1797 = L. orbiculata.
flavicans Mett. Mém. soc. sc. nat. Cherb. **10**. 317. 1864 (nomen); Fourn. Ann. sc. nat. V. **18**. 334. 1873 — N. Caledonia.
Fraseri Hk. sp. **1**. 221 t. 70 B. 1846 = Schizoloma.
fumarioides Mett. Ann. sc. nat. V. **2**. 217. 1864 = Odontosoria.
Galeottii Fée, Gen. 107. 1850-52 = L. quadrangularis.
Gardneri Hk. sp. **1**. 213 t. 65 C. 1846 = L. virescens.
genkofolia St. Hil. Voy. Dist. Diam. **1**. 379. 1833 = L. virescens.
gomphophylla Bak. Ann. Bot. **5**. 204. 1891. NPfl. 221 — Borneo.
Goudotiana (Kze.) Mett.; Kuhn. Fil. Afr. 68. 1868 — Madagascar.
Davallia Kze. Anal. 35 t. 22 f. 2. 1837. HB. 101; *Stenoloma*? Fée 1850-52; Odontoloma Mett. 1856; Acrophorus Moore 1857.
gracilis Bl. Enum. 217. 1828 — Asia trop. N. Caledonia.
— L. concinna J. Sm. 1841. HB. 105. L. securifolia Pr. 1836 — L. Lobbiana Hk. 1846. L. crenulata Fée 1850-52 — L. colobodes Kze. 1846.

gracilis Kl. Linn. **18**. 549. 1844 = L. stricta.
gracillima Copeland in Perkins. Fragm. 181. 1905 — Luzon.
Griffithiana Hk. sp. **1**. 219 t. 68 B. 1846 = Schizoloma ensifolium.
Gueriniana Desv. Prod. 312. 1827 = Schizoloma.
guianensis (Aubl.) Dry. Tr. Linn. Soc **3**. 42. 1797. HB. 107. Chr. 293 — Amer. trop. — Adiantum Aubl. Hist. pl. Guian. **2**. 963 t. 365. 1775 — Lindsaya javitensis H.B.Willd. 1810. — (conf. L. stricta).
Harveyi Carr. in Seem. Fl. Vit. 338. 1873 — Fiji. Aneiteum.
heterophylla Dry. Tr. Linn. Soc. **3**. 41 t. 8 f. 1. 1797 = Schizoloma.
heterophylla Bedd. Ferns S. Ind. t. 206. 1864 = Schizoloma.
heterophylla Prentice. JoB. **1873**. 295 = L. dimorpha.
horizontalis Hk. sp. **1**. 214 t. 62 B. 1846 = L. lancea var.
Hosei C. Chr. Ind. 394. 1906 — Borneo.
L. trilobata Bak. JoB. **1891**. 107 (non Col. 1884).
humilis Kuhn, Ann. Lugd. Bat. **4**. 278. 1869 — Celebes. Ins. Philipp.
L. adiantoides J. Sm. 1841 (nomen); Hk. sp. **1**. 204 t. 61 C. 1846. HB. 104. NPfl. 221 (non supra).
hymenophylloides Bl. Enum. 218. 1828. Chr 294. NPfl. 221 — Malesia. Polynesia. — Acrophorus Moore 1857; Davallia Bak. 1867. HB. 93: Odontoloma J. Sm. 1875 — Davallia Lapeyrousii Hk. 1861; Lindsaya Bak. 1867. [HB. 106.
imbricata Desv. Berl. Mag. **5**. 326. 1811 = L. linearis.
imbricata Liebm. Vid. Selsk. Skr. V. **1**. 269 (seors. 117). 1849 — Mexico.
impressa Christ. Ann. Jard. Buit. II. **5**. 132. 1905 — Borneo.
incisa Prentice. JoB. **1873**. 295 — Queensland. N. S. Wales. N. Zealand.
indurata Bak. JoB. **1888**. 324 = Schizoloma.
intermedia Hk. sp. **1**. 222 t. 67 B. 1846 = L. decomposita.
interrupta Wall. List n. 2195. 1829 = L. orbiculata.
jamesonioides Bak. JoB. **1879**. 39; Ic. pl. t. 1626. NPfl. 221 — Borneo. Celebes.
japonica (Bak.) Diels. Nat. Pfl. **1**^{4}. 221. 1899 — Japonia.
L. cultrata β japonica Bak. Syn. 105. 1867.
javanensis Bl. Enum. 219. 1828 = L. orbiculata.
javitensis H.B.Willd. **5**. 424. 1810 = L. guianensis.
Kirkii Hk. Syn. 108. 1867. Chr. 293 — Ins. Sechellae.
L. Pervillei Mett.; Kuhn 1867. NPfl. 221.
Klotzschiana Moritz. Bot. Zeit. **1854**. 855 (nomen); Ettingsh. Farnkr. 121 t. 145 f. 1. 2. 1865 = L. lancea.
Knudsenii Hill. Fl. Haw. 623. 1888 = Diellia.
Kunzei Moritz; Ettingsh. Farnkr. 121 t. 147 f. 7. 1865 = L. lancea.
laciniata Hill. Fl. Haw. 621. 1888 = Diellia.
lancea (L.) Bedd. Ferns br. Ind. Suppl. 6. 1876. Chr. 292. NPfl. 221 — Amer. trop. Ceylon. Malesia. — Adiantum L. sp. ed. II. **2**. 1557. 1763; Lindsaya trapeziformis Dry. Tr. Linn. Soc. **3**. 42 t. 9. 1797. HB. 107 (pt.) — L. horizontalis Hk. 1846 — L. arcuata Kze. 1834 — L. montana Fée 1866 — L. consanguinea Fée 1866 — L. caudata Hk. 1846 (Ceylon).
lanceolata Lab. N. Holl. Pl. Spec. **2**. 98 t. 248 f. 1. 1806 = Schizoloma ensifolium var.
lanuginosa Hk. sp. **1**. 210 t. 69 A. 1846 = Nephrolepis acutifolia.
?lanuginosa Wall. List n. 154. 1828 = Nephrolepis radicans.
Lapeyrousii Bak. Syn. 106. 1867 = L. hymenophylloides var.
laxa Kze.; (Bak. Syn. 107. 1867) = L. lancea var.

Leprieurii Hk. sp. **1**. 208 t. 62 D. 1846 = L. falcata.
leptophylla Bak. JoB. **1884**. 141: Ic. pl. t. 1628. NPfl. 221 — Madagascar.
Lessonii Bory, Dup. Voy. Bot. **1**. 278 t. 37 f. 2. 1828 = L. cuneata var.
L'Herminieri Fée, 11 mém. 15 t. 6 f. 1. 1866 — Guadeloupe.
linearis Sw. Schrad. Journ. **1800**². 78. 1801. HB. 104. Chr. 292. NPfl. 221 — Australia. Tasmania. N. Caledonia. N. Zealand. — Adiantum Poir. 1810; Lindsaya imbricata Desv. 1811 — L. trilobata Col. 1884.
lobata Poir. Enc. Suppl. **3**. 448. 1813 = L. decomposita.
Lobbiana Hk. sp. **1**. 205 t. 62 C. 1846 = L. gracilis.
Loheriana Christ, Bull. Boiss. **6**. 144 t. 4 f. 6. 1898 — Luzon.
longipinna Wall. List n. 2194. 1829 = Schizoloma ensifolium.
longissima Christ, Ann. Jard. Buit. II. **5**. 131. 1905 — Borneo.
Lowei ht. = Arthropteris obliterata.
lucida Bl. Enum. 216. 1828 = L. gracilis.
lunata Willd. sp. **5**. 421. 1810 = L. linearis.
Macgillivrayi Carr. in Seem. Fl. Vit. 337. 1873. HB. 109 — N. Caledonia. Ins. Pinorum.
macrophylla Klf. Enum. 218. 1824 = Schizoloma.
macrophylla Kze. Anal. 37 (non t. 25). 1837 = Adiantum dolosum.
madagascariensis Bak. Journ. Linn. Soc. **16**. 198. 1877; Ic. pl. t. 1629. NPfl. 221 — Madagascar. — Schizoloma Kuhn 1882.
malabarica Bak. Syn. Index 545. 1874 = L. decomposita var.
Mannii Hill. Fl. Haw. 624. 1888 = Davallia.
mazaruniensis Jenm. W. Ind. and Guiana Ferns 75. 1899 — Guiana.
media R.Br. Prod. Fl N. Holl. 156. 1810. HB. 111 — Australia trop. c. ins. Schizoloma Kuhn 1882.
medioeris Fourn. Ann. sc. nat. V. **18**. 336. 1873 — N. Caledonia.
meifolia Mett.; Kuhn, Linn. **36**. 79. 1868 = Odontosoria.
membranacea Kze. Linn. **18**. 121. 1844 = Schizoloma ensifolium.
Merrilli Copeland in Perkins, Fragm. 181. 1905 — Ins. Philipp.
Mertensiana Kze. Bot. Zeit. **1846**. 446 = Schizoloma ensifolium.
?**Michleriana** Eat. Mem. Amer. Acad. n s. **8**. 213. 1860 — HB. 113 — Columbia. Dictyoxiphium Moore 1861.
microphylla Sw. Schrad. Journ. **1800**². 79. 1801. HB. 110. Chr. 295 — Austr. N. Zealand. N. Caledonia. — Adiantum Poir. 1810 (non Sw. 1788); Odontosoria J. Sm. 1875; Schizoloma Kuhn 1882. Stenoloma Lindsayoides Fée 1850-52.
microphylla Pr. Rel. Haenk. **1**. 59 t. 10 f. 2. 1825 = L. stricta.
montana Fée, 11 mém. 17 t. 6 f. 2. 1866 = L. lancea.
montana Copeland in Perkins, Fragm. 182. 1905 = L. Copelandi.
Moorei Fourn. Ann. sc. nat. V. **18**. 336. 1873 = Davallia.
Moritziana Kl. Linn. **18**. 548. 1844 = L. guianensis.
multifrondulosa Fée, Gen. 105 t. 28 f. 1. 1850-52 = L. stricta.
natunae Bak. Kew Bull. **1896**. 40 — Ins. Natuna.
nervosa Mett. Ann. sc. nat. IV. **15**. 62. 1861. HB. 471 — N. Caledonia. Schizoloma Kuhn 1882.
nervosa Lindm. Ark. f. Bot. **1**. 199 t. 8 f. 5. 1903 = L. rigidiuscula.
nitens Bl. Enum. 217. 1828 = L. decomposita.
nitidissima Rich.; Willd. sp. **5**. 423. 1810 = L. lancea.
oblongifolia Reinw.; J. Sm. JoB. **3**. 415. 1841; Hk. sp. **1**. 206 t. 61 D. 1846 = L. pectinata.

obtusa Hk. sp. 1. 224. 1846 = L. decomposita.
odorata Roxb. Calc. Journ. 4. 511. 1844 = L. cultrata.
oligoptera Kze. Bot. Zeit. 1846. 445 — Schizoloma ensifolium.
orbiculata (Lam.) Mett.; Kuhn, Ann. Lugd. Bat. 4. 279. 1869. NPfl. 221 — Asia trop. Austr. trop. — Adiantum Lam. Enc. 1. 41. 1783; Schizoloma Kuhn 1882; Lindsaya flabellulata Dry. Tr. Linn. Soc. 3. 41 t. 8 f. 2. 1797. HB. 107. Chr. 292 — L. tenera Dry. 1797; Adiantum triangulare Poir. 1810 — Lindsaya javanensis Bl. 1828 — Davallia trichomanoides Bedd. 1866 (non Bl. 1828); D. shizophylla Bak. 1874. HB. 468.
ovata J. Sm.: Hk. sp. 1. 204 t. 64 A. 1846 = Schizoloma Guerinianum var.
ovoidea Fée, Cr. vasc. Br. 2. 21 t. 87 f. 1. 1872-73 — Brasilia.
oxyphylla Bak. JoB. 1891. 3 — Madagascar.
pallens Wall. List n. 148. 1828 = L. cultrata.
pallida Kl. Linn. 18. 547. 1844 = L. quadrangularis.
panamensis Mett. Fil. Lips. 105. 1856 = Dictyoxiphium.
parasitica Wall. List n. 2196. 1829 = L. lancea.
? Parishii Bak. Syn. ed. I. 109. 1867 = Stenochlaena sorbifolia var. 18.
Parkeri (Hk.) Kuhn, Chaetopt. 346. 1882 — Guiana.
Davallia Hk. sp. 1. 176 t. 53 C. 1846. HB. 93; Odontoloma Pr. 1849; Acrophorus Moore 1857.
parvifolia Pr. Tent. 131. 1836 = L. stricta.
parvula Fée, 11 mém. 17 t. 7 f. 1. 1866. HB. 471. NPfl. 221 — Trinidad. Guiana.
pectinata Bl. Enum. 217. 1828. HB. 106. Chr. 293. NPfl. 221 — Asia trop. L. oblongifolia Reinw.; J. Sm 1841. — (veris. = L. repens var.).
pelecophylla Pr. Tent. 131. 1836 = L. pectinata.
pellaeiformis Christ, Verh. Nat. Ges. Basel 11. 430. 1896 = Schizoloma.
pendula Kl. Linn. 18. 548. 1844. HB. 108. Chr. 293. NPfl. 221 — Guiana. Brasilia.
pentaphylla Hk. sp. 1. 219 t. 67 A. 1846 = Schizoloma ensifolium.
Pervillei Mett.; Kuhn, Fil. Deck. 17. 1867 = L. Kirkii.
Pickeringii (Brack.) Mett.; Kuhn, Linn. 36. 81. 1869 — Polynesia.
Synaphlebium Brack. Expl. Exp. 16. 223 t. 30 f. 2. 1854; Schizoloma Moore 1857.
pinnata Mett.; Kuhn, Ann. Lugd. Bat. 4. 279. 1869 = Tapeinidium.
plicata Bak. Journ. Linn. Soc. 25. 350. 1889 — Madagascar.
polita Wall. List pag. 62. 1829 = Schizoloma ensifolium.
polymorpha Wall. List n. 147. 1828; Hk. et Grev. Ic. Fil. t. 75. 1829 = L. orbiculata.
portoricensis Desv. Berl. Mag. 5. 326. 1811 — Amer. trop.
L. rubescens Kze. 1848.
prolongata Fourn. Ann. sc. nat. V. 18. 334. 1873 — N. Caledonia.
Schizoloma Kuhn 1882.
propinqua Hk. in Night. Ocean. Sketch. App. 130. 1835; sp. 1. 223 t. 66 B. 1846 = L. decomposita var.
pteroides Desv. Prod. 312. 1827; Wall. List n. 2193. 1829 = Schizoloma ensifolium.
pulchella Mett.; Kuhn, Linn. 36. 81. 1869 = L. adiantoides.
pulchra Carr. in Seem. Fl. Vit. 337. 1873 = L. decomposita var.
pulchra var. Mett. Ann. sc. nat. IV. 15. 65. 1861 = L. gracilis.
pumila Kl. Linn. 18. 545. 1844 = L. guianensis.
pumila Hk. Syn. 112. 1867 = Diellia.
pusilla Splitgb. Tijdschr. Nat. Gesch. 7. 423. 1840 = L. stricta.

quadrangularis Raddi. Opusc. sci. Bol. **3**. 294. 1819; Pl. bras. **1**. 55 t. 74. 1825 — Amer. trop. — L. trapeziformis pt. HB. 107.
Raddiana Kl. Linn. **18**. 549. 1844 = L. stricta.
recurvata Wall. List n. 149. 1828; Hk. sp. **1**. 220 t. 70 A. 1846 = L. decomposita var.
reniformis Dry. Tr. Linn. Soc. **3**. 40 t. 7 f. 1. 1797 = Schizoloma.
repanda Kze. Bot. Zeit. **1848**. 511 — Ins. Bonin Sima. (= L. orbiculata?).
repens (Bory) Bedd. Ferns S. Ind. 72 t. 209. 1863-65; Kuhn, Fil. Afr. 68. 1868. Chr. 294 — Asia trop. Polynesia. Mauritius.
Dicksonia Bory, Voy. **2**. 323. 1804; Davallia Desv. 1827. HB. 93; Odontoloma Pr. 1849; Acrophorus Moore 1857 — Davallia Boryana Pr. 1825; Saccoloma Pr. 1836; Odontoloma J. Sm. 1841 — Davallia hemiptera Bory 1833; Saccoloma? Pr. 1836; Odontoloma Pr. 1849; Acrophorus Moore 1861 — Davallia Macraeana Hk. et Arn. 1832; Odontoloma Brack. 1854; Acrophorus Carr. 1873; Odontoloma Hookeri J. Sm. 1842.
repens Thwait. Enum. pl. Zeyl. 388. 1864 = L. pectinata.
repens var. laciniata Mett.; Kuhn, Ann. Lugd. Bat. **4**. 277. 1869 = L. hymenophylloides.
retusa Mett. Fil. Lips. 105. 1856 = Odontosoria.
rigescens Willd. sp. **5**. 421. 1810 = L. stricta.
rigida J. Sm. JoB. **3**. 415. 1841 (nomen); Hk. sp. **1**. 217 t. 63 A. 1846. HB. 108. NPfl. 221 — Malacca. Perak. Celebes. Ins. Philipp. N. Guinea.
Lindsaynium Fée 1850-52.
rigidiuscula Lindm. Hedwigia **43**. 308. 1904 — Venezuela.
L. nervosa Lindm. Ark. f. Bot. **1**. 199 t. 8 f. 5. 1903 (non Mett. 1861).
ripidopteris Kze. Bot. Zeit. **1850**. 328 — Brasilia.
rubescens Kze. Linn. **21**. 227. 1848 } = L. portoricensis.
rufescens Kze.; J. Sm. Lond. JoB. **1**. 200. 1842; Kze. Bot Zeit. **1850**. 385 } = L. portoricensis.
sagittata Dry. Tr. Linn. Soc. **3**. 40. 1797 = Schizoloma.
Sarasinorum Christ, Verh. Nat. Ges. Basel **11**. 429. 1896; Ann. Jard. Buit. **15**. 101 t. 14 f. 13. 1897 — Celebes. Ins. Philipp.
scandens Hk. sp. **1**. 205 t. 63 B. 1846. HB. 106. Chr. 293. NPfl. 221 — Malesia.
Schizoloma Ettingsh. Farnkr. 213 t. 145 f. 4; t. 146 f. 6. 1865 = Schizoloma ensifolium.
Schlechtendalii Christ, Farnkr. 296. 1897 = Odontosoria.
Schomburgkii Kl. Linn. **18**. 545. 1844 — Guiana.
scoparia Mett. Ann. sc. nat. IV. **15**. 64. 1861 = Odontosoria.
sectorifolia Goldm. Nova Acta **19**. Suppl. I. 464. 1843 (patria indic. erronea) = Adiantum flabellulatum.
securifolia Pr. Tent. 131. 1836; Goldm. Nova Acta **19**. Suppl. 464. 1843; Kze. Farnkr. **1**. 204 t. 85 f. 1. 1846 = L. gracilis.
Seemannii J. Sm. Bot. Voy. Herald 239. 1854. HB. 105 — Panama.
Seemanni Carr. in Seem. Fl. Vit. 338. 1873 = L. decomposita var.
semilunata C. Chr. Ind. 397. 1906 — Brasilia.
L. lancea var. semilunata C. Chr. Bot. Tidsskr. **25**. 81. 1902.
serpens Wall. List n. 150. 1828 = L. decomposita.
serrata Goldm. Nova Acta **19**. Suppl. I. 464. 1843 = ? L. stricta.
Spruceana Mett.; Kuhn, Linn. **36**. 79. 1869 = L. tarapotensis.
Sprucei Hk. Syn. ed. I. 108. 1867 = *Odontosoria meifolia*.
stolonifera Mett.; Kuhn, Linn. **36**. 81. 1869 — Polynesia.
Davallia Bak. 1874. HB. 468 (an eadem ac Synaphlebium pulchrum Brack. 1854; Lindsaea Carr. 1873? vide L. decomposita).
striata Bl. Enum. 220. 1828 = L. orbiculata.

stricta (Sw.) Dry. Tr. Linn. Soc. **3**. 42. 1797. HB. 108. Chr. 293. NPfl. 221 — Amer. trop. — Adiantum Sw. Prod. 135. 1788. Lindsaya rigescens Willd. 1810 — L. elegans Hk. 1837 — L. pusilla Splitgb. 1840 — (f. Lindm. Ark. f. Bot. **1**. 199 = L. guianensis forma).

sublobata Kze. Linn. **18**. 121. 1844 = Schizoloma ensifolium.

tarapotense C. Chr. Ind. 398. 1906 — Peru. — L. Spruceana Mett.; Kuhn, Linn. **36**. 79. 1869. HB. 108. Chr. 293. NPfl. 221 (non L. Sprucei Hk. 1867).

tenera Dry. Tr. Linn. Soc. **3**. 42 t. 10. 1797 = L. orbiculata var.

tenera Klf. Enum. 219. 1824 = L. dubia.

tenuifolia Bl. Enum. 219. 1828 — Malesia. Polynesia. — Odontoloma Pr. 1849; Acrophorus Moore 1857 (pt.); Davallia triquetra Bak. 1867. HB. 93; Odontoloma J. Sm. 1875; Lindsaya Mett.; Chr. 1897. 294. NPfl. 222.

tenuifolia Mett. Ann. sc. nat. IV. **15**. 64. 1861 = Odontosoria chinensis.

tenuis Kl. Linn. **18**. 550. 1844. NPfl. 221 — Guiana.
L. filiformis Hk. 1846. HB. 106.

Thwaitesii Mett.; Kuhn, Fil. Deck. 18. 1867 (nomen) = L. venusta.

trapeziformis Dry. Tr. Linn. Soc. **3**. 42 t. 9. 1797 = L. lancea.

trichomanoides Dry. Tr. Linn. Soc. **3**. 43 t. 11. 1797 = L. cuneata.

tricrenata Bak. JoB. **1890**. 106 — N. Guinea.

trilobata Col. Tr. N. Zeal. Inst. **16**. 345. 1884 = L. linearis var.

trilobata Bak. JoB. **1891**. 107 = L. Hosei.

tripartita Bl. Enum. 219. 1828 = L. decomposita.

triquetra Mett.; Christ, Farnkr. 294. 1867 = L. tenuifolia.

truncata Pr. Tent. 131. 1836 = Schizoloma Guerinianum.

Ulei Hier.; Christ, Hedwigia **44**. 365. 1905 — Amazonas.

uncinella Krug, Engl. Jahrb. **24**. 92. 1897 = Odontosoria.

variabilis Hk. et Arn. Bot. Beech. Voy. 257 t. 52. 1836-40 = Schizoloma heterophyllum.

venusta Klf.; Kuhn, Linn. **36**. 79. 1869. HB. 471 — Ceylon.
L. Thwaitesii Mett.; Kuhn 1867 (nomen).

Vieillardii Mett. Ann. sc. nat. IV. **15**. 65. 1861 = Schizoloma.

virescens Sw. Vet. Ak. Handl. **1817**. 73 t. 4 f. 4. HB. 106. NPfl. 221 — Brasilia.
L. genkofolia St. Hil. 1833. L. Gardneri Hk. 1846 — L. Catharinae Hk. 1846. HB. 106.

viridis Col. Tasm. Journ. **2**. 174. 1844; Bak. JoB. **1875**. 110 — N. Zealand.
Odontosoria Kuhn 1882.

vittata Zoll. Nat. et Geneesk. Arch. **1**. 400. 1844 = Schizoloma ensifolium.

Walkerae Hk. sp. **1**. 209 t. 69 A. 1846 = Schizoloma.

LINDSAYNIUM Fée, Gen. Fil. 333. 1850-52 = **Lindsaya.**

rigidum Fée, Gen. 333 t. 27 bis f. C. 1850-52.

LINDSAYOPSIS Kuhn, Die Chaetopterides 347. 1882 = **Odontosoria.**

aculeata Kuhn, Chaetopt. 347. 1882.

divaricata Kuhn, Chaetopt. 347. 1882 = O. Schlechtendalii.

scandens Kuhn, Chaetopt. 347. 1882.

LITOBROCHIA Presl, Tent. Pterid. 148. 1836 = **Pteris.**

aculeata Pr. Tent. 149. 1836 = P. spinosa.

affinis Fée, 11 mém. 25. 1866 — Guadeloupe.

Aghardii Fée, 11 mém. 25. 1866 — India occ.

alata Pr. Tent. 149. 1836.

altissima Fée. Gen. 135. 1850-52 = ? P. woodwardioides.

ampla Pr. Tent. 148. 1836 = ? P. grandifolia vel P. Haenkeana.
angustata Fée, Cr. vasc. Br. 1. 49 t. 11 f. 1. 1869 = P. macroptera.
appendiculata Gay, Fl. chil. 6. 491. 1853 = Histiopteris incisa.
arborescens Fée, Gen. 135. 1850-52 = P. Kunzeana.
areolata Moore, Proc. Hort. Soc. 1860; Lowe, New Ferns t. 57. 1861 — India. (an = P. comans?).
articulata Pr. Tent. 149. 1836 = Pellaea angulosa.
aurita J. Sm. JoB. 3. 405. 1841 = Histiopteris incisa var.
Balbisiana Fée, Gen. 135. 1850-52.
Beecheyana Moore, Ind. XLIV. 1857.
Berteroana Fée, Gen. 135. 1850-52 = P. comans var.
biaurita J. Sm. Cat. cult. ferns 37. 1857.
biformis Fée, Cr. vasc. Br. 1. 47. 1869 = P. spinosa.
Borbonica Fée, Gen. 136. 1850-52.
brasiliensis Pr. Tent. 149. 1836 = P. denticulata.
breviloba Moore, Ind. 220. 1861 = P. Luschnathiana.
brevinervis Fée, 11 mém. 27 t. 8 f. 2. 1866.
Brunoniana Pr. Tent. 149. 1836 = Histiopteris incisa.
camptocarpa Fée, Gen. 137. 1850-52 = P. podophylla.
chrysodioides Fée, Gen. 135. 1850-52.
cognata Pr. Tent. 150. 1836 (nomen) — ?
comans Pr. Tent. 149. 1836.
contracta (Lam.) Fée, Gen. 135. 1850-52 (nomen) — ? — Pteris Lam.
cordifolia Moore, Ind. 342. 1862 = Doryopteris c. (J. Sm.).
crassipes Fée, Gen. 135. 1850-52 = P. gigantea.
Cumingiana Pr. Tent. 149. 1836 (nomen) — ?
Currori J. Sm. Ferns br. and for. ed. II. 312. 1876 = Lonchitis C.
decipiens Moore, Ind. 342. 1862 = Doryopteris d.
decora Moore, Ind. 342. 1862 = Doryopteris d.
decurrens Pr. Tent. 149. 1836.
denticulata Pr. Tent. 149 t. 5 f. 20. 1836.
dimidiata Pr. Tent. 149. 1836 = ? P. semipinnata.
dissimilis Fée, Cr. vasc. Br. 2. 31 t. 90. 1872-73 = P. denticulata var.
divaricata Brack. Expl. Exp. 16. 108. 1854 = P. tripartita.
dura Moore, Ind. XLIV. 1857 = Pellaea d.
elata Fée, Gen. 135. 1850-52 = P. altissima.
elegans Fée, Gen. 135. 1850-52 = Histiopteris incisa.
Endlicheriana Fée, Gen. 134. 1850-52 = P. comans var.
Galeottii Fée, 8 mém. 75. 1857 = P. biaurita.
Gardneri Fée, Gen. 139. 1850-52.
gigantea Pr. Tent. 149. 1836.
grandifolia J. Sm. JoB. 4. 163. 1841.
grandis Fée, 8 mém. 75. 1857 = P. Kunzeana.
grandis Moore; (Hk. Bak. Syn. 167. 1867) = Doryopteris nobilis.
Haenkeana Pr. Tent. 149 t. 5 f. 24. 1836.
hastata Pr. Tent. 148. 1836 = Doryopteris sagittifolia.
hederacea Pr. Tent. 149. 1836 = Doryopteris varians.
hemipteris Fée, 8 mém. 76. 1857 = P. orizabae.
horizontalis Fée, Cr. vasc. Br. 1. 48 t. 12 f. 1. 1869 = P. decurrens.
inaequalis Fée, 11 mém. 26 t. 8 f. 1. 1866.

incisa Pr. Tent. 149. 1836 = Histiopteris i.
intermedia Pr. Tent. 149. 1836 = P. macroptera.
intermedia J. Sm. JoB. **3**. 405. 1841 = P. tripartita.
Karsteniana Fée, Gen. 135. 1850-52 = P. Kunzeana.
Kleiniana Fée, Gen. 135. 1850-52 = P. geminata.
Kunzeana Fée, Gen. 135. 1850-52.
lactuca Pr. Tent. 149. 1836.
lanceofolia J. Sm. Hist. Fil. 291. 1875 = P. lanceifolia.
leptophylla J. Sm. Bot. Mag. **72** Comp. 22. 1846.
lonchophora Moore, Ind. 246 1861 = Doryopteris l.
longibrachiata Fée, 11 mém. 25. 1866.
ludens Bedd. Ferns br. Ind. t. 27. 1865 = Doryopteris l.
macilenta J. Sm. JoB. **4**. 163. 1841.
macroptera J. Sm. JoB. **4**. 163. 1841.
Mannii J. Sm. Hist. Fil. 291. 1875.
marginata Pr. Tent. 149. 1836 = P. tripartita.
Mexicana Fée, Gen. 136. 1850 52.
microdictyon Fée, Gen. 138. 1850-52.
Milneana Carr. in Seem. Fl. Vit. 350. 1873.
Montbrisonis Fée, Gen. 137. 1850-52.
nobilis Moore, Gard. Chr. **1862**. 932 = Doryopteris n.
organensis Fée, Cr. vasc. Br. **1**. 240. 1869 = P. splendens.
Orizabae Fée, Gen. 135. 1850-52.
pallida Pr. Tent. 149. 1836 = Histiopteris incisa.
palmata Moore, Ind. 342. 1862 = Doryopteris pedata var.
papyracea Fée, Gen. 136. 1850-52 = P. splendens.
patens Gay, Fl. chil. **6**. 490. 1853 = Histiopteris incisa.
pedata Pr. Tent. 149 t. 5 f. 25. 1836 = Doryopteris p.
pedata Bedd. Ferns br. Ind. t. 26. 1865 = Doryopteris ludens.
pellucida Pr. Tent. 149. 1836 = P. woodwardioides.
podophylla Pr. Tent. 149. 1836.
polita J. Sm. JoB. **4**. 163. 1841.
prealta Fée, Cr. vasc. Br. **1**. 46 t. 11 f. 2. 1872 = P. splendens var.
prealta Moore, Ind. 342. 1862 = Doryopteris p.
propinqua J. Sm. Bot. Voy. Herald. 234. 1854.
Pseudo-Lonchitis Fée, Gen. 135. 1850-52 = P. tripartita var.
pulchra Pr. Tent. 149. 1836 = ? P. podophylla var.
Raddiana Pr. Tent. 149. 1836 = Doryopteris R.
Rottleriana Fée, Gen. 135. 1850-52 = P. biaurita.
sagittaefolia Pr. Tent. 148. 1836 = Doryopteris s.
Schiedeana Pr. Tent. 148. 1836 (nomen) = P. Kunzeana.
Sellowiana Pr. Tent. 149. 1836 (nomen) — ?
sericea Fée, Cr. vasc. Br. **1**. 48 t. 11 f. 3. 1869.
setifera Fée. Gen. 138. 1850-52 = P. podophylla var.
sinuata Brack. Expl. Exp. **16**. 110 t. 14. 1854 = Histiopteris incisa var.
Smithii Moore, Ind. 342. 1862 = Doryopteris ludens.
spinulifera J. Sm. Ferns br. and for. 193. 1866 = P. atrovirens.
spinulosa J. Sm. JoB. **4**. 163. 1841 = P. leptophylla.
splendens Pr. Tent. 149. 1836.
trifoliata Moore, Ind. 342. 1862 = Doryopteris t.

tripartita Pr. Tent. 150. 1836.
triplicata Fée, Gen. 135. 1850-52 = P. heteroclita.
tristicula Pr. Tent, 149. 1836 = P. denticulata.
Tussaci Fée, Gen. 137. 1850-52.
undulata Moore, Gard. Chr. **1867**. 348 = P. comans.
varians Pr. Tent. 149. 1836 = Doryopteris v.
varians Fée, Cr. vasc. Br. **1**. 49 t. 12 f. 2. 1869 = P. spinosa.
vespertilionis Pr. Tent. 149. 1836 = Histiopteris incisa.
Vieillardii Carr. in Seem. Fl. Vit. 349. 1873.
Wallichiana Pr. Tent. 150. 1836 = P. biaurita.
Wallichiana Fée, Gen. 135. 1850-52.
woodwardioides Pr. Tent. 149. 1836.

LITOLOBIUM Newman, The Phytologist **5**. 236. 1854 = **Dennstaedtia.**
punctilobulum Newm. l. c.

LLAVEA Lagasca, Gen. et Sp. 16. 1816 — (Polypodiaceae Gen. **74**).
cordifolia Lag. Gen. et Sp. 16. 1816. HB. 144. Chr. 155. NPfl. 279 — Mexico-Guatemala. — Allantodia Desv. 1827; Asplenium Spr. 1827; Allosorus Keys. 1873; Cryptogramme Prantl 1882; Ceratodactylis osmundoides J. Sm. 1839; Allosorus Karwinskii Kze. 1839; Botryogramme Fée 1850-52.

Lobium Keyserling, Pol. Cyath. Hb. Bung. 15. 1873 *(Asplenium §)* = Asplenium.

Lobochlaena Fée, Gen. Fil. 315. 1850-52 *(Cardiochlaena §)* = Aspidium.

LOMAGRAMMA J. Smith, Journ. Bot. **3**. 402. 1841; **4**. 152. 1841.
(an gen. propr.? species: lomarioides et perakensis).
lomarioides J. Sm. Hist. Fil. 143. 1875 = Leptochilus.
perakensis Bedd. Handb. Suppl. 107. 1892 = Leptochilus.
?polyphylla Brack. Expl. Exp. **16**. 83 t. 12 f. 3. 1854 = Polybotrya.
praestantissima Gris. Fl. br. W. Ind. 678. 1864 = Acrostichum.
pteroides J. Sm. JoB. **3**. 402. 1841; Hk. Gen. t. 98. 1842 = Leptochilus lomarioides.

LOMAPHLEBIA J. Smith, Hist. Fil. 182. 1875 = **Polypodium.**
linearis J. Sm. Hist. Fil. 183. 1875 = P. gramineum.

LOMARIA Willdenow, Mag. Ges. Nat. Fr. Berlin **3**. 160. 1809 = **Blechnum.**
acrodonta Fée, 8 mém. 70. 1857.
acrostichoides Klf.; Spr. Syst. **4**. 63. 1827 = Stenochlaena sorbifolia var.
aculeata Bl. Enum. 205. 1828 = Stenochlaena sorbifolia var.
acuminata Desv. Berl. Mag. **5**. 326. 1811 = B. attenuatum.
?acuminata Pr. Rel. Haenk. **1**. 52. 1825 = Stenochlaena palustris.
acuminata Bak. Syn. 481. 1874 = B. norfolkianum.
acuta Desv. Prod. 290. 1827.
adnata Bl. Enum. 205. 1828 = Plagiogyria a.
aggregata Col. Tr. N. Zeal. Inst. **20**. 223. 1888 = B. lanceolatum.
alata Kze.; Fée, Gen. 68. 1850-52 (nomen) — ?
alpina Spr. Syst. **4**. 62. 1827 = B. penna marina.
alta Hew.; Moore, Ind. XXIV. 1857 = B. lineatum.
alternans Col. Tr. N. Zeal. Inst. **28**. 616. 1896.
ambigua Fée, Gen. 68. 1850-52 = B. laevigatum.
andicola Phil. Anal. Univ. chil. **94**. 353. 1896.
andina Bak. Syn. 482. 1874.
angustifolia HBK. Nov. Gen. et Sp. **1**. 18. 1815 = B. polypodioides.
antarctica Carm. Tr. Linn. Soc. **12**. 513. 1818 = B. penna marina.
apodophylla Bak. JoB. **1885**. 104 = B. Hancockii.

araucana Phil. Anal. Univ. chil. **94**. 354. 1896.
arborescens Kl. et Karst. Linn. **20**. 347. 1844 = B. lineatum.
areolaris Harr. Journ. Linn. Soc. **16**. 28. 1877.
arguta Fée, 8 mém. 70. 1857 = Plagiogyria semicordata.
articulata F. Muell. Fragm. **5**. 187. 1866 = Plagiogyria euphlebia.
aspera Kl. Linn. **20**. 344. 1844.
attenuata Willd. sp. **5**. 290. 1810.
attenuata auctt. quoad pl. americ. trop. = B. meridense.
aurata Fée, 8 mém. 71. 1857 = ? B. Schomburgkii.
aurea Wall. List n. 38. 1828 = Onychium siliculosum.
auriculata Desv. Berl. Mag. **5**. 330. 1811 = B. punctulatum.
auriculata Bl. Enum. 202. 1828 = B. Blumei.
auriculata Bak. Syn. 481. 1874 = B. Fullagari.
australis Kze. Linn. **9**. 57. 1834 = B. penna marina.
australis Link, Fil. sp. 82. 1841.
Banksii Hk.fil. Fl. N. Zeal. **2**. 31 t. 76. 1854.
bella Phil. Linn. **29**. 104. 1857 = B. attenuatum.
Bibreae Sturm (Bibra, Beitr. Naturg. Chile 42. —) = B. arcuatum.
biformis Bak. Journ. Linn. Soc. **15**. 415. 1876.
binervata Desv. Berl. Mag. **5**. 315. 1811 = B. polypodioides.
biserrata Mert. et Lind.; Hk. sp. **3**. 19. 1860 = Plagiogyria semicordata.
?blechnoides Desv. Prod. 289. 1827 = B. auriculatum.
blechnoides Bory, Dup. Voy. Bot. **1**. 273. 1828 = B. valdiviense.
borealis Link, Hort. Berol. **2**. 80. 1833 = B. spicant.
Boryana Willd. sp. **5**. 292. 1810 = B. tabulare.
brasiliensis Raddi, Pl. bras. **1**. 50 t. 72, 72 bis 1825 = ? B. striatum.
brasiliensis Fourn. Mex. pl. **1**. 113. 1872 = B. ornifolium.
Bredemeyeriana Kl. Linn. **20**. 346. 1844 = Leptochilus serratifolius.
callosa Fée, Gen. 70. 1850-52 = Plagiogyria pycnophylla
campylotis Kze. Linn. **17**. 567. 1843 = B. occidentale var.
capensis Willd. sp. **5**. 291. 1810.
caruifolia Wall. List n. 39. 1828 = Onychium siliculosum.
castanea Mak. Bot. Mag. Tokyo **6**. 45. 1892 (nomen) — Japonia.
caudata Bak. Syn. 179. 1867 = B. Sprucei.
chilensis Klf. Enum. 154. 1824.
ciliata Moore, Gard. Chr. **1866**. 290 = B. Moorei.
cinnamomea Klf. Enum. 153. 1824 = B. tabulare.
circinnata Boj. Hort. Maur. 406. 1837 = B. attenuatum.
Colensoi Hk.fil. Ic. pl t. 627-628. 1844 = B. Patersoni*.
concinna Bak. JoB. **1885**. 103 = Plagiogyria stenoptera.
confinis Cord. Fl. Réun. 58. 1895.
confusa Fourn. Ann. sc. nat. V. **18**. 316. 1873 = B. capense.
contigua Carr. in Seem. Fl. Vit. 350. 1873; Fourn. Ann. sc. nat V. **18**. 315. 1873.
cordata Desv. Berl. Mag. **5**. 330. 1811 —?
coriacea Schrad. Gött. gel. Anz. **1818** 916. = B. tabulare.
coriacea Brack. Expl. Exp. **16**. 122. 1854 = B. Patersoni*.
costaricensis Bak. Syn. 482. 1874 = Plagiogyria c.
costaricensis Christ, Bull. Boiss. II. **4**. 1092. 1904.
crenata Pr. Rel. Haenk. **1**. 51. 1825 = B. spicant.
crenulata Moore (Hk. Bak. Syn. 180. 1867) = B. amabile.

Cumingiana Hk. Fil. Exot. sub t. 43. 1859 = B. Patersoni.
cuspidata Kze. Linn. **9**. 59. 1834 = B. acutum.
cycadifolia Colla, Mem. Ac. Torino **39**. 43 t. 71. 1836.
cycadoides Pappe et Raws. Syn. Fil. Afr. austr. 28. 1858.
Dalgairnsiae Pappe et Raws. Syn. Fil. Afr. austr. 27. 1858.
danaeacea Kze. Linn. **18**. 326. 1844.
decipiens Pappe et Raws. Syn. Fil. Afr. austr. 29. 1858 = B. attenuatum.
decomposita Don, Prod. Fl. Nep. 14. 1825 — Nepal.
Lomariobotrys Fée 1850-52. — (= Onychium japonicum ?).
decrescens Fée, Gen. 68. 1850-52; 7 mém. 24 t. 9 f. 1 = B. polypodioides.
deflexa Col. Tasm. Journ. **2**. 178. 1844 = B. vulcanicum.
deflexa Liebm. Vid. Selsk. Skr. V. **1**. 236 (seors. 84). 1849 = B. danaeaceum.
deflexa Bak. JoB. **1888**. 226 = B. Faberi.
deltoides Col. Tasm. Journ. **1**. 376. 1841; **2**. 177. 1844 = B. vulcanicum.
dendrophila Sod. Cr. vasc. quit. 118. 1893.
densa Klf. Enum. 151. 1824 = B. punctulatum.
dentata Kuhn, Forschr. Gazelle **4** Farne 6. 1889.
Deplanchei Bak. Syn. 481. 1874.
discolor Willd. sp. **5**. 293. 1810.
discolor var. natalensis Bak. Syn. 481. 1874 = B. inflexum.
distans Col. Tr. N. Zeal. Inst. **28**. 615. 1896 = B. Hillii.
divergens Kze. Linn. **9**. 57. 1834 = B. Plumieri.
diversifolia Bak. Syn. 181. 1867.
doodioides Brack. Expl. Exp. **16**. 124. 1854 = B. lanceolatum vel B. obtusatum.
Dregeana Fée, 10 mém. 9. 1865 = B. punctulatum.
duplicata Potts, Tr. N. Zeal. Inst. **9**. 491. 1877 = B. capense.
dura Moore, Gard. Chr. **1866**. 290.
egenolfioides Bak. Kew Bull. **1894**. 7.
elliptica Boj. Hort. Maur. 406. 1837 = Stenochlaena pollicina.
elongata Bl. Enum. 201. 1828 = B. Patersoni*.
emarginata Carr. in Seem. Fl. Vit. 350. 1873 = B. opacum.
ensiformis Liebm. Vid. Selsk. Skr. V. **1**. 234 (seors. 82). 1849.
eriopus Kze. Linn. **13**. 152. 1839 = Stangeria paradoxa (Cycadaceae).
euphlebia Kze. Bot. Zeit. **1848**. 521 = Plagiogyria e.
exaltata Fée, 11 mém. 10 t. 3. 1866 = B. L'Herminieri var.
eximia F. Muell. Fragm. **5**. 188. 1866 = B. capense.
falcata Spr. Syst. **4**. 62. 1827 = B. discolor.
falciformis Liebm. Vid. Selsk. Skr. V. **1**. 234 (seors. 82). 1849.
Fauriei Christ, Bull. Boiss. **4**. 666. 1896 = Plagiogyria F.
Feei Jenm. Bull. Dep. Jam. **1**. n. 43. 7. 1893.
fernandeziana Phil. Anal. Univ. chil. **1873**. 105 = B. attenuatum.
Fialhoi Fée et Glaz.; Fée, Cr. vasc. Br. **1**. 239 t. 7 f. 2. 1869 = Plagiogyria [semicordata.
filiformis A. Cunn. in Hk. Comp. Bot. Mag. **2**. 363. 1836.
Floresii Sod. Cr. vasc. quit. 115. 1893.
fluviatilis Spr. Syst. **4**. 64. 1827.
fragilis Liebm. Vid. Selsk. Skr. V. **1**. 232 (seors. 80). 1849 = B. polypodioides.
Fraseri A. Cunn. in Hk. Comp. Bot. Mag. **2**. 364. 1836.
fraxinea Willd. sp. **5**. 294. 1810 = Stenochlaena pollicina.
? fraxinifolia Raddi, Pl. bras. **1**. 51 t. 73. 1825 = Stenochlaena marginata.
Fullagari F. Muell. Fragm. **8**. 157. 1874.

Gayana Fée; Gay, Fl. chil. 6. 481. 1853; Fée, 7 mém. 25 t. 10 f. 1 = B. penna [marina.
Germainii Hk. sp. 3. 32 t. 152. 1860.
Gheisbreghtii Bak. Syn. 481. 1874 = B. Ghiesbreghtii.
gibba Lab. Sert. Austr. Cal. 3 t. 4, 5. 1824.
gigantea Klf. Enum. 150. 1824 = B. attenuatum.
Gilliesii Hk. et Grev. Ic. Fil. t. 207. 1831.
glauca Bl. Enum. 204. 1828. a. b. = Plagiogyria glauca.
c. = Plagiogyria pycnophylla.
Glaziovii Christ, Ann. Jard. Cons. bot. Genève 3. 42. 1899.
gracilis Bl. Enum. 204. 1828 = Stenochlaena g.
grandis Boj. Hort. Maur. 407. 1837 = Stenochlaena tenuifolia.
Gueinzii Moug.; Fée, Gen. 69. 1850-52 = B. tabulare.
? Haenkeana Pr. Rel. Haenk. 1. 53. 1825 = Stenochlaena palustris.
hamata Klf. Enum. 150. 1824 = B. attenuatum.
Hancockii Bak. Ann. Bot. 5. 220. 1891.
hastata Kze. Linn. 10. 508. 1836 = B. auriculatum.
heterophylla Desv. Berl. Mag. 5. 330. 1811 = B. attenuatum.
heterophylla Col. Tasm. Journ. 2. 175. 1844 = B. Patersoni*.
imperialis Fée et Glaz.; Fée, Cr. vasc. Br. 1. 21 t. 7 f. 1. 1869.
inflexa Kze. Linn. 18. 117. 1844.
integrifolia Klf.; Spr. Syst. 4. 63. 1827 = Stenochlaena pollicina.
intermedia Col. Tr. N. Zeal. Inst. 19. 274. 1887 = B. Hamiltonii.
?juglandifolia Pr. Rel. Haenk. 1. 52. 1825 = Stenochlaena palustris.
Krebsii J. Sm. Hist. Fil. 305. 1875 = B. punctulatum var.
lanceolata Spr. Syst. 4. 62. 1827.
lanceolata Colla, Mem. Ac. Torino 39. 45. 1836 = B. L'Herminieri.
lanuginosa Kze. Anal. 19. 1837 = B. cycadifolium.
latifolia Col. Tasm. Journ. 2. 175. 1844 = B. capense.
Lechleri Moore, Gard. Chr. 1866. 634 = B. valdiviense
Lenormandi Bak. Syn. 181. 1867.
Leyboldtiana Phil. Anal. Univ. chil. 1861. 68; Linn. 33. 303. 1864.
L'Herminieri Bory; Kze. Farnkr. 1. 173 t. 73. 1845.
? limonifolia Wall. List n. 35. 1828 = Stenochlaena palustris.
leariaefolia Pr. Rel. Haenk. 1. 52. 1825 — ? Peru.
linearis Col. Tasm. Journ. 1. 376. 1841; 2. 176. 1844 = B. penna marina.
lineata Willd. sp. 5. 290. 1810.
longifolia Klf. Enum. 153. 1824 = Stenochlaena sorbifolia var. 2.
longifolia Hk. et Arn. Bot. Beech. Voy. 257. 1836-40 — China.
longifolia Schlecht.; Mart. et Gal. Mém. Ac. Brux. 15. 49. 1842 = B. ornifolia.
loxensis HBK. Nov. Gen. et Sp. 1. 18. 1815 — Peru.
? *lucida* Pr. Rel. Haenk. 1. 52. 1825 — Luzon. — (certe Stenochlaena sp.).
magellanica Desv. Berl. Mag. 5. 330. 1811 = B. tabulare.
marattiaefolia Desv. Prod. 291. 1827 = Stenochlaena sorbifolia var. 1.
marginata Schrad. Gött. gel. Anz. 1824. 871 = Stenochlaena.
marginata Fée, Gen. 71. 1850-52 = B. Montbrisonis.
martinicensis Spr. Neu. Entdeck. 3. 5. 1822 = B. polypodioides.
Matsumureana Mak. Bot. Mag. Tokyo 8. 90 (pars jap.). 1894 = Plagiogyria M.
Mayi ht. May; Gard. 65. 330. 1904 — Hort.
melanocaulon Brack. Expl. Exp. 16. 122. 1854 = B. Patersoni*.
membranacea Col.; Hk. sp. 3. 34 t. 145. 1860.

meridensis Kl. Linn. **20**. 345. 1847.
mexicana Fée, 8 mém. 70. 1857 = B. polypodioides.
Meyeriana Kze. Linn. **10**. 509. 1836 = Stenochlaena tenuifolia.
microbasis Bak. JoB. **1880**. 328.
microphylla Goldm. Nova Acta **19** Suppl. I. 460. 1843 = B. penna marina.
Milnei Carr. in Seem. Fl. Vit. 351. 1873.
minor Spr. Syst. **4**. 65. 1827 = B. capense.
minor Link, Hort. Berol. **2**. 80. 1833 = B. australe.
mollis Zoll. et Moritz; Zoll. Syst. Verz. 40, 49. 1854 = Photinopteris speciosa.
Moritziana Kl. Linn. **20**. 347. 1847 = ? B. Schomburgkii.
mucronata Fée, Cr. vasc. Br. **1**. 20 t. 8 f. 3. 1869 = B. Plumieri.
mucronulata Fée, Gen. 72. 1850-52 — India orient. — Struthiopteris Trev. 1869.
neo-caledonica Lind. et Fourn. L'Ill. Hort. **23**. 132 t. 251. 1876.
nigra Col. Tasm. Journ. **1**. 375. 1841; **2**. 176. 1844.
niponica Kze. Bot. Zeit. **1848**. 508 = B. spicant var.
norfolkiana Hew. Lond. JoB. **1**. 122 nota. 1842.
nuda Willd. sp. **5**. 289. 1810 = B. discolor var.
obtusata Lab. Sert. Austr. Cal. 4 t. 6. 1824.
obtusifolia Pr. Tent. 143. 1836 = B. Schomburgkii.
oligoneuron Col. Tr. N. Zeal. Inst. **16**. 346. 1884 = B. membranaceum.
Olivieriana Fée, Gen. 70. 1850-52 — India.
onocleoides Spr. Syst. **4**. 62. 1827 = B. polypodioides.
opaca Bak. Syn. 176. 1867.
orbiculata Desv. Prod. 289. 1827 = Stenochlaena pollicina.
ornifolia Pr. Rel. Haenk. **1**. 51. 1825.
palmaeformis Desv. Prod. 290. 1827 = B. tabulare.
palustris Schrad. Gött. gel. Anz. **1824**. 871 = B. lineatum.
parvifolia Col. Tr. N. Zeal Inst. **20**. 224. 1888.
Patersonii Spr. Syst. **4**. 62. 1827.
paucijuga Col. Tr. N. Zeal. Inst. **20**. 222. 1888 = B. vulcanicum.
pectinata Liebm. Vid. Selsk. Skr. V. **1**. 233 (seors. 81). 1849 = Plagiogyria semicordata.
pedunculata Goldm. Nova Acta **19** Suppl. I. 460. 1843 = Angiopteris evecta.
Penna marina Trev. Atti Ist. Veneto III. **14**. 570. 1869.
petiolaris Sod. Cr. vasc. quit. 114. 1893.
pilosa Brack. Expl. Exp. **16**. 125 t. 15. 1854 = B. vulcanicum.
pimpinellifolia Hk.fil. Lond. JoB. **3**. 412. 1844 = B. filiforme.
Plumieri Desv. Berl. Mag. **5**. 325. 1811.
Plumieri Fée, 11 mém. 10 t. 4. 1866 = B. Feei.
polymorpha Reinw.; Zoll. Nat. Geneesk. Arch. **2**. 204. 1845; Hassk. Flora **1847**. 319 = Stenochlaena aculeata.
polypodioides Gaud. Ann. sc. nat. **5**. 98. 1825 = B. penna marina.
polypodioides Desv. Prod. 288. 1827.
procera Spr. Syst. **4**. 65. 1827 = B. capense.
propinqua A. Cunn. in Hk. Comp. Bot. Mag. **2**. 363. 1836 = B. filiforme.
propinqua Trev. Atti Ist. Veneto II. **2**. 167. 1851 (nomen) — Rio.
pteropus Kze. Farnkr. **1**. 97 t. 46. 1842.
pubescens Kze. Farnkr. **1**. 122 t. 55 f. 2. 1843 = B. auriculatum var.
pubescens Bak. Journ. Linn. Soc. **15**. 415. 1876 = B. Bakeri.
pumila Klf. Enum. 151. 1824 = B. australe.

pumila Raoul, Ann. sc. nat III. **2**. 115. 1844 = B. parvifolium.
punctata Bl. Enum. 201. 1828 = B. Patersoni*.
punctulata Kze. Linn. **10**. 507. 1836.
pycnophylla Kze. Bot. Zeit. **1848**. 143 = Plagiogyria p.
pygmaea Col. Tr. N. Zeal. Inst. **25**. 322. 1893.
pyrophila Bl. Enum. 204. 1828 = ? B. vestitum.
Reedei Phil. Anal. Univ. chil. **1873**. 581 = B. chilense.
Regnelliana Kze. Linn. **20**. 576. 1847 = B. lineatum.
rigida Fée, Gen. 68. 1850-52 = B. punctulatum.
rigida J. Sm. Ferns br. and for. 290. 1866 = B. durum.
Rimbachii Sod. Cr. vasc. quit. 117. 1893.
Rinconii Trev. Atti Ist. Veneto II **2**. 166. 1851 (nomen) — Guatemala.
robusta Carm. Tr. Linn. Soc. **12**. 512. 1818 = B. tabulare.
robusta Fée, Gen. 69. 1850-52 = B. tabulare.
? **rotundifolia** Bl. Enum. 204. 1828 = ? B. capense.
rotundifolia Col. Tasm. Journ. **1**. 377. 1841; **2**. 179. 1844; Raoul, Ann. sc. nat. III. **2**. 115. 1844 = B. fluviatile.
rufa Spr. Nova Acta **10**. 230. 1821 = ? B. Schomburgkii.
Ryani Klf. Enum. 155. 1824 = ? B. Schomburgkii.
salicifolia Kze. Linn. **9**. 58. 1834 = B. tabulare.
scabra Klf.; Sieb. Syn. Fil. exs. n. 107 = B. laevigatum.
scandens Willd. sp. **5**. 293. 1810 = Stenochlaena palustris.
scandens Raddi, Opusc. sci. Bol. **3**. 290. 1819 = Stenochlaena marginata.
Schiedeana Pr. Tent. 143. 1836 = B. ornifolium.
Schomburgkii Kl. Linn. **20**. 346. 1847.
Schottii Colla, Mem. Ac. Torino **39**. 44 t. 72. 1836 = ? B. tabulare.
secunda Wall. List n. 34. 1828 = Stenochlaena pollicina.
Sellowiana Pr. Tent. 142. 1836 = B. capense.
semicordata Bak. Syn. 182. 1867 = Plagiogyria s.
? **serpens** Wall. List n. 32. 1828 = Leptochilus axillaris.
serrulosa Desv. Prod. 290. 1827 — Peru.
sessilifolia Kl.; Christ, Bull. Boiss. II. **4**. 1092. 1904.
setigera Gaud. Ann. sc. nat. **5**. 98. 1825 = B. tabulare.
Sieberiana Pr. Tent. 142. 1836 (nomen) — ?
simillima Bak. JoB. **1884**. 141.
socialis Sod. Cr. vasc. quit. 122. 1893.
sorbifolia Klf. Enum. 151. 1824 = Stenochlaena s.
speciosa Bl. Enum 202. 1828 = Dryostachyum s.
spectabilis Kze. Bot. Zeit. **1848**. 144 = Stenochlaena sorbifolia var. 8.
spectabilis Liebm. Vid. Selsk. Skr. V. **1**. 235 (seors. 83). 1849 = B. ornifolium.
Spicant Desv. Berl. Mag. **5**. 325. 1811.
spicata Willd. sp. **5**. 289. 1810 = Hymenolepis s.
spissa Fée, Gen. 71. 1850-52 = B. tabulare.
spissa Christ. Bull. Boiss. II. **4**. 1092. 1904 = B. Christii.
splendens Hort.; Kze. Linn. **23**. 261. 1850 — Hort.
? **spondiaefolia** Wall. List n. 37. 1828 = Stenochlaena sorbifolia var. 3.
squamulosa Desv. Prod. 290. 1827.
stenophylla Kl. Linn. **20**. 346. 1847.
stenophylla Bak. JoB. **1884**. 142 = B. Humblotii.
stenoptera Bak. Ann. Bot. **5**. 220. 1891 = Plagiogyria s.

stipitellata Sod. Rec. 29. 1883.
stolonifera Fourn. Mex. pl. **1**. 113. 1872.
striata Willd. sp. **5**. 291. 1810.
subcordata Fourn. Ann. sc. nat. V. **18**. 316. 1873 = B. filiforme.
tabularis Mett.; Bak. Fl. bras. **1**^{2}. 418. 1870.
tenuifolia Desv. Berl. Mag. **5**. 326. 1811 = Stenochlaena t.
trichomanoides Desv. Prod. 287. 1827 = B. penna marina.
triloba Fée, Gen. 68. 1850-52 = B. auriculatum var.
triquetra Moore, Ind. 14. 1857 = Plagiogyria euphlebia.
tuberculata J. Sm. Cat. Kew Ferns 1856 = B. striatum.
uliginosa Phil. Linn. **29**. 105. 1857.
unguiculata Fée, Gen. 70. 1850-52.
variabilis Willd. sp. **5**. 294 1810 = Stenochlaena sorbifolia var. 1.
varians Fourn. Mex. pl. **1**. 113. 1872.
vestita Bl. Enum. 203. 1828.
Vieillardii Bak. Syn. 175. 1867.
villosa Fée, Gen. 68. 1850-52 = B. vulcanicum.
violacea Fée, 11 mém. 11 t. 5. 1866.
vittata Ad.Br.; Fourn. Ann sc. nat. V. **18**. 313. 1873.
volubilis Hk. sp. **3**. 39 t. 150. 1860 = B. volubile var.
vulcanica Bl Enum. 202. 1828.
Werckleana Christ. Bull. Boiss. II. **4**. 1091. 1904.
Wightiana Wall. List n. 2169. 1829 (nomen) —?
woodwardioides (Bernh.) Moore, Ind. 11. 1857 —?
Asplenium Bernh. 1802; Acrostichum lineare Spr. Schrad. Journ. **1799**^{2}. 267.
xiphophylla Bak. JoB. **1884**. 142.

LOMARIDIUM Presl, Epim. Bot. 154. 1849 = **Blechnum.**
Herminieri Pr. Epim. 263. 1849 = B. L'Herminieri.
Plumieri Pr. Epim 155. 1849.
?semicordatum Pr. Epim. 155. 1849 = Plagiogyria s.

LOMARIOBOTRYS Fée, Gen. Fil. 45. 1850-52 = **Stenochlaena.**
decomposita Fée, Gen. 46. 1850-52 = Lomaria d.
Meyeriana Fée, Gen. 46 t. 5 A. 1850-52 } = S. tenuifolia.
tenuifolia Fée, Gen. 46. 1850 52 } = S. tenuifolia.

Lomariocycas J. Smith, Hist. Fil. 305. 1875 (*Lomaria §*) = Blechnum.

LOMARIOPSIS Fée, Hist. Acrost. 10. 66. 1845 = **Stenochlaena.**
acuminata Fée, Gen. 45. 1850-52 = L. Pervillei.
Balansae Fourn. Ann. sc. nat. V. **18**. 271 1873 = S. sorbifolia var. 15.
Boryana Fée, Acrost. 68. 1845 = S. pollicina.
Brackenridgei Carr. in Seem. Fl. Vit. 373. 1873 = S. sorbifolia var. 5.
Brightiae Bail. Hdb. Queensl. Ferns 10. 1874 = S. sorbifolia var. 13.
buxifolia Fée, Acrost. 69. 1845 = S. pollicina var.
cochinchinensis Fée, Acrost. 66 t. 26. 1845 = S. sorbifolia var. 3.
cuspidata Fée. Acrost. 68 t. 27. 1845 = S. sorbifolia var. 1.
decrescens Kuhn, Verh. zool bot. Ges. **19**. 571. 1869.
elongata Fée, Acrost. 67. 1845 = S. sorbifolia var. 4.
erythrodes Fée, Acrost. 67. 1845 = S. marginata.
Fendleri Eat. Mem. Amer. Ac. n. s. **8**. 195. 1860 = S. sorbifolia var. 10.
fraxinea J. Sm. Ferns br. and for. 284. 1866 = S. pollicina.

guineensis Kuhn, Fil. Afr. 53. 1868 (nomen) — Africa occ. trop.
Acrostichum Carr. 1901.
Hamiltoniana Mett.; Salom. Nom. 221. 1883 = Polybotrya appendiculata var.
heteromorpha Moore, Ind. XVII. 1857 = Blechnum filiforme.
Horsfieldii Mett. Fil. Lips. 22. 1856 = Photinopteris speciosa.
Hügelii (Fée) Pr. Epim. 263. 1849 — N. Zealand. — Stenochlaena Fée hb.
leptocarpa Fée, Acrost. 69 t. 29. 1845 = S. sorbifolia var. 5.
longifolia J. Sm. Cat. cult. Ferns 25. 1857 = S. sorbifolia var. 7.
ludens Fée, Acrost. 70 t. 30. 1845 = S. sorbifolia var. 20.
macrophylla Mett.; Salom. Nom. 221. 1883 (nomen) — ?
marginata Kuhn, v. Deck. Reis. **3**[3]. Bot. 22. 1879.
Novae-Caledoniae Mett Ann. sc. nat. IV. **15**. 58. 1861 = S. sorbifolia var. 12.
oleandraefolia Mett.; Kuhn, Verh. zool. bot. Ges. **19**. 571. 1869 = S. sorbifolia var. 9.
palustris Mett.; Kuhn, Fil. Afr. 53. 1868 = Elaphoglossum p.
palustris Kuhn, Ann. Lugd. Bat. **4**. 294. 1869.
Pervillei Mett.; Kuhn, Fil. Afr. 53. 1868 — Ins. Sechellae.
phlebodes Fée, Acrost 66. 1845 = S. marginata.
Pittieri Christ, Prim. Fl. Costar. **3**. 9. 1901.
pollicina Mett.; Kuhn, Fil. Afr. 53. 1868.
polyphylla Kuhn, Fil. Afr. 54. 1868 = S. decrescens.
polyphylla Kuhn, Verh. zool. bot. Ges. **19**. 571. 1869 = Polybotrya p.
Prieuriana Fée, Acrost. 66 t. 25 f. 1. 1845 = S. marginata.
Raddiana Mett. Fil. Lips. 22. 1856 = Leptochilus guianensis.
recurvata Fée, Acrost. 68 t. 28. 1845 = S. sorbifolia var. 6.
Requiniana Kuhn, Verh. zool. bot. Ges. **19**. 571. 1869 = Platytaenia R.
scandens Mett. Fil. Lips. 22. 1856 = S. palustris.
Seemanni Carr. in Seem. Fl. Vit. 373. 1873 = S. sorbifolia var. (Fiji. Aneiteum).
Smithii Fée, Arost. 71 t. 33 f. 2, t. 53. 1845 = S. sorbifolia var. 7.
sorbifolia Fée, Acrost. 69. 1845.
spectabilis Mett. Fil. Lips. 22. 1856 = S. sorbifolia var. 8.
spinescens Fée, Acrost. 71 t. 33 f. 1. 1845 = S. aculeata.
spondiaefolia Moore, Ind. 4. 1857 = S. sorbifolia var. 20.
tenuifolia Christ, Farnkr. 42. 1897.
triquetra Ettingsh. Farnkr. 13 t. 5 f. 1. 6, t. 6. f. 5. 1865 = Plagiogyria euphlebia.
undulata Mett. Fil. Lips. 22. 1856 = Leptochilus scalpturatus.
variabilis Fée, Acrost. 70 t. 31, 32. 1845 = S. sorbifolia var. 1.
vestita Fourn. Bull. Soc. Fr. **19** 250. 1872 = S. sorbifolia var 14.
Wrightii Mett.; Eat. Mem. Amer. Ac. n. s. **8**. 195. 1860 = S. sorbifolia var. 11.
Yapurensis J. Sm. Hist. Fil. 140. 1875 = S. marginata.

Lonchitidium Fée, Gen. Fil. 126. 1850-52 (*Pteris* §) = Lonchitis.

LONCHITIS Linné (Gen. Plant. 781); Spec. pl. **2**. 1078. 1753 — (Polypodiaceae Gen. **85**).

?Adscensionis Forst. Comm. soc. reg. Gött. **9**. 72. 1789 = Pteris dentata var.
anthriscifolia Bory in Willd. sp. **5**. 461. 1810 (syn.); Desv. Prod. 306 1827 [= Dennstaedtia.
aurita L. sp. **2**. 1078. 1753 — Amer. trop.
L. Lindeniana Hk. sp. **2**. 56 t. 89 A. 1851. NPfl. 295; Pteris Christ 1901.
Lonchitis pubescens pt. HB. 128.
bipinnata Forsk. Fl. Aegypt.-Arab. 184. 1775 = ? Asplenium achilleifolium.
caffrorum Bernh. Schrad. Journ. **1800**[2]. 124. 1801 = Mohria.

Currori (Hk.) Mett.; Kuhn, v. Deck. Reis. 3^3. Bot. 10. 1879. NPfl. 295 — Africa trop. Madagascar. — Pteris Hk. sp. 2. 232 t. 140. 1858. HB. 168. Chr. 174; Litobrochia J. Sm. 1876.

Ghiesbreghtii Linden, Cat. —. = L. hirsuta.

glabra Bory, Voy. 1. 321. 1804 — Africa austr. Réunion.
Pteris Mett. 1856 — Lonchitis stenochlamys Fée 1850-52 — L. natalensis Hk. 1851; Pteris Kuhn 1868. L. pubescens β HB. 128.

hirsuta L. sp. 2. 1078. 1753. NPfl. 295 — Amer. trop.
Pteris J. Sm. 1841. Chr. 173; Antiosorus Kuhn 1882; Pteris lacinata Willd. 1810. HB. 160; Anisosorus Trev. 1851. ? Pteris villosa Sw. 1801.

hirsuta Bory, Voy. 1. 238, 321. 1804 = L. pubescens.

hirsuta Bak. Fl. bras. 1^2. 385. 1870 = L. aurita.

Javanica Desr. in Lam. Enc. 3. 594. 1789 = ? L. pubescens.

Lindeniana Hk. sp. 2. 56 t. 89 A. 1851 = L. aurita.

macrochlamys Fée, Gen. 142. 1850-52 = L. aurita.

macrophylla ht. Wendl. (Salom. Nom. 222. 1883) — Hort. (Réunion).

madagascariensis Hk. sp. 2. 58 t. 87 B. 1851 — Madagascar. Ins. Comorae.
Pteris Kuhn 1868.

Natalensis Hk. sp. 2. 57 t. 89 B. 1851 = L. glabra.

occidentalis Bak. Syn. 128. 1867. NPfl. 295 — Africa trop.
Pteris Kuhn 1868; Antiosorus Kuhn 1882; Pteris glaberrima Kuhn 1868.

pedata L. sp. ed. II. 2. 1536. 1763 = Pteris podophylla.

polypodioides Vell. Fl. flum. 11. t. 100. 1827; Arch. Mus. Nac. Rio Jan. 5. 456. 1881 = Dryopteris reticulata.

polypus Bak. Journ. Linn. Soc. 15. 414. 1876 — Madagascar.

pubescens Willd.; Klf. Enum. 195. 1824. HB. 128 (pt.) NPfl. 295 — Afr. trop. et austr. c. ins. — Pteris Kuhn 1868. Chr. 174; Phegopteris Keys. 1873; Lonchitis hirsuta Bory, Voy. 1. 321. 1804 (non L. 1753). L tomentosa Fée 1850-52 — ? L. Javanica Desr. 1789.

repens L. sp. 2. 1078. 1753 = Hypolepis.

stenochlamys Fée, Gen. 142. 1850-52 = L. glabra.

tenuifolia Forst. Prod 80. 1786 = Hypolepis.

tomentosa Fée, Gen. 143. 1850 52; 7 mém. 32 t. 23 f. 3 = L. pubescens.

LOPHIDIUM Richard, Act. Soc. hist. nat. Paris 1. 114. 1792 = **Schizaea.**

elegans Pr. Suppl. 77. 1845.

Flabellum Pr. Suppl. 77. 1845 = S. elegans var.

latifolium Rich. Act. soc. hist. nat. Paris 1. 114. 1792 = S. elegans.

pacificans Pr. Suppl. 77. 1845.

spectabile Pr. Suppl. 76. 1845 = S. pacificans var.

LOPHODIUM Newman, The Phytologist 4. 371, app. XVI. 1851 = **Dryopteris.**

abbreviatum Newm. Phyt. 4. app. XXI = D. filix mas.

Callipteris Newm. l. c. 421 = D. cristata.

collinum Newm. l. c. app. XVIII = D. spinulosa* dilatata.

erosum Newm. l. c. app. XXI = D. filix mas.

Filix mas Newm. l. c. app. XX.

Foenisecii Newm. l. c. app. XVI = D. aemula.

glanduliferum Newm. Phytol. 4. 371. 1851 }
glandulosum Newm. l. c app. XVIII. } = D. spinulosa* dilatata.
multiflorum Newm. l. c. 371. }

recurvum Newm. l. c. 371 = D. aemula.

rigidum Newm. l. c. app. XXI.
spinosum Newm. l. c. 371 = D. spinulosa.
uliginosum Newm. l. c. 371 = L. cristata × spinulosa.

LOPHOLEPIS J. Smith, Lond. Journ. Bot. 1. 195. 1842 = **Polypodium.**
Goniophlebium § Lopholepis J. Sm. in Hook. Gen. t. 51. 1840.
ciliata J. Sm. Lond. JoB. **1**. 195. 1842.
piloselloides J. Sm. Bot. Voy. Herald 229. 1854.
vacciniifolia J. Sm. Ferns br. and for. 85. 1866.

LOPHOSORIA Presl, Abh. böhm. Ges V. **5**. 344. 1848 = **Alsophila.**
acaulis Fée, Cr. vasc. Br. **1**. 174. 1869
affinis Pr. Abh. böhm. Ges V. **5**. 345. 1848
brasiliensis Kl. Allg. Gartenzeit. **1856**. 107. } = A. quadripinnata.
caesia Fée, Cr. vasc. Br. **1**. 172. 1869.
densa Kl. Allg. Gartenzeit. **1856**. 107
discolor Pr. Abh. böhm. Ges. V. **5**. 345. 1848
dorsalis Fée, Cr. vasc. Br. **1**. 173 t. 51 f. 3. 1869.
frigida Kl. Allg. Gartenzeit. **1856**. 107
glauca Kuhn, Engl. Jahrb. **24**. 83. 1897 (syn.)
glaucescens Kl. Allg. Gartenzeit. **1856**. 107.
polypodioides Pr. Abh. böhm. Ges. V. **5**. 345, 346. 1848 } = A. quadripinnata.
prostrata Fée, Cr. vasc. Br. **1** 173. 1869
pruinata Pr. Abh. böhm. Ges. V. **5**. 345. 1848
Schaffneriana Fourn. Mex. pl. **1**. 134. 1872.
Warscewiczii Kl.: Fourn. Mex. pl. **1**. 134. 1872 = A. quadripinnata.

LORINSERIA Presl, Epim. Bot. 72. 1849 = **Woodwardia.**
areolata Pr. Epim. 72. 1849.
Harlandii J. Sm. Hist. Fil. 311. 1875.
thelypteroides Pr. Epim. 72. 1849 = W. virginica.

LOTZEA Klotzsch et Karsten. Linnaea **20**. 358. 1847 = **Diplazium.**
diplazioides Kl. et Karst Linn. **20**. 358. 1847 = D. Klotzschii.

Loxochlena J. Smith, Hist. Fil. 304. 1875 *(Lomaria §)* = Blechnum (punctulatum *Krebsii).

LOXOGRAMME (Blume) Presl, Tent. Pterid. 214. 1836 = **Polypodium.**
Antrophyum § Loxogramme Bl. Fl. Jav. Fil. 73. 1828.
avenia Pr. Tent. 215. 1836 = P. Blumeanum.
Blumeana Pr. Tent. 215. 1836.
conferta Copeland, Polyp. Philipp. 68. 1905 = P. Copelandii.
coriacea Pr. Tent. 215. 1836 = P. loxogramme.
elongata Moore, Ind. LXV. 1857 = P. lanceolatum*.
flavescens Pr. Tent. 215. 1836 = ? P. scolopendrinum.
involuta Pr. Tent. 215. 1836 = P. scolopendrinum.
lanceolata Pr. Tent. 215 t. 9 f. 8. 1836 = P. loxogramme.
lineata Pr. Tent. 215. 1836 = Antrophyum l.
macrophylla Moore, Ind. LXV. 1857 = P. scolopendrinum.
parallela Copeland in Perkins, Fragm. 182. 1905.
?scolopendrina Pr. Tent. 215. 1836.

LOXOMA vide Loxsoma.

LOXOSCAPHE Moore, Journ. Bot. **5**. 227. 1853 = **Asplenium.**
brachycarpum Kuhn, v. Deck. Reis. 3^3. Bot. 36. 1879.
concinnum Moore, JoB. **5**. 227. 1853 = A. theciferum*.

foeniculaceum Moore, Ind. 295. 1861 = A. stenolobum.
gibberosum Moore, JoB. **5**. 227. 1853.
Lindeni Moore, JoB. **5**. 227. 1853 = A. theciferum var.
Mannii Kuhn, v. Deck. Reis. **3**[3]. Bot. 37. 1879 = Davallia M.
nigrescens Moore, Ind. 297. 1861 = A. hypomelas.
Schimperi Moore, JoB. **5**. 227. 1853 = A. theciferum*.
theciferum Moore, Ind. 302. 1861.

LOXSOMA R. Brown; A. Cunn. in Hook. Comp. to Bot. Mag. **2**. 366. 1836 (Loxoma) — (? Hymenophyllaceae Gen. **3**).
Cunninghami R. Br.; A. Cunn. Comp. Bot. Mag. **2**. 366 t. 31-32. 1836. HB. 56. Chr. 306. NPfl. 113 — N. Zealand.

LOXSOMOPSIS Christ, Bull. Boiss. II. **4**. 399. 1904 — (? Hymenophyllaceae Gen. 4).
costaricensis Christ, Bull. Boiss. II. **4**. 399 t. 1. 1904 — Costa Rica.
Lehmannii Hier. Engl. Jahrb. **34**. 435. 1904 — Ecuador.

LUERSSENIA Kuhn; Luerss. Bot. Centralbl. **11**. 77. 1882 — (Polypodiaceae Gen. **10**).
kehdingiana (Luerss.) Kuhn, Bot. Centralbl. **11**. 77. 1882. NPfl. 181 — Sumatra. *Aspidium* Luerss. msc. l. c. Chr. 235.

LYGODICTYON J. Smith in Hook. Gen. Fil. t. 111 B. 1842 = **Lygodium.**
Forsteri J. Sm. in Hk. Gen. t. 111 B. 1842 = L. reticulatum.
heterodoxum J. Sm. Ferns br. and for. 259. 1866.
lanceolatum J. Sm. Ferns br. and for. ed. II. 323. 1877.
reticulatum J. Sm. Hist. Fil. 352. 1875.
Schkuhrii J. Sm. Lond. JoB. **2**. 384. 1843 = L. reticulatum.

LYGODIUM Swartz, Schrad. Journ. **1800**[2]. 106. 1801 — (Schizaeaceae Gen. **2**).
acuminatum Sturm. Fl. bras. **1**[2]. 174 t. 14 f. 12. 1859 = L. volubile.
articulatum A. Rich. Fl. N. Zél. 96 t. 15. 1832. HB. 437. Prantl, Schiz. 63. Chr. 354. NPfl. 364 — N. Zealand.
Boivini Kuhn, Fil. Afr. 168. 1868. HB. 525. Prantl. Schiz. 84 — Ins. Comorae.
Brycei Bak. Kew Bull. **1901**. 138 — Afr. austr. (Rhodesia).
chaerophylloides Desv. Prod. 205. 1827 = L. japonicum.
circinnatum (Burm.) Sw. Syn. 153. 1806. Prantl. Schiz. 64. Chr. 353. NPfl. 366 — Asia trop. Queensland. — Ophioglossum Burm. Fl. Ind. 228. 1768; Hydroglossum Willd. 1802. Ophioglossum pedatum Burm. 1768; Lygodium Sw. 1801; Hydroglossum Willd. 1802. Ugena macrostachya Cav. 1801. U. dichotoma Cav. 1801; Lygodium Sw. 1806. HB. 437. Hydroglossum Willd. 1810. Ophioglossum flexuosum L. fil. 1781 (non L. 1753); Lygodium Sw. 1801 pt. — L. longifolium Sw. 1801; Hydroglossum Willd. 1802.
cochinchinense Desv. Prod. 206. 1827 = L. japonicum.
commutatum Pr. Suppl. 110. 1845 = L. mexicanum.
cubense HBK. Nov. Gen. et Sp. **1**. 31. 1815. Prantl, Schiz. 74. Chr. 356. NPfl. 366 — Cuba. — Hydroglossum Steud. 1824.
dichotomum Sw. Syn. 154. 1806 = L. circinnatum.
digitatum Pr. Rel. Haenk. **1**. 73. 1825. Prantl. Schiz. 66 — Ins. Philipp. Malacca.
digitatum Eat. Mem. Amer. Ac. n. s. **8**. 217. 1860 = L. radiatum.
dissectum Desv. Berl. Mag. **5**. 308. 1811 = L. japonicum.
elegans Desv. Berl. Mag. **5**. 308. 1811 = L. flexuosum.
expansum Desv. Prod. 204. 1827 = L. volubile var.
Finlaysonianum Wall. List n. 2002. 1829; Pr. Suppl. 112. 1845 = L. japonicum.

flexuosum (L.) Sw. Schrad. Journ. **1800**². 106 (pt.). 1801; Syn. 153. 1806. Prantl, Schiz. 72. Chr. 355. NPfl. 366 — China austr.-Malesia-Ins. Philipp. Queensland. Ophioglossum L. sp. **2**. 1063. 1753; Ramondia Mirb. 1801; Hydroglossum Willd. 1802. Ugena semihastata Cav. 1801; Lygodium Desv. 1827. HB. 437 — Hydroglossum pinnatifidum Willd. 1802 pt.; Lygodium Sw. 1803. HB. 438 pt. — Hydroglossum auriculatum Willd. 1802 — Lygodium semibipinnatum R. Br. 1810; Hydroglossum semipinnatum Poir. 1813 — Lygodium microstachyum et elegans Desv. 1811; Hydroglossum Steud. 1824 — L. pilosum Desv. 1827 — L. serrulatum Bl. 1828 — L. Meyerianum, Rottlerianum, tenue Pr. 1845.

Forsteri Lowe, New and rare ferns t. 35. 1861 = L. reticulatum.

gracile Bak. JoB. **1888**. 35 = L. oligostachyum.

gracilescens Col. Tr. N. Zeal. Inst. **28**. 620. 1896 = L. reticulatum.

hastatum Desv. Prod. 204. 1827 = L. volubile.

heptaphyllum Schrad. Gött. gel. Anz. **1824**. 863 = L. volubile.

heterodoxum Kze. Farnkr. **2**. 32 t. 113. 1849. HB. 439. Prantl, Schiz. 75. Chr. 353. NPfl. 366 — Mexico-Venezuela. — Hydroglossum Moore 1857; Lygodictyon J. Sm. 1866. Hydroglossum spectabile Liebm. 1849.

heterophyllum Pr. Rel. Haenk. **1**. 73. 1825 = L. circinnatum.

hians Fourn. Ann. sc. nat. V. **18**. 355. 1873; Diels, Hedwigia **44**. 133 c. fig. 1905 — N. Caledonia.

hirtum Klf. Enum. 47. 1824 = L. volubile var.

intermedium Mett.; Kuhn, Linn. **36**. 168. 1869 = L. volubile var.

japonicum (Thbg.) Sw. Schrad. Journ. **1800**². 106. 1801. HB. 439. Prantl, Schiz. 68. Chr. 356. NPfl. 366 — Japonia. China. Asia, Austr. trop. Ophioglossum Thbg. Fl. Jap. 328. 1784; Hydroglossum Willd. 1802 — Lygodium pubescens Klf. 1824 — L. dissectum Desv. 1811 — L. chaerophylloides et cochinchinense Desv. 1827 — L. tenue Bl. 1828.

Kerstenii Kuhn, Fil. Deck. 28. 1867. HB. 525 — Afr. orient. trop. Ins. Comorae. Madagascar. — L. subalatum Bojer; Kuhn, Fil. Afr. 170. 1868. HB. 438. Prantl, Schiz. 69. NPfl. 366.

laciniatum Lodd. Cat. 1849 — Hort.

lanceolatum Desv. Berl. Mag. **5**. 307. 1811. HB. 439. Prantl, Schiz. 80. NPfl. 366 Ins. Comorae. Madagascar. — Hydroglossum Steud. 1824; Lygodictyon J. Sm. 1877. Hydroglossum madagascariense Poir. 1813.

longifolium Sw. Schrad. Journ. **1801**². 305. 1803 = L. circinnatum.

lucens Klf. Enum. 47. 1824 = L. volubile.

maculatum Bory, Bél. Voy. Bot. **2**. 15. 1833 = L. flexuosum.

mexicanum Pr. Rel. Haenk. **1**. 72. 1825. (HB. 438). Prantl, Schiz. 70. NPfl. 366 — Amer. trop. — L. palmatilobum Sturm 1859.

Meyenianum Pr. Suppl. 106. 1845 = L. flexuosum.

micans Sturm, Fl. bras. **1**². 178. 1859. Prantl, Schiz. 79 — Ind. occ. Guinea.

microphyllum R. Br. Prod. Fl. Nov. Holl. 162. 1810 = L. scandens.

microphyllum Link, Hort. Berol. **2**. 141. 1833 — Hort.

microstachyum Desv. Berl. Mag. **5**. 308. 1811 = L. flexuosum.

mucronulatum Sturm, Fl. bras. **1**². 171 t. 14 f. 9. 1859 = L. polymorphum.

oligostachyum (Willd.) Desv. Prod. 205. 1827 — Ind. occ. Hydroglossum Willd. sp. **5**. 81. 1810. Lygodium gracile Bak. 1888.

palmatilobum Sturm, Fl. bras. **1**². 170 t. 14 f. 6. 1859 = L. mexicanum.

palmatum (Bernh.) Sw. Syn. 154. 1806. HB. 436. Prantl, Schiz. 64. Chr. 353. NPfl. 365 — U. S. A. atlant. — Gisopteris Bernh. Schrad. Journ. **1800**². 129 t. 2 f. 1. 1801; Hydroglossum Willd. 1802; Cteisium paniculatum Michx. 1803.

pedatum Sw. Schrad. Journ. **1800**². 106. 1801 = L. circinnatum.

pedatum Goldm. Nova Acta **19**. Suppl. I. 467. 1843 = L. volubile.

pilosum Desv. Prod. 205. 1827 = L. flexuosum.

pinnatifidum Sw. Schrad. Journ. **1801**². 303. 1803 = L. flexuosum et L. polystachyum.

pinnatifidum Prantl, Schiz. 83. 1881 = L. polystachyum.

Poeppigianum Pr. Suppl. 103. 1845 = L. cubense.

Pohlianum Pr. Suppl. 105. 1845 = L. polymorphum.

polycarpum Desv. Prod. 204. 1827 = L. reticulatum.

polymorphum (Cav.) HBK. Nov. Gen. et Sp. **1**. 31. 1815 — Amer. trop. Ugena Cav. Icon. **6**. 75 t. 595 f. 1. 1801; Lygodium venustum Sw. 1803. HB. 438. Prantl, Schiz. 71. Chr. 356. NPfl. 366 — Hydroglossum hastatum Willd. 1810 — Lygodium Pohlianum Pr. 1845 — L. mucronulatum Sturm 1859 — Hydroglossum hirsutum Willd. 1810.

polystachyum Wall. List n. 177. 1828; Moore, Gard. Chr. **1859**. 671. HB. 438. Chr. 355 — Malacca. Tenasserim.
L. pinnatifidum Prantl, Schiz. 83. 1881. NPfl. 366 (Sw. 1803 pt.).

puberulum Sturm, Fl. bras. **1**². 176. 1859 = L. volubile.

pubescens Klf. Enum. 47. 1824 = L. japonicum.

pubescens Wall. List n. 2200. 1829 = L. flexuosum.

radiatum Prantl, Schiz. 66. 1881. Chr. 353. NPfl. 366 — Guatemala-Columbia.
L. digitatum Eat. Mem. Amer. Ac. n. s. 8. 217. 1860. HB. 437 (non Pr. 1825).

reticulatum Schkuhr, Kr. Gew. **1**. 139 t. 139. 1809. HB. 439. Prantl, Schiz. 82. Chr. 355. NPfl. 366 — Polynesia. Australia.
Lygodictyon J. Sm. 1875. Ophioglossum scandens Forst. Prod. 78. 1786 (non L. 1753); Hydroglossum Pr. 1845. H. polycarpum Willd. 1810; Lygodium Desv. 1827. Lygodictyon Forsteri J. Sm. 1842; Lygodium Lowe 1861. Lygodictyon Schkuhrii J. Sm. 1843 — Lygodium gracilescens Col. 1896.

Rottlerianum Pr. Suppl. 108. 1845 = L. flexuosum.

salicifolium Pr. Suppl. 102. 1845. Prantl. Schiz 79 — India.
L. pinnatifidum pt. HB. 438.

scandens (L.) Sw. Schrad. Journ. **1800**². 106. 1801. HB. 437 (pt.); Prantl. Schiz. 81. Chr. 354. NPfl. 366 — Africa, Asia, Polynesia. Australia trop.
Ophioglossum L. sp. **2**. 1063. 1753; Odontopteris Bernh. 1801; Ramondia Mirb. 1801; Hydroglossum Willd. 1802 — Ugena microphylla Cav. 1801; Lygodium R.Br. 1810.

scandens Schkuhr, Kr. Gew. **1**. 138 t. 138. 1809 = L. volubile.

Schiedeanum Pr. Suppl. 110. 1845 = L. mexicanum.

semibipinnatum R.Br. Prod. Fl. N. Holl. 162. 1810 = L. flexuosum.

semihastatum Desv. Prod. 203. 1827 = L. flexuosum.

serrulatum Bl. Enum. 254. 1828 = L. flexuosum.

Smithianum Pr. Suppl. 112. 1845 (nomen); Prantl. Schiz. 80. 1881. Chr. 355. NPfl. 366 — Africa occ. trop. — L. pinnatifidum pt. HB. 438.

subalatum Bojer; Kuhn. Fil. Afr. 170. 1868 = L. Kerstenii.

surinamense Miq. »Diar. Inst. Bat. **1843**. 8« = ? L. volubile.

tenue Bl. Enum. 254. 1828 = L. japonicum.

tenue Pr. Suppl. 108. 1845 = L. flexuosum.

trifurcatum Bak. Syn. 437. 1868. Prantl, Schiz. 67. Chr. 353. NPfl. 366 — [Melanesia.
varium Link, Hort. Berol. **2**. 140. 1833 = L. volubile.
venustum Sw. Schrad. Journ. **1801**². 303. 1803 = L. polymorphum.
volubile Sw. Schrad. Journ. **1801**². 304. 1803. HB. 438 (pt.). Prantl, Schiz. 76. Chr. 355. NPfl. 366 — Amer. trop. — Hydroglossum Willd. 1810; Osmunda scandens Aubl. Hist. pl. Guiane **2**. 961. 1775; Lygodium Schkuhr 1809 (non supra) — L. lucens Klf. 1824 — L. hastatum Mart. 1834 — L. hirtum Klf. 1824 — Hydroglossum expansum Poir. 1813; Lygodium Desv. 1827.
Wrightii Mett.; Eat. Mem. Amer. Ac. n. s. **8**. 217. 1860 (nomen); Prantl, Schiz. 78. 1881 — Cuba.

Lytoneuron Klotzsch, Linnaea **20**. 343. 1847 *(Doryopteris §)* = Doryopteris.
Macroglena Presl, Abh. böhm. Ges. Wiss. V. **5**. 333. 1848 *(Trichomanes §)* = [Trichomanes.
MACROPLETHUS Presl, Epim. 141. 1849 = **Hymenolepis.**
platyrhynchos Pr. Epim. 142. 1849.
Macropteris Webb et Berth. Hist. nat. Canar. **3**². Phyt **3**. 450. 1847 *(Pteris §)* = Pteris.
MACROSTOMA Griffith, Asiatic Researches **19**. 110. 1836; Ic. plant. asiat. **2**. t. 137. 1849 = **Christensenia.**

MARATTIA Swartz, Prod. Fl. Ind. occ. 128. 1788 — (Marattiaceae Gen. **3**).
acuminata Willd.; Klf. Enum. 33 t. 1 f. 2. 1824 = M. attenuata.
alata Sw. Prod. 128. 1788. HB. 442. Chr. 358. NPfl. 142 — Amer. trop. Myriotheca Poir. 1797; Discostegia Pr. 1845 — Marattia laevis Sm. 1790. NPfl. 442; Myriotheca Poir. 1797; Discostegia Pr. 1845.
alata Raddi, Pl. Bras. **1**. 74 t. 83, 84. 1825 = M. Kaulfussii.
alata Hk. et Arn. Bot. Beech. Voy. 102. 1832 = M. Douglasii.
arguta Mett.; Kuhn, Linn. **36**. 168. 1869. HB. 525 — ? (Hort.).
Ascensionis J. Sm. Cat. cult. Ferns 79. 1857 = M. purpurascens.
attenuata Lab. Sert. Austr. Cal. 9 t. 13, 14. 1824. HB. 441. NPfl. 441 — N. Caledonia.
Boivini Mett.; Ettingsh. Farnkr. 246 t. 178 f. 15-16. 1865; Kuhn, Fil. Afr. 174. 1868. NPfl. 441 — Madagascar.
Burkei Bak. Gard. Chr. III. **22**. 425 f. 129. 1897 — ? Columbia.
calliodous de Vriese, Mon. 6 t. 3 f. 20, t. 4. f. 20. 1853 = M. fraxinea.
cicutifolia Klf. Enum. 32. 1824. HB. 441. Chr. 358. NPfl. 441 — Brasilia. Gymnotheca Pr. 1845.
Cooperi Moore; ht. Veitch, (Gartenfl. **13**. 154. 1864) — N. Caledonia.
cristata J. Sm. Cat. Kew Ferns 8. 1856 = M. purpurascens.
cuneiformis de Vriese, Mon. 7 t. 3 f. 21, t. 4 f. 21. 1853 = M. fraxinea.
Douglasii (Pr.) Bak. Syn. 441. 1868. Chr. 359. NPfl. 441 — Ins. Hawaii. Fiji. Stibasia Pr. Suppl. 16. 1845; Gymnotheca Moore 1857.
Dregeana Pr. Suppl. 9. 1845 = M. salicifolia.
elegans Endl. Prod. Fl. Norf. 6. 1833 = M. salicina.
fraxinea Sm. Pl. Ic. ined. 2. t. 48. 1790; Gmel. Syst. Nat. 2². 1294. 1791; HB. 440 (pt.). Chr. 358. NPfl. 441 — Asia, Africa, Austr. trop. Myriotheca Poir. 1797; M. fraxinifolia Bory 1804. M. sorbifolia Bory 1804; Marattia Sw. 1806.
fraxinea Raddi, Opusc. sci. Bol. **3**. 280. 1819; Pl. Bras. 1. 74 t. 82. 1825 = [M. Raddii.
? Haenkeana Pr. Suppl. 11. 1845 = ? M. pellucida.
Juergensii Rosenst. Festschr. Alb. v. Bamberg 68. 1905 — Brasilia austr.

Kaulfussii J. Sm. in Hk. Gen. t. 26. 1838. HB. 442. Chr. 359. NPfl. 442 — Amer. trop. — Eupodium J. Sm. 1842; Marattia laevis Klf. Enum. 31. 1824 (non Sm. 1790); M. alata Raddi 1825 (non Sw. 1788).

laevis Sm. Pl. Ic. ined. **2**. t. 47. 1790; Gmel. Syst. Nat. **2**^{2}. 1294. 1791 = [M. alata var.

laevis Klf. Enum. 31. 1824 = M. Kaulfussii.

Laucheana Blass. Hamb. Gartenzeit. (1858?); Gartenfl. **7**. 287. 1858 = M. alata.

laxa Kze. Linn. **18**. 306. 1844. HB. 441. Chr. 358 — Mexico.

Gymnotheca Pr. 1845.

macrophylla de Vriese, Mon. 3 t. 3 f. 17. 1853 = M. fraxinea.

melanesica Kuhn, Forschr. Gazelle **4** Farne 15 t. 3. 1889. NPfl. 442 — [N. Hannover.

Mertensiana (Pr.) C. Chr. Ind. 415. 1906 — Ins. Carolinae.

Gymnotheca Pr. 1848; Epim. 9 t. 1. 1849.

microcarpa Mett.; Ettingsh. Farnkr. 246 t. 117 f. 7, 8. 1865; Kuhn, Fil Afr. 175. 1868 — Ins. Comorae ? Afr. occ. trop.

microphylla Lodd. Cat. 1849 — Hort.

microphylla (Fée) Fourn. Mex. pl. **1**. 136. 1872 — Mexico.

Discostegia Fée, 9 mém. 43. 1857; Marattia Schaffneri Moore 1862.

natalensis Pr. Suppl. 9. 1845 = M. salicifolia.

obtusidens Sturm, Fl. bras. **1**^{2}. 151. 1859 = M. cicutifolia.

oppositifolia Sm. in Rees, Cycl. **22**. 1812 = M. fraxinea.

pellucida Pr. Suppl. 10. 1845. (HB. 441). NPfl. 441 — Ins. Philipp. N. Caledonia.

an = M. silvatica?

pinnata Roxb. Calc. Journ. **4**. 519. 1844 = ? Angiopteris crassipes.

podolepis (de Vriese) Sturm, Fl. bras. **1**^{2}. 151. 1859. (HB. 441) — Brasilia.

Gymnotheca de Vriese, Mon. 10. 1853.

polyodon (Pr.) Sturm, Fl. bras. **1**^{2}. 150. 1859 — Brasilia.

Gymnotheca Pr. Suppl. 14. 1845.

purpurascens de Vriese, Mon. 7 t. 3 f. 19, t. 4 f. 19. 1853. (HB. 441). NPfl. 441 [— Ascension.

M. Ascensionis J. Sm. 1857.

Raddiana Schott, Gen. t. 15. 1835 = M. Raddii.

Raddii Desv. Prod. 207. 1827 — Brasilia.

M. fraxinea Raddi, Opusc. sci. Bol. **3**. 280. 1819; Pl. Bras. **1**. 74 t. 82. 1825 (non Sm. 1790); M. Raddiana Schott 1834; Gymnotheca Pr. 1845.

salicifolia Schrad. Gött. gel. Anz. **1818**. 920. (HB. 440). NPfl. 441 — Afr. austr.

M. Dregeana Pr. 1845.

salicina Sm. in Rees, Cycl. **22**. —. 1812; de Vriese, Mon. 5 t. 3 f. 18, t. 4 f. 18. 1853. (HB. 440). NPfl. 441 — Norfolk. Ins. Societatis. N. Zealand.

M. elegans Endl. 1833.

sambucina Bl. Enum. 256. 1828. (HB. 441). NPfl. 441 — Malesia.

Schaffneri Moore, Ind. 341. 1862 = M. microphylla.

silvatica Bl. Enum. 256. 1828. (HB. 441) — Java.

Smithii Mett.; Kuhn, Verh. zool. bot. Ges. **19**. 584. 1869 — Polynesia.

sorbifolia Sw. Syn. 168. 1806 = M. fraxinea.

sorbifolia Brack. Expl. Exp. **16**. 312. 1854 = M. Smithii.

Stanleyana ht.; Rev. Hort. **1903**. 227 — Congo.

sylvatica = silvatica.

ternatea de Vriese, Mon. 4. 1853 — Ins. Molucc. ? N. Caledonia.

Verschaffeltiana (de Vriese) Sturm, Fl. bras. **1**^{2}. 153 t. 10. 1859 — Brasilia.

Gymnotheca de Vriese, Mon. 10. 1853.

weinmanniifolia Liebm. Vid. Selsk. Skr. V. **1**. 308 (seors. 156). 1849. NPfl. 442 [— Mexico.

Gymnotheca de Vriese 1853.

MARGINARIA Bory, Dict. class. d'hist. nat. **6**. 587. 1824; **10**. 176. 1826 = [**Polypodium.**
amoena Pr. Tent. 188. 1836.
angustifolia Pr. Tent. 188. 1836.
arguta Hk. Gen. ad t. 14. 1838.
argyrata (argentea) Bory, Dict. class **10**. 177. 1826.
auriseta Pr. Tent. 187. 1836 = P. tectum.
blepharoides Pr. Tent. 189. 1836 (nomen) — ? — Goniophlebium Moore 1862.
californica Pr. Tent. 188. 1836.
cardiophylla Pr. Tent. 188. 1836 = ? P. vacciniifolium.
Catharinae Pr. Tent. 188. 1836.
ceteracina Bory, Dict. class. **10**. 176. 1826 = P. polypodioides.
chnoodes Pr. Tent. 189. 1836.
Cumingiana Pr. Tent. 188. 1836 = ? P. tridens.
dasypleura Pr. Tent. 188 t. 7 f. 2. 1836.
dimorpha Link, Fil. sp. 119. 1841 = P. angustifolium.
distans Pr. Tent. 189. 1836 = P. fraxinifolium.
diversifolia Pr. Tent. 188. 1836 = P. lycopodioides.
ensifolia Pr. Tent. 188. 1836 = P. angustifolium var.
ensiformis Pr. Tent. 188. 1836.
falcata Fée, 1 mém. t. 2 f. 6. 1844 — ?
furfuracea Pr. Tent. 188. 1836.
gladiata Pr. Tent. 189. 1836 = P. attenuatum.
Haenkeana Pr. Tent. 187. 1836 = P. ciliatum Pr. (non Willd.).
harpeodes Pr. Tent. 188. 1836 = P. latipes.
incana Pr. Tent. 188. 1836 = P. polypodioides.
Karwinskyana Pr. Tent. 188. 1836 = P. plebeium.
laeta Pr. Tent. 188. 1836 = P. loriceum var.
latebrosa Pr. Tent. 189. 1836 = ? P. argutum.
latipes Pr. Tent. 188. 1836.
loricea Pr. Tent. 188. 1836.
marginata (scolopendrina) Bory, Dict. class. **10**. 176. 1826 = P. lanceolatum.
meniscifolia Pr. Tent. 189. 1836 = P. brasiliense var.
minima Bory, Dict. class. **10**. 177. 1826 = P. polypodioides.
neriifolia Pr. Tent. 189. 1836 = P. brasiliense.
nummularia Pr. Tent. 188. 1836 = P. pyrolifolium.
oleandrifolia Pr. Tent. 188. 1836 = P. attenuatum.
pellita Pr. Tent. 187. 1836.
piloselloides Pr. Tent. 187. 1836.
rhoifolia Pr. Tent. 189. 1836 = P. brasiliense.
rufula Pr. Tent. 189. 1836 = P. lepidopteris
serpens Pr. Tent. 188. 1836 = P. Swartzii.
squamata Pr. Tent. 188. 1836.
taeniosa Pr. Tent. 188. 1836 = P. angustifolium var.
vacciniifolia Pr. Tent. 188. 1836.
verrucosa Hk. Gen. t. 14. 1838.

Marginariopsis Moore. Index LXXIII. 1857 (*Phlebodium* §) = Polypodium.

MARSILEA Linné (Gen. pl. 799) Spec. plant. 2. 1099. 1753 — (Marsileaceae Gen. 1).
aegyptiaca Willd. sp. 5. 540. 1810. Bak. Hdb. 145. NPfl. 419 — Aegyptus. Zalusianskya O.Ktze. 1891. [Astrachan.
aegyptiaca Wall. List n. 7096. 1829 = M. minuta.

ancylopoda A.Br. Monatsber. Akad. Berl. **1863**. 434. 1864. Bak. Hdb. 146. NPfl. 420 — Ecuador. — Zalusianskya O.Ktze. 1891.

angustifolia R.Br. Prod. Fl. N. Holl. 167. 1810. Bak. Hdb. 141. NPfl. 420 — Austr. bor. — Zalusianskya O.Ktze. 1891.

Berteroi A.Br. Monatsber. Ak. Berl. **1870**. 747. Bak. Hdb. 146. NPfl. 420 — San Domingo. (-) Argentina. — Zalusianskya O.Ktze. 1891.

biloba Willd. sp. **5**. 540. 1810. Bak. Hdb. 144. NPfl. 419 — Africa austr. Zalusianskya O.Ktze. 1891. Marsilea glomerata Pr. 1845.

brachycarpa A.Br. Monatsber. Ak. Berl. **1863**. 420. 1864. NPfl. 418. (Bak. Hdb. 140) — Pegu.

brachypus A.Br. Monatsber. Ak. Berl. **1863**. 421. 1864. NPfl. 418. (Bak. Hdb. 140) — India austr.

brasiliensis Mart. Ic. Cr. Bras. 122 t. 73 f. 2. 1834 = M. polycarpa.

brevipes Nutt.; A.Br. Monatsber. Ak. Berl. **1870**. 741 = M. vestita.

Brownii A.Br. Monatsber. Ak. Berl. **1863**. 418. 1864. Bak. Hdb. 140. NPfl. 418 — N. S. Wales. — Zalusianskya O.Ktze. 1891; Marsilea quadrifolia R.Br. Prod. Fl. N. Holl. 167. 1810 (non L. 1753).

Burchellii (Kze.) A.Br. Monatsber. Ak. Berl. **1863**. 429. 1864. Bak. Hdb. 144. NPfl. 419 — Africa austr. — Zalusianskya O.Ktze. 1891; Marsilea quadrifolia γ Burchellii Kze. Linn. **10**. 556. 1836.

capensis A.Br. Monatsber. Ak. Berl. **1863**. 428. 1864. Bak. Hdb. 144. NPfl. 419 — Africa austr. — Zalusianskya O.Ktze. 1891.

concinna Bak. JoB. **1886**. 279. Hdb. 144. (NPfl. 420) — Paraguay. Uruguay. Argentina. — Zalusianskya O.Ktze. 1891.

condensata Bak. JoB. **1886**. 281. Hdb. 145. (NPfl. 420) — Scinde (India). Zalusianskya O.Ktze. 1891.

cormandelica Burm. vide coromandelica.

cornuta A.Br. Monatsber. Ak. Berl. **1870**. 728 = M. diffusa var.

coromandeliana auctt. vide coromandelica.

coromandelica Burm. Fl. Ind. t. 62 f. 3. 1768 (cormandelica). Bak. Hdb. 147 — India. Socotra. — Zaluzianskya O.Ktze. 1891; Marsilea quadrifolia Burm. Fl. Ind. 237. 1768 (non L. 1753); M. minuta β coromandeliana L. 1771; M. coromandelina Willd. 1810; M. coromandeliana auctt. NPfl. 420.

coromandelina Willd. sp. **5**. 539. 1810 = M. coromandelica.

crenata Pr. Rel. Haenk. **1**. 84 t. 12 f. 13. 1825 = ? M. minuta.

crenata A.Br. Monatsber. Ak. Berl. **1863**. 420. 1864 (pt.?) = M. crenulata.

crenulata Desv. Prod. 179. 1827. Bak. Hdb. 141. NPfl. 418 — Ins. Mascar. var. Senegambia. — Zalusianskya O.Ktze. 1891. Marsilea microcarpa A.Br. 1839 — M. senegalensis A.Br. 1839. Bak. Hdb. 141; Zalusianskya O.Ktze. 1891.

deflexa A.Br. Monatsber. Ak. Berl. **1863**. 421. 1864. Bak. Hdb. 140. NPfl. 418 Brasilia. var. Columbia. — Zalusianskya O.Ktze. 1891 — Marsilea striata Mett. 1865.

diffusa Leprieur; A.Br. Flora **1839**. 300; Monatsber. Ak. Berl. **1863**. 419. 1864. Bak. Hdb. 141. NPfl. 418 — Ins. Canar. Algeria-Africa trop. Ins. Mascar. Zalusianskya O.Ktze. 1891 — Marsilia cornuta A.Br. 1870.

distorta A.Br. Monatsber. Ak. Berl. **1863**. 433. 1864. Bak. Hdb. 147. NPfl. 420 — Senegambia. — Zalusianskya O.Ktze. 1891.

Dregeana A.Br. Monatsber. Ak. Berl. **1863**. 428. 1864 = M. macrocarpa.

Drummondii A.Br. Linn. **25**. 721. 1852. Bak. Hdb. 146 (pt.). NPfl. 419 — Australia. Zalusianskya O.Ktze. 1891; Marsilia macropus Hk. 1854 (non Engelm. 1848).

elata A.Br. App. Ind. Sem. Berol. 1867. 3; Monatsber. Ak. Berl. **1870**. 739. NPfl. 419 — Australia. — M. Drummondii var. Bak. Hdb. 146.

Ernesti A.Br. (Sitzber. Ges. nat. Freunde **1870**. 46); Monatsber. Ak. Berl. **1870**. 746. Bak. Hdb. 143. NPfl. 420 — Caracas.
Zalusianskya O.Ktze. 1891.

erosa Willd. sp. **5**. 540. 1810 = M. minuta.

europaea Desv. Prod. 178. 1827 = M. quadrifolia.

exarata A.Br. Monatsber. Ak. Berl. **1870**. 732. NPfl. 419. (Bak. Hdb. 143) — Queensland.

Fabri Dunal; St. Hil. Ann. sc. nat. II. **6**. 378. 1836 = M. pubescens.

fimbriata Schum. Vid. Selsk. Afh. **4**. 235 (seors. 461). 1827. Bak. Hdb. 142 — Guinea. — Zalusianskya O.Ktze. 1891.

Fischeri Hier. in Engl. Pflw. Ostafr. C. 90. 1895 — Africa orient.

Fournieri C. Chr. Ind. 418. 1906 — Mexico.
M. minuta Fourn. Bull. Soc. Fr. **27**. 329. 1880 (non L. 1771).

gibba A.Br. Monatsber. Ak. Berl. **1870**. 745. Bak. Hdb. 145. NPfl. 420 — Africa austr. — Zalusianskya O.Ktze. 1891.

glomerata Pr. Abh. böhm. Ges. V. **3**. 580. 1845 = M. biloba.

gracilenta A.Br. Monatsber. Ak. Berl. **1863**. 421. 1864. NPfl. 418. (Bak. Hdb. 141) — India.

gymnocarpa Leprieur; A.Br. Flora **1839**. 300 (nomen); Monatsber. Ak. Berl. **1863**. 422. 1864 = M. pygmaea.

hirsuta R.Br. Prod. Fl. N. Holl. 167. 1810. Bak. Hdb. 142. NPfl. 149 — Australia.
Zalusianskya O.Ktze. 1891.

hirsutissima A.Br. Monatsber. Ak. Berl. **1870**. 736. NPfl. 419 — Australia.
M. Drummondii Bak. Hdb. 146 pt.

Howittiana A.Br. Monatsber. Ak. Berl. **1870**. 734. NPfl. 419 — Australia.
M. Drummondii Bak. Hdb. 146 pt.

macra A.Br. App. Ind. Sem. Berol. 1867. 3; Monatsber. Ak. Berl. **1870**. 735. NPfl. 419 — Australia. — M. Drummondii Bak. Hdb. 146 pt.

macrocarpa Pr. Abh. böhm. Ges. V. **3**. 580. 1845. Bak. Hdb. 144. NPfl. 419 — Afr. austr. — Zalusianskya O.Ktze. 1891; Marsilia Dregeana A.Br. 1864.

macropoda Engelm.; A.Br. Amer. Journ. Sci. II. **3**. 56. 1847 (nota); apud Kze. l. c. II. **6**. 88. 1848 — Texas. New Mexico. — M. macropus A.Br. 1864. Bak. Hdb. 140. NPfl. 418 (non Hk. 1854); Zalusianskya O.Ktze. 1891.

macropus Hk. Ic. pl. t. 909. 1854 = M. Drummondii.

macropus A.Br. Monatsber. Ak. Berl. **1863**. 418. 1864 = M. macropoda.

mexicana A.Br. Monatsber. Ak. Berl. **1870**. 747. Bak. Hdb. 143. NPfl. 420 — Mexico. — Zalusianskya O.Ktze. 1891.

microcarpa A.Br. Flora **1839**. 300 = M. crenulata.

minuta L. Mant. 308. 1771 (excl. β). Bak. Hdb. 140 (pt.) — India. Java. Ins. Philipp. — Lemma Desrouss. 1789; Zalusianskya O.Ktze 1891. Marsilea erosa Willd. 1810. NPfl. 418 — ? M. crenata Pr. 1825.

minuta β coromandeliana L. Mant. 308. 1771 = M. coromandelica.

minuta Fourn. Bull. Soc. Fr. **27**. 329. 1880 = M. Fournieri.

mollis Robins. et Fernald. Pr. Amer. Ac. **30**. 123. 1895 — Mexico.

mucronata A.Br. Am. Journ. Sci. II. **3**. 55 f. 2. 1847 = M. vestita.

Muelleri A.Br. Linn. **25**. 721. 1852. NPfl. 419 — Australia.
M. Drummondii var. Bak. Hdb. 146 — M. salvatrix Hanst. 1863. NPfl. 419.

muscoides Lepr.; A.Br. Monatsber. Ak. Berl. **1863**. 422. 1864. Bak. Hdb. 147. NPfl. 420 — Africa occ. — Zalusianskya O.Ktze. 1891; Marsilea muscosa Lepr. 1839 (nomen).

muscosa Leprieur; A.Br. Flora **1839**. 300 (nomen) = M. muscoides.

mutica Mett. Ann. sc. nat. IV. **15**. 88. 1861. Bak. Hdb. 146. NPfl. 420 — N. Caledonia. — Zalusianskya O.Ktze. 1891.

nardu A.Br. Monatsber. Ak. Berl. **1870**. 736. NPfl. 419 — Australia. M. Drummondii Bak. Hdb. 146 pt.

natans L. sp. **2**. 1099. 1753 = Salvinia.

natans Vell. Fl. flum. **11** t. 111. 1827; Arch. Mus. Nac. Rio Jan. **5**. 460. 1881 = Salvinia auriculata.

nubica A.Br. Monatsber. Ak. Berl. **1863**. 432. 1864. Bak. Hdb. 142. NPfl. 420 — Kordofan. — Zalusianskya O Ktze. 1891.

oligospora Goodd. Bot. Gaz. **33**. 66. 1902 — Wyoming.

oxaloides A.Br. Monatsber. Ak. Berl. **1870**. 736. NPfl. 419 — Australia. M. Drummondii Bak. Hdb. 146 pt.

patens Gmel. Syst. Nat. 2^{2}. 1322. 1791 = Salvinia natans.

picta Fée, 9 mém. 47. 1857 = M. vestita var.

polycarpa Hk. et Grev. Ic. Fil. t. 160. 1829. Bak. Hdb. 139. NPfl. 418 — Amer. trop. Ins. Societatis. — Zalusianskya O.Ktze. 1891; Marsilea brasiliensis Mart. 1834.

pubescens Tenore, Fl. Napoli Prod. Suppl. I. 70 —; Cat. pl. hort. Neapol. ed. II. App. I. 67. 1819. Bak. Hdb. 142. NPfl. 418 — Mediterranea. Zalusianskya O.Ktze. 1891. Marsilea Fabri Dunal 1836.

pygmaea A.Brongn. Dict. class. **10**. 199. 1826 (pygmea) — Senegambia. M. gymnocarpa Leprieur 1839; A.Br. Monatsber. Ak. Berl. **1863**. 432. 1864. Bak. Hdb. 145. NPfl. 420; Zalusianskya O.Ktze. 1891.

quadrata A.Br. Monatsber. Ak. Berl. **1870**. 745. Bak. Hdb. 146. NPfl. 419 — Borneo. — Zalusianskya O.Ktze. 1891.

quadrifolia L. sp. **2**. 1099. 1753. NPfl. 418 — Europa centr. et austr.-India bor. Japonia. Connecticut. — Lemma Desrouss. 1789; Zalusianskya O.Ktze. 1891; Z. marsilioides Neck. 1775; Pteris quadrifoliata L. 1759; Marsilea L. 1763. Bak. Hdb. 139; M. europaea Desv. 1827.

quadrifolia Burm. Fl. Ind. 237. 1768 = M. coromandelica.

quadrifolia R.Br. Prod. Fl. N. Holl. 167. 1810 = M. Brownii.

quadrifolia γ Burchellii Kze. Linn. **10**. 556. 1836 = M. Burchellii.

quadrifoliata L. sp. ed. II. **2**. 1563. 1763 = M. quadrifolia.

rotundata A.Br. in Kuhn, Fil. Afr. 200. 1868. Bak. Hdb. 144. NPfl. 419 — Angola. Zalusianskya O.Ktze. 1891.

salvatrix Hanstein, Monatsber. Ak. Berl. **1862**. 105 t. 21. 1863 = M. Muelleri.

salvinioides Neck. Acta Theod. pal. phys. **3**. 297 t. 21. 1775 = Salvinia natans.

senegalensis A.Br. Flora **1839**. 300 (nomen) = M. crenulata var.

sericea A.Br. Monatsber. Ak. Berl. **1870**. 734. NPfl. 419 — Australia. M. Drummondii Bak. Hdb. 146 pt.

stratiotes A.Br. Monatsber. Ak. Berl. **1870**. 723. (Bak. Hdb. 139) — Amazonas.

striata Mett. Ann. sc. nat. V. **3**. 310. 1865 = M. deflexa.

strigosa Willd. sp. **5**. 539. 1810. Bak. Hdb. 142. NPfl. 418 — Rossia austr. Sibiria occ. — Zalusianskya O.Ktze. 1891.

subangulata A.Br. (Sitzber. Ges. Naturf. Freunde Berl. **1870**. 46); Monatsber. Ak. Berl. **1870**. 723. NPfl. 418. (Bak. Hdb. 139) — Caracas. ?Jamaica.

subterranea Leprieur; A.Br. Flora **1839**. 301 (nomen); Monatsber. Ak. Berl. **1863**. 433. 1864. Bak. Hdb. 145. NPfl. 418 — Senegambia.
Zalusianskya O.Ktze. 1891.
tenuifolia Engelm. apud Kze. Amer. Journ. Sci. II. **6**. 89. 1848. Bak. Hdb. 141. NPfl. 419 — Texas. — Zalusianskya O.Ktze. 1891.
tetraphylla Thbg. Fl. Jap. 340. 1784 = M. quadrifolia.
trichopoda Leprieur; A.Br. Flora **1839**. 300 (nomen); Monatsber. Ak. Berl. **1863**. 422. 1864. Bak. Hdb. 147 — Senegambia.
Zalusianskya O.Ktze. 1891; Marsilia trichopus Sad. NPfl. 420.
trichopus Sadeb. Nat. Pfl. **1**^{4}. 420. 1900 = M. trichopoda.
Trifolia Blanco, Fl. Filip. 835. 1837 — Ins. Philipp.
uncinata A.Br. Flora **1839**. 300 (nomen); Am. Journ. Sci. **3**. 55 f. 1. 1847. NPfl 419 — Louisiana. Texas. — Zalusianskya O.Ktze. 1891; M. vestita var. Bak. Hdb. 143.
vestita Hk. et Grev. Ic. Fil. t. 159. 1829. Bak. Hdb. 143 (pt.) NPfl. 419 — U. S. A. pacif.-Mexico. — Zalusianskya O.Ktze. 1891 — Marsilea mucronata A.Br. 1847. NPfl. 419 — M. picta Fée 1857.
villosa Klf. Enum. 272. 1824. Bak. Hdb. 143. NPfl. 419 — Ins. Hawaii.
Zalusianskya O.Ktze. 1891.
villosa Burchell msc. (Bak. JoB. **1886**. 280) — Africa austr.
vulgaris Bory, Bél. Voy. Crypt. **2**. 3. 1833 = { M. crenulata. M. quadrifolia.
MASCHALOSORUS van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 320. 1861
Mougeotii v.d.B. l. c. = T. osmundoides var. [**Trichomanes**.
MATONIA R.Brown in Wallich, Plant. Asiat. Rar. **1**. t. 16. 1829 — (Matoniaceae).
pectinata R.Br. in Wall. Pl. As. Rar. **1**. t. 16. 1829. HB. 45. Chr. 335. NPfl. 347 — Malacca. Borneo. — Prionopteris Farquhariana Wall. 1828 (nomen).
sarmentosa Bak. Journ. Linn. Soc. **24**. 256. 1887; Ann. Bot. **5**. 191. t. 14. 1891. Chr. 336. NPfl. 347 — Borneo.
MATTEUCCIA Todaro, Syn. Pl. Acot. vasc. Sicilia 30. 1866 — (Polypodiaceae [Gen. **7**).
Cavaleriana (Christ) C. Chr. Ind. 420. 1906 — China.
Struthiopteris Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 118 c. figg.
orientalis (Hk.) Trev. Atti Ist. Veneto III. **14**. 586. 1869 — Himalaya orient. China. Japonia. — Struthiopteris Hk. 2 Cent. t. 4. 1860. Chr. 285. NPfl. 166; Onoclea Hk. 1862. HB. 46; Pterinodes O.Ktze. 1891.
struthiopteris (L.) Todaro. Syn. Pl. Acot. vasc. Sicilia 30. 1866 — Europa bor. et centr. Sicilia. Asia temp. Amer. bor. atlant.
Osmunda L. sp. **2**. 1066. 1753; Onoclea Hoffm. 1795. HB. 46; Pterinodes O.Ktze. 1891; Struthiopteris germanica Willd. 1809. Chr. 285. NPfl. 164 — S. pensylvanica Willd. 1810 — Onoclea nodulosa Schkuhr 1809.
MECODIUM Presl, Epim. Bot. 258. 1849 = **Hymenophyllum.**
sanguinolentum Pr. Epim. 258. 1849.
MECOSORUS Klotzsch, Linnaea **20**. 404. 1847 = **Polypodium.**
elongatus Kl. Linn. **20**. 406. 1847 = P. lanceolatum*.
marginellus Kl. Linn. **20**. 405. 1847.
nudus Kl. Linn. **20**. 405. 1847 = P. gramineum.
persicariaefolius Kl. Linn. **20**. 406. 1847.
Schomburgkii Kl. Linn. **20**. 406. 1847 = P. megalophyllum.
trilobus Kl. Linn. **20**. 406. 1847 = P. synammia.
Megalastrum J. Smith. Hist. Fil. 216. 1875 (*Lastrea* §) = Dryopteris.

MELANOPTERIS J. Smith msc. (Hk. Bak. Syn. 258. 1867) = **Aspidium** (plantagineum).

MENISCIUM Schreber, Linn. Genera plant. ed. VIII. **2**. 757. 1791 = **Dryopteris.**

acrostichoides Desv. Prod. 223. 1827 = D. reticulata.

affine Pr.; Ettingsh. Denkschr. Ak. Wien **23**. 94 t. 13 f. 3. 1864 — Brasilia.

Andreanum Sod. Rec. 71. 1883.

angustifolium Willd. sp. **5**. 133. 1810.

arborescens H.B.Willd. sp. **5**. 133. 1810 = ? D. reticulata.

Beccarianum Cesati, Rend. Ac. Napoli **16**. 27, 30. 1877 = D. Cesatiana.

chrysodioides Fée, Gen. 225. 1850-52 = D. reticulata.

cristatum Desrouss. in Lam. Enc. **4**. 94. 1797 — Martinique.

Asplenium Mett. Salom. 1883. (veris. Diplazium sp.).

Cumingii Fée, Gen. 222. 1850-52 = D. urophylla.

cuspidatum Bl. Enum. 114. 1828 = D. urophylla.

deltigerum Wall. List n. 59. 1828 = Leptochilus scalpturatus.

dentatum Pr. Del. Prag. **1**. 162. 1822 = D. serrata.

dentatum Kl. Linn. **14**. 286. 1840 = D. reticulata *macrophylla.

elongatum Fée, Cr. vasc. Br. **1**. 83 t. 25 f. 1. 1869 = D. reticulata.

erosum Wall. List n. 62. 1828 = D. triphylla.

falcatum Liebm. Vid. Selsk. Skr. V. **1**. 183 (seors. 31). 1849 = D. reticulata var.

Fendleri Eat. Bot. Gaz. **3**. 89. 1878 = Leptochilus F.

fragile Kze. Bot. Zeit. **1848**. 259 = D. lineata.

giganteum Mett. Fil. Lechl. **1**. 19. 1856.

grande hort.; Bak. Ann. Bot. **5**. 487. 1891 — Hort.

Guyanense Fée, Gen. 224. 1850-52 = D. reticulata *macrophylla.

Hosei Bak. Journ. Linn. Soc. **22**. 230. 1886.

Jungersenii Fée, Gen. 223. 1850-52 = D. reticulata *falcata.

Kapplerianum Fée, Gen. 224. 1850-52 = D. reticulata *sorbifolia.

Kennedyi F. Muell. Fragm. **4**. 165. 1864 = D. urophylla.

lineatum Kze. Bot. Zeit. **1848**. 259.

longifolium Desv. Prod. 223. 1827; Fée, Cr. vasc. Br. **1**. 84 t. 25 f. 2. 1869 [= D. reticulata.

longifrons Wall. List n. 60. 1828 = D. urophylla.

macrophyllum Kze. Flora **1839**[1]. Beibl. 44 = D. reticulata var.

molle Bak. Syn. 392. 1868 = D. reticulata*.

oligophyllum Ht. Lind.; Bak. Ann. Bot. **5**. 487. 1891 — Hort.

opacum Bak. JoB. **1877**. 166 = D. Christii.

palustre Raddi, Opusc. Bol. **3**. 284. 1819; Pl. bras. **1**. 9 t. 20. 1825 = D. serrata.

Parishii Bedd. Ferns br. Ind. t. 184. 1866 = D. triphylla var.

pauciflorum Hk. sp. **5**. 164. 1864.

proliferum Sw. Syn. 19, 207. 1806.

pubescens Lind. Cat. 1858 = D. Ghiesbreghtii.

puncta lunulatum Rich. Act. Soc. Hist. nat. **1**. 114. 1792 — Cayenne.

reticulatum Sw. Schrad. Journ. **1801**[2]. 274. 1803.

rostratum Fée, Gen. 224 t. 18 B. f. 3. 1850-52 = D. serrata.

salicifolium Wall. List n. 63. 1828.

Salzmanni Fée, Gen. 223. 1850-52 = D. reticulata.

Screberi Rich. Act. Soc. Hist. nat. **1**. 114. 1792 (nomen) = D. serrata.

serratum Cav. Descr. 548. 1803.

sessilifolium Pohl; Fée, Cr. vasc. Br. **1**. 84. 1869 = D. reticulata.

simplex Hk. Lond. JoB. **1**. 294 t. 11. 1842.

sorbifolium Desrouss. in Lam. Enc. **4**. 93. 1797 = D. reticulata*.
stenophyllum Bak. JoB. **1891**. 108 = D. brevipinna.
Thwaitesii Hk. Fil. exot. t. 83. 1859; Syn. 391. 1868.
Thwaitesii Thwait. En. pl. Zeyl. 382. 1864 = D. prolifera.
triphyllum Sw. Schrad. Journ. **1800**². 16. 1801.
villosum J. Sm. (HB. 392) = ? D. triphylla.

MERINGIUM Presl, Hymenoph. 24. 1843.
?Blumeanum Pr. Hym. 24. 1843 = Hymenophyllum.
Meyenianum Pr. Hym. 24 t. 8 f. B. 1843 = Trichomanes.

MERTENSIA Willdenow, Vet. Akad. Nya Handl. **1804**. 163 = **Gleichenia.**
acutifolia Gay, Fl. chil. **6**. 540. 1853 = G. quadripartita.
alpina Poir. Enc. Suppl. **3**. 670. 1813.
angusta Kl.; Sturm, Fl. bras. **1**². 225. 1859 = G. revoluta.
arachnoides Hassk. JoB. **7**. 322. 1855 = G. glauca var.
Bancroftii Kze. Linn. **18**. 307. 1844.
Beyrichiana Sturm, Fl. bras. **1**². 235. 1859 = G. nervosa.
bifida Willd. Vet. Ak. Nya Handl. **1804**. 168 t. 5 f. B.
bifurcata Kze. Bot. Zeit. **1846**. 418 = G. laevigata.
brasiliana Desv. Berl. Mag. **5**. 329. 1811 = G. pectinata.
canescens Klf. Enum. 38. 1824 = G. pectinata.
cinnamomea Beyr.; Sturm, Fl. bras. 1². 234. 1859 = G. nervosa.
circinnata Poir. Enc. Suppl. **3**. 669. 1813.
coeruleo-glauca Poir. Enc. Suppl. **3**. 669. 1813 = G. polypodioides var.
commutata Trev. Atti Ist. Veneto II. **2**. 164. 1851 (nomen) — Malacca.
crassifolia Pr. Abh. böhm. Ges. V. **5**. 339. 1848; Epim. 23 t. 13. 1849 = [G. linearis.
cryptocarpa Gay, Fl. chil. **6**. 539. 1853.
Cumingiana Pr. Tent. 51. 1836 = G. linearis.
Cunninghamii J. Sm. Lond. JoB. **2**. 381. 1843.
decurrens Raddi, Pl. Bras. **1**. 73 t. 7. 1825 = G. bifida vel G. trifurcans.
dicarpa Poir. Enc. Suppl. **3**. 670. 1813.
dichotoma Willd. Vet. Ak. Nya Handl. **1804**. 167 = G. linearis.
discolor Schrad. Gött. gel. Anz. **1824**. 863 = G. linearis.
elata Desv. Prod. 201. 1827 = G. pectinata.
emarginata Raddi, Pl. Bras. **1**. 76 t. 6. 1825 = G. pectinata.
emarginata Brack. Expl. Exp. **16**. 297 t. 42. 1854 = G. linearis var.
excelsa J. Sm. Lond. JoB. **2**. 381. 1843 = G. glauca.
farinosa Klf. Wes. d. Farnkr. 37. 1827; Kze. Anal. 6 t. 3. 1837 = G. bifida.
ferruginea Desv. Berl. Mag. **5**. 307. 1811 = ? G. bifida.
flabellata Poir. Enc. Suppl. **3**. 670. 1813.
flagellaris Bory; Willd. sp. **5**. 74. 1810.
flagellaris Brack. Expl. Exp. **16**. 294. 1854 = G. Brackenridgei.
flexuosa Schrad. Gött. gel. Anz. **1824**. 863.
fulva Desv. Prod. 201. 1827 = G. furcata.
furcata Willd. Vet. Ak. Nya Handl. **1804**. 166.
gigantea Wall.; Pr. Tent. 51. 1836 = G. glauca var.
glabra Brack. Expl. Exp. **16**. 292. 1854 = G. glauca var. pinnata.
glauca Sw. Vet. Ak. Nya Handl. **1804**. 177.
glaucescens H.B.Willd. sp. **5**. 72. 1810 = G. pectinata.
gleichenioides Liebm. Vid. Selsk. Skr. V. **1**. 296 (seors. 144). 1849 = G. Lieb-
gracilenta Trev. Atti Ist. Veneto II. **2**. 164. 1851 (nomen) — Luzon. [manni.

gracilis Mart. Ic. Cr. Bras. 107 t. 59 f. 2. 1834.
gracilis Mart. Ic. Cr. Bras. 106. 1834 = G. pennigera.
grandis Fée, 9 mém. 31. 1857; 11 mém. 120. 1866 — G. Bancroftii.
hawaiensis Brack. Expl. Exp. 16. 295. 1854 = G. owhyhensis.
Hermanni Poir. Enc. Suppl. 3. 670. 1813 = G. linearis.
Hookeri J. Sm. Lond. JoB. 2. 381. 1843 = G. pectinata.
immersa Klf. Enum. 38. 1824 = G. bifida.
Klotzschii Brack. Expl. Exp. 16. 297. 1854 = G. linearis var.
laevigata Willd. sp. 5. 75. 1810.
lanuginosa Moric.; Fée, Cr. vasc. Br. 1. 202. 1869 = G. gracilis.
latissima Fée, Cr. vasc. Br. 1. 203 t. 73 f. 1. 1869 = G. nervosa.
Lessonii A. Rich. Sert. Astrol. XXVII. 1834 = G. linearis var.
linearis Fritsch, Bull. Boiss. II. 1. 1092. 1901.
littoralis Phil. Anal. Univ. chil. 1873. 104.
longipes Fée, Cr. vasc. Br. 2. 87 t. 105 f. 2. 1872-73.
longipinnata Kl. Linn. 18. 537. 1844.
longissima Kze. Bot. Zeit. 1846. 418 = G. glauca.
magellanica Desv. Prod. 201. 1827 = G. quadripartita.
Mathewsii Liebm. Vid. Selsk. Skr. V. 1. 296 (seors. 145). 1849.
microphylla Klf. Enum. 41. 1824 = G. circinnata.
muricata Klf. in Sieb. Syn. n 18 = G. flagellaris.
neglecta Sturm, Fl. bras. 1^{2}. 235. 1859 = G. nervosa.
nervosa Klf. Enum. 37. 1824.
nigropaleacea Sturm, Fl. bras. 1^{2}. 222. 1859 = ? G. furcata.
nitida Pr. Tent. 51. 1836.
nuda Moritz; Reichardt, Denkschr. Ak. Wien 17^{2}. 28 t. 1 f. 1-4. 1859.
obtusa Desv. Berl. Mag. 5. 307. 1811 = G. laevigata.
oligocarpa Phil. Anal. Univ. chil. 94. 361. 1896.
palmata Schaffn.; Fée, 9 mém. 32. 1857.
pectinata Willd. Vet. Ak. Nya Handl. 1804. 168 t. 4.
pedalis Klf. Enum. 39. 1824.
pennigera Mart. Ic. Cr. Bras. 130 t. 59 f. 1. 1834.
pinnata Kze. Anal. 6. 1837 = G. glauca var.
plumaeformis Pr. Abh. böhm. Ges. V. 5. 338. 1848: Epim. 24 t. 15. 1849 = [G. laevigata.
pruinosa Mart. Ic. Cr. Bras. 109. 1834.
pteridifolia Pr. Abh. böhm. Ges. V. 5. 339. 1848: Epim. 23 t. 14. 1849 = [G. linearis var.
pubescens H.B.Willd. sp. 5. 73. 1810 = G. furcata.
pumila Mart. Ic. Cr. Bras. 111 t. 60 f. 2. 1834 = G. linearis.
quadripartita Poir. Enc. Suppl. 3. 669. 1813.
remota Klf. Enum. 39. 1824.
revoluta Desv. Prod. 200. 1827.
rigida Kze. Linn. 9. 16. 1834 = G. flexuosa.
rigida J. Sm. Lond. JoB. 2. 381. 1843 = G linearis var. Lessonii.
rubescens Mart.; Fée, Cr. vasc. Br. 1. 199. 1869 — Brasilia.
rufinervis Mart. Ic. Cr. Bras. 111. 1834 = G. linearis var.
rupestris Poir. Enc. Suppl. 3. 670. 1813.
Sartorii Fée, 9 mém. 32. 1857.
scalpturata Fée. Cr. vasc. Br. 1. 199 t. 72 f. 1. 1889 = G. linearis.
Schaffneri Fée, 9 mém. 40. 1857 — Mexico.
Schomburgkiana Sturm. Fl. bras. 1^{2}. 233. 1859 = G. linearis var. rufinervis.

seminuda Kl. Linn. **18**. 538. 1844.
Sieberi Pr. Tent. 51. 1836 = G. linearis.
simplex Desv. Dict. sc. nat. t. 91; Prod. 200. 1827.
spectabilis Trev. Atti Ist. Veneto II. **2**. 164. 1851 (nomen) — Luzon.
speluncae Poir. Enc. Suppl. **3**. 670. 1813 = G. circinnata.
spissa Fée, Cr. vasc. Br. **1**. 200 t. 73 f. 2. 1869 = G. linearis.
squamosa Fée, Cr. vasc. Br. **1**. 202 t. 72 f. 2. 1869 = G. furcata.
squamulosa Desv. Journ. de Bot. appl. **1**. 268. 1813.
subflabellata Brack. Expl. Exp. **16**. 294. 1854 = G. pruinosa.
subtrisperma Fée, 11 mém. 122 t. 32 f. 2. 1866.
tenera Poir. Enc. Suppl. **3**. 670. 1813.
tenuis Pr. Tent. 51 t. 1 f. 7. 1836.
tomentosa Cav.; Sw. Vet. Ak. Nya Handl. **1804**. 177 t. 5 f. 4; Syn. 164, 392. [1806 = G. furcata.
trifurcans Fée, Cr. vasc. Br. **1**. 201 t. 74 f. 1. 1869.
truncata Willd. Vet. Ak. Nya Handl. **1804**. 169 t. 5 f. A. = ? G. laevigata.
umbraculifera Kze. Linn. **18**. 114. 1844.
velata Kze. Linn. **9**. 15. 1834.
vestita Kze. Bot. Zeit. **1848**. 100.

MESOCHLAENA R.Brown in Horsf. Pl. jav. rar. **1**. 5. 1838 — (Polypodiaceae [Gen. **12**).
asplenioides J. Sm. JoB. **3**. 18. 1840 = M. polycarpa.
javanica R.Br. in Horsf. Pl. jav. rar. **1**. 5. 1838 = M. polycarpa.
moluccana R.Br.; J. Sm. JoB. **3**. 18. 1840 = M. polycarpa.
polycarpa (Bl.) Bedd. Ferns br. Ind. Suppl. 13. 1876. NPfl. 181 — Malesia-Polynesia. — Aspidium Bl. Enum. 156. 1828. Chr. 249; Didymochlaena Bak. 1867. HB. 248; Nephrodium Keys. 1873. Mesochlaena javanica R.Br. 1838; Stegnogramme Fée 1850-52; Aspidium Mett. 1856; Nephrodium Hk. 1859. Stegnogramme Mesochlaena Fée 1850-52. Sphaerostephanos asplenioides J. Sm. 1839; Mesochlaena J. Sm. 1840 — Lastrea microchlamys de Vriese 1848. Nephrodium Bak. 1876.

Mesopleuria Moore, Ind. Fil. XXXVI. 1857 *(Adiantum §)* = Adiantum.

MESOSORUS Hasskarl, Observ. Bot. Fil. Bogor. **1**. 2. 1856 = **Gleichenia.**
arachnoides Hassk. Obs. **1**. 6. 1856 = G. glauca var.
bifurcatus Hassk. Obs. **1**. 8. 1856 = G. laevigata.
dichotomus Hassk. Obs. **1**. 9. 1856 = G. linearis.
excelsus Hassk. Obs. **1**. 3. 1856 = G. glauca var.
giganteus Hassk. Obs. **1**. 2. 1856 = G. glauca var.
glaucus Hassk. Obs. **1**. 15. 1856.
longissimus Hassk. Obs. **1**. 3. 1856 = G. glauca.
vestitus Hassk. Obs. **1**. 13. 1856.

MESOTHEMA Presl, Epim. Bot. 111. 1849 = **Blechnum.**
appendiculatum Pr. Epim. 112, 261. 1849.
auriculatum Pr. Epim. 112. 1849.
australe Pr. Epim. 112. 1849.
campylotis Pr. Epim. 112. 1849 = B. occidentale var.
hastatum Pr. Epim. 111. 1849 = B. auriculatum.
javanicum Pr. Epim. 262. 1849 = B. Blumei.
plantagineum Pr. Epim. 111. 1849 = B. lanceola.
punctulatum Pr. Epim. 113. 1849.
remotum Pr. Epim. 111. 1849 = B. auriculatum var.
rigidum Pr. Epim. 113. 1849 = B. australe.
trilobum Pr. Epim. 112. 1849 = B. auriculatum var.

METAXYA Presl, Tent. Pterid. 59. 1836 = **Alsophila.**
Parkeri J. Sm. Lond. JoB. **1**. 668. 1842 = A. blechnoides var.
rostrata Pr. Tent. 60 t. 1 f. 5. 1836 = A. blechnoides.
Micrasplenium Keyserling, Pol. Cyath. Hb. Bung. 15. 1873 (*Asplenium* §) = [Asplenium.
MICROBROCHIS Presl, Epim. Bot. 51. 1849 = **Aspidium.**
apiifolia Pr. Epim. 52. 1849 = A. cicutarium*.
Microdictyon Fée, 11 mém. 66. 1866 (*Goniopteris* §) = Dryopteris.
MICROGONIUM Presl, Hymenoph. 19. 1843 = **Trichomanes.**
Berteroanum Pr. Hym. 20, 46 t. 6 f. B. 1843 = T. Hookeri.
bimarginatum v.d.B. Hym. Jav. 7. 1861.
cuspidatum Pr. Hym. 20. 1843.
erosum Pr. Abh. böhm. Ges. V. **5**. 335. 1848.
Henzariense v.d.B. Hym. Jav. 7. 1861 = T. Henzaianum.
Hookeri Pr. Abh. böhm. Ges. V. **5**. 335. 1848.
Motleyi v.d.B. Hym. Jav. 5 t. 1. 1861.
Petersii v.d.B. Hym. Jav. 7. 1861.
Schaffneri Fée, 9 mém. 38. 1857 = T. hymenoides.
sublimbatum v.d.B. Hym. Jav. 6 t. 2. 1861.
MICROGRAMMA Presl, Tent. Pterid. 213. 1836 = **Polypodium.**
persicariaefolia Pr. Tent. 214 t. 9 f. 7. 1836.
MICROLEPIA Presl, Tent. Pterid. 124. 1836 — (Polypodiaceae Gen. **33**).
aculeata Mett. Fil. Lips. 103. 1856 = Odontosoria.
adiantoides J. Sm. Lond. JoB. **1**. 427. 1842 = Saccoloma domingense.
affinis Pr. Epim. 97. 1849 = Davallia hymenophylloides.
alata J. Sm. JoB. **3**. 416. 1841 = Saccoloma moluccanum.
alata J. Sm. Lond. JoB. **1**. 427. 1842 = Saccoloma inaequale.
arcuata Fourn. Mex. pl. **1**. 131. 1872 = Saccoloma inaequale var.
aspidioides Pr. Epim. 96. 1849 = M. speluncae.
biflora Mett. Fil. Lips. 103. 1856 = Odontosoria.
biserrata Pr. Epim. 97. 1849 = Tapeinidium pinnatum.
Blumei Moore, Ind. 297. 1861 = Saccoloma moluccanum.
Boryana Fourn. Mex. pl. **1**. 131. 1872 = Saccoloma inaequale var.
brasiliensis Pr. Tent. 125. 1836; Epim. 95. 1849 = Saccoloma inaequale.
calobodon Mett.; Kuhn, Linn. **36**. 147. 1869 = M. madagascariensis.
calcescens Pr. Epim. 95. 1849 = M. marginata var.
campylura Moore, Ind. 296. 1861 = Saccoloma moluccanum var.
?*caudata* Fée, Gen. 327. 1850-52 — Mexico.
caudigera Moore, Ind. 303. 1861 — Bhotan.
Davallia urophylla Hk. Fil. Exot. t. 19 nota 1859 (nomen?); Bak. Syn. 99. 1867 (non Wall. 1828 nomen); Microlepia Bedd. 1883 (non 1866 cum descr.).
chaerophylla Ettingsh. Farnkr. 208 t. 142 f. 2. 1865 = Davallia pulchra.
chinensis Mett. Fil. Lips. 103. 1856 = Odontosoria.
ciliata Copeland Polyp. Philipp. 55. 1905 = M. hirsuta.
coniifolia Fée, 11 mém. 93. 1866 = Balantium.
cristata J. Sm. JoB. **3**. 416. 1841 = M. strigosa.
cystopteroides Pr.; Ettingsh. Farnkr. 209 t. 143 f. 2. 1865 — Guatemala.
decurrens Fée, Gen. 326. 1850-52 = Davallia.
Denhami Moore, Ind. 292. 1861 = Tapeinidium.
distans J. Sm. Lond. JoB. **1**. 427. 1842 = ? Saccoloma inaequale.
domingensis Fée, 11 mém. 93. 1866 = Dicksonia.

effusa Fée, 8 mém. 109. 1857 — Pulo Penang.
elegans Mett. Fil. Lips. 103. 1856 = Saccoloma.
exserta Mett.; Kuhn, Linn. **36**. 148. 1869 — Fiji.
Davallia Speluncae var. HB. 470.
firma Mett.; Kuhn, Linn. **36**. 146. 1869 — India bor.
firmula (Bak.) C. Chr. Ind. 210. 1905; 426. 1906 — Sumatra.
Davallia Bak. Ann. Bot. **8**. 123. 1894.
flaccida J. Sm. Lond. JoB. **1**. 427. 1842 = M. speluncae.
flaccida Fée, Gen. 327. 1850-52 = Dennstaedtia.
flexuosa Ettingsh. Farnkr. 208 t. 138 f. 4. t. 140 f. 5. 1865 = Davallia.
fluminensis Fée, Cr. vasc. Br. **1**. 151 t. 51 f. 1. 1869 = Dennstaedtia.
Forsteri Armstr. Tr. N. Zeal. Inst. **13**. 364. 1881 = Odontosoria angustifolia.
Galeottii Fée, Gen. 327. 1850-52 = Saccoloma inaequale var.
gracilis J. Sm. Lond. JoB. **1**. 427. 1842 = Tapeinidium pinnatum var.
Hancei Prantl, Arb. bot. Gart. Breslau **1**. 35. 1892 — Hongkong. Khasia.
Davallia polypodioides Benth. 1861 (non Don 1825); Microlepia Hance 1873.
Henriettae Kuhn. v. Deck. Reis. **3**^{3}. Bot. 62. 1879 = Dennstaedtia.
hirsuta (J. Sm.) Pr. Epim. 97. 1849 — Ins. Philipp. Celebes. Borneo.
Leucostegia J. Sm. 1841 (nomen); Davallia ciliata Hk. sp. **1**. 184 t. 60 A. 1846. HB. 98 (non Pr. 1822); Microlepia Copeland 1905.
hirta (Klf.) Pr. Tent. 125. 1836 — Asia trop. Ins. Hawaii.
Davallia Klf. Enum. 223. 1824. HB. 100. D. villosa Don 1825. Dicksonia Kaulfussiana Gaud. 1827.
Hookeriana (Wall.) Pr. Epim. 95. 1849. Chr. 308. NPfl. 215 — China merid.-India bor. — Davallia Wall. 1828 (nomen); Hk. sp. **1**. 172 t. 47 B. 1846. HB. 97; Saccoloma Fée 1850-52; Scypholepia J. Sm. 1875.
humilis Pr. Tent. 125. 1836 = Cystopteris fragilis.
hypolepidoides Kuhn, v. Deck. Reis **3**^{3}. Bot. 62. 1879 = M. madagascariensis.
inaequalis Pr. Tent. 125. 1836 = Saccoloma.
inaequalis var. minor Brack. Expl. Exp. **16**. 235 t. 33 f. 1. 1854 = Saccoloma [moluccanum.
incisa Fée, Gen. 328. 1850-52 — Guadeloupe.
Jamaicensis Fée, 11 mém. 92. 1866 = M. speluncae.
Jamaicensis Hill. Fl. Haw. 626. 1888 (part.?) = Leptolepia Andersoni.
japonica Pr. Epim. 95. 1849 = M. strigosa.
khasiyana Pr. Epim. 95. 1849 = M. strigosa.
Kurzii (Clarke) Bedd. Handb. 66. 1883 — Birma.
Davallia Clarke, Tr. Linn. Soc. II. Bot. **1**. 446. 1880.
L'Herminieri Moore, Ind. 296. 1861 = Davallia.
Lindeni Fée, Gen. 327. 1850-52 = Saccoloma domingense var.
Lindeni Mett. Fil. Lips. 104 t. 27 f 17. 18. 1856 = Asplenium theciferum var.
lindsayaeformis Fée, Cr. vasc. Br. **1**. 152 t. 51 f. 2 1869 = Dennstaedtia.
lonchitidea Pr. Tent. 125. 1836 = M. platyphylla.
madagascariensis (Kze.) Pr. Epim. 95. 1849 — Madagascar.
Dicksonia? Kze. Anal. 38. 1837; Davallia Kze. 1850. Microlepia calabodon Mett.; Kuhn 1869; Davallia Bak. 1874. HB. 470 — Dicksonia hypolepidoides Bak. 1877; Microlepia Kuhn 1879.
majuscula (Lowe) Moore, Ind. 297. 1861 — Ceylon.
Davallia Lowe, Ferns 8 t. 33. 1859(?). HB. 99. D. proxima? Thwait. Enum. pl. Zeyl. 438. 1864 (non Bl. 1828); Microlepia Bedd. 1863-65; Davallia Thwaitesii Bak. 1867; Microlepia Bedd. 1870.

manilensis Pr. Tent. 125. 1836 (nomen) — Java. Celebes. Ins. Philipp.
Davallia Goldm. Nova Acta **19**. Suppl. I. 465. 1843. D. Sarasinorum Christ [1896.
Mannii Eat. apud Mann, Proc. Amer. Ac. **7**. 212. 1868 = Davallia.
marginalis Bedd. Ferns br. Ind. Correct. 1870 = M. marginata.
marginalis × strigosa Prantl, Arb. bot. Gart. Breslau **1**. 38. 1892 = M. marginata × strigosa.
marginata (Houtt.) C. Chr. Ind. 212. 1905; 427. 1906 — Japonia. China-Ceylon.
Polypodium Houtt. Pfl. Syst. **13**[1]. 199. 1786; P. marginale Thbg. Fl. Jap. 337. 1784 (non L. 1753); Dicksonia Sw. 1801; Davallia Bak. 1868. HB. 98; Microlepia Bedd. 1870. Chr. 307. NPfl. 215. Davallia scabra Don 1825; Microlepia J. Sm. 1842 — Davallia calvescens Hk. 1846; Microlepia Pr 1849. Davallia urophylla Wall. 1829 (nomen); Microlepia Moore 1861; Bedd. 1866 (non 1883).
marginata × strigosa — Japonia.
M. marginalis × strigosa Prantl, Arb. bot. Gart. Breslau **1**. 38. 1392.
mollis Fée, Gen. 328. 1850-52 = Dryopteris crenata.
mollis Moore Ind. 297. 1861 = M. speluncae.
moluccana Pr. Epim. 95. 1849 = Tapeinidium amboynense.
multifida Pr. Epim. 96. 1849 = M. speluncae.
nigricans Pr. Epim. 95. 1849 = Saccoloma inaequale var.
nipponica (Miq.) C. Chr. Ind. 427. 1906 — Japonia.
Davallia Miq. Ann. Lugd. Bat. **3**. 180. 1867.
Novae-Zealandiae J. Sm. Cat. Kew Ferns 1856 = Leptolepia.
?oedematosa Fée, Gen. 328. 1850-52 = Dennstaedtia exaltata.
papillosa Brack. Expl. Exp. **16**. 237 t. 34. 1854 = Saccoloma moluccanum.
phanerophlebia (Bak.) C. Chr. Ind. 213. 1905; 427. 1906 — Tonkin.
Davallia Bak. JoB. **1890**. 262.
philippinensis (Harr.) Copeland, Polyp. Philipp. 56. 1905 — Luzon.
Davallia Harr. Journ. Linn. Soc. **16**. 27. 1877.
pilosella (Hk.) Moore, Ind. 298. 1861 — Japonia.
Davallia Hk. 2 Cent. t. 96. 1861. Trichomanes hirsutum Thbg. Fl. Jap. 339. 1784 (non L. 1753); Davallia Sw. 1801. HB. 98; Humata Desv. 1827; Dennstaedtia Mett. 1867; Trichomanes japonicum Poir. 1808 (non Thbg. 1784).
pilosula (Wall.) Pr. Tent. 125. 1836; Prantl, Arb. bot. Gart. Breslau **1**. 36. 1892 — Asia trop. — Davallia Wall. 1828 (nomen); D. polypodioides γ hispida Hk. sp. **1**. 182. 1846.
Pinkneyi (Col.) C. Chr. Ind. 213. 1905; 427. 1906 — N. Zealand.
Davallia Col. Tr. N. Zeal. Inst. **29**. 415. 1897.
pinnata J. Sm. JoB. **3**. 416. 1841 = Tapeinidium.
platyphylla (Don) J. Sm. Lond. JoB. **1**. 427. 1842. Chr. 307. NPfl. 215 — India bor.-Ceylon. — Davallia Don, Prod. Fl. Nepal. 10. 1825; D. lonchitidea Wall. 1828; Microlepia Pr. 1836.
Plumieri Fée, Gen. 327. 1850-52 = Saccoloma domingense.
Pohliana Ettingsh. Farnkr. 210 t. 142 f. 11, t. 141 f. 2, 4. 1865 = Saccoloma [inaequale.
polyantha Fée, Gen. 327. 1850-52 = Davallia divaricata.
polypodioides Pr. Tent. 125. 1825 = M. speluncae.
polypodioides Bedd. Ferns S. Ind. t. 15. 1863 = M. trapeziformis.
polypodioides Hance, Journ. Linn. Soc. **13**. 143. 1873 = M. Hancei.
proxima (Bl.) Pr. Epim. 95. 1849 — Malesia.
Davallia Bl. Enum. 238. 1828 (excl. β). D. strigosa β (part.). HB. 99.

proxima Bedd. Ferns S. Ind. t. 254. 1863-65 = M. majuscula.
pteropus Bedd. Ferns br. Ind. t. 313. 1869 = Davallia multidentata.
rhomboidea Pr. Tent. 125. 1836 = M. trapeziformis.
rhomboidea J. Sm. JoB. **3**. 416. 1841 = M. trichosticha.
scaberula Mett.; Kuhn, Linn. **36**. 148. 1869 — Ins. Societatis. N. Guinea.
scabra J. Sm. Lond. JoB. **1**. 427. 1842 = M. marginata.
Schimperi Mett. Fil. Lips. 104. 1856 = Asplenium theciferum*.
Schlechtendalii Mett. Fil. Lips. 104. 1856 = Odontosoria.
scoparia Carr. in Seem. Fl. Vit. 340. 1873 = Odontosoria.
speluncae (L.) Moore, Ind. XCIII. 1857. Chr. 308. NPfl. 215 — Trop. et subtrop. Polypodium L. sp. **2**. 1093. 1753; Aspidium Willd. 1810; Davallia Bak. 1867. HB. 100 (excl. varr. p. 470). ? Dicksonia polypodioides Sw. 1801; ? Humata Desv. 1827 (an = M. strigosa ?); Davallia Hk. 1846 (part.); Microlepia Pr. 1836. Davallia flaccida R.Br. 1810; Microlepia J. Sm. 1842 — Davallia jamaicensis Hk. 1846; ? Acrophorus Moore 1854; Microlepia [Fée 1866.
splendens Moore, Ind. XCIII. 1857 = ? Davallia solida.
stenoloba Prantl, Arb. bot. Gart. Breslau **1**. 36. 1892 — Sikkim.
strigosa (Thbg.) Pr. Epim. 95. 1849. Chr. 308. NPfl. 215 — Japonia. China-Asia-Polynesia trop. — Trichomanes Thbg. Fl. Jap. 339. 1784; Dicksonia Thbg. 1794; Davallia (Sw.) Kze. 1848. HB. 98 (excl. β); Dennstaedtia J. Sm. 1875; Polypodium cristatum Houtt. 1786 (non L. 1753); Microlepia J. Sm. 1841. Dicksonia japonica Sw. 1801; Davallia Kze. 1848; Microlepia Pr. 1849. Davallia polypodioides Don 1825 (an Dicksonia Sw. 1801?; vide M. speluncae) — D. Khasiyana Hk. 1846; Microlepia Pr. 1849.
subvolubilis Moore, Ind. 301. 1861 = M. speluncae.
tenuifolia Pr. Epim. 97. 1849 = Davallia hymenophylloides.
tenuifolia Mett. Fil. Lips. 104 t. 27 f. 1-4. 1856 = Odontosoria chinensis.
tenuis Brack. Expl. Exp. **16**. 236. 1854 = Saccoloma moluccanum.
Thwaitesii Bedd. Ferns br. Ind. Corret. II. 1870 = M. majuscula.
trapeziformis (Roxb.) Kuhn, Chaetopt. 347. 1882 — Asia trop. Davallia Roxb. Calc. Journ. **4**. 516. 1844 — D. rhomboidea Wall. 1828 (nomen); Kze. 1850. HB. 99 (D. strigosa var.); Microlepia Pr. 1836 (nomen).
trichosticha J. Sm. JoB. **3**. 416. 1841 (nomen) — Malesia-Samoa. Davallia Hk. sp. **1**. 183. 1846; D. speluncae var. HB. 470. D. divergens [Kze. 1847.
uncinella Mett. Fil. Lips. 103. 1856 = Odontosoria.
urophylla Moore, Ind. 290. 1861 (nomen); Bedd. Ferns br. Ind. t. 103. 1866 [= M. marginata.
urophylla Bedd. Handb. 64. 1883 = M. caudigera.
venusta Mett. Fil. Lips. 103. 1856 = Odontosoria clavata.
villosa Pr. Epim. 95. 1849 = M. marginata.
Wilfordii Moore, Ind. 299. 1861 — Japonia. Korea. China. Davallia Bak. 1867. HB. 98. Chr. 305. D. rhomboidea Hk. 2 Cent. t. 48. 1860 (non Wall. 1828; Kze. 1850).
?zeylanica Pr. Epim. 96. 1849 = Dennstaedtia flaccida.

MICROPODIUM Mettenius, Ann. Mus. Lugd. Bat. **2**. 232. 1866.
cardiophyllum Hance, JoB. **1883**. 268 = Asplenium.
Durvillei Mett. Ann Lugd. Bat. **2**. 233. 1866 = Stenochlaena sorbifolia var. 16.
Fijeense J. Sm. Hist. Fil. 323. 1875 = Asplenium amboinense.
lanceum J. Sm. Hist. Fil. 323. 1875 = Diplazium.
longifolium Mett. Ann. Lugd. Bat. **2**. 233. 1866 = Triphlebia.
pinnatum Mett. Ann. Lugd. Bat. **2**. 233. 1866 = Triphlebia.

scolopendrioides J. Sm. Hist. Fil. 323. 1875 = Asplenium.
sundense Mett. Ann. Lugd. Bat. **2**. 232. 1866 = Asplenium vittaeforme.
vittaeforme J. Sm. Hist. Fil. 323. 1875 = Asplenium.

MICROPTERIS Desvaux, Prodrome in Mém. Soc. Linn. Paris **6**. 217. 1827.
blechnoides Desv. Prod. 217. 1827 = Monogramma seminuda.
orientalis Desv. Prod. 217. 1827 = Polypodium serrulatum.
pectinata Desv. Prod. 217. 1827 = Acrostichum serratum Poir.
serrulata Desv. Prod. 217. 1827 = Polypodium.

MICROPTERIS J. Smith, Hist. Fil. 186. 1875 = **Polypodium.**
fallax J. Sm. Hist. Fil. 186. 1875.
***graminoides* J. Sm. Hist. Fil. 186. 1875 — Jamaica.**
monticola J. Sm. Hist. Fil. 186. 1875.

MICROSORIUM Link, Hort. Berol. **2**. 110. 1833 = **Polypodium.**
irioides Fée, Gen. 268. 1850-52 = P. punctatum.
irregulare Link, Hort. Berol. **2**. 110. 1833 = P. punctatum.
longissimum Fée, Gen. 268 t. 20 B. f. 2. 1850-52 = P. myriocarpum.
?palmatum Fée, Gen. 269. 1850-52 = P. insigne.
sessile Fée, Gen. 268. 1850-52 = P. punctatum.
?trifidum Fée, Gen. 269. 1850-52 = Aspidium irriguum.

MICROSTAPHYLA Presl, Epim. bot. 160. 1849 = **Elaphoglossum.**
Bangii Hier. Engl. Jahrb. 34. 539. 1904 = E. Moorei.
bifurcata Pr. Epim. 161. 1849 = E. furcatum.
dimorphum J. Sm. Hist. Fil. 131. 1875.
furcata Fée, 7 mém. 45 t. 13. 1857.
Moorei Und. Torreya 5. 88. 1905.

MICROSTEGIA Presl, Epim. bot. 90. 1849 = **Diplazium.**
ambigua Pr. l. c. 91 = D. esculentum. dilatata Pr. l. c. 91 = D. maximum.
Kunzeana Pr. l. c. 92 = D. asperum. marginata Pr. l. c. 92 = ? D. polypodioides.
pubescens Pr. l. c. 260 = D. esculentum. serrulata Pr. l. c. 91 = D. manilense.
aspera Pr. l. c. 92 — esculenta Pr. l. c. 91 — microphylla Pr. l. c. 91 — polypodioides Pr. l. c. 91 — prionophylla Pr. l. c. 92 — sylvatica Pr. l. c. 90 — **varia Pr. l. c. 92 = D. sp. homonym.**

MICROSTEGNUS Presl, Abh. böhm. Ges. Wiss. V. **5**. 353. 1848 = **Hemitelia.**
grandifolius Pr. Abh. böhm. Ges. V. 5. 353. 1848.

MICROTERUS Presl, Epim. bot. 124. 1849 = **Polypodium.**
neglectus Pr. Epim. 124. 1849.

Microtrichomanes Mett.; Prantl. Hymenoph. 51. 1875 *(Gonocormus §)* = Trichomanes.

MILDELLA Trevisan, Rendic. Istituto Lombardo sci. et lett. II. **9**. 810. 1876 [= **Pellaea.**
intramarginalis Trevis. l. c.

MOHRIA Swartz, Syn. Fil. 6. 159. 1806 — (Schizaeaceae Gen. **3**).
achilleaefolia Lowe, New Ferns t. 42 B. 1862 = M. caffrorum.
aurita J. Sm. Lond. JoB. 2. 388. 1843 = Aneimia.
burniifolia J. Sm. Lond. JoB. 2. 388. 1843 = Aneimia.
caffrorum (L.) Desv. Prod. 198. 1827. HB. 436. Prantl, Schiz. 84. Chr. 352. NPfl. 367 — Afr. orient. et austr. Madagascar. Ins. Mascarenae.
Polypodium L. Mantissa 307. 1771; Adiantum L.fil. 1781; Lonchitis Bernh. 1801; Osmunda marginalis Sav. 1797; O. thurifera Sw. 1801; O. thurifraga Bory 1804; Mohria Sw. 1806 — M. crenata Desv. 1811 — **Cheilanthes fuscata Bl. 1828 (patria erronea).**
crenata Desv. Berl. Mag. **5**. 307. 1811 = M. caffrorum.

intermedia J. Sm. Lond. JoB. 2. 387. 1843 = Aneimia cicutaria.

lepigera Bak. Ann. Bot. 5. 498. 1891 — Zambesia.

Nothochlaena Bak. JoB. 1884. 53.

millefolium J. Sm. Lond. JoB. 2. 388. 1843 = Aneimia.

thurifraga Sw. Syn. 159, 385 t. 5. 1806 = M. caffrorum.

vestita Bak. Tr. Linn. Soc. II. Bot. 2. 355. 1887; Hk. Ic. pl. t. 1696. 1887 — Kilimanjaro.

MONACHOSORUM Kunze, Bot. Zeit. 1848. 119 — (Polypodiaceae Gen. 37).

davallioides Kze. Bot. Zeit. 1848. 119 = M. subdigitatum.

Henryi Christ, Bull. Boiss. 6. 869. 1898. NPfl. 218 — Yunnan.

subdigitatum (Bl.) Kuhn, Chaetopt. 345. 1882. Chr. 76. NPfl. 218 — Asia trop.

Aspidium Bl. Enum. 171. 1828; Polypodium Bl. Fl. Jav. Fil. 196 t. 93. 1829. HB. 340; Gymnogramme Keys. 1873; Phegopteris Bedd. 1883. Monachosorum davallioides Kze. 1848; Anogramme Fée 1850-52; Polypodium Mett. 1856. P. coniifolium Wall. 1828; Desmopodium J. Sm. msc.

MONOCHLAENA Gaudichaud, Freyc. Voy. Bot. 340. 1827 = **Didymochlaena.**

sinuosa Gaud. l. c. = D. truncatula.

MONOGONIA Presl, Tent. Pterid. 146. 1836 = **Dryopteris.**

palustris Pr. Tent. 146 t. 5 f. 15. 1836 = D. tetragona.

MONOGRAMMA Schkuhr, Krypt. Gewächse 1. 82. 1809; Desvaux, Berl. Mag. 5. 302. 1811 — (Polypodiaceae Gen. 88).

dareicarpa Hk. sp. 5. 121 t. 288 A. 1864. HB. 375. NPfl. 299 — Borneo. [N. Guinea. Ins. Philipp.

?extensa Fée, Gen. 98. 1850-52 = M. paradoxa.

?furcata Desv. Berl. Mag. 5. 303. 1811 = M. graminoides.

graminea (Poir.) Schkuhr, Kr. Gew. 1. 82. 1809; Desv. Berl. Mag. 5. 302. 1811. HB. 375. NPfl. 299 — Ins. Mascar. Ins. Sechellae. ? Afr. austr.

Pteris Poir. Enc. 5. 708. 1804; Caenopteris Schkuhr 1809 (t. 87); Grammitis pumila Sw. 1806; Monogramme linearis Klf. 1820.

graminifolia Hk. sp. 5. 125. 1864 = M. seminuda.

graminoides (Sw.) Bak. Syn. 375. 1868. Chr. 54. NPfl. 298 — Amer. trop.-Chile. St. Helena. — Acrostichum Sw. Prod. 128. 1788; Grammitis Sw. 1801; Asplenium Sw. 1806; Cochlidium Klf. 1820; Pleurogramme Fée 1851-52; Taenitis Mett. 1856. Monogramma furcata Desv. 1811; Pleurogramma J. Sm. 1875.

gyroflexa (Christ) C. Chr. Ind. 430. 1906 — Costa Rica.

Pleurogramme Christ, Prim. Fl. costaric. 3. 10. 1901.

immersa Hk. sp. 5. 125. 1864 = M. linearifolia.

interrupta Bak. Ann. Bot. 5. 482. 1891. NPfl. 299 — N. Guinea.

Pleurogramme Christ 1897. 55.

Junghuhnii Hk. sp. 5. 123 t. 289 B. 1864 = M. paradoxa.

linearifolia Desv. Berl. Mag. 5. 302 t. 7 f. 5. 1811 — Cuba. Guiana. Venezuela.

Grammitis Steud. 1824; Pleurogramme Moore 1857. P. immersa Fée 1851-52. Chr. 55; Monogramme Hk. 1864. HB. 376.

linearis Klf. Jahrb. Pharm. Berlin 1820. 44; Enum. 196. 1824 = M. graminea.

linearis Bl. Enum. 198. 1828 = M. paradoxa.

minor Jenm. Bull. Dep. Jam. 4. 212. 1897 — Jamaica.

myrtillifolia (Fée) Hk. sp. 5. 125. 1864. (HB. 376) — ? Mexico.

Pleurogramme Fée, 3 mém. 58. 1851-52.

paradoxa (Fée) Bedd. Ferns br. Ind. Suppl. 24. 1876. Chr. 54. NPfl. 298 — Ceylon. Malesia. Polynesia. Austr. trop. — Pleurogramme? Fée. 3 mém.

38 t. 4 f. 4. 1851-52; Vaginularia Mett. 1868-69. V. Junghuhnii Mett. 1856; Monogramme Hk. 1864. HB. 375 — Diclidopteris angustissima Brack. 1854; Vaginularia Mett. 1868-69.

robusta (Christ) C. Chr. Ind. 431. 1906 — China. Thibet orient.
Pleurogramme Christ, Bull. Boiss. **6**. 867. 1898.

rostrata Hk. sp. **5**. 122 t. 288 B. 1864. HB. 375. NPfl. 298 — Nicaragua. Cuba. Para.

Rudolfii Rosenst. Festschr. Alb. v. Bamberg 63. 1905 — Brasilia austr.

seminuda (Willd.) Bak. Syn. 375. 1868. NPfl. 299 — Amer. trop. Guinea.
Blechnum Willd. Phyt. 13 t. 8 f. 2. 1794; Grammitis Willd. 1810; Taenitis Kuhn 1868; Pleurogramma J. Sm. 1875. Chr. 55; Micropteris blechnoides Desv. 1827. Taenitis pumila Klf. 1824; Pleurogramme Pr. 1836. Taenitis linearis Klf. 1824; Pleurogramme Pr. 1836. Taenitis graminifolia Hk. 1825; Pleurogramme Pr. 1836; Monogramme Hk. 1864 — Grammitis Iserti Sw. 1829 (Guinea).

subfalcata Hk. sp. **5**. 122 t. 289 A. 1864. HB. 375 — N. Hebridae.

trichoidea J. Sm. JoB. **3**. 394. 1841 (nomen). HB. 375. NPfl. 298 — Ins. Philipp. Borneo. — Vaginularia Fée, 3 mém 54. 1851-52.

Monophlebia Moore, Index LXXXIX. 1857 *(Lastrea §)* = Dryopteris.

MÜLLERIA Schaffner msc. (Fourn. Mex. pl. **1**. 59. 1872) = **Trichomanes.**

MYRIOPTERIS Fée, Genera Filicum 148. 1850-52 = **Cheilanthes.**

Bradburii J. Sm. Hist. Fil. 280. 1875 = C. tomentosa.
cheiloglyphis Fée, 8 mém. 77. 1857.
contracta Fée, Gen. 149. 1850-52 = C. hirta.
elegans J. Sm. Cat. cult. ferns 29. 1857 = C. myriophylla.
Fendleri Fourn. Mex. pl. **1**. 125. 1872.
frigida J. Sm. Cat. cult. ferns 29. 1857 = C. lendigera.
gracilis Fée, Gen. 150 t. 29 f. 6. 1850-52 = C. Feei.
gracillima J. Sm. Hist. Fil. 280. 1875.
hirta J. Sm. Ferns br. and for. 174. 1866.
induta Fée, Gen. 149 t. 12 A f. 3. 1850-52.
intermedia Fée, Gen. 149. 1850-52 = C. hirta.
intermedia Fourn. Bull. Soc. Fr. **27**. 328. 1880 = C Fournieri
lanuginosa Fourn. Mex. pl. **1**. 125. 1872 = C. lendigera.
lanuginosa J. Sm. Hist. Fil. 280. 1875 = C. Feei.
lendigera J. Sm. Cat. cult. ferns 28. 1857.
lentigera Fée, Gen. 149 t. 12 A. f. 4. 1850-52 = C. lendigera.
Lindheimeri J. Sm. Bot. Voy. Herald 340. 1854.
Mac Leanii J. Sm. Hist. Fil. 280. 1875 = C. pilosa.
marsupianthes Fée, Gen. 149 t. 12 A. f. 1. 1850-52 = C. lendigera.
minor Fée, Gen. 149. 1850-52 = C. lendigera.
myriophylla J. Sm. Bot. Voy. Herald 340. 1854.
paleacea Fée, Gen. 149. 1850-52 = C. myriophylla.
rufa Fée, 8 mém. 77. 1857 = C. Schaffneri.
scariosa Fée, Gen. 149. 1850-52.
Szovitzii J. Sm. Hist. Fil. 281. 1875 = C. persica.
tomentosa Fée, Gen. 149 t. 12 A f. 2. 1850-52.
vestita J. Sm. Cat. cult. ferns 29. 1857 = C. lanosa.
villosa Fée, Gen. 149 t. 28 f. 3. 1850-52 = C. lendigera.

MYRIOTHECA Commerson in Jussieu, Gen. pl. 15. 1789 = **Marattia.**
alata Poir. Enc. **4**. 403. 1797.
arborescens Blanco, Fl. Filip. 831 1837 = ? Cyathea integra.
frascinea Poir. Enc. **4**. 404. 1797 = M. fraxinea.
fraxinifolia Bory, Voy. **1**. 266. 1804 = M. fraxinea.
laevis Poir. Enc. **4**. 403. 1797 = M. alata var.
sorbifolia Bory, Voy. **1**. 267. 1804 = M. fraxinea.

Myrmecophila Christ, Farnkr. 112. 1897 *(Polypodium §)* = Polypodium.

MYRMECOSTYLUM Presl, Hymenoph. 27. 1843 = **Hymenophyllum.**
clavatum Pr. Hym. 28. 1843 = H. polyanthos*.
? dichotomum Pr. Hym. 28. 1843.
tortuosum Pr. Hym. 28 t. 10 f. A. 1843.

Nebroglossa Presl, Epim. Bot. 166. 1849; Fée, Gen. 52 *(Aconiopteris §)* — Elaphoglossum.

NEMATOPERA Kunze, Bot. Zeit. **1845**. 797 (corr. nom.) = **Peranema.**

NEOCHEIROPTERIS Christ, Bull. Soc. bot. France **52** Mém. I. 21. 1905 — (Polypodiaceae Gen. **16**).
Cheiropteris Christ, Bull. Boiss. **6**. 876. 1898.
palmatopedata (Bak.) Christ, Bull. Soc. Fr. **52** Mém. I. 21. 1905 — Yunnan. Polypodium Bak. Kew Bull. **1898**. 232; Cheiropteris Christ, Bull. Boiss. **7**. 21 t. 1. 1899. NPfl. 189; C. Henryi Christ, Bull. Boiss. **6**. 876. 1898.

Neottopteridastrum Fée, Gen. Fil. 190. 1850-52 *(Asplenium §)* = Asplenium.

NEOTTOPTERIS J. Smith, Journ. Bot. **3**. 409. 1841; **4**. 175. 1841 = **Asplenium.**
australasica J. Sm. Cat. cult. ferns 49. 1857 = A. nidus var.
elliptica Fée, Gen. 203. 1850-52 = A. nidus var. ficifolium.
Grevillei J. Sm. JoB. **4**. 176. 1841.
Mauritiana Fée, Gen. 203. 1850-52 = A. nidus.
musaefolia J. Sm. JoB. **3**. 409. 1841 = A, nidus var.
Nidus J. Sm. in Hk. Gen. t. 113 B. 1842.
orientalis Fée, Gen. 203. 1850-52 = A. nidus.
ovata J. Sm.; Fée, Gen. 203. 1850-52 = A. nidus.
pachyphylla Fée, Gen. 203. 1850-52.
Phyllitidis J. Sm. JoB. **3**. 409. 1841 = A. nidus var.
rigida Fée, Gen. 203. 1850-52 = A. nidus.
Simonsiana J. Sm. Hist. Fil. 330. 1875.
simplex Fée, Gen. 203. 1850-52 = A. nidus.
squamulata Fée, Gen. 203. 1850-52.
stenocarpa Fée, Gen. 203. 1850-52 = A. nidus var. ficifolium.
stipitata J. Sm. JoB. **3**. 409. 1841 = A. squamulatum.
taeniosa Fée, Gen. 203. 1850-52 = A. amboinense.
vulgaris J. Sm. JoB. **3**. 409. 1841 = A. nidus.

NEPHRODIUM Richard in Marthe, Cat. Jard. Méd. Paris 120. 1801; Michaux. Fl. bor. Am. **2**. 266. 1803 = **Dryopteris.**
abbreviatum Fée, Gen. 306 1850-52 = Polystichum a.
abortivum J. Sm.; Moore, Ind. 84. 1858.
abruptum J. Sm. JoB. **3**. 411. 1841; Hk. sp. **4**. 77 t. 241 B. 1862 = D. truncata.
abruptum Bedd. Ferns S. Ind. t. 86. 1863-65 = D. megaphylla.
abruptum Bak. Syn. 263. 1867 = D. Kunzeana.
aciculatum Bak. Journ. Linn. Soc. **22**. 226. 1886.
acrosorum Hier. Engl. Jahrb. **34**. 446. 1904.

acrostichoides Michx. Fl. bor. Am. **2**. 267. 1803 = Polystichum a.
acrostichoides Desv. Prod. 255. 1827 — Timor.
acrostichoides J. Sm. JoB. **3**. 411. 1841 = D. canescens*.
acuminatum Pr. Rel. Haenk. **1**. 31. 1825 } = Nephrolepis biserrata.
acutum Pr. Rel. Haenk. **1**. 31. 1825 }
acutum Hk. sp. **4**. 147. pt. 1862.
acutum Hk. sp. **4**. 147. pt. t. 271. 1862 = D. firmifolia.
Adenopteris Moore, Ind. 85. 1858 = D. pennigera.
aemulum Bak. Syn. 279. 1867.
affine Lowe, JoB. **1**. 25. 1834 (nomen); Tr. Cambr. Phil. Soc. **6**. 525. 1838 = D. filix mas.
agatholepis Kiærsk. Cat. Fil. hb. ht. Haun. 62. 1874 = D. patula.
alatellum Christ in Schum. et Laut. Fl. deut. Südsee 112. 1901.
albescens Desv. Prod. 258. 1827 — Jamaica. — Aspidium Mett.; Salom. 1883.
albopunctatum Desv. Prod. 255. 1827 = D. orientalis.
alleopterum Diels, Nat. Pfl. **1**^{4}. 168. 1899 = D. alloëoptera.
alpinum Opiz (? Böh. phan. krypt. Gew. —. 1823) = Cystopteris fragilis* regia.
alsophilaceum Bak. Fl. bras. **1**^{2}. 474. 1870 = D. tenuifolia.
amblyotis Bak. Ann. Bot. **5**. 331. 1891 = Aspidium Veitchianum.
amboinense Pr. Tent. 81. 1836 = D. parasitica*.
amboinense var. evoluta Clarke, Journ. Linn. Soc. **24**. 417. 1888 = D. evoluta.
amphioxypteris Sod. Rec. 51. 1883.
amplissimum Hk. sp. **4**. 145. 1862.
amplum Bory, Bél. Voy. Bot. **2**. 62. 1833 = D. inaequalis.
amplum Hk. sp. **4**. 264. 1862 = D. ameristoneura.
amplum Bak. Syn. 285. 1867.
anateinophlebium Bak. Journ. Linn. Soc. **16**. 202. 1877.
andinum Bak. Ann. Bot. **5**. 331. 1891 = Aspidium a.
androgynum Diels, Nat. Pfl. **1**^{4}. 179. 1899 = D. tetragona.
aneitense Bak. Ann. Bot. **5**. 326. 1891 = D. Lenormandi.
angustifolium Pr. Epim. 48. 1849 = D. hispidula.
angustifrons Bak. Syn. 283. 1867.
antarcticum Bak. Journ. Linn. Soc. **14**. 479. 1875 = D. filix mas*.
antioquianum Bak. JoB. **1881**. 205 = Aspidium a.
aoristisorum Copeland, Polyp. Philipp. 28. 1905.
apicale Bak. Syn. 499. 1874.
apiciflorum Hk. sp. **4**. 112 t. 248. 1862.
apiculatum Pr. Epim. 258. 1849 (nomen) — China.
apifolium Hk. et Arn. Bot. Beech. Voy. 105. 1832 = ? Aspidium cicutarium*. (A. Gaudichaudii Mett.).
appendiculatum Schott, Gen. ad t. 10. 1834; Hk. sp. **4**. 105. 1862 = ? D. prolixa (vel D. cana).
appendiculatum Pr. Epim. 47. 1849 = D. parasitica.
appendiculatum Racib. Pterid. Buitenz. 186. 1898.
aquilinoides Desv. Prod. 261. 1827.
aquilinum Hemsl. Chall. Exp. Bot. **1**^{2}. 167 t. 39. 1885.
arborescens Bak. Syn. 286. 1867 = D. ludens.
arbuscula Desv. Prod. 253. 1827.
arcuatum Desv. Prod. 257. 1827 = D. serra.
argutum Diels, Nat. Pfl. **1**^{4}. 173. 1899 = D. rigida*.

aridum J. Sm. JoB. 4. 188. 1841.
aristatum Pr. Rel. Haenk. 1. 37. 1825 = Polystichum a.
aristatum Pr. Fl. Sicul. XLVI. 1826 = D. rigida *australis.
aristatum Hk. sp. 4. 62 t. 238. 1862 = D. otaria.
articulatum J. Sm. Cat. cult. ferns 54. 1857 = D. megaphylla.
artinexum Clarke, Tr. Linn. Soc. II. Bot. 1. 536. 1880 = Aspidium a.
Ascensionis Hk. sp. 4. 124 t. 257. 1862.
ascendens Donn. Smith, Bot. Gaz. 16. 14. 1891 = Polystichum apiifolium.
asperulum Copeland, Polyp. Philipp. 25. 1905.
asplenioides Michx. Fl. bor. Am. 2. 268. 1803 = Athyrium filix femina.
asplenioides Bak. Syn. 293. 1867.
asplenioides Diels, Nat. Pfl. 1^{4}. 172. 1899 = D. aspidioides.
assamense Hope, JoB. 1890. 326 = D. filix mas*.
athamanticum Hk. sp. 4. 125 t. 258. 1862.
athyrioides Bak. JoB. 1884. 363 = Aspidium costaricanum.
atomiferum Sod. Rec. 48. 1883 — Ecuador. (non in Cr. vasc. quit.).
attenuatum Fée, Gen. 305. 1850-52 = D. stenobasis.
attenuatum Moore, Ind. 86. 1858 = D. invisa.
augescens J. Sm. Bot. Mag. 72 Comp. 32. 1846 = D. serra.
aureo-vestitum Hk. sp. 4. 101 t. 246. 1862 = D. vellea.
auriculare Pr. Epim. 258. 1849 = ? D. lineata.
auriculatum A. Rich. Sert. Astrol. XLIV. 1834 = Polystichum a.
auriculatum Fée, 10 mém. 42 t. 42 f. 2. 1865 = D. arbuscula.
auriculatum Diels, Nat. Pfl. 1^{4}. 171. 1899 = D. squamaestipes.
Bakeri Harr. Journ. Linn. Soc. 16. 29. 1877 = D. canescens*.
? banksiaefolium Pr. Rel. Haenk. 1. 34. 1825 = Osmunda b.
barbigerum Hk. sp. 4. 113. 1862.
Barclayi Bak. Syn. 504. 1874 = Aspidium B.
Baromez Sweet, Hort. Britt. 580. 1827 = Cibotium B.
Barteri Bak. Syn. 299. 1867 = Aspidium B.
basiattenuatum Jenm. Gard. Chr. III. 15. 330. 1894.
basilare Pr. Epim. 258. 1849.
Beccarianum Ces. Atti Ac. Napoli 7^{8}. 23. 1876.
Beddomei Bak. Syn. 267. 1867.
Bedieri Cordem. Bull. soc. sci. et arts Réun. 1890-91. —; Fl. Réun. 77. 1895.
Benoitianum Pr. Tent. 81. 1836 = Polystichum B.
Bergianum Bak. Syn. 269. 1867 = D. prolixa et ? D. Bergiana.
bermudianum Bak. ap. Hemsl. Chall. Exp. Bot. 1^{1}. 86 t. 13. 1885.
biauritum Bedd. Handb. Suppl. 68. 1892.
bibrachiatum Jenm. Gard. Chr. III. 15. 230. 1894.
bidentatum Pr. Rel. Haenk. 1. 32. 1825 = Nephrolepis b.
biforme Diels, Nat. Pfl. 1^{4}. 178. 1899.
biseriale Diels, Nat. Pfl. 1^{4}. 170. 1899.
biserratum Pr. Rel. Haenk. 1. 31. 1825 = Nephrolepis b.
bissetianum Bak. JoB. 1877. 366.
Blanfordii Hope, Journ. Bomb. Nat. Hist. Soc. 12. 624 t. 11. 1899.
Blumei J. Sm. JoB. 3. 411. 1841 = D. canescens.
Blumei Hk. sp. 4. 135. 1862 (part.) = D. rhodolepis.
Boivini Bak. Syn. 501. 1874.
Bojeri Bak. Syn. 280. 1867 = D. filix mas *elongata.

Boottii (Bak. Syn. 275. 1867) = D. cristata × spinulosa.
borbonicum Bak. Ann. Bot. **5**. 317. 1891.
Borneense Hk. sp. **4**. 111. 1862.
Boryanum Hk. sp. **4**. 126. 1862 = D. filix mas *elongata.
Boryanum Bak. Syn. 284. 1867.
brachyodon Hk. sp. **4**. 83 1862.
brachypodum Bak. Tr. Linn. Soc. II. Bot. **2**. 290. 1887.
brachypus Sod. Rec. 43. 1883; Cr. vasc. quit. 228. 1893 = D. opposita var.
Brackenridgei Bak. Syn. 494. 1874.
braineoides Diels, Nat. Pfl. 1^4. 171. 1899.
Braunianum Hk. sp. **4**. 84. 1862.
?bromeliaefolium Pr. Rel. Haenk. **1**. 33. 1825 = Osmunda banksiifolia.
Brownii Desv. Prod. 252. 1827 = Nephrolepis exaltata.
Brunonianum Hk. sp. **4**. 113 t. 251. 1862.
Buchanani Bak. Syn. 498. 1874.
Buckholzii Bak. Ann. Bot **5**. 319. 1891 = D. Buchholzii.
bulbiferum Michx. Fl. bor. Am. **2**. 268. 1803 = Cystopteris b.
caespitosum Diels, Nat. Pfl. 1^4. 172. 1899.
calanthum Endl. Prod. Fl. Norfolk. 9. 1833 — Ins. Norfolk.
calcaratum Hk. sp. **4**. 93. 1862.
calcareum Jenm. JoB. **1886**. 271 = D. tetragona var.
callosum Keys. Pol. Cyath. Hb. Bung. 48. 1873.
Cañadasii Sod. Rec. 48. 1883.
canariense Kiærsk. Cat. Fil. hb. hort. Haun. 62. 1874 = D. filix mas*.
canescens Pr. Tent. 81. 1836; ? Epim. 49. 1849 = D. unita vel D. parasitica.
canescens Christ, Verh. Nat. Ges. Basel **11**. 208. 1895.
canum Bak. Syn. 267. 1867.
carazunense = corazonense.
caribaeum Jenm. JoB. **1886**. 270.
Caripense Hk. sp. **4**. 99. 1862 = D. submarginalis.
Carrii Diels, Nat. Pfl. 1^4. 170. 1899.
catocarpum Hk. sp. **4**. 259. 1862.
catopteron Hk. sp. **4**. 137. 1862 = D. lanuginosa.
caucaense Hier. Engl. Jahrb. **34**. 444. 1904.
caudatum Diels, Nat Pfl. 1^4. 170. 1899.
caudiculatum Sieb.; Pr. Tent. 81. 1836; Epim. 46. 1899 = D. truncata.
caudiculatum J. Sm. JoB. **3**. 411. 1841 = D. basilaris.
chaerophylloides Desv. Prod. 261. 1827 — ? Puerto Rico. (= ? D. pubescens var.).
cheilocarpum Moore, Ind. 88. 1858 = ? D. oligocarpa.
chinense Bak. Syn. 278. 1867 = D. subtripinnata.
chontalense Hemsl. Biol. Centr.-Amer. **3**. 645. 1885.
chrysolobum Fée, Gen. 305. 1850-52.
chrysotrichum Bak. Ann. Bot. **5**. 328. 1891.
cicutarium Bak. Syn. 299. 1867 = Aspidium c.
cicutarium var.? dubia Clarke et Bak. Journ. Linn. Soc. **24**. 417. 1888 =
ciliatum Desv. Prod. 257. 1827 = Microlepia marginata. [Aspidium dubium.
ciliatum Clarke, Tr. Linn. Soc. II. Bot. **1**. 514. 1880 = D. calcarata var.
ciliatum Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874 = D. excelsa.
cirrhosum Bak. Syn. ed. II. 263. 1874.
Clarkei Bak. Syn. 497. 1874.

clypeolutatum Desv. 258. 1827 — Jamaica.
Aspidium (clypeolatum) Mett.; Salom. 1883.
coadunatum Moore, Ind. 88. 1858 = Aspidium c. (Klf.).
coarctatum Hier. Engl. Jahrb. **34**. 444. 1904.
cochleatum Don, Prod. Fl. Nepal. 6. 1825.
cognatum Trevis. Atti Ist. Veneto II. **2**. 165. 1851 (nomen) — Rio Janeiro.
cognatum Hk. sp. **4**. 123 t. 256. 1862.
concinnum Bak. Syn. 268. 1867.
confertum J. Sm. JoB. **4**. 188. 1841 = Cyclodium meniscioides.
confluens F. Müll.; Bak. Syn. 504. 1874 = Aspidium Mülleri.
conforme Sod. Rec. 45. 1883.
coniifolium Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 720. 1903 = Polystichum carvifolium.
conionevron Fée, Gen. 308. 1850-52 = D. extensa.
connexum Kuhn; Bak. Fl. bras. 1^2. 489. 1870 = D. Martini.
consanguineum Moore, Ind. 101. 1858 = ? D. gongylodes.
conspersum Schrad. Gött. gel. Anz. **1824**. 869 = D. patens.
conterminum Desv. Prod. 255. 1827 = D. opposita.
conterminum v. Thouarsianum Hk. sp. **4**. 91. 1862 = D. tomentosa.
contiguum Schott, Gen. ad t. 10. 1834 = D. gongylodes.
continuum Desv. Prod. 254. 1827 = D. gongylodes.
?contractum Moore, Ind. 88. 1858 = Aspidium c.
corazonense Bak. JoB. **1877**. 163 = D. corazonensis.
cordatum Diels, Nat. Pfl. 1^4. 168. 1899.
coriaceum Hope, JoB. **1890**. 328 = Polystichum varium.
?coriandrifolium Desv. Prod. 259. 1827 = Aspidium cicutarium.
costale Bak. Syn. 495. 1874 = D. densisora.
costata Bedd. Ferns br. Ind. t. 220. 1867 = D. penangiana.
costata Diels, Nat. Pfl. 1^4. 179. 1899 = D. pennigera var.
costulare Bak. Journ. Linn. Soc. **16**. 203. 1877.
crassifolium Hk. sp. **4**. 114. 1862.
crassipes Sod. Cr. vasc. quit. 234. 1893.
Creaghii Bak. Kew Bull. **1898**. 230.
crenatum Desv. Prod. 252. 1827 = Nephrolepis exaltata.
crenatum Keys. Pol. Cyath. Hb. Bung. 49. 1873 = D. Poiteana.
crenatum Bak. Fl. Maur. 497. 1877.
crenulaeum Jenm. Bull. Dep. Jam. **3**. 68. 1896.
crinibulbon Hk. sp. **4**. 92 t. 244. 1862 = D. cirrhosa.
crinicaule (Kuhn) Diels, Nat. Pfl. 1^4. 177. 1899 (nomen) — Madagascar.
crinipes Hk. sp. **4**. 71. 1862.
crinitum Desv. Prod. 257. 1827.
crinitum Sod. Cr. vasc. quit. 250, 642. 1893 = ? D. longipetiolata.
crinitum var. nudatum Bak. Fl. Maur. 495. 1877 = D. humida.
cristatum Michx. Fl. bor. Am. **2**. 269. 1803.
cristatum β uliginosum Bak. Syn. 273. 1867 = D. cristata × spinulosa.
Ctenitis Bak. Syn. 265. 1867.
cubense Bak. Syn. 501. 1874.
cucullatum Bak. Syn. 290. 1867 = D. unita.
Cumingianum J. Sm. Bot. Herald 237 t. 50. 1854.
Cumingii J. Sm. JoB. **3**. 411. 1841 = D. extensa.
Currori Bak. Syn. 500. 1874.

?cuspidatum Pr. Rel. Haenk. **1**. 31. 1825 = Leptochilus c.
cuspidatum Bak. Syn. 260. 1867 = D. khasiana.
cyatheoides Pr. Tent. 81. 1836.
?davallioides Desv. Prod. 254. 1827 = Nephrolepis acuminata.
davallioides Bak. Syn. 287. 1867.
Dayi Bedd. JoB. **1887**. 323.
debile Bak. JoB. **1880**. 212 = D. pseudoreptans.
debile Diels, Nat. Pfl. **1**^{4}. 178. 1899.
decadens Bak. JoB. **1886**. 183.
decipiens Hk. sp. **4**. 86 t. 243. 1862.
decipiens Fée, 10 mém. 45. 1865 = D. dicarpa.
decompositum R.Br. Prod. Fl. N. Holl. 149. 1810.
decompositum β microphyllum Hk. sp. **4**. 146. 1862 = D. davallioides.
decrescens Bak. Syn. 497. 1874.
decurrens Bak. Syn. 299. 1867 = Aspidium d.
decurrenti-alatum Diels, Nat. Pfl. **1**^{4}. 171. 1899.
decursivo-pinnatum Bak. Syn. 259. 1867.
decurtatum Fée, Gen. 305. 1850-52 = D. abortiva.
decussatum Diels, Nat. Pfl. **1**^{4}. 171. 1899.
deflexum Pr. Rel. Haenk. **1**. 36 t. 5 f. 2. 1825 = D. Lindigii.
deflexum J. Sm.; Hk. & Bk. Syn. 292 in adnota 1867 = ? D. refracta.
dejectum Jenm. Gard. Chr. III. **18**. 640. 1895.
delicatulum Dcne. Jacq. Voy. **4**. 178 t. 179. 1844 = Nephrolepis cordifolia.
deltoideum Desv. Prod. 259. 1827.
denticulatum Fée, Gen. 305. 1850-52 = Polystichum d.
deparioides Bory, Bél. Voy. Bot. **2**. 59. 1833 = Nephrolepis acuminata.
deparioides Hk. (Fil. Exot. t. 53 et t. 60 in obs. 1859); sp. **4**. 139. 1862.
deversum Rosenst. Hedwigia **43**. 224. 1904 = D. patens.
devolvens Bak. JoB. **1885**. 217.
dicarpum Fée, Gen. 305. 1850-52.
Dickinsii Bak. in Hk. Ic. pl. t. 1659. 1886.
didymosorum Parish, Bedd. Ferns br. Ind. t. 200. 1866.
diffractum Bak. Kew Bull. **1898**. 230.
dilatatum Desv. Prod. 261. 1827 = D. spinulosa*.
dimidiatum Bak. Syn. 500. 1874.
dimorphum Moore, Ind. 90. 1858 = D. heterocarpa.
diplazioides Hk. sp. **4**. 99. 1862 = D. Moritziana.
diplazioides Hier. Engl. Jahrb. **34**. 445. 1904.
disjunctum Moore, Ind. 90. 1858 (nomen) — ? — Aspidium Wall. hb.
dissectum Desv. Prod. 259. 1827.
dissectum Bak. Syn. ed. I. 290. 1867 = D. invisa.
dissidens Hk. sp. **4**. 66. 1862.
dissimile Schrad. Gött. gel. Anz. **1824**. 869 = D. patens.
dissitifolium Bak. Kew Bull. **1899**. 120 = D. papuana.
distans Hk. sp. **4**. 76 in obs. 1862; Bak. Syn. 288. 1867.
distans Diels, Nat. Pfl. **1**^{4}. 170. 1899 = D. brunnea.
diversifolium Diels, Nat. Pfl. **1**^{4}. 179. 1899 = D. vivipara.
diversilobum Pr. Epim. 47. 1849 = D. parasitica vel D. canescens.
divisum Hk. sp. **4**. 133. 1862 = D. Boryana.
domingense Bak. Ann. Bot. **5**. 329. 1891 = Aspidium d.

Dryopteris Michx. Fl. bor. Am. 2. 270. 1803 = D. Linnaeana.
duale Donn. Smith, Bot. Gaz. 15. 29 t. 4. 1890 = Polystichum apiifolium.
Dubrueillianum Hk. et Arn. Bot. Beech. Voy. 105. 1832 = D. cyatheoides.
?duriusculum Pr. Rel. Haenk. 1. 38. 1825 = Polystichum d.
Eatoni Bak. Syn. 276. 1867.
echinatum Bak. Syn. 266. 1867.
Ecklonianum Pappe et Raws. Syn. Fil. Afr. austr. 14. 1858 = D. gongylodes.
Ecklonii Pr. Epim. 49. 1849 = D. gongylodes.
edentulum Bak. Syn. 279. 1867.
edule Don, Prod. Fl. Nepal. 5. 1825 = Nephrolepis cordifolia.
effusum Bak. Syn. 287. 1867.
Eggersii Hier. Engl. Jahrb. 34. 441. 1904.
elatum Desv. Prod. 262. 1827 = D. lanuginosa.
elatum Bak. Syn. 298. 1867 = Aspidium chimborazense.
elatum Bak. Syn. 502. 1874 = D. mauritiana var.
elegans Moore, Ind. 1. 1857 = Abacopteris e.
elegantulum Sod. Cr. vasc. quit. 243. 1893.
elongatum Hk. et Grev. Ic. Fil. t. 234. 1831 = D. filix mas*.
Elwesii Bak. Syn. 497. 1874.
eminens Bak. JoB. 1880. 213.
Endresi Bak. Ann. Bot. 5. 329. 1891 = Aspidium E.
enneaphyllum Bak. JoB. 1887. 170.
eriocarpum Dcne. Arch. Mus. 2. 185. 1841 = D. crenata.
eriocaulon Bak. Syn. 495. 1874.
erubescens Diels, Nat. Pfl. 1^4. 171. 1899.
erythrosorum Hk. sp. 4. 120 t. 253. 1862.
eurostotrichum Bak. JoB. 1880. 329.
eusorum Bedd. Ferns br. Ind. t. 130. 1866 = D. truncata.
Everettii Bak. Kew Bull. 1896. 41 = Aspidium E.
evolutum Bedd. Hdb. Suppl. 76. 1892.
evolutum β Bedd. Hdb. Suppl. 77. 1892 = D. Gustavi.
exaltatum R.Br. Prod. Fl. N. Holl. 148. 1810 = Nephrolepis e.
excellens Hk. sp. 4. 63. 1862 = Aspidium e.
excisum Fée, 10 mém. 44 t. 44 f. 2. 1865 — Ceylon.
excultum Hk. sp. 4. 149. 1862 = D. effusa*.
exiguum Hk. sp. 4. 92. 1862 = D. philippina.
expansum Pr. Rel. Haenk. 1. 38. 1825 = D. spinulosa *dilatata.
expansum Desv. Prod. 261. 1827 = Aspidium martinicense.
extensum Moore, Ind. 91. 1858.
extensum β minor Bedd. Ferns br. Ind. t. 201. 1866 = D. parasitica.
extensum var. microsora Clarke, Tr. Linn. Soc. II. Bot. 1. 530. 1880 = D. [molliuscula.
Faberi Bak. Ann. Bot. 5. 316. 1891.
falciculatum Desv. Prod. 259. 1827.
falcilobum Hk. sp. 4. 108. 1862 = D. calcarata var.
Falconeri Hk. sp. 4. 123 t. 254. 1862 = D. barbigera.
fasciculatum Bak. Ann. Bot. 5. 320. 1891.
Fauriei Christ, Bull. Boiss. 4. 671. 1896 = Athyrium F.
Fendleri Hk. sp. 4. 82. 1862.
ferox Moore, Ind. 91. 1858.
ferrugineum Bak. Syn. 283. 1867 = D. peranemiformis.

fibrillosum Bak. Journ. Linn Soc. 15. 418. 1876 = D. Poolii.
Fijiense Hk. 2 Cent. t. 67. 1861.
Filix femina Michx. Fl. bor. Am. 2. 268. 1803 = Athyrium f.
Filix mas Rich. in Marthe, Cat. jard. méd. Paris 129. 1801.
Filix mas var. odontoloma Bak. Syn. 498. 1874 = D. filix mas *serrato-dentata.
firmifolium Bak. Syn. 501. 1874.
firmum Bak.; Jenm. JoB. 1879. 260.
Fischeri Bak. Fl. bras. 1^2. 479. 1870.
flaccidum Hk. sp. 4. 133 t. 263. 1862.
flexuosum Bak. Fl. bras. 1^2. 483. 1870.
floridanum Hk. Fil. exot. t. 99. 1859.
Foenisecii Lowe, Tr. Cambr. Phil. Soc. 4. 7. 1830; 6. 524. 1838 = D. aemula.
Fordii Bak. JoB. 1889. 177 = D. crenata.
Fournieri Bak. Ann. Bot. 5. 317. 1891.
Fournieri Bak. Ann. Bot. 5. 328. 1891 = Aspidium novae caledoniae.
fragile Bak. Ann. Bot. 5. 322. 1891.
fragrans Richards. in Franklin Narr. of a Journ. 754. 1823.
Fraseri Bak. Syn. 495. 1874.
Fraserianum Gaud. Freyc. Voy. Bot. 339. 1827 — Australia.
(prob. = D. decomposita).
funestum Hk. sp. 4. 129 t. 259. 1862 = D. protensa var.
Funkii Bak. Syn. 496. 1874 = D. Funckii.
furcatum Hk. sp. 4. 136. 1862.
fuscipes Clarke, Tr. Linn. Soc. II. Bot. 1. t. 75. 1880 = Aspidium membranifolium.
Gaimardianum Gaud. Freyc. Voy. Bot. 335 t. 12 f. 1. 1827 = Humata G.
Gamblei Hope, Journ. Bomb. Nat. Hist. Soc. 12. 533 t. 7. 1899.
Gardneri Bak. Ann. Bot. 5. 330. 1891 = Aspidium organense.
Gardnerianum Bak. Fl. bras. 1^2. 474. 1870 = D. densiloba.
geropogon Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874.
Ghiesbreghtii Diels, Nat. Pfl. 1^4. 179. 1899.
gibbosum Gaud. Freyc. Voy. Bot. 338. 1827 = Nephrolepis biserrata.
giganteum Bak. Syn. ed. I. 300. 1867 = Aspidium Thwaitesii.
giganteum Bak. Syn. ed. II. 503. 1874 = Aspidium g.
giganteum Ces. Rend. ac. sci. Napoli 16. 15. 1877 = Aspidium andaiense.
giganteum Diels, Nat. Pfl. 1^4. 177. 1899.
giganteum Diels, Engl. Jahrb. 29. 189. 1900 = Diplazium maximum var.
Gilberti Clute, Fern Bull. 10. 90. 1902.
glabellum A. Cunn. in Hk. Comp. Bot. Mag. 2. 367. 1836.
glaberrimum Moore, Ind. 93. 1858 = Aspidium g.
glabratum Bak. Syn. 500. 1874.
glabrum Bak. Syn. 278. 1867.
glanduliferum Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874.
glanduligerum (Bak. Syn. 262. 1867) = D. gracilescens*.
glandulosum J. Sm. JoB. 3. 411. 1841; Pr. Epim. 45. 1849 = D. urophylla.
glandulosum Moore, Ind. 1. 1857.
glandulosum Hk. sp. 4. 76. 1862; Bak. Syn. 291. 1867 = { D. glandulosa. D. lineata. D. urophylla. }
glaucostipes Bedd. Handb. Suppl. 80. 1892.
globuliferum Hk. sp. 4. 96. 1862.

Goldianum Hk. et Grev. Ic. Fil. t. 102. 1829.
gongyloides Schott, Gen. Fil. ad t. 10. 1834 = D. gongylodes.
gracilescens Hk. sp. **4**. 93. 1862.
grammitoides Copeland, Polyp. Philipp. 21. 1905.
grande Bak. Ann. Bot. **5**. 330. 1891 = Aspidium grandifolium.
granulosum J. Sm. Bot. Voy. Herald 428. 1857.
granulosum Bak. JoB. **1891**. 4 = D. subgranulosa.
grenadense Jenm. Gard. Chr. III. **15**. 198. 1894.
Griffithii Bak. Syn. 300. 1867 = Aspidium multicaudatum.
Grisebachii Bak. Syn. 285. 1867 = D. ameristoneura.
griseum Bak. Syn. 271. 1867.
Grunowii Moore, Ind. 93. 1858.
guadalupense Fée, 11 mém. 89 t. 24 f. 3. 1866.
guatemalense Bak. Syn. 498. 1874.
Gueintzianum Hier. Engl. Jahrb. **28**. 341. 1900.
guianense Desv. Prod. 253. 1827 = ? Nephrolepis biserrata.
Gustavi Bedd. JoB. **1893**. 227.
gymnogrammoides Diels, Nat. Pfl. **1**[4]. 170. 1899.
gymnophyllum Bak. JoB. **1887**. 170.
gymnopodum Bak. Tr. Linn. Soc. II. Bot. **4**. 252. 1894.
gymnosorum Mak. Bot. Mag. Tokyo **13**. 64. 1899.
Haenkeanum Pr. Epim. 46. 1849.
Harrisoni Bak. Ann. Bot. **5**. 326. 1891.
Harveyi Bak. Syn. 497. 1874.
hastaefolium Diels, Nat. Pfl. **1**[4]. 168. 1899 = D. sagittata.
hastatum Jenm. JoB. **1879**. 261.
hederaefolium Bak. Journ. Linn. Soc. **19**. 295. 1882 = Aspidium h.
Hendersoni Bak. Ann. Bot. **5**. 323. 1891 = D. hirsutifrons.
heptaphyllum Bak. JoB. **1879**. 296 = Aspidium h.
heterocarpum Moore, Ind. 93. 1858.
heterodon Moore, Ind. 93. 1858 = Mesoschlaena polycarpa.
heterophyllum Hk. sp. **4**. 62. 1862 = D. canescens*.
heteropterum Desv. Prod. 256. 1867.
heterosorum Bak. Syn. 504. 1874 = Aspidium h.
hexagonopterum Diels, Nat. Pfl. **1**[4]. 170. 1899.
Hilsenbergii Pr. Tent. 81. 1836; Epim. 47. 1869 = D. parasitica var.
Hippocrepis Pr. Rel. Haenk. **1**. 36. 1825 = Aspidium h.
hirsutulum Pr. Rel. Haenk. **1**. 32. 1825 = Nephrolepis h.
hirsutum Don, Prod. Fl. Nepal. 6. 1825 = D. crenata.
hirsutum Bory, Dup. Voy. Bot. **1**. 268. 1828 = ? Nephrolepis hirsutula.
hirsutum J. Sm. JoB. **3**. 412. 1841 (nomen); Pr. Epim. 48. 1849 = D. adenophora.
hirsutum Diels, Nat. Pfl. **1**[4]. 177. 1899 = D. hirtula.
hirtipes Hk. sp. **4**. 115 t. 249. 1862.
hirtum Hk. sp. **4**. 128. 1862.
hirtum Copeland, Polyp. Philipp. 25. 1905 = D. setosa.
hispidulum Bak. Syn. 293. 1867.
hispidum Hk. sp. **4**. 150. 1862 = Polystichum h.
Hoffmanseggii Desv. Prod. 252. 1827 = Nephrolepis biserrata.
Holmei Bak. Ann. Bot. **5**. 317. 1891.

Hookeri Moore et Houlst. Gard. Mag. Bot. **3**. —. 1851; J. Sm. Cat. cult. Ferns [54. 1857 = D. arbuscula.
Hopeanum Bak. Syn. 494. 1874.
Hopei Bak. Ann. Bot. **5**. 326. 1891 = D. molliuscula.
Hornei Bak. Syn. 500. 1874.
Hosei Bak. Ann. Bot. **5**. 330. 1891 = Aspidium H.
Hudsonianum Brack. Expl. Exp. **16**. 188 t. 25. 1854 = D. truncata.
Humblotii Bak. Bull. Soc. Linn. Paris **1**. 534. 1885.
humidum Cord. Fl. Réun. 74. 1895.
imbricatum Boj. Hort. Maur. 392. 1837 = Nephrolepis cordifolia.
immersum Hk. sp. **4**. 112. 1862.
impressum Desv. Prod. 259. 1827 — Timor.
impressum Fée, Gen. 305. 1850-52 = D. immersa.
Imrayanum Hk. sp. **4**. 86. 1862 = D. rotundata var.
inaequale Schrad. Gött. gel. Anz. **1824**. 869 — Brasilia. (Nephrolepis sp.).
inaequale Hk. sp. **4**. 125. 1862.
inaequilaterale Bak. Syn. ed. I. 454. 1868 = D. Brackenridgei.
inaequilaterum Col. Tr. N. Zeal. Inst. **20**. 229. 1888 = D. gongylodes.
incisum Bak. Syn. 288. 1867.
indecorum Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874.
ingens Atk.; Clarke, Tr. Linn. Soc. II. Bot. **1**. 526 t. 73. 1880 = D. dissecta.
inquinans Fée, 10 mém. 43 t. 44 f. 1. 1865 — Réunion.
Aspidium Kuhn 1868 (non Fée 1857).
insculptum Desv. Prod. 254. 1827 = D. gongylodes.
insigne Bak. Syn. 262. 1867.
intermedium Pr. Rel. Haenk. **1**. 38. 1825 = D. spinulosa*.
intermedium Bak. Syn. 283. 1867 = D. rhodolepis.
invisum Desv. Prod. 257. 1827 = D. patens*.
invisum Carr. in Seem. Fl. Vit. 362. 1873.
irregulare Bak. Syn. ed. I. 298. 1867 = Aspidium latifolium.
irriguum Bak. Syn. 297. 1867 = Aspidium i.
isolatum Bak. Gard. Chr. n. s. **14**. 494. 1880 = D. Sheareri.
jamaicense Bak.; Jenm. JoB. **1877**. 264.
Jamesoni Hk. sp. **4**. 66. 1862 = D. parasitica var.
japonicum Bak. Ann. Bot. **5**. 318. 1891.
javanicum Hk. Fil. exot. t. 61. 1859 = Mesochlaena polycarpa.
Jenmani Bak.; Jenm. JoB. **1877**. 263.
Juergensii Rosenst. Festschr. Alb. v. Bamberg 63. 1905.
juglandifolium Bak. JoB. **1879**. 296 — Samoa.
Aspidium Christ 1890 (non Kze. 1847). (= ? Aspidium pachyphyllum).
kanakorum Bak. Ann. Bot. **5**. 331. 1891 = Aspidium k.
Karwinskyanum Bak. Syn. 279. 1867.
Kaulfussii Hk. sp. **4**. 97. 1862 = D. oligocarpa*.
Keckii Bak. Ann. Bot. **5**. 330. 1891 = Aspidium K.
Keraudrenianum Diels, Nat. Pfl. 1^4. 177. 1899.
kilmense Bak. Syn. 498. 1874 = D. kilemensis.
Kingii Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 621 t. 9. 1899 = D. filix mas*.
Krameri Diels, Nat. Pfl. 1^4. 170. 1899 = D. oyamensis.
Kuhnii Hier. Engl. Jahrb. **34**. 440. 1904.
Kunthii Desv. Prod. 258. 1897 = D. patens
Kunzeanum Hk. sp. **4**. 102. 1862.

Labrusca Racib. Bull. Acad. Cracovie **1902**. 61 = Aspidium L.
lacerum Bak. Syn. 273. 1867.
laetum Komarov, Fl. Mansh. 124. 1901.
laeve Moore, Ind. 94. 1858.
Lagerheimii Sod. Cr. vasc. quit. 252. 1893.
lancastriense Desv. Prod. 260 = D. cristata.
lanceolatum Bak. Syn. 498. 1874.
lanceum Moore, Ind. 95. 1858 = Athyrium macrocarpum.
lancilobum Bak. Syn. 499. 1874.
lanosum Michx. Fl. bor. Am. **2**. 270 1803 = Cheilanthes l.
lanuginosum Desv. Prod. 262. 1827.
lanuginosum Hassk. Tijdschr. Nat. Gesch. **10**. 116. 1843 — Java.
larutense Bedd. Handb. Suppl. 73. 1892.
lasiopteris Sod. Rec. 45. 1883.
Lastii Bak. Ann. Bot. **5**. 324. 1891.
latebrosum Moore, Ind. 95. 1858.
latifolium Pr. Epim. 45. 1849 = D. urophylla.
latifolium Bak. Syn. ed. I. 297. 1867; ed. II. 298. 1874 = Aspidium l.
latifrons Hk. sp. **4**. 138. 1862.
latipinna Hk. Syn. 292. 1867 = D. parasitica var.
Laurentiorum Christ, Mission E. Laurent 5. 1905.
lawrenceanum Bak. JoB. **1884**. 143 = Aspidium L.
laxum Diels, Engl. Jahrb. **29**. 189. 1900.
Layardi Bak. Ann. Bot. **5**. 323. 1891.
Lazazaburii Bak. Ann. Bot. **5**. 331. 1891 = Aspidium Lazarzaburoi.
Lechleri Hier. Engl. Jahrb. **34**. 448. 1904.
lepidum Bak. Ann. Bot. **5**. 318. 1891.
lepigerum Bak. Syn. 284. 1867.
Lepinei Bak. Syn. 501. 1874.
Leprieurii Hk. sp. **4**. 106. 1862.
leuconevron Fée, Gen. 306 t. 18 C. f. 3. 1850-52 — Réunion.
Aspidium Kuhn 1868.
leucorhachis Cheesem. Tr. Linn. Soc. II. Bot. **6**. 309. 1903.
leucostipes Bak. JoB **1885**. 105.
Leuzeanum Hk. sp. **4**. 61. 1862 = Aspidium L.
Levyi Bak. Syn. 502. 1874.
ligulatum Hk. sp. **4**. 112. 1862.
limbatum Desv. Prod. 260. 1827.
Lindeni Bak. Syn. 493. 1874.
lineatum Pr. Epim. 48. 1849.
lineatum Bedd. Ferns br. Ind. t. 133. 1866 = D. urophylla.
Linkianum Diels, Nat. Pfl. **1**[4]. 172. 1899 = D. diplazioides.
Lizarzaburui (oi ex err.) Sod. Rec. 55. 1883 = Aspidium L.
Lobbii Bak. Syn. 297. 1867 = Aspidium L.
loherianum Copeland, Polyp. Philipp. 30. 1905.
lonchodes Hk. sp. **4**. 99. 1862.
longicaule Bak. JoB. **1881**. 204.
longicuspe Bak. Journ. Linn. Soc. **16**. 202. 1877.
longifolium Hier. Engl. Jahrb. **34**. 449. 1904 = D. reticulata.
longipes Moore, Ind. 95. 1858.

longissimum Diels, Nat. Pfl. 1^{4}. 179. 1899.
lucens Boj.; Fée, Gen. 305. 1850-52 = D. gongylodes.
?lucidulum Pr. Rel. Haenk. **1**. 39. 1825 = Davallia solida.
lucidum Bak. Gard. Chr. n. s. **8**. 456. 1877.
ludens Bak. Ann. Bot. **5**. 325. 1891.
Luerssenii Harr. Journ. Linn. Soc. **16**. 29. 1877
lugubre Rosenst. Hedwigia **43**. 224. 1904; Hier. Engl. Jahrb. **34**. 442. 1904.
lustratum Hier. Engl. Jahrb. **34**. 443. 1904.
Macarthyi Bak. Ann. Bot. **5**. 321. 1891.
macradenium Sod. Rec. 47. 1883.
macrophyllum Bak. Syn. 300. 1867 = Aspidium martinicense.
macrophyllum Keys. Pol. Cyath. Hb. Bung. 48. 1873 = D. reticulata*.
macrosorum Bak. Journ. Linn. Soc. **19**. 295. 1882 = Aspidium m.
macrostegium Hk. sp. **4**. 148. 1862.
macrotis Hk. sp. **4**. 86 t. 242 B. 1862.
macrourum Schott, Gen. Fil. ad t. 10. 1834 = D. patens*.
macrourum Hk. sp. **4**. 96. 1862 = D. patens.
magnum Bak. JoB. **1884**. 142.
malabariense Fée, 10 mém. 43. 1865 — »Malabar«.
mamillosum Bak. Ann. Bot. **5**. 330. 1891 = Aspidium m.
Mannii Hope, JoB. **1890**. 145 = D. rhodolepis.
marginale Michx. Fl. bor. Am. **2**. 267. 1803.
marginale β umbilicatum Desv. Prod. 260. 1827 = D. filix mas.
marginatum Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 740 t. 33. 1903 = D.
mascaranense Bak. Syn. 503. 1874 = D. mascarensis. [filix mas*.
Matsumurae Mak. Bot. Mag. Tokyo **13**. 48. 1899.
mauritianum Desv. Prod. 252. 1827 = Nephrolepis biserrata.
Mauritianum Fée, Gen. 308. 1850-52.
Maximowiczii Bak. Syn. 499. 1874.
maximum Bak. JoB. **1886**. 182.
megaphyllum Bak. Journ. Linn. Soc. **22**. 227. 1886 = D. subarborea.
melanocaulon Bak. Syn. 296. 1867 = Aspidium m.
melanopus Hk. sp. **4**. 110. 1862 = D. sagenioides.
melanorachis Bak. JoB. **1888**. 325 = Aspidium m.
membranifolium Pr. Rel. Haenk. **1**. 36 t. 5 f. 3. 1825 = Aspidium m.
membranifolium Hk. sp. **4**. 131. 1862 = { Aspidium m. (t. 261).
Dryopteris dissecta.
menisciinerve Moore, Ind. 96. 1858 = Aspidium m.
meniscioides J. Sm. JoB. **4**. 188. 1841 = Cyclodium m.
meniscioides Clarke, Tr. Linn. Soc. II. Bot. **1**. 532. 1880 = D. glandulosa.
mexicanum Pr. Rel. Haenk. **1**. 38. 1825 = D. patula.
Meyanthidis Bak. Syn. 297. 1867 = Aspidium menyanthidis.
microbasis Bak. Syn. 496. 1874.
microcarpon Fée, Gen. 307. 1850-52 = D. unita.
microcarpum Pr. Tent. 81. 1836 = D. gongylodes.
micuochlamys Bak. Journ. Linn. Soc. **15**. 107. 1876 = Mesochlaena polycarpa.
microsorum Endl. Prod. Fl. Norf. 9. 1833 = D. decomposita.
microsorum Hk. sp. **4**. 106. 1862 = D. leptosora.
microsorum Bedd. Hdb. 270. 1883 = D. molliuscula.
microstegium Hk. sp. **4**. 119 t. 250. 1862 = D. brunnea.

Milnei Hk. 2 Cent. t. 62. 1861.
Miquelianum Yabe, Bot. Mag. Tokyo **17**. 65. 1903.
mohrioides Desv. Prod. 261. 1827 = Polystichum m.
molle R.Br. Prod. Fl. N. Holl. 149. 1810 = D. parasitica.
molliculum Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874.
molliusculum Bedd. Handb. Suppl. 68. 1892.
mongolicum Bak. Ann. Bot. **5**. 317. 1891.
monocarpum Cordem. Fl. Réun. 74. 1895.
montanum Bak. Syn. 271. 1867 = D. oreopteris.
monticola Mak. Bot. Mag. Tokyo **13**. 80. 1899.
Motleyanum Hk. Syn. 266 in adnota 1867.
moulmeinense Bedd. Ferns br. Ind. Corrections 1870.
mucronatum J. Sm. JoB. **3**. 412. 1841 = D. arida, Haenkeana, hispidula, parasitica, f. auctt. divers.
multicaudatum Clarke, Tr. Linn. Soc. II. Bot. **1**. 540 t. 77. 1880 = Aspidium.
multifidum Rich. Sert. Astrol. XXXIX. 1834 — Vanikoro. — (Nephrolepis spec. [dub.).
multijugum Bak. Syn. 291. 1867 = D. ochthodes.
multilineatum Pr. Tent. 81. 1836; Diels in Laut. et Schum. Fl. deut. Südsee 113. 1901 = D. Haenkeana.
multisetosum Bak. Ann. Bot. **5**. 324. 1891 } = D. multiseta.
multisetum Bak. Journ. Linn. Soc. **22**. 227. 1886 } = D. multiseta.
muricatum Diels, Nat. Pfl. 1^4. 179. 1899 = D. subspinosa.
myriolepis Bak. JoB. **1888**. 34.
namburense Bedd. Handb. Suppl. 69. 1892.
Napoleonis Bory, Bél. Voy. Bot. **2**. 61 t. 6. 1833.
nebulosum Bak. JoB. **1880**. 213 = Aspidium n.
negligens Jenm. Bull. Dep. Jam. **3**. 21. 1896.
nemorale Sod. Cr. vasc. quit. 267. 1893.
nemorosum Desv. Prod. 261. 1827.
nervosum Hier. Engl. Jahrb. **34**. 445. 1904.
nevadense Bak. Ann. Bot. **5**. 320. 1891 = D. oregana.
Nidus Clarke, Journ. Linn. Soc. **15**. 156. 1876 = D. filix mas*.
nigrescens Bak. Syn. 504. 1874 = Aspidium n.
nigritanum Bak. Syn. 499. 1874 = D. tomentella.
nigrovenium Christ; Donn. Smith, Bot. Gaz. **20**. 545. 1895.
nimbatum Jenm. Gard. Chr. III. **15**. 264. 1894.
nitidulum Pr. Epim. 46. 1849 = Aspidium n.
nitidulum Bak. Fl. bras. 1^2. 597. 1870 = D. anoptera.
nitidum Desv. Prod. 256. 1827 = D. crinita.
Nockianum Jenm JoB. **1886**. 270.
novaeboracense Desv. Prod. 257. 1827.
nudum Bak. JoB. **1879**. 41 = Aspidium n.
nymphale Desv. Prod. 258. 1827 = D. parasitica.
obliquatum Bak. Syn. 264. 1867.
obliteratum R.Br. Prod. Fl. N. Holl. 148. 1810 = Arthropteris o.
obliteratum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
obovatum Bak. JoB. **1890**. 265.
obscurum Moore, Ind. 98. 1858; Racib. Pterid. Buit. 185. 1898 = D. arida.
obscurum Diels, Nat. Pfl. 1^4. 173. 1899 = D. sagenioides*.
obtusatum Pr. Rel. Haenk. **1**. 35. 1825 = D. gongylodes.

obtusatum Desv. Prod. 254. 1827 = D. pteroides.
obtusatum Diels, Nat. Pfl. $\mathbf{1}^4$. 171. 1899 = D. opaca.
obtusifolium Pr. Rel. Haenk. **1**. 32. 1825 = Nephrolepis radicans.
obtusilobum Bak. Syn. 284. 1867 = D. peranemiformis var.
ochropteroides Bak. Ann. Bot. **5**. 325. 1891.
ochrorachis Bak. JoB. **1884**. 142.
ochthodes Hk. sp. **4**. 109. 1862.
occultum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 627 t. 13. 1899.
odontocarpos Rich. Sert. Astrol. XLIV. 1834 — Ins. Mariannae.
odontoloma Clarke, Tr. Linn. Soc. II. Bot. **1**. 521. 1880 = D. filix mas *serrato-dentata.
odontoloma Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 736 t. 31. 1903 = D.
odoratum Bak. Syn. 280. 1867 = D. crenata. [filix mas*.
oligocarpum Desv. Prod. 256. 1827.
oligodictyon Bak. Ann. Bot. **5**. 328. 1891 = Aspidium megalocarpum.
oligodontum Desv. Prod. 261. 1827 = D. filix mas*.
oligophlebium Bak. JoB. **1875**. 291.
omeiense Diels, Nat. Pfl. $\mathbf{1}^4$. 171. 1899.
oosorum Bak. Kew Bull. **1896**. 41.
oppositum Fée, 8 mém. 108. 1857 — Martinique.
oppositum Hk. sp. **4**. 136 t. 266. 1862 = D. mascarenarum.
oppositum Diels, Nat. Pfl. $\mathbf{1}^4$. 172. 1899.
opulentum Pr. Tent. 81. 1836 = D. pteroides.
oreopteris Desv. Prod. 257. 1827.
Oreopteris Fée, Gen. 306. 1850-52 = D. pteroides.
orientale Desv. Prod. 260. 1827 — »in Oriente«.
ornatum Christ, Verh. Nat. Ges. Basel **11**. 235. 1895.
Otaria Bak. Syn. 288. 1867.
oxyodon Copeland, Polyp. Philipp. 23. 1905.
pachyphyllum Bak. Syn. 299. 1867 = Aspidium p.
pachyrachis Hk. sp. **4**. 100. 1862.
Palatanganum Hk. sp. **4**. 260. 1862 = D. ampla.
paleaceum Diels, Nat. Pfl. $\mathbf{1}^4$. 177. 1899 = D. furcata*.
pallidivenium Bak. Syn. 290. 1867 = D. striata.
pallidum Bory, Exp. scient. de Morée $\mathbf{3}^2$. 287 t. 36. 1832 = D. rigida *australis.
pallidum Hemsl. Biol. Centr.-Amer. **3**. 648. 1885 = D. chlorophylla.
palmoides Rich. Sert. Astrol. XXXIX. 1834 = Nephrolepis biserrata.
paludosum Liebm. Vid. Selsk. Afh. V. **1**. 275 (seors. 123). 1849 = D. gongylodes.
palustre Bak. Syn. 270. 1827.
panamense Pr. Rel. Haenk. **1**. 35. 1825 = D. opposita.
pandum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 623. 1899 = D. filix mas*.
Papilio Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 625 t. 12. 1899.
papyraceum Bedd. Handb. Suppl. 69. 1892.
parallelogrammum Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 728. 1903 = D. filix mas *khasiana et *patentissima.
parallelum Bak. Journ. Linn. Soc. **15**. 417. 1876.
parasiticum Desv. Prod. 260. 1827.
Parishii Hk. sp. **4**. 131 t. 260. 1862.
patens Desv. Prod. 258. 1827.
patens J. Sm. Cat. cult. ferns 54. 1857 = D. parasitica.

patens J. Sm. Hist. Fil. 208. 1875 = D. silvatica (t. Syn. cit.).
patentissimum Clarke, Journ. Linn. Soc. **15**. 156. 1876 = D. filix mas*.
patulum Bak. Fl. bras. 1². 480. 1870.
pauciflorum Kiærsk. Cat. Fil. hb. ht. Haun. 63. 1874 = D. crinita.
paucijugum Jenm. JoB. **1886**. 270 = D. chartacea.
Pavonianum Hier. Engl. Jahrb. **34**. 445. 1904.
pectinatum Link, Hort. Berol. **2**. 125. 1833 — Nephrolepis p.
pectinatum Hier. Engl. Jahrb. **28**. 341. 1899 = D. orientalis.
pedatum Hk. sp. 4. 84. 1862.
pellitum Desv. Prod. 255. 1827 = ? D. parasitica.
pellucidum Diels, Engl. Jahrb. 29. 191. 1900.
pendulum Desv. Prod. 252. 1827 = Nephrolepis cordifolia*.
pennigerum Pr. Rel. Haenk. 1. 35. 1825.
pennigerum Moore, Ind. 100. 1858 = D. megaphylla.
pentangularum Col. Tasm. Journ. **2**. 169. 1844 = D. decomposita.
Pentheri Krasser, Ann. Hofm. Wien 15. 5 t. 2 f. 1-5. 1900.
perakense Bak. Ann. Bot. 5. 319. 1891.
Peripae Sod. Rec. 52. 1883.
phanerophlebium Bak. Syn. 494. 1874.
Phegopteris Prantl, Exkursionsflora v. Baiern 23. 1884.
philippinense Bak. Ann. Bot. **5**. 327. 1891 = D. basilaris.
Pica Bak. Syn. 298. 1867 = Aspidium p.
pilosiusculum Racib. Pterid. Buit. 189. 1898.
piloso-hispidum Hk. sp. **4**. 105. 1862 = D. rudis.
pilosulum Hk. sp. **4**. 102. 1862 = ? D. mollicula.
Pittsfordensis Woolson, Fern. Bull. 12. 106. 1904.
Plantianum Pappe et Raws. Syn. Fil. austr. 14. 1858 — Natal.
Aspidium Kuhn 1868.
platyphyllum Desv. Prod. 261. 1827 = Polystichum aculeatum var. 23.
platypus Hk. sp. **4**. 149. 1862 = Polystichum aristatum var.
plectochlaena Fée, Gen. 307. 1850-52 = D. unita.
pleiopodum Bak. Ann. Bot. **5**. 328. 1891 (err.) = N. pteropodum.
pleiotomum Bak. Syn. ed. II. 297. 1874 = Aspidium p.
plumiferum Desv. Prod. 254. 1827 = D. arida.
Plumula Pr. Rel. Haenk. **1**. 33. 1825 = Polystichum munitum.
podophyllum Hk. sp. 4. 87. 1862.
Pohlianum Pr. Tent. 81. 1836 = D. gongylodes.
***Poiretii* Desv. Prod. 258. 1827 — Martinique.**
Polypodium pubescens Poir. Enc. **5**. 546. 1804 (»non L.« 1759).
polycarpum Keys. Pol. Cyath. Hb. Bung. VII. 1873 = Mesochlaena p.
polycarpum Diels, Nat. Pfl. 1⁴. 179. 1899 = D. stegnogrammoides.
polylepis Bak. Ann. Bot. 5. 320. 1891.
polylepis Sod. Cr. vasc. quit. 260. 1893 = D. fusca.
polymorphum Opiz, Kratos **1819**. IV = Cystopteris fragilis.
polymorphum Bak. Syn. 297. 1867 = Aspidium p.
polyphyllum Pr. Rel. Haenk. **1**. 37. 1825 = Polystichum aculeatum var. 24.
polypodiforme Mak. Bot. Mag. Tokyo **13**. 58. 1899.
polytrichum Schrad. Gött. gel. Anz. **1824**. 869 = D. patens.
polytrichum Bak. JoB. **1891**. 107 = D. trichopoda.
popayanense Hier. Engl. Jahrb. 34. 447. 1904.

Powellii Bak. Ann. Bot. **5**. 325. 1891.
praetervisum Bak. Syn. 495. 1874.
Prenticei Bak. Syn. 494. 1874.
Preslii Bak. Syn. 272. 1867.
Prionitis Diels, Nat. Pfl. **1**^{4}. 169. 1899.
prionophyllum Moore, Ind. 101. 1858 = Aspidium p. (Wall.).
prismaticum Desv. Prod. 256. 1827 = D. truncata.
procerum Don. Prod. Fl. Nepal. 6. 1825 — Nepal. — Aspidium Spr. 1827.
procerum Bak. Syn. 502. 1874 = D. mauritiana.
procurrens Bak. Syn. 290. 1867.
productum Pr. Tent. 81. 1836.
proliferum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
prolificum Diels, Nat. Pfl. **1**^{4}. 173. 1899 = D. erythrosora f. monstr.
prolixum Desv. Prod. 256. 1827.
propinquum R.Br. Prod. 148. 1810 = D. gongylodes*.
propinquum Pr. Rel. Haenk. **1**. 34. 1825 = D. arida.
protensum Bak. Fl. bras. **1**^{2}. 481. 1870.
pseudomontanum Rosenst. Hedwigia **43**. 225. 1904.
pseudotetragonum Hier. Engl. Jahrb. **34**. 442. 1904.
pseudothelypteris Rosenst. Hedwigia **43**. 225. 1904.
Ptarmica Bak. Fl. bras. **1**^{2}. 479. 1870.
pterifolium Bak. Syn. 497. 1874.
pteroides J. Sm. Cat. cult. ferns 54. 1857.
pteroideum Diels, Nat. Pfl. **1**^{4}. 171. 1899.
pteropodum Bak. JoB. **1888**. 325 = Aspidium p.
puberulum Bak. Syn. 495. 1874 = D. Feei.
puberulum Bak. JoB. **1875**. 201 = D. Macarthyi.
puberum J. Sm. Cat. cult. ferns 54. 1857 = D. arbuscula.
pubescens Don, Prod. Fl. Nepal. 6. 1825 = D. prolixa.
pubescens Desv. Prod. 261. 1827.
pubescens Brack. Expl. Exp. **16**. 186. 1854 — Ins. Philipp.
pubirachis Bak. JoB. **1876**. 344.
pulchrum Desv. Prod. 256. 1827 = D. strigosa.
pulvinuliferum Bak. Syn. 500. 1874.
punctatum Parish, Bedd. Ferns br. Ind. t. 131. 1866 = D. extensa.
punctatum Bak. Syn. 503. 1874 = D. ochthodes.
punctatum Diels, Nat. Pfl. **1**^{4}. 177. 1899.
punctilobulum Michx. Fl. bor. Am. **2**. 268. 1803 = Dennstaedtia p.
punctulatum Desv. Prod. 253. 1827 = Nephrolepis biserrata*.
punctulatum Bak. Syn. 261. 1867 = D. orientalis.
purpurascens Hk. sp. **4**. 132 t. 262 = D. sparsa*.
pusillum Bak. Syn. 260. 1867.
quadrangulare Fée, Gen. 308. 1850-52 — Guiana.
quinquefidum Bak. JoB. **1890**. 265 = Aspidium q.
raddi Desv. Prod. 258. 1827 = D. patens.
Raddianum Hk. sp. **4**. 98 t. 245. 1862 = D. deflexa.
ramentaceum Bak. Fl. bras. **1**^{2}. 473. 1870 = D. eriocaulis.
ramosum Hope, JoB. **1896**. 126.
rampans Bak. JoB. **1889**. 177.
recedens Hk. sp. **4**. 135 t. 265. 1862.

refractum Hk. sp. **4**. 162 t. 252. 1862.
refulgens Diels, Nat. Pfl. 1^4. 170. 1899.
regulare Desv. Prod. 252. 1827 = Nephrolepis hirsutula.
regulare Bak. JoB. **1875**. 200 = D. peregrina.
remotum Hew. Lond. JoB. **1**. 121 nota. 1842 — Ins. Norfolk. Timor.
remotum Hk. Brit. Ferns t. 22. 1861.
remotum Clarke, Tr. Linn. Soc. II. Bot. **1**. 523. 1880 = D. spinulosa var.
repandum Moore, Ind. 102. 1858 = D. urophylla.
repens Bail. Handb. Queensl. Ferns 50. 1874 = Arthropteris obliterata.
repens Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 535 t. 8. 1899.
reptans Diels, Nat. Pfl. 1^4. 168. 1899.
resiniferum Hk. et Arn. Bot. Beech. Voy. 105. 1832 = D. gongylodes.
resino-foetidum Hk. sp. **4**. 105. 1862.
reticulatum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
retroflexum Desv. Prod. 256. 1827 = ? D. refracta.
retrorsum Sod. Rec. 51. 1883.
rhizophyllum Pr. Rel. Haenk. **1**. 31. 1825 = Polystichum r.
rhodolepis Clarke, Tr. Linn. Soc. II. Bot. **1**. 526 t. 72. 1880.
Richardsi Bak. Syn. 494. 1874.
Riedlianum Moore, Ind. 102. 1858 = Polystichum Riedleanum.
rigescens Sod. Cr. vasc. quit. 239. 1893.
rigidulum Bak. Syn. 496. 1874 = D. scalpturoides.
rigidum Desv. Prod. 261. 1827.
riparium Desv. Prod. 256. 1827 = D. tomentosa.
Rivoirei Diels, Nat. Pfl. 1^4. 179. 1899 = D. Poiteana.
Robertianum Prantl, Excursionsfl. v. Bayern 24. 1884.
Rodigasianum Moore, L'Ill. hort. **29**. 27 t. 442. 1882.
Rosthornii Diels, Engl. Jahrb. **29**. 190. 1900.
rotundatum Desv. Prod. 253. 1827.
rubidum Copeland, Polyp. Philipp. 27. 1905.
rubiginosum Hk. sp. **4**. 143. 1862.
rubrinerve Diels, Nat. Pfl. 1^4. 178. 1899.
rude Diels, Nat. Pfl. 1^4. 171. 1899.
rufescens Schrad. Gött. gel. Anz. **1824**. 869 = Nephrolepis hirsutula.
rufescens Diels, Nat. Pfl. 1^4. 175. 1899.
rufidulum Michx. Fl. bor. Am. **2**. 269. 1803 = Woodsia ilvensis.
rufum Desv. Prod. 257. 1827.
rugulosum Copeland, Polyp. Philipp. 26. 1905 = D. punctata *rugosula.
sagenioides Bak. Syn. 271. 1867.
sagittaefolium Moore, Ind. 103. 1858.
sakayense Zeiller, Bull. Soc. Fr. **32**. 75. 1885.
salicifolium Diels, Nat. Pfl. 1^4. 178. 1899.
Salvini Bak. Syn. 274. 1867.
Sancti-Gabrieli Bak. Fl. bras. 1^2. 469. 1870.
sanctum Bak. Syn. 267. 1867.
Sangnellii Bak. Ann. Bot. **5**. 318. 1891.
sarawakense Bak. Journ. Linn. Soc. **22**. 225. 1886 = D. rhodolepis.
savaiense Bak. Ann. Bot. **5**. 318. 1891.
scabriusculum Dav. Bot. Gaz. **21**. 256. 1896 (syn.) = D. Davenportii.
scabrosum Bak. Syn. 283. 1867.

scandicinum Bory, Bél. Voy. Bot. **2**. 63. 1833 = Athyrium s.
Schaffneri Fée, 8 mém. 108. 1857.
Schimperianum Hope, Journ. Bomb. Nat. Hist. Soc. **14**. 733. 1903.
schizotis Hk sp. **4**. 107. 1862 = D. patens.
Schkuhrii Link, Fil. sp. 109. 1841 = Nephrolepis pectinata.
sclerophyllum Pr. Tent. 81. 1836 = D. asplenioides.
scolopendrioides Hk. sp. **4**. 65. 1862.
securidiforme Diels, Nat. Pfl. **1**^{4}. 175. 1899.
Seemanni Bak. Ann. Bot. **5**. 320. 1891.
semibipinnatum Bak. Syn. 297. 1867 = Aspidium s.
semicordatum Pr. Rel. Haenk. **1**. 32. 1825 = Cyclopeltis s.
semihastatum Hk. sp. **4**. 87. 1862.
semilunatum Sod. Rec. 46. 1883.
serra Desv. Prod. 253. 1827.
Serra Hk. sp. **4**. 98. 1862 = D. patens *invisa.
serrato-dentatum Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 622 t. 10. 1899 = [D. filix mas*.
serratum Pr. Rel. Haenk. **1**. 34. 1825 = D. Haenkeana.
serratum Desv. Prod. 254. 1827 = ? D. gongylodes.
serratum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
serrulatum Jenm. Bull. Dep. Jam. **3**. 189. 1896.
setigerum Pr. Rel. Haenk. **1**. 37. 1825 = Polystichum aculeatum.
setigerum Bak. Syn. 284. 1827 = D. setigera et D. ornata.
setosum Bak. Syn. 274. 1867 = D. Millettii.
setosum Keys. Pol. Cyath. Hb. Bung. 48. 1873 = D. pseudotetragona.
setulosum Bak. JoB. **1890**. 265.
Sewellii Bak. Journ. Linn. Soc. **15**. 418. 1876.
Sheareri Bak. JoB. **1875**. 200.
Shepherdi Fée, Gen. 305. 1850-52 = ? D. decomposita.
Sherringiae Jenm. JoB. **1887**. 99 = Aspidium psammiosorum.
Sherringii Jenm. JoB. **1879**. 261 = D. Balbisii var.
shikokianum Mak. Bot. Mag. Tokyo **13**. 62. 1899.
Sieboldii Hk. Fil. Exot. t. 31. 1859.
siifolium Bak. Syn. 299. 1867 = Aspidium s.
sikkimense Clarke, Tr. Linn. Soc. II. Bot. **1**. 525. 1880.
Simonsii Bak. Syn. 504. 1874 = Aspidium S.
simplex Diels, Nat. Pfl. **1**^{4}. 177. 1899.
simplicifolium J. Sm. JoB. **3**. 411. 1841 = D. canescens*.
simulans Bak. Syn. ed. II. 300. 1874 = { Aspidium Thwaitesii. Aspidium cicutarium.
simulans Bak. JoB. **1888**. 325 = D. simillima.
simulans Bak. JoB. **1890**. 106 = D. conterminoides.
simulatum Dav.; Diels, Nat. Pfl. **1**^{4}. 172. 1899.
singalanense Bak. JoB. **1880**. 212.
singaporianum Bak. Syn. 296. 1867 = Aspidium s.
Sintenesii Bak. Ann. Bot. **5**. 327. 1891 = D. Sintenisii.
Skinneri Moore, Ind. 104. 1858.
Sloanei Pr. Rel. Haenk. **1**. 33. 1825 = Polystichum triangulum.
Sloanei Bak. Syn. ed. II 263. 1874 = D. patens *invisa.
Smithianum Pr. Epim. 46. 1849 = D. hispidula.
Sodiroi Bak. JoB. **1877**. 164 = Aspidium S.

sophoroides Desv. Prod. 256. 1827.
sorbifolium Pr. Rel. Haenk. **1**. 31. 1825 = ? Polybotrya polyphylla.
sorbifolium Hier. Engl. Jahrb. **34**. 449. 1904 = D. reticulata*.
sparsum Desv. Prod. Fl. Nepal. 6. 1825.
sparsum var. gracilis Bak. Syn. 498. 1874 = Polystichum Thomsoni.
sparsum var. membranacea Bak. Syn. 498. 1874 = Athyrium macrocarpum.
spectabile Hk. sp. **4**. 115. 1862 = D. syrmatica.
spectabile Clarke, Tr. Linn. Soc. II. Bot. **1**. 526. 1880 = D. Hendersoni.
spectabile Hier. Engl. Jahrb. **34**. 446. 1904 = D. subincisa var.
Spekei Bak. Syn. 263. 1867.
sphaerocarpum Hk. sp. **4**. 139. 1862 = D. athyrioides.
spinulosum Strempel, Fil. Berol. Syn. 30. 1822.
splendens Desv. Prod. 253. 1827 = Nephrolepis biserrata.
splendens Hk. sp. **4**. 126. 1862 (excl. var.).
splendens var. angustifrons Hk. sp. **4**. 126. 1862 = D. angustifrons.
sporodicum (»Fée«) Diels, Nat. Pfl. 1^4. 168. 1899 (nomen) — ?
Sprengelii Hk. sp. **4**. 94. 1862 = D. Balbisii.
Sprucei Bak. Syn. 269. 1867.
squamigerum Hk. et Arn. Bot. Beech. Voy. 106. 1832.
squamigerum (Mett.) Rosenst. Hedwigia **43**. 224. 1904 (nomen) — Brasilia austr. (Aspidium Mett. HB. 265 sub N. vestito). (= D. deflexa?).
squamisetum Hk. sp. **4**. 140 t. 268. 1862.
squamosissimum Sod. Cr. vasc. quit. 256. 1893.
squamulosum Hk.fil. Fl. N. Zeal. **2**. 39. 1854 = D. thelypteris*.
Stegnogramme Diels, Nat. Pfl. 1^4. 181. 1899.
stenophyllum Sod. Rec. 44. 1883.
stenophyllum Bak. JoB. **1884**. 363 = D. Harrisonii.
stenophyllum Bak. Journ. Linn. Soc. **22**. 227 t. 11. 1886 = Aspidium Hosei.
stenopteris Eat. Amer. Journ. Sci. II. **27**. 199. 1859 = D. incisa.
stenopteron Bak. JoB. **1890**. 265 = Aspidium Balansae.
stenopterum Diels, Nat. Pfl. 1^4. 170. 1899 = Dryopteris Dielsii.
stipellatum Moore, Ind. 105. 1858.
stipitatum Bak. Syn. 278. 1867.
stipulare Desv. Prod. 256. 1827 = D. patens*.
stramineum Sod. Rec. 43. 1883; Cr. vasc. quit. 239. 1893 = D. opposita.
strigilosum Dav. Bot. Gaz. 21. 257 (syn.).
strigosum Desv. Prod. 256. 1827.
strigosum Jenm. Bull. Dep. Jam. **3**. 141. 1896 = D. scolopendrioides.
subbiauritum Hk. sp. **4**. 85. 1862 = D. orientalis var.
subconfluens Clarke, Tr. Linn. Soc. II. Bot. **1**. 536. 1880 = Aspidium s.
subcrenulatum Bak. Journ. Linn. Soc. **16**. 202. 1877.
subcuneatum Bak. Fl. bras. 1^2. 487. 1870.
subdigitatum Bak. Journ. Linn. Soc. **24**. 259. 1887 = Aspidium psilopodum.
subdimorphum Christ, Verh. Nat. Ges. Basel **11**. 16. 1894 = D. canescens.
subfuscum Bak. Syn. 267. 1867.
subglabrum Sod. Cr. vasc. quit. 259. 1893.
subglandulosum Bak. Syn. 285. 1867 = D. Helliana.
subincisum Christ, Bull. Boiss. II. **3**. 612. 1903.
subintegrum Sod. Rec. 54. 1883.
subjunctum Bak. Ann. Bot. **5**. 321. 1891.

subobliquatum Bak. Syn. 261. 1867.
subpectinatum Bedd. Handb. Suppl. 66. 1892 = N. otaria.
subpedatum Harr. Journ. Linn. Soc. **16**. 30. 1877 = Aspidium s.
subquinquefidum Hk. sp. **4**. 130. 1862 = D. protensa et D. securidiformis.
subsericeum Bak. Syn. 281. 1867.
subtriangulare Hope, JoB. **1890**. 327.
subtriphyllum Bak. Syn. 296. 1867 = Aspidium s.
subtripinnatum Bak. Syn. ed. I. 455. 1868.
sulcatum Kiærsk. Cat. Fil. hb. ht. Haun. 64. 1874 = D. crinita.
supinum Sod. Cr. vasc. quit. 241. 1893.
syrmaticum Desv. Prod. 259. 1827.
Tarapotense Hk. sp. **4**. 107. 1862 = D. submarginalis.
tectum Bedd. Handb. Suppl. 79. 1892 = D. didymosora.
tenebricum Jenm. JoB. **1882**. 326.
tenericaule Hk. sp. **4**. 142. 1862 = D. setigera.
tenericaule Hk. sp. **4**. t. 269. 1862 = D. ornata.
tenerum R.Br. Prod. Fl. N. Holl. 149. 1810.
tenue Michx. Fl. bor. Am. **2**. 269. 1803 = Cystopteris fragilis.
tenuifolium Hk. sp. **4**. 144. 1862 = D. tenuifrons.
terminans J. Sm. Bot. Mag. **72** Comp. 32. 1846 = D. pteroides.
ternatum Bak. Syn. 296. 1867 = Aspidium t.
tetragonum Pr. Rel. Haenk. **1**. 35. 1825 = D. pseudotetragona.
tetragonum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
tetragonum Boj. Hort. Maur. 393. 1837 = D. prolixa.
Thelypteris Strempel, Fil. Berol. Syn. 32. 1822.
thelypteroides Michx. Fl. bor. Am. **2**. 267. 1803 = D. noveboracensis.
thelypteroides Hk. sp. **4**. 95. 1862 = D. tomentosa.
Thwaitesii Bak. Syn. 277. 1867 = D. deparioides var.
tibeticum Bak. Ann. Bot. **5**. 318. 1891 = D. thibetica.
tijuccanum Diels, Nat. Pfl. **1**⁴. 170. 1899.
timoriense Desv. Prod. 253. 1827 = Nephrolepis biserrata.
tokyoense Mak. Bot. Mag. Tokyo **12**. 87. 1898.
tomentosum Desv. Prod. 256. 1827.
Totta Diels, Nat. Pfl. **1**⁴. 170. 1899 = D. africana.
transversarium Brack. Expl. Exp. **16**. 187. 1854 = ? D. pennigera.
trapezoides Pr. Rel. Haenk. **1**. 37. 1825 = Polystichum mohrioides.
Trianae Bak. Syn. 286. 1867.
tricholepis Bak. ap. Hemsl. Biol. Cent. Amer. Bot. **3**. 651. 1885.
trichophlebium Bak. Journ. Linn. Soc. **16**. 201. 1877.
trichophorum Bak. Syn. 265. 1867.
trifoliatum Bory, Bél. Voy. Bot. **2**. 59. 1833 = Aspidium pica.
tripartitum Bak. JoB. **1879**. 296 = Aspidium t.
triphyllum Diels, Nat. Pfl. **1**⁴. 178. 1899.
triste Hk. sp. **4**. 104. 1862.
truncatum Pr. Tent. 81. 1836.
tuberculatum Ces. Rend. Ac. Napoli **16**. 26, 29. 1877.
tuberosum Desv. Prod. 252. 1827 = Nephrolepis cordifolia.
Tuerckheimii Donn. Smith. Bot. Gaz. **12**. 133 t. 11. 1887.
tylodes (Bak. Syn. 268. 1867) = D. ochthodes.
uliginosum (Bak. Syn. 273. 1867) = D. cristata × spinulosa.

undulatum Bak. Syn. 276. 1867 = D. sparsa*.
unidentatum Diels, Nat. Pfl. **1**[4]. 174. 1899.
unifurcatum Bak. JoB. **1888**. 228.
unitum R.Br. Prod. Fl. N. Holl. 148. 1810 = D. gongylodes.
unitum Bory, Bél. Voy. Bot. **2**. 61. 1833.
Urbani Sod. Cr. vasc. quit. 638. 1893.
urophyllum Keys. Pol. Cyath. Hb. Bung. 49. 1873.
usitatum Jenm. JoB. **1879**. 261.
valdepilosum Bak. JoB. **1881**. 204.
variabile Hk. sp. **4**. 140. 1862 = D. Vogelii.
varians Fée, 11 mém. 88 t. 24 f. 2. 1866.
variolarium Desv. Prod. 255. 1827 = D. orientalis.
variolosum Bak. Syn. 298. 1867 = Aspidium v.
varium Pr. Rel. Haenk. **1**. 36. 1825 = Sagenia v.
varium Desv. Prod. 259. 1827 = Polystichum v.
vastum Bak. Syn. 296. 1867 = Aspidium v.
vastum Hier. Engl. Jahrb. **34**. 446. 1904 = D. subincisa*.
velatum Hk. sp. **4**. 101 t. 247. 1862.
velleum Desv. Prod. 261. 1827.
velutinum Hk.fil. Fl. N. Zeal. **2**. 39 t. 80. 1854.
venulosum Desv. Prod. 255. 1827 = D. gongylodes.
venulosum Hk. sp. **4**. 71. 1862 = D. elata.
venustum Hew.; Moore, Gard. Chr. **1855**. 677 c. fig.
Vescoi Drake, Fl. Polyn. franç. 301. 1893.
vestitum Bak. Syn. 265. 1867 = D. deflexa.
Vieillardii Bak. Syn. 282. 1867.
vile Bak. Syn. 499. 1874.
villosum Pr. Rel. Haenk. **1**. 38. 1825.
violascens Fée, Gen. 305. 1850-52 = D. parasitica var.
viridescens Bak. Syn. 275. 1867.
? *viscidum* Watson, Proc. Amer. Ac. **26**. 163. 1891 — Ascension.
viscosum Bak. Syn. 264. 1867.
Vitis Racib. Bull. int. Ac. Cracovie **1902**. 60 = Aspidium v.
Vogelii Hk. sp. **4**. 130. 1862.
vulcanicum Bak. Ann. Bot. **8**. 127. 1894.
Wakefieldii Bak. Ann. Bot. **5**. 326. 1891.
Walkerae Bak. Syn. 493. 1874.
Warburgii Diels in Schum. et Laut. Fl. deut. Südsee 112. 1901 = D. debilis.
Wardii Bak. Syn. 500. 1874.
Webbianum Hk. sp. **4**. 85. 1862 = D. orientalis.
Welwitschii Bak. Syn. 274. 1867.
Wightii Clarke. Tr. Linn. Soc. II. Bot. **1**. 538 t. 76. 1880 = Aspidium polymorphum.
Wilsoni Bak. Ann. Bot. **5**. 321. 1891.
Wrightii Hk. sp. **4**. 64 t. 239. 1862.
Wrightii Diels, Nat. Pfl. **1**[4]. 168. 1899 = D. Sauvallei.
xanthotrichium Sod. Rec. 52. 1883 = D. effusa var.
xylodes Hope. Journ. Bomb. Nat. Hist. Soc. **14**. 724. 1903 = D. ochthodes var.
Zambesiacum Bak. Ann. Bot. **5**. 318. 1891.
Zeylanicum Fée, 10 mém. 42. 1865 — Ceylon.
Zollingerianum Bak. Syn. 298. 1867 = Aspidium brachiatum.

NEPHROLEPIS Schott, Gen. Fil. t. 3. 1834 — (Polypodiaceae Gen. 27).

abrupta (Bory) Mett. Fil. Lips. 99. 1856. Chr. 291. NPfl. 208 — Madagascar. Ins. Mascarenae. — Dicksonia Bory, Voy. 2. 187 t. 30. 1804. HB. 52; Leptopleuria Pr. 1836.

acuminata (Houtt.) Kuhn, Ann. Lugd. Bat. 4. 286. 1869 — Perak. Malesia. Ophioglossum Houtt. Nat. Hist. 14. 94. 1783; ed. germ. Pfl. Syst. 13¹. 53 t. 94 f. 3. 1786. Aspidium davallioides Sw. Schrad. Journ. 1800². 33. 1801; Nephrodium? Desv. 1827; Nephrolepis Kze. 1846. HB. 302. Chr. 289. [NPfl. 207.

acuminata Pr. Tent. 79. 1836 = N. biserrata.

acuta Pr. Tent. 79. 1836 = N. biserrata.

acutangula Pr. Tent. 79. 1836; Epim. 43. 1849 = N. hirsutula.

acutifolia (Desv.) Christ, Verh. Nat. Ges. Basel 11. 243. 1895. Chr. 291. NPfl. 206 — Birma-Malesia-Australia. Africa trop.
Lindsaya Desv. Prod. 312. 1827. Isoloma lanuginosa J. Sm. 1842; Lindsaya Hk. 1846. HB. 109.

altescandens Bak. Syn. 301. 1867 = Arthropteris.

altescandens Bail. Handb. Ferns Queensl. 51. 1874 = Arthropteris obliterata.

barbata Copeland in Perkins, Fragm. 178. 1905 — Ins. Philipp. (Mindanao).

biaurita Pr. Epim. 43. 1849 — Mauritius. — Lepidonevron Fée 1850-52.

bidentata Pr. Tent. 79. 1836 — Ins. Mariannae. — Nephrodium Pr. Rel. Haenk. 1. 32. 1825; Aspidium Spr. 1827; Lepidonevron Fée 1850-52.

biserrata (Sw.) Schott, Gen. Fil. ad t. 3 1834 — Trop.
Aspidium Sw. Schrad. Journ. 1800². 32. 1801; Nephrodium Pr. 1825; Hypopeltis Bory 1833; Lepidonevron Fée 1850-52. Aspidium acutum Schkuhr 1806; Nephrodium Pr. 1825; Nephrolepis Pr. 1836. HB. 301. Chr. 289. NPfl. 207 — Aspidium splendens Willd. 1810; Nephrodium Desv. 1827; Nephrolepis Pr. 1836.
— Aspidium gibbosum Willd. 1810; Nephrodium Gaud. 1828; Nephrolepis Pr. 1836 — Aspidium acuminatum Willd. 1810; Nephrodium Pr. 1825; Nephrolepis Pr. 1836; Lepidonevron Fée 1850-52.
— Aspidium paraënse Willd. 1810; Nephrolepis Pr. 1836 — Polypodium salignum Sol. msc.; Nephrolepis Carr. 1873.
—* Polypodium punctulatum Poir. 1804; Aspidium Sw. 1806; Nephrodium Desv. 1827; Nephrolepis Pr. 1836; Lepidonevron Fée 1850-52.

celebica Zoll. Syst. Verz. 44, 49. 1854 — Celebes.

cordifolia (L.) Pr. Tent. 79. 1836. HB. 300. Chr. 288. NPfl. 206 — Trop.-Japonia. N. Zealand. — Polypodium L. sp. 2. 1089. 1753. Aspidium Sw. 1801. A. tuberosum Bory; Willd. 1810; Nephrodium Desv. 1827; Nephrolepis Pr. 1836. — Aspidium undulatum Afz.; Sw. 1801; Nephrolepis J. Sm. 1846. — Aspidium pendulum Raddi 1819; Nephrodium Desv. 1827; Nephrolepis J. Sm. 1841 — N. pluma Moore 1878 (Madagascar).

crenata Fée, 1 mém. 14 t. 1 f. 19. 1844 = N. exaltata.

cultrifolia Pr. Tent. 79. 1836; Epim. 44. 1849 = N. exaltata.

davallioides Kze. Bot. Zeit. 1846. 460 = N. acuminata.

delicatula Dcne. (HB. 301. 1867) = N. cordifolia.

dentata Goldm. Nova Acta 19. Suppl. I. 463. 1843 = ? N. exaltata.

depauperata De Vriese, Ned. Kr. Arch. 1. 9. 1846 = N. cordifolia.

dicksonioides Christ, Verh. Nat. Ges. Basel 11. 241 t. 3 f. 8-10. 1895. Ann Jard. Buit. 15. 140 t. 15 f. 18. 1897. Chr. 291. NPfl. 208 — Celebes.
Dicksonia nephrolepioides Christ 1895.

Duffii Moore, Gard. Chr. n. s. **9**. 622 f. 113. 1878 — Malesia. Austr. bor.
ensifolia Pr. Tent. 79. 1836 = N. biserrata.
exaltata (L.) Schott, Gen. Fil. t. 3. 1834. HB. 301. Chr. 289. NPfl. 207 — Trop. Polypodium L. Syst. Nat. ed. X. **2**. 1326. 1759; Aspidium Sw. 1801; Nephrodium R.Br. 1810.
falciformis J. Sm. Ferns br. and for 287. 1866 = N. exaltata.
flexuosa Col. Tr. N. Zeal. Inst. **20**. 231. 1888 = N. cordifolia.
floccigera (Bl.) Moore, Ind. 92. 1858. HB. 302. Chr. 291. NPfl. 207 — Java, Celebes. — Aspidium Bl. Enum. 147. 1828.
Fosteri ht. Hill; Gard. Chr. III. **34**. 64. 1903 — Hort.
Gaimardiana Pr. Tent. 79. 1836 = Humata.
gibbosa Pr. Tent. 79. 1836 = N. biserrata.
Hestoni ht.; Gard. Chr. III. **34**. 384. 1903 — Hort.
hirsutula (Forst.) Pr. Tent. 79. 1836 — Trop.
Polypodium Forst. Prod. 81. 1786; Aspidium Sw. 1801; Polystichum Bernh. 1802; Nephrodium Pr. 1825; Lepidonevron Fée 1850-52 — Nephrodium rufescens Schrad. 1824; Aspidium Kze. 1839; Lepidonevron Fée 1850-52; Nephrolepis Wawra 1866.
imbricata Pr. Tent. 79. 1836 = N. cordifolia.
intermedia Fée, 9 mém. 32. 1857 = N. cordifolia.
intermedia Sod. Rec. 57. 1883 = N. exaltata.
intramarginalis Kze. Linn. **23**. 268, 312. 1850 = ? N. exaltata.
Kuroiwae Mak. Bot. Mag. Tokyo **9**. 6. 1895 — Liu Kiu.
lanceolata Goldm. Nova Acta **19**. Suppl. I. 462. 1843 — Manila.
Lauterbachii Christ, Bull. Boiss. II. **1**. 456. 1901 — N. Guinea.
N. cordifolia var. Lauterbachii Christ in Schum. et Laut. Fl. deut. Südsee [118. 1901.
lindsayae Christ, Bull. Boiss. **6**. 837. 1898 — Sumatra.
lunulata Keys. Pol. Cyath. Hb. Bung. 40. 1873 = Didymochlaena truncatula.
macrophylla Pr. Epim. 43. 1849 — Luzon.
? mauritiana Moore, Ind. 96. 1858 = N. biserrata.
Mayii ht. May; Gard. Mag. **1903**. 688, 705 c. fig. — Hort.
neglecta Kze. Linn. **13**. 148. 1839 = N. rivularis.
obliterata J. Sm. JoB. **4**. 197. 1841 = Arthropteris.
obtusifolia Pr. Tent. 80. 1836 = N. acuminata.
occidentalis Kze. Linn. **18**. 343. 1844 = N. pectinata.
paleacea Jungh. et de Vriese. Ned. Kr. Arch. **1**. 4. 1846 = N. cordifolia.
paraënsis Pr. Tent. 79. 1836 = N. biserrata.
pectinata (Willd.) Schott, Gen. Fil. ad t. 3. 1834. Chr. 289. NPfl. 207 — Amer. trop. — Aspidium Willd. sp. **5**. 223. 1810; Nephrodium Link 1833. Aspidium trapezoides Schkuhr 1806 (non Sw. 1801).
pendula J. Sm. JoB. **4**. 197. 1841 = N. cordifolia var.
Piersoni ht.; Gard. Mag. **1902**. 230 — Hort.
pilosa Pr. Tent 79. 1836 = N. hirsutula.
platyotis Kze. Linn. **23**. 268, 312. 1850 = N. biserrata.
Pluma Moore, Gard. Chron. n. s. **9**. 588 f. 68. 1878 = N. cordifolia var.
punctulata Pr. Tent. 79 t. 2 f. 14. 1836 = N. biserrata*.
radicans (Burm.) Kuhn, Ann. Lugd. Bat. **4**. 285. 1869 — Asia trop.
Polypodium Burm. Fl. Ind. 233 t. 66 f. 3. 1768. Nephrolepis volubilis J. Sm. 1841 (nomen); Pr. Epim. 44. 1849. Chr. 288. NPfl. 207. (HB 301);

Lepidonevron Fée 1850-52 — ? Aspidium obtusifolium Willd. 1810; Nephrodium Pr. 1825; Nephrolepis Pr. 1836.
ramosa Moore, Ind. 102. 1858 = Arthropteris obliterata.
repens Brack. Expl. Exp. **16**. 209. 1854 = Arthropteris obliterata.
rhizodes Kze. Bot. Zeit. **1848**. 236 = N. cordifolia.
rivularis (Vahl) Mett.; Krug, Engl. Jahrb. **24**. 122. 1897 — Amer. trop.
Polypodium Vahl, Ecl. Amer. **3**. 51. 1807. Aspidium sesquipedale Willd. 1810; Nephrolepis Pr. 1836; Lepidonevron Fée 1869 — Nephrolepis neglecta Kze. 1839 — N. valida Kze. 1848. 1850.
rufescens Wawra, Maxim. Reise 200 t. 101. 1866 = N. hirsutula.
saligna Carr. in Seem. Fl. Vit. 361. 1873 = N. biserrata.
Schkuhrii Fée, Gen. 319. 1850-52 = N. pectinata.
sesquipedalis Pr. Tent. 79. 1836 = N. rivularis.
Sieberi Pr. Tent. 79. 1836 = N. biserrata.
splendens Pr. Tent. 79. 1836 = N. biserrata.
subcordata Pr. Tent. 79. 1836; Epim. 43. 1849 = N. biserrata.
trichomanoides J. Sm. JoB. **3**. 413. 1841 = Arthropteris obliterata.
tuberosa Pr. Tent. 79. 1836 = N. cordifolia.
undulata J. Sm. Bot. Mag. **72**. Comp. 35. 1846 = N. cordifolia var.
valida Kze. Linn. **21**. 229. 1848 (nomen); Linn. **23**. 268. 1850 = N. rivularis.
volubilis J. Sm. JoB. **3**. 413. 1841 = N. radicans.
Wallichiana Pr. Tent. 79. 1836 = N. hirsutula.
Westoni ht. Craig; Gard. Chr. III. **34**. 309. 1903 — Hort.
Zollingeriana de Vriese, Ned. Kr. Arch. **1**. 10. 1846 = N. biserrata.

NEUROCALLIS Fée, 2 mém. Hist. Acrostich. 19, 89. 1845.
aureo-nitens Moore, Ind. XIX. 1857 = Trachypteris pinnata.
bipinnata Moore, Ind. XIX. 1857 = Leptochilus lomarioides.
lomarioides Pr. Epim. 177. 1849 = Leptochilus.
pinnata Moore, Ind. XIX. 1857 = Leptochilus lomarioides.
polyphylla Moore, Ind. XIX. 1857 = Polybotrya.
praestantissima Fée, Hist. Acrost. 89 t. 52. 1845 = Acrostichum.
Requiniana Fée, Hist. Acrost. 90. 1845 = Platytaenia.
rivularis Moore, Ind. 276. 1861 = Leptochilus.
scandens Moore, Ind. XIX. 1857 = Leptochilus guianensis.

NEURODIUM Fée, 3 mémoire 11, 28. 1851-52 = **Paltonium.**
lanceolatum Fée, 3 mém. 28. 1851-52.
sinense Christ, Bull. Boiss. **6**. 880. 1898.

NEUROGRAMMA (Pr.) Link, Fil. spec. 138. 1841.
Gymnogramma § Neurogramma Presl, Tent. Pterid. 218. 1836.
alismaefolia Christ, Farnkr. 64. 1897 = Syngramma.
calomelanos Diels, Nat. Pfl. **1**⁴. 264. 1899 = Ceropteris.
cartilagidens Christ, Farnkr. 63. 1897 = Syngramma.
Delavayi Diels, Nat. Pfl. **1**⁴. 262. 1899 = Gymnopteris.
elaphoglossoides Christ, Farnkr. 63. 1897 = Syngramma.
ferruginea Diels, Nat. Pfl. **1**⁴. 263. 1899 = Gymnopteris.
fraxinea Christ, Farnkr. 63. 1897 = Coniogramme.
hispida Diels, Nat. Pfl. **1**⁴. 264. 1899 = Gymnopteris.
marginata Christ, Farnkr. 63. 1897 = Syngramma.
Muelleri Diels, Nat. Pfl. **1**⁴. 262. 1899 = Gymnopteris.
pedata Link, Fil. sp. 139. 1841 = Gymnopteris.

peruviana Hier. Engl. Jahrb. 34. 482. 1904 = Ceropteris.
quinata Christ, Farnkr. 64. 1897 = Syngramma.
Reynoldsii Diels, Nat. Pfl. 1^4. 262. 1899 = Gymnopteris.
rufa Link, Fil. sp. 138. 1841 = Gymnopteris.
scandens Fée. Cr. vasc. Br. **1**. 263. 1869; **2**. 39 t. 92. 1872-73 = Gymnogramma.
tartareum Diels, Nat. Pfl. 1^4. 264. 1899 = Ceropteris.
tomentosa Link, Fil. sp. 139. 1841 = Gymnopteris.
triangularis Diels, Nat. Pfl. 1^4. 264. 1899 = Ceropteris.
vestita Diels, Nat. Pfl. 1^4. 262. 1899 = Gymnopteris.
xerophila Diels, Nat. Pfl. 1^4. 263. 1899 = Gymnogramma.

NEUROMANES Trevisan, Atti Istituto Veneto II. **2**. 163. 1851 = **Trichomanes.**
abruptum v.d.B. Ned. kr. Arch. **4**. 350. 1859 = T. Hostmannianum.
affine Trev. Atti Ist. Veneto II. **2**. 163. 1851 = T. Hostmannianum.
Hedwigii v.d.B. Ned. kr. Arch. **4**. 348. 1859 = T. pinnatum.
immersum v.d.B. Ned. kr. Arch. **4**. 349. 1859 = T. pinnatum var.
Kaulfussii v.d.B. Ned. kr. Arch. **4**. 348. 1859 = T. pennatum.
Schomburgkianum v.d.B. Ned. kr. Arch. 5^2. 139. 1861.
Vittaria v.d.B. Ned. kr. Arch. **4**. 349. 1859.

NEURONIA Don, Prod. Fl. Nepal. 6. 1825 = **Oleandra.**
asplenioides Don, Prod. 7. 1825 = O. Wallichii.

NEUROPHYLLUM Presl, Hymenoph. 18. 1843 = **Trichomanes.**
abruptum Fée, 1 mém. 14 t. 1 f. 5. 1844 = T. Hostmannianum.
Hedwigianum Fée, 10 mém. 151. 1865 } T. pinnatum.
Hedwigii Fée, 11 mém. 104. 1866. }
Hostmannianum Kl. Linn. **18**. 532. 1844.
pennatum Pr. Hym. 19. 1843.
pinnatum Pr. Hym. 19 t. 4 f. C. 1843.
rhizophyllum Fée, 11 mém. 105. 1866 = T. pinnatum.
thecaphyllum Fée, 11 mém. 104 t. 28 f. 1. 1866 = T. pinnatum.
Vittaria Pr. Hym. 19. 1843.

NEUROPLATYCEROS (Endl.) Fée, 2 mémoire: Hist. d'Acrost. 24, 102. 1845 = **Platycerium.**
Acrostichum § Neuroplatyceros Endl. Gen. pl. **1**. 59. 1836.
aethiopicus Fée, Acrost. 103 t. 64. 1845 = P. stemaria.
alcicornis Fée, Acrost. 102. 1845 = P. bifurcata.
biformis Fée, Acrost. 104. 1845 = P. coronarium.
grandis Fée, Acrost. 103. 1845.

NEUROPTERIS Desvaux, Prodrome 292. 1827 = **Saccoloma.**
elegans Desv. Prod. 293 t. 8 f. 2. 1827.

?**NEUROSORIA** Mett.; Kuhn, Bot. Zeit. **1869**. 437 — (Polypodiaceae Gen. **76 a**).
pteroides (R.Br.) Mett.; Kuhn, Bot. Zeit. **1869**. 438 — Australia trop.
Acrostichum R.Br. Prod. Fl. N. Holl. 144. 1810. (HB. 424, 524); Phorolobus Desv. 1827.

NEUROSORUS Trevisan, Atti Ist. Veneto II. **2**. 168. 1851 = **Coniogramme.**

NIPHIDIUM J. Smith, Hist. Fil. 99. 1875 = **Cyclophorus.**
americanum J. Sm. Hist. Fil. 99. 1875.

NIPHOBOLUS Kaulfuss, Enum. Fil. 124. 1824 = **Cyclophorus.**
abbreviatus Zoll. Nat. en Gen. Arch. **1**. 397. 1844.
acrocarpus Christ et Gies. Bull. Ac. Géogr. Bot. Mans **1902**. 221.
acrostichoides Klf. Enum. 127. 1824 = C. confluens.

adnascens Klf. Enum. 124. 1824
africanus Kze. Linn. **10**. 501. 1839.
albicans Bl. Enum. 107. 1828.
americanus Diels, Nat. Pfl. 1⁴. 325. 1899.
anetioides Christ, Ann. Jard. Buit. II. **4**. 38. 1904.
angustatus Spr. Syst. **4**. 44. 1827.
angustissimus Gies.; Diels, Nat. Pfl. 1⁴. 326. 1899 = C. taeniodes.
annamensis Christ, Journ. de Bot. **19**. 74. 1905.
antillarum Spr. Syst. **4**. 44. 1827 = Polypodium piloselloides.
assimilis Diels, Engl. Jahrb. **29**. 207. 1900.
atropunctatus Keys. Pol. Cyath. Hb. Bung. 39. 1873 = Polypodium lineare.
Beddomeanus Gies. Niph. 101. 1901.
bicolor Klf. Enum. 128. 1824 = C. serpens.
Blumeanus Kze. Bot. Zeit. **1848**. 120.
Bonii Christ; Gies. Niph. 120. 1901.
Boothii Bedd. Ferns br. Ind. t. 258. 1868.
calvatus Diels, Engl. Jahrb. **29**. 207. 1900.
carnosus Bl. Enum. 105. 1828 = C. adnascens.
caudatus Klf. Enum. 127. 1824 = C. adnascens.
Cavalerianus Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 107.
ceylanicus Gies. Niph. 216 f. 19. 1901.
Chamissoanus Pr. Tent. 127. 1836 = C. adnascens.
Christii Gies. Niph. 140. 1901.
confluens Spr. Syst. **4**. 44. 1827.
confluens Bail. Handb. Ferns Queensl. 43. 1874 = C. rupestris.
costatus Pr. Tent. 202. 1836 = C. Beddomeanus.
cuneatus Kuhn, Bot. Zeit. **1868**. 40.
Davidii Gies.; Diels, Nat. Pfl. 1⁴. 325. 1899 = C. pekinensis.
detergibilis Bedd. Ferns br. Ind. t. 162. 1866 = C. flocculosus.
distichocarpus Gies.; Diels, Nat. Pfl. 1⁴. 325. 1899.
Drakeanus Diels, Engl. Jahrb. **29**. 206. 1900.
elongatus Bl. Enum. 105. 1828 = C. adnascens.
fissus Bl. Enum. 106. 1828 = C. acrostichoides.
fissus Bedd. Ferns br. Ind. Correct. II. 1870 = C. porosus.
flocciger Bl. Enum. 107. 1828.
flocculosus Spr. Syst. **4**. 45. 1827.
Gardneri Kze.; J. Sm. Cat. cult. ferns 12. 1857.
Giesenhagenii Christ, Ann. Cons. Jard. Bot. Genève **7-8**. 330. 1904.
glaber Klf. Enum. 127. 1824 = C. albicans.
gralla Gies. Niph. 128. 1901.
hastatus Kze. Bot. Zeit. **1848**. 505.
Heteractis (es) J. Sm. Ferns br. and for. ed. II. 296. 1877.
heterophyllus Spr. Syst. **4**. 44. 1827 = C. spissus.
inaequalis Christ, Bull. Soc Fr. **52** Mém. I. 25. 1905.
Koenigii Bl. Fl. Javae, Fil. 50 (nota). 1828 — Ceylon.
laevis J. Sm. Bedd. Ferns br. Ind. t. 161. 1866.
lanceolatus Keys. Pol. Cyath. Hb. Bung. 39. 1873 = Polypodium l.
lanuginosus Gies. Niph. 190. 1901.
Lauterbachii Christ in Schum. et Laut. Fl. deut. Südsee 142. 1901.

linearifolius Hk. 2 Cent. t. 58. 1861.
linearis Klf. Wes d. Farrnkr. 113. 1827 (nomen) — ?
linearis Keys. Pol. Cyath. Hb. Bung. 39. 1873 = Polypodium l.
Lingua Spr. Syst. **4**. 45. 1827.
Lingua Bedd. Ferns S. Ind. t. 240. 1863-65 = C. pannosus.
Lingua Bedd. Ferns br. Ind. Suppl. t. 385. 1876 = C heteractis.
lingulatus Klf. Enum. 125. 1824 — Nova Brittania.
longifolius Spr. Syst. **4**. 45. 1827.
longifolius Keys. Pol. Cyath. Hb. Bung. 38. 1873 = Polypodium revolutum.
lycopodioides Keys. Pol. Cyath. Hb. Bung. 38. 1873 = Polypodium l.
macrocarpus Hk. et Arn. Bot. Beech. Voy. 74 t. 18. 1832 = C. angustatus.
Mannii Gies. Niph. 107. 1901 = C. porosus.
Martini Christ, Bull. Soc. Fr. **52** Mém. I. 23. 1905.
mollis Kze. Bot. Zeit. **1848**. 121.
?neglectus Fée, Gen. 263. 1850-52 = Polypodium n.
nudus Gies. Niph. 149. 1901.
nummularifolius Bedd. Ferns S. Ind. t. 186. 1863-65 = Drymoglossum heterophyllum.
nummularifolius J. Sm. JoB. **3**. 396. 1841.
obovatus Kze. Bot. Zeit. **1848**. 120 = C. nummularifolius.
pannosus Bedd. Ferns br. Ind. Suppl. 22. 1876.
penangianus Hk. Ic. pl. t. 203. 1840.
percussus Keys. Pol. Cyath. Hb. Bung. 38. 1873 = Polypodium p.
pertusus Spr. Syst. **4**. 44. 1827 = C. adnascens.
petiolosus Diels. Engl. Jahrb. **29**. 207. 1900.
piloselloides Keys. Pol. Cyath. Hb. Bung. 38. 1873 = Polypodium p.
polycarpus Spr. Syst. **4**. 45. 1827 = ? Polypodium punctatum.
polydactylon Gies.; Diels, Nat. Pfl. 1^4. 325. 1899.
porosus Pr. Tent. 202. 1836.
princeps Gies.; Diels, Nat. Pfl. 1^4. 325. 1899.
puberulus Bl. Enum. 108. 1828 = C. acrostichoides.
Rasamalae Gies. Niph. 187. 1901.
rotundifolius Fée, Gen. 262. 1850-52 = C. nummularifolius.
rupestris Spr. Syst. **4**. 44. 1827.
salicifolius Keys. Pol. Cyath. Hb Bung. 38. 1873 = Polypodium s.
Samarensis Fée, Gen. 263. 1850-62.
Sarasinorum Gies. Niph. 162. 1901.
Schimperianus Gies.; Diels, Nat. Pfl. 1^4. 325. 1899.
Schmidianus Kze. Bot. Zeit. **1848**. 121 = C. sticticus.
Schraderi Keys. Pol. Cyath. Hb. Bung. 39. 1873 = Polypodium S.
Scytopteris Fée, Gen. 262. 1850-52 = C. confluens.
serpens Endl. Prod. Fl. Norf. 8. 1883 (?); Carr. in Seem. Fl. Vit. 367. 1873.
sesquipedalis Keys. Pol. Cyath. Hb. Bung. 39. 1873 = Polypodium excavatum.
Sheareri Diels, Nat. Pfl. 1^4. 325. 1899.
sinensis Loud. Hort. brit. Suppl. —. 1839 = C. lingua.
Smithianus Moore, Ind. LXXVI. 1857 = C. acrostichoides.
spathulifer Bory. Bél. Voy. Bot. **2**. 24. 1833 = ? C. adnascens.
sphaerocephalus Hk. et Grev. Ic. Fil. t. 94. 1829 = C. angustatus.
sphaerostichus Copeland, Polyp. Philipp. 114. 1905.
spissus Klf. Enum. 126. 1824.

splendens J. Sm. JoB. **3**. 396. 1841.
stenophyllus Gies. Niph. 131. 1901.
sticticus Kze. Linn. **24**. 257. 1851.
stigmosus Moore, Ind. 276. 1861.
subfurfuraceus Bedd. Ferns br. Ind. t. 259. 1868.
subvelutinus Christ, Ann. Cons. Jard. bot. Genève **7-8**. 331. 1904.
tener Gies. Niph. 211. 1901.
tonkinensis Gies. Niph. 144. 1901
tricholepis Carr. in Seem. Fl. Vit. 367. 1873 = C. blepharolepis.
tricuspe J. Sm. Hist. Fil. 98. 1875 = C. hastatus.
vaccinifolius Keys. Pol. Cyath. Hb. Bung. 38 1873 = Polypodium v.
varius Klf. Enum. 125. 1824.
venosus Bl. Fl. Javae, Fil. 63 t. 28 f. 1. 1828 = C. stigmosus.
vittarioides Pr. Tent. 202. 1836 = C. spissus.
Warburgii Gies. Niph. 163. 1901.
xiphioides Christ, Bull. Soc. Fr. **52** Mém. I. 25. 1905.

NIPHOPSIS J. Smith, Cat. cult. Ferns 6. 1857 = **Cyclophorus.**
angustatus J. Sm. Cat. cult. Ferns 6. 1857.

NOTHOLAENA R.Brown, Prod. Fl. Nov. Holl. 145. 1810 — (Polypodiaceae (Nothochlaena auctt.). Gen. **71**).

affinis (Mett.) Moore, Ind. 233. 1861. HB. 273. NPfl. 274 — Mexico. Guatemala. Cheilanthes Mett. Cheil. 19. n. 2. 1859; Aleuritopteris Fourn. 1872.
albida Prantl, Engl. Jahrb. **3**. 405. 1882 (nomen) = N. cretacea.
albidula Sturm, Enum. pl. vasc. cr. chil. 16. 1858 = Pellaea nivea.
andromedaefolia Keys. Pol. Cyath. Hb. Bung. 29. 1873 = Pellaea.
argentea Lowe, Ferns **1** t. 55. 1856 = N. sulphurea var. candida.
argyrostigma J. Sm. JoB. **4**. 50. 1841 = Cheilanthes.
Aschenborniana Kl. Linn. **20**. 417. 1847. HB. 371. NPfl. 274 — Texas-Arizona. Mexico. — Cheilanthes Mett. n. 7. 1859. Nothochlaena bipinnata Liebm. 1849.
?*asplenioides* Christ, Engl. Jahrb. **24**. 132. 1897 — Guadeloupe.
atropurpurea Keys. Pol. Cyath. Hb. Bung. 30. 1873 = Pellaea.
aurantiaca Eat. Proc. Am. Acad. **22**. 462. 1887 — Mexico.
aurea (Poir.) Desv. Prod. 219. 1827 — Peru. — Pteris Poir. Enc. **5**. 710 1804.
Balansae Bak. JoB. **1878**. 301 — Paraguay.
bipinnata Liebm. Vid. Selsk. Skr. V. **1**. 214 (seors. 62). 1849 = N. Aschenborniana.
bonariensis (Willd.) C. Chr. Ind. 6. 1905; 459. 1906 — Amer. trop. et subtrop. (Texas-Chile). — Acrostichum Willd. sp. **5**. 114. 1810. Cincinalis ferruginea Desv. 1811; Nothochlaena Desv. 1813. HB. 370. Chr. 150. NPfl. 272; Cheilanthes Willd.: Link 1822. Mett. n. 12. Pellaea Nees 1847; Nothochlaena rufa Pr. 1825.
brachypus (Kze.) J. Sm. Ferns br. and for. 172 1866. (HB. 371) — Mexico. Cheilanthes squamosa var.? brachypus Kze. Linn. **18**. 340. 1844; C. brachypus Kze. Linn. **23**. 307. 1850. Mett. n. 8.
Brackenridgei Bak. Syn. ed. I. 371. 1868 = N. doradilla.
Brownii Desv. Prod. 220 1827 — Australia.
Gymnogramme Kuhn 1869; Notholaena vellea R.Br. Prod. Fl. N. Holl. 146. 1810. NPfl. 274 (non Desv. 1827); N. lanuginosa auctt. quoad pl. austral. HB. 370.
bryopoda Maxon, Proc. Biol. Soc. Wash. **18**. 205. 1905 — Mexico.
Buchanani Bak. Syn. 373. 1868. NPfl. 274 — Natal.

Bureaui Christ. Bull. Soc. Fr. **52** Mém. I. 59. 1905 — Yunnan.
californica Eat. Bull. Torr. Cl. **10**. 27. 1883 = N. cretacea.
calomelanos Keys. Pol. Cyath. Hb. Bung. 29. 1873 = Pellaea hastata.
canariensis Desv. Prod. 220. 1827 = N. Marantae.
candida Hk. sp. **5**. 110. 1864 = N. sulphurea var.
candida var. quinquefido-palmata Hk. sp. **5**. 111. 1864 = N. Hookeri.
canescens Kze. Pl. Preiss. **2**. 110. 1847 — Australia.
Cheilanthes Mett. n. 18. 1859 (non Kze. 1839).
capensis Spr. Tent. Suppl. Syst. 32. 1828 = Cheilanthes hirta.
carnosa Wall. List n. 138. 1828 = Drymoglossum.
chalcolepis A.Br. (Kze. Linn. **13**. 135. 1839) = N. sinuata.
cheilanthoides Spr. Nova Acta **10**. 227 t. 17 f. 3-4. 1821 = Cheilanthes microphylla.
Chilensis Sturm, Enum. pl. vasc. cr. chil. 16. 1858 = Pellaea.
chinensis Bak. Gard. Chr. n. s. **14**. 494. 1880. Chr. 152. NPfl. 274 — China.
chrysophylla Kl. Allg. Gartenzeit **1855**. 265 = Pellaea nivea var.
ciliata Desv. Journ. de Bot. appl. **1**. 92. 1813 = N. hirsuta.
cinnamomea Bak. Syn. 515. 1874 — Guatemala.
cordata Desv. Journ. de Bot. appl. **1**. 92. 1813 = Ceterach.
cordata Keys. Pol. Cyath. Hb. Bung. 29. 1873 = Pellaea.
crassifolia Moore et Houlst.; Lowe, Ferns **1**. t. 14 A. 1856 = N. sinuata.
cretacea Liebm. Vid. Selsk. Skr. V. **1**. 216 (seors. 64). 1849 — California-Mexico. — Cheilanthes Mett. n. 5. 1859; Aleuritopteris Fourn. 1872. Notholaena californica Eat. 1883.
dealbata Kze. Amer. Journ. Sci. II. **6**. 82. 1848 = Pellaea.
deltoidea Bak. Syn. 514. 1874 — Mexico.
densa J. Sm. JoB. **3**. 395. 1841; Kze. Farnkr. **1**. 149 t. 65 f. 2. 1844 = N. hirsuta.
distans R.Br. Prod. Fl. N. Holl. 146. 1810. HB. 372. Chr. 151. NPfl. 274 — Australia. Polynesia. N. Zealand. Celebes. — Cheilanthes Mett. n. 19. 1859.
doradilla Colla, Mem. Ac. Torino **39**. 46 t. 73. 1836. HB. ed. II. 371 — Chile. Peru. — N. sinuata Brack. 1854 (non Klf. 1824); N. Brackenridgei Bak. 1868. HB. ed. I. 371.
Eckloniana Kze. Linn. **10**. 501. 1836. HB. 371. Chr. 150. NPfl. 274 — Africa austr. — Cheilanthes Mett. n. 10. 1859. Notholaena Krebsiana Pr. 1836 (nomen).
elegans Pr. Tent. 224. 1836 (nomen) — ?
eriophora Fée, Gen. 159 t. 13 f. 3. 1850-52. HB. 372. Chr. 152. NPfl. 274 — Brasilia. — Polypodium Hk. 1854; Cheilanthes Mett. n. 14. 1859.
Fendleri Kze. Farnkr. **2**. 87 t. 136. 1851 = Pellaea.
ferruginea Desv. Journ. de Bot. appl. **1**. 92. 1813 = N. bonariensis.
ferruginea Eat. Mem. Amer. Acad. n. s. **8**. 201. 1860 = Gymnopteris.
flavens Moore, Ind. LXX. 1857 = Pellaea.
fragilis Hk. sp. **5**. 114 t. 287 A. 1864. HB. 372 — Austr. trop.
Cheilanthes fragillima F. Muell. 1866.
Fraseri (Mett.) Bak. Syn. 514. 1874 — Argentina.
Cheilanthes Mett.; Kuhn, Linn. **36**. 83. 1869.
Galeottii Fée, Gen. 159. 1850-52 = N. Aschenborniana.
Gilliesii Fée, Gen. 159. 1850-52 = N. scariosa.
glabra Brack. Expl. Exp. **16**. 21. 1854 = N. hirsuta.
goyazensis Taubert, Engl. Jahrb. **21**. 421. 1896 — Brasilia centr.
Grayi Dav. Bull. Torr. Cl. **7**. 50. 1880 — Arizona. Texas.

hirsuta (Poir.) Desv. Journ. de Bot. appl. **1**. 93. 1813. HB. 372. NPfl. 274 — Asia. Austr. Polynesia trop. — Pteris Poir. Enc. **5**. 719. 1804 (non Bory 1804); Cincinalis Desv. 1811; Cheilanthes Mett. n. 17. 1859 — Pteris nudiuscula β R.Br. 1810; Notholaena Desv. 1827; Pellaea? Hk. 1858; Cheilanthes Moore 1861 — Acrostichum javense Willd. 1810; Cincinalis Desv. 1811; Notholaena Desv. 1827; Cheilanthes Moore 1861 (an ad Cheilanthem tenuifoliam pertinens?) — Nothochlaena sulcata Link 1833. N. densa J. Sm. 1841; Kze. 1844; Cheilanthes Fée 1850-52. Mett. n. 16.

hirta J. Sm. JoB. **4**. 50. 1841 = Cheilanthes.

hirta Goldm. Nova Acta **19**. Suppl. I. 455. 1843 = N. hirsuta.

Hookeri Lowe, Ferns **1**. 59. 1856 (t. 13) = Pellaea nivea.

Hookeri Eat. U. S. Geogr. Surv. W. 100 Merid. **6**. 308 t. 30. 1878 — California-Mexico. — N. candida var. quinquefido-palmata Hk. 1864. — (prob. = N. sulphurea var.).

hypoleuca Kze. Linn. **9**. 54. 1834. HB. 372. NPfl. 274 — Chile. Cheilanthes Mett. n. 11. 1859.

inaequalis Kze. Farnkr. **1**. 146 t. 64 f. 1. 1844. HB. 371. NPfl. 274 — Afr austr. Cheilanthes Mett. n. 15. 1859.

incana Pr. Rel. Haenk. **1**. 19 t. 1 f. 2. 1825 = Pellaea nivea.

interrupta Klf. Enum. 137. 1824 = ? N. distans.

javensis Desv. Prod. 221. 1827 = N. hirsuta.

Krebsiana Pr. Tent. 224. 1836 = N. Eckloniana.

laevis Mart. et Gal. Mém. Ac. Brux. **15**. 46. 1842 = N. sinuata var.

lanuginosa Desv.; Poir. Enc. Suppl. **4**. 110. 1816 = N. vellea.

lasiopteris F. Muell. JoB. **5**. 106. 1853 = N. Brownii.

Lemmoni Eat. Bull. Torr. Cl. **7**. 63. 1880 — Arizona.

lendigera J. Sm. Bot. Mag. **72** Comp. 10 1846 = Cheilanthes.

lentigera J. Sm. JoB. **4**. 50. 1841 = Cheilanthes lendigera.

lepigera Bak. JoB. **1884** 53 = Mohria.

lutea (Desv.) Moore. Ind. 11. 1857 — Amer. austr. Acrostichum Desv. Prod. 211. 1827. — (certe Pellaea (Cincinalis) sp.).

Marantae (L.) R.Br. (Prod. Fl. N. Holl. 146. 1810); Desv. Journ. de Bot. appl. **1**. 92. 1813. HB. 371. Chr. 151. NPfl. 274 — Ins. Atlant. Mediterr.-Himalaya. Yunnan. Acrostichum L. sp. **2**. 1071. 1753; Ceterach DC. 1805; Cincinalis Desv. 1811; Gymnogramme Mett. 1856. Acrostichum canariense Willd. 1810; Notholaena Desv. 1827.

Matthewsii Gris. Abh. Ges. Wiss. Gött. **19**. 276. 1874 = Cheilanthes pruinata.

microphylla Keys. Pol. Cyath. Hb. Bung. 28. 1873 = Cheilanthes.

microphylla Bolle, Sitzber. Nat. Fr. Berlin — 1858 = Cheilanthes persica.

micropteris Keys. Pol. Cyath. Hb. Bung. 28. 1873 = Cheilanthes.

mollis Kze. Linn. **9**. 54. 1834. HB. 372. NPfl. 274 — Chile. Guatemala. Cheilanthes Pr. 1836. Mett. n. 28.

mollis ht.; Gard. Chr. III. **18**. 638. 1893 — Hort.

monosticha Moore. Ind. 231. 1861 = N. sulphurea var.

myriophylla J. Sm. JoB. **4**. 50. 1841 = Cheilanthes.

Nealleyi Seaton. Contr. U. S. Nat. Herb. **1**. 61. 1890 = N. Schaffneri.

Newberryi Eat. Bull. Torr. Cl. **4**. 12. 1873. HB. 515 — California.

nigricans (Willd.) Desv. Prod. 221. 1827 — Hispaniola. Pteris Willd. sp. **5**. 396. 1810. (Plum. t. 42).

nivea Desv. Journ. de Bot. appl. **1**. 93. 1813 = Pellaea.

nudiuscula Desv. Prod. 221. 1827 = N. hirsuta.
obducta (Mett.) Bak. Syn. 515. 1874 — Argentina. Bolivia. Columbia.
Cheilanthes Mett.; Kuhn, Linn. **36**. 83. 1869.
palmatifida Kze. Farnkr. **1**. 148. 1844 (nomen) = N. eriophora.
Palmeri Bak. in Hk. Ic. pl. t. 1678. 1887 — Mexico. (an Pellaea sp.?).
Parryi Eat. Amer. Nat. **9**. 351. 1875. NPfl. 274 — Utah-California-Arizona.
paucijuga Bak. Syn. 515. 1874 = N. pumilio.
persica Bory, Bél. Voy. Bot. **2**. 23. 1833 = Cheilanthes.
peruviana Desv. Prod. 220. 1827 = N. scariosa.
pilosa Hk. et Arn. Beech. Voy. Bot. 74. 1832 = N. hirsuta.
piloselloides Klf. Enum. 133. 1824 = Drymoglossum heterophyllum.
Plukenetii Fée, Gen. 159. 1850-52; 7 mém. 39 t. 23 f. 4 = N. vellea.
Pohliana Kze. Farnkr. **1**. 45. 1840. HB. 372 — Brasilia.
Cheilanthes Mett. Cheil. 23. n. 13. 1859. — (an Adiantopsis sp.?).
Prenticei (Luerss.) Bak. Ann. Bot. **5**. 481. 1891 — Queensland. Austr. bor.
Cheilanthes Luerss. Bot. Centralbl. **9**. 442. 1882.
Pringlei Dav. Bull. Torr. Cl. **13**. 132 t. 58. 1886 — Mexico.
profusa Pr. Tent. 224. 1836 = ? Pellaea involuta.
pruinosa Fée, 8 mém. 78. 1857: 10 mém. 20 t. 34 f. 2 = N. sinuata.
?**pteridiformis** (Ces.) Bak. Malesia **3**. 49. 1886 — N. Guinea.
Gymnogramme Ces. Rend. Ac. Napoli **16**. 30. 1877.
pulchella Kze. Bot. Zeit. **1843**. 633; Linn. **17**. 567. 1843 = Pellaea dealbata.
pulveracea Kze. Linn. **13**. 135. 1839 = N. sulphurea.
pumila Link, Fil. sp. 146. 1841 = ? N. distans vel Pellaea involuta.
pumilio R.Br. Prod. Fl. N. Holl. 146. 1810 — Queensland.
Cheilanthes F. Muell. 1882; Nothochlaena paucijuga Bak. 1874. HB. 515.
Rawsoni Pappe; Pappe et Raws. Syn. Fil. Afr. austr. 42. 1858. HB. 370. NPfl. 274 — Africa austr. — Cheilanthes Mett.; Kuhn 1868.
remota Klf. Enum. 138. 1824 = N. hirsuta.
Reynoldsii F. Muell. Fragm. **8**. 175. 1874 = Gymnopteris.
rigida Dav. Gard. and For. **4**. 519 f. 80. 1891 — Mexico.
rufa Pr. Rel. Haenk. **1**. 19. 1825 = N. bonariensis.
rufidula Desv. Prod. 221. 1827 = Woodsia ilvensis.
scariosa (Sw.) Bak. Fl. bras. **1**^{2}. 540. 1870 — Amer. trop.
Acrostichum Sw. Syn. 16. 1806. Cheilanthes Desv. 1827 (non Pr. 1825). A. lanuginosum Willd. Schr. Ak. Erfurt **1802**. 31 t. 3 f. 4. 1802 (non Desf. 1800). Cheilanthes squamosa Gill.; Hk. et Grev. t. 151. 1829. Mett. n. 9; Notholaena Fée 1857. HB. 371. Chr. 150. NPfl. 274 — N. peruviana Desv. 1827.
Schaffneri (Fourn.) Und.; Dav. Gard. and For. **4**. 519. 1891 — Mexico. Texas.
Aleuritopteris Fourn. Bull. Soc. Fr. **27**. 328. 1880. Notholaena Nealleyi Seaton 1890.
semiglabra Kze. Farnkr. **2**. 59 t. 124 f. 2. 1850 = Cheilanthes tenuifolia.
setigera Desv. Journ. de Bot. appl. **1**. 93. 1813 = Woodsia ilvensis.
sinuata (Lag.) Klf. Enum. 135. 1824. HB. 370. Chr. 150. NPfl. 272 — Arizona-Chile. — Acrostichum Lag.; Sw. Syn. 14. 1806; Gymnogramma Pr. 1836 — Notholaena laevis Mart. et Gal. 1842 — N. pruinosa Fée 1857.
sinuata Brack. Expl. Exp. **16**. 19. 1854 = N. doradilla.
squamata Moore et Houlst.; Gard. Mag. Bot. **3**. 20. 1851 = N. brachypus.
squamosa Lowe, Ferns **1**. t. 17 B. 1856 = N. brachypus.

squamosa Fée, 9 mém. 10. 1857 = N. scariosa.
stellapilis J. Sm. JoB. 4. 50. 1841 (nomen) —?)
Streetiae Bak. Journ. Linn. Soc. 16. 204. 1877 = Cheilanthes.
subcordata Desv. Journ. de Bot. appl. 1. 92. 1813 = N. Marantae.
sulcata Link, Hort. Berol. 2. 367. 1833 = N. hirsuta var.
sulphurea (Cav.) J. Sm. Bot. Voy. Herald. 1. 233. 1854. HB. 373. Chr. 153. NPfl. 274 — California-Chile. — Pteris Cav. Descr. 269. 1802; Allosorus Pr. 1836; Pteris Brack. 1854; Cheilanthes pulveracea Pr. 1825. Mett. n. 70; Allosorus Pr. 1836; Notholaena Kze. 1839; Aleuritopteris Fée 1857. A. Preslii Fourn. 1872 — Ceropteris monosticha Fée 1857; Cheilanthes Mett. n. 4. 1859; Nothochlaena Moore 1861 — Cheilanthes Borsigiana Reichb. f. et Warz. 1858. Mett. n. 6 — C. candida Mart. et Gal. 1842. Mett. n. 3; Aleuritopteris Fée 1850-52; Nothochlaena Hk. 1864.
Szovitsii Keys. Pol. Cyath. Hb. Bung. 28. 1873 = Cheilanthes persica.
tectaria Desv. Prod. 219. 1827 = N. sinuata.
tenera Gill.; Hk. Bot. Mag. t. 3055. 1831 = Pellaea.
ternifolia Keys. Pol. Cyath. Hb. Bung. 29. 1873 = Pellaea.
tomentosa Desv. Journ. de Bot. appl. 1. 92. 1813 = N. bonariensis.
tomentosa J. Sm. JoB. 4. 50. 1842 = Cheilanthes.
tricholepis Bak. JoB. 1883. 245 — Africa orient. trop.
trichomanoides (L.) R.Br. (Prod. Fl. N. Holl. 145. 1810); Desv. Journ. de Bot appl. 1. 92. 1813. HB. 373. Chr. 150. NPfl. 272 — Ind. occ. Mexico. Guatemala. Pteris L. sp. 2. 1074. 1753; Cincinalis Desv. 1811; Cheilanthes Mett. n. 1. 1859; Acrostichum pterioides Bernh. 1802.
undulata Wall. List n. 140. 1828 = Leptochilus scalpturatus var.
vellea (Ait.) Desv. Journ. de Bot. appl. 1. 93. 1813 — Ins. Atlant.-Mediterr. Nubia-Himalaya. Afghanistan. — Acrostichum Ait. Hort. Kew. 3. 457. 1789; Cincinalis Desv. 1811; Gymnogramme Kuhn 1868; Cosentinia Todaro 1866; Acrostichum lanuginosum Desf. 1800; Notholaena Desv.; Poir. 1816. HB. 370. Chr. 151. NPfl. 274; Gymnogramme A.Br. 1854 — Nothochlaena Plukenetii Fée 1850-52.
vellea R.Br. Prod. Fl. N. Holl. 146. 1810 = N. Brownii.
vestita Desv. Journ. de Bot. appl. 1. 93. 1813 = Cheilanthes lanosa.

Notocarpia Presl. Tent. Pterid. 54. 1836 *(Cyathea §)* = Cyathea.

NOTOGRAMME Presl, Epim. Bot. 263. 1849 = **Coniogramme.**
japonica Pr. Epim. 263. 1849.

NOTOLEPEUM Newman, Hist. Brit. Ferns ed. II. 9. 1844 = **Ceterach.**
Ceterach Newm. Hist. Brit. Ferns ed. II. 9. 1844 = C. officinarum.

Notophoria Presl, Abh. böhm. Ges. Wiss. V. 5. 349. 1848 *(Hemitelia §)* = Hemitelia.

Nymphopteris Webb et Berth. Hist. nat. Canar. 3². Phyt. 3. 449. 1849 *(Pteris §)* [= Pteridium.

OCHLOGRAMMA Presl, Epim. Bot. 93. 1849 = **Diplazium.**
Cumingii Pr. Epim. 94. 1849.

OCHROPTERIS J. Smith, Journ. Bot. 4. 158. 1841 — (Polypodiaceae Gen. 82).
angusta Fée, Gen. 120 t. 29 f. 3. 1850-52 = O. pallens.
pallens (Sw.) J. Sm. JoB. 4. 158. 1841. HB. 127. Chr. 141. NPfl. 290 — Madagascar. Ins. Mascar. — Adiantum Sw. Schrad. Journ. 1800². 84. 1801; Cheilanthes Desv. 1827; Pteris Mett. 1856; Cryptogramme Prantl 1882; Cheilanthes davallioides Bory; Willd. 1810 — Ochropteris angusta Fée 1850-52 — O. peltigera Fée 1850-52.
peltigera Fée, Gen. 102 t. 29 f. 4. 1850-52 = O. pallens.

ODONTOLOMA J. Smith, Journ. Bot. **3**. 415. 1841; Lond. JoB. **1**. 424. 1842 = [**Lindsaya.**
adiantoides Pr. Epim. 97. 1849.
bifidum Mett. Fil. Lips. 104. 1856.
Blumeanum Mett. Fil. Lips. 104. 1856.
Boryanum J. Sm. JoB. **3**. 415. 1841 = L. repens.
cuneifolium Pr. Epim. 97. 1849 = L. adiantoides.
fijiensis J. Sm. Hist. Fil. 269. 1875 = Davallia solida var.
Goudotianum Mett. Fil. Lips. 104. 1856.
hemipterum Pr. Epim. 98. 1849 = L. repens var.
Hookeri J. Sm. Lond. JoB. **1**. 424. 1842 = L. repens.
hymenophylloides J. Sm. Hist. Fil. 269. 1875.
Macraeanum Brack. Expl. Exp. **16**. 226. 1854 = L. repens var.
Parkeri Pr. Epim. 97. 1849.
pulchellum J. Sm. JoB. **3**. 415. 1841; Field. et Gardn. Sert. pl. t. 51. 1844 [= L. adiantoides.
repens Pr. Epim. 97. 1849.
Schlechtendalii Fourn. Mex. pl. **1**. 132. 1872 = Odontosoria S.
tenuifolium J. Sm. JoB. **3**. 415. 1841 = L. Blumeana.
tenuifolium Pr. Epim. 98. 1849.
triquetra J. Sm. Hist. Fil. 269. 1875 = L. tenuifolia.

ODONTOMANES Presl, Epim. Bot. 20. 1849 = **Trichomanes.**
Hostmannianum Pr. Epim. 21. 1849.

ODONTOPTERIS Bernhardi, Schrad. Journ. **1800²**. 127. 1801 = **Lygodium.**
scandens Bernh. l. c.

ODONTOSORIA (Pr.) Fée, Gen. Fil. 325. 1850-52 — (Polypodiaceae Gen. **34**).
Davallia § Odontosoria Presl, Tent. Pter. 129. 1836.

aculeata (L.) J. Sm. Cat. cult. ferns 67. 1857. NPfl. 216 — Ind. occ.
Adiantum L. sp. **2**. 1096. 1753; Davallia Sm. 1793. HB. 102; Trichomanes Poir. 1808; Stenoloma Fée 1850-52; Microlepia Mett. 1856; Lindsaya Mett. 1861. Chr. 296; Lindsayopsis Kuhn 1882.

angustifolia (Bernh.) C. Chr. Ind. 24. 1905; 464. 1906 — N. Zealand.
Lindsaea Bernh. Schr. Ak. Erfurt **1802**. 14 f. 6; Adiantum clavatum Forst. Prod. 83. 1786 (non L. 1753); Davallia Forsteri Carr. 1873 (nomen); Bak. Syn. 470. 1874; Microlepia Armstr. 1881.

bifida (Klf.) J. Sm. Hist. Fil. 264. 1875. NPfl. 215 — Brasilia.
Davallia Klf. Enum. 222 1824. HB. 101; Odontoloma Mett. 1856; Acrophorus Moore 1857; Lindsaya Mett.; Kuhn 1882. Chr. 296. Stenoloma Glaziovii Fée 1869.

biflora (Klf.) C. Chr. Ind. 207. 1905; 464. 1906 — Luzon.
Davallia Klf. Enum. 221. 1824; Microlepia Mett. 1856.

chinensis (L.) J. Sm. Bot. Voy. Herald 430. 1857. NPfl. 215 — Japonia. China. Asia trop. Polynesia. Ins. Comorae et Mascar. Madagascar.
Trichomanes L. sp. **2**. 1099. 1753; Adiantum Burm. 1768; Davallia Sm. 1793; Microlepia Mett. 1856; Lindsaya Mett.; Kuhn 1868; Stenoloma Bedd. 1883 — Davallia tenuifolia Sw. 1801. HB. 102; Adiantum Poir. 1810; Stenoloma Fée 1850-52; Microlepia Mett. 1856; Odontosoria J. Sm. 1857; Lindsaya Mett. 1861. Chr. 296 (non Bl. 1828) — Adiantum chusanum L. 1753; Davallia Willd. 1810 — Trichomanes cuneiforme Forst. 1786; Davallia Sw. 1801.

clavata (L.) J. Sm. Hist. Fil. 264. 1875. NPfl. 215 — Ind. occ. Florida.
Adiantum L. sp. **2**. 1096. 1753; Davallia Sm. 1793. HB. 102; Stenoloma Fée 1850-52; Lindsaya Mett. 1861. Chr. 295; Schizoloma Kuhn 1882.

decomposita (Bak.) C. Chr. Ind. 209. 1905; 465. 1906 — Madagascar.
Davallia Bak. JoB. **1884**. 141.

divaricata J. Sm. Hist. Fil. 264. 1875 = O. Schlechtendalii.

ferruginea Diels, Nat. Pfl. **1**^{4}. 215. 1899 = O. Melleri.

flabellifolia (Bak.) C. Chr. Ind. 210. 1905; 465. 1906 — Madagascar.
Davallia Bak. Journ. Linn. Soc. **15**. 414. 1876; Lindsaya Kuhn 1879.

fumarioides (Sw.) J. Sm. Hist. Fil. 264. 1875 — Ind. occ. Columbia.
Davallia Sw. Schrad. Journ. **1800**^{2}. 89. 1801. HB. 102; Trichomanes Poir. 1808; Stenoloma Fée 1850-52; Lindsaya Mett. 1864. Acrostichum aculeatum L. 1759; Trichomanes Sw. 1788; Davallia Hedw. 1803 (non Odontosoria supra).

Lenormandi (Bak.) C. Chr. Ind. 211. 1905; 465. 1906 — N. Caledonia.
Davallia Bak. Syn. 471. 1874.

meifolia (HBK.) C. Chr. Ind. 164. 1905; 465. 1906 — Panama-Brasilia.
Davallia HBK. Nov. Gen. et Sp. **1**. 23. 1815. HB. 470; Lindsaya Mett. 1869; Darea fumarioides H.B.Willd. sp. **5**. 299. 1810; Caenopteris Desv. 1827 (non Odontosoria supra). Lindsaya Sprucei Hk. 1867; Davallia Bak. 1870; Odontosoria Diels 1899. NPfl. 215. Stenoloma gratissima Fée 1869.

Melleri (Hk.) C. Chr. Ind. 212. 1905; 465. 1906 — Ins. Comorae. Madagascar. Réunion. — Davallia Hk.; Bak. Syn. ed. I. 102. 1867. D. ferruginea Desv. 1827. HB. ed. II. 102 (non Cav. 1802); Lindsaya Kuhn 1868. Chr. 297; Schizoloma Kuhn 1882; Odontosoria Diels 1899. NPfl. 215. Davallia thalictroides Pr. 1836 (nomen).

microphylla J. Sm. Hist. Fil. 264. 1875 = Lindsaya.

Moorei Kuhn, Chaetopt. 346. 1882 = Davallia.

odontolabia (Bak.) Diels, Nat. Pfl. **1**^{4}. 215. 1899 — Madagascar.
Davallia Bak. JoB. **1884**. 140.

parvula Fée, 10 mém. 46 t. 36 f. 2. 1865 — Ceylon.

retusa (Cav.) J. Sm. Bot. Herald 430. 1857. NPfl. 215 — Malesia. Melanesia.
Davallia Cav. Descr. 278. 1802; Stenoloma Fée 1850-52; Lindsaya Mett. 1856. HB. 110. Chr. 296; Schizoloma Kuhn 1882. Lindsaya cuneifolia Pr. 1825 — Davallia decipiens Ces. 1877.

scandens (Desv.) C. Chr. Ind. 354. 1905; 465. 1906 — Peru.
Humata Desv. Prod. 324. 1827; Davallia Moore 1861; Lindsayopsis Kuhn 1882. (an adhuc O. uncinella?).

Schlechtendalii (Pr.) C. Chr. Ind. 209. 1905; 465. 1906 — Mexico-Guatemala.
Davallia Pr. 1836. HB. 103; Stenoloma Fée 1850-52; Microlepia Mett. 1856; Odontoloma Fourn. 1872; Lindsaya Christ 1897. 296; Davallia divaricata Schlecht. et Cham. Linn. **5**. 617. 1830 (non Bl. 1828); Odontosoria J. Sm. 1875. NPfl. 215; Lindsayopsis Kuhn 1882.

scoparia (Mett.) Diels, Nat. Pfl. **1**^{4}. 215. 1899 — N. Caledonia.
Lindsaya Mett. Ann. sc. nat. IV. **15**. 64. 1861; Davallia Hk. 1867. HB. 101; Microlepia Carr. 1873; Schizoloma Kuhn 1882.

Sprucei Diels, Nat. Pfl. **1**^{4}. 215. 1899 = O. meifolia.

tenuifolia J. Sm. Cat. cult. ferns 67. 1857 = O. chinensis.

? triloba Fée, 11 mém. 92. 1866 = Davallia.

uncinella (Kze.) Fée, Gen. 326 t. 27 B. f. 1. 1850-52. NPfl. 215 — Ind. occ.
Davallia Kze. Bot. Zeit. **1850**. 213; Farnkr. **2**. 96 t. 140. 1851. HB. 102; Microlepia Mett. 1856; Lindsaya Krug 1897. Chr. 297. Prosaptia bipinnata Pr. 1836 (nomen); Davallia? Hk. 1846. — (an = O. scandens supra?).

viridis Kuhn, Chaetopt. 346 t. 1 f. 2. 1882 = Lindsaya.

OEOSPORANGIUM Visiani, Atti Istituto Veneto III. **12**. 656. 1867 = **Cheilanthes.**
persica Vis. Fl. dalm. suppl. 1872 } = C. persica.
Szovitsii Vis. Atti Ist. Veneto III. **12**. 656 t. 10-11. 1867 }
OETOSIS Necker, Element. bot. **3**. 318. 1790 —? (vide seqq.).
OETOSIS O.Ktze., Rev. Gen. Plant. **2**. 817. 1891.
angustifolia O.Ktze. l. c. = Vittaria.
blechnodes O.Ktze. l. c. = Taenitis blechnoides.
carnosa O.Ktze. l. c. = Drymoglossum.
furcata O.Ktze. l. c. = Eschatogramme.
lanceolata O.Ktze. l. c. = Paltonium.
obtusa O.Ktze. l. c. = Taenitis blechnoides var.
?ophioglossodes O.Ktze. l. c. = Hymenolepis spicata var.
pilosellodes O.Ktze. l. c. = Drymoglossum heterophyllum.
rigida O.Ktze. l. c. = Drymoglossum.
OETOSIS Greene, Pittonia **4**. 103. 1900 = **Vittaria.**
elongata, ensiformis, isoetifolia, lineata, zosteraefolia Greene, l. c. = V. sp.
filiformis Greene l. c. = V. lineata. [homonym.
OLEANDRA Cavanilles, Anales de Historia Natural **1**. 115. 1799; Descr. plant. 252. 1802 — (Polypodiaceae Gen. **25**).
articulata (Sw.) Pr. Tent. 78 t. 2 f. 12. 1836. HB. 302. Chr. 280. NPfl. 204 — Africa trop. et austr. Ins. Mascar. Ins. Sechellae.
Aspidium Sw. Schrad. Journ. **1800²**. 30. 1801 (pt.); Polypodium Juss.; Poir. Enc. **5**. 514. 1804; Hypopeltis Bory 1833 — Oleandra Welwitschii [Bak. 1867.
bantamensis Kze. Farnkr. **2**. 72. 1850 = O. neriiformis.
chinensis Hance, Ann. sc. nat. IV. **18**. 238. 1862 = O. Cumingii.
ciliata Kl.; Kuhn, Linn. **36**. 126. 1869 — N. Hebridae. Ins. Pinorum.
colubrina (Blanco) Copeland in Perkins, Fragm. 179. 1905 — Luzon.
Blechnum Blanco, Fl. Filip. 834. 1837.
Cumingii J. Sm. JoB. **3**. 413. 1841 (nomen); Pr. Epim. 41. 1849. HB. 303. NPfl. 204 — China merid. Assam. India austr. Malesia. Luzon. *Tahiti.
O. chinensis Hance 1862.
—* O. Sibbaldii Grev. 1848 (Tahiti).
cuspidata Bak. Malesia **3**. 44. 1886 — N. Guinea.
densifrons Kze. Bot. Zeit. **1851**. 347 = O. articulata.
distenta Kze. Bot. Zeit. **1851**. 347 = O. articulata.
hirta Brack. Expl. Exp. **16**. 214 t. 29. 1854 = O. neriiformis var.
hirtella Miq.; Kze. Farnkr. **2**. 70 t. 129. 1850 = O. neriiformis.
lomatopus Pr. Epim. 43. 1849 = O. neriiformis.
macrocarpa Pr. Epim. 41. 1849 — Luzon.
micans Kze. Bot. Zeit. **1851**. 346 = O. nodosa.
micrantha Kze. Bot. Zeit. **1851**. 349 = O. neriiformis.
mollis Pr. Epim. 41. 1849 = O. neriiformis.
Moritzii Pr. Epim. 42. 1849 = O. musifolia.
musifolia (Bl.) Pr. Epim. 42. 1849. HB. 302 — Ceylon. Malesia.
Aspidium Bl. Enum. 141. 1828.
neriifolia Pr. Tent. 78. 1836 = O. neriiformis.
neriiformis Cav. Anal. Hist. Nat. **1**. 115. 1799; Descr. 253. 1802. HB. 302. Chr. 280. NPfl. 204 — Ind. bor.-Malesia-Polynesia. Amer. austr. ? Afr. occ. trop.
Aspidium Sw. 1806. A. pistillare Sw. 1801; Ophiopteris verticillata Reinw. 1824 — Oleandra pilosa Hk. 1842 — O. hirta Brack. 1854.

neriiformis Bedd. Ferns S. Ind. t. 91. 1863-65 = O. musifolia.

nodosa (Willd.) Pr. Tent. 78. HB. 303. Chr. 280. NPfl. 204 — Amer. trop. Ins. Fernando Po (Guinea). — Aspidium Willd. sp. **5**. 211. 1810. A. articulatum Schkuhr 1806 (non Sw. 1806); Polypodium Vahl 1807 (non Poir. [1804).

ornata Christ, Verh. Nat. Ges. Basel **11**. 240. 1895 = O. Whitmeei.

parallela Keys. Pol. Cyath. Hb. Bung. 41. 1873 = Humata Gaimardiana.

phyllarthron Pr. Epim. 42. 1849 = O. neriiformis.

pilosa Hk. Gen. t. 45 B. 1840 = O. neriiformis.

Sibbaldii Grev. Ann. Nat. Hist. II. **1**. 327. 1848 = O. Cumingii*.

tricholepis Kze. Bot. Zeit. **1851**. 349 — Borneo.

Trujillensis Karst. Fl. Col. **1**. 147 t. 73. 1860 = O. neriiformis.

Wallichii (Hk.) Pr. Tent. 78. 1836. HB. 303 — Ind. bor. Aspidium Hk. Exot. Fl. **1** t. 5. 1823. Neuronia asplenioides Don 1825.

Welwitschii Bak. (Syn. 303. 1867) = O. articulata var.

Whitmeei Bak. JoB. **1876**. 11. NPfl. 204 — Celebes. Ins. Philipp. Samoa.

OLFERSIA Raddi, Opusc. sci. Bologna **3**. 283. 1819 = **Polybotrya**; Presl, Tent. Pterid. 232. 1836 = **Elaphoglossum et Stenochlaena.**

acrocarpa, angulata, auricoma, callaefolia, ciliata, conformis, crinita, dissimilis, foeniculacea, laurifolia, longifolia, obducta, peltata, pilosa, Sellowiana, stricta, succisaefolia, tecta, undulata, vestita Pr. Tent. 234. 1836 — aemula, cuspidata, decorata, decurrens, dimorpha, flabellata, gorgonea, hybrida, Lingua, pachyphylla, petiolosa, scolopendrifolia, Sieberi, simplex Pr. l. c. 235 — apoda, muscosa, spatulata, tomentosa, villosa Pr. l. c. 233 = Elaphoglossum sp. homonym.

aculeata Bl. Fl. Javae Fil. Suppl. t. 96. —. = Stenochlaena a.

angustata Pr. Tent. 234. 1836 = Elaphoglossum conforme.

bifurcata Pr. Tent. 234. 1836 = Elaphoglossum furcatum.

Blumeana Pr. Tent. 235. 1836 = Elapoglossum decurrens.

caudata Kze. Linn. **21**. 206. 1848 = Polybotrya.

cervina Kze. Flora **1824**. 312 = Polybotrya.

corcovadensis Raddi, Opusc. sci. Bol. **3**. 283 t. 11 f. b. 1819 = Polybotrya cervina var.

glabrescens Pr. Tent. 234. 1836 = Elaphoglossum tomentosum.

gracilis Bl. Fl. Javae Fil. Suppl. t. 95. —. = Stenochlaena g.

horridula Pr. Tent. 233. 1836 = Elaphoglossum spathulatum var.

integrifolia Pr. Tent. 235. 1836 = Stenochlaena pollicina.

japurensis Pr. Tent. 234. 1836 = Stenochlaena marginata.

Kaulfussiana Pr. Tent. 290. 1836 }
Kunzeana Pr. Tent. 235. 1836 } = Stenochlaena sorbifolia var. 2.

Langsdorffii Pr. Tent. 234. 1836 = Elaphoglossum muscosum.

lepidota Pr. Tent. 233. 1836 = Elaphoglossum plicatum.

?lineata Pr. Tent. 235. 1836 = Stenochlaena pollicina.

lomarioides Pr. Tent. 235. 1836 = Stenochlaena pollicina.

marginata Pr. Tent. 234. 1836 = Elaphoglossum conforme.

neriifolia Pr. Tent. 235. 1836 = Elaphoglossum petiolatum var.

nervosa Moore, Ind. 1. 1857 = Elaphoglossum.

obtusa Moore, Ind. 1. 1857 = Elaphoglossum gorgoneum.

paleacea Pr. Tent. 234. 1836 = Elaphoglossum hirtum.

piloselloides Pr. Tent. 233. 1836 = Elaphoglossum spathulatum var.

Richardi Moore, Ind. 1. 1857 = Elaphoglossum.

salicifolia Pr. Tent. 234. 1836 = Elaphoglossum petiolatum var.
scandens Pr. Tent. 235. 1836 = Stenochlaena palustris.
sorbifolia Pr. Tent. 234. 1836 = Stenochlaena s.
sphenophylla Pr. Tent. 235. 1836 = Elaphoglossum flabellatum.
splendens Pr. Tent. 233. 1836 = Elaphoglossum hirtum.
squamata Pr. Tent. 233. 1836 = Acrostichum squamosum Cav.
subdiaphana Moore, Ind. XVIII. 1857 = Elaphoglossum nervosum.
tripartita Pr. Tent. 234. 1836 = Elaphoglossum flabellatum.
triquetra Pr. Tent. 235. 1836 = Plagiogyria euphlebia.
variabilis Pr. Tent. 235. 1836 = Stenochlaena sorbifolia var. 1.
Vieillardii Carr. in Seem. Fl. Vit. 373. 1873 = Elaphoglossum.
viscosa Pr. Tent. 234. 1836 = Elaphoglossum petiolatum.

OLIGOCAMPIA Trevisan, Atti Istituto Veneto II. 2. 165. 1851 = **Dryopteris.**
microcarpa Trev. l. c. = D. otaria.

Oligolepidum Moore, Ind. Fil. XVI. 1857 *(Elaphoglossum §)* = Elaphoglossum.

ONOCLEA Linné, Spec. plant. 2. 1062. 1753 (Polypodiaceae Gen. 8).
attenuata Sw. Schrad. Journ. **1800**². 73. 1801 = Blechnum.
augescens Link, Fil. sp. 37. 1841 = O. sensibilis.
Boryana Sw. Syn. 111, 306. 1806 = Blechnum tabulare.
capensis Thbg. Prod. Fl. Cap. 171. 1800 = Blechnum.
crispa Hoffm. Deut. Fl. 2. 11. 1795 = Cryptogramma.
discolor Sw. Syn. 111. 1806 = Blechnum.
lineata Sw. Schrad. Journ. **1800**². 73. 1801 = Blechnum.
nodulosa Michx. Fl. bor. Am. 2. 272. 1803 = Woodwardia areolata.
nodulosa Schkuhr, Krypt. Gew. 1. 95 t. 104. 1809 = Matteuccia struthiopteris.
nuda Lab. Nov. Holl. Pl. Spec. 2. 96 t. 246. 1806 = Blechnum discolor var.
obtusilobata Schkuhr. Kr. Gew. 1. 95 t. 103. 1809 = O. sensibilis.
orientalis Hk. sp. 4. 161. 1862 = Matteuccia.
polypodioides L. Mant. 306. 1771 = Gleichenia.
polypodioides Sw. Fl. Ind. occ. 3. 1585. 1806 = Blechnum.
procera Spr. Schrad. Journ. **1799**². 267 = Blechnum capense.
quercifolia Willd. Schrift. Ak. Erfurt **1802**. 27 = Leptochilus zeylanicus.
scandens Sw. Syn. 112. 309. 1806 = Stenochlaena palustris.
sensibilis L. sp. 2. 1062. 1753. HB. 46. Chr. 284. NPfl. 166 — Amer. bor. atlant. Asia orient. — Calypterium Bernh. 1802; Riedlea Mirb. 1803; Pterinodes O.Ktze. 1891. Ragiopteris onocleoides Pr. 1836. Onoclea augescens Link 1841.
sorbifolia Sw. Syn. 112. 1806 = Stenochlaena.
Spicant Hoffm. Deutschl. Fl. 2. 11. 1795 = Blechnum.
spicata Sw. Schrad. Journ. **1801**². 299. 1803 = Hymenolepis.
striata Sw. Syn. 304. 422. 1806.
Struthiopteris Hoffm. Deutschl. Fl. 2. 11. 1795 = Matteuccia.
variabilis Poir. Enc. Suppl. 4. 149. 1816 = Stenochlaena sorbifolia var. 1.

ONOPTERIS Necker, Element. Bot. 3. 316. 1790 = **Asplenium.**

ONYCHIUM Kaulfuss. Jahr. d. Pharmacie, Berlin **1820**. 45; Enum. 144. 1824 — (Polypodiaceae Gen. 75).
angustifolium Kze. Farnkr. 2. 11. 1848 = Cheilanthes.
auratum Klf. Enum. 144. 1824 = O. siliculosum.
capense Klf. Enum. 145 t. 1 f. 8. 1824 = O. japonicum.
carnosum Reinw. Syll. Pl. 2. 3. 1824 = Lecanopteris.
carvifolium Fée. Gen. 132. 1850-52 = O. japonicum.

chaerophyllum Fée, Gen. 132. 1850-52 = Cheilanthes marginata.
chinense Fée, Gen. 132. 1850-52 (nomen) — (an Phorolobus Desv.?).
contiguum Hope, Journ. Bomb. Nat. Hist. Soc. **13**. 444. 1901 = O. japonicum var.
cornutum Hassk. Tijdschr. Nat. Gesch. **10**. 115. 1843 = Ceratopteris thalictroides.
densum Brack. Expl. Exp. **16**. 120 t. 13 f. 2. 1854 = Pellaea.
heterophyllum Kuhn, Engl. Jahrb. **24**. 98. 1897 = Anopteris hexagona.
japonicum (Thbg.) Kze. Bot. Zeit. **1848**. 507. HB. 143. Chr. 155 — Japonia. China. Himalaya. Java. Luzon. — Trichomanes Thbg. Fl. Jap. 340. 1784; Caenopteris Thbg. 1795; Darea Willd. 1810; Pteris Mett. 1856; Cryptogramme Prantl 1882. NPfl. 279. Onychium capense Klf. 1824 (patria erron.); Allosorus Pr. 1836 — Leptostegia lucida Don 1825; Onychium Spr. 1827: Cheilanthes Wall. 1828. C. contigua Wall. 1828; Onychium Hope [1901.
Krebsii Kze. Linn. **10**. 504. 1836 = Blechnum punctulatum var.
lucidum Spr. Syst. **4**. 66. 1827 = O. japonicum var.
melanolepis (Dcne.) Kze. Farnkr. **2**. 9 t. 104 f. 2. 1848. HB. 143. Chr. 155 — Abyssinia. Arabia. Persia. — Allosorus Dcne. Arch. Mus. **2**. 189. 1841; Pteris Ettingsh. 1865; Cryptogramme Prantl 1882. NPfl. 279. Allosorus cuspidatus Hochst. 1847.
micropterum Hk. Gen. Fil. ad t. 11. 1838 (nomen) —?
multifidum Fée, 8 mém. 74. 1857 — Cuba. — (an = O. strictum?).
?robustum Fée, Gen. 132. 1850-52 = Doryopteris.
siliculosum (Desv.) C. Chr. Ind. 20. 1905; 469. 1906 — Asia trop. Pteris Desv. Berl. Mag. **5**. 324. 1811; Phorolobus Desv. 1827; ? Acrostichum viviparum Cav. 1802 (non L.fil. 1781). Onychium auratum Klf. 1824. HB. 143. Chr. 155; Allosorus Pr. 1836; Pteris Mett. 1856; Cryptogramme Prantl 1882; NPfl. 279. Pteris chrysocarpa (chrysosperma) Hk. et Grev. 1829.
strictum Kze. Farnkr. **2**. 11. 1848. HB. 143. Chr. 155 — Cuba. Haïti. Puerto Rico. Cryptogramme Prantl; Salom. 1883. NPfl. 279.
tenue Christ, Bull. Boiss. II. **1**. 451. 1901 — N. Guinea. Java. Luzon.

ONYCHIUM Reinwardt, Sylloge plant. **2**. 2. 1824 = **Lecanopteris.**
carnosum Reinw. l. c. 3.

OOCHLAMYS Fée, Gen. Fil. 297. 1850-52 = **Dryopteris.**
Rivoirei Fée, Gen. 297. 1850-52 = D. opposita.

OPHIALA Desvaux, Prodrome 195. 1827 = **Helminthostachys.**
zeylanica Desv. Prod. 195. 1827.

OPHIODERMA (Bl.) Endlicher Gen. Plant. 66. 1836 = **Ophioglossum.**
Ophioglossum § Ophioderma Blume, Enum. pl. Jav. 259. 1828.
pendulum Pr. Suppl. 56. 1845.

OPHIOGLOSSUM Linné, (Gen. plant. 503) Spec. plant. **2**. 1062. 1753 — (Ophioglossaceae Gen. **1**).
Prantl = Prantl in Jahrb. Bot. Gart. u. Mus. Berlin **3**. 1884.
acuminatum Houtt. Nat. Hist. **14**. 94. 1783 = Nephrolepis.
alaskanum E. Britt. Bull. Torr. Cl. **24**. 556 t. 319 f. 5. 1897. NPfl. 469 — Alaska.
alpinum Rouy, Bull. Soc. Fr. **44**. 437. 1897 — Gallia alpina.
arenarium E. Britt. Bull. Torr. Cl. **24**. 555 t. 318, 319 f. 3. 1897. NPfl. 467 — N. Jersey. N. Hampshire.
azoricum Pr. Suppl. 49. 1845 = O. vulgatum.
Bergianum Schlecht. Adumbr. 10. 1825. HB. 477. Prantl 310 t. 7 f. 3. Chr. 364. NPfl. 466 — Africa austr. — Rhizoglossum Pr. 1845.

Braunii Prantl, Ber. deut. bot. Ges. **1**. 351. 1883; Jahrb. Bot. Gart. **3**. 313 t. 7 f. 8. 1884. NPfl. 466 — Ins. Caboverdicae. ? Yunnan.

brevipes Bedd. Ferns S. Ind. 23 t. 72. 1863-65 = O. fibrosum.

bulbosum Michx. Fl. Bor. Am. **2**. 276. 1803 = O. crotalophoroides.

bulbosum Bedd. Ferns br. Ind. Suppl. 28. 1876 = O. fibrosum.

californicum Prantl, Ber. deut bot. Ges. **1**. 351. 1883; Jahrb. bot. Gart. **3**. 315 t. 7 f. 11. 1884. NPfl. 467 — California-Mexico.

capense Sw. Schrad. Journ. **1801**[2]. 308. 1803 — Afr. austr.

O. nudicaule Thbg. Prod. Fl. cap. 171. 1800 (vix L.fil. 1781).

capense Schlecht. Adumbr. 9 t. 1 f. 2. 1825 { α regulare = O. regulare. β nudicaule = O. capense.

circinnatum Burm. Fl. Ind. 228. 1768 = Lygodium.

cognatum Pr. Suppl. 53. 1845 = O. pedunculosum.

concinnum Brack. Expl. Exp. **16**. 315 t. 44 f. 1. 1854 — Ins. Hawaii.

cordifolium Roxb.; Wall. List n. 47. 1828; Roxb. Calc. Journ. **4**. 475. 1844 = O. pedunculosum.

coriaceum A. Cunn. in Hk. Comp. Bot. Mag. **2**. 361. 1836; Prantl 314 t. 7. f. 9, 10, 15. NPfl. 466 — Australia. Tasmania. N. Zealand. Bolivia.

O. gramineum R.Br. 1810 (non Willd. 1802).

costatum R.Br. Prod. Fl. N. Holl. 163. 1810. (Chr. 364 pt.) — Austr. N. Zealand.

crotalophoroides Walt. Fl. Carol. 256. 1788. Prantl 323 t. 8 f. 25. NPfl. 467 Carolina. Alabama-Andes-Chile-Argentina. — O. bulbosum Michx. 1803. HB. 445. Chr. 363. O. tuberosum Hk. et Arn. 1832. O. stipatum Colla 1836.

Cumingianum Pr. Suppl. 52. 1845 = O. pedunculosum.

cuspidatum Milde, Bot. Zeit. **1864**. 107 = O. capense et O. vulgatum.

decipiens Poirault, Ann. sc. nat. VII. **18**. 146. 1893. NPfl. 467 — Mexico.

Dietrichiae Prantl, Ber. deut. bot. Ges. **1**. 352. 1883; Jahrb. bot. Gart. **3**. 320 t. 8 f. 18. 1884. NPfl. 467 — Queensland.

ellipticum Hk. et Grev. Ic. Fil. t. 40 A. 1828. Prantl 324 t. 8 f. 27. 28. NPfl. 468 — Amer. trop.

ellipticum Welw.; Carr. Cat. Welw. Pl. **2**[2]. 279. 1901 = O. Gomezianum.

elongatum A. Cunn. in Hk. Comp. Bot. Mag. **2**. 361. 1836 = O. pedunculosum.

Engelmanni Prantl, Ber. deut. bot. Ges. **1**. 351. 1883; Jahrb. bot. Gart. **3**. 318 t. 8 f. 17. 1884. NPfl. 467 — U. S. A.

fibrosum Schum. Vid. Selsk. Afh. **4**. 226 (seors. 452). 1827. Prantl 324 t. 8 f. 26 (pt.). NPfl. 468 — India. Africa trop.

O. Wightii Grev. et Hk. 1833. O. brevipes Bedd. 1863-65.

filiforme Roxb. Calc. Journ. **4**. 476 t. 26. 1844 = Lygodium scandens.

flexuosum L. sp. **2**. 1063. 1753 = Lygodium.

flexuosum L.fil. Suppl. 443. 1781 = Lygodium circinnatum.

furcatum Roxb. Calc. Journ. **4**. 478. 1844 = Lygodium circinnatum.

furcatum J. Sm. Ferns br. and for. 272. 1866 = O. pendulum.

Gomezianum Welw.; A.Br. apud Kuhn, Fil. Afr. 176. 1868. Prantl 315 t. 7 f. 13. NPfl. 467 — Africa trop. et austr.

O. nudicaule auctt. plur. (an L.fil. 1781?) — O. ellipticum Welw.; Carr. 1801 (non Hk. et Grev. 1828) = var. latifolium Prantl 316.

gracillimum Welw.; Bak. Syn. 445. 1868; (A.Br. in Kuhn, Fil. Afr. 177. 1868. var.) = O. gramineum var.

gramineum Willd. Schrift. Ak. Erfurt **1802**. 18 t. 1 f. 1. Prantl 311 t. 7 f. 4. NPfl. 466 — India. N. Guinea. Afr. occ. — O. gracillimum Welw. (Africa).

gramineum R.Br. Prod. Fl. Nov. Holl. 163. 1810 = A. coriaceum.
intermedium Hk. Ic. pl. t. 995. 1854. HB. 446. NPfl. 469 — Java. Borneo.
japonicum Prantl, Ber. deut. bot. Ges. 1. 353. 1883; Jahrb. bot. Gart. 3. 327 t. 8 f. 29. 1884. NPfl. 468 — Japonia. China centr.
japonicum Thbg. Fl. Jap. 328. 1784 = Lygodium.
lanceolatum Watelet, Bull. Soc. Fr. 5. 16. 1858 — Gallia. (prob. = O. vulgatum forma).
lanceolatum Prantl, Ber. deut. bot. Ges. 1. 352. 1883 = O. Prantlii.
lancifolium Pr. Suppl. 50. 1845. Prantl 327 t. 8 f. 30. NPfl. 468 — Madagascar. [Mauritius.
Loureirianum Pr. Suppl. 55. 1845 — China. Cochinchina.
O. lusitanicum Lour. Fl. Coch. 2. 825. 1790 (non L. 1753).
Luersseni Prantl, Ber. deut. bot. Ges. 1. 352. 1883; Jahrb. bot. Gart. 3. 321 t. 8 f. 20. 1884. NPfl. 467 — Australia.
lusitanicum L. sp. 2. 1063. 1753. HB. 445 (pt.). Prantl 312 t. 7 f. 6. 7. Chr. 363. NPfl. 466 — Europa atlant. Mediterr. Ins. Canar.
lusitanicum v. gracillimum A.Br. apud Kuhn, Fil. Afr. 177. 1868 = O. gramineum.
lusoafricanum Welw.; Prantl, Ber. deut. bot. Ges. 1. 351. 1883; Jahrb. bot. Gart. 3. 312 t. 7 f. 5. 1884. NPfl. 466 — Africa occ.
macrorrhizum Kze. Anal. 2. 1837; Farnkr. 1. 57 t. 29 f. 1. 1840. Prantl 322 t. 8 f. 22. NPfl. 467 — Brasilia austr. Argentina.
melipillense Remy; Gay, Fl. Chil. 6. 541. 1853 — Chile.
microstichum Ach. Vet. Ak. Handl. 1899. 59 t. 3 A = O. vulgatum.
minimum (Hk.fil.) Armstr. Tr. N. Zeal. Inst. 13. 342. 1881 — N. Zealand.
O. vulgatum ε minimum Hk.fil. Fl. N. Handb. Fl. N. Zeal. 386. 1864.
moluccanum Schlecht. Adumbr. 9. 1825 = O. pedunculosum.
nudicaule L.fil. Suppl. 443. 1781 — Africa austr. — (an = O. Gomezianum?).
nudicaule auctt. = spec. plur. Prantlianae. (vide Prantl).
obovatum Miq. Ann. Lugd. Bat. 4. 93. 1868-69 — Timor.
opacum R.Br.; Carm. Tr. Linn. Soc. 12. 509. 1818. Prantl 324. NPfl. 467 — Tristan d'Acunha. St. Helena.
ovatum Salisb. Prod. 401. 1796 (nomen); Opiz, Kratos 4. —. 1819 = O. vulgatum.
ovatum Bory, Voy. 2. 206. 1804. Prantl 327 t. 8 f. 31. NPfl. 468 — Ins. Mascar.
O. sarcophyllum Desv. 1827.
palmatum L. sp. 2. 1062. 1753. HB. 446. Chr. 364. Prantl 332. NPfl. 469 — Florida. Amer. trop. Réunion. Madagascar. — Cheiroglossa Pr. 1845.
parvifolium Grev. et Hk. Bot. Misc. 3. 218. 1833 — Ind. austr. Malacca.
pedatum Burm. Fl. Ind. 227. 1768 = Lygodium circinnatum.
pedunculosum Desv. Berl. Mag. 5. 306. 1811. Prantl 328 t. 8 f. 33. 34. NPfl. 469 — Japonia. Asia trop. Austr. N. Zeal. N. Caledonia.
O. moluccanum Schlecht. 1825. O. petiolatum Hk. 1825 — O. elongatum A. Cunn. 1836. O. reticulatum auctt. quoad. pl. asiat.
pendulum L. sp. ed. II. 2. 1518. 1763. HB. 446. Prantl 331. Chr. 364. NPfl. 469 Asia. Austr. Polynesia trop. — Ophioderma Pr. 1845.
pennatum Lam. Fl. franç. 1. 9. 1778 = Botrychium lunaria.
peruvianum Pr. Suppl. 52. 1845 = O. reticulatum.
petiolatum Hk. Exot. Flora 1 t. 56. 1823 = O. pedunculosum.
polyphyllum A.Br. apud Seubert, Fl. Azor. 17. 1844 = O. vulgatum var.
Prantlii C. Chr. Ind. 471. 1906 — Australia. — O. vulgatum v. lanceolatum Luerss. Journ. Mus. Godeffr. 3. 247 c. figg. 1875; O. lanceolatum Prantl 1883. 320 t. 8 f. 19. NPfl. 467 (non Watelet 1858).

pubescens Rafinesque, Journ. de Bot. **4**. 273. 1814 — U. S. A.
pusillum Nutt. Gen. **2**. 248. 1818. NPfl. 467 — U. S. A. merid.
O. tenerum Mett.; Prantl 1883. 322 t. 8 f. 23.
regulare (Schlecht.) C. Chr. Ind. 471. 1906 — Africa austr.
O. capense *α* regulare Schlecht. Adumbr. 9 t. 1 f. 2. 1825; O. capense auctt. Prantl 316 t. 7 f. 14. NPfl. 467 (non Sw. 1803).
reticulatum L. sp. **2**. 1063. 1763. HB. 446 (pt.). Chr. 363. Prantl 330 t. 8 f. 32. NPfl. 469 — Africa trop. (Caboverd.-Mascar.). Amer. trop. Samoa. Ins. Carolinae.
reticulatum auctt. quoad. pl. asiat. = O. pedunculosum.
rubellum Welw.; A.Br. apud Kuhn, Fil. Afr. 179. 1868. HB. 445. Prantl 321 t. 8 f. 21. NPfl. 467 — Africa occ.
sarcophyllum Desv. Prod. 193. 1827 = A. ovatum.
scandens L. sp. **2**. 1063. 1753 = Lygodium.
scandens Forst. Prod. 78. 1786 = Lygodium reticulatum.
scandens Vell. Fl. flum. **11** t. 53. 1827; Arch. Mus. Nac. Rio Jan. **5**. 443. 1881 = Lygodium volubile.
Schmidii Kze. Linn. **24**. 246. 1851 — India austr.
simplex Ridley; Bower, Nature **64**. 617. 1901; Ann. Bot. **18**. 205 t. 15. 1904 — Sumatra.
Spruceanum Fée, Cr. vasc. Br. **1**. 218 t. 52 f. 3. 1869 = O. ypanemense.
stipatum Colla, Mem. Ac. Torino **39**. 52 t. 75 f. 1. 1836 = O. crotalophoroides.
tenerum Mett.; Prantl, Ber. deut. bot. Ges. **1**. 352. 1883 = O. pusillum.
timorense Miq. Ann. Lugd. Bat. **4**. 93. 1868-69 — Timor.
tuberosum Hk. et Arn. Bot. Beech. Voy. 53. 1832 = O. crotalophoroides.
unifolium Gilib. Exerc. Phytol. **2**. 554. 1792 = O. vulgatum.
valdivianum Phil. Linn. **33**. 306. 1865 — Chile.
vulgatum L. sp. **2**. 1062. 1753. HB. 445 (pt.). Prantl 318 t. 7 f. 12, 16. Chr. 363. NPfl. 467 — Europa. Madeira. Amer. bor. Asia occ.
O. microstichum Ach. 1809 — O. polyphyllum A.Br. 1844.
vulgatum auctt. plur. = spec. plur. Prantlianae (vide Prantl).
vulgatum var. australasiaticum Luerss. Journ. Mus. Godeff. **3**. 246 c. figg. 1875 = O. pedunculosum.
vulgatum var. lanceolatum Luerss. Journ. Mus. Godeff. **3**. 247 c. figg. 1875 = O. Prantlii.
vulgatum var. surinamense Luerss. Journ. Mus. Godeff. **3**. 247 c. figg. 1875 = O. ellipticum.
Wightii Grev. et Hook. Bot. Misc. **3**. 218. 1833 = O. fibrosum.
ypanemense Mart. Ic. Cr. Bras. 39, 130 t. 73 f. 1. 1834. Prantl 323 t. 8 f. 24. NPfl. 467 — Brasilia. Columbia. — O. Spruceanum Fée 1869.
zeylanicum Houtt. Nat. Hist. **14**. 43 t. 94 f. 1. 1783 = Leptochilus.

OPHIOPTERIS Reinw. Syll. Plant. **2**. 3. 1824 = **Oleandra.**
verticillata Reinw. l. c. = O. neriiformis.

Oreopteris J. Smith, Cat. cult. ferns 57. 1857 *(Lastrea §)* = Dryopteris.

ORMOPTERIS J. Smith, Hist. Fil. 281. 1875 = **Pellaea.**
gleichenioides J. Sm. Hist. Fil. 281. 1875.

ORNITHOPTERIS Bernhardi in Schrad. neu. Journ. **1**². 40. 1806 = **Aneimia.**
adiantifolia Bernh. Schrad. neu. Journ. **1**². 50 t. 3 f. 15 a. 1806.
cicutaria Und. Mem. Torr. Cl. **12**. 15. 1902.
hirsuta Bernh. Schrad. neu. Journ. **1**². 50 t. 3 f. 15 b. 1806.
mexicana Und. Our. nat. Ferns ed. VI. 76. 1900.
Wrightii Millsp. Field Col. Mus. Bot. Ser. **3**. 1. 1903.

ORNITHOPTERIS (Agardh) J. Smith, Hist. Fil. 297. 1875.
Pteris § Ornithopteris Agardh, Rec. Gen. Pter. 45. 1839.
aquilina J. Sm. Hist. Fil. 298. 1875 = Pteridium.
caudata J. Sm. Hist. Fil. 298. 1875 = Pteridium aquilinum*.
esculenta J. Sm. Hist. Fil. 298. 1875 = Pteridium aquilinum*.
glutinosa J. Sm. Hist. Fil. 299. 1875 = Paesia viscosa.
rugulosa J. Sm. Hist. Fil. 299. 1875 = Paesia rugosula.
scaberula J. Sm. Hist. Fil. 299. 1875 = Paesia.
ORTHOGRAMMA Presl, Epim. bot. 121. 1849 = **Blechnum.**
Gilliesii Pr. Epim. 121. 1849.
laevigata Pr. Epim. 121. 1849.
OSMUNDA Linné, (Gen. pl. ed. II. 502) Spec. plant. **2**. 1063. 1753 — (Osmundaceae Gen 3).
adiantifolia L. sp. **2**. 1065. 1753 = Aneimia.
alata Goldie, Edinb. Phil. Journ. **6**. 332. 1822 = O. cinnamomea.
asplenifolia Sav. in Lam. Enc. **4**. 652. 1797 = Aneimia adiantifolia.
aurita Sw. Prod. 127. 1788 = Aneimia.
banksiifolia (Pr.) Kuhn, Ann. Lugd. Bat. **4**. 299. 1869 — Kamtschatka. Japonia. Asia trop. — Nephrodium Pr. Rel. Haenk. **1**. 34. 1825. Plenasium Pr. 1836. Asplenium? aureum Bl. 1828; Plenasium Pr. 1836. Osmunda Presliana J. Sm. 1841. NPfl. 379 — Nephrodium? bromeliaefolium Pr. 1825; Plenasium? Pr. 1836. Osmunda Haenkeana Pr. 1845; O. zeylanica Kze. 1850. O. javanica pt. HB. 426.
barbara Thbg. Prod. Fl. Cap. 171. 1800 = Todea.
basilaris Spr. Anleit. **3**. 160. 1804 = O. Claytoniana.
bavarica Schmid in Hoppe, Bot. Taschenb. **1803**. 7 = Botrychium matricariae.
bifurcata Jacq. Coll. **3**. 282 t. 20 f. 4. 1789 = Elaphoglossum furcatum.
bipinnata L. sp. **2**. 1065. 1753 = O. cinnamomea.
bipinnata Hk. Fil. Exot. t. 9. 1857 = O. Mildei.
biternata Sav. in Lam. Enc. **4**. 650. 1797 = Botrychium.
blechnoides Zoll. Syst. Verz. 49. 1854 — Java.
borealis Salisb. Prod. 402. 1796 = Blechnum spicant.
brasiliensis Vell. Fl. flum. **11** t. 54. 1827; Arch. Mus. Nac. Rio Jan. **5**. 444. 1881 = Aneimia phyllitidis.
Brayeri Watelet, Bull. Soc. Fr. **5**. 16. 1858 = O. regalis.
capensis L. Mant. 306. 1771 = Blechnum.
capensis Pr. Suppl. 63. 1845 = O. regalis var.
caroliniana Walt. Fl. Carol. 287. 1788 = Woodwardia areolata.
cervina L. sp. **2**. 1065. 1753 = Polybotrya.
cicutaria Sav. in Lam. Enc. **4**. 650. 1797 = Botrychium virginianum.
cinnamomea L. sp. **2**. 1066. 1753. HB. 426. Chr. 332. NPfl. 379 — Amer. bor. atlant.-Mexico-Brasilia. Ind. occ. Amur. Manschuria. Japonia. Yunnan. Struthiopteris Bernh. 1801; Osmundastrum Pr. 1848. Osmunda bipinnata L. 1753 — O. imbricata Kze. 1849 (Brasilia).
Claytoniana L. sp. **2**. 1066. 1753. HB. 426. Chr. 333. NPfl. 380 — Amer. bor. atlant. Himalaya. China. — Struthiopteris Bernh. 1801; Plenasium Pr. 1848 — Osmunda interrupta Michx. 1803; Plenasium Pr. 1848.
coronaria König; Müll. Naturforsch. Halle **21**. 107 t. 3. 1785 = Platycerium.
crispa L. sp. **2**. 1067. 1753 = Cryptogramma.
deltoidea Cav. Icon. **6**. 70 t. 593 f. 1. 1801 = Aneimia tomentosa.
dichotoma Spr. Schrad. Journ. **1799**[2]. 268 = Schizaea.

discolor Forst. Prod. 78. 1786 = Blechnum.
dissecta Poir. Enc. Suppl. **4**. 233. 1816 = Botrychium.
filiculifolia L. sp. **2**. 1065. 1753 = Aneimia.
filiformis Sav. in Lam. Enc. **4**. 652. 1797 = Aneimia.
flexuosa Sav. in Lam. Enc. **4**. 652. 1797 = Aneimia.
Fraseri F. Muell. Cens. 137. 1882 = Leptopteris.
fulva Cav. Icon. **6**. 70 t. 593 f. 2. 1801 = Aneimia.
glaucescens Link, Fil. sp. 20. 1841 = O. regalis var.
gracilis Link, Hort. Berol. **2**. 145. 1833 — Brasilia.
Haenkeana Pr. Suppl. 67. 1845 = O. banksiifolia.
Hilsenbergii Grev. et Hk. in Hk. Bot. Misc. **3**. 230. 1833 = O. regalis.
hirsuta L. sp. **2**. 1064. 1753 = Aneimia.
hirta L. sp. **2**. 1064. 1753 = Aneimia.
Hügeliana Pr. Suppl 64. 1845 = O. regalis.
humilis Cav. Icon. **6**. 69 t. 592 f. 3. 1801 = Aneimia.
hymenophylloides Armstr. Tr. N. Zeal. Inst. **13**. 368. 1881 = Leptopteris.
imbricata Kze. Farnkr. **2**. 29 t. 112. 1849 = O. cinnamomea var.
interrupta Michx. Fl. Bor. Am. **2**. 273. 1803 = O. Claytoniana.
japonica Thbg. Fl. Jap. 330. 1784 = O. regalis var.
japonica Houtt. Pfl. Syst. **13**[1]. 62 t. 95 f. 1. 1786 (an etiam in Houtt. Nat. Hist. **14**. 1783?) = O. lancea.
javanica Bl. Enum. 252. 1828. HB. 426 (pt.). Chr. 333 (pt.). NPfl. 378 — Borneo. Java. Sumatra. Celebes. — Plenasium Pr. 1848.
javanica auctt. quoad pl. ex Asia orient.-Ceyl. = O. banksiifolia.
lancea Thbg. Fl. Jap 330. 1784. HB. 427. NPfl. 380 — Japonia.
Osmundastrum Pr. 1848; Osmunda japonica Houtt. 1786 (1783?).
lanceolata Gmel. Nov. Comm. Ac. Petr. **12**. 516 t. 11 f. 2. 1768 = Botrychium.
lanceolata Roxb. Calc. Journ. **4**. 479 t. 27. 1844 = Leptochilus decurrens.
leptophylla Sav. in Lam. Enc. **4**. 657. 1797 = Anogramma.
Leschenaultii Wall. List n. 51. 1828 = O. regalis.
lineata Sw. Prod. 127. 1788 = Blechnum.
Lunaria L. sp. **2**. 1064. 1753 = Botrychium.
Lunaria var. Beckeana L. Amoen. Acad. **8**. 105. 1771 = Botrychium matricariae.
lunata Salisb. Prod. 401. 1796 = Botrychium lunaria.
marginalis Sav. in Lam. Enc. **4**. 655. 1797 = Mohria caffrorum.
Matricariae Schrank, Baier. Fl. **2**. 419. 1789 = Botrychium.
matricarioides Poir. Enc. Suppl. **4**. 233. 1816 = Botrychium matricariae.
mexicana Fée, 9 mém. 43. 1857 = O. regalis var.
Mildei C. Chr. Ind. 474. 1906 — Hongkong.
O. bipinnata Hk. Fil. Exot. t. 9. 1857 (non L. 1753). HB. 426. NPfl. 380.
monticola Wall. List n. 52. 1828 = O. Claytoniana.
Moorei F. Muell. Cens. 137. 1882 = Leptopteris.
multifida Gmel. Nov. Comm. Ac. Petr. **12**. 517 t. 11 f. 1. 1768 = Botrychium virginianum.
obliqua Poir Enc. Suppl. **4**. 233. 1816 = Botrychium.
oblongifolia Cav. Icon. **6**. 69 t. 592 f. 2. 1801 = Aneimia.
obtusa Banks. et Sol.; Forst. Prod. 93. 1786 (nomen) = Blechnum Banksii.
obtusifolia Willd.; Klf. Enum. 43. 1824 = O. regalis var.
oxyodon Miq. Ann. Lugd. Bat. **3**. 182. 1867 = O. banksiifolia.
palustris Schrad. Gött. gel. Anz. **1824**. 866 = O. regalis var.

peltata Sw. Prod. 127. 1788 = Elaphoglossum.
Phyllitidis L. sp. **2**. 1064. 1753 = Aneimia.
pilosa Wall.; Grev. et Hk. in Hk. Bot. Misc. **3**. 229. 1833 = O. Claytoniana.
Plumieri Tausch, Flora **1836**². 426 = O. regalis.
pollicina Willem. in Usteri Ann. **18**. 61. 1796 = Stenochlaena.
polypodioides Sw. Prod. 127. 1788 = Blechnum.
Presliana J. Sm. JoB. **3**. 420. 1841 = O. banksiifolia.
procera Forst. Prod. 78. 1786 = Blechnum capense.
ramosa Roth, Fl. Germ. **1**. 444. 1788 = Botrychium.
regalis L. sp. **2**. 1065. 1753. HB. 427. Chr. 332. NPfl. 380 — Europa. India. Asia orient. Amer. bor. atlant Mexico.-India occ.-Uruguay. Afr. austr. cum ins. ? N. Zealand. — Struthiopteris Bernh. 1801; Aphyllocalpa Lag., Garcia et Clem. 1802 — Osmunda japonica Thbg. 1784; Osmundastrum Pr. 1848 (Asia; Afr. austr.) — Osmunda Hügeliana Pr. 1845 (India) — O. obtusifolia Willd.; Klf. 1824 (Ins. Mascar. Madagascar) — O. capensis Pr. 1845 (Afr. austr.) — O. spectabilis Willd. 1810. O. glaucescens Link 1833 (Texas) — O. palustris Schrad. 1824. O. mexicana Fée 1857 (Mexico-Brasilia) — O. Plumieri Tausch 1836 (Mediterranea).
reptans Banks et Sol.; Forst. Prod. 93. 1786 (nomen) = Blechnum filiforme.
rupestris Salisb. Prod. 402. 1796 = Cryptogramma crispa.
rutacea Poir. Enc. Suppl. **4**. 232. 1816 = Botrychium ramosum.
scandens Aubl. Hist. pl. Guian. **2**. 961. 1775 = Lygodium volubile.
speciosa Wall. List n. 50. 1828 = O. regalis var. japonica.
spectabilis Willd. sp. **5**. 98. 1810 = O. regalis var.
Spicant L. sp. **2**. 1066. 1753 = Blechnum.
spicata L. v. Buch, Beschr. Can. Ins. 189. 1825 — Madeira.
(an = Spicant L.; err. typ.?).
Struthiopteris L. sp. **2**. 1066. 1753 = Matteuccia.
superba Armstr. Tr. N. Zeal. Inst. **13**. 368. 1881 = Leptopteris.
tenella Cav. Icon. **6**. 69 t. 592 f. 1. 1801 = Aneimia.
ternata Thbg. Fl. Jap. 329 t. 32. 1784 = Botrychium.
thurifera Sw. Schrad. Journ. **1800**². 105. 1801 } = Mohria caffrorum.
thurifraga Bory, Voy. **1**. 348. 1804 }
tomentosa Sav. in Lam. Enc. **4**. 652. 1797 = Aneimia.
totta Sw. Schrad. Journ. **1800**². 105. 1801 = Todea barbara.
trifida Jacq. Coll. **3**. 281 t. 20 f. 3. 1789 = Leptochilus zeylanicus.
trifrons Bory. Voy. **3**. 163. 1804 = Stenochlaena pollicina.
Vachellii Hk. Ic. pl. t. 15. 1837. NPfl. 379 — China merid.
Plenasium Pr. 1848.
verticillata L. sp. **2**. 1065. 1753 = Aneimia.
villosa Poir. Enc. Suppl. **4**. 231. 1816 = Aneimia flexuosa*.
virginiana L. sp. **2**. 1064. 1753. } = Botrychium virginianum.
virginica Houtt. Pfl. Syst. **13**¹. 57. 1786 }
zeylanica L. sp. **2**. 1063. 1753 = Helminthostachys.
zeylanica Kze. Farnkr. **2**. 52 t. 122. 1850 = O. banksiifolia.

OSMUNDASTRUM Presl, Abh. böhm. Ges. Wiss. V. **5**. 326. 1848 = **Osmunda.**
Osmunda § Osmundastrum Pr. Suppl. 68. 1845.
cinnamomeum Pr. Abh. böhm. Ges. V. **5**. 326. 1848.
japonicum Pr. l. c. = O. regalis var.
lanceum Pr. l. c.

Osmundopteris Milde, Fil. Europae 209. 1867 *(Botrychium §)* = Botrychium (virginianum).

OTHONOLOMA Link, Hort. Berol. **2**. 41. 1833 = **Cheilanthes.**

OXYGONIUM Presl, Tent. Pterid. 117. 1836 = **Diplazium.**

alismaefolium Pr. Tent. 118 t. 4 f. 14. 1836 = D. Cumingii vel Syngramma al.

alismaefolium J. Sm. JoB. **4**. 178. 1841 = D. Cumingii.

cordifolium J. Sm. Hist. Fil. 332. 1875.

elegans J. Sm. JoB. **4**. 178. 1841 = D. fraxinifolium.

integrifolium Moore, Ind. LVI. 1857 = D. cordifolium.

ovatum Pr. Tent. 118 t. 4 f. 7. 1836 = Syngramma alismifolia.

vittaeformis J. Sm. JoB. **4**. 178 1841 = Syngramma v.

Pachychaetum Presl, Hymenoph. 16. 1843 *(Trichomanes §)* = Trichomanes.

PACHYDERRIS J. Smith msc. (Moore, Ind. CXLVI. 1857) = Dryopteris (pedata).

Pachyglossa Christ, Mon. Elaph. 20. 1899 *(Elaphoglossum §)* = Elaphoglossum.

PACHYLOMA Van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 318. 1861 = **Hymenophyllum** (marginatum).

PACHYPLEURIA Presl, Epim. bot. 98. 1849 = **Humata.**

Davallia § Pachypleuria Pr. Tent. 128. 1836.

alata Pr. Epim. 98. 1849 = Davallia a.

angustata Pr. Epim. 98. 1849 = H. heterophylla*.

intermarginalis Pr. Epim. 98. 1849 = H. Gaimardiana.

lepida Pr. Epim. 99. 1849.

parallela Pr. Epim. 98. 1849 = H. Gaimardiana.

pectinata Pr. Epim. 98. 1849.

pedata Pr. Epim. 98. 1849 = H. repens.

sessilifolia Pr. Epim. 98. 1849.

subimbricata Pr. Epim. 261. 1849 = H. repens.

trifoliata Pr. Epim. 98. 1849 = H. repens.

vestita Pr. Epim. 261. 1849.

PAESIA St. Hilaire, Voy. Distr. Diamans **1**. 381. 1833 — (Polypodiaceae Gen 87).

acclivis (Kze.) Kuhn, Chaetopt. 347. 1882 — Venezuela. Columbia.
Allosorus Kze. Farnkr. **2**. 6. 1848; Pteris Mett. 1856. Üb. Pteris 10 t. 5 (16) f. 9-12. HB. 479.

amazonica (Christ) C. Chr. Ind. 476. 1906 — Peru.
Pteris Christ, Hedwigia **44**. 364. 1905.

anfractuosa (Christ) C. Chr. Ind. 333. 1905; 476. 1906 — Costa Rica.
Gymnogramme Christ, Bull. Soc. bot. Belg. **35**. 235. 1896. Chr. 73 c. fig. NPfl. 259; Pteris Christ, Bull. Boiss. II. **5**. 257. 1905 — Gymnogramme amaurophylla Christ 1904.

aquilina Keys. Pol. Cyath. Hb. Bung. 22. 1873 = Pteridium.

glandulosa Kuhn, Chaetopt. 347. 1882 = Gymnogramma.

radula (Bak.) C. Chr. Ind. 476. 1906 — Sumatra.
Pteris Bak. JoB. **1880**. 211.

rugosula (Lab.) Kuhn, Chaetopt 347. 1882 — N. Caledonia. ? Tahiti. Luzon.
Pteris Lab. Sert. Austr. Cal. 6 t. 8. 1824; Cheilanthes Fourn. 1873. Pteris rugulosa auctt. HB. 480. ? Allosorus? Guill. 1836. ? Hypolepis Hk. 1852 (non J. Sm. 1846). ? Ornithopteris J. Sm. 1875. Paesia Diels 1899. NPfl. 297. ? Cheilanthes divaricatissima Dry. msc.; Pteris Bak. 1874. HB. 163.

rugulosa Diels, Nat. Pfl. 1^4. 297. 1899 = P. rugosula.

scaberula (A. Rich.) Kuhn, Chaetopt. 347. 1882. NPfl. 297 — N. Zealand.
Pteris A. Rich. Fl. N. Zél. 82 t. 11. 1832. Mett. Pteris n. 1. HB. 163. Chr. 164; Allosorus Pr. 1836; Ornithopteris J. Sm. 1875 — Pteris microphylla A. Cunn. 1836 (non Sw. 1806 nec Colla 1836).

scalaris Kuhn, Chaetopt. 347. 1882 = P. viscosa.

viscosa St. Hil. Voy. Distr. Diamans **1**. 381. 1833. NPfl. 297 — Jamaica. Andes. Brasilia centr. — Pteris Moore 1858. HB. 163. Chr. 164. Allosorus scalaris Moritz msc.; Pteris Moritz 1854; Mett. Pteris n. 2 t. 5; Paesia Kuhn 1882; Pteris resistens Mett. 1856. Ornithopteris glutinosa J. Sm. 1875.

PALMA FILIX Adanson, Fam. des plantes **2**. 21. 1763 —?

PALTONIUM Presl, Epim. bot. 156. 1849 — (Polypodiaceae Gen. **98**).

lanceolatum (L.) Pr. Epim. 156. 1849 — Florida. Ind. occ. Guiana. Guatemala.
Pteris L. sp. **2**. 1073. 1753; Taenitis Klf. 1824. HB. 397; Pteropsis Desv. 1827; Drymoglossum J. Sm. 1841; Neurodium Fée 1851-52. Chr. 129; Vittaria Bak. 1870; Elaphoglossum Keys. 1873; Oetosis O.Ktze. 1891; Heteropteris Diels 1899. NPfl. 305; Cheilogramma Maxon 1901 (an Und. 1900?).

sinense (Christ) C. Chr. Ind. 477. 1906 — China merid.
Neurodium Christ, Bull. Boiss. **6**. 880. 1898.

PANICULARIA Colla, Mem. Acad. Torino **39**. 33. 1836 = **Thyrsopteris.**

Berteri Colla, Mem. Ac. Torino **39**. 35 t. 64. 1836 = T. elegans.

PARABLECHNUM Presl, Epim. bot. 109. 1849 = **Blechnum.**

ambiguum Pr. Epim. 109. 1849 = B. laevigatum.

ciliatum Pr. Epim. 109. 1849.

procerum Pr. Epim. 109. 1849 = B. capense.

salicifolium Pr. Epim. 110. 1849 = B. tabulare.

Paraceterach F. Mueller, Fragmenta **5**. 138. 1866 *(Grammitis §)* = Gymnopteris.

PARAGRAMMA (Bl.) Moore, Index XXXII. 1857 = **Polypodium.**
Grammitis § Paragramma Blume, Enum. 114. 1828.

decurrens Moore, Ind. XXXII. 1857 = P. revolutum.

longifolia Moore, Ind. XXXII. 1857 = P. revolutum.

stenophylla J. Sm. Hist. Fil. 115. 1875.

Paralindsaya Keyserling, Pol. Cyath. Hb. Bung. 3. 1873 *(Lindsaya §)* = Lindsaya.

Parallosorus Keyserling, Pol. Cyath. Hb. Bung. 5. 1873 *(Cheilanthes §)* = Cheilanthes.

Paralomaria Fée, Gen. Fil. 68. 1850-52 *(Lomaria §)* = Blechnum.

Paraneura Prantl, Ber. deut. bot. Ges. **1**. 350. 1833 *(Ophioglossum §)* = Ophioglossum.

Parapolystichum Keyserling, Pol. Cyath. Hb. Bung. 11. 1873 *(Polystichum §)* = Dryopteris.

Parapteris Keyserling, Pol. Cyath. Hb. Bung. 4. 1873 *(Pteris §)* = Pteris.

Parasaccoloma Keyserling, Pol. Cyath. Hb. Bung. 4. 1873 *(Saccoloma §)* = Saccoloma.

Parasplenium Keyserling, Pol. Cyath. Hb. Bung. 15. 1873 *(Asplenium §)* = Asplenium.

Parenchymaria K. Müller, Bot. Zeit. **1854**. 545 *(Vittaria §)* = Vittaria.

PARESTIA Presl, Epim. bot. 99. 1849 = **Davallia.**

caudata Pr. Epim. 100. 1849 = D. solida.

elata Pr. Epim. 100. 1849 }
elegans Pr. Epim. 99. 1849 } = D. denticulata.
epiphylla Pr. Epim. 100. 1849 }

PARKERIA Hooker, Exotic Flora 2 t. 147. 1825 = **Ceratopteris.**

Lockharti Hk. et Grev. Ic. Fil. t. 97. 1829.

pteridoides Hk. Exot. Flora 2 t. 147. 1825 = C. thalictroides.

PATANEMA J. Smith (Moore, Index XCVIII. 1857) = **Aspidium** (Moorei).

PATANIA Presl. Tent. Pterid. 137. 1836 = **Dennstaedtia.**

adiantoides Pr. Epim. 102. 1849 = Saccoloma domingense.

appendiculata Bedd. Ferns br. Ind. Suppl. 2. 1876.

concinna Pr. Tent. 138. 1836 = D. Mathewsii.

consanguinea Pr. Epim. 102. 1849.

Elwesii Bedd. Ferns br. Ind. Suppl. 2 t. 347. 1876 = D. appendiculata var.

erosa Pr. Tent. 138 t. 5 f. 12-13. 1836 = ? D. cicutaria.

Lindeni Pr. Epim. 103. 1849 = Saccoloma domingense.

obtusifolia Pr. Tent. 138 t. 5 f. 14. 1836.

scabra Bedd. Ferns br. Ind. Suppl. 2. 1876.

triangularis Pr. Epim. 261. 1849 = D. consanguinea.

PELLAEA Link, Fil. Spec. 48, 59. 1841 — (Polypodiaceae Gen. 67).

abyssinica Fée, 6 mém. sub t. 3. 1853 = Cheilanthes coriacea.

acutiloba Prantl, Engl. Jahrb. 3. 425. 1882 = Doryopteris.

adiantoides J. Sm. Cat. Kew Ferns 4. 1856 = P. viridis.

adiantoides Prantl, Engl. Jahrb. 3. 418. 1882 = P. Boivini.

alabamensis Bak. Syn. 148. 1867 = Cheilanthes.

alcicornis Prantl, Engl. Jahrb. 3. 429. 1882 = Doryopteris ornithopus.

ambigua (Fée) Bak. Syn. 147. 1867. NPfl. 266 — Columbia. Synochlamys Fée. 7 mém. 35 t. 20 f. 4. 1857; Cheilanthes Mett. 1859 (non A. Rich. 1832 nec Brack. 1854); Allosorus O.Ktze. 1891.

andromedifolia (Klf.) Fée, Gen. 129. 1850-52. HB. 150. Chr. 159. NPfl. 267 — California-Chile (?) — Pteris Klf. Enum. 188. 1824; Allosorus Klf.; Kze. 1834; Platyloma J. Sm. 1841; Nothochlaena Keys. 1873. (Crypteris divaricata Nutt. — C. pubescens Nutt.).

andromedifolia Gay, Fl. chil. 6. 493. 1853 = P. myrtillifolia.

angulosa (Bory) Bak. Syn. ed. II. 153. 1874 — Madagascar. Ins. Mascar. Pteris Bory; Willd. sp. 5. 377. 1810; Pteridella Mett.; Kuhn 1879; Allosorus O.Ktze. 1891; Pteris articulata Klf.; Spr. 1827; Litobrochia Pr. 1836; Doryopteris Fée 1850-52; Pellaeopsis J. Sm. 1875; Pellaea Bak. Syn. ed. I. 153. 1867. NPfl. 267.

angustifolia Bak. Syn. 150. 1867 = Cheilanthes.

?arabica Fée, Gen. 130. 1850-52; 6 mém. 4 t. 3 f. 1 = Cheilanthes coriacea.

articulata Bak. Syn. ed. I. 153. 1867 = P. angulosa.

aspera Bak. Syn. 148. 1867 = P. scabra.

atropurpurea (L.) Link, Fil. sp. 59. 1841. HB. 147. Chr. 158. NPfl. 266 — Canada-U. S. A.-Mexico. — Pteris L. sp. 2. 1076. 1753; Asplenium Bernh. 1806; Allosorus Kze.; Pr. 1836; Platyloma J. Sm. 1841; Nothochlaena Keys. 1873 — Pellaea glabella Mett.; Kuhn 1869. NPfl. 266.

atropurpurea occidentalis Nelson, Fern Bull. 7. 30. 1899 = P. occidentalis.

auriculata (Thbg.) Fée, Gen. 129. 1850-52. HB. 145. Chr. 158. NPfl. 268 — Afr. austr. — Adiantum Thbg. Prod. Fl. Cap. 173. 1800; Pteris Sw. 1801; Cheilanthes Link 1833; Allosorus Pr. 1836; Cassebeera J. Sm. 1846 — Pteris confluens Thbg. 1800.

Barklyae (Hk.) Bak. Syn. 151. 1867 — Ins. Sechellae. — Cheilanthes Hk. msc.; Allosorus O.Ktze. 1891; Pteris Barkleyae Mett.; Kuhn, Fil. Afr. 77. 1868.

bella Bak. Syn. 477. 1874 = P. brachyptera var.

Boivini Hk. sp. **2**. 147 t. 118 A. 1858. HB. 149. Chr. 160 — Africa austr. cum ins. India austr. Ceylon. — Pteris Bedd. 1863; Allosorus O.Ktze. 1891; Pteris adiantoides Desv. 1827 (non Bory 1810); Pteridella Kuhn 1879; Pellaea Prantl 1882. NPfl. 267.

Bojeri Hk. sp. **2**. 145 t. 119 A. 1858 = P. quadripinnata.

Bongardiana Bak. Fl. bras. 1². 397 t. 55 f. 2. 1870 = P. flavescens.

brachyptera (Moore) Bak. Syn. 477. 1874 — California. Oregon. Platyloma Moore, Gard. Chr. **1873**. 141; Allosorus O.Ktze. 1891 — Platyloma bellum Moore 1873; Pellaea Bak. 1874. HB. 474; Allosorus O.Ktze. 1891.

brasiliensis Bak. Engl. Jahrb. **17**. 522. 1893 — Brasilia.

Breweri Eat. Pr. Amer. Acad. **6**. 555. 1865. HB. 145 — U. S. A. occ.

Bridgesii Hk. sp. **2**. 238. 1858, **3** t. 142 B. 1860. HB. 151. NPfl. 266 — California. — Platyloma J. Sm. 1875; Allosorus Luerss. 1876.

Burkeana Bak. Syn. 153. 1867 = P. dura.

calomelanos Link, Fil. sp. 61. 1841 = P. hastata.

cambodiensis Bak. Ann. Bot. **5**. 213. 1891 — Cambodia. Allosorus O.Ktze. 1891.

candida Prantl, Engl. Jahrb. **3**. 417. 1882 = P. nivea var.

chilensis Fée, Gen. 129. 1850-52 (nomen) — Juan Fernandez. Cincinalis Fée; Gay, Fl.chil. **6**. 497. 1853; Nothochlaena Sturm 1858. HB. 374.

collina Prantl, Engl. Jahrb. **3**. 428. 1882 = Doryopteris.

columbina Hk. Syn. 146. 1867 = Doryopteris.

columbina β vestita Bak. Syn. 146. 1867 = Doryopteris paradoxa.

concolor Bak. Fl. bras. 1². 396. 1870 = Doryopteris.

consobrina Hk. sp. **2**. 145 t. 117 A. 1858 = P. quadripinnata.

contracta Fée, Gen. 129. 1850-52 = P. quadripinnata var.

cordata (Cav.) J. Sm. Cat. Kew Ferns 4. 1856. HB. 152 (excl. β). NPfl. 267 — Mexico-Peru. — Pteris Cav. Descr. 267. 1802; Allosorus Pr. 1836; Platyloma J. Sm. 1841; Nothochlaena Keys. 1873 — Pteris sagittata Cav. 1802; Allosorus Pr. 1836; Platyloma J. Sm. 1841; Pellaea Link 1841. [NPfl. 266.

cordata Fée, Gen. 130. 1850-52 = P. paradoxa.

crenulans Fée, Cr. vasc. Br. **2**. 27 t. 87 f. 3. 1872-73 = Doryopteris lomariacea.

crispatula Bak. Ann. Bot. **5**. 215. 1891 = Doryopteris.

cuneata J. Sm. Ferns br. and for. 180. 1866 = Cheilanthes marginata var.

dealbata (Pursh) Prantl, Engl. Jahrb. **3**. 417. 1882 — U. S. occid. Cheilanthes Pursh, Fl. Am. sept. **2**. 671. 1814; Nothochlaena Kze. 1848. HB. 374; Cincinalis Fée 1850-52; Gymnogramme Nutt.; Mett. 1859 (non Pr. 1825); Nothochlaena pulchella Kze. 1843.

decomposita Hk. sp. **2**. 151 t. 119 B. 1858 = Cheilanthes angustifolia.

deltoidea Bak. Syn. 146. 1867 = Doryopteris.

densa (Brack.) Hk. sp. **2**. 150 t. 125 B. 1858. HB. 149. Chr. 158 — Amer. bor. pacif.-Costa Rica. — Onychium Brack. Expl. Exp. **16**. 120 t. 13 f. 2. 1854; Allosorus O.Ktze. 1891: Cryptogramme Diels 1899. NPfl. 280.

depilata Bak. Syn. 478. 1874 — Fiji. — Allosorus O.Ktze. 1891.

dichotoma Prantl, Engl. Jahrb. **3**. 420. 1882 = Adiantopsis.

Doniana (J. Sm.) Hk. sp. **2**. 137 t. 125 A. 1858. HB. 152. Chr. 160. NPfl. 267 Africa trop. Ins. Sechellae. — Platyloma J. Sm. msc.; Pteris Kuhn 1868; Pteridella Mett.; Kuhn 1879; Allosorus O.Ktze. 1891.

dura Hk. sp. **2**. 139 t. 113 A. 1858; Syn. ed. I. 147. 1867 (pt.) = P. Goudotii.

dura (Willd.) Bak. JoB. **1880**. 327. Chr. 160. NPfl. 267 — Africa austr. c. ins. Pteris Willd. sp. **5**. 376. 1810; Allosorus Pr. 1836; Litobrochia Moore 1857; Pteridella Mett.; Kuhn 1879. Pteris Burkeana Hk. 1858; Pellaea Bak. 1867. HB. 153; Pellaeopsis J. Sm. 1875.

falcata (R.Br.) Fée, Gen. 129. 1850-52. HB. 151. Chr. 160. NPfl. 266 — Asia trop.-Austr. Tasmania. N. Zealand. N. Caledonia.
Pteris R.Br. Prod. Fl. N. Holl. 154. 1810; Platyloma J. Sm. 1841; Allosorus Kze. 1850 — Pteris seticaulis Hk. 1840.

fasciata Fourn. Bull. Soc. Fr. **16**. LXIX. 1869 — Venezuela.

Fauriei Christ, Bull. Boiss. II. **4**. 612. 1904 — Formosa.

Fendleri (Kze.) Prantl, Engl. Jahrb. **3**. 417. 1882. NPfl. 267 — Wyoming-Arizona. — Nothochlaena Kze. Farnkr. **2**. 87 t. 136. 1851. HB. 374; Gymnogramme Mett. 1859.

ferruginea Nees, Linn. **19**. 654. 1847 = Notholaena bonariensis.

flavens (Sw.) C. Chr. Ind. 9. 1905; 480. 1906 — Amer centr.
Acrostichum Sw. Syn. 16, 204. 1806; Cincinalis? Desv. 1811; Gymnogramme Klf. 1824; Nothochlaena Moore 1857. HB. 374. — (an P. nivea var.?).

flavescens Fée, Cr. vasc. Br. **1**. 44 t. 22 f. 2. 1869. HB. 477. NPfl. 269 — Brasilia. — Allosorus O.Ktze. 1891; Pellaea Bongardiana (Kze.) Bak. 1870; Allosorus Kze. msc.

flavescens Fourn. Mex. pl. **1**. 119. 1872 = Doryopteris Fournieri.

flexuosa (Klf.) Link, Fil. sp. 60. 1841. Chr. 159. NPfl. 267 — N. Mexico-Peru. S. Domingo. — Pteris Klf.; Schlecht. et Cham. Linn. **5**. 614. 1830; Allosorus Klf.; Kze. 1839; Platyloma J. Sm. 1846; Pellaea cordata β HB. 153.

Fournieri Bak. Syn. 476. 1874 = Doryopteris.

fumariaefolia Phil.; Bak. Ann. Bot. **5**. 214. 1891 = Cryptogramma crispa.

geraniifolia Fée, Gen. 130. 1850-52 = Doryopteris concolor.

glabella Mett.; Kuhn, Linn. **36**. 87. 1869 = P. atropurpurea var.

glauca J. Sm. Ferns br. and for. 180. 1866 = Cheilanthes.

Glaziovii Bak. Fl. bras. 1^2. 595. 1870 = Doryopteris quinquelobata.

Glaziovii β minor Bak. Fl. bras. 1^2. 595. 1870 = Doryopteris microphylla.

gleichenioides (Gardn.) Christ, Bull. Boiss. II. **2**. 545. 1902 — Brasilia.
Cassebeera Gardn. in Hk. Ic. pl. t. 507. 1843. HB. 143. Chr. 154. NPfl. 288; Ormopteris J. Sm. 1875; Bakeriopteris O.Ktze. 1891.

Goudotii (Kze.) C. Chr. Ind. 480. 1906 — Afr. austr. cum ins.
Pteris Kze. msc. (Kuhn, Fil. Afr. 87. 1867); P. pectiniformis Godet; Mett. in Kuhn, Fil. Afr. 87. 1868 (non Goldm. 1843); Pellaea Bak. 1874. HB. 147. Chr. 158. NPfl. 267; Pteridella Mett.; Kuhn. 1879; Allosorus O.Ktze. 1891. Pellaea dura Hk. 1858 t. 113 A. (non supra); Platyloma J. Sm. 1875.

gracilis Hk. sp. **2**. 138 t. 133 B. 1858 = Cryptogramma Stelleri.

Greggii Mett.; Kuhn, Linn. **36**. 86. 1869. HB. 477 — Mexico.
Allosorus Mett. msc. (l. c.). O.Ktze. 1891.

hastata (Thbg.) Prantl, Engl. Jahrb. **3**. 418. 1882. NPfl. 267 — Abyssinia. Africa austr. Réunion. Himalaya. Yunnan. — Pteris Thbg. Prod. Fl. Cap. 172. 1800 (non Sw. 1801); Pteridella Mett.; Kuhn 1879; Pteris calomelanos Sw. 1801; Allosorus Pr. 1836; Pellaea Link 1841. HB. 152. Chr. 161; Platyloma J. Sm. 1846; Notholaena Keys. 1873.

hastata Link, Fil. sp. 60. 1841 = P. viridis.

hederacea Prantl, Engl. Jahrb. **3**. 428. 1882 = Doryopteris varians.

Henryi Christ, Bull. Boiss. 7. 7. 1899 — Yunnan.
hirsuta Hk. sp. 2. 152. 1858 = Cheilanthes glauca.
Holstii Hier.; Diels, Nat. Pfl. 1^4. 267. 1899 — Africa orient. trop.
Pteridella Hier. in Engl. Pflwelt Ostafr. C. 78. 1895.
integricuspis Fourn. Ann. sc. nat. V. 18. 324. 1873 — N. Caledonia.
Allosorus O.Ktze. 1891.
intermedia Mett.; Kuhn, Linn. 36. 84. 1869 — Texas. New Mexico. Mexico.
intramarginalis (Klf.) J. Sm. Cat. Kew Ferns 4. 1856. HB. 148. Chr. 161 — Mexico-Costa Rica. — Pteris Klf.; Schlecht. et Cham. Linn. 5. 613. 1830 (nomen); Kze. Anal. 21. t. 17 f. 1. 1837; Allosorus Pr. 1836; Cassebeera (infram.) J. Sm. 1841; Cheilanthes Hk. 1852; Platyloma Lowe 1857; Mildella Trevis. 1876 — Pteris fallax Mart. et Gal. 1842 (= β HB. 148).
involuta (Sw.) Bak. Syn. ed. II. 148. 1874 — Africa austr.
Pteris Sw. Schrad. Journ. 1800^2. 69. 1801; Allosorus? Pr. 1836; Pteridella Mett.; Kuhn 1879 — ? Cheilanthes profusa Kze. 1836 (excl. var.); Notholaena Pr. 1836; Pellaea J. Sm. 1866.
itatiaiensis Fée. Cr. vasc. Br. 2. 26 t. 88 f. 1. 1872-73 = Doryopteris lomariacea var.
Kitchingii Bak. JoB. 1880. 327 — Madagascar. — Allosorus O.Ktze. 1891.
lancifolia Bak. JoB. 1874. 199; Ann. Bot. 5. 214. 1891 — Namaqualand.
Allosorus O.Ktze. 1891.
lanuginosa Fée, 8 mém. 115. 1857 = P. ternifolia.
leucomelas (Mett.) Bak. Syn. 478. 1874 — Africa austr. — Pteris Mett.; Kuhn, Fil. Afr. 83. 1868; Pteridella Mett.; Kuhn 1879; Allosorus O.Ktze. 1891.
lomariacea Hk. sp. 2. 133. 1858 = Doryopteris.
lomariacea b columbina Hk. sp. 2. 133 t. 112 A. 1858 = Doryopteris columbina.
lomarioides Bak. Kew Bull. 1895. 229 — Somaliland.
lonchophora Bak. Fl. bras. 1^2. 397. 1870 = Doryopteris.
longimucronata Hk. sp. 2. 143 t. 115 A. 1858 = P. mucronata var.
Lorentzii Hier. Engl. Jahrb. 22. 392. 1896 = Doryopteris.
ludens Prantl. Engl. Jahrb. 3. 419. 1882 = Doryopteris.
macrophylla Fée, Gen. 129. 1850-52 = P. viridis.
marginata Bak. Syn. 151. 1867 = Cheilanthes.
membranacea Dav. Bot. Gaz. 21. 262 t. 18 f. 5-6. 1896 — Mexico.
microphylla Fée. Cr. vasc. Br. 1. 43 t. 4 f. 2. 1869 = Doryopteris.
microphylla Mett.; Kuhn, Linn. 36. 86. 1869 = P. pulchella var.
mucronata Eat. U. S. and Mex. Bound. Surv. Bot. 233. 1859 (pt.). HB. 148 — Kansas-Texas-California. — Allosorus Eat. Amer. Journ. Sci. II. 22. 138. 1856; Cheilanthes Moore 1861; Pellaea Wrightiana Hk. 1858. NPfl. 266 — P. longimucronata Hk. 1858.
mucronata Fée, 9 mém. 8. 1857 = P. atropurpurea.
myrtillifolia Mett.; Kuhn, Linn. 36. 85. 1869. HB. 477. NPfl. 267 — Chile.
Allosorus myrtifolius O.Ktze. 1891.
namaquensis Bak. JoB. 1874. 199; Ann. Bot. 5. 214. 1891 — Namaqualand.
Allosorus O.Ktze. 1891.
nitida (R.Br.?) Bak. Syn. 478. 1874 — Austr. trop.
? Pteris R.Br. Prod. Fl. N. Holl. 155. 1810; Allosorus O.Ktze. 1891.
— Pteris nitida R.Br. est verisim. Cheilanthes caudata R.Br. (t. Benth.).
nitidula (Wall.) Bak. Syn. 149. 1867 — Himalaya. Yunnan. Thibet orient.
Pteris Wall. 1828 (nomen); Allosorus Pr. 1836 (nomen); Cheilanthes Hk. sp. 2. 112. 1852.

nivea (Poir.) Prantl, Engl. Jahrb. 3. 417. 1882. NPfl. 267 — Arizona. New Mexico-Peru-Chile. — Pteris Poir. Enc. 5. 718. 1804. Cincinalis Desv. 1811; Notholaena Desv. 1813; HB. 374. Chr. 153 (excl. syn.); Gymnogramme Mett. 1859. Acrostichum albidulum Cav.; Sw. 1806; Nothochlaena Sturm 1858 — N. incana Pr. 1825 — Gymnogramme candida Mett. 1859; Pellaea Prantl 1882 — Nothochlaena chrysophylla Kl. 1855.

?nudiuscula Hk. sp. 2. 151. 1858 = Notholaena hirsuta.

occidentalis (Nelson) Rydberg, Mem. N. Y. Bot. Gard. 1. 466. 1900 — S. Dakota-Washington. — P. atropurpurea occidentalis Nelson, Fern Bull. 7. 30. 1899; P. pumila Rydb. 1900.

ornithopus Hk. sp. 2. 143 t. 116 A. 1858. HB. 149. Chr. 158. NPfl. 266 — California. — Cheilanthes Moore 1861; Allosorus O.Ktze. 1891.

pallida Bak. Syn. 478. 1874 — Mexico. — Allosorus O.Ktze. 1891.

palmescens Fée, Gen. 130. 1850-52 = Doryopteris concolor.

paradoxa (R.Br.) Hk. sp. 2. 135 t. 111 A. 1858. HB. 152. NPfl. 266 — Austr. Adiantum R.Br. Prod. Fl. N. Holl. 155. 1810; Pteris Desv. 1827; Allosorus Kze. 1850; Platyloma Brownii J. Sm. 1841.

paradoxa Fée, Cr. vasc. Br. 1. 43. 1869 = Doryopteris.

patula Prantl, Engl. Jahrb. 3. 426. 1882 = Doryopteris.

Pearcei Bak. Syn. 476. 1874 — Columbia. — Allosorus O.Ktze. 1891.

pectiniformis Bak. Syn. ed. II. 147. 1874 = P. Goudotii.

pedata Fée, Gen. 130. 1850-52 = Doryopteris.

pilosa Hk. sp. 2. 132 t. 114 B. 1858 = Doryopteris.

pinnata (Klf.) Prantl, Engl. Jahrb. 3. 418. 1882 — Brasilia. Cassebeera Klf. Enum. 217 t. 1 f. 11. 1824. HB. 142. Chr. 154. NPfl. 288; Pteris Mett. 1856; Bakeriopteris O.Ktze. 1891 — Cassebeera petiolata Fée 1857; Bakeriopteris O.Ktze. 1891.

Pringlei Dav. Gard. and Forest 4. 555 f. 88. 1891 — Mexico. Allosorus O.Ktze. 1891.

profusa J. Sm. Ferns brit. and for. 180. 1866; Bak. Syn. ed. I. 148. 1867 = P. involuta var.

pteroides (L.) Prantl, Engl. Jahrb. 3. 420. 1882. NPfl. 268 — Afr. austr. et occ. Adiantum L. Mant. 130. 1767; Cheilanthes Sw. 1806. HB. 131. Chr. 143; Cassebeera Pr. 1836; Adiantopsis Moore 1857; Choristosoria Mett.; Kuhn 1879.

pulchella (Mart. et Gal.) Fée, Gen. 129. 1850-52. HB. 150. Chr. 159. NPfl. 267 — Arizona. New Mexico-Peru. — Allosorus Mart. et Gal. Mém. Ac. Brux. 15. 47 t. 10 f. 1. 1842; Platyloma Moore 1857; Cincinalis J. Sm. 1866 — Allosorus formosus Liebm. 1849 — Pellaea microphylla Mett.; Kuhn 1869. (HB. 477) (non Fée 1869).

pumila Rydberg, Mem. N. Y. Bot. Gard. 1. 4. 1900 = P. occidentalis.

quadripinnata (Forsk.) Prantl. Engl. Jahrb. 3. 420. 1882. Chr. 161. NPfl. 269 — Africa (Ins. Caboverd.-Ins. Mascar.). — Pteris Forsk. Fl. Aegypt.-Arab. 186. 1775; Allosorus Pr. 1836; Cheilanthes Kuhn 1868; Pteridella Mett.; Kuhn 1879; Pteris consobrina Kze. 1836; Pellaea Hk. 1858. HB. 150; Allosorus Pappe et Raws. 1858 — Cheilanthes Atherstoni Hk. 1852 — C. triangula Kze. 1836 — C. contracta Kze. 1850; Pellaea Fée 1850-52; Pteris Mett. 1859; Pellaea Bojeri Hk. 1858; Cheilanthes Moore 1861 (an ad P. involutam pertinens?).

quinquelobata Fée, Cr. vasc. Br. 1. 42 t. 10 f. 1. 1869 = Doryopteris.

Raddiana Prantl, Engl. Jahrb. 3. 427. 1882 = Doryopteris.

reflexa Fourn. Bull. Soc. Fr. **16**. LXIX. 1869 — Columbia.

Regnelliana Prantl, Engl. Jahrb. **3**. 420. 1882 = Cheilanthes.

Riedelii Bak. Ann. Bot. **5**. 213. 1891 — Brasilia. — Allosorus O.Ktze. 1891.

rigida Hk. sp. **2**. 144. 1858 = Doryopteris.

robusta Hk. sp. **2**. 147. 1858 = Doryopteris.

rotundifolia (Forst.) Hk. sp. **2**. 136. 1858. HB. 151. Chr. 160. NPfl. 266 — N. Zealand. Norfolk Ins. ? N. S. Wales. — Pteris Forst. Prod. 79. 1786; Platyloma J. Sm. 1841; Allosorus Kze. 1850.

sagittata Link, Fil. sp. 60. 1841 = P. cordata var.

sagittifolia Prantl, Engl. Jahrb. **3**. 426. 1882 = Doryopteris.

scabra C. Chr. Ind. 172. 1905; 483. 1906 — Texas-New Mexico. Cheilanthes aspera Hk. sp. **2**. 111 t. 108 A. 1858 (non Klf. 1831); Pellaea Bak. 1867. HB. 148; Allosorus O.Ktze. 1891.

Schweinfurthii Hier.; Diels, Nat. Pfl. 1⁴. 267. 1899 — Africa orient. trop. Pteridella Hier. in Engl. Pflwelt Ostafr. C. 78. 1895.

Seemanni Hk. sp. **2**. 141 t. 117 B. 1858. HB. 145 — Mexico. Allosorus O.Ktze. 1891.

Skinneri Hk. sp. **2**. 141 t. 118 B. 1858 = Doryopteris.

spiculata Fée, Gen. 129. 1850-52 ≕ P. atropurpurea.

squamosa Hope et Wright, Journ. Linn. Soc. **35**. 518. 1903 = Doryopteris.

Stelleri Bak. Syn. ed. I. 453. 1868 = Cryptogramma.

subsimplex Fée, Cr. vasc. Br. **1**. 44 t. 4 f. 3. 1869 = Doryopteris.

Tamburii Hk. sp. **2**. 134 t. 129 A. 1858 = Cheilanthes argentea.

tenera (Gill.) Prantl, Engl. Jahrb. **3**. 417. 1882. NPfl. 267 — Utah. California. Arizona. Mexico. Bolivia. Chile. — Nothochlaena Gill; Hk. Bot. Mag. t. 3055. 1831. HB. 373. (Chr. 153); Cincinalis Fée 1850-52; Gymnogramme Mett. 1859.

ternifolia (Cav.) Link, Fil. sp. 59. 1841. HB. 148. Chr. 158. NPfl. 266 — Texas-Bolivia-Chile. S. Domingo. Ins. Hawaii. — Pteris Cav. Descr. 266. 1802; Platyloma J. Sm. 1846; Allosorus Kze. 1847; Cheilanthes Moore 1861; Nothochlaena Keys. 1873; Pteris subverticillata Sw. 1806; Allosorus Pr. 1836 — Pellaea Weddelliana Fée 1857; Cheilanthes Moore 1861.

triphylla Prantl, Engl. Jahrb. **3**. 418. 1882 = Doryopteris.

tripinnata Bak. Journ. Linn. Soc. **25**. 350. 1889 — Madagascar. Allosorus O.Ktze. 1891.

vestita Prantl, Engl. Jahrb. **3**. 423. 1882 = Doryopteris paradoxa.

viridis (Forsk.) Prantl, Engl. Jahrb. **3**. 420. 1882. NPfl. 269 — Africa. (Ins. Cabo-verd.-Abyssinia-Ins. Mascar.). — Pteris Forsk. Fl. Aegypt.-Arab. 186. 1775; Adiantum Vahl 1794; Cheilanthes Sw. 1806; Pteridella Mett.; Kuhn 1879; Allosorus O.Ktze. 1891; Adiantum hastatum L.fil. 1781; Pteris Sw. 1801 (non Thbg. 1800); Cheilanthes Kze. 1836; Allosorus Pr. 1836; Pellaea Link 1841. HB. 152. Chr. 160 (non Pellaea supra); Cassebeera J. Sm. 1841; Platyloma Lowe 1857. Pteris adiantoides Bory; Willd. 1810; Allosorus Pr. 1836; Pellaea J. Sm. 1856; Platyloma Lowe 1857 — Pteris hastaefolia Schrad. 1818; Cheilanthes Kze. 1850 — C. macrophylla Kze. 1850.

Weddelliana Fée, 8 mém. 74. 1857 = P. ternifolia var.

Wrightiana Hk. sp. **2**. 142 t. 115 B. 1858 = P. mucronata.

Zippelii (Miq.) Bak. Syn. 477. 1874 — N. Guinea. Allosorus Miq. Ann. Lugd. Bat. **4**. 98. 1868-69.

Pellaeastrum Fournier, Ann. sc. nat. V. **18**. 331. 1873 (*Lindsaya* §) = Schizoloma.

PELLAEOPSIS J. Smith, Hist. Fil. 289. 1875 = **Pellaea.**
articulata J. Sm. Hist. Fil. 290. 1875 = P. angulosa.
Burkeana J. Sm. Hist. Fil. 290. 1875 = P. dura.
PELTAPTERIS Link, Fil. spec. 147. 1841 = **Elaphoglossum.**
peltata Link, Fil. sp. 147. 1841.
PELTOCHLAENA Fée. Gen. Fil. 289. 1850-52 = **Polystichum.**
nephrodioides Fée, Gen. 289. 1850-52 = P. abbreviatum*.
PERANEMA Don, Prod. Fl. Nepal. 12. 1825 — (Polypodiaceae Gen. 2).
aspidioides Mett. Fil. Lechl. **2**. 33. 1859 = Diacalpe.
cyatheoides Don, Prod. Fl. Nepal. 12. 1825. Chr. 286. NPfl. 160 — Himalaya. India austr. — Sphaeropteris barbata Wall. 1828. HB. 49.
PERICOPTIS Wallich, List sub n. 92. 1828 = **Schizoloma.**
PERIPTERIS Rafinesque, Journ. Phys. **89**. 262. 1819 = **Pteris.**
Perrinia Hooker, Spec. Fil. **1**. 62. 1844 *(Woodsia §)* = Woodsia.
PHANEROPHLEBIA Presl, Tent. Pter. 84. 1836 — (Polypodiaceae Gen. 18).
auriculata Und. Bull. Torr. Cl. **26**. 212 t. 359 f. 3, 4 t. 360 f. 2. 1899 — Texas-Arizona-Mexico bor.
aurita Fée. Cr. vasc. Br. **2**. 70 t. 100 f. 1. 1872-73 — Brasilia.
Aspidium Bak. 1891.
guatemalensis Und. Bull. Torr. Cl. **26**. 214. 1899 — Guatemala.
juglandifolia (H.B.Willd.) J. Sm. JoB. **4**. 187. 1841 — Mexico-Venezuela.
Polypodium H.B.Willd. sp. **5**. 195. 1810; Amblia Pr. 1836; Aspidium Kze.; Kl. 1847. Mett. n. 79. HB. 257. Chr. 233; Cyrtomium Moore 1857; Dryopteris O.Ktze. 1891; Polystichum Diels 1899. NPfl. 193.
Lindeni Fourn. Mex. pl. **1**. 100 t. 4. 1872 — Mexico.
macrosora (Bak.) Und. Bull. Torr. Cl. **26**. 213. 1899 — Costa Rica.
Aspidium Christ 1904; A. juglandifolium var. macrosorum Bak. JoB. **1887**. 25.
nobilis (Schlecht. et Cham.) Pr. Tent. 85 t. 2 f. 19. 1836 — Mexico.
Aspidium Schlecht. et Cham. Linn. **5**. 610. 1830; Mett. n. 82; Cyrtomium Moore 1857.
pumila (Mart. et Gal.) Fée, Gen. 282. 1850-52 — Mexico.
Aspidium Mart. et Gal. Mém. Ac. Brux. **15**. 64 t. 17 f. 1. 1842.
remotispora Fourn. Mex. pl. **1**. 100. 1872 — Mexico. Guatemala.
(serrata vide Amblia).
umbonata Und. Bull. Torr. Cl. **26**. 211 t. 359 f. 3, t. 360 f. 4. 1899 — Mexico bor.
PHEGOPTERIS (Presl) Fée, Gen. Fil. 242. 1850-52 = **Dryopteris.**
Polypodium § Phegopteris Pr. Tent. 179. 1836.
abbreviata Mett. Pheg. u. Asp. 23 n. 45. 1858 = ? D. brachyodus.
adenochrysa Fée, Gen. 245. 1850-52 = D. concinna.
adnata Fée, Cr. vasc. Br. **1**. 103. 1869 = D. grandis *macroptera.
alloeoptera Mett. Pheg. u. Asp. 19 n. 34. 1858.
alpestris Mett. Fil. Lips. 83. 1856 = Athyrium a.
alsophiloides Fourn. Ann. sc. nat. V. **18**. 289. 1873.
amaurophylla Christ, Bull. Boiss. **7**. 14. 1899.
ammifolia Fée, Gen. 243. 1850-52 = Polystichum aculeatum var. 1.
ampla Fée, Gen. 243. 1850-52.
amplificata Fée, 8 mém. 89. 1857.
Andreana Christ, Prim. Fl. Costaric. **3**. 35. 1901.
angelicaefolia Mett.; Kuhn, Fil. Afr. 121. 1868 = Aspidium a.
Aneitensis Kuhn, Verh. zool. bot. Ges. **19**. 577. 1869.
angustifolia Mett. Fil. Lechl. **2**. 22. 1859.

anomala Mett.; Salom. Nom. 273. 1883 = Polystichum aculeatum var. 37.
appendiculata J. Sm. Hist. Fil. 233. 1875 = D. squamaestipes.
appendiculata Mett.; Salom. Nom. 273. 1883.
aquilina Mett.; Kuhn, Fil. Afr. 121. 1868.
arborescens Mett. Fil. Lechl. 2. 24. 1859 = D. reticulata*.
aspidioides Mett. Fil. Lips. 82 t. 17 f. 1-4. 1856.
aspidioides Fée, Gen. 243. 1850-52 = ? D. tristis.
aspleniifolia Fée, Gen. 243. 1850-52 = Polypodium a.
asplenioides Mett. Fil. Lips. 82. 1856 = D. aspidioides.
asterothrix Mett. Pheg. u. Asp. 21 n. 40. 1858 = D. reptans*.
aureo-vestita Fée, 11 mém. 52. 1866 = D. vellea.
auriculata J. Sm. Hist. Fil. 233. 1875 = D. squamaestipes.
aurita J. Sm. Cat. cult. Ferns 17. 1857.
axillaris Fée, Gen. 243. 1850-52 = Alsophila leucolepis.
barbata Mett. Pheg. u. Asp. 24 n. 48. 1858 = D. ferox.
Barberi Mett.; Kuhn, Linn. **36**. 125 in obs. 1869 = Aspidium B.
Barteriana Mett.; Kuhn, Fil. Afr. 121. 1868.
biformis Mett.; Kuhn, Fil. Afr. 121. 1868
bivestita Mett. Pheg. u. Asp. 27 n. 57. 1858.
Blanchetiana Fée, Gen. 245. 1850-52 — Brasilia.
blanda Fée, 8 mém. 91. 1857.
blepharodes Fée, 8 mém. 89. 1857 — Réunion.
Boivini Mett.; Ettingsh. Farnkr. 168 t. 111 f. 4. 1865 = Aspidium puberulum.
Boryana Mett.; Ettingsh. Farnkr. 162 t. 98 f. 2, t. 102 f. 2, t. 103 f. 4. 1865 [= D. punctata.
brachyodus Mett. Pheg u. Asp. 21 n. 38. 1858.
brevinervis Fée, 10 mém. 32. 1865 = ? D. villosa.
brevinervis Fée, Cr. vasc. Br. **1**. 243 t. 77 f. 2. 1869 = D. tijuccana.
Brongniartii Mett. Pheg. u. Asp. 31 n. 66. 1858 = Aspidium B.
brunea J. Sm. Hist. Fil. 233. 1875 = D. brunnea.
caespitosa Fourn. Mex. pl. **1**. 89. 1872.
calcarea Fée, Gen. 243. 1850-52 = D. Robertiana.
camptocavlon Fée. Cr. vasc. Br. **2**. 60 t. 98 f. 1. 1872-73.
cana Mett. Ann. Lugd. Bat. **1**. 223. 1864 } = D. subincia*.
canescens Mett. Pheg. u. Asp. 30 n. 64. 1858 } = D. subincia*.
? canescens Mett. Ann. Lugd. Bat. **1**. 223. 1864.
caudata Fée, Gen. 243. 1850-52.
cheilosora Mett.; Salom. Nom. 273. 1883 = D. diplazioides.
ciliata Fée, Gen. 248. 1850-52 — Cuba.
cochleata Mett. Pheg. u. Asp. 11 n. 6. 1858 = Polystichum aculeatum var. 26.
concinna Fée, Gen. 243. 1850-52.
connexa Fée, Gen. 243. 1850-52.
cordata Fée, Gen. 244. 1850-52.
crenata Mett. Fil. Lips. 84. 1856 = D. Poiteana.
crenulato-serrulata Mak. Bot. Mag. Tok. **17**. 78. 1903.
crinalis Mann, Pr. Amer. Ac. **7**. 218. 1868.
crinita Fée, Gen. 243. 1850-52.
cruciata Mett.; Kuhn, Fil. Afr. 122. 1868.
ctenoides Fée, 11 mém. 54 t. 14 f. 2. 1866.
Cunninghami Mett. Fil. Lips. 84 t. 17 f. 5-6. 1856 = D. pennigera.
cuspidata Mett. Pheg. u. Asp. 25 n. 52. 1858 = ? D. urophylla.

cyatheaefolia Mett.; Kuhn, Fil. Afr. 122. 1868.
Dahlii Hier. Notizbl. bot. Gart. Berl. **2**. 84. 1898 = Aspidium D.
Davidi Bedd. Handb. Suppl. 82. 1892 = Athyrium D.
debilis Mett. Ann. Lugd. Bat. **1**. 223 t. 6 f. 1. 1864.
decurrenti-alata Christ, Farnkr. 274. 1897.
decursive-pinnata Fée, Gen. 242 t. 20 A. f. 1. 1850-52.
decussata Mett. Fil. Lips. 83 t. 17 f. 8. 1856.
deflexa Mett. Ann. sc. nat. V. **2**. 241. 1864 = D. Lindigii.
delicatula Fée, 11 mém. 51 t. 20 f. 1. 1866 — Guadeloupe.
denticulata Fée, Cr. vasc. Br. **1**. 100 t. 32 f. 2. 1869 — Brasilia.
dicksonioides Mett.; Kuhn, Linn. **36**. 118. 1869 = D. glabra.
dictyophylla Mett.; Kuhn, Linn. **36**. 106. 1869 = Polystichum d.
difformis Mett. Fil. Lips. 84 t. 25 f. 8. 1856 = Aspidium irregulare.
dilatata Fée, 11 mém. 52 t. 14 f. 1. 1866 = D. subincisa.
diplazioides Mett. Ann. sc. nat. V. **2**. 241. 1864.
distans Mett. Pheg. u. Asp. 16 n. 23. 1858 = D. brunnea.
divergens Fée, Gen. 243. 1850-52 = D. effusa*.
diversifolia Kuhn, v. Deck. Reis. **3**^{3}. Bot. 44. 1879 = D. vivipara.
draconopteron Mett. Ann. sc. nat. V. **2**. 243. 1864 = Aspidium d.
drepanum J. Sm. Cat. cult. Ferns 16. 1857 = Polystichum.
Dryopteris Fée, Gen. 243. 1850-52 = D. Linnaena.
dubia Karst. Fl. Col. **1**. 169 t. 84. 1861 = Polystichum d.
Duchassaigniana Fée, 11 mém. 57 t. 14 f. 3. 1866 = D. diplazioides.
effusa Fée, Gen. 243. 1850-52.
elata Fée, Gen. 247. 1850-52 = D. Boryana.
elongata Fourn. Mex. pl. **1**. 89. 1872 — Mexico.
elongata J. Sm. Hist. Fil. 233. 1875 = D. khasiana.
emarginata Fée; Salom. Nom. 274. 1883 = D. subobliquata.
epieroides Fée, Gen. 248. 1850-52 = D. villosa.
eriopodia Fée, Cr. vasc. Br. **1**. 102 t. 31 f. 1. 1869 = D. grandis *macroptera.
erubescens J. Sm. Hist. Fil. 233. 1875.
eximia Mett.; Kuhn, Linn. **36**. 107. 1869 = Polystichum aculeatum var. 30.
extensa Mett.; Salom. Nom. 274. 1883 = D. subincisa.
falcata Mett. Fil. Lechl. **2**. 23. 1859 = D. reticulata*.
falciculata Fée, Cr. vasc. Br. **1**. 99. 1869.
faucia Salomon, Nom. 275. 1883 = D. obliterata.
ferox Mett. Ann. Lugd. Bat. **1**. 224. 1864.
ferruginea Mett. Ann. Lugd. Bat. **1**. 224 t. 6 f. 2. 1864 = Aspidium Zippelianum.
flagellaris Mak. Bot. Mag. Tok. **9**. —. 1895 = Polystichum f.
flavopunctata Fée, Gen. 243. 1850-52 = D. rotundata.
flexilis J. Sm. Cat. cult. Ferns 16. 1857 = Athyrium alpestre.
fluminensis Fée. Cr. vasc. Br. **1**. 97. 1869 = D. deflexa.
formosa Fée, Gen. 243. 1850-52 = D. grandis.
fragilis Kuhn, v. Deck. Reis. **3**^{3}. Bot. 66. 1879.
fulgens Mett.; Schenck, Hedw. **35**. 166. 1896 (nomen) — Brasilia.
Galeottii Fée. Gen. 243. 1850-52.
Galeottii A.Br.: Mett. Fil. Lechl. **2**. 26. 1859 = D. subincisa.
Germaniana Fée, 11 mém. 55 t. 13 f. 2. 1866.
gigantea Mett. Fil. Lechl. **2**. 22. 1859 = D. simplicifrons.
glanduligera Fourn. Mex. pl. **1**. 90. 1872 (err.) = D. glandulifera.

gracilis Mett. Pheg. u. Asp. 17 n. 27. 1858 = D. heteroclita.
grandis Mett. Novara's Reise Bot. **1**. 219. 1870.
grandis Salomon, Nom. 275. 1883 = Diplazium striatum.
grossa Christ, Bull. Boiss. **7**. 13. 1899.
Hasseltii Mett. Pheg. u. Asp. 13 n. 14. 1858.
hastaefolia J. Sm. Cat. cult. Ferns 16. 1857 = D. sagittata.
Helenensis Kuhn, Fil. Afr. 122. 1868 = D. punctata.
Helliana Fée, Gen. 247. 1850-52.
heterocarpa Fée, Cr. vasc. Br. **1**. 100 t. 30 f. 2. 1869 = D. tijuccana.
heteroclita Kuhn, Engl. Jahrb. **24**. 133. 1897 (syn.).
heterosora Salom. Nom. 275. 1883 = Aspidium vastum var.
hexagonoptera Fée, Gen. 243. 1850-52.
Hillebrandii Luerss. Flora **1875**. 436 = D. honolulensis.
Hillii Luerss.; Salom. Nom. 275. 1883.
hirsuta Fée, Gen. 248. 1850-52 = D. villosa an D. subincisa?
hirsuta Fée, Cr. vasc. Br. **2**. 61. 1872-73 = D. hirtula.
hirta Christ, Bull. Boiss. **6**. 195. 1898 = D. setosa.
honesta Mett. Pheg. u. Asp. 28 n. 59. 1858.
Honolulensis Mann, Proc. Amer. Acad. **7**. 218. 1868.
hydrophila Fée, 11 mém. 56 t. 13 f. 3. 1866.
impressa Fée, 8 mém. 90. 1857 = D. rudis.
Imrayana J. Sm. Hist. Fil. 233. 1875 = D. rotunda var.
?inaequalis Fée, 8 mém. 91. 1857 = D. excelsa.
incisa Keys. Pol. Cyath. Hb. Bung. 51. 1873 = Histiopteris i.
incrassata Christ, Bull. Boiss. **6**. 963. 1898.
irregularis Mett. Ann. Lugd. Bat. **1**. 225. 1864 = Aspidium i.
irregularis Luerss. Fil. Graeff. 172. 1871 = Aspidium macrodon.
Karsteniana Mett. Pheg. u. Asp. 30 n. 63. 1858 = D. subincisa*.
Kennedyi Luerss. Journ. Mus. Godeffr. **3**. 18. 1873 = D. urophylla.
Keraudreniana Mann, Pr. Amer. Ac. **7**. 218. 1868.
Kingi Bedd. Handb. Suppl. 84. 1892.
Krameri Mak. Phan. et Pter. Jap. Ic. t. 30. 1899 = D. oyamensis.
lachnopodia J. Sm. Cat. cult. Ferns 16. 1857 = D. ampla.
laevigata Mett.; Kuhn, Linn. **36**. 112. 1869.
lanata Fée, Gen. 246. 1850-52 = D. cruciata.
laserpitiifolia Bedd. Handb. Suppl. 84. 1892.
lasiernos Mett. Pheg. u. Asp. 28 n. 58. 1858 = D. hirtula.
latealata Christ. Ann. Jard. Bot. Genève **3**. 36. 1899.
Lechleri Mett. Fil. Lechl. **2**. 25. 1859 = D. subincisa*.
leptoptera Fée, Gen. 244. 1850-52 — Haïti.
lineata Mett.; Salom. Nom. 276. 1883 = D. penangiana.
Linkiana Mett. Fil. Lips. 82. 1856 = D. diplazioides.
lonchodes Fée, 11 mém. 52. 1866.
lunulata Fée, Gen. 243. 1850-52 = Alsophila extensa.
luxurians Mett. Pheg. u. Asp. 25 n. 51. 1858 = D. prolifera.
macrodonta Mett. Pheg. u. Asp. 31 n. 68. 1858 = Aspidium m.
macrophylla Mett. Fil. Lechl. **2**. 22. 1859 = D. reticulata var.
macrophylla J. Sm. Hist. Fil. 233. 1875 = D. pellucido-punctata.
macroptera Fée, Gen. 243. 1850-52 = D. grandis.
macrosora Fée, Gen. 243. 1850-52 = ? D. subincisa.

macrotheca Fée, 11 mém. 56. 1866 = ? D. Germaniana.
manipurensis *Bedd. Handb.* Suppl. 83. 1892.
Manniana Kuhn, Fil. Afr. 123. 1868.
marginans Fée, Cr. vasc. Br. **1.** 104 t. 61 f. 1. 1869 = D. villosa an D. subincisa?
Martinicensis Fourn. Mex. pl. **1.** 90. 1872 = D. subincisa.
Maximowiczii Christ, Farnkr. 271. 1897 = Polystichum M.
megalocarpa Salom. Nom. 276. 1883 = Aspidium m.
megalodus Mett. Pheg. u. Asp. 24 n. 47. 1858.
? *melanorachis* Fée, 8 mém. 91. 1857 — Mexico.
membranacea Mett. Fil. Lechl. **2.** 22. 1859 = D. reticulata.
meniscioides Ettingsh. Farnkr. 167 t. 109 f. 3, t. 110 f. 2. 1865 = D. prolifera.
meniscioides Mett.; Salom. Nom. 276. 1883.
microdendron Eat.; Mann, Pr. Amer. Ac. 7. 218. 1868 = D. stegnogrammoides.
Mischlichii Christ, Farnkr. 274. 1897 = D. obtusiloba var.
mollicula J. Sm. Cat. cult. Ferns 16. 1857 = D. concinna.
mollis Mett. Ann. sc. nat. V. **2.** 242. 1864 = D. reticulata*.
mollis Kuhn, Fil. Afr. 123. 1868 = D. Dianae.
mollivillosa Fée, 10 mém. 32. 1865 = D. villosa an D. subincisa?
Montbrisoniana Fée, Gen. 247. 1850-52 — Réunion.
montevidensis Mett.; Salom. Nom. 276. 1883 = Polystichum aculeatum var. 9.
multilineata Luerss. Fil. Graeff. 173. 1871 = D. moulmeinensis.
munita Mett. Pheg. u. Asp. 14 n. 18. 1858 = D. rufescens.
neglecta Mett.; Kuhn, Linn. **36.** 124. 1869 = Aspidium n.
nervosa Fée, Gen. 244. 1850-52; 6 mém. 13 t. 2 f. 4 = {D. philippina vel Polystichum auriculatum var.}
nicaraguensis Fourn. Bull. Soc. Fr. **19.** 252. 1872.
nitens Fée, Gen. 246. 1850-52 = Polystichum n.
obliterata Mett. Pheg. u. Asp. 23 n. 46. 1850-52.
obscura Fée, Gen. 243. 1850-52 = D. sagenioides var.
obtusata Christ, Farnkr. 274. 1897 = D. opaca.
obtusiloba Christ, Farnkr. 273. 1897.
oligophlebia Salom. Nom. 276. 1883.
opaca Mett. Pheg. u. Asp. 15 n. 21. 1858.
opaca Christ, Farnkr. 270. 1897 = D. Christii.
oppositifolia Kuhn, Fil. Afr. 123. 1868.
Oreopteridastrum Fée, Cr. vasc. Br. **1.** 97. 1869 = D. submarginalis.
Oreopteris Fée, Gen. 243. 1850-52.
ornata Fée, Gen. 243. 1850-52.
organensis Fée, Cr. vasc. Br. **1.** 105. 1869 — Brasilia.
paludosa J. Sm. Hist. Fil. 233. 1875 = D. brunnea.
palustris Mett. Fil. Lechl. **2.** 24. 1859 = D. serrata.
pauciflora Mett.; Kuhn, Fil. Afr. 123. 1868.
Pavoniana Mett.; Kuhn, Linn. **36.** 112 (in obs.). 1869.
perforata Fée, Gen. 248. 1850-52 — ?
persicaefolia Christ in Warb. Monsunia **1.** 82. 1900 = ? Polypodium p.
petrophya Mett. Ann. Lugd. Bat. **1.** 225. 1864 = Aspidium saxicola.
Phegopteris Keys. Pol. Cyath. Hb. Bung. 50. 1873.
philippinensis Mett. Pheg. u. Asp. 26 n. 55. 1858. var. 1 = Aspidium ambiguum. var. 2 = D. sagenioides* obscura.
pilosa Fée, 8 mém. 127. 1857 = D. rudis.

pilosa Mett. Pheg. u. Asp. 18 n. 30. 1858.
pilosula Fée, 8 mém. 90. 1857 = D. oligocarpa.
platyphylla Mett. Pheg. u. Asp. 12 n. 9. 1858 = Polystichum aculeatum var. 23.
? *Plumierii* J. Sm. Bot. Voy. Herald 228. 1854 — Ind. occ.
plumosa J. Sm. Ferns br. and for. 289. 1866 = Athyrium filix femina.
poecilophlebia Luerss. Journ. Mus. Godeffr. **3**. 18. 1873.
Poeppigii Fée; Gay, Fl. Chil. **6**. 506. 1853 = D. punctata.
polycarpa Mett.; Kuhn, Linn. **36**. 124. 1869 = Aspidium Kuhnii.
polycarpa Hill. Fl. Haw. 560. 1888 = D. stegnogrammoides.
polypodioides Fée, Gen. 243. 1850-52 = D. phegopteris.
polypodioides Mett. Fil. Lips. 82. 1856.
polystichiformis Fée, Gen. 243. 1850-52 = Polystichum aculeatum var. 23.
polystichops Wawra, Maxim. Reise 201 t. 102. 1866 = Polystichum aculeatum
Portoricensis Fée, Gen. 243. 1850-52 = D. pubescens var. [var. 23.
Prionitis Fée, Gen. 243. 1850-52.
procera Mann, Pr. Amer. Ac. **7**. 218. 1868 = D. Keraudreniana var.
procurrens Mett.; Salom. Nom. 277. 1883 = D. pilosa var.
prolifera Mett. Fil. Lips. 84. 1856 = D. vivipara.
prolifera Kuhn, v. Deck. Reise 3^{3}. Bot. 44. 1879.
propinqua Fée, Cr. vasc. Br. **1**. 103. 1869 = D. grandis *macroptera.
pteroidea Mett. Pheg. u. Asp. 9 n. 3. 1858.
pubescens Fée, Cr. vasc. Br. **1**. 98. 1869 = D. oligocarpa.
pubescens Keys. Pol. Cyath. Hb. Bung. 51. 1873 = Lonchitis p.
punctata Mett. Ann. Lugd. Bat. **1**. 222. 1864.
pycnolepis Mett. Pheg. u. Asp. 11 n. 7. 1858 = Polystichum aculeatum var. 27
refulgens Mett. Ann. sc. nat. V. **2**. 240. 1864.
refracta Fourn. Bull. Soc. Fr. **19**. 254. 1872.
reptans Eat. Bull. Torr. Cl. **10**. 101. 1883.
reticulata Mett. Fil. Lechl. **2**. 24. 1859.
rhaetica Pérard, Bull. Soc. Fr. **16**. 258. 1869 = Athyrium alpestre.
rigida Mett. Pheg. u. Asp. 10 n. 5. 1858 = Polystichum aculeatum var. 24.
rivulorum Fée, Cr. vasc. Br. **1**. 98. 1869 = D. oligocarpa.
Robertiana A.Br.; Aschers. Fl. Brand. **2**. 198. 1859.
rotundata Fée, 11 mém. 52. 1866.
rubrinervis Mett.; Kuhn, Linn. **36**. 116. 1869.
rudis Mett. Fil. Lips. 83. 1856.
rufescens Mett. Ann. Lugd. Bat. **1**. 223. 1864.
rugosula Mett. Ann. Lugd. Bat. **1**. 222. 1864 } = D. punctata *rugosula.
rugulosa Fée, Gen. 243. 1850-52 }
rupestris Mett. Fil. Lips. 82. 1856.
rustica Fée, 11 mém. 55 t. 13 f. 1. 1866.
salicifolia Mett. Fil. Lechl. **2**. 22. 1859.
sancta Fée, Gen. 243. 1850-52.
Sandwicensis Mann, Pr. Amer. Ac. **7**. 218. 1868.
saxatilis Mett.; Salom. Nom. 278. 1883 = Polystichum aculeatum var. 24?.
scalpturata Fée, Gen. 243. 1850-52 = D. heteroptera.
scalpturoides Fée, 11 mém. 51. 1866.
Scottii Bedd. Ferns br. Ind. Suppl. 19. 1876 = ? D. hirtipes.
scrobiculata Fée, Cr. vasc. Br. **1**. 102 t. 31 f. 2. 1869 = D. grandis *macroptera.
Seemanni J. Sm. Bot. Voy. Herald 228 t. 49. 1854 = D. brachyodus.

sericea Mett.; Eat. Mem. Amer. Ac. n. s. **8**. 208. 1860 = D. pubescens var.
serrata Mett. Ann. sc. nat. V. **2**. 243. 1864.
Sieberiana Fée, Gen. 243. 1850-52 = D. cyatheifolia.
simplex Mett. Fil. Lechl. **2**. 21. 1859.
Sloanei Fée, Gen. 243. 1850-52 = D. ampla.
sorbifolia Mett. Fil. Lechl. **2**. 23. 1859 = D. reticulata*.
sparsisora Keys. Pol. Cyath. Hb. Bung. 51. 1873 = Hypolepis s.
speciosissima Mett. Cheil. 4 in nota 1859 = Plecosorus s.
spectabilis Fée, Gen. 243. 1850-52 = D. subincisa var.
sphaeropteroides Christ, Bull. Boiss. **7**. 14. 1899.
spinulosa Hill. Fl. Haw. 566. 1888 = D. acutidens.
splendida Fée, Gen. 243. 1850-52 = D. grandis.
Stegnogramme Mett. Fil. Lips. 84. 1856.
stegnogrammoides Luerss. Flora **1875**. 435.
stenolepis Fée, 8 mém. 89. 1857 = D. Galeottii.
stenoptera Christ, Bull. Boiss. **6**. 963. 1898 = D. Dielsii.
straminea Fée, Gen. 246. 1850-52 = D. Helliana.
straminea Salom. Nom. 279. 1883.
Sturmii Phil. Anal. Univ. Chil. **94**. 358. 1896.
subconnexa Christ in Warb. Monsunia **1**. 83. 1900.
subdecurrens Luerss. Bot. Centralbl. **11**. 30. 1882 = Aspidium s.
subdigitata Bedd. Hdb. 295. 1883 = Monachosorum s.
subdryopteris Christ, Prim. Fl. Costaric. **3**. 36. 1901.
subincisa Fée, Gen. 243. 1850-52.
submarginalis J. Sm. Cat. cult. Ferns 16. 1857.
subobscura Christ, Bull. Boiss. **6**. 836. 1898.
subsimilis Mett.; Kuhn, Fil. Afr. 123. 1868.
subsimilis Christ, Ann. Jard. bot. Genève **3**. 36. 1899 = D. paraguayensis.
Tatei Salom. Nom. 279. 1883 = Aspidium T.
tenella Fée, Gen. 243. 1850-52; 7 mém. 62 t. 25 f. 2. 1857 = D. sancta.
tenuifolia Keys. Pol. Cyath. Hb. Bung. 51. 1873 = Hypolepis t.
tenuis Fée. Cr. vasc. Br. **1**. 99 t. 30 f. 1. 1869 = D. rotundata.
tetragona Fée, Gen. 243. 1850-52 = ? D. oligocarpa.
tetragona Mett. Fil. Lips. 84. 1856.
thelipteroides Fée, Gen. 243. 1850-52 = D. tomentosa.
Thwaitesii Mett. Fil. Lechl. **2**. 24. 1859.
tijuccana Fée, Gen. 243. 1850-52.
tomentosa Mett.; Kuhn, Fil. Afr. 123. 1868 = D. arthrothrix.
totta Mett. Pheg. u. Asp. 18 n. 31. 1858 = D. africana.
trichodes J. Sm. Cat. cult. Ferns 17. 1857 = D. setigera.
tricholepis Fée, Cr. vasc. Br. **1**. 98 t. 32 f. 1. 1869 = D. tijuccana.
triphylla Mett. Fil. Lechl. **2**. 21. 1859.
triphylla Kuhn, Fil. Afr. 124. 1868 = Aspidium puberulum.
Ulei Christ in Schwacke, Pl. Nov. Mineiras **2**. 29. 1900.
unidentata J. Sm. Ferns br. and for. 170. 1866.
unita Mett. Pheg. u. Asp. 22 n. 42. 1858 = D. silvatica.
urophylla Mett. Pheg. u. Asp. 26 n. 53. 1858.
vasta Mett. Fil. Lechl. **2**. 27. 1859 = D. subincisa*.
vestita Phil. Linn. **29**. 107. 1857 = D. subincisa*.
villosa Mett. Fil. Lips. 82. 1856 = D. dasyphylla.

villosa Fée, 11 mém. 53. 1866 = D. pubescens.
Vogelii Kuhn, Fil. Afr. 124. 1868 = D. subcoriacea.
vulgaris Mett. Fil. Lips. 83. 1856 = D. phegopteris.
Walkerae J. Sm. Ferns br. and for. 169. 1866.

PHLEBIOGONIUM Fée, Gen. Fil. 314. 1850-52 = **Aspidium.**
impressum Fée, Gen. 314 t. 24 A. f. 2. 1850-52 = A. brachiatum.

Phlebiolepicystis Diels, Nat. Pfl. 1[4]. 321. 1899 (*Lepicystis §*) = Polypodium.

PHLEBIOPHYLLUM van den Bosch, Versl. Ak. Wet. Amsterdam **11**. 321. 1861 = **Trichomanes** (venosum).

Phlebodiopsis Moore, Ind. Fil. LXXVII. 1857 (*Pleopeltis §*) = Polypodium.

PHLEBODIUM (R.Br.) J. Smith, Journ. Bot. **4**. 58. 1841 = **Polypodium.**
Polypodium § Phlebodium R.Brown in Horsf. Pl. Jav. rar. 7. 1838.
angustatum Brack. Expl. Exp. **16**. 41. 1854 (err.) = P. angustum.
angustum J. Sm. JoB. **4**. 59. 1841.
areolatum J. Sm. JoB. **4**. 59. 1841 = P. aureum*.
aureum J. Sm. JoB. **4**. 59. 1841.
decumanum J. Sm. JoB. **4**. 59. 1841.
dictyocallis J. Sm. Cat. cult. Ferns 4. 1857 = P. decumanum var.
elegans ht. de Smet-Duvivier; Gard. Chr. III. **33**. 266. 1903 — Hort.
elongatum J. Sm. JoB. **4**. 59. 1841 = P. lanceolatum*.
ensiforme J. Sm. JoB. **4**. 59. 1841.
fulvum Moore, Ind. 346. 1862 = P. aureum *areolatum.
grande Moore, Ind. 258. 1861 = P. aureum.
inaequale Moore, Gard. Chr. **1855**. 660 = P. Lowei.
lanceolatum J. Sm. JoB. **4**. 59. 1841.
lanosum Moore, Ind. 258. 1861 = P. aureum *areolatum.
latum J. Sm. JoB. **4**. 59. 1841 (nomen) —?
lycopodioides J. Sm. Bot. Mag. **72**. Comp. 12. 1846.
macrocarpum J. Sm. JoB. **4**. 59. 1841 = P. lanceolatum.
Mayii ht. May.: Gard. Chr. III. **23**. 332 f. 121. 1898 = P. aureum var. hort.
microdictyum Moore, Ind. 258. 1861 = P. aureum.
multiseriale Moore et Houlst. Gard. Chr. **1855**. 469 = P. decumanum var.
nigripes J. Sm. Ferns br. and for. ed. II. 296. 1877 = P. plebejum var.
nitidum J. Sm. Bot. Mag. **72**. Comp. 13. 1846 — Honduras.
percussum J. Sm. JoB. **4**. 59. 1841.
pulvinatum J. Sm. Bot. Mag. **72**. Comp. 13. 1846 = P. aureum*.
salicifolium J. Sm. JoB. **4**. 59. 1841 = P. lycopodioides*.
serpens J. Sm. JoB. **4**. 59. 1841 = P. Swartzii.
sporadocarpum J. Sm. Bot. Mag. **72**. Comp. 13. 1846 = P. aureum *areolatum.
squamulosum J. Sm. Bot. Mag. **72**. Comp. 13. 1846.
venosum Moore et Houlst. Gard. Mag. Bot. **3**. —. 1851 = P. stigmaticum.

PHOROLOBUS Desvaux, Prodrome 291. 1827.
acrostichoides Fée, Gen. 131. 1850-52 = Cryptogramma.
Brunonianus Fée, Gen. 131. 1850-52 = Cryptogramma.
chinensis Desv. Prod. 291. 1827 = Pteris ensiformis.
crispus Desv. Prod. 291. 1827 = Cryptogramma.
domingensis Desv. Prod. 291. 1827 (Plum. t. 73) — Hispaniola.
pteroides Desv. Prod. 291. 1827 = Neurosoria.
siliculosus Desv. Prod. 291. 1827 = Onychium.

PHOTINOPTERIS J. Smith, Journ. bot. **3**. 403. 1841; **4**. 155. 1841 — (Polypodiaceae Gen. **105**).
Cumingii Pr. Epim. 192. 1849 = P. speciosa.
drynarioides Bedd. Ferns br. Ind. t. 325. 1869 = Dryostachyum.
Horsfieldii J. Sm. JoB. **3**. 403. 1841 = P. speciosa.
Humboldtii Pr. Epim. 192. 1849 (patria erronea) = P. speciosa.
rigida Bedd. Ferns br. Ind. t. 211. 1867 = P. speciosa.
simplex J. Sm. JoB. **3**. 403. 1841 = P. speciosa.
speciosa Bl.; Pr. Epim. 264. 1849. NPfl. 328 — Ins. Philipp. Malesia.
Lomaria Bl. Enum. 202. 1828; Dryostachyum Kuhn 1869; Polypodium Chr. 1897. 121 (non Bl. 1828); Acrostichum rigidum Wall. 1828. HB. 424; Photinopteris Bedd. 1867. P. Horsfieldii J. Sm. 1841; Lomariopsis Mett. 1856 — Photinopteris simplex J. Sm. 1841 — P. Cumingii Pr. 1849.
Phyllitidis J. Smith, Journ. Bot. **3**. 397. 1841 *(Drynaria §)* = Polypodium.
PHYLLITIS Ludwig, Instit. Hist. Phys. Regn. Veg. ed. II. 142. 1757 — (Polypodiaceae Gen. **46**).
Balansae (Bak.) C. Chr. Ind. 492. 1906 — Paraguay.
Scolopendrium Bak. in Hk. Ic. pl. t. 1653. 1886. NPfl. 233.
brasiliensis (Sw.) O.Ktze. Rev. Gen. Pl. **2** 818. 1891 — Brasilia.
Asplenium Sw. Vet. Ak. Handl. **1817**. 65 t. 3 f. 1; Scolopendrium Kze. 1850. HB. 247. Chr. 212. NPfl. 233; Antigramma Moore 1857; Scolopendrium ambiguum Raddi 1819; S repandum Pr. 1822; Antigramma Pr. 1836 — A. subsessilis Fée 1850-52; Scolopendrium Diels 1899. NPfl. 233.
Delavayi (Franch.) C. Chr. Ind. 492. 1906 — Birma septent. China austr.
Scolopendrium Franch. Bull. Soc. Fr. **32**. 29. 1885. Chr. 213. NPfl. 233.
D'Urvillei O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 = Stenochlaena sorbifolia var. 16.
hemionitis (Lag.) O.Ktze. Rev Gen. Pl. **2**. 818. 1891 — Mediterranea.
Scolopendrium Lag. Garcia et Clem. Anal. cienc. **5**. 150 t. 41 f. 2. 1802. HB. 247. Chr. 211. NPfl. 230; S. sagittatum DC. 1815 — S. cordatum Fée 1850-52.
hybrida (Milde) C. Chr Ind. 492. 1906 — Ins. Lussin.
Scolopendrium Milde, Verh. zool bot. Ges. **14**. 235 t. 18. 1864. Chr. 211. NPfl. 230.
Ikenoi (Mak.) C. Chr. Ind 492. 1906 — Japonia.
Scolopendrium Mak. Bot. Mag. Tokyo **13**. 130. 1899.
Lindeni Maxon, Fernwort Paper **2**. 30. 1900 = P. scolopendrium var.
longifolia O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 = Triphlebia.
nigripes (Fée) O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 — Mexico. Guatemala.
Schaffneria Fée, 7 mém. 56 t. 17 f. 1. 1857: Asplenium Hk. 1857; Scolopendrium Hk. 1862. HB. 247. Chr. 212. NPfl. 233; Antigramma J. Sm. 1875.
pinnata O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 = Triphlebia.
plantaginea (Schrad.) O.Ktze. Rev Gen. Pl **2**. 818. 1891 — Brasilia.
Scolopendrium Schrad. Gött. gel. Anz. **1824**. 870. HB. 247. Chr. 211. NPfl. 233; Antigramma Pr. 1836; Asplenium Douglasii Hk. et Grev. 1829; Hemidictyum Pr. 1836; Antigramma Hk. 1840; Scolopendrium Fisch.; Hk. 1862. Camptosorus rumicifolius Link 1841 — Antigramme Lageana Fée et Glaz. 1869.
rhizophylla O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 = Camptosorus.
scolopendrium (L.) Newm. Hist. brit. Ferns ed. II. 10. 1844 — Europa. Ins. Atlant. Asia minor-Persia. Japonia. Amer. bor. atlant. Mexico.
Asplenium L. sp. **2**. 1079. 1753; Scolopendrium Karst. 1880-83; S. vulgare Sm. 1793. HB. 246. Chr. 211. NPfl. 230; Phyllitis Bubani 1901; Scolopendrium officinarum Sw. 1801; S. Phyllitis Roth 1799; S. Lingua Cav.

1802; S. officinale Lam. et DC. 1805 — S. Lindeni Hk. 1842; Phyllitis Maxon 1900 (Mexico).
sibirica O.Ktze. Rev. Gen. Pl. **2**. 818. 1891 = Camptosorus.
vulgaris Bubani, Fl. Pyrenea **4**. 423. 1901 = P scolopendrium.
PHYLLITIS Moench, Methodus plant. 724. 1794 = **Asplenium.**
heterophylla Moench, Method. 724 = A. germanicum.
lancifolia Moench, Meth. Suppl. 316. 1802 = A. adiantum nigrum.
rotundifolia Moench, Method. 724. 1794 = A. trichomanes.
ruta muraria Moench, Method. 724. 1794.
Phyllitis J. Smith, Ferns br. and for. 209. 1866 *(Asplenium §)* = Asplenium.
Phyllotrichium Prantl, Ber. deut. bot. Ges. **1**. 349. 1883 *(Botrychium §)* = Botrychium.
PHYMATODES Presl. Tent. Pter. 195. 1836 = **Polypodium.**
acrosora Pr. Tent. 197. 1836 = P. crassifolium.
alata J. Sm. Bonplandia **9**. 261. 1861 = P. Wilkesii.
albosquamata J. Sm. Ferns br. and for. 94. 1866 = P. albidosquamatum.
alternifolia Pr. Tent. 196. 1836.
angustiloba Pr. Tent. 198. 1836 = Drynaria a.
arguta Pr. Tent. 197. 1836 (nomen) —?
Billardieri Pr. Tent. 196. 1836 = P. diversifolium.
brancaefolia Pr. Tent. 198. 1836 = Drynaria quercifolia.
capitellata Pr. Tent. 197. 1836 = P. Wallichianum.
conjugata Pr. Tent. 198. 1836 = Dipteris c.
coronans Pr. Tent. 198. 1836.
crassifolia Pr. Tent. 197. 1836.
crassinervis Pr. Tent. 197. 1836 = P. platyphyllum.
crenulata Pr. Tent. 197. 1836 = P. palmatum.
cuspidata J. Sm. Cat. cult. Ferns 10. 1857 = P. leiorhizum.
elongata Pr. Tent. 196. 1836 = P. lineare.
elongata Pappe et Raws. Syn. Fil. Afr. austr. 41. 1858 = P. Schraderi.
excavata Pr. Tent. 196. 1836.
exiguum Und. Torreya **3**. 18. 1903 = P. Swartzii.
Gaudichaudii Pr. Tent. 198. 1836 = Drynaria rigidula.
geminata J. Sm. Cat. cult. Ferns 9. 1857.
glauca J. Sm. Ferns br. and for. 94. 1866 = P. glauco-pruinatum.
grandifolia Pr. Tent. 198. 1836 = P. membranaceum var.
grossa Pr. Tent. 196. 1836 = P. phymatodes.
immersa Pr. Tent. 196. 1836.
incurvata J. Sm. Ferns br. and for. 94. 1866.
irioides Pr. Tent. 198. 1836 = P. punctatum.
laciniata Pr. Tent. 197. 1836 = P. macrochasmum var.
latifolia Pr. Tent. 196. 1836 = Aspidium l.
leiorhiza Pr. Tent. 196. 1836.
lingulata Pr. Tent. 198. 1836 = P. punctatum.
longifolia J. Sm. Cat. cult. Ferns 9. 1857 = P. revolutum.
longipes J. Sm. Cat. cult. Ferns 9. 1857 = P. phymatodes.
longissima J. Sm. Cat. cult. Ferns 10. 1857.
loriformis Pr. Tent. 196. 1836 = P. excavatum var.
lycopodioides Millsp. Field Col. Mus. Bot. Ser. **3**. 12. 1903.
Meyeniana Pr. Tent. 197. 1836 = P. palmatum.
morbillosa Pr. Tent. 198. 1836 = Drynaria quercifolia.

myriocarpa Pr. Tent. 196 t. 8 f. 12. 1836.
nigrescens J. Sm. Ferns br. and for. 94. 1866.
normalis Pr. Tent. 196. 1836.
nuda J. Sm. Bot. Voy. Herald 425. 1857 = P. lineare.
Ottoniana Pr. Tent. 197. 1836 (nomen) — ?
Drynaria Fée 1850-52 (nomen); Pleopeltis Moore 1862 (nomen).
ovata Pr. Tent. 197. 1836 = P. phyllomanes var.
oxyloba Pr. Tent. 196. 1836 = P. hastatum var.
peltidea J. Sm. Cat. cult. Ferns 9. 1857 = P. phymatodes.
phymatodes Maxon, Contr. U. S. Nat. Mus. 9. 352 t. 62. 1905.
plantaginea Pr. Tent. 197. 1836 = Aspidium p.
Plukenetii Pr. Tent. 196. 1836.
polycarpa Pr. Tent. 198 t. 8 f. 19. 1836 = P. punctatum.
porrecta Pr. Tent. 197. 1836 = P. crassifolium.
propinqua Pr. Tent. 198. 1836 = Drynaria p.
pustulata Pr. Tent. 196. 1836.
quercifolia Pr. Tent. 198. 1836 = Drynaria q.
rupestris Moore, Ind. 348. 1862 = P. triquetrum var.
saccata J. Sm. Cat. cult. Ferns 9. 1857 = P. nigrescens.
scandens Pr. Tent. 196. 1836 = P. pustulatum.
Schomburgkiana J. Sm. Lond. JoB. 1. 196. 1842 = P. megalophyllum.
secunda Pr. Tent. 197. 1836 = Dryopteris gongylodes*.
sessilis Pr. Tent. 198. 1836 = P. punctatum.
simplex Pr. Tent. 196. 1836 = P. excavatum.
sinuata Pr. Tent. 197. 1836.
sinuosum J. Sm. Ferns br. and for. ed. II. 296. 1877.
spectrum Pr. Tent. 197. 1836.
sphaerocephala Pr. Tent. 196. 1936 = Cyclophorus angustatus.
stenophylla J. Sm. Bonplandia 9. 261. 1861.
Swartzii Und. Our. Nat. Ferns ed. VI. 84. 1900.
sylvatica Pr. Tent. 198. 1836 = Drynaria quercifolia.
terminalis J. Sm. Cat. cult. Ferns 9. 1857 = P. phymatodes.
Thouarsii Pr. Tent. 198. 1836 = Drynaria Willdenowii.
tridactyla Pr. Tent. 196. 1836 = P. pteropus.
Vieillardii Fourn. Ann. sc. nat. V. 18. 287. 1873.
vulgaris Pr. Tent. 196. 1836 = P. phymatodes.
Wallichiana Pr. Tent. 197. 1836 = Drynaria propinqua.
zosteraeformis Pr. Tent. 196. 1836.

PHYMATOPSIS J. Smith. Hist. Fil. 104. 1875 = **Polypodium.**
accedens J. Sm. Hist. Fil. 104. 1875.
ebenipes J. Sm. Hist. Fil. 105. 1875.
Griffithiana J. Sm. Hist. Fil. 104. 1875.
incurvata J. Sm. Hist. Fil. 105. 1875.
laciniata J. Sm. Hist. Fil. 105. 1875 = P. macrochasmum var.
oodes J. Sm. Hist. Fil. 104. 1875.
palmata J. Sm. Hist. Fil. 105. 1875.
rhynchophylla J. Sm. Hist. Fil. 104. 1875.
rostrata J. Sm. Hist. Fil. 104. 1875 = P. subrostratum.
trifida J. Sm. Hist. Fil. 105. 1875 = P. hastatum var.

Physapteris Presl. Tent. Pterid. 160. 1836 (*Cheilanthes §*) = Cheilanthes.

PHYSEMATIUM Kaulfuss, Flora 1829[1]. 341 = **Woodsia.**
aspidioides Kze. Anal. 43. 1837 = Diacalpe a.
canescens Trev. Nu. Giorn. bot. It. 7. 161. 1875 = W. crenata.
Cumingianum Kze. Anal. 43. 1837 = W. crenata.
elongatum Trev. Nu. Giorn. bot. It. 7. 161. 1875.
europolepis Trev. Nu. Giorn. bot. It. 7. 155. 1875.
fragile Kze. Anal. 42. 1837.
incisum Pr. Tent. 66. 1836 = W. montevidensis.
molle Klf. Flora 1829[1]. 341.
obtusum Hk. Fl. bor. Am. 2. 259. 1840.
oreganum Trev. Nu. Giorn. bot. It. 7. 162. 1875.
Perrinianum Pr. Tent. 66. 1836 = W. obtusa.
philippinum Pr. Epim. 34. 1849 = Dryopteris p.
polystichoides Trev. Nu. Giorn. bot. It. 7. 161. 1875.
scopulinum Trev. Nu. Giorn. bot. It. 7. 161. 1875.
vestitum Pr. Epim. 66. 1849 = W. obtusa.

PILULARIA Linné, Spec. plant. 2. 1100. 1753 — (Marsileaceae Gen. 3).
americana A.Br. Monatsber. Ak. Berl. **1863**. 435. 1864. Bak. Hdb. 148. NPfl. 421. — California. Oregon. Arkansas. Chile. — Calamistrum O.Ktze. 1891.
globulifera L. sp. 2. 1100. 1753. Bak. Hdb. 149. NPfl. 421 — Europa. Calamistrum O.Ktze. 1891.
Mandoni A.Br. Monatsber. Ak. Berl. **1870**. 752. Bak. Hdb. 149. NPfl. 421 — Bolivia. — Calamistrum O.Ktze. 1891.
minuta Durieu; A.Br. Monatsber. Ak. Berl. **1863**. 435. 1864. Bak. Hdb. 148 NPfl. 421 — Mediterranea. — Calamistrum O.Ktze. 1891.
natans Mérat, Fl. Paris. ed. II. 2. 283. 1821 = P. globulifera.
novae-hollandiae A.Br. Monatsber. Ak. Berl. **1863**. 435. 1864. Bak. Hdb. 148. NPfl. 421 — Austr. Tasmania. — Calamistrum O.Ktze. 1891.
novae-zelandiae Kirk, Tr. N. Zeal. Inst. 9. 547 t. 29. 1877. Bak. Hdb. 148. NPfl. 421 — N. Zealand. — Calamistrum O.Ktze. 1891.
valdiviana Philippi (Bak. Hdb. 148) = P. americana.

PINONIA Gaudichaud. Ann. sc. nat. 3. 507. 1824 = **Cibotium.**
splendens Gaud. l. c.; Freyc. Voy. 96. 369 t. 21. 1827 = C. Chamissoi.

PLAGIOGYRIA (Kunze) Mettenius, Farng. II Plagiogyria 1858 — (Polypodiaceae Gen. 77).
Lomaria § Plagiogyria Kze. Farnkr. 2. 61. 1850.
adnata (Bl.) Bedd. Ferns br. Ind. t. 51. 1865. NPfl. 282 — Japonia. Yunnan. Khasia. Java. Ins. Philipp. — Lomaria Bl. Enum. 205. 1828. HB. 182.
? *aequidentata* Fourn. Mex. pl. 1. 133. 1872 — Mexico.
assurgens Christ. Bull. Soc. bot. It. **1901**. 293 — China.
biserrata Mett. Plagiog. 8 n. 1. t. 4. 1858 = P. semicordata.
costaricensis Mett. Kuhn. Linn. **36**. 149. 1869. — Costa Rica. Lomaria Bak. 1874. HB. 482.
euphlebia (Kze.) Mett. Plagiog. 10. n. 6. 1858. NPfl. 282 — Japonia. China. Ind. bor. Australia trop. — Lomaria Kze. Bot. Zeit. **1848**. 521. HB. 183. Acrostichum triquetrum Wall. 1828 (nomen); Olfersia Pr. 1836 (nomen); Stenochlaena J. Sm. 1841 (nomen); Plagiogyria Mett. n. 5. 1858; Lomariopsis Ettingsh. 1865 — Lomaria articulata F. Muell. 1866. — (an Acrostichum seetacoonense Roxb. 1844?).
Fauriei (Christ) C. Chr. Ind. 497. 1906 — Japonia. Lomaria Christ, Bull. Boiss. 4. 666. 1896.

glauca (Bl.) Mett. Plagiog. 9. n. 3. 1858. Chr. 175. NPfl. 282 — Ind. bor. Yunnan. Malesia. Luzon. — Lomaria (a. b.) Bl Enum. 204. 1828. HB. 182.

Henryi Christ, Bull. Boiss. 7. 8. 1899. NPfl. 282 — Yunnan.

Matsumureana Mak. Bot. Mag. Tokyo 8. 333. 1894 — Japonia. Lomaria Mak. Bot. Mag. Tokyo 8. 90. 1894 (pars japonica).

pycnophylla (Kze.) Mett. Plagiog. 8. n. 2. 1858. Chr. 175. NPfl. 282 — Yunnan. Ind. bor. Malacca. Java. Borneo. Ins. Philipp. — Lomaria Kze. Bot. Zeit. **1848**. 143. HB. 183; Stenochlaena? Pr. 1849. Lomaria callosa Fée 1850-52 — Plagiogyria scandens Mett. n. 4. 1858. NPfl. 282.

scandens Mett. Plagiog. 9. n. 4. t. 4 f. 21-22. 1858 = P. pycnophylla.

semicordata (Pr.) Christ, Farnkr. 176. 1897. NPfl. 282 — Amer. trop Lomaridium? Pr. Epim. 155. 1849; Lomaria Bak. 1867. HB. 182. L. pectinata Liebm. 1849 (nom. optim.?). L. arguta Fée 1857. L. biserrata Mert. et Lind. hb.; Hk. 1860; Plagiogyria Mett. 1858. n. 1. t. 4 — Lomaria Fialhoi Fée et Glaz. 1869.

stenoptera (Hance) Diels, Nat. Pfl. 1⁴. 282. 1899 — Formosa. China. Blechnum Hance JoB. **1883**. 268; Lomaria Bak. 1891; L. concinna Bak 1885.

triquetra Mett. Plagiog. 10. n. 5. 1858 = P. euphlebia.

PLATYCERIUM Desvaux, Prodrome 213. 1827 — (Polypodiaceae Gen. **111**).

Aethiopicum Hk. Gard. Ferns t. 9. 1862 = P. stemaria.

alcicorne Desv. Prod. 213. 1827 = P. bifurcatum et P. stemaria.

alcicorne Gaud. Freyc. Voy. Bot. 307. 1827 et auctt. = P. bifurcatum.

andinum Bak. Ann. Bot. **5**. 496. 1891. Chr. 125. NPfl. 339; Christ, Hedwigia **44**. 363. 1905 — Peru. Bolivia. — Alcicornium Und. 1905.

angolense Welw.; Bak. Syn. 425 in adnota 1868. Chr. 127. NPfl. 339 — Africa trop. — Alcicornium Und. 1905; Platycerium elephantotis Schweinf. 1871; Kuhn, v. Deck. Reis. **3**³. Bot. 54. 1879.

angustatum Desv. Prod. 213. 1827 = P. bifurcatum.

biforme Bl. Fl. Jav. Fil. 14. t. 18. 1828 = P. coronarium.

biforme Hk. Gen. t. 80 B. 1842 = P. grande.

bifurcatum (Cav.) C. Chr. Ind. 498. 1906 — Australia. N. Caledonia. Ins. Lord Howe. ? Africa orient. — Acrostichum Cav. Anal. Hist. Nat. **1**. 105. 1799; Descr. 240. 1802; Alcicornium Und. 1905; Platycerium angustatum Desv. 1827; Acrostichum alcicorne pt. auctt. vet. (non Willem. 1796); Platycerium Gaud. 1827. HB. 425. Chr. 124. NPfl. 339; Neuroplatyceros Fée 1845; Elaphoglossum Keys. 1873.

elephantotis Schweinf. Bot. Zeit. **1871**. 361 c. fig. = P. angolense.

Ellisii Bak. Journ. Linn. Soc. **15**. 421. 1876; Ic. pl. t. 1695. Chr. 127. NPfl. 339 — Madagascar. — Alcicornium Und. 1905.

grande (A. Cunn.) J. Sm. JoB. **3**. 402. 1841 (nomen). HB. 425. Chr. 127. NPfl. 339 — Malesia. Austral. trop. — Acrostichum A. Cunn. msc.; Neuroplatyceros Fée. 2 mém. 103. 1845; Alcicornium Und. 1905.

Hillii Moore, Gard. Chr. n. s. **10**. 51 f. 6; 429 f. 74-75. 1878 — Queensland. Alcicornium Und. 1905.

madagascariense Bak. Journ. Linn. Soc. **15**. 421. 1876; **16**. 205. 1877. NPfl. 339 — Madagascar. — Alcicornium Und. 1905.

stemaria (Beauv.) Desv. Prod. 213. 1827 (stemmaria). Chr. 126. NPfl. 339 — Africa trop. Madagascar. — Acrostichum Beauv. Fl. d'Oware **1**. 2. t. 2. 1804; Alcicornium Und. 1905; Neuroplatyceros aethiopicus Fée 1845; Platycerium Hk. 1862. HB. 425. — an Acrostichum alcicorne Willem. 1796?

sumbawense Christ. Farnkr. 125. 1897 et in Warb. Monsunia **1**. 64. 1900. NPfl. 339 — Sumbawa. — Alcicornium Und. 1905.

Veitchii (Und.) C. Chr. Ind. 497. 1906; (Gard. Chr. III. **19**. 652. 1896 (nomen)) — Australia. — Alcicornium Und. Bull. Torr. Cl. **32**. 596. 1905.

Wallichii Hk. Gard. Chr. **1858**. 765; Fil. Exot. t. 97. 1859. HB. 425. Chr. 127. NPfl. 339 — Tenasserim. Malacca. — Alcicornium Und. 1905.

Wandae Racib. Bull. int. Ac. Cracovie **1902**. 58 — N. Guinea.

Willinckii Moore, Gard. Chr. n. s. **3**. 301 f. 56. 1875 – Java. Alcicornium Und. 1905.

Platyglossa Christ, Mon. Elaph. 20. 1899 *(Elaphoglossum §)* = Elaphoglossum.

PLATYLOMA J. Smith, Journ. Bot. **4**. 160. 1841 = **Pellaea.**

adiantoides Lowe, Ferns **3** t. 33. 1857 = P. viridis.

albida (Cunn.) J. Sm. JoB. **4**. 160. 1841 (nomen) — ? — Pteris Cunn. msc.

andromedæfolia J. Sm. JoB. **4**. 160. 1841.

atropurpurea J. Sm. JoB. **4**. 160. 1841.

bellum Moore, Gard. Chr. **1873**. 213 = P. brachyptera var.

brachypterum Moore, Gard. Chr. **1873**. 141.

Bridgesii J. Sm. Hist. Fil. 165. 1875.

Brownii J. Sm. JoB. **4**. 160. 1841 = P. paradoxa.

calomelanos J. Sm. Bot. Mag. **72** Comp. 21. 1846 = P. hastata.

cordata J. Sm. JoB. **4**. 160. 1841.

dura J. Sm. Hist. Fil. 165. 1875 = P. Goudotii.

falcata J. Sm. JoB. **4**. 160. 1841.

flexuosa J. Sm. Bot. Mag. **72**. Comp. 21. 1846.

geraniifolium Lowe, Ferns **3** t. 27. 1857 = Doryopteris concolor.

hastatum Lowe, Ferns **3** t. 32. 1857 = P. viridis.

intramarginale Lowe, Ferns **3** t. 31. 1857.

pulchellum Moore, Ind. LXVI. 1857.

rotundifolia J. Sm. JoB. **4**. 160. 1841.

sagittata J. Sm. JoB. **4**. 160. 1841 = P. cordata var.

ternifolia J. Sm. Bot. Mag. **72** Comp. 21. 1846.

PLATYTAENIA Kuhn, Die Chætopterides 330. 1882 — (Polypodiaceae Gen. **97**).

Requiniana (Gaud.) Kuhn, Chaetopt. 331. 1882. NPfl. 305 — Singapore. Ins. Moluccae. Ins. Philipp. N. Guinea. N. Hebridae.

Acrostichum Gaud. Freyc. Voy. Bot. 304 t. 4. 1827. (HB. 423); Neurocallis Fée 1845; Lomariopsis Kuhn 1869; Poecilopteris Carr. 1873.

PLATYZOMA R. Brown, Prod. Fl. Nov. Holl. 160. 1810 = **Gleichenia**.

alpinum Desv. Prod. 199. 1827.

dicarpum Desv. Prod. 199. 1827.

ferrugineum Desv. Prod. 199. 1827 = G. circinnata.

latum Desv. Prod. 199. 1827 = G. circinnata.

microphyllum R. Br. Prod. Fl. N. Holl. 160. 1810 = G. platyzoma.

recurvum Desv. Prod. 199. 1827 = G. circinnata.

rupestre Desv. Prod. 199. 1827.

speluncae Desv. Prod. 199. 1827 = G. circinnata.

PLECOSORUS Fée, Gen. Fil. 150. 1850-52 — (Polypodiaceae Gen. **19**).

?**leptocladon** Fée, 10 mém. 19 t. 34. f. 1. 1865 — Columbia.

Mexicanus Fée, Gen. 151 t. 13 A. f. 1. 1850-52 = P. speciosissimus.

peruvianus Fée, Gen. 151 t. 13 A. f. 2. 1850-52 — Peru. Mexico. Cheilanthes Bak. 1891.

speciosissimus (A. Br.) Moore, Ind. XL. 1857. NPfl. 195 — Mexico. Guatemala. Cheilanthes A. Br.; Kunze, Anal. 35 t. 23. 1837. HB. 141; Phegopteris Mett. 1859; Plecosorus Mexicanus Fée 1850-52. Chr. 279.

PLECTOPTERIS Fée, Gen. Fil. 230. 1850-52 = **Polypodium.**

Calymmodon Fée, 8 mém. 124. 1857 } = P. cucullatum.
gracilis Fée, Gen. 230 t. 19 B. 1850-52 }

PLENASIUM Presl, Tent. Pterid. 109. 1836 = **Osmunda.**

aureum Pr. Tent. 249 t. 3 f. 13. 1836 ± O. banksiifolia.
banksiaefolium Pr. Tent. 110. 1836.
?bromeliaefolium Pr. Tent. 110. 1836 = O. banksiifolia.
Claytonianum Pr. Abh. böhm. Ges. V. **5**. 325. 1848.
interruptum Pr. l. c. 325. 1848 = O. Claytoniana.
javanicum Pr. l. c. 325. 1848.
pilosum Pr. l. c. 325. 1848 = O. Claytoniana.
Vachellii Pr. l. c. 325. 1848.

PLEOCNEMIA Presl, Tent. Pterid. 183. 1836 = **Aspidium.**

aristata Bedd. Ferns S. Ind. t. 83. 1863-65 = Dryopteris otaria.
Blumei Moore, Ind. LXXXVII. 1857 = Dryopteris canescens.
Clarkei Bedd. Ferns br. Ind. Suppl. 15 t. 368. 1876 = A. artinexum.
conjugata Pr. Epim. 259. 1849 } = A. Leuzeanum.
Cumingiana Pr. Epim. 50. 1849 }
dissidens Moore, Ind. 90. 1858.
gigantea Pr. Epim. 259. 1849.
javanica Pr. Epim. 50. 1849 = A. Leuzeanum.
Leuzeana Pr. Tent. 184 t. 7 f. 12. 1836.
megalocarpa Bedd. Handb. Suppl. 48. 1892.
membranacea Bedd. Ferns br. Ind. Suppl. 15. 1876 = A. devexum.
membranacea Bedd. Handb. 225 f. 115. 1883.
Milnei Fourn. Ann. sc. nat. V. **18**. 299. 1873 = A. viridans.
stenopteris Moore, Ind. 105. 1858 = Dryopteris incisa.
Thwaitesii Bedd. Handb. 223. 1883.
Trimeni Bedd. Handb. 223. 1883 = A. giganteum.
Vieillardii Fourn. Ann. sc. nat. V. **18**. 299. 1873 = A. novae caledoniae.

PLEOPELTIS Humb. et Bonpl.; Willd. Spec. plant. **5**. 211. 1810 = **Polypodium.**

accedens Moore, Ind. LXXVII. 1857.
acuminata Moore, Ind. 344. 1862 = P. accedens.
adfinis Moore, Ind. 344. 1862 = P. commutatum.
alata Moore, Ind. 344. 1862 = P. Wilkesii.
albido-squamata Pr. Tent. 193. 1836.
alternifolia Moore, Ind. 344. 1862.
angusta H. B. Willd. sp. **5**. 211. 1810.
angusta Boj. Hort. Maur. 389. 1837 = P. furcatum.
?angustata Pr. Tent. 193. 1836 = Cyclophorus a.
angustata Moore, Ind. LXXVIII. 1857 = P. palmatum var.
angustifolia Eat. Amer. Journ. Sci. II. **27**. 198. 1859 = P. lanceolatum*.
apoda Moore, Ind. LXXVIII. 1857 — ? — Polypodium sessile Wall.?
araneosa Moore, Ind. 345. 1862 = P. aureum *areolatum.
areolata Pr. Tent. 193. 1836 = P. aureum*.
astrolepis Fourn. Mex. pl. **1**. 87. 1872.
aurea Pr. Tent. 193. 1836.

Billardieri Moore, Ind. LXXVIII. 1857 = P. diversifolium.
Brownii Fourn. Bull. Soc. Fr. **16**. 424. 1869.
capitellata Bedd. Ferns br. Ind. t. 12. 1865 = P. Wallichianum.
caudiformis Moore, Ind. 349. 1862 = P. Feei var.
chilensis (Fée) Moore, Ind. 345. 1862 —? — Drynaria »Fée«.
clathrata Bedd. Handb. 348. 1883.
coadunata Moore, Ind. 345. 1862 = Aspidium macrodon.
corcovadensis Pr. Tent. 193. 1836 = P. lanceolatum.
crassifolia Moore, Ind. LXXVIII. 1857.
crassinervata Moore, Ind. 345. 1862 = P. lanceolatum*.
crassinervia Moore, Ind. LXXVIII. 1857 = P. platyphyllum.
crenato-pinnata Bedd. Handb. Suppl. 96. 1892.
cuspidata Pr. Tent. 193. 1836 = P. percussum.
decumana Pr. Tent. 193. 1836.
decurrens Bl. Enum. addend. 1828 = P. revolutum.
decurrens Moore, Ind. 345. 1862 = P. insigne.
dilatata Bedd. Ferns br. Ind. t. 122. 1866 = P. euryphyllum.
ebenipes Bedd. Ferns br. Ind. t. 138. 1866.
elegans Fée, 1 mém. 14 t. 2 f. 13. 1844 (nomen) —?
elongata Klf. Enum. 246. 1824 = P. lineare.
elongata J. Sm. Cat. cult. Ferns 5. 1857 = P. lanceolatum*.
ensata Moore, Ind. 346. 1862.
ensifolia Carm. Tr. Linn. Soc. **12**. 510. 1818 = P. lanceolatum.
excavata Moore, Ind. 347. 1862.
fossa Moore, Gard. Chr. n. s. **18**. 586. 1882.
geminata Moore, Ind. 346. 1862.
glabra Fourn. Bull. Soc. Fr. **16**. 424. 1869 = Cyclophorus confluens.
glauca Moore, Ind. LXXVIII. 1857 = P. glauco-pruinatum.
glaucina Moore, Ind. 346. 1862 = P. aureum *areolatum.
glaucopsis Bedd. Handb. Suppl. 96. 1892.
grandifolia Moore, Ind. LXXVIII. 1857 = P. membranaceum var.
Grevilleana Moore, Ind. LXXVIII. 1857 = P. excavatum.
Griffithiana Moore, Ind. LXXVIII. 1857.
hastata Moore, Ind. 346. 1862.
Helenae Pr. Tent. 193. 1836 = P. lanceolatum.
hemionitidea Moore, Ind. 346. 1862.
heterocarpa Moore, Ind. LXXVIII. 1857 = P. Zollingerianum.
himalayensis Bedd. Ferns br. Ind. t. 318. 1869.
inaequalis Pr. Tent. 193. 1836 (nomen) —?
?incana Wall. List n. 55. 1828 = P. polypodioides.
incurvata Moore, Gard. Chr. **1860**. 1105.
irioides Moore, Ind. LXXVIII. 1857 = P. punctatum.
insignis Bedd. Ferns br. Ind. t. 214. 1867.
iteophylla Link, Fil. sp. 120. 1841 = P. geminatum.
juglandifolia Moore, Ind. LXXVIII. 1857 = P. Wallichianum.
Kaulfussiana Pr. Tent. 193. 1836 = P. lanceolatum.
laciniata Bedd. Handb. Suppl. 97. 1892 = P. macrochasmum.
lanceola Carr. in Seem. Fl. Vit. 367. 1873.
lanceola Bailey, Hdb. Ferns Queensl. 45. 1874 = P. simplicissimum.
lanceolata Klf. Enum. 245. 1824.

latifolia Moore, Ind. 346. 1862 = Aspidium l.
Lehmanni Bedd. Ferns br. Ind. t. 260. 1868.
leiorhiza Moore. Ind. 346. 1862.
lepidopoda Moore, Ind. 258. 1861 = P. diversifolium.
lepidota Pr. Tent. 193. 1836 = P. lanceolatum.
Leprieurii Moore, Ind. 262. 1861 = P. lycopodioides*.
leucospora Moore, Ind. LXXVII. 1857.
linearis Klf. Enum. 246. 1824 = P. lanceolatum.
linearis Moore, Ind. 346. 1862.
lomarioides Moore, Ind. LXXVIII. 1857.
longifolia Bl. Enum. addend. 1828 = P. revolutum.
longifolia Moore, Ind. 346. 1862 = P. punctatum.
longifrons Moore, Ind. 346. 1862 = P. normale var.
longissima Moore, Ind. LXXVIII. 1857.
longissima Bedd. Ferns S. Ind. t. 176. 1863-65 = P. nigrescens.
loriformis Moore, Ind. 346. 1862 = P. excavatum var.
lycopodioides Pr. Tent. 193. 1836.
macrocarpa Klf. Enum. 245. 1824 = P. lanceolatum.
malacodon Bedd. Ferns br. Ind. Suppl. 23 t. 387. 1876 = P. hastatum.
malacodon Bedd. Handb. 363. 1883. (Ferns br. Ind. Correct. 1870).
marginalis Klf. Enum. 246. 1824 = P. lanceolatum.
maxima Moore, Ind. LXXVIII. 1857.
membranacea Moore, Ind. 191. 1860.
moulmeinensis Bedd. Ferns br. Ind. t. 205. 1867 =? P. Lehmanni.
musaefolia Moore, Ind. LXXVIII. 1857.
myriocarpa Moore, Ind. LXXVIII. 1857.
neglecta Moore, Ind. 262. 1861.
nigrescens Carr. in Seem. Fl. Vit. 368. 1873.
normalis Moore, Ind. 347. 1862.
nuda Hk. Exot. Fl. **1** t. 63. 1823 = P. lineare.
nummularia Moore, Ind. 394. 1862 = P. pyrolifolium.
obtusata Moore, Ind. 347. 1862 = P. punctatum.
oodes Moore, Ind. LXXVII. 1857.
Ottoniana Moore, Ind. 347. 1862 = Phymatodes O.
ovata Moore. Ind. LXXVIII. 1857 = P. phyllomanes var.
oxyloba Bedd. Ferns S. Ind. t. 175. 1863-65 = P. hastatum var.
palmata Moore, Ind. 347. 1862.
Parishii Bedd. Ferns br. Ind. t. 125. 1866 = Drynaria propinqua.
percussa Hk. et Grev. Ic. Fil. t. 67. 1828.
persicariaefolia Moore, Ind. LXXVIII. 1857.
Phymatodes Moore, Ind. LXXVIII. 1857.
picta Moore, Gard. Chr. n. s. **15**. 331. 1881.
pinnatifida Gill.; Hk. et Grev. Ic. Fil. t. 157. 1829 = P. pycnocarpum.
plantaginea Moore, Ind. 349. 1862 = Aspidium p.
platyphylla Bedd. Handb. Suppl. 94. 1892.
polycarpa Moore. Ind. LXXVIII. 1857 = P. punctatum.
polylepis Moore, Ind. 348. 1862.
Powellii Carr. in Seem. Fl. Vit. 367. 1873.
pteropus Moore, Ind. LXXVIII. 1857.
pulverulenta Moore. Ind. 347. 1862 = P. punctatum.

punctata Bedd. Ferns br. Ind. Suppl. 22. 1876.
Purdiei Moore, Ind. 349. 1862 = Aspidium draconopterum.
pustulata Moore, Ind. LXXVIII. 1857.
Raddiana Gaud. Freyc. Voy. Bot. 350. 1827 = P. angustum.
rhynchophylla Moore, Ind. LXXVIII. 1857.
rostrata Bedd. Ferns br. Ind. t. 159. 1867 = P. subrostratum.
rupestris Moore, Ind. LXXVIII. 1857 = P. triquetrum var.
salicifolia Pr. Tent. 193. 1836 = P. lycopodioides*.
Schomburgkiana Moore, Ind. 348. 1862 = P. megalophyllum.
Scortechinii Bedd. Handb. Suppl. 95. 1892.
serpens Kze.; Pr. Tent. 193. 1836 = ? P. Swartzii.
sessilis Moore, Ind. LXXVIII. 1857 = P. punctatum.
simplex Bedd. Handb. 347. 1883 = P. excavatum.
sinuosa Bedd. Ferns br. Ind. t. 8. 1865.
spectrum Moore, Ind. 348. 1862.
sphaerocephala Bl. Fl. Jav. Fil. 64. 1828 = Cyclophorus angustatus.
squamata J. Sm. Hist. Fil. 114. 1875.
squamulosa Pr. Tent. 193. 1836.
stenoloma Moore, Ind. 348. 1862 = P. angustum.
stenophylla Moore, Ind. LXXVII. 1857.
Stewartii Bedd. Ferns br. Ind. t. 204. 1867.
stigmatica Pr. Tent. 193. 1836.
subspathulata Moore, Ind. 348. 1862 = P. lineare.
superficialis Bedd. Ferns br. Ind. t. 75. 1865.
surinamensis Pr. Tent. 193. 1836 = P. lycopodioides*.
tenuiloris Moore, Ind. LXXVIII. 1857.
Thunbergiana Klf. Wes. d. Farrnkr. 113. 1827 = P. lineare.
torulosa Moore, Ind. 348. 1862 = P. angustum.
tridactyla Moore, Ind. LXXVIII. 1857 = P. pteropus.
tridens J. Sm. Hist. Fil. 114. 1875.
trifida Bedd. Ferns br. Ind. Correct. II. 1870 = P. hastatum var.
ussuriensis Regel et Maack, Mém. Ac. Pétersb. VII. **4**^{4}. 175. 1861 = P. lineare.
varians Moore, Ind. 344. 1862 = P. albidosquamatum.
Wardii Bedd. Handb. Suppl. 98. 1892.
Wightiana Bedd. Ferns S. Ind. t. 180. 1863-65 = P. lineare.
Wrayi Bedd. Handb. Suppl. 93. 1892.
xiphias Moore, Gard. Chr. n. s. **15**. 331. 1881.
zeylanica Moore, Ind. 349. 1862 = P. lineare.
Zippelii Moore, Ind. 348. 1862.
zosteraeformis Bedd. Ferns br. Ind. t. 123. 1866.

PLEURIDIUM (Pr.) Fée, Gen. Fil. 273. 1850-52 = **Polypodium.**
Polypodium § Pleuridium Presl, Tent. 196. 1836.
albo-punctatissimum J. Sm. Ferns br. and for. 95. 1866 = P. crassifolium.
angustatum J. Sm. Ferns br. and for. 96. 1866 = P. palmatum var.
angustum Fée, 8 mém. 98. 1857 = P. crassifolium var.
crassifolium Fée, Gen. 274. 1850-52.
crassinervium J. Sm. Ferns br. and for. 95. 1866 = P. platyphyllum.
cuspidiflorum J. Sm. Bonplandia **9** 261. 1861 = P. accedens.
juglandifolium J Sm. Cat. cult. Ferns 10. 1857 = P. Wallichianum.
oxyloba J. Sm. Ferns br. and for. 96. 1866 = P. hastatum var.

palmatum J. Sm. Cat. cult. Ferns 10. 1857.
rupestre Fée, Gen. 274. 1850-52 = P. triquetrum var.
triquetrum J. Sm. Ferns br. and for. 95. 1866.
venustum J. Sm. Cat. cult. Ferns 10. 1857 = P. himalayense.
vulcanicum J. Sm. Bonplandia **9**. 261. 1861 = P. Feei.

Pleurogonium Presl, Tent. Pterid. 187. 1836 (*Marginaria* §) = Polypodium.

PLEUROGRAMMA R.Brown in Horsf. Pl. Jav. rar. 4. 1838 = **Dryopteris.**

PLEUROGRAMME (Bl.) Presl, Tent. Pterid. 223. 1836 = **Monogramma.**
Antrophyum § Pleurogramme Bl. Fl. Javae Fil. 69. 1828.
furcata J. Sm. Hist. Fil. 178. 1875 = M. graminoides
graminifolia Pr. Tent. 223. 1836 = M. seminuda.
graminoides Fée, 3 mém. 37. 1851-52
gyroflexa Christ, Prim. Fl. Costar. **3**. 10. 1901.
immersa Fée. 3 mém. 37 t. 4 f. 5. 1851-52 = M. linearifolia.
interrupta Christ, Farnkr 55. 1897.
linearifolia Moore, Ind. XXVII. 1857.
linearis Pr. Tent. 223 t. 10 f. 2. 1836 = M. seminuda.
myrtillifolia Fée, 3 mém. 38 1851-52.
?paradoxa Fée, 3 mém. 38 t. 4 f. 4. 1851-52.
pumila Pr. Tent. 223. 1836 = M. seminuda.
robusta Christ, Bull. Boiss. **6**. 867. 1898.
seminuda J. Sm. Hist. Fil. 178. 1875.

PLEUROMANES Presl, Epim. Bot. 258. 1849 = **Trichomanes.**
acutum Pr. Epim. 258. 1849 } = T. pallidum.
pallidum Pr. Epim. 258. 1849 }

PLEUROSORUS Fée, Gen. Fil. 179. 1850-52 — (Polypodiaceae Gen. **50**).
cuneatus Fée, Gen. 180. 1850-52 = P. rutifolius.
immersus Fée. Gen. 180 t. 16 C. 1850-52 = P. papaverifolius.
papaverifolius (Kze.) Fée. Gen. 180. 1850-52. NPfl. 275 — Chile.
Gymnogramme Kze. Anal. 12 t. 8 f. 2. 1837; Ceterach Mett. 1856; Gymnogramme Pozoi γ HB. 379 — Pleurosorus immersus Fée 1850-52; Ceterach Sturm 1858; Gymnogramme Moore 1861.
Pozoi (Lag.) Diels, Nat. Pfl. **1**[4]. 245. 1899 — Hispania.
Hemionitis Lag. Gen. et Sp. 33. 1816; Gymnogramma Desv. 1827. HB. 379 (excl. varr.). Chr. 71 (pt.); Ceterach A.Br.; Milde 1866; Grammitis hispanica Coss. —.; Ceterach Mett. 1856.
rutifolius (R.Br.) Fée, Gen. 180. 1850-52. NPfl. 245 — Australia. N. Zealand.
Grammitis R.Br. Prod. Fl. N. Holl. 146. 1810; Gymnogramma Desv. 1827; Ceterach Mett. 1856; Gymnogramme Pozoi β HB 379 — G. alpina Potts 1879.

PODEILEMA R.Brown msc. (Wall. Ic. pl. as. rar. **1**. 42. 1830) = **Peranema.**

PODOPELTIS Fée, Gen. Fil. 286. 1850-52 = **Aspidium.**
plantaginea Fée, Gen. 9. 1850-52.
Singaporiana Fée, Gen. 286. 1850-52.
sinuata J. Sm. Hist. Fil. 199. 1875 = A. plantagineum var.

POECILOPTERIS Presl. Tent. Pter. 241. 1836 = **Leptochilus.**
Brackenridgei Carr. in Seem. Fl. Vit. 374. 1873 = ? L. lomarioides.
Blumeana Bedd. Ferns br. Ind. ad finem 1870 = L. lomarioides.
brunea Pr. Tent. 242. 1836 = L. alienus.
contaminans Moore, Ind. 8. 1857 = L. virens.
costata Moore, Ind. 8. 1857 = L. scalpturatus.

crenata Pr. Epim. 174. 1849 = L. serratifolius.
crispatula J. Sm. Cat. cult. Ferns 23. 1857 = L. virens var.
diversifolia Pr. Tent. 242. 1836.
flagellifera J. Sm. Cat. cult. Ferns 23. 1857 = L. heteroclitus.
fluviatilis J. Sm. Hist. Fil. 137. 1875.
fraxinifolia Pr. Tent. 242 t. 11 f. 1. 1836 = L. serratifolius.
heteroclita Pr. Tent. 242 1836.
Hookeriana Moore, Ind. XX. 1857 = L. subcrenatus.
lobulosa Pr. Epim. 174. 1849 — Guatemala. — Heteronevron Fée 1850-52.
lonchophora Moore, Ind. 11. 1857 = L. cuspidatus.
ludens Moore, Ind. 11. 1857 = Polybotrya appendiculata var.
?menisciоides Pr. Epim. 175. 1849 = L. nicotianifolius.
polyphylla Carr. in Seem. Fl. Vit. 374. 1873 = ? Polybotrya p.
Presliana Moore, Ind. XX. 1857.
prolifera Pr. Epim. 172. 1849.
punctulata Pr. Tent. 242. 1836 = L. auriculatus.
Quoyana Pr. Epim. 173. 1849 = L. cuspidatus var.
repanda Pr. Epim. 172. 1849 = L. cuspidatus.
repanda Bedd. Ferns S. Ind. t. 202. 1863-65 = L. virens.
Requiniana Carr. in Seem. Fl. Vit. 374. 1873 = Platytaenia R.
rivularis Carr. in Seem. Fl. Vit. 374. 1873.
scandens Pr. Epim. 175. 1849 = L. guianensis.
semicordata Moore; Bedd. Ferns br. Ind. t. 270. 1868 = L. virens.
sinuosa Pr. Epim. 173. 1849 = L. cuspidatus.
Smithii Carr. in Seem. Fl. Vit. 374. 1873 = L. cuspidatus.
subcrenata Moore, Ind. XX. 1857.
subrepanda Pr. Epim. 171. 1849 = L. Zollingeri.
terminans Moore, Ind. 15. 1857 = L. virens.
terminans Bedd. Ferns S. Ind. t. 203. 1863-65 = L. subcrenatus.
virens Moore, Ind. XX. 1857.

POIKILOPTERIS Eschweiler, Linnaea 2. 117. 1827 = Poecilopteris.

POLYAETNIUM Desvaux, Prodrome 218. 1827 (err.) = Polytaenium.

POLYBOTRYA Humb. et Bonpl.; Willd. sp. 5. 99. 1810 — (Polypodiaceae Gen. **21**).
acrostichoides Mett.; Kuhn. Fil. Afr. 52. 1868 = Leptochilus.
acuminata Klf.; Link, Hort. Berol. **2**. 135. 1833 = P. scandens
altescandens C. Chr. Ind. 7. 1905; 503. 1906 — Ecuador.
Acrostichum chrysolepis Sod. Cr. vasc. quit. 485. 1893 (non Fée 1869).
andina C. Chr. Ind. 11. 1905; 504. 1906 — Ecuador.
Acrostichum insigne Bak. JoB. **1877**. 167 (non Fée 1872-73).
apiifolia J. Sm JoB. **3**. 401. 1841 (nomen); Kze. Farnkr. **1**. 142 t. 62. 1844. NPfl. 197 — Ins. Philipp. — Psomiocarpa Pr. 1849; Acrostichum Hk. 1864. HB. 413; (Amphisoria Trevis. 1851). (an Acrostichum tripinatum Blanco 1837?).
appendiculata (Willd.) J. Sm. JoB. **4**. 150. 1841. Chr. 43. NPfl. 196 — Asia trop. Acrostichum Willd. sp. **5**. 114. 1810. HB. 415; Lacaussadea Gaud. 1846; Egenolfia J. Sm. 1866 — Gymnogramme auriculata Klf. 1824 — Polybotrya marginata Bl. 1828 — Gymnogramme rhizophylla Klf. 1824; Polybotrya Pr. 1836; Lacaussadea Gaud. 1846; Egenolfia Fée 1850-52 — Acrostichum Hamiltonianum Wall. 1828; Egenolfia Schott 1834; Polybotrya Pr. 1836; Lomariopsis Mett.; Salom. 1883 — Acrostichum ludens Wall. 1829; Poecilopteris Moore 1857 — etc. vide Egenolfia et Lacaussadea.

articulata J. Sm. JoB. **3**. 401. 1841 (nomen): Fée. Acrost. 74 t. 37. 1845. Chr. 42. NPfl. 198 — Celebes-Ins. Societatis. — Acrostichum Hk. 1864. HB. 413; Teratophyllum Mett.; Kuhn 1869; Arthrobotrya J. Sm. 1875.

aspidioides Gris. Cat. Pl. Cub. 276. 1866 — Cuba.
Acrostichum Bak. 1868. HB. 414.

asplenifolia Pr. Tent. 231. 1836 = P. appendiculata.

aurita Bl. Fl. Javae Fil. 15 t. 1. 1828 = Stenosemia.

bifurcata J. Sm. JoB. **4**. 150. 1841 = Elaphoglossum furcatum.

Blumeana Mett. Fil. Lips. 24 t. 2 f. 10. 1856 = Leptochilus lomarioides.

botryoides (Bak.) C. Chr. Ind. 6. 1905; 504. 1906 — Columbia.
Acrostichum Bak. JoB. **1881**. 206.

canaliculata Kl. Linn. **20**. 429. 1847 — Venezuela. Costa Rica.
Acrostichum Hk. 1864. HB. 415.

caudata Kze. Linn. **9**. 23. 1834 — Amer. trop. — Olfersia Kze. 1848; Psomiocarpa Pr. 1849; Acrostichum Hk. 1864. HB. 414; (Amphisoria Trevis. 1851).

cervina (L.) Klf. Enum. 55. 1824. Chr. 45. NPfl. 198 — Amer. trop.
Osmunda L. sp. **2**. 1065. 1753; Acrostichum Sw. 1806. HB. 416; Olfersia Kze. 1824; Dorcapteris Pr. 1849 — Olfersia corcovadensis Raddi 1819; Polybotrya Spr. 1827. P. Raddiana Link 1833. Acrostichum linearifolium Pr. 1822.

cicutaria Bl. Enum. 100. 1828 = Stenosemia aurita var.

Coenopteris Christ, Farnkr. 44. 1897 = P. serratifolia.

corcovadensis Spr. Syst. **4**. 33. 1827 = P. cervina var.

cyathifolia Fée et L'Herm.; Fée, 11 mém. 6 t. 2. 1866 — Guadeloupe.

cylindrica Klf. Enum. 56. 1824 = P. osmundacea.

exaltata Brack. Expl. Exp. **16**. 78. 1854 = P. appendiculata.

flabellata Mett. Ann. sc. nat. V. **2**. 206. 1864 = Elaphoglossum.

foeniculacea J. Sm. JoB. **4**. 150. 1841 = Elaphoglossum.

fractiserialis (Bak.) J. Sm. Hist. Fil. 133. 1875 — Peru.
Acrostichum Bak. Syn. 414. 1868.

fraxinifolia Mett. Fil. Lechl. **2**. 8. 1859 = Leptochilus serratifolius.

frondosa Fée, Cr. vasc. Br. **1**. 15. 1869 = P. osmundacea var.

fulvostrigosa Christ. Bull. Boiss. II. **1**. 70. 1901 — Alto Amazonas.

furcata Mett.; Kuhn. Fil. Afr. 52. 1868 = Elaphoglossum.

Gaudichaudiana Moore, Ind. XV. 1857 = P. appendiculata.

glandulosa Mett.; Kuhn. Linn. **36**. 65. 1869 — Brasilia.

Hamiltoniana Pr. Tent. 231. 1836 = P. appendiculata var.

Hartii (Bak.) C. Chr. Ind. 11. 1905; 504. 1906. — Trinidad.
Acrostichum Bak. JoB. **1888**. 371.

Helferiana Kze. Farnkr. **2**. 35 t. 114. 1848 = P. appendiculata var.

incisa Link, Hort. Berol. **2**. 135. 1833 = P. osmundacea var.

intermedia J. Sm. JoB. **3**. 401. 1841; Fée Acrost. 76 t. 40 f. 1. 1845 = P. appendiculata.

juglandifolia Christ, Bull. Boiss. II. **4**. 965. 1904 = P. Kalbreyeri.

Kalbreyeri C. Chr. Ind. 12. 1905; 504. 1906 — Columbia. Costa Rica.
Acrostichum juglandifolium Bak. JoB. **1881**. 207; Ic. pl. t. 1691 (non Klf. 1824); Polybotrya Christ 1904.

Kunzei Ettingsh. Denkschr. Ak. Wien **22**. 66 f. 2. 1864 = P. serratifolia.

Lechleriana Mett. Fil. Lechl. **1**. 4 t. 1 f. 1-5. 1856. Chr. 42. NPfl. 198 — Ecuador. Peru. — Acrostichum Hk. 1864. HB. 415 (non Mett. 1856).

lomarioides Mett. Fil. Lechl. **2**. 3. 1859 — Peru.

lomarioides Kuhn, Ann. Lugd. Bat. **4**. 295. 1869 = Leptochilus.
marattioides Brack. Expl. Exp. **16**. 79. 1854 = Asplenium multilineatum.
marginata Bl. Enum. 100. 1828 = P. appendiculata.
Meyeriana Mett. Fil. Lips. 24 t. 1. f. 4, 7. 1856 = Stenochlaena tenuifolia.
montana Moore, Ind. 350. 1862 = P. appendiculata.
nana Fée, Acrost. 75 t. 38 f. 1. 1845 = Blechnum lanceolatum.
neglecta Fée, Acrost. 76 t. 39 f. 2. 1845 = P. appendiculata.
Nieuwenhuisenii Racib. Bull. int. Ac. Cracovie **1902**. 57 — Borneo.
nodiflora Bory, Bél. Voy. Bot. **2**. 17. 1833 = P. appendiculata.
nutans Kze. Linn. **9**. 24. 1834 — Peru. — Acrostichum Hk. 1864.
orientalis Bl. Enum. 99. 1828 = Stenosemia aurita.
osmundacea H.B.Willd. sp. **5**. 99. 1810 Chr. 42. NPfl. 198 — Amer. trop.
Acrostichum Hk. 1864. HB. 415 — Polybotrya cylindrica Klf. 1824; Acrostichum Hk. 1864 — Polybotrya incisa Link 1833: Psomiocarpa Pr. 1849; Acrostichum Hk. 1864 — ? Osmunda filiculifolia L. 1753; Anemia [Sw. 1806.
peltata J. Sm. JoB. **4**. 150. 1841 = Elaphoglossum.
plumbicaulis (Bak.) J. Sm. Hist. Fil. 133. 1875 — Peru.
Acrostichum Bak. Syn. 413. 1868.
? Plumieri Moore, Ind. 14. 1857 = Acrostichum serrulatum Willd.
polybotryoides (Bak.) Christ. Bull. Boiss. II. **1**. 70. 1901 — Columbia.
Acrostichum Bak JoB. **1881**. 207; Ic. pl. t. 1690.
polyphylla (Brack.) C. Chr. Ind. 15. 1905; 505. 1906 — Polynesia.
Lomagramma Brack Expl. Exp. **16**. 83 t. 12 f. 3. 1854; Neurocallis Moore 1857; Acrostichum Hk. 1864. HB. 424; Lomariopsis Kuhn 1869: Chrysodium Luerss. 1871; ? Poecilopteris Carr. 1873 — ? Aspidium sorbifolium Willd. 1810; Nephrodium Pr. 1825; Sagenia Pr. 1836.
prolifera Bory, Bél. Voy. Bot. **2**. 18. 1833 = Leptochilus.
prolifera Mett. Fil. Lips. 24. 1856 = Leptochilus subcrenatus.
pterioides Kuhn, Ann. Lugd. Bat. **4**. 295. 1869 = Leptochilus lomarioides var.
pubens Mart.; Kze. Linn. **9**. 22. 1834; Mart. Ic. Cr. Bras. 87 t. 25. 1834 — Guiana. Brasilia. Peru. — Acrostichum Bak. 1868. HB. 414.
quercifolia Mett. Fil. Lechl. **2**. 12. 1859 = Leptochilus zeylanicus.
Raddiana Klf.; Link, Hort Berol. **2**. 134. 1833 = P. cervina var. corcovadensis.
rhizophylla Pr. Tent. 231. 1836 = P. appendiculata var.
scandens (Raddi) Christ, Bull. Boiss. II. **4**. 965. 1904 — Brasilia. Costa Rica.
Aspidium Raddi, Pl. Bras. **1**. 34 t. 49. 1825; Polybotrya acuminata Klf.; Link 1833; Psomiocarpa Pr. 1849; Acrostichum Hk. 1864. HB. 414.
scandens Fée, Gen. 47. 1850-52 — Mexico.
semipinnata Fée, Cr. vasc. Br. **1**. 16. 1869 = P. caudata.
serratifolia (Fée) Kl. Linn. **20**. 430. 1847 — Amer. trop.
Soromanes Fée, Acrost. 82 t. 43 (»dentata«). 1845; Acrostichum Coenopteris Kze.; Hk. 1864. HB. 417; Polybotrya Christ 1897. 44. NPfl. 198. Soromanes Christ 1903. Botryothallus Kunzei Kl. 1846; Polybotrya Ettingsh. 1864 — P. sorbifolia Mett.; Kuhn 1869.
serrulata J. Sm. JoB. **3**. 401. 1841; Fée, Acrost. 76. 1845 = P. appendiculata.
sorbifolia Mett.; Kuhn, Linn. **36**. 64. 1869 = P. serratifolia var.
sorbifolia Keys. Pol. Cyath. Hb. Bung. 32. 1873 = Stenochlaena.
speciosa Schott, Gen. t. 7. 1834 = P. osmundacea.
stenosemioides (Bak.) Copeland, Polyp Philipp. 40. 1905 — Borneo. Ins. Philipp.
Acrostichum Bak. Journ. Linn. Soc. **22**. 230. 1887.

suberecta (Bak.) C. Chr. Ind. 19. 1905; 506. 1906 — Columbia.
Acrostichum Bak. JoB. **1881**. 207; Ic. pl. t. 1692.
subquinquefida Mett. Fil. Lechl. 12. 1859 = Leptochilus latifolius.
taccaefolia Mett. Fil. Lechl. 12. 1859 = Leptochilus latifolius.
tenuifolia Kuhn, Fil. Afr. 52. 1868 = Stenochlaena.
trapezoides Link. Fil. sp. 164. 1841; Kze. Linn. **16**. 310. 1842 = P. osmundacea.
trilobata Mett. Fil. Lips. 24. 1856 = Leptochilus latifolius.
tripartita J. Sm. JoB **4**. 150. 1841 = Elaphoglossum flabellatum.
vivipara Hk. Exot. Flora **2** t. 107. 1825 = P. appendiculata var.
Wilkesiana Brack. Expl. Exp. **16**. 80 t. 10. 1854 — Polynesia.
Acrostichum Hk. 1864. HB. 413.

POLYCAMPIUM Presl, Epim. Bot. 135. 1849 = **Cyclophorus.**
hastatum Pr. Epim. 137. 1849.
Lingua Pr. Epim. 136. 1849.
penangianum Pr. Epim. 136. 1849.

POLYDICTYUM Presl, Epim. Bot. 52. 1849 = **Aspidium.**
heterophyllum Pr. Epim 259. 1849 = A. menyanthidis vel A. cicutarium.
hippocrepis Pr. Epim. 259. 1849.
menyanthidis Pr. Epim. 53. 1849.

POLYGRAMMA Presl, Epim. Bot. 156. 1849 = **Plagiogyria** (semicordata).

Polylepidum Moore, Ind. Fil. XVI. 1857 *(Elaphoglossum §)* = Elaphoglossum.

POLYPODIUM Linné (Gen. pl. 784); spec. plant. **2**. 1082. 1753 — (Polypodiaceae Gen. **100**).

Subgenera: C = **Campyloneurum.**
G = **Goniophlebium.**
Gr = **Grammitis.**
L = **Lomagramme.**
M = **Marginaria.**
My = **Myrmecophila.**
P = **Eupolypodium.**
Ph = **Phlebodium.**
Pl = **Pleopeltis.**
S = **Selliguea.**

abbreviatum Sieb. Syn. n. 168 = ? Dryopteris brachyodus.
abbreviatum Mett. Fil. Lips. 33. 1856; Pol. 130 n. 266. 1857 = Cyclophorus.
(P) **abietinum** Eat. ap. Mann, Proc. Amer. Acad. 7. 219. 1868 — Ins. Hawaii.
(P) **abitaguae** Hk. sp. **4**. 206. 1862. HB 333. NPfl. 310 — Ecuador.
abruptum Desv. Prod. 239. 1827 — Ind. occ. — (Dryopteris sp.).
(Pl) **accedens** Bl. Enum. 121. 1828. Mett. n. 165. HB. 353. Chr. 102. NPfl. 316 — Malesia. Polynesia. — Pleopeltis Moore 1857; Phymatopsis J. Sm. 1875. Polypodium lyciaefolium Bory 1828 (an nomen opt.?). P. cuspidiflorum Reinw. hb.; Drynaria J. Sm. 1841; Craspedaria Fée 1850-52; Pleuridium J. Sm. 1861.
(P) **achilleifolium** Klf. Enum. 116. 1824. Mett. n. 59. HB. 338. Chr. 86. NPfl. 311 — Brasilia. Ecuador. — Ctenopteris J. Sm. 1875. Polypodium piligerum Hk. 1841.
acrodontium Fée. Cr. vasc. Br. **2**. 58 t. 97 f. 2. 1872-73 = P. pilosissimum.
S **acroscopum** Christ. Journ. de Bot. **19**. 75. 1905 — Annam.
acrosorum Kze. Linn. **9**. 39. 1834 = P. crassifolium.

acrostichoides Forst. Prod. 81. 1786 = Cyclophorus.
acrostichoides Poir. Enc. **5**. 529. 1804 = Polystichum.
acrostichoides Willd. sp. **5**. 156. 1810 = Cyclophorus albicans.
aculeatum L. sp. **2**. 1090. 1753 = Polystichum.
aculeatum Raddi, Opusc. sci. Bol. 288. 1819; Pl. Bras. **1**. 27 t. 42. 1825 = Alsophila microdonta.
aculeatum Vell. Fl. flum. **11** t. 65. 1827; Arch. Mus. Nac. Rio Jan. **5**. 446 1881 = Cyatheacearum sp.
acuminatum Houtt. Pfl. Syst. **13**[1]. 204 t. 99 f. 2. 1786 = Dryopteris unita.
acuminatum Roxb.; Don, Prod. Fl. Nepal. 2. 1825 — Nepal.
acuminatum Sod. Cr. vasc. quit. 354. 1893 = P. brasiliense var.
acutangulum Pr. Tent. 181. 1836 (nomen) — ? — (Dryopteris sp.).
(Pl) ***acutilobum*** (Pr.) Ettingsh. Farnkr. 53 t. 27 f. 1. 1865 — Asia.
Pleuridium Pr. msc.
acutissimum Wall. List pag. 250 (n. 4727). 1829 = P. lineare.
acutum Burm. Fl. Ind. 232. 1768 = P. phymatodes.
acutum Roxb. Calc. Journ. **4**. 492. 1844 — Amboyna. — (Dryopteris sp.).
(P) **adelphum** Maxon. Contr. U. S. Nat. Herb. 8. 275 t. 62. 1903 — Mexico. Guatemala.
(P) **adenophorus** Hk. et Arn. Bot. Beech. Voy. 104 t. 32. 1832. ? Mett. n. 66. HB. 328. NPfl. 310 — Ins. Hawaii. Peru. ? Guadeloupe.
Ctenopteris J. Sm. 1875. Adenophorus pinnatifidus Gaud. 1827 (non [Polypodium Gilib. 1792).
adfine Bl. Enum. 126. 1828 = P. affine.
adhaerens Wall. List n. 2223. 1829 (nomen) — India.
adianthifolium Poir. Enc. **5**. 540. 1804 = Asplenium lanceolatum.
adianthiforme Blanco. Fl. Filip. ed. II. 573. 1845 = ? Nephrolepis acuminata.
adianthoides Burm. Fl. Ind. 234. 1768 — India.
adianthoides Aubl. Hist. Pl. Guian. **2**. 962. 1775 — Guiana.
adiantiforme Forst. Prod. 82. 1786 = Polystichum.
adnascens Sw. Syn. 25. 222 t. 2 f. 2. 1806 = Cyclophorus.
adnatum Wall. List n. 328. 1828 (nomen) = Dryopteris brunnea.
(G) **adnatum** Kze.; Kl. Linn. **20**. 395. 1847. Mett. n. 126. HB. 345. NPfl. 314 — Guatemala-Guiana-Ecuador. Ins. Galapagos.
Goniophlebium Moore 1862 — Polypodium Richardii Kl. 1847.
adspersum Schrad. Gött. gel. Anz. **1818**. 915 = ? P. lanceolatum.
adspersum Bl. Enum. 113. 1828 = P. hirtellum var.
adspersum Bak. vide aspersum.
aemulum Ait. Hort. Kew. **3**. 466. 1789 = Dryopteris.
(P) **aequale** Maxon, Contr. U. S. Nat. Herb. 8 273 t. 61 f. 1. 1903 — Guatemala.
affine Forst. Prod. 83. 1786 = Cyathea.
affine Bl. Fl. Jav. Fil. 166 t. 69. 1829 = P. commutatum.
affine Mart. et Gal. Mém. Ac. Brux. **15**. 37 t. 8 f. 1. 1842 = P. Martensii.
affine Lowe. Ferns **2** t. 50. 1858 = Dryopteris obliterata.
africanum Desv. Prod. 239. 1827 = Dryopteris.
africanum Mett. Pol. 131 n. 268 t. 3 f. 11-12. 1857 = Cyclophorus.
alatellum Christ. Verh. Nat. Ges. Basel **11**. 441. 1896 = P. congenerum.
alatum L. sp. **2**. 1086. 1753 = Dryopteris scolopendrioides.
alatum Hk. sp. **5**. 85. 1863 = P. Wilkesii.
Alberti Regel, Acta Hort. Petr. **7**. 622. 1881 = P. lineare var.
albicans Mett. Pol. 127 n. 257. 1857 = Cyclophorus.

(Pl) **albidosquamatum** Bl. Enum. 132. 1828. Mett. n. 207. Chr. 114 — Malesia. N. Guinea — Pleopeltis Pr. 1836; Drynaria J. Sm. 1841; Polypodium albosquamatum Hk. 1863. HB. 369; Phymatodes J. Sm. 1866 — Polypodium varians Bl. 1828. Mett. n. 206; Pleopeltis Moore 1862.

(P) **albidulum** Bak. Fl. bras. 1[2]. 598. 1870. HB. 509 — Brasilia.

albidum Pr. Del. Prag 1. 165. 1822 = P. polypodioides.

(P) **albobrunneum** Bak. Fl. Maur. 505. 1877 — Ins. Sechellae.

albopunctatum Raddi. Opusc. sci. Bol. 287. 1819; Pl. Bras. 1. 21 t. 30. 1825 = P. brasiliense var.

albopunctatum Bak.; Jenm. JoB. 1877. 265 = ? P. subtile.

(Gr) **albosetosum** Bailey. Rep. Govt. Sci. Exp. Bellenden-Ker 78. 1889; Syn. Queensl. Fl. Suppl. III. 94. c. fig. 1890 — Queensland.

albosquamatum Hk. sp. 5. 92. 1863 = P. albidosquamatum.

(Pl) **albulum** Christ in C. Chr. Ind. 508. 1906 — Celebes.
P. lagopodioides Christ, Ann. Jard. Buit. II. 4. 37. 1904 (non Desv. 1811).

album Lam. Fl. franç. 1. 21. 1778 = Cystopteris fragilis.

alcicorne Bak. JoB. 1888. 229 = Polystichum.

(P) **Alfari** Donn, Smith, Bot. Gaz. 33. 262. 1902 — Costa Rica.

alleganum Poir. Enc. 5. 545. 1804 = Dryopteris noveboracensis.

alloeopterum Kze. Linn. 25. 748. 1853 = Dryopteris

alpestre Spenn. Fl. Frib. 1. 12. 1825 = Athyrium.

alpestre Bl. Enum. 123. 1828 = P. hirtellum.

alpinum Lam. Fl. franç. 1. 22. 1778 = Asplenium fontanum.

alpinum Jacq. Coll. 2. 171. 1788 = Cystopteris fragilis *regia.

alsophiloides Liebm. Vid. Selsk. Skr. V. 1. 208 (seors. 56). 1849 = Dryopteris subincisa *martinicensis.

alsophiloides Bak. JoB. 1879. 297 = Dryopteris waiwaiensis.

alsophilum Link, Hort. Berol. 2. 106. 1833 = Alsophila paleolata.

alternans Wall. List n. 329. 1828 = Cyathea.

(P) **alternidens** Ces. Atti Ac. Napoli 7[8]. 25 t. 2 f. 4. 1876 — Borneo. Celebes.

(Pl) *alternifolium* Willd. sp. 5. 168. 1810 — India.
Phymatodes Pr. 1836; ? Drynaria Brack. 1854; Pleopeltis Moore 1862. (prob. = P. longissimum Bl. vel P. nigrescens).

alternifolium Link, Hort. Berol. 2. 93. 1833 = P. phymatodes.

alternifolium Mett. Pol. 107 n. 204 b. t. 1 f. 35. 1857 = P. palmatum.

alternifolium Hk. sp. 4. 222 t. 277 A. 1862 = P. longum.

altescandens Colla, Mem. Acad. Torino 39. 48. 1836 = Arthropteris.

altum Boj. Hort. Maur. 417. 1837 = P. punctatum.

ambiguum Desv. Prod. 242. 1827 — Brasilia. — (Alsophila? sp.).

ambiguum Bl. Enum. addend. 1828 = P. punctatum.

ambiguum Mett.; Kuhn, Linn. 36. 134. 1869 = P. ursipes.

Americanum Hk. sp. 5. 54. 1863 = Cyclophorus.

ammifolium Poir. Enc. 5. 554. 1804 = Polystichum aculeatum var. 1.

(G) **amoenum** Wall. List n. 290. 1828 (nomen); Mett. Pol. 80 n. 131. 1857. HB. 341. Chr. 93. NPfl. 311 — India bor. China. Formosa.
Marginaria Pr. 1836; Goniophlebium J. Sm. 1840.

(S) **ampelideum** Christ, Journ. de Bot. 19. 78. 1905 — Annam.

(P) **amphidasyon** Kze.; Mett. Pol. 49 n. 51. 1857 — Merida.

amphostenon Kze.; Kl. Linn. 20. 399. 1847 = P. angustifolium var.

amplum H.B.Willd. sp. 5. 207. 1810 = Dryopteris.

amplum Col. Tr. N. Zeal. Inst. **24**. 396. 1892 = ? Dryopteris punctata.
(S) **anceps** (Christ) C. Chr. Ind. 509. 1906 — Yunnan.
Selliguea Christ, Bull. Boiss. **6**. 879. 1898.
andaiense Bak. Malesia **3**. 45. 1886 = Aspidium.
(P) **andinum** Hk. 2 Cent. t. 6. 1860. HB. 323. NPfl. 308 — Costa Rica-Ecuador.
andinum Karst. Fl. Col. **1**. 171 t. 85. 1861 = P. surucuchense.
androgynum Poir. Enc. **5**. 535. 1804 = Dryopteris tetragona.
Aneitense Hk. sp. **4**. 268. 1862 = Dryopteris.
(P) **anfractuosum** Kze.: Kl. Linn. **20**. 375. 1847. Mett. n. 39 — Columbia.
angelicaefolium Schum. Vid. Selsk. Afh. **4**. 228 (seors. 454). 1827 = Aspidium.
angulare Fries, Nov. Fl. Suec. Mant. **1**. 20. 1832 = Polystichum Braunii.
angustilobum Ettingsh. Farnkr. 51 t. 36 f. 8. 1865 — Australia.
(Drynaria Pr. hb.).
angulatum Willd. sp. **5**. 185. 1810 = Aspidium.
angustatum Sw. Syn. 27, 224. 1806 = Cyclophorus.
angustatum Bl. Enum. 133. 1828 = P. palmatum var.
(C) **angustifolium** Sw. Prod. 130. 1788. HB. 347. Chr. 95. NPfl. 314 — Amer. trop.
Marginaria Pr. 1836; Grammitis Hew. 1838; Cyrtophlebium J. Sm. 1846; Campylonevron Fée 1850-52; Goniophlebium Brack. 1854 — Polypodium ensifolium Willd. 1810; Marginaria Pr. 1836; Goniophlebium Brack. 1854; Campyloneuron J. Sm. 1857 — Polypodium taeniosum H.B.Willd. 1810. Mett. n. 136; Marginaria Pr. 1836; Campylonevron Fée 1850-52 — Polypodium dimorphum Link 1833; Marginaria Link 1841; Campyloneuron Fourn. 1872 — Polypodium solutum Kl. 1847; Goniophlebium Fée 1850-52; Campylonevron Fée 1850-52 — Polypodium amphostenon Kze.; Kl. 1847; Campylonevron Fée 1850-52 — C. Jamesoni Fée 1850-52; Polypodium Mett.; Salom. 1883.
—* Campyloneurum lucidum Moore 1861 (nomen); Polypodium Beyr.: Hk. 1863. HB. 348 (non Roxb. 1844).
angustifrons Kze. Linn. **13**. 134. 1839 — Mexico.
Lastrea Moore 1858. (Dryopteris? sp.).
angustissimum Fée, Cr. vasc. Br. **2**. 55 t. 96 f. 3. 1872-73 = P. moniliforme.
angustissimum Bak. Ann. Bot. **5**. 472. 1891 = Cyclophorus taeniodes.
(Pl) **angustum** (H.B.Willd.) Liebm. Vid. Selsk. Skr. V. **1**. 186 (seors. 34). 1849. Mett. n. 158. HB. 363. Chr. 110 — Amer. trop — Pleopeltis H.B.Willd. sp. **5**. 211. 1810; Phlebodium J. Sm. 1841; Drynaria Fée 1850-52; Lepicystis Diels 1899. NPfl. 323; Pteropsis angustifolia Fourn. 1872 (non alior.) — Polypodium sectifrons Kze.: Mett. 1857. n. 181.
anisomeron Fourn. Mex. pl. **1**. 81. 1872 = P. plesiosorum.
anisopterum Kze. Bot. Zeit. **1848**. 118 = Dryopteris Hasseltii.
(Pl) **Annabellae** Forbes, JoB. **1888**. 33 t. 280. NPfl. 316 — N. Guinea.
(S) **annamense** Christ, Journ. de Bot. **19**. 77. 1905 — Annam.
(Pl) **annuifrons** Mak. Bot. Mag. Tokyo **13**. 48. 1899; Phan. et Pterid. Jap. Ic. **1** t. 7. 1899 — Japonia. — P. japonense Mak. Bot. Mag. Tokyo **12**. 165. 1898 (nomen?).
anocarpos Kze. Linn. **9**. 40. 1834 = P. crassifolium.
anomalum Hk. et Arn.; Hk. JoB. **8**. 360 t. 11. 1856 = Polystichum aculeatum var. 37.
anomalum Christ, Bull. Boiss. **6**. 201 t. 3 f. 3. 1898 = P. heterolobum.
anthriscifolium Hoffm. in Röm. et Usteri Mag. **9**. 11 t. 14 f. C. 1790 = Cystopteris fragilis.

(P) **antioquianum** Bak. JoB. **1881**. 205 — Columbia.
aoristisorum Harr. Journ. Linn. Soc. **16**. 30. 1877 = Dryopteris.
aphanophlebium Kze. Bot. Zeit. **1845**. 288 = P. laevigatum.
apicidens Bak. Kew Bull. **1895**. 54 = Dryopteris.
(P) **apiculatum** Kze.; Kl. Linn. **20**. 378. 1847. Mett. n. 36. HB. ed. II. 332. NPfl. 309 — Guiana-Brasilia-Peru. — P. leptophyllum Moritz: Eat. 1860 (non L. 1753) — P. confluens Fée 1869 (non alior.).
appendiculatum Poir. Enc. **5**. 533. 1804 = Dryopteris deltoidea.
appendiculatum Sw. Fl. Ind. occ. **3**. 1677. 1806 = Aspidium cicutarium.
appendiculatum Wall.; Bak. Syn. 306. 1867 = ? Dryopteris cana.
appendiculatum Kl. Allg. Gartenzeit. **1855**. 377 = P. plesiosorum var.
appendiculatum Bedd. Ferns br. Ind. t. 256. 1868 = Dryopteris squamaestipes.
aquilinum Thouars, Fl. Trist. d'Ac. 32. 1804 = Dryopteris.
Araguato Moritz, Bot. Zeit. **1854**. 855 (nomen) = Aspidium.
araneosum Mart. et Gal. Mém. Ac. Brux. **15**. 33 t. 5 f. 2. 1842 = P. aureum.
arborescens Roxb. Calc. Journ. **4**. 495. 1844 = ? Cyathea crenulata.
arboreum L. sp. **2**. 1092. 1753 = Cyathea.
arboreum Lour. Fl. Cochinch. **2**. 831. 1790 = Cyathea Rumphii.
arboreum Roxb.; Wall. List n. 2226. 1829 = ? Cyathea crenulata.
arcuatum Poir. Enc. **5**. 528. 1804 = Dryopteris serra.
arcuatum Moritz; Mett. Pol. 56 n. 67 b. 1857 (nomen) = P. suspensum.
(Pl) **arenarium** Bak. Kew Bull. **1895**. 56. — Yunnan.
areolatum H.B.Willd. sp. **5**. 172. 1810 = P. aureum*.
arfakianum Bak. Malesia **3**. 45. 1886 = Dryopteris.
argentatum Jacq. Hort. Schoenbr. **4**. 23 t. 447. 1804 = Polystichum adiantiforme.
(G) **argutum** Wall. List n. 308. 1828 (nomen); Hk. sp. **5**. 32. 1863. HB. 511. NPfl. 312 — Himalaya. — Marginaria Hk. 1838; Goniophlebium J. Sm. 1840; Schellolepis J. Sm. 1875.
(P) **argyratum** Bory; Willd. sp. **5**. 175. 1810. Mett. n. 70. HB. 331. Chr. 84. NPfl. 311 — Ins Mascarenae. — Marginaria Bory 1826.
argyratum var. brasiliana Fée, Cr. vasc. Br. **1**. 265. 1869 = P. albidulum.
(G) **argyrolepis** Sod. Cr. vasc. quit. 349. 1893 — Ecuador.
argyrophanes Spr. Syst. **4**. 51. 1827 = P. argyratum.
aristatum Forst. Prod. 82. 1786 = Polystichum.
aristatum Vill. Hist. Pl. Dauph. **3**. 844. 1789 = Dryopteris spinulosa *dilatata.
armatum Sw. Prod. 134. 1788 = Alsophila.
(P) **aromaticum** Maxon, Proc. U. S. Nat. Mus. **27**. 743. 1904; Bull. Dep. Agric. Jam. **2**. 177. 1904 — Jamaica.
(P) **arthropodium** Fée, 8 mém. 88. 1857 — Mexico.
arthrothrix Hk. sp. **5**. 14. 1863 = Dryopteris.
articulatum Juss.; Poir. Enc. **5**. 514. 1804 = Oleandra.
articulatum Vahl, Ecl. Amer. **3**. 51. 1807 = Oleandra nodosa.
articulatum Desv. Prod. 236. 1827 = P. fraxinifolium.
arvonicum Wither. Bot. Arrang. ed. III. **3**. 774. 1796 = Woodsia ilvensis.
(P) **ascensionense** Hier. Hedwigia **44**. 93. 1905 — Ascension.
asininum (Spruce) Salom. Nom. 289. 1883 (nomen) — Peru. — Grammitis Spruce.
aspersum Bak. Kew Bull. **1898**. 231 = P. mengtzeense.
asperulum J. Sm. JoB. **3**. 394. 1841 = Dryopteris.
asperum L. sp. **2**. 1093. 1753 = Alsophila.
asperum Pr. Rel. Haenk. **1**. 24 t. 3 f. 4. 1825 = Dryopteris urophylla.

aspidioides Pr. Del. Prag. **1**. 170. 1822 = ? Dryopteris tenuifolia.

aspidioides Bail. Proc. Linn. Soc. N. S. Wales **5**. 32. 1881; Lithogr. t. 145-146 — Queensland. (Dryopteris? sp.).

(M) **aspidiolepis** Bak. JoB. **1887**. 26 — Costa Rica.

aspidiolepis Bak. Ann. Bot. **5**. 474. 1891 (err. script.) = P. excavatum var.

(P) **asplenifolium** L. sp. **2**. 1084. 1753. Mett. n. 68. (HB. 329) — Amer. trop. Fernando Po. — Phegopteris Fée 1850-52.

asplenifolium β L. sp. **2**. 1084. 1753 = P. cultratum.

asplenioides Sw. Schrad. Journ. **1800**[2]. 26. 1801 = Dryopteris.

asplenioides Bory, Bél. Voy. Bot. **2**. 33. 1833 — Java.

assimile? Kze. Linn. **23**. 275. 1850 (nomen) = P. chnoodes.

assimile Bak. JoB. **1875**. 201 = Cyclophorus.

asterolepis Bak. JoB. **1888**. 230 = P. excavatum var.

asterosorum Bak. JoB. **1880**. 214 = Cyclophorus.

asterothrix Hk. sp. **5**. 6. 1863 = Dryopteris reptans*.

(M) *astrolepis* Liebm. Vid. Selsk. Skr. V. **1**. 185 (seors. 33). 1849 — Mexico Pleopeltis Fourn. 1872. (P. lanceolatum var.?).

(M) **atacamense** Bak. Journ. Linn. Soc. **22**. 64. 1887 (nomen) — Peru. Atacama. P. squamatum Phil. Linn. **29**. 107. 1857 (nomen); Anal. Univ. Chil. **94**. 357. 1896 (non L. 1753).

(P) **athyrioides** Hk. sp. **4**. 224. t. 277 B. 1862. HB. 339 — Peru.

(G) **Atkinsoni** C. Chr. Ind. 511. 1906 — Himalaya. — P. Hendersoni Atkins.; Bak. Syn. 511. 1874 (non Lowe 1858); Goniophlebium Bedd. 1876.

atropunctatum Gaud. Freyc. Voy. Bot. 346. 1827 = P. lineare.

atrovirens Langsd. et Fisch. Ic. Fil. 12 t. 14. 1810 = Alsophila.

(G) **attenuatum** H.B.Willd. sp. **5**. 191. 1810. HB. 345. NPfl. 314 — Amer. trop. Goniophlebium Pr. 1836; Polypodium dissimile Schkuhr. Kr. Gew. **1**. 14 t. 14. 1806 (non L. 1753) — P. gladiatum Kze. 1834; Mett. n. 122. Marginaria Pr. 1836; Goniophlebium Fée 1850-52 — Polypodium xiphophoron Kze.; Mett. 1857. n. 107; Goniophlebium Fée 1869.

attenuatum R.Br. Prod. Fl. N. Holl. 146. 1810 = P. Brownii.

attenuatum Rich. Fl. N. Zél. 62. 1832 = P. dictyopteris.

attenuatum Roxb. Calc. Journ. **4**. 482. 1844 — Pulo Penang.

attenuatum Kze.; Ettingsh. Farnkr. t. 32 f. 2. 1865 = P. aureum.

(P) **aturense** Maury, Jour. de Bot. **3**. 129 f. 3. 1889 — Orinoco.

Aubertii Desv. Prod. 243. 1827 = ? Dryopteris cruciata.

auratum Vell. Fl. flum. **11**. t. 74. 1827; Arch. Mus. Nac. Rio Jan. **5**. 448. 1881 = P. aureum.

(Ph) **aureum** L. sp. **2**. 1087. 1753. Mett. n. 182. HB. 347. Chr. 94. NPfl. 314 — Amer. trop. ? Australia (F. Müller).

Pleopeltis Pr. 1836; Chrysopteris Link 1841; Phlebodium J. Sm. 1841 — ? Polypodium dulce Poir. 1804; Chrysopteris Fée 1850-52; Goniophlebium Moore 1862 — Polypodium leucatomos Poir. 1804.

— *P. areolatum H.B.Willd. 1810. Mett. n. 183. NPfl. 314; Goniophlebium Pr. 1836; Pleopeltis Pr. 1836; Phlebodium J. Sm. 1841; Chrysopteris Fée 1857. Polypodium fulvum Mart. et Gal. 1842; Drynaria Fée 1850-52; Phlebodium Moore 1862. Chrysopteris lanosa Fée 1857; Phlebodium Moore 1862 — Polypodium sporadocarpum Willd. 1810. Mett. n. 185; Chrysopteris Link 1841 (= areolatum verum); Phlebodium J. Sm. 1846. Chrysopteris glauca Fée 1850-52 — Polypodium araneosum Mart. et Gal.

1842; Drynaria Fée 1850-52; Pleopeltis Moore 1862; Chrysopteris Fourn. 1872 — Polypodium glaucinum Mart. et Gal. 1842; Drynaria Fée 1850-52; Pleopeltis Moore 1862; Chrysopteris Fourn. 1872.

—* Polypodium pulvinatum Link 1833. Mett. n. 184; Chrysopteris Link 1841; Phlebodium J. Sm. 1846.

— vide Chrysopteris et Phlebodium.

auriculatum L. sp. **2**. 1088. 1753 = Polystichum.

auriculatum Raddi, Opusc. sci. Bol. **3**. 288. 1819 = Dryopteris grandis.

auriculatum Pr. Del. Prag. **1**. 167. 1822 = P. Preslii.

auriculatum Wall. List n. 314. 1828 = Dryopteris squamaestipes.

aurisetum Raddi, Pl. Bras. **1**. 12 t. 23 f. 1. 1825 = P. tectum.

auritum Lowe, Ferns **2**. t. 51. 1858 = Dryopteris.

australe Fée, Gen. 236 t. 20 A. f. 2. 1850-52 = P. vulgare var.

australe Mett. Pol. 36 n. 15. 1857 = P. Billardieri.

austriacum Jacq. Obs. Bot. **1**. 45. 1764 = ? Dryopteris spinulosa.

(Pl) **austrosinicum** Christ in C. Chr. Ind. 512. 1906 — China merid.
P. Henryi Christ, Bull. Boiss. **6**. 873. 1898 (non infra).

avenium Desv. Berl. Mag. **5**. 314. 1811 = P. percussum.

avenium Mett. Fil. Lips. 37. 1856 = P. Blumeanum.

axillare Ait. Hort. Kew. **3**. 466. 1789 = Athyrium filix femina*.

axillare Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 27 t. 41. 1825 = Alsophila leucolepis.

(P) **azuayense** Sod. Cr. vasc. quit. 323. 1893 — Ecuador.

bacciferum Poir. Enc. **5**. 554. 1804 = Dennstaedtia cicutaria.

(S) **Baileyi** (Bak.) C. Chr. Ind. 334. 1905; 512. 1906 — Queensland.
Gymnogramme Bak. Kew Bull. **1892**. 86; G. membranacea Bail. Syn. Queensl. Fl. Suppl. III. 94. 1890 (non Hk. 1864).

(Pl) **Bakeri** Luerss. Abh. Nat. Ver. Bremen **7**. 48. 1882 — Sumatra.
P. torulosum Bak. JoB. **1880**. 215 (non 1877).

Bakeri Lindm. Ark. för Bot. **1**. 240 t. 11 f. 9. 1903 = P. truncorum.

Balansæ Bak. Ann. Bot. **5**. 459. 1891 = Dryopteris alsophiloides.

(M) **balaonense** Hier. Engl. Jahrb. **34**. 529. 1904 — Ecuador.

Balbisii Spr. Nova Acta **10**. 228. 1821 = Dryopteris.

(Gr) **Baldwinii** Bak. Ann. Bot. **5**. 463. 1891 — Ins. Hawaii.

(M) **Bangii** Bak. Kew Bull. **1901**. 145 — Bolivia.

(P) **barathrophyllum** Bak. JoB. **1891**. 107 — Borneo.

(P) **barbatulum** Bak. Syn. 323. 1867. NPfl. 308 — Réunion. — P. ciliatum Bojer 1837 (nomen, non Willd. 1810) — P. Beaumontii Leperv.; Kuhn 1868.

barbatum Kze.; Spr. Syst. **4**. 57. 1827; Kze. Linn. **9**. 52. 1834 = Dryopteris hirta.

barbatum Desv. Prod. 242. 1827 = Dryopteris subincisa *Karsteniana.

barbatum Hk. sp. **5**. 10. 1863 = Dryopteris ferox.

Barberi Hk. sp. **5**. 100. 1864 = Aspidium.

Barometz L. sp. **2**. 1092. 1753 = Cibotium.

Baroni Bak. Journ. Linn. Soc. **22**. 535. 1887 = Dryopteris.

Baronii Christ. Nu. Giorn. bot. It. n. s. **4**. 100 t. 2. 1897 = Drynaria reducta.

Barterianum Hk. sp. **4**. 253. 1862 = Dryopteris.

(P) **basiattenuatum** Jenm. Bull. Dep. Jam. **4**. 114. 1897 — Jamaica.

Baudouini Bak. Ann. Bot. **5**. 480. 1891 = Drynaria.

Beaumontii Leperv.; Kuhn. Afr. 146. 209. 1868 = P. barbatulum.

Beccarianum Ces. Rend. Ac. Napoli **16**. 27. 30. 1877 = Aspidium.

Beckleri Hk. sp. 4. 224. 1862 = Arthropteris obliterata.

(G) **Beddomei** Bak. Syn. ed. II. 344. 1874. NPfl. 312 — Birma.
Goniophlebium molle Bedd. Ferns br. Ind. t. 206. 1867 (non Polypodium Schreb. 1771 nec Jacq. 1789); Polypodium puberulum Bak. Syn. ed. I. 344. 1867 (non Schlecht. et Cham. 1830).

Bergianum Schlecht. Adumbr. 20 t. 9. 1825 = Dryopteris.

(M) **Bernoullii** Bak. Syn. 510. 1874 (Bernouillii) — Guatemala.

Berterianum Spr. Syst. 4. 59. 1827 = Polystichum adiantiforme.

Berteroanum Hk. sp. 4. 269. 1862 = Dryopteris subincisa*.

Beyrichianum Pr. Tent. 178. 1836 = P. trichomanoides.

(P) **biauriculatum** Hk. Ic. pl. t. 121. 1837. HB. 336 — Peru.
Goniophlebium Moore 1862.

bicolor Mett. Pol. 125 n. 251. 1857 = Cyclophorus serpens.

bicuspe Bl. Enum. 125. 1828 = Cheiropleuria.

bifidum Hoffm. in Röm. et Usteri Mag. 9. 10. 1790 = Athyrium filix femina.

biforme Lour. Fl. Cochinch. 2. 827. 1790 — Cochinchina. — (Drynaria sp.?).

biforme Bak. Syn. 505. 1874 = Dryopteris.

(Pl) **bifrons** Hk. Fil. exot. t 52. 1858. HB. 361 — Ecuador.

bifurcatum Bak. Syn. ed. II. 362. 1874 = Dipteris Lobbiana.

(Gr) **Billardieri** (Willd.) C. Chr. Ind. 513. 1906 — Africa austr. N. Guinea. Australia. Amer. antarct. Ins. antarct. — Grammitis Willd. sp. 5. 139. 1810; G. australis R Br. 1810; Polypodium Mett. n. 15. 1857. HB. 322. Chr. 78. NPfl. 308 (non Fée 1850-52) — P. gramineum Poir. 1804 (non Sw. 1788) — Grammitis magellanica Desv. 1811 — Polypodium Poeppigianum Mett. n. 17. 1857 — (Asplenium angustifolium Jacq. 1786 verisim. adhuc pertinet). — vide Grammitis.

Billardieri R.Br. Prod. Fl. Nov. Holl. 147. 1810 = P. diversifolium.

Billardieri Fée, Gen. 236. 1850-52 = P. grammitidis.

binervatum Poir. Enc 5. 521. 1804 = Blechnum polypodioides.

? binerve Hk. sp. 4. 175 t. 273 B. 1862 = ? Stenochlaena pollicina.

(P) **bipinnatifidum** Bak. JoB. 1890. 109 — N. Guinea.

biseriale Bak. Syn. 309. 1867 = Dryopteris.

biserratum Mart. et Gal. Mém. Ac. Brux. 15. 38 t. 9 f. 1. 1842 = P. subpetiolatum.

(Gr) **bisulcatum** Hk. sp. 4. 164. 1862. HB. 321. Chr. 77. NPfl. 308 — Borneo.
Grammitis Hk. Ic. pl. t. 998. 1854; Holcosorus pentagonus Moore 1857.

bivestitum Tausch; Pr. Tent. 181. 1836 = Dryopteris.

(P) **Blanchetii** C. Chr. Bot. Tidsskr. 25. 78. 1902 — Brasilia.
P. exiguum Fée, Cr. vasc. Br. 1. 89 t. 37 f. 1. 1869 (non Hew. 1838).

blandum Fée, 7 mém. 59 t. 22 f. 5. 1857 = P. capillare.

blandum Bak. Ann. Bot. 5. 455. 1891 = Dryopteris.

blechnoides Rich. Act. Soc. Nat. Hist. Paris 1. 114. 1791 (nomen) = Alsophila rostrata.

(P) **blechnoides** (Grev.) Hk. sp. 4. 180. 1862. HB. 331. Chr. 85. NPfl. 310 — Polynesia. Queensland. Java. — Grammitis Grev. Ann. Mag. Nat. Hist. II. 1. 320 t. 17. 1848; Cryptosorus J. Sm. 1875. C. Seemanni J. Sm. 1861 — Polypodium contiguum Brack. 1854 (non Hk. 1854).

Blumeanum Pr. Tent. 290. 1836 = P. diplosorum.

(L) **Blumeanum** (Pr.) C. Chr. Ind. 60. 1905; 513. 1906 — Japonia. Malesia.
Loxogramme Pr. Tent. 215. 1836 (nomen); Selliguea Fée 1850-52; S.

Blumei Kze. 1846; Polypodium Mett. n. 219. 1857; Gymnogramme Franch. et Sav. 1876; Grammitis coriacea Bl. Enum. 117. 1828 (non Klf. 1827); Antrophyum Bl. Fl. Jav. Fil. 85 t. 37 f. 1 (non Polypodium Sw. 1788 nec Raddi 1819) — Grammitis avenia Bl. Enum. 117. 1828; Antrophyum Bl. Fl. Jav. Fil. 86 t. 37 f. 2. 1828; Loxogramme Pr. 1836; Selliguea Kze. 1846; Polypodium Mett. 1856. Pol. n. 220. Chr. 107 (non Desv. 1811); Gymnogramme Bak. 1868. HB. 388.

Blumei Mett. Pol. 113 n. 219 t. 3 f. 27-28. 1857 = P. Blumeanum.

(G) **Bodinieri** Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 203 — China.

(S) **Boisii** Christ, Journ. de Bot. **19**. 76. 1905 — Annam.

(P) **Boivini** Mett.; Kuhn, Fil. Afr. 146. 1868. HB. 507 — Madagascar.

Bojeri Hk. sp. **4**. 290. 1862 = Dryopteris cruciata.

(Pl) **boninense** Christ in Warb. Monsunia **1**. 61. 1900 — Bonin.

Bonplandii Desv. Prod. 241. 1827 = Dryopteris submarginalis.

Boothii Hk. sp. **5**. 53. 1863 = Cyclophorus.

boreale Salisb. Prod. —. 1796 = P. vulgare.

Borneense Hk. sp. **5**. 11. 1863 = Dryopteris labuanensis.

brachylepis Bak. Gard. Chr. n. s. **14**. 494. 1880 = P. Buergerianum.

brachyodus Kze. Linn. **9**. 48. 1834 = Dryopteris.

(P) **brachyphlebium** Bak. Ann. Bot. **8**. 129. 1894 — Sumatra.

Brackenridgei Hk. sp. **5**. 9. 1863 = Dryopteris subspinosa.

braineoides Bak. JoB. **1888**. 229 = Dryopteris.

?brancaefolium Pr. Rel. Haenk. **1**. 22. 1825 = Drynaria quercifolia.

(G) **brasiliense** Poir. Enc. **5**. 525. 1804. Chr. 94. NPfl. 314 — Amer. trop.

P. neriifolium Schkuhr 1806. Mett. n. 124. HB. 345; Marginaria Pr. 1836; Goniophlebium Hk. 1841 — Polypodium menisciifolium Langsd. et Fisch. 1810. Mett. n. 123. HB. 346; Marginaria Pr. 1836; Goniophlebium J. Sm. 1840 — Polypodium albopunctatum Raddi 1819; Goniophlebium J. Sm. 1846 — Polypodium longifolium Pr. 1822; P. Preslianum Link 1833. Mett. n. 125 — Goniophlebium acuminatum Fée 1866; Polypodium Sod. 1893 (non Houtt. 1786).

breve J. Sm.; Bak. Ann. Bot. **5**. 472. 1891 (ex errore?) = Cyclophorus laevis.

brevifolium Lodd.; Link, Hort. Berol. **2**. 90. 1833 = P. repens.

(P) **brevistipes** Mett.; Kuhn, Linn. **36**. 131. 1869 — Brasilia.

P. tenuiculum Fée 1869 (non 1850-52). HB. 509.

Brongniartii Bory, Dup. Voy. Bot. **1**. 263 t. 34. 1828 = Aspidium.

Brownianum Spr. Syst. **4**. index 1827; Link, Hort. Berol. **2**. 89. 1833 = P. Brownii.

(Pl) **Brownii** Wikstr. Vet. Ak. Handl. **1825**. 442. 1826. Mett. n. 148. HB. 355. NPfl. 315 — Australia. N. Caledonia. Fiji.

Drynaria (Browniana) Fée 1850-52; Pleopeltis Fourn. 1869; Polypodium attenuatum R.Br. Prod. Fl. N. Holl. 146. 1810 (non Willd. 1810); Dictyopteris Pr. 1836; Dictymia J. Sm. 1846.

(P) **brunneo-viride** Bak. ap. Jenm. JoB. **1877**. 265 — Jamaica.

brunneum Wall. List n. 333. 1828 = Dryopteris.

(Pl) **Buergerianum** Miq. Ann. Lugd. Bat. **3**. 170. 1867 — Japonia. China.

P. brachylepis Bak. 1880.

bulbiferum L. sp. **2**. 1091. 1753 = Cystopteris.

(Pl) **bullatum** Bak. Journ. Linn. Soc. **15**. 420. 1876 — Madagascar.

(P) **Burbidgei** Bak. JoB. **1879**. 42 — Borneo.

(S) **Büttneri** Kuhn, Verh. Bot. Ver. Brand. 31. 68. 1889 — Congo.
(G) **Caceresii** Sod. Cr. vasc. quit. 360. 1893 — Ecuador. Columbia.
(S) **Cadieri** Christ, Journ. de Bot. 19. 76. 1905 — Annam.
caducum H.B.Willd. sp. 5. 193. 1810 = Cyclopeltis semicordata.
? caesium Pr. Rel. Haenk. 1. 27. 1825 = Alsophila quadripinnata.
(Gr) **caespitosum** (Bl.) Mett. Ann. Lugd. Bat. 2. 219. 1866. (HB. 321) — Java. Ins. Philipp. — Grammitis Bl. Enum. 115. 1828; Fl. Jav. Fil. 106 t. 46 f. 1.
caespitosum Lodd.; Link, Hort. Berol. 2. 91. 1833 = P. repens.
caespitosum Bak. Syn. ed. II. 305. 1874 = Dryopteris.
caffrorum L. Mant. 307. 1771 = Mohria.
cajanense Desv. Berl. Mag. 5. 314. 1811 = P. ciliatum.
Calaguala Ruiz, Mem. leg. Calaguala c. tab. 1805 = P. angustifolium.
calcareum Sm. Brit. Fl. 1117. 1804 = Dryopteris Robertiana.
(G) **californicum** Klf. Enum. 102. 1824. HB. 341. Chr. 83. NPfl. 312 — U. S. pacif. Marginaria Pr. 1836; Goniophlebium Moore 1862. Polypodium intermedium Hk. et Arn. 1840 (non Willd. 1810).
californicum Schlecht. et Cham. Linn. 5. 606. 1830 = P. plesiosorum.
californicum Mett. Pol. 71 n. 104. 1857 = P. translucens.
Callipteris Ehrh Hannöv. Mag. 8-9. 127, 138. 1784 = Dryopteris cristata.
callolepis Fée, 8 mém. 86. 1857 = P. Martensii.
calvatum Bak. JoB. 1879. 304 = Cyclophorus.
cambricum L. sp. 2. 1086. 1753 = P. vulgare var.
Cameroonianum Hk. sp 5. 104. 1864 = Aspidium varians.
camporum Lindm. Ark. för. Bot. 1. 228 t. 10 f. 13. 1903 = Dryopteris.
(M) **camptocarpum** (Fée) C. Chr. Ind. 186. 1905; 515. 1906 — Brasilia. Craspedaria Fée, Cr. vasc. Br. 2. 65 t. 98 f. 1. 1872-73.
camptocaulon Bak. Ann. Bot. 5. 459. 1891 = Dryopteris.
camptonevron Fée, Gen. 237. 1850-52; 7 mém. 60 t. 23 f. 1 = P. tenuifolium.
(P) **camptophyllarium** Fée, 8 mém. 86. 1857 — Columbia.
(Pl) **campyloneuroides** Bak. Journ. Linn. Soc. 20. 229. 1883 — Borneo.
campylopodum Kl. Linn. 20. 395. 1847 = P. brasiliense.
canariense Willd.; Pr. Tent. 179. 1836 = P. vulgare *serratum.
cancellatum Fée, Gen. 242. 1850-52; 6 mém. 12 t. 7 f. 2. 1853 = P. Lindenianum.
canescens Bl. Enum. 133. 1828 = Dryopteris.
canescens Kze.; Hk. sp. 4. 262. 1862 = Dryopteris subincisa*.
(S) **cantoniense** Bak. JoB. 1879. 304. NPfl. 316 — Canton. Gymnogramme Bak. Ic. pl. t. 1685. 1887. — (an Cheiropleuria sp.?).
capense L.fil. Suppl. 445. 1781 = Hemitelia.
(P) **capillare** Desv. Berl. Mag. 5. 316. 1811. HB. 329. Chr. 81 — Amer. trop. P. pilipes Kl. 1847. Mett. n. 56 (non Hk. 1840) — P. blandum Fée 1857 — P. decipiens Hk. 1862; Ctenopteris J. Sm. 1875.
capitellatum Wall. List n. 306. 1828 = P. Wallichianum.
cardiophyllum Ettingsh. Farnkr. 41 t. 23 f. 23, 24. 1865 = P. vacciniifolium.
caribaeum Desv. Berl. Mag. 5. 319. 1811 = Dryopteris nemorosa.
caripense H.B.Willd. sp. 5. 202. 1810 = Dryopteris submarginalis.
(P) **carnosulum** Christ. Bull. Boiss. II. 4. 1102. 1904 — Costa Rica.
carnosum Mett. Pol. 124 n. 249. 1857 = Cyclophorus adnascens.
carnosum Kellogg. Proc. Calif. Acad. Sci. 2. 88 f. 24. 1861 = P. Scouleri.
carnosum Christ, Verh. Nat. Ges. Basel 11. 26. 1894 = Lecanopteris.
Carrii Bak. Fl. bras. 1^2. 597. 1870 = Dryopteris.

carthusianum Vill. Hist. Pl. Dauph. 1. 292. 1786 = ? Dryopteris rigida.
cartilagineum Pr.: Ettingsh. Denkschr. Ak. Wien **22**. 79. 1864; Farnkr. 38 t. 15 f. 8. 1865 = P. plebejum.
(G) **catharinae** Langsd. et Fisch. Ic. Fil. 9 t. 9. 1810. Mett. n. 108. HB. 343 — Brasilia. Mexico Costa Rica. — Marginaria Pr. 1836; Goniophlebium J. Sm. 1840 — Polypodium glaucum Raddi 1825 (non Thbg 1784); Goniophlebium J. Sm. 1857; Chrysopteris Raddiana Fée 1869 — Goniophlebium pictum Fée 1869.
(P) **caucanum** Hier. Engl. Jahrb. **34**. 503. 1903 — Columbia.
caudatum Raddi. Opusc. sci. Bol. **3**. 288. 1819; Klf. Enum. 113. 1824 = Dryopteris.
caudatum Mett. Pol. 126 n. 255. 1857 = Cyclophorus adnascens.
caudatum Reinw.; Salom. Nom. 292. 1883 = P. Feei var.
(G) **caudiceps** (Moore) Bak. Ann. Bot. **5**. 469. 1891 — Formosa. Goniophlebium Moore, Gard. Chr. n. s. **25**. 234. 1886.
caudiforme Bl. Enum. 122. 1828 = P. Feei var.
caudigerum Wall. List n. 298. 1828 = Dryopteris unita.
cayennense Mett. Pol. 93 n. 166. 1857 = P. ciliatum.
(P) **celebicum** Bl. Enum. 127. 1828; Fl. Jav. Fil. 179 t. 84 B. HB. 330 — Malesia. Ins. Philipp. — Cryptosorus J. Sm. 1875.
(P) **Cesatianum** Bak. JoB. **1879**. 42 — Borneo. P. papillosum Ces. Atti Ac. Napoli 7^{8}. 25 1876 (non Bl. 1828).
ceteraccinum Michx. Fl. bor. Am. **2**. 271. 1803 = P. polypodioides.
(G) **chacapoyense** Hk. sp. **5**. 29 t. 281. 1863. HB. 343 — Peru.
chaerophylloides Poir. Enc. **5**. 542. 1804 = Dryopteris pubescens var.
(G) **chartaceum** Bak. JoB. **1877**. 166 — Ecuador.
chattagrammicum Clarke, Tr. Linn. Soc. II. Bot. **1**. 548 t. 81. 1880 = Aspidium membranifolium.
cheilanthoides Bak. Malesia **3**. 45. 1886 = Dryopteris brunneo-villosa.
(P) **cheilostictum** Fée, 8 mém. 87. 1857 — Mexico.
(Pl) **chenopus** Christ, Bull. Soc. Fr. **52** Mém. I. 20. 1905 — Thibet orient.
(Pl) **chinense** Mett.; Kuhn, JoB. **1868**. 270 — China. Formosa.
(Pl) **chionolepis** Sod. Cr. vasc. quit. 375. 1893 — Ecuador.
(G) **chnoodes** Spr. Neu. Entdeck. **3**. 6. 1822. Mett. n. 121. HB. 344. NPfl. 314 — Ind. occ. Venezuela-Ecuador. — Marginaria Pr. 1836; Goniophlebium Fée 1850-52. — (an Polypodium dissimile L. 1759?).
(P) **chnoophorum** Kze. Flora **1839**[1]. Beibl. 34. Mett. n. 80. HB. 333 — Brasilia austr. Paraguay. — P. sublanosum Hk. 1862. HB. ed. I. 330.
(G) **chondrocheilon** (Fée) C. Chr. Ind. 325. 1905; 516. 1906 — Columbia. Goniophlebium Fée, 8 mém. 92. 1857.
(Gr) **Christii** Copeland, Polyp. Philipp. 118. 1905 — Mindanao. (an P. mediale var. parasiticum?).
(S) **christovalense** C. Chr. Ind. 328. 1905; 516. 1906 — Ins Salomonis. Gymnogramme palmata Bak. Syn. 389. 1868: Polypodium Sal. 1883 (non Bl. 1828).
chrysoconion Spr. Syst. **4**. 51. 1827. (Plum. t. 77) — Hispaniola.
(P) **chrysolepis** Hk. Ic. pl. t. 721. 1845. HB. 321 — Ecuador. Bolivia. Lepicystis Diels 1899. NPfl. 322.
chrysopodum Kl. Linn. **20**. 401. 1847 = P. repens.
cicutarium L. Syst. Nat. ed. X. **2**. 1326. 1759 = Aspidium.

cicutarium Poir. Enc. **5**. 534. 1804 = Dryopteris hexagonoptera.
(P) **ciliare** Fée, Cr. vasc. Br. **1**. 94 t. 27 f. 2. 1869 — Brasilia.
P. ovalescens Fée 1869.
(M **ciliatum** Willd. sp. 5. 144. 1810 — Amer. trop. — Goniophlebium J. Sm. 1840; Craspedaria Link 1841; Lopholepis J. Sm. 1842. Polypodium cajanense Desv. 1811. Mett. n. 166; P. piloselloides β HB. 340.
ciliatum Spr. Nova Acta **10**. 229. 1821 = Dryopteris megalodus.
ciliatum Pr. Del. Prag. **1**. 169. 1822 — Brasilia.
Lastrea Pr. 1849. (Dryopteris sp.).
ciliatum Pr. Rel. Haenk. **1**. 20. 1825 — Mexico.
(vix Willd.) Marginaria Haenkeana Pr. 1836.
ciliatum Boj. Hort. Maur. 416. 1837 = P. barbatulum.
cincinalis Güldenst. Reis. d. Russl. **2**. 343. 179 (nomen) — Rossia austr.
(P) **cinerascens** Lindm. Ark. för. Bot. **1**. 238 t. 11 f. 6. 1903 — Brasilia.
cinereum Cav. Descr. 248. 1802 = Alsophila quadripinnata.
(P) **circinatum** Sod. Cr. vasc. quit. 333. 1893 — Ecuador.
Clarkei Bak. Ann. Bot. **5**. 457. 1891 = Dryopteris mauiensis.
(Pl) **clathratum** Clarke, Tr. Linn. Soc. II. Bot. **1**. 559 t. 82 f. 1. 1880 — India bor. China — Pleopeltis Bedd. 1883.
(P) **clavifer** Hk. sp. **4**. 176. 1862. HB. 334 — Borneo. N. Guinea.
Grammitis Hk. 2 Cent. t. 5. 1860; Calymmodon Moore 1861.
coalescens Bak. JoB. **1877**. 164 = Dryopteris.
(C) **coarctatum** Kze. Linn. **9**. 39. 1834. Mett. n. 145. HB. 348. NPfl. 314 — Costa Rica-Peru-Paraguay. — Campylonevron Fée 1850-52.
coarctatum Kl. Linn. **20**. 382. 1847 = Dryopteris.
cochleatum Kl. Linn. **20**. 388. 1847 = Polystichum aculeatum var. 26.
colpodes Kze. Linn. **23**. 276. 316. 1850 = P. plesiosorum.
colpothrix Kze. Linn. **23**. 276. 316. 1850 = P. persicifolium.
colubrinum Vell. Fl. flum. **11** t. 71. 1827; Arch. Mus. Nac. Rio Jan. **5**. 448. 1881 = Polystichum abbreviatum.
(G) **columbianum** Bak. Ann. Bot. **5**. 470. 1891 — Columbia.
commune DC. Fl. franç **5**. 242. 1815 = P. vulgare.
(Pl) **commutatum** Bl. Enum. addend. 1828; Fl. Jav. Fil. 165 t. 73 — Malesia. Ins. Philipp. — P. adfine Bl. 1828 (affine Bl. 1829. Mett. n. 221. HB. 364 non Forst. 1786); Drynaria J. Sm. 1841; Pleopeltis Moore 1862; Colysis J. Sm. 1875.
(P) **comorense** Bak. JoB. **1877**. 72 — Ins. Johanna.
comosum L. sp. **2**. 1084. 1753 = P. phyllitidis.
comosum Wall. List n. 319. 1828 = Alsophila squamulata.
compositum Link, Hort. Berol. **2**. 106. 1833 = Dryopteris asplenioides.
comptoniaefolium Desv. Berl. Mag. **5**. 316. 1811 } = P. trifurcatum.
comptonioides Desv. Prod. 231. 1827. }
concinnum Willd. sp. **5**. 201. 1810 = Dryopteris.
concinnum Mett.; Kuhn, Linn. **36**. 132. 1869 = ? P. lanigerum.
confertum Roxb. Calc. Journ **4**. 493. 1844 — Chittagong.
confluens R.Br. Prod. Fl. N. Holl. 146. 1810 = Cyclophorus.
confluens Wall. List n. 325 1828 = Aspidium irregulare.
confluens Colla, Mem. Ac. Torino **39**. 50. 1836 — Chile.
confluens Roxb. Calc. Journ. **4**. 494. 1844 — Chittagong.
confluens Liebm. Vid. Selsk. Skr. V. **1**. 191 (seors. 39). 1849 = P. conterminans.
confluens Hk. sp. **5**. 46. 1863 = Cyclophorus rupestris.

confluens Fée, Cr. vasc. Br. **1.** 89 t. 28 f. 3. 1869 = P. apiculatum.
conforme Brack. Expl. Exp. **16.** 4 t. 1 f. 2. 1854 = P. Hookeri.
confusum J. Sm. Lond. JoB. **1.** 194. 1842 = P. jubaeforme var.
(Gr) **congenerum** (Bl.) Pr. Tent. 180. 1836 — Malesia.
Grammitis Bl. Enum. 115. 1828; Fl. Jav. Fil. 108 t. 46 f. 3 a. b.
coniifolium Wall. List n. 326. 1828 = Monachosorum subdigitatum.
conjugatum Poir. Enc. **5.** 516. 1804 = P. phymatodes et Drynaria querci-[folia.
conjugatum Klf. Wes. d. Farrnkr. 104. 1827 = Dipteris.
conjugatum Bak. Syn. 366. 1868 = P. coronans.
(P) **conjunctisorum** Bak. Ann. Bot. **8.** 129. 1894 — N. Guinea.
conjunctum Schur, Enum. Pl. Trans. 831. 1866 = ? Dryopteris Linnaeana.
connectile Michx. Fl. bor. Am. **2.** 271. 1803 = Dryopteris phegopteris.
(Gr) **Connellii** Bak.; Wright, Tr. Linn Soc. II. Bot. **6.** 82. 1901 — Guiana.
connexum Klf. Enum. 120. 1824 = Dryopteris.
consanguineum Kl. Linn. **20.** 387. 1847 = ? Dryopteris oligocarpa.
(P) **consimile** Eat. Mem. Amer. Acad. n. s. **8.** 198. 1860 (nomen); Mett. Ann. sc. nat. V. **2.** 253. 1864 — Mexico-Columbia.
contaminans Wall. List n. 320. 1828 = Alsophila glauca.
(G) **conterminans** Liebm. Vid. Selsk. Skr. V. **1.** 201 (seors. 49). 1849 — Mexico-[Guatemala.
P. confluens Liebm. 1849 (non R.Br. 1810 nec alior.).
contiguum Wall. List n. 285. 1828 = P. lineare.
contiguum J. Sm. JoB. **3.** 394. 1841 = Davallia.
contiguum Hk. Ic. pl. t. 987. 1854 = P. revolutum.
contiguum Brack. Expl. Exp. **16.** 6 t. 2 f. 1. 1854 = P. blechnoides.
contractum Desv. Prod. 242. 1827 — Peru. (non Polypodium sp.).
(L) **Copelandii** Christ in C. Chr. Ind. 518. 1906 — Ins. Philipp.
Drymoglossum confertum Copeland in Perkins, Fragm. 187. 1905; Loxogramme Copeland, Polyp. Philipp. 68. 1905 (non Polypodium Roxb. 1844).
corcovadense Raddi. Opusc. sci. Bol. **3.** 288. 1819 = Alsophila.
(M) *cordatum* Desv. Prod. 226. 1827 — Brasilia
cordatum Kze. Linn. **9.** 44. 1834 = P. Kunzeanum.
cordatum Moore, Ind. LXXI. 1857 = Dryopteris.
(P) **Cordemoyi** C. Chr. Ind. 518. 1906 — Réunion. — P. crassifrons Cordem. Fl. Réun. 86. 1895 (non Bak. 1867). (an P. leucosorum HB. 324?).
cordifolium L. sp. **2.** 1089. 1753 = Nephrolepis.
cordifolium Mart. et Gal. Mém. Ac. Brux. **15.** 31 t. 4 f. 2. 1842 = Aspidium trifoliatum juv.
coriaceum Sw. Prod. 133. 1788 = Polystichum adiantiforme.
coriaceum Raddi, Opusc. sci. Bol. **3.** 286. 1819; Pl. Bras. **1.** 16 t. 25. 1825 = P. crassifolium.
coriaceum Roxb. Calc. Journ. **4.** 481 t. 28. 1844 = Cyclophorus angustatus.
coriaceum Mett. Pol. 112 n. 217 t. 3 f. 12. 1857 = P. loxogramme.
(G) *coriaceum* (Fée) Kuhn, Fil. Afr. 146. 1868 — Réunion.
Goniophlebium Fée, 10 mém. 33 t. 37 f. 2. 1865.
coriandrifolium Sw. Fl. Ind. occ. **3.** 1675. 1806 = Aspidium cicutarium [vel A. hippocrepis.
(P) **cornigerum** Bak. Syn. 508. 1874 — Ceylon.
(Pl) **coronans** Wall. List n. 288. 1828 (nomen); Mett. Pol. 121 n. 242 t. 3 f. 40, 41. 1857. Chr. 117. NPfl. 319 — India bor. China merid. Formosa. Malacca.
Phymatodes Pr. 1836 (nomen); Drynaria J. Sm. 1841 (nomen); Polypodium conjugatum Bak. 1868. HB. 366 (non Poir. 1804); Drynaria Bedd. 1870.

coronopus Poir. Enc. 5. 527. 1804 = Aspidium cicutarium.

(P) **corticolum** C. Chr. Ind. 188. 1905; 519. 1906 — Ceylon.
P. glandulosum Hk. sp. 4. 193 t. 276 A. 1862. HB. 327. NPfl. 309 (non Desv. 1811); Ctenopteris J. Sm. 1875.

costale Jenm. Bull. Dep. Jam. 4. 140. 1897 (err. script.) = P. costatum.

costaricanum Hier. Engl. Jahrb. 34. 530. 1904 = P. Wendlandii.

(G) **costaricense** Christ, Bull. Boiss. 4. 660. 1896; Farnkr. 96 — Costa Rica.

costatum Wall. List n. 265. 1828 (nomen) = Cyclophorus Beddomeanus et C. flocculosus.

(C) **costatum** Kze. Linn. 9. 38. 1834. Mett. n. 143 — Amer. centr. Jamaica.
Campyloneurum Pr. 1836; Cyrtophlebium J. Sm. 1842. Campyloneuron immersum J. Sm. 1854.

costatum Mett. Pol. 131 n. 267 t. 3 f. 14. 1857 = Cyclophorus Beddomeanus.

costatum Hk. sp. 5. 7. 1863 = Dryopteris pennigera*.

(Pl) **costulatum** (Ces.) Bak. JoB. **1880**. 215 — Sumatra. — Acrostichum Ces. Rend. Ac. Napoli **16**. 27, 30. 1877. — (an P. triquetrum f. monstr.?).

(Pl) **crassifolium** L. sp. 2. 1083. 1753. Mett. n. 210. HB. 360. Chr. 105. NPfl. 315.
Anaxetum Schott 1834; Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Dipteris J. Sm. 1842; Pleuridium Fée 1850-52; Pleopeltis Moore 1857 — Polypodium acrosorum Kze. 1834 — P. anocarpos Kze. 1834; Phymatodes Pr. 1836 — Pleuridium angustum Fée 1857 — Polypodium porrectum Willd. 1810; Phymatodes Pr. 1836 — Polypodium Vittaria Mett. 1856.

(P) **crassifrons** Bak. Syn. 325. 1867. NPfl. 309 — N. Caledonia.

crassifrons Cordem. Fl. Réun. 86. 1895 = P. Cordemoyi.

crassimargo Kze.; Mett. Pol. 71 n. 103 b. 1857 = P. lepidopteris.

crassinervatum Kiærsk. Cat. Fil. herb. ht. Haun. 49. 1874 = P. lanceolatum*.

crassinerve Schum. Vid. Selsk. Afh. 4. 227 (seors. 453). 1827 = P. punctatum var.

crassinerve Bl. Enum. 125. 1828 = P. platyphyllum.

crassum Kirk, Tr. N. Zeal. Inst. 17. 232. 1885 = P. Billardieri.

(P) **craterisorum** Harr. Journ. Linn. Soc. **16**. 31. 1877 — Ins. Philipp.

crenato-dentatum Kl. Linn. **20**. 384. 1847 — Peru.
(Polystichum vel Dryopteris sp.).

(Pl) **crenato-pinnatum** Clarke, Journ. Linn. Soc. **25**. 99 t. 42. 1888 — Munipore. Yunnan. — Pleopeltis Bedd. 1892. (conf. P. pseudo-serratum).

crenatum Forsk. Fl. Aegypt. arab. 185. 1775 = Dryopteris.

crenatum Sw. Prod. 132. 1788 = Dryopteris Poiteana.

crenatum Schrank, (Naturhist. Briefe 2. 295. 1785); Baier. Flora 2. 426. 1789 = Athyrium filix femina.

crenatum Hk. Fil. Exot. t. 84. 1858 = Dryopteris Ghiesbreghtii.

crenulatum Gmel. Syst. Nat. 2^2. 1313. 1791 = Dryopteris crenata.

crenulatum Kze.; Mett. Pol. 110 n. 210 b. 1857 = P. palmatum.

crinale Hk. et Arn. Bot. Beech. Voy. 105. 1832 = Dryopteris.

crinitum Poir. Enc. 5. 528. 1804 = Dryopteris.

crispatum L. sp. 2. 1080. 1753. (Plum. t. 102 B.) — Martinique.

crispatum Hk. sp. 5. 1. 1863 = P. goniopteroides.

(P) **crispulum** Christ, Bull. Boiss. II. 4. 1102. 1904 — Costa Rica.

crispum Gouan, Illustr. obs. bot. 81. 1773 = Cystopteris fragilis *regia.

crispum Roth, Usteri neu. Ann. 2. 10 Stück. 56. 1794 = Cryptogramma.

cristatum L. sp. 2. 1090. 1753 = Dryopteris.

cristatum Houtt. Pfl. Syst. 13[1]. 208 t. 99 f. 3. 1786 = Microlepia strigosa.
Crossii Bak. Ann. Bot. 5. 455. 1891 = Dryopteris.
cruciatum Klf. Enum. 117. 1824 = Dryopteris hexagonoptera.
cruciatum Bak. Fl. Maur. 503. 1877 = Dryopteris.
cryptocarpon Fée, 8 mém. 88. 1857 = P. furfuraceum.
(Gr) **cryptophlebium** Bak. JoB. **1880**. 370 — Madagascar.
(P) **cryptosorum** C. Chr. Ind. 519. 1906 — Malesia.
P. decipiens Mett.; Kuhn, Linn. **36**. 129. 1869. HB. 508 (non Hk. 1862).
(?) **cryptum** Und. et Maxon, Bull. Torr. Cl. **29**. 579 c. fig. 1092 — Cuba.
crystallinum Kze. Farnkr. **2**. 85 t. 135. 1851 = Dryopteris.
ctenoides Jenm. Bull. Dep. Jam. **4**. 129. 1897 = Dryopteris.
cubanum Bak. Syn. 304. 1867 = Dryopteris cordata.
Cubense Fée, Gen. 241. 1850-52; 7 mém. 61 t. 26 f. 1 = P. subpetiolatum.
Cubense Christ, Engl. Jahrb. **24**. 131. 1897 = P. vexatum.
(P) **cucullatum** Nees et Bl. Nova Acta **11**. 121 t. 12 f. 3. 1823. Mett. n. 7. HB. 324. Chr. 80. NPfl. 309 — Ceylon. Malesia-Fiji-N. Caledonia.
Xiphopteris Spr. 1827; Grammitis Bl. 1828; Calymmodon Pr. 1836; Plectopteris gracilis Fée 1850-52; P. Calymmodon Fée 1857.
(P) **cuencanum** Hier. Engl. Jahrb. **34**. 505. 1904 — Ecuador.
(P) **cultratum** Willd. sp. **5**. 187. 1810. Mett. n. 46. HB. 327. Chr. 81. NPfl. 310 — Amer. trop. Madagascar. Ins. Mascar. — P. asplenifolium β L. sp. **2**. 1084. 1753. — P. reclinatum Brack. 1854. HB. 335. — P. elasticum Bory; Willd 1810. Mett. n. 44; (Réunion); P. leveilleanum Desv. 1827. P. flexile Fée 1853.
Cumingianum Hk. sp. **5**. 103. 1864 = Aspidium macrodon.
cuneatum Kuhn, Bot. Zeit. **1868**. 41 (syn.) = Cyclophorus.
Cunninghami Hk. Gard. Ferns in obs. ad t. 30. 1862 = P. dictyopteris.
(Pl) **curtidens** Christ, Ann. Jard. Buit. II. **5**. 122. 1905 — Borneo.
(P) **Curtisii** Bak. JoB. **1881**. 367 — Sumatra.
curvans Mett. Ann. sc. nat. V. **2**. 253. 1864 = ? P. curvatum.
(P) **curvatum** Sw. Schrad. Journ. **1800**[2]. 24. 1801. HB. 329 Chr. 86. NPfl. 310 — Ind. occ. Columbia-Peru.
— P. inaequale Fée 1866. HB. 325. NPfl. 309 (non Link 1833). ? P. curvans Mett. 1864 = P. curvatum Mett. Pol. 59. n. 75; Hk. sp. **4**. 201.
curvivenium Wall. List n. 292 1828 = Drynaria.
cuspidatum Don, Prod. Fl. Nepal. 2. 1825 = P. leiorhizum.
cuspidatum Pr. Rel. Haenk. **1**. 20 t. 1 f. 3. 1825 = P. percussum.
cuspidatum Bl. Enum. 132. 1828 = P. persicifolium.
cuspidatum Roxb. Calc. Journ. **4**. 491. 1844 = Dryopteris urophylla.
cuspidatum Moritz. Syst. Verz. — 1845-46 — Drynaria rigidula.
cuspidatum Mett. Pol. 81. n. 134. 1857 = P. verrucosum.
cyatheaefolium Desv. Prod. 239. 1827 = Dryopteris.
cyathoides Sw. Syn. 37. 1806 = Cyathea percussa.
(Pl) **cyclobasis** Bak. Kew Bull. **1896**. 42 — N. Guinea. — (an P. Annabellae?).
cyclocolpon Christ, Bull. Boiss. **4**. 659. 1896 = Dryopteris.
(Pl) **cyclophyllum** Bak. Ann. Bot. **5**. 473. 1891. NPfl. 316 — China.
cymatodes Kze. Linn. **23**. 277, 317. 1850 = P. fraxinifolium.
cynapifolium Hoffm. Deutschl. Fl. **2**. 9. 1795 = Cystopteris fragilis.
(Pl) **cyrtolobum** Clarke, Tr. Linn. Soc. II. Bot. **1**. 563 t. 83. 1880 — India bor.
(Pl) **dactylinum** Christ, Bull. Soc. Fr. **52** Mém. I. 20. 1905 — China. [Yunnan.

(P) **daguense** Hier. Engl. Jahrb. **34**. 504. 1904 — Columbia.
(P) **dareiforme** Hk. 2 Cent. t. 24. 1860. HB. 339. Chr. 88 — India bor.
Leucostegia Bedd. 1876; Davallia Clarke 1880 (pt.). — (conf. Davallia
(G) **dasypleuron** Kze. Linn **9**. 43. 1834. Mett. n. 119 — Peru. Columbia. [Clarkei).
Marginaria Pr. 1836; Goniophlebium Moore 1857. (an P. lanuginosum
daucifolium Poir. Enc. **5**. 542. 1804 = Dryopteris effusa. [Schrad. 1824?).
(P) **davalliaceum** F. Muell. et Bak. JoB. **1890**. 108. NPfl. 310 — N. Guinea.
(an Davallia (Prosaptia) sp.?)
davallioides Mett. Fil. Lips. 30. 1856 = Monachosorum subdigitatum.
Davidi Franch. Nouv. Arch. Mus. II. **10**. 119. 1887 = Athyrium.
Davidii Bak. Ann. Bot. **5**. 472. 1891 = Cyclophorus pekinensis.
dealbatum Forst. Prod. 83. 1786 = Cyathea.
debile Bak. Syn. 313. 1867 = Dryopteris.
decipiens Hk. sp. **4**. 231 t. 279 B. 1862 = P. capillare.
decipiens Mett.; Kuhn, Linn. **36**. 129. 1869 = P. cryptosorum.
(P) **decorum** Brack. Expl. Exp. **16**. 7 t. 2 f. 2. 1854 HB. 331. Chr. 85. NPfl. 310
— Ceylon-Malesia-Polynesia. — P. nutans J. Sm. 1841; Mett. n. 28 (non
(P) **decrescens** Christ. Ann. Jard. Buit. II. **4**. 35. 1904 — Celebes. [Bl. 1828).
(Ph) **decumanum** Willd. sp. **5**. 170. 1810. Mett. Pol. n. 180. HB. 347. Chr. 95.
NPfl. 314 — Amer. trop. — Pleopeltis Pr. 1836; Phlebodium J. Sm. 1841;
Chrysopteris Fée 1850-52; Campyloneuron Fée 1869 — Chrysopteris
dictyocallis Fée 1850-52; Polypodium Mett. n. 186 b. 1857; Phlebodium
J. Sm. 1857; P. multiseriale Moore 1855; Chrysopteris Fée 1866.
(C) **decurrens** Raddi. Opusc. sci. Bol. **3**. 287. 1819; Pl. Bras. **1**. 23 t. 33. 1825.
Mett. n. 146. HB. 349. Chr. 96. NPfl. 315 — Ind. occ.-Brasilia. Peru.
Campyloneurum Pr. 1836: Cyrtophlebium J. Sm. 1841 — Campylo-
nevron juglandifolium Fée 1869. — (an Polypodium simile L. 1759?).
decurrens Kze. Bot. Zeit. **1846**. 421 = P. revolutum.
decursive-pinnatum van Hall, Nieuwe Verhdl. Nederl. Inst. **5**. 204 c. tab. 1836
decussatum L. sp. **2**. 1093. 1753 = Dryopteris. [= Dryopteris.
deflexum Klf. Enum. 114. 1824 = Dryopteris.
deflexum Lowe, Ferns **1** t. 45. 1856 = P. fraxinifolium.
deflexum Bak. Syn. 303. 1867 = Dryopteris Lindigii.
(P) **delicatulum** Mart. et Gal. Mém. Ac. Brux. **15**. 35 t. 7 f. 1. 1842. Mett. n. 35.
HB. 326. NPfl. 309 — Mexico-Ecuador. — Ctenopteris J. Sm. 1854.
(P) **delitescens** Maxon, Bull. Torr. Cl. **32**. 74. 1905 — Jamaica.
Grammitis muosuroides Schkuhr, Kr. Gew. **1**. 9 t. 7. 1804 (non Sw. 1801).
(P) **deltodon** Bak. Journ. Linn. Soc. **15**. 419. 1876 — Madagascar.
(P) **deltoideophyllum** Bak. JoB. **1876**. 345 — Samoa.
deltoideum Sw. Prod. 133. 1788 = Dryopteris.
deltoideum Liebm. Vid. Selsk. Skr. V. **1**. 190 (seors. 38). 1849 = P. catha-
deltoideum Bak. JoB. **1888**. 230 = P. hemitomum. [rinae.
demeraranum Bak. Tr. Linn. Soc. II. Bot. **2**. 290. 1887 = Dryopteris.
(G) **demissum** (Fée) C. Chr. Ind. 326. 1905; 521. 1906 — Brasilia.
Goniophlebium Fée, Cr. vasc. Br. **2**. 63. 1872-73.
(P) **dendricolum** Jenm. Gard. Chr. III. **16**. 467. 1894 — Grenada. Jamaica.
dentatum Forsk. Fl. Aegypt. arab. 185. 1775 = Dryopteris parasitica.
dentatum Schrank (Naturh. Briefe **2**. 295. 1785); Baier. Flora **2**. 426 1789
= Athyrium filix femina.
dentatum Dick. Fl. Crypt. fasc. 3 t. 7 f. 1. 1793 = Cystopteris fragilis.

dentatum Bak. Ann. Bot. **5**. 456. 1891 = Dryopteris longipetiolata.
denticulatum Sw. Prod. 134. 1788 = Polystichum.
(P) **denticulatum** (Bl.) Pr. Tent. 178. 1836 — Java.
Grammitis Bl. Fl. Jav. Fil. 121 t. 50 f. 4. 1828; Calymmodon Moore 1861.
dentigerum Wall. List n. 334. 1828 = Athyrium filix femina.
deparioides Bak. JoB. **1879**. 297 = Aspidium depariopsis.
deparioides Christ, Farnkr. 116. 1897 = Lecanopteris.
(P) **dependens** Bak. Syn. 335. 1867 — Mexico. Ecuador. Peru.
(Gr) **Deplanchei** Bak. Syn. 322. 1867 — N. Caledonia.
Austrogramme Fourn. 1873; Grammitis athroosperma Fourn. 1869.
(P) **depressum** C. Chr. Ind. 522. 1906 — Brasilia.
P. immersum Fée, Cr. vasc. Br. **1**. 88 t. 27 f. 1. 1869 (non Vahl 1794).
detergibile Hk. sp. **5**. 49. 1863 = Cyclophorus flocculosus.
(P) **devolutum** Bak. Journ. Linn. Soc. **15**. 419. 1876 — Madagascar.
Dianae Hk. sp. **4**. 234. 1862 = Dryopteris.
diaphanum Bory, Voy. **1**. 328. 1804 = Cystopteris fragilis.
dichotomum Houtt. Nat. Hist. **14**. 181. 1783; Pfl. Syst. **13**^{1}. 204 t. 99 f. 2. 1786 = Dryopteris unita.
dichotomum Thbg. Fl. Jap. 338 t. 37. 1784 = Gleichenia linearis.
dicksonifolium Roxb. in Beats. St. Helena 318. 1816 = Dryopteris punctata.
(Gr) **dicranophyllum** C. Chr. Ind. 331. 1905; 522. 1906 — Grenada. Guiana. Brasilia. — Grammitis furcata Hk. et Grev. Ic. Fil. t. 62. 1828; Polypodium Mett. 1857. n. 9. HB. 322. NPfl. 308 (non Desv. 1827).
dictyocallis Mett. Pol. 100. n 186 b. 1857 = P. decumanum var.
dictyophyllum Kze.; Mett. Pol 96. n. 173. 1857 = P. lycopodioides *surinamense.
(Pl) **dictyopteris** Mett. Ann. sc. nat. IV. **15**. 72. 1861 — N. Zealand. N. Hebridae.
P. attenuatum A. Rich. Fl. N. Zél. 62. 1832 (non Willd. 1810 nec R.Br. 1810); Dictyopteris Hk. 1841; D. lanceolata J. Sm. 1841; Dictymia Hk.fil. 1854 (non Polypodium L. 1753); Polypodium Cunninghami Hk. 1862. HB. 354. NPfl. 315.
(G) **Dielseanum** C. Chr. Ind. 522. 1906 — China.
P. leuconeuron Diels, Engl. Jahrb. **29**. 203. 1900 (non Christ 1900).
difforme Bl. Enum. 135. 1828 = Aspidium irregulare.
(C) *difforme* (Loud.) Kze. Linn. **23**. 277. 1850 — Hort.
Cyrtophlebium Loud. Cat. 1849; Campyloneurum Moore 1861.
diffundens Kze. Bot. Zeit. **1846**. 422 = P. insigne.
(S) **digitatum** (Bak.) C. Chr. Ind. 522. 1906 — Cochinchina. Tonkin. Yunnan.
Gymnogramme Bak. JoB. **1890**. 267; Grammitis Finlaysoniana Wall. 1829 (nomen); Selliguea Moore 1857 (nomen); Gymnogramme Bak. 1891. Colysis tridactylis Fée 1850-52 (nomen).
dilatatum Hoffm. Deutschl. Fl. **2**. 7. 1795 = Dryopteris spinulosa*.
dilatatum Wall. List n. 295. 1828 = P. euryphyllum.
dilatatum Liebm. Vid. Selsk. Skr. V. **1**. 208 (seors. 56). 1849 = Dryopteris effusa.
dilatatum var. grandidentatum Ces. Atti Acad. Napoli **7**^{8}. 27. 1876 = P. [grandidentatum.
dimidiatum Pr. Tent. 179. 1836 (nomen) —?
(Gr) **diminutum** Bak. Syn. 507. 1874 — Ins. Lord Howe.
dimorphum Link, Hort. Berol. **2**. 88. 1833 = P. angustifolium var.
dimorphum Zoll. Syst. Verz. 37, 48. 1854 = Drynaria propinqua.
dimorphum Bak. Ann. Bot. **5**. 477. 1891 = P. hainanense.

(Gr) **diplosorum** Christ, Verh. Nat. Ges. Basel **11**. 440. 1896; Ann. Jard. Buit. **15**. 146. 1897 — Java. Borneo. Celebes. Ins. Philipp.
Grammitis hirta Bl. Enum. 115. 1828; Fl. Jav. Fil. t. 47 f. 1 (non HBK. 1815); Polypodium Pr. 1836. HB. 320 pt. (non Sw. 1788 nec Pr. 1825) — Grammitis setosa Bl. 1828; Fl. Jav. Fil. t. 48 f. 3; Polypodium Pr. 1836 (non Thbg. 1784 nec Mett. 1857).
Dipteris Bl. Enum. 135. 1828 = Dipteris conjugata.
(P) **discolor** Hk. Ic. pl. t. 386. 1841. HB. 325. NPfl. 309 — Guiana.
Ctenopteris J. Sm. 1875.
disjunctum Rupr. Distr. Cr. vasc. Ross. 52. 1845 = Dryopteris Linnaeana.
dissectum Forst. Prod. 81. 1786 = Dryopteris.
dissectum Sw. Prod. 134. 1788 = Dennstaedtia.
(G) *dissimile* L. Syst. Nat. ed. X. **2**. 1325. 1759 (Pluk. t. 288 f. 1) — Ind. occ.
? Goniophlebium J. Sm. 1841. (an eadem ac P. chnoodes Spr.?).
dissimile Houtt. Pfl. Syst. **13**[1]. 190 t. 99 f. 1. 1786 = Blechnum capense.
dissimile Schkuhr, Kr. Gew. **1**. 14 t. 14. 1806 = P. attenuatum.
dissimile Hk. sp. **5**. 25. 1863 = P. chnoodes.
dissimile Gris. Fl. br. W. Ind. 679. 1864 = P. sororium.
dissitifolium Bak. Kew Bull. **1895**. 54 = Dryopteris.
distans Klf. Enum. 113. 1824 = Dryopteris submarginalis.
distans Don, Prod. Fl. Nepal. 2. 1825 = Dryopteris brunnea.
distans Raddi, Pl. Bras. **1**. 21 t. 31. 1825 = P. fraxinifolium.
distichocarpum Mett. Ann. Lugd. Bat. **2**. 231. 1866 = Cyclophorus.
(G) **divaricatum** Fourn. Mex. pl. **1**. 80. 1872 — Mexico.
divergens Willd.; Schkuhr. Kr. Gew. **1**. 27 t. 26 b. 1806 = Dryopteris effusa.
(Pl) **diversifolium** Willd. sp. **5**. 166. 1810 — Austr. Tasm. Ins. Norfolk. N. Caledonia. N. Zealand. — P. scandens Lab. N. Holl. pl. spec. **2**. 91 t. 240. 1806 (non Forst. 1786); P. Billardieri R.Br. 1810 (non supra). Mett. n. 189. HB. 364. Chr. 111. NPfl. 318; Phymatodes Pr. 1836; Chrysopteris Link 1841; Drynaria J. Sm. 1846; Pleopeltis Moore 1857. (vide tamen Carr. in Seem. Fl. Vit 368. 1873).
diversifolium R.Br. Prod. Fl. N. Holl. 147. 1810 = Drynaria rigidula.
diversifolium Sw. Vet. Ak. Handl. **1817**. 60 = Dryopteris vivipara.
diversifrons Kl.; Kze. Linn. **23**. 278. 1850 = Dryopteris parasitica.
(Pl) **dolichopodum** Diels, Engl. Jahrb. **29**. 205. 1900 — China.
(P) **dolorense** Hier. Engl. Jahrb. **34**. 512. 1904 — Columbia.
domingense Spr. Syst. **4**. 51. 1827 = Dryopteris scolopendrioides.
Donianum Spr. Syst. **4**. 54. 1827 — Nepal.
P. tenellum Forst. Prod. Fl. Nepal. 2. 1825 (non Forst. 1786).
(Gr) **dorsipilum** Christ in Warb. Monsunia **1**. 59. 1900 — China.
draconopterum Hk. sp. **5**. 86. 1863 = Aspidium.
Drakeanum Franch. Nouv. Arch. Mus. II. **7**. 165. 1883 = Cyclophorus.
drepanopteron Kze. Linn. **23**. 278, 318. 1850 = Athyrium.
Drepanum Lowe, Tr. Cambr. Phil. Soc. **6**. 523. 1838 = Polystichum.
(Pl) **drymoglossoides** Bak. JoB. **1887**. 170 — China.
drynarioides Gris. Abh. Ges. Wiss. Gött. **7**. 284. 1857 = P. aureum *areolatum.
drynarioides Christ, Farnkr. 117. 1897 = Dryostachyum.
Dryopteris L. sp. **2**. 1093. 1753 = Dryopteris Linnaeana.
dubium Roxb. Calc. Journ. **4**. 496. 1844 = Microlepia strigosa.

dubium Hk. sp. 5. 15. 1863 α = Polystichum dubium.
β = Polystichum dictyophyllum.

dubium Kuhn, Fil. Afr. 147. 1868 = ? Cyclophorus spissus.

dulce Poir. Enc. 5. 523. 1804 = ? P. aureum.

(Pl) **dulitense** Bak. Kew Bull. **1893**. 211 — Borneo.

(P) **duriusculum** Christ, Verh. Nat. Ges. Basel **11**. 213 t. 2 f. 5-7. 1895; Ann. Jard. Buit. **15**. 151 t. 16 f. 23. 1897 — Celebes.

(G) **Eatoni** Bak. Syn. 511. 1874 — Mexico — P. Ghiesbreghtii Eat. Proc. Amer. Ac. **8**. 618. 1873 (non Linden; Bak. 1867).

(Pl) **ebenipes** Hk. sp. 5. 88. 1863. HB. 365 — Ind. bor. Yunnan. Borneo. Pleopeltis Bedd. 1866; Phymatopsis J. Sm. 1875. Polypodium melanopus R.Br.; Wall. 1829 (nomen).

echinatum Gmel. Syst. Nat. **2**[2]. 1309. 1791 = Polystichum.

(P) **echinolepis** Fée, 8 mém. 87. 1857 — Mexico.

Eckloni Kze. Linn. **10**. 498. 1836 = P. polypodioides.

(P) **ecostatum** Sod. Cr. vasc. quit. 308. 1893 — Ecuador. Costa Rica.

(P) **ecuadorense** C. Chr. Ind. 524. 1906 — Ecuador. — P. subscabrum Hk. sp. **4**. 183 t. 274 A. 1862 (non Kl. 1847); P. pichinchense Hier. Engl. Jahrb. **34**. 506. 1904 (non P. pichinchae Sod. 1893).

effusum Sw. Prod. 134. 1788 = Dryopteris.

efulense Bak. Kew Bull. **1897**. 299 = Dryopteris.

(P) **Eggersii** Bak. in Hk. Ic. pl. t. 1671. 1886 — Dominica. Jamaica.

(P) **Ehrenbergianum** Kl. Linn. **20**. 381. 1847. Mett. n. 82 — Mexico. P. olivaceum Liebm. 1849.

(Pl) **eilophyllum** Diels, Engl. Jahrb. **29**. 204. 1900 — China. — P. involutum Bak. JoB. **1889**. 177. NPfl. 315 (non Desv. 1811 nec Mett. 1856).

elasticum Rich. Act. Soc. hist. nat. Paris **1**. 114. 1792 (nomen) = P. plumula.

elasticum Bory; Willd. sp. 5. 183. 1810 = P. cultratum var.

elasticum γ Glaziovii Bak. Fl. bras. **1**[2]. 517 t. 64 f. 2. 1870 = P. pectinatum var.

elatius Schrad. Gött. Gel. Anz. **1824**. 868 = P. brasiliense.

elatum Roxb. Calc. Journ. **4**. 495. 1844 — Amboyna.

eleagnifolium Bory, Dup. Voy. Bot. **1**. 259 t. 31 f. 1. 1828 = Cyclophorus serpens.

elegans Cav. Hort. Matr. **1**. —. t. —; Sw. Syn. 35. 1806 = P. loriceum.

elegans Poir. Enc. **5**. 519. 1804 = P. plumula.

(G) **eleutherophlebium** (Fée) Mett. Pol. 75. n. 115. 1857. HB. 343 — Venezuela. Columbia. — Goniophlebium Fée, Gen. 255. 1850-52.

(P) **ellipsoideum** Fée, 7 mém. 57 t. 21 f. 1. 1857. (HB. ed. I. 333) — Mexico.

ellipticosorum Fée, Gen. 239. 1850-52 = P. xanthotrichium.

(S) **ellipticum** Thbg. Fl. Jap. 335. 1784. Chr. 107. NPfl. 318 — Japonia. Formosa. China. Ind. bor. Ins. Philipp. Queensland.
Gymnogramme Bak. 1868. HB. 388; Selliguea Bedd. 1870. Hemionitis pothifolia Don 1825; Gymnogramme Spr. 1827; Selliguea J. Sm. 1841; Colysis Pr. 1849; Polypodium Mett. 1856. Pol. n. 193. Grammitis decurrens Wall.; Hk. et Grev. 1827; Selliguea Pr. 1836; Gymnogramme Hk. 1864 — G. pentaphylla Bak. 1898 — Grammitis ampla F. Muell. 1866; Benth. 1878; Gymnogramme Bak. 1891 (Queensland).

(S) **Elmeri** Copeland in Perkins, Fragm. 191. 1905 — Luzon.

elongatum Ait. Hort. Kew **3**. 465. 1789 = Dryopteris filix mas*.

elongatum Schrad. Gött. gel. Anz. **1818**. 915 = P. Schraderi.

elongatum Desv. Prod. 228. 1827 — Brasilia.

elongatum Wall. List n. 309. 1828; Hk. sp. **4**. 234. 1862 = Dryopteris khasiana.
elongatum Mett. Pol. 88 n. 154 t. 2 f. 8, 9. 1857 = P. lanceolatum*.
elongatum Goldm. Nova Acta **19**. Suppl. I. 454. 1843 = P. lineare.
(P) **Endresi** Bak. Ann. Bot. **5**. 468. 1891 — Costa Rica.
enerve Cav. Descr. 245. 1802 — Ins. Mariannae. — P. evenium Spr. 1827.
(Pl) **Engleri** Luerss. Engl. Jahrb. **4**. 361. 1883 — Japonia.
(Pl) **ensatum** Thbg. Tr. Linn. Soc. **2**. 341. 1794. Mett. n. 225. HB. 361. NPfl. 315 — Japonia. China. — Pleopeltis Moore 1862; Polypodium Phyllitidis Thbg. Fl. Jap. 335. 1784 (non L. 1753).
ensifolium Wild. sp. **5**. 152. 1810 = P. angustifolium var.
(G) **ensiforme** Thbg. Prod. Fl. Cap. 172. 1800. Mett. n. 106. HB. 341. Chr. 92. NPfl. 312 — Africa austr. — Marginaria Pr. 1836; Phlebodium J. Sm. 1841; Goniophlebium Fée 1850-52 (vix Polypodium triphyllum Jacq. 1789).
ensiforme Vell. Fl. flum. **11** t. 61. 1827; Arch. Mus. Nac. Rio Jan. **5**. 445 1881 = P. decurrens.
eriophorum Hk. Ic. pl. t. 991. 1854 = Notholaena.
erubescens Wall. List n. 330. 1828 = Dryopteris.
(G) **erythrocarpum** Mett.; Kuhn, Linn. **36**. 135. 1869. HB. 511. NPfl. 312 — Sikkim. China. — Goniophlebium Bedd. 1876.
euchlorum Sod. Rec. 58. 1883 = Dryopteris.
euchlorum Kl.; Hier. Engl. Jahrb. **34**. 509. 1904 = P. subsessile.
(Pl) **euryphyllum** C. Chr. Ind. 525. 1906 — China merid. Ind. bor. Ceylon. Malacca. Samoa. — P. dilatatum Wall. 1828 (nomen, non Hoffm. 1795 nec Liebm. 1849); Hk. sp. **5**. 85. 1863. HB. 365. Chr. 112. NPfl. 318; Pleopeltis Bedd. 1866; Colysis J. Sm. 1875 — Drynaria acuminata Brack. 1854 (p. 47 non p. 41); D. sylvatica Brack. 1854; Polypodium Mett. 1870 (non Schkuhr 1806); P. expansum Bak. 1876 (non Poir. 1804) [(Samoa).
evectum Forst. Prod. 81. 1876 = Angiopteris.
evenium Spr. Syst. **4**. 46. 1827 = P. enerve.
exaltatum L. Syst. Nat. ed. X. **2**. 1326. 1759 = Nephrolepis.
(Pl) **excavatum** Bory; Willd. sp. **5**. 158. 1810 — Africa trop. Madagascar. Ins. Mascar. Himalaya. Japonia. — Phymatodes Pr. 1836; Drynaria Fée 1850-52; Pleopeltis Moore 1862. Polypodium simplex Sw. Schrad. Journ. **1800**². 19. 1801 (non Burm. 1768): Phymatodes Pr. 1836; Pleopeltis Bedd. 1883; Polypodium lineare β HB. 354 — P. scolopendrium Don 1825. P. sesquipedale Wall. 1828. Mett. n. 162; Drynaria J. Sm. 1846.; Niphobolus Keys. 1873 — Polypodium loriforme Wall. 1828. Mett. n. 164; Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Pleopeltis Moore 1862 — Drynaria phlebodes Fée 1850-52; Polypodium Kze.; Mett. 1857. n. 163 — P. asterolepis Bak. 1888 (non P. astrolepis Liebm. 1849); P. aspidiolepis Bak. 1891 et auctt. (non Bak. 1887).
excavatum Roxb. Calc. Journ. **4**. 485. 1844 = P. longissimum.
excelsum Desv. Prod. 243. 1827 = Dryopteris.
excelsum Bak. Syn. 505. 1874 = Dryopteris tannensis.
exiguum Hew. Mag. Nat. Hist. II. **2**. 458. 1838 = P. Swartzii.
exiguum Gris. Fl. br. W. Ind. 701. 1864 = P. Grisebachii.
exiguum Fée, Cr. vasc. Br. **1**. 89 t. 37 f. 1. 1869 = P. Blanchetii.
eximium Kze. Bot. Zeit. **1846**. 424 = Aspidium irregulare.
exosum Gmel. Syst. Nat. **2**². 1311. 1791 = Dryopteris invisa.
expansum Poir. Enc. **5**. 523. 1804 = ? Aspidium martinicense.

expansum Bak. JoB. **1876**. 12 = P. euryphyllum var.
exsudans Liebm. Vid. Selsk. Skr. V. **1**. 206 (seors. 54). 1849 (nomen) = P. litigiosum
(P) **exsudans** Christ, Bull. Boiss. II. **4**. 1103. 1904 — Costa Rica. [Liebm. l. c.
(Pl) **exsul** Mett.; Kuhn, Linn. **36**. 138. 1869. HB. 513 — Brasilia. — Drynaria
extensum Forst. 83. 1786 = Alsophila. [Fée 1872-73.
extensum Pr. Rel. Haenk. **1**. 26. 1825 = Dryopteris subincisa.
extensum Fée, Cr. vasc. Br. **1**. 85 t. 28 f. 3. 1869 = P. recurvatum.
externatum Bory, Bél. Voy. Bot. **2**. 28. 1833 = P. nigrescens.
(Pl) **Faberi** Christ, Bull. Soc. Fr. **52**. Mém. I. 17. 1905 — China.
(G) **falcaria** Kze. Linn. **18**. 316. 1844 Mett. n. 117 — Mexico. Ecuador. Goniophlebium Fée 1850-52. G. Calaguala Fée 1857 — G. invertens Fée 1857 — ? G. arcuatum Fée 1850-52 — Polypodium punctulatum Hk. 1845. (HB. 343) (non Poir. 1804); Goniophlebium Moore 1862.
falcatum L. fil. Suppl. 446. 1781 = Polystichum.
falcatum Kellog. Proc. Calif. Ac. **1**. 20. 1854 = P. glycyrrhiza.
falciculatum Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 24 t. 36 bis 1825
falciforme Sieb. Fl. Mart. exsicc. n. 61 = P. chnoodes. [= Dryopteris.
(G) **falcoideum** Kuhn; Hier. Engl. Jahrb. **34**. 533. 1904 — Costa Rica.
(M) **fallax** Schlecht. et Cham. Linn. **5**. 609. 1830. Mett. n. 91. HB. 337. Chr. 88 — Mexico-Costa Rica. — Micropteris J. Sm. 1875; Lepicystis Diels 1899. NPfl. 323.
(P) **farinosum** Hk. Ic. pl. t. 947. 1854. HB. 329 — Columbia-Ecuador.
fasciale H.B.Willd. sp. **5**. 156. 1810 = P. laevigatum.
(Gr) **fasciatum** (Bl.) Pr. Tent. 180. 1836. Mett. n. 19. HB. 322. NPfl. 308 — Malesia. Ins. Philipp. — Grammitis Bl. Enum. 116. 1828; Fl. Jav. Fil. 117 t. 49 f. 1 — G. longa Fée 1850-52 (an = P. congenerum?).
fasciculatum Pr. Tent. 180. 1836 = P. setigerum.
faucium Liebm. Vid. Selsk. Skr. V. **1**. 209 (seors. 57). 1849 = Dryopteris obliterata.
(P) **Fauriei** Christ, Bull. Boiss. **4**. 672. 1896. NPfl. 312; Farnkr. 83. 1897 — Japonia. — P. vulgare var. japonicum Franch. et Sav. 1876; P. japonicum Maxon 1902; Mak. 1905 (non Houtt. 1783).
(P) **Fawcettii** Bak. JoB. **1889**. 270 — Jamaica.
fecundum Hassk. Cat. pl. Bogor. 5. 1844. = Aspidium irregulare.
(S) **Feei** (Bory) Mett. Pol. 110. n. 212. 1857. NPfl. 316 — Malesia-Polynesia. Selliguea Bory, Dict. class. **6**. 588. 1824; **17** t. 41; Gymnogramme Hk. 1864 pt. ? HB. 389 — Grammitis vulcanica Bl. 1828; Polypodium Bl. 1828. Mett. n. 212 b; Pleuridium J. Sm. 1861; Gymnogramme Hk. 1864 — Polypodium caudiforme Bl. 1828. Chr. 108. NPfl. 316; Grammitis Hk. 1862; Pleopeltis Moore 1862; Gymnogramme Hk. 1864. HB. 388; Selliguea J. Sm. 1866; Dryostachyum caudatum J. Sm. 1841; Polypodium Reinw.; Salom 1883. — conf. Kuhn, Verh. zool. bot. Ges. **25**. 596.
Feei Fourn. Mex. pl. **1**. 82. 1872 = P. plesiosorum var.
felinum Roxb. Calc. Journ. **4**. 495. 1844 — Amboyna. Penang.
(C) **Fendleri** Eat. Mem. Amer. Acad. n. s. **8**. 199. 1860. HB. 349. Chr. 97. NPfl. 315 — Venezuela-Ecuador. — Campyloneurum J. Sm. 1875 (non Moore 1861); C. magnificum Moore 1861; Polypodium Hier. 1904.
fernandense Bak. Ann. Bot. **5**. 462. 1891 = Aspidium.
ferox Hassk. Cat. pl. Bog. 4. 1844 = Dryopteris.
(P) *ferrugineum* Mart. et Gal. Mém. Ac. Brux. **15**. 36 t. 7 f. 2. 1842. Mett. n. 32 — Mexico. — (an P. regiscens var.).
ferrugineum Roxb. Calc. Journ. **4**. 487. 1844 — ? Nephrolepis hirsutula.

ferrugineum Bak. Syn. 318. 1867 = Aspidium Zippelianum.
fibrillosum Bak. Syn. 307. 1867 = Dryopteris.
Fieldingianum Kze.; Mett. Pol. 75. n. 114. 1857 = ? P. lachnopus.
(P) **filicula** Klf. Enum. 275. 1824. Mett. n. 73. (HB. 332) — Amer. trop.
filipendulaefolium Fée, Gen. 240. 1850-52; 6 mém. 11 t. 5 f. 2. 1853 = P. subfalcatum.
filipes Moore, Gard. Chr. **1855**. 388 = ? Arthropteris tenella juv.
filipes Christ in Schwacke, Pl. Nov. Mineiras **2**. 20. 1900 = P. oxylepis.
Filix femina L. sp. **2**. 1090. 1753 = Athyrium.
Filix mas L. sp. **2**. 1090. 1753 = Droypteris.
Finlaysonianum Wall. List n. 2221. 1829 = Alsophila.
firmulum Bak. Kew Bull. **1893**. 211 = Dryopteris.
firmulum Maxon, Contr. U. S. Nat. Herb. **8**. 274 t. 61 f. 3. 1903 = P. Maxonii.
firmum Klf. Wes. d. Farrnkr. 100. 1827 (nomen) = P. Brownii.
(P) **firmum** Kl. Linn. **20**. 378. 1847. Mett. n. 37. HB. 335. NPfl. 309 — Amer. trop. — Ctenopteris J. Sm. 1875. — (= P. rigescens var.?).
Fischerianum Bak. Syn. ed. I. 455. 1868 = Dryopteris grandis*.
(P) **fissidens** Maxon, Contr. U. S. Nat. Herb. **8**. 275. t. 61 f. 4. 1903 — Mexico.
fissum Bak. Syn. 351. 1867 = Cyclophorus porosus et spec. plur.
flabellatum Kl. Linn. **20**. 382. 1847 = Woodsia crenata.
flabelliferum Goldm. Nova Acta **19**. Suppl. I. 455. 1843 = P. Meyenianum.
(P) **flabelliforme** Poir. Enc. **5**. 519. 1804. Mett. n. 41. HB. 326. — Ind. occ. Venezuela-Peru. — Ctenopteris J. Sm. 1875 — Polypodium oligosorum Moritz; Mett. 1869 (non Kl. 1847).
flabellinervum Bak. Journ. Linn. Soc. **22**. 228. 1886 (err. script.) = P. flabellivenium.
(P) **flabellivenium** Bak. Syn. 322. 1867. Chr. 78. NPfl. 308 — Borneo. P. holophyllum Bak. 1879.
(G) **flagellare** Christ, Bull. Boiss. **4**. 660. 1896; Bull. Boiss. **4**. 660. 1896 — Costa Rica.
flagelliferum Roxb. Calc. Journ. **4**. 487 t. 31. 1844 = Nephrolepis exaltata.
flavopunctatum Klf. Enum. 108. 1824 = Dryopteris rotundata.
flexile Fée, 6 mém. 9 t. 2 f. 3. 1853 = P. cultratum var. elasticum.
flexile Moore, Handb. Brit. Ferns ed. II. app. 225. 1853 = Athyrium alpestre var.
(S) **flexilobum** Christ, Bull. Ac. Géogr. Bot. **1904**. 107 — China. Selliguea elliptica var. flagellaris Christ, Bull. Boiss. **7**. 6. 1899 (non Polypodium flagellare Christ 1896).
floccigerum Mett. Pol. 129. n. 262. 1857 = Cyclophorus.
flocculosum Don, Prod. Fl. Nepal. 1. 1825 = Cyclophorus.
fluminense Vell. Fl. flum. **11** t. 66. 1827; Arch. Mus. Nac. Rio Jan. **5**. 447. 1881 = P. fraxinifolium.
foecundum Vell. Fl. flum. **11**. t. 75. 1827; Arch. Mus. Nac. Rio Jan. **5**. 449. 1881 — Brasilia. — (Dryopteris sp. forte = D. alloeoptera.)
foetidum Poir. Enc. **5**. 534. 1804 = Aspidium cicutarium.
foliolosum Wall. List pag. 64 (n. 359). 1829 = Acrophorus stipellatus.
fontanum L. sp. **2**. 1089. 1753 = Asplenium.
(G) **formosanum** Bak. JoB. **1885**. 105 — Formosa.
formosum Raddi, Pl. Bras. **1**. 25 t. 38. 1825 = Dryopteris grandis.
(P) **Forsythianum** Bak. Kew Bull. **1897**. 300 — Madagascar.
Fortunei Kze.; Mett. Pol. 121. n. 241 t. 3 f. 42-45. 1857 = Drynaria.
Fortunii Lowe, Ferns **1**. t. 42 B. 1856 = P. lineare.

(Pl) **fossum** (Moore) Bak. Ann. Bot. 5. 479. 1891 — Easter Island.
Pleopeltis Moore, Gard. Chr. n. s. 18. 586. 1882.

(P) **Fournieri** C. Chr. Ind. 528. 1966 — Mexico. — P. pubescens Fée. 8 mém 87. 1857. HB. 509 (non L. 1759 nec alior.)

fragile L. sp. 2. 1091. 1753 = Cystopteris.

fragile Bak. Journ. Linn. Soc. 16. 203. 1877 = Dryopteris.

fragrans L. sp. 2. 1089. 1753 = Dryopteris.

fragrans L. Mant. 307. 1771 = Cheilanthes pteridioides.

fragrans Vill. Hist. pl. Dauph. 3. 843. 1789 = Dryopteris rigida.

fragrans Lour. Fl. Cochinch. 829. 1790 — Cochinchina.

(M) **Fraseri** Mett.: Kuhn. Linn. 36. 137. 1869. (HB. 510) — Ecuador.

(P) **fraternum** Schlecht. et Cham. Linn. 5. 608. 1830. Mett. n. 87. HB. 337 NPfl. 311 — Mexico. — P. Henchmani J. Sm. 1851.

(G) **fraxinifolium** Jacq. Coll. 3. 187. 1789; Ic. pl. rar. t. 639. 1793. Mett. n. 127. HB. 346. Chr. 94. NPfl. 314 — Mexico-Peru. Brasilia. — Goniophlebium Moore 1857. Polypodium distans Raddi 1825 (non Klf. 1824); Marginaria Pr. 1836; Goniophlebium J. Sm. 1842 — P. triseriale Sw. 1801 — P. rhizocaulon Willd. 1810; Goniophlebium Pr. 1836 — P. articulatum Desv. 1827. Mett. n. 128 (non Poir. 1804); Goniophlebium Pr. 1836.

fraxinifolium Hk. sp. 5. 4. 1863 = Dryopteris vivipara.

Freyreisii Spr. Syst. 4. 55. 1827 = Dryopteris vivipara.

(P) **Friderici et Pauli** Christ. Verh. Nat. Ges. Basel 11. 439. 1896; Ann. Jard. Buit. 15. 148 t. 16 f. 21. 1897 — Celebes.

Friderici et Pauli Christ, Ann. Jard. Buit. II. 4. 37. 1904 = Davallia.

(P) **Friedrichsthalianum** Kze. Farnkr. 2. 55 t. 123. 1850. Mett. n. 93. HB. 339.
(M?) Chr. 87 — Mexico-Costa Rica. — Lepicystis Diels 1899. NPfl. 323.

(P) **fucoides** Christ, Bull. Boiss. II. 5. 2. 1905 — Costa Rica.

fulvescens Hk. et Grev.; Hk. Bot. Misc. 2. 239. 1831 = Dryopteris punctata.

fulvum Mart. et Gal. Mém. Ac. Brux. 15. 33 t. 6. 1842 = P. aureum *areolatum.

fumarioides Weis, Pl. Cr. Fl. Gott. 319. 1770 = Cystopteris fragilis.

(G) **Funckii** Mett. Pol. 57 n. 72. 1857 — Venezuela. Columbia.

funiculosum Desv. Prod. 226. 1827 = P. lycopodioides *surinamense.

(P) **funiculum** Fée, Gen. 241. 1850-52; 6 mém. 12 t. 8 f. 2. Mett. n. 58. HB. 338. Chr. 87. NPfl. 311 — Cuba. — Ctenopteris J. Sm. 1875.

furcatum Sw. Schrad. Journ. 1800². 28. 1801 = Gleichenia.

(P) **furcatum** Desv. Prod. 230. 1827 — Ins. Mascarenae. ? Ins. Comorenses. P. multifidum Bory, Voy. 2. 193 t. 20 f. 2. 3. 1804. Mett. n. 8. HB. 322. Chr. 77. NPfl. 308 (non Jacq. 1789) — ? Grammitis obtusa Willd.: Klf. 1824; Polypodium Kuhn 1868 (non Poir. 1804 nec Spr. 1804).

furcatum Roxb. Calc. Journ. 4. 493. 1844 = ? Gleichenia glauca.

furcatum Mett. Pol. 34. n 9. 1857 = P. dicranophyllum.

furfuraceum Wall. List n. 278. 1828 (nomen) = Cyclophorus acrostichoides.

(M) **furfuraceum** Schlecht. et Cham. Linn. 5. 607. 1830. Mett. n. 98. HB. 332. Chr. 90. — Mexico-Costa Rica. — Marginaria Pr. 1836; Goniophlebium Moore 1857; Lepicystis Diels 1899. NPfl. 323. — Polypodium cryptocarpon Fée 1857 — P. nivosum Fée 1857.

furfuratum Sessé et Mociño, Fl. Mex. 262. 1896 — Cuba.

(P) **fuscatum** Bl. Enum. 129. 1828; Fl. Jav. Fil. 185 t. 86 C. HB. 331. Chr. 81 — Perak. Malesia. — Ctenopteris Kze. 1846. — (an P. mollicomum var.?).

(P) **fuscopilosum** F. Muell. et Bak. JoB. 1887. 169 — Queensland.

(Pl) **fuscopunctatum** Hk. sp. **5**. 69 t. 285 A. 1863. HB. 357 — Ecuador.
(Pl) **galatheae** C. Chr. Bot. Tidsskr. **25**. 79. 1902 — Brasilia.
Craspedaria crispata Fée, Cr. vasc. Br. **1**. 119 t. 36 f. 1. 1869 (non Polypodium L 1753).
Galeottii Mart.; Mart. et Gal. Mém. Ac. Brux. 43 t. 7 f. 3. 1842 = Dryopteris.
(P) *Galeottii* Mett.; Kuhn, Linn. **36**. 135. 1869. HB. 509 — Mexico.
Gardneri Mett. Pol. 129. n. 263. 1857 = Cyclophorus.
Gaudichaudii Bory, Ann. sc. nat. **5**. 471 t. 14. 1825 = Drynaria rigidula.
(G) **Gauthieri** (Fée) C. Chr. Ind. 326. 1905; 529. 1906 — Brasilia.
Goniophlebium Fée, Cr. vasc. Br. **1**. 112 t. 34 f. 3. 1869.
(?) **gedeanum** Racib. Pterid. Buit. 96. 1898 — Java.
(Pl) **geminatum** Schrad. Gött. gel. Anz. **1824**. 867. Mett. n. 175. HB. 353 — Brasilia. — Phymatodes J. Sm. 1857; Pleopeltis Moore 1862; Anapeltis J. Sm. 1866. Polypodium iteophyllum Link 1833; Pleopeltis Link 1841; Drynaria J. Sm. 1846.
geraniifolium Poir. Enc. **5**. 540. 1804 = ? Asplenium cristatum.
Germanianum Bak. Syn. 306. 1867 = Dryopteris.
gestasianum Mett.; Salom. Nom. 299. 1883 = P. tectum.
Gheisbreghtii Linden, Cat. 1858; Bak. Syn. 315. 1867 = Dryopteris Ghiesbreghtii.
Ghiesbreghtii Eaton. Proc. Amer. Acad. **8**. 618. 1873 = P. Eatoni.
(P) **gibbosum** Fée, 6 mém. 8 t. 2 f. 2. 1854; Hier. Hedwigia **44**. 100. 1905 — Amer. trop. — ? P. undulatum Fourn. 1872 (non Willd. 1810).
(G) *giganteum* Desv. Prod. 236. 1827 — Brasilia. — Goniophlebium Moore 1862.
giganteum Wall. List n. 321. 1828 = Alsophila glabra.
giganteum Zoll. Syst. Verz. 37. 1854 = P. heracleum.
(G) **Gilliesii** C. Chr. Ind. 327. 1905; 529. 1906 — Peru. Argentina.
P. pubescens Gill.; Hk. et Grev. Ic. Fil. t. 182. 1830. HB. 342 (non L. 1759 nec alior.); Goniophlebium Moore 1862.
(Gr) **Gilpinae** Bak. Journ. Linn. Soc. **16**. 204. 1877 — Madagascar.
glabellum Hew. Mag. Nat. Hist. II. **2**. 458. 1838 — Jamaica. Trinidad.
glabrum Burm. Fl. Ind. 235. 1768 — Java. Ceylon.
glabrum Roxb.; Wall. List n. 281. 1828; Roxb. Calc. Journ. **4**. 482 t. 28. 1844 = P. punctatum.
glabrum Mett. Pol. 123. n. 245 t. 3 f. 6-8. 1857 = Cyclophorus confluens.
gladiatum Vell. Fl. flum. **11** t. 59. 1827; Arch. Mus. Nac. Rio Jan. **5**. 445. [1881 = P. phyllitidis.
gladiatum Wall. List n. 279. 1828 = P. lineare.
gladiatum Kze. Linn. **9**. 45. 1834 = P. attenuatum.
glanduliferum Liebm. Vid. Selsk. Skr. V. **1**. 206 (seors. 54). 1849 = Dryopteris.
glandulosum Desv. Berl. Mag. **5**. 317. 1811 — Ind. occ. (Dryopteris tetragona?).
glandulosum Hk. sp. **4**. 193 t. 276 A. 1862 = P. corticolum.
glaucescens Bory, Dup. Voy. Bot. **1**. 260. 1828 = P. synammia.
glaucinum Mart. et Gal. Mém. Ac. Brux. **15**. 32 t. 5 f. 1. 1842 = P. aureum [*areolatum.
glaucistipes Wall. List n. 297. 1828 = Drynaria rigidula.
(G) **glaucophyllum** Kze.; Kl. Linn. **20**. 393. 1847; Kze. Farnkr. **1**. 227 t. 93. 1847. Mett. n. 129. HB. 340. Chr. 91. NPfl. 311 — Amer. trop.
Goniophlebium Fée 1850-52 — G. semipinnatifidum Fée 1850-52; Polypodium Mett. n. 130. 1857.

(Pl) **glauco-pruinatum** C. Chr. Ind. 248. 1905; 530. 1906 — Ins. Philipp. Drynaria glauca J. Sm. 1841 (nomen); Brack. Expl. Exp. **16**. 54. 1854. Polypodium Kze.; Mett. Pol. 102. n. 190 t. 2 f. 12, 13. 1857. HB. 365 (non Thbg. 1784 nec Raddi 1825); Pleopeltis Moore 1857; Phymatodes [J. Sm. 1866.

(Pl) **glaucopsis** Franch. Bull. Soc. Fr. **32**. 29. 1885 — Yunnan. Pleopeltis Bedd. 1892.

glaucum Thbg. Fl. Jap. 338. 1784; Houtt. Nat. Hist. **14**. 177. 1783; Pfl. Syst. **13**^{1}. 200. 1786 = Gleichenia.

glaucum Sw. Prod. 134. 1788 = Alsophila pruinata.

? *glaucum* Pr. Rel. Haenk. **1**. 26. 1825 — Ins. Philipp. — (non Polypodium).

glaucum Raddi, Pl. Bras. **1**. 20 t. 29 f. 1. 1825 = P. catharinae.

glaucum Kze.; Mett. Pol. 102. n. 190 t. 2 f. 12, 13. 1857 = P. glauco-pruinatum.

(P) **Glaziovii** Bak. Ann. Bot. **5**. 465. 1891 — Brasilia.

Glaziovii Christ in Schwacke, Pl. Nov. Mineiras **2**. 20. 1900 = P. pectinatum [var.

globuliferum Poir. Enc. **5**. 554. 1804 = ? Dennstaedtia adiantoides.

(Pl) **glossipes** Bak. Ann. Bot. **5**. 476. 1891 — Cambodia.

glycirhiza Fée, 11 mém. 49. 1866 = P. sororium.

(P) **glycyrrhiza** Eat. Amer. Journ. Sci. II. **22**. 138. 1856 — Alaska-California. P. falcatum Kellogg 1854 (non L.fil. 1781); P. vulgare var. HB. 334; P. vulgare var. occidentale Hk. 1840; P. occidentale Maxon 1904.

gonatodes Kze. Linn. **23**. 278, 318. 1850 = P. plesiosorum.

(P?) **goniopteroides** C. Chr. Ind. 188. 1905; 530. 1906 — Panama. Ctenopteris crispata J. Sm. Bot. Voy. Herald 227 t. 48. 1854; Goniopteris Moore 1857; Polypodium Hk. 1863; HB. 361. NPfl. 310 (non L. 1753); Glyphotaenium J. Sm. 1875.

Gordoni Bak. JoB. **1879**. 297 = Dryopteris.

(P) **gracile** Hk. Bot. Misc. **2**. 239. 1831; Hk. et Grev. Ic. Fil. t. 222. 1831. HB. 338 — Peru. — Ctenopteris J. Sm. 1875.

gracile Lowe, Ferns **2** t. 9 A. 1858 = Dryopteris reptans var.

gracilentum Jenm. Bull. Dep. Jam. **4**. 129. 1897 = Dryopteris.

(P) **gracillimum** Copeland in Perkins Fragm. 189. 1905 — Ins. Philipp. (Mindanao).

(P) **gradatum** Bak. Fl. bras. **1**^{2}. 513. 1870 — Brasilia. P. hirsutulum Fée 1869. HB. 509 (non Raddi 1825).

(Gr) **graminellum** C. Chr. Ind. 530. 1906 — Samoa. P. simplex Bak. JoB. **1876**. 12 (non Sw. 1801).

(Gr) **gramineum** Sw. Prod. 130. 1788. Mett. n. 12. HB. 321. NPfl. 308 — Ind. occ. Guiana. — Grammitis linearis Sw. 1801; Lomaphlebia J. Sm. 1875 — Mecosorus nudus Kl. 1847; Polypodium Mett. n. 18. 1857 (non Forst. 1786) — (Asplenium angustifolium Jacq. 1786 vix adhuc pertinet).

gramineum Poir. Enc. **5**. 511. 1804 = P. Billardieri.

gramineum Ces. Atti Acad. Napoli **7**^{8}. 24. 1876 = Vittaria lineata.

grammicum Spr. Neu. Entdeck. **3**. 6. 1822 = Dryopteris decussata.

(P) **grammitidis** R.Br. Prod. Fl. N. Holl. 147. 1810. Mett. n. 60. HB. 327. Chr. 82. NPfl. 310 — Australia-Tasmania. — Grammitis Keys. 1873; Ctenopteris J. Sm. 1875; Grammitis heterophylla Lab. Fl. Holl. pl. spec. **2**. 91 t. 239. 1806; Xiphopteris Spr. 1827 (non Polypodium L. 1753); Gymnogramma Billardieri Klf. 1827; Polypodium Fée 1850-52 (non R.Br. 1810).

(S) **grammitoides** (Bak.) Diels, Engl. Jahrb. **29**. 205. 1900 — China. Gymnogramme Bak. JoB. **1889**. 178; Selliguea Christ 1903.

grande Pr. Del. Prag. 1. 171. 1822 = Dryopteris.
grandidens Kze. Bot. Zeit. **1846**. 423 = P. persicifolium var.
(Pl) **grandidentatum** (Ces.) Bak. Ann. Bot. **5**. 479. 1891 — Borneo.
P. dilatatum var. grandidentatum Ces. Atti Ac. Napoli 7^8. 27. 1876.
grandifolium Wall. List n. 282. 1828 = P. membranaceum.
granulosum Pr. Rel. Haenk. 1. 24 t. 4 f. 2. 1825 = Dryopteris.
granulosum Benth. Fl. Hongk. 459. 1861 = Dryopteris urophylla.
gratum Fée, Cr. vasc. Br. 1. 242 t. 76 f. 2. 1869 = P. taxifolium.
(P) **graveolens** Bak.; Jenm. JoB. **1877**. 265 — Jamaica. Columbia.
(P) **grenadense** Jenm. Gard. Chr. III. **15**. 134. 1894 — Grenada.
Grevilleanum Wall. List pag. 250 (n. 5169). 1829 = P. excavatum.
(Pl) **Griffithianum** Hk. Ic. pl. t. 951. 1854. HB. 359. NPfl. 316 — China merid. India bor. Borneo. — Pleopeltis Moore 1857; Phymatopsis J. Sm. 1875.
Griffithii Hk. sp. **4**. 236. 1862 = Dryopteris brunnea.
(P) **Grisebachii** Und. in C. Chr. Ind. 531. 1906—Jamaica. Guadeloupe. Martinique.
P. exiguum Gris. Fl. br. W. Ind. 701. 1864. HB. 326 (non Hew. 1838).
(Pl) **griseo-nigrum** Bak. Kew Bull. **1895**. 55 — Yunnan.
griseum Schkuhr, Kr. Gew. 1. 25 t. 25 b. 1806 = Alsophila quadripinnata.
griseum Liebm. Vid. Selsk. Skr. V. 1. 198 (seors. 46). 1849 = P. Liebmanni.
grossum Langsd. et Fisch. Ic. Fil. 9 t. 8. 1810 = P. phymatodes.
guadalupense Wikstr. Vet. Ak. Handl. **1825**. 435. 1826 — Guadeloupe.
guadalupense Fée, Cr. vasc. Br. 1. 95. 1869 (nomen) — Guadeloupe.
(Ph?) *guatemalense* Kl. Allg. Gartenzeit. **1855**. 33 — Guatemala.
Goniophlebium Moore 1862.
Guatemalense Hk. sp. **5**. 29. 1863 = P. Lowei.
Gueintzii Mett. Pol. 91. n. 161 t. 3 f. 18, 19. 1857 = P. lineare.
guianense Aublet, Hist. pl. Guian. **2**. 962. 1775 = Leptochilus.
(P) **Guilleminianum** Fourn. Mex. pl. 77 t. 2. 1872. HB. 510 — Mexico.
gymnogrammoides Bak. JoB. **1888**. 229 = Dryopteris.
(Pl) **gyroflexum** Christ, Bull. Boiss. **6**. 994. 1898 — Brasilia. (Amazonas).
(P) **Haalillioanum** Brack. Expl. Exp. **16**. 5 t. 1 f. 4. 1854. NPfl. 309 — Ins. Hawaii.
Haenkeanum Spr. Syst. **4**. 46. 1827 = P. stigmaticum.
Haenkeanum Pr.; Ettingsh. Farnkr. 42 t. 24 f. 1. 2. 1865 = P. vacciniifolium.
Haenkei Liebm. Vid. Selsk. Skr. V. 1. 204. 1849 = P. attenuatum.
(P) **Hahnii** Fourn. Mex. pl. 1. 79. 1872 — Mexico.
(Pl) **hainanense** C. Chr. Ind. 531. 1906 — Hainan. (Formosa?).
P. dimorphum Bak. Ann. Bot. **5**. 477. 1891 (non Link 1833).
Hamiltonianum Diels, Nat. Pfl. 1^4. 316. 1899 = P. pedunculatum.
(Pl) **hammatisorum** Harr. Journ. Linn. Soc. **16**. 32. 1877 — Luzon.
(Pl) **Hancockii** Bak. JoB. **1885**. 106 — Formosa.
Hancockii Bak. Ann. Bot. 8. 127. 1894 = Aspidium sumatranum.
harpeodes Link, Hort. Berol. **2**. 97. 1833 = P. latipes.
(P?) ?**harpophyllum** Zenker; Kze. Linn. **24**. 256. 1851. Mett. n. 88. HB. 511 — India austr. Celebes. — (est f. Beddome = Polystichum auriculatum).
(Pl) **Harrisii** Jenm. Gard. Chr. III. **27**. 241. 1900 — Jamaica.
(P) **Hartii** Jenm. JoB. **1886**. 272; Hier. Hedw. **44**. 95. 1905 — India occ.
(P) **Hartwegianum** Hk. in Benth. Pl. Hartw. 54. 1840; Ic. pl. t. 340. 1840. HB. 336 — Mexico. — (an = P. puberulum?).
Hasseltii Bl. Fl. Jav. Fil. 195 t. 92. 1829 = Dryopteris.
hastaefolium Sw. Schrad. Journ. **1800**^{2}. 25. 1801 = Dryopteris sagittata.

(Pl) **hastatum** Thbg. Fl. Jap. 335. 1784. Mett. n. 201. HB. 361. Chr. 109. NPfl. 318 — Japonia. Formosa-China-Himalaya-Ceylon. Ins. Philipp.
Drynaria Fée 1850-52; Pleopeltis Moore 1862 — Polypodium trifidum Don 1825. HB. 363. NPfl. 318 (non Hoffm. 1790 nec With. 1796); Pleopeltis Bedd. 1870; Phymatopsis J. Sm. 1875 — Polypodium oxylobum Wall. 1828. Mett. n. 202. Phymatodes Pr. 1836; Pleopeltis Bedd. 1863; Pleuridium J. Sm. 1866.
hastatum Sw. Fl. Ind. occ. **3**. 1653. 1806 = Dryopteris sagittata.
hastatum Vell. Fl. flum. **11** t. 68. 1827; Arch. Mus. Nac. Rio Jan. **5**. 447. 1881 = Aspidium martinicense.
(Gr) **Havilandii** Bak. Tr. Linn. Soc. II. Bot. **4**. 253. 1894 — Borneo.
(P) **hawaiiense** Und. ap. Heller, Minn. bot. St. **1**. 784. 1897 — Ins. Hawaii.
Haynaldii Sod. Rec. 61. 1883 = Aspidium.
(Pl) **hederaceum** Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 215 c. fig. — China.
Heleopteris Borck. Röm. Arch. 1^3. 19. 1798 = Dryopteris filix mas.
(P) **Helleri** Und. ap. Heller, Minn. bot. St. **1**. 785. 1897 — Ins. Hawaii.
(Pl) **Hellwigii** Diels in Schum. et Laut. Fl. deut. Südsee 140 t. 3 f. A, B. 1901 — N. Guinea.
(Pl) **hemionitideum** Wall. List n. 284. 1828 (nomen). Mett. n. 215. HB. 360. Chr. 105. NPfl. 315 — China merid. Ind. bor.-austr. (? Malesia).
Selliguea Pr. 1836 (nomen); Drynaria J. Sm. 1841 (nomen); Colysis Pr. Epim. 147. 1849; Pleopeltis Moore 1862.
hemionitis Cav. Descr. 248. 1802 — Ins. Mariannae.
(Pl) **hemitomum** Hance, JoB. **1883**. 269 — China. — P. deltoideum Bak. 1888 (non Sw. 1788 nec Liebm. 1849). — (an P. phyllomanes forma?).
Hemsleyanum Bak.; Hemsl. Biol. Centr. Amer. **3**. 660 t. 108. 1885 = Dryopteris.
Henchmani J. Sm.; Moore et Houlst. Gard. Mag. Bot. **3**. —. 1851; Lowe, Ferns **1** t. 30. 1856 = P. fraternum.
Hendersoni Lowe, Ferns **2** t. 37 B. 1858 = P. angustifolium.
Hendersoni Atkins.; Bak. Syn. 511. 1874 = P. Atkinsoni.
Henriquesii Bak. Bol. Soc. Brot. **4**. 154 t. 1. 1886 = Dryopteris.
Henryi Christ, Bull. Boiss. **6**. 873. 1898 = P. austrosinicum.
(S) **Henryi** (Bak.) Diels, Nat. Pfl. 1^4. 316. 1899 — China.
Gymnogramme Bak. JoB. **1887**. 171; Selliguea Christ 1898.
(My) **heracleum** Kze. Bot. Zeit. **1848**. 117. Mett. n. 231. HB. 366. Chr. 116. NPfl. 319 — Malesia. N. Guinea. — Drynaria Moore 1862.
(C) **herbaceum** Christ in Schwacke, Pl. Nov. Mineiras **2**. 22. 1900; Bull. Boiss. II. **2**. 375. 1902 — Brasilia.
(P) **hesperium** Maxon, Proc. Biol. Soc. Wash. **13**. 200. 1900 — U. S. occ.
Heteractis Mett.; Kuhn, Linn. **36**. 140. 1869 = Cyclophorus.
(S) **heterocarpum** (Bl.) Mett. Fil. Lips. 37 t. 25 f. 24-25. 1856. Pol. n. 211. NPfl 316 — Java. Celebes. — Grammitis Bl. Enum. 118. 1828; Selliguea Bl. Fl. Jav. Fil. 125 t. 52 f. 1. 1828; Gymnogramme Hk. 1864. HB. 389; Polypodium Mettenianum Ces. 1876.
heterocarpum Bl. Fl. Jav. Fil. 167 t. 75. 1829 = P. Zollingerianum.
heteroclitum Desv. Berl. Mag. **5**. 318. 1811 = Dryopteris.
heteroclitum Fée, Cr. vasc. Br. **1**. 93 t. 26 f. 4. 1869 = P. pectinatum var.
(Pl) **heterolobum** C. Chr. Ind. 532. 1906 — Luzon. — P. anomalum Christ, Bull. Boiss. **6**. 201 t. 3 f. 3. 1898 (non Hk. et Arn. 1856).

(P) **heteromorphum** Hk. et Grev. Ic. Fil. t. 108. 1829. Mett. n. 48. HB. 334. NPfl. 310 — Mexico. Ecuador. — Arthropteris J. Sm. 1875 — Polypodium variabile Mett.; Kuhn 1869.

heterophlebium Bak. JoB. **1884**. 363 = Dryopteris.

heterophyllum L. sp. **2**. 1083. 1753. (Plum. t. 120) — Ind. occ.

heterophyllum Pr. Tent. 179. 1836 (nomen) — ?

heterosorum Bak. Syn. 506. 1874 = Aspidium vastum.

(P) **heterotrichum** Bak.; Jenm. JoB. **1879**. 262 — Jamaica.

Hewardii Gris. Fl. br. W. Ind. 696. 1864 = Dryopteris heteroclita.

hexagonopterum Michx. Fl. bor. Am. **2**. 271. 1803 = Dryopteris.

(P) **Hieronymusii** C. Chr. Ind. 533. 1906 — India posterior. — P. sertularioides J. Sm 1841 (nomen, non Bak. 1876); Hier. Hedwigia **44**. 98. 1905.

(P) **Hildebrandtii** Hier. Hedwigia **44**. 91. 1905 — Madagascar.

(P) **Hillebrandii** Hk. sp. **4**. 228 t. 279 A. 1862. NPfl. 311 — Ins. Hawaii.

Hillebrandii Hk. sp. **4**. 254. 1862 = Dryopteris honolulensis.

Hillii Bak. Syn. 505. 1874 = Dryopteris.

(Pl) **himalayense** Hk. sp. **5**. 91. 1863. HB. 369. Chr. 114. NPfl. 321 — Himalaya. China. — Pleopeltis Bedd. 1869; Polypodium venustum Wall. 1828; Clarke 1880 (non Desv. 1811); Pleuridium J. Sm. 1857; Arthromeris [J. Sm. 1875.

Hippocrepis Jacq. Coll. **3**. 186. 1789 = Aspidium.

hirsutissimum Raddi, Opusc. sci. Bol. **3**. 286. 1819; Pl. Bras. **1**. 17 t. 26. 1825 = P. lepidopteris.

hirsutulum Forst. Prod. 81. 1786 = Nephrolepis.

(G) *hirsutulum* Raddi, Pl. Bras. **1**. 21 t. 29 f. 2. 1825 — Brasilia. Goniophlebium Moore 1862.

hirsutulum Fée, Cr. vasc. Br. **1**. 87 t. 36 f. 2. 1869 = P. gradatum.

hirsutum Desv. Berl. Mag. **5**. 317. 1811 — Guiana. — (Dryopteris sp.).

hirsutum Sw. Vet. Akad. Handl. **1817**. 61 = Dryopteris hirtula.

(Gr) **hirtellum** Bl. Enum. 122. 1828. Mett. n. 14. HB. 320 — China merid. Perak. Malesia. Ins. Philipp. N. Caledonia. — Grammitis pusilla Bl. Fl. Jav. Fil. 109 t. 46 f. 4-6. 1828; Polypodium Christ 1897 — P. alpestre Bl. 1828 — P. lasiosorum Hk. 1862 — P. adspersum Bl. 1828 (non Schrad. 1818); Grammitis Bl. 1828 — ? G. nana Fée 1856.

hirtisorum Desv. Berl. Mag. **5** 313. 1811 = P. piloselloides.

hirtulum Kze.; Steud. Nom. **2**. 341. 1824 = Dryopteris.

hirtum Sw. Prod. 133. 1788 = Dryopteris.

hirtum Schkuhr, Kr. Gew. **1**. 25 t. 25 c. 1806 — Jamaica.

hirtum Pr. Rel. Haenk. **1**. 27. 1825 = Dryopteris setosa.

hirtum Vell. Fl. flum. **11** t. 79. 1827; Arch. Mus. Nac. Rio Jan. **5**. 450. 1881 = Dryopteris parasitica.

hirtum Pr. Tent. Pter. 180. 1836 = P. diplosorum.

hirtum Mett. Pol. 38 n. 22. 1857 = P. diplosorum et P. pubinerve.

hirtum Hk. sp. **4**. 170 t. 273 A. 1862. HB. 320 = P. diplosorum, P. Reinwardtii, P. pubinerve.

(Gr) **holophlebium** Bak. JoB. **1880**. 370 — Madagascar.

holophyllum Bak. JoB. **1879**. 43 = P. flabellivenium.

holophyllum Bak. JoB. **1888**. 325 = Dryopteris.

honestum Kze. Linn. **9**. 49. 1834 = Dryopteris.

Honolulense Hk. sp. **4** 288 (nota). 1862 = Dryopteris.

(Gr) **Hookeri** Brack. Expl. Exp. **16**. 4. 1854. HB. 319. Chr. 78. NPfl. 308 — Polynesia. Austr. — P. setigerum Hk. et Arn. Bot. Beech. Voy. 103 t. 21 A. 1832 (non Bl. 1828); Grammitis J. Sm. 1875 — P. conforme Brack. 1854; Grammitis J. Sm. 1875.

Hookeri Cord. Fl. Réun. 84. 1895 = P. barbatulum.

(P) **Hornei** Bak. JoB. **1879**. 298 — Fiji.

horridum L. sp. **2**. 1092. 1753 = Hemitelia.

Horsfieldii R.Br.; Wall. List n. 286. 1828; R.Br. in Horsf. Pl. Jav. rar. t. 1. 1838 = Dipteris conjugata.

(S) **Hosei** C. Chr. Ind. 334. 1905; 534. 1906 — Borneo. Perak. Gymnogramme campyloneuroides Bak. Journ. Linn. Soc. **24**. 261. 1887 (non Polypodium Bak. 1883); Selliguea Bedd. 1892.

Hostmanni Kl. Linn. **20**. 397. 1847 = Dryopteris reticulata* sorbifolia.

hoyaefolium Wall. List n. 2230. 1829 (nomen) — Martaban.

Huegelii Fée; Ettingsh. Farnkr. 35 f. 14-16 t. 22 f. 8. 1865 = P. superficiale.

(Pl) **Humblotii** C. Chr. Ind. 534. 1906 — Madagascar. P. inconspicuum Bak. JoB. **1884**. 143 (non Bl. 1828).

Humboldtii Poir. Enc. Suppl. **4**. 497. 1816 = Alsophila blechnoides.

(P) **humile** Mett. Ann. sc. nat. V. **2**. 251. 1864. (HB. 325) — Columbia.

hydrophilum Bak. Ann. Bot. **5**. 456. 1891 = Dryopteris.

(P) *hygrometricum* Splitgb. Tijdschr. Nat. Gesch. **7**. 409. 1840 — Para.

hymenodes Wall. List n. 283. 1828 = Leptochilus axillaris.

hymenodes Kze. Linn. **23**. 279, 319. 1850; Mett. Fil. Lips. 37 t. 25. f. 40-41. 1856 = P. superficiale.

(Pl) **hymenolepioides** Christ, Journ. de Bot. **19**. 75. 1905 — Annam.

(P) **hymenophylloides** Klf. Enum. 118. 1824. Mett. n. 1. HB. 337. Chr. 89. NPfl. 311 — Ins. Hawaii. — Adenophorus Hk. et Grev. 1830; A. minutus Gaud. 1825; Amphoradenium Desv. 1827.

hyperboreum Sw. Schrad. Journ. **1800**². 27. 1801 = Woodsia alpina.

ichtiosmum Sod. Rec. 59. 1883 = Dryopteris.

ilvense Vill. Hist. pl. Dauph. **3**. 848. 1789 = Woodsia.

imbricatum Liebm. Vid. Selsk. Skr. V. **1**. 210 (seors. 58). 1849 = Dryopteris.

imbricatum Karst. Ann. Jard. Buit. **12**. 168 t. 15 f. 29. 30, t. 16 f. 49. t. 19. 1895 = P. mirabile.

(Pl) *immersum* Vahl, Symb. **3**. 104. 1794 — India. — Phymatodes Pr. 1836.

immersum Fée, Cr. vasc. Br. **1**. 88 t. 27 f. 1. 1869 = P. depressum.

imponens Ces. Rend. Ac. Napoli **16**. 27. 29. 1877 = Dryopteris.

impuber Roxb. Calc. Journ. **4**. 494. 1844 — Amboyna.

inaequale Klf.; Link, Hort. Berol. **2**. 107. 1833 = Dryopteris subincisa.

inaequale Lowe, Ferns **2**. t. 28. 1858 = P. Lowei.

inaequale (Pr.) Ettingsh. Denkschr. Akad. Wien **22**. 93 t. 20 f. 4. 1864 — Nepal. — Drynaria Pr. hb.

inaequale Fée, 11 mém. 47 t. 12 f. 3. 1866 = P. curvatum.

inaequalifolium Colla, Mem. Ac. Torino **36**. 49. 1836 — Juan Fernandez.

incanoides Fée, 8 mém. 88. 1857 = P. polypodioides.

incanum Sw. Prod. 131. 1788 (incannum) = P. polypodioides.

incisum Sw. Prod. 131. 1788 = Dryopteris.

incisum Hoffm. in Roem. et Ust. Bot. Mag. **9**. 10 f. 13 b. 1790 = Athyrium filix femina.

(P) **inconspicuum** Bl. Enum. 130. 1828; Mett. Ann. Lugd. Bat. **2**. 221. 1866. (HB. 327) — Java. Ins. Philipp.
P. parvulum Bl. Fl. Jav. Fil. 187 t. 37 C. 1828 (non Bory 1810).
inconspicuum Bak. JoB. **1884**. 143 = P. Humblotii.
(Pl) **incurvatum** Bl. Enum. 126. 1828. Fl. Jav. Fil. 151 t. 65. 1828. Mett. n. 200. HB. 363. Chr. 108. NPfl. 318 — Malacca. Java. Borneo.
Pleopeltis Moore 1860; Phymatodes J. Sm. 1866; Phymatopsis J. Sm. 1875. (an P. trilobum Houtt. 1786?).
(P) **induens** Maxon, Bull. Torr. Cl. **32**. 75. 1905 — Jamaica.
P. saxicolum Bak. JoB. **1877**. 264 (non Sw. 1817).
(Pl) **induratum** Bak. Ann. Bot. **5**. 475. 1891. NPfl. 315 — N. Guinea.
(Pl) **insigne** Bl. Enum. 127. 1828. Fl. Jav. Fil. 166 t. 74. 1829. Mett. n. 228. HB. 363. NPfl. 318 — Malacca. Java. Ins. Philipp. — Pleopeltis Bedd. 1867; Colysis J. Sm. 1875; Drynaria decurrens Brack. 1854; Pleopeltis Moore 1862. Polypodium tridactylon Mett. n. 196. 1857 (non Hk. et Grev. 1831).
insigne Liebm. Vid. Selsk. Skr. V. **1**. 193 (seors. 41). 1849 = P. plectolepis.
integerrimum Noronha, Prod. Phyt. ex Thouars, Mél. bot. et voy. 1811 — Madagascar.
integerrimum Vell. Fl. flum. **11**. t. 60. 1827; Arch. Mus. Nac. Rio Jan. **5**. 445. 1881 = Nephrolepis biserrata*.
intermedium Colla, Mem. Ac. Torino **39**. 51 t. 74. 1836 = P. translucens.
intermedium Hk. et Arn. Bot. Beech. Voy. 405. 1841 = P. californicum.
intermedium Rupr. Distr. Cr. vasc. Ross. 51. 1845 = Dryopteris phegopteris.
(S) **interruptum** C. Chr. Ind. 333. 1905; 535. 1906 — Borneo. — Gymnogramme acuminata Bak. JoB. **1888**. 326 (non Klf. 1824 nec Polypodium Don 1825).
(Pl) **intramarginale** Bak.; Christ, Bull. Boiss. II. **3**. 509. 1903 — China.
(Gr) **intromissum** Christ, Verh. Nat. Ges. Basel **11**. 440. 1896; Ann. Jard. Buit. **15**. 148 t. 15 f. 20. 1897 — Celebes.
inversum Vell. Fl. flum. **11** t. 72. 1827; Arch. Mus. Nac. Rio Jan. **5**. 448. [1881 = P. plumula.
invisum Forst. Prod. 81. 1786 = Dryopteris.
invisum Sw. Prod. 133. 1788 = Dryopteris patens*.
involucratum Roxb. Calc. Journ. **4**. 491. 1844 — Amboyna. (Dryopteris sp.).
involutum Desv. Berl. Mag. **5**. 318. 1811 = Dryopteris heteroclita.
involutum Mett. Fil. Lips. 37 t. 25 f. 26. 27. 1856 = P. scolopendrinum.
involutum Bak. JoB. **1889**. 177 = P. eilophyllum.
(S) **iridifolium** (Christ) Diels, Nat. Pfl. **1**^{4}. 316. 1899 — Celebes. Java.
Gymnogramme Christ, Verh. Nat. Ges. Basel **11**. 248. 1895; Ann. Jard. Buit. **15**. 170 t. 16 f. 26. 1897.
irioides Poir. Enc. **5**. 513. 1804 = P. punctatum.
irregulare Pr. Rel. Haenk. **1**. 25 t. 4 f. 3. 1825 = Aspidium.
(G) **Irvingii** Kuhn, Fil. Afr. 147. 1868 (nomen). NPfl. 311 — Africa occ. trop.
P. glaucophyllum β Hk. sp. **5**. 18. 1863.
(G) **isomerum** Fourn. Mex. pl. **1**. 80 t. 3. 1872 — Mexico.
iteophyllum Link, Hort. Berol. **2**. 88. 1833 = P. geminatum.
(Gr) **Jagorianum** Mett.; Kuhn, Linn. **36**. 128. 1869. HB. 507 — Ins. Philipp.
jaintense Clarke, Tr. Linn. Soc. II. Bot. **1**. 552 t. 82 f. 4. 1880 = Cyclophorus laevis.
jamaicense Desv. Berl. Mag. **5**. 318. 1811 — Jamaica. — (Dryopteris sp.).
Jamesoni Mett.; Salom. Nom. 302. 1883 = P. angustifolium.
Jamesoni Jenm. Bull. Dep. Jam. **4**. 112. 1897 = P. myosuroides.

(P) **jamesonioides** Fée, 7 mém. 59 t. 21 f. 4. 1857 — Columbia.
japonense Mak. Bot. Mag. Tokyo **12**. 165. 1898 = P. annuifrons.
japonicum Hoult. Nat. Hist. **14**. 167 t. 98 f. 3. 1783; Pfl. Syst. **13**. 187 t. 98 f. 3. 1786 = Polystichum falcatum.
japonicum Maxon, Fern Bull. **10**. 42. 1902; Mak. Bot. Mag. Tokyo **19**. 139. 1905 = P. Fauriei.
(P) **jubiforme** Klf. Flora **1823**[1]. 364. Mett. n. 40. HB. 327. NPfl. 310 — Ind. occ. Panama. Brasilia. Guiana. — Ctenopteris J. Sm. 1875 — Polypodium saccatum 1850-52 — P. confusum J. Sm. 1842; Ctenopteris J. Sm. 1875.
juglandifolium H.B.Willd. sp. **5**. 195. 1810 = Phanerophlebia.
juglandifolium Don, Prod. Fl. Nepal. 3. 1825 = P. Wallichianum.
(P) **jungermannioides** Kl. Linn. **20**. 373. 1847. Mett. n. 13. HB. 320. NPfl. 308 — Guatemala-Chile.
(P) **Kalbreyeri** Bak. Tr. Linn. Soc. II. Bot. **2**. 291. 1897 — Guiana.
Karstenianum Kl. Linn. **20**. 390. 1847 = Dryopteris subincisa*.
Karwinskianum A.Br.; Kze. Linn. **23**. 279. 319. 1850; Mett. Pol. 66. n. 95. 1857 = P. plebejum et P. leucosticton.
Karwinskianum Lowe, Ferns **2** t. 40. 1858 = P. loriceum.
Kaulfussii Wikstr. Vet. Ak. Handl. **1825**. 443. 1826 = Dryopteris caudata.
Kaulfussii Pr. Tent. 178. 1836 = P. pseudo-grammitis.
(P) **Kegelianum** Kze. Lin. **21**. 210. 1848. Mett. n. 45 — Guiana.
Kennedyi F. Muell. Fragm. **4**. 165. 1864 (syn.) = Dryopteris urophylla.
Keraudrenianum Gaud. Freyc. Voy. Bot. 362 t. 7. 1827 = Dryopteris.
(P) **khasyanum** Hk. Ic. pl. t. 949. 1854. HB. 325. NPfl. 309 — Khasya. Assam. Perak. — Cryptosorus Fée 1865.
Knightiae ht. Sander, Gard. Chr. III. **33**. 245 f. 99. 1903 — Hort. (Australia).
(P) **Knudsenii** Hier. Hedwigia **44**. 79. 1905 — Ins. Hawaii.
(Gr) **knutsfordianum** Bak. JoB. **1890**. 107 — N. Guinea.
Koenigii Bl. Fl. Jav. Fil. 50. 1828 = ? Cyclophorus adnascens.
(P) **Kookenamae** Jenm.; Bak. Tr. Linn. Soc. II Bot. **2**. 292. 1887 — Guiana.
(Gr) **Koordersii** Christ, Ann. Jard. Buit. **15**. 144 t. 15 f. 19. 1897. NPfl. 308. — Celebes.
(G) **Korthalsii** Mett. Ann. Lugd. Bat. **2**. 223. 1866. HB. 345. NPfl. 314 — Perak. Sumatra. — Goniophlebium Bedd. 1892. (an P. lampongense?).
Krameri Franch. et Sav. Enum. pl. Jap. **2**. 244 (Kremeri). 1876 (nomen); 641. 1879 = Dryopteris oyamensis.
(G) **Kuhnii** Fourn. Bull. Soc. Fr. **19**. 251. 1872 — Nicaragua.
Kunthii Desv. Prod. 226. 1827 = P. rosmarinifolium.
(G) **Kunzeanum** C. Chr. Ind. 325. 1905; 536. 1906 — Peru. Brasilia.
P. cordatum Kze. Linn. **9**. 44. 1834. Mett. n. 147 t. 2 f. 1-2. HB. 345. NPfl. 314 (non Desv. 1827); Goniophlebium Moore 1862.
Labrusca Hk. sp. **5**. 73 t. 285 B. 1863 = Aspidium.
lacerum Thbg. Fl. Jap. 337. 1784 = Dryopteris.
(P) **lachniferum** Hier. Engl. Jahrb. **34**. 515. 1904 — Columbia-Ecuador.
lachnopodium J. Sm. Bot. Mag **72** Comp. 8. 1846 = Dryopteris ampla.
(G) **lachnopus** Wall. List n. 310. 1828 (nomen); Hk Ic. pl. t. 952. 1854. Mett. n. 113. HB. 342. NPfl. 312 — Himalaya.
Goniophlebium J. Sm. 1840; Schellolepis J. Sm. 1875 — Polypodium Fieldingianum Kze.; Mett. n. 114. 1857; Goniophlebium Moore 1862.
laciniatum Burm. Fl. Ind. 231. 1768 (Rheede t. 20, 21) — Java. India. (vix filix).

laciniatum Lam. Fl. franç. 1. 14. 1778 = P. vulgare var. cambricum.
laciniatum Gmel. Syst. Nat. 2[2]. 1311. 1791 = Dryopteris dissecta.
laciniatum Noronha, Prod. phyt. ex Thouars, Mél. Voy. et Bot. —. 1811 — [Madagascar.
laciniatum Bl. Enum. 131. 1828 = P. macrochasmum.
laetum Salisb. Prod. 403. 1796 = Athyrium filix femina.
laetum Raddi, Opusc. sci. Bol. 3. 287. 1819; Pl. Bras. 1. 19. t. 28. 1825 = [P. loriceum var.
laeve Mett.; Kuhn, Linn. 36. 139. 1869 = Cyclophorus.
(C) **laevigatum** Cav. Descr. 244. 1802. HB. 348. Chr. 96. NPfl. 314 — Amer. trop. Campyloneurum Pr. 1836. Polypodium lapathifolium Poir. 1804 — P. fasciale H.B.Wild. 1810. Mett. n. 137; Campyloneurum Pr. 1836 — C. Fendleri Moore 1862 — C. minus Fée 1850-52 — C. acrocarpon Fée [1869.
laevigatum Bak. Syn. 505. 1874 = Dryopteris.
laevigatum var. crispatum C. Chr. Bot. Tidsskr. 25. 79. 1902 = P. vulpinum.
lagopodioides Desv. Berl. Mag. 5. 313. 1811 = P. lycopodioides.
lagopodioides Christ, Ann. Jard. Buit. II. 4. 37. 1904 = P. albulum.
(Pl) **lagunense** Christ, Bull. Boiss. 6. 201 t. 3 f. 4. 1898 — Luzon.
(G) **lampongense** Zoll. Syst. Verz. 37, 48. 1854 — Java. — (conf. P. Korthalsii).
(S) **lanceola** Mett. Ann. sc. nat. IV. 15. 78. 1861. HB. 358 — N. Caledonia. Pleopeltis Carr. 1873; Selliguea Fourn. 1873; S. peltatisquama Fourn. 1869.
lanceola F. Muell. Fragm. 7. 120. 1870 = P. simplicissimum.
lanceolato-cristatum Hoffm. in Röm. et Ust. Mag. 9. 9. 1790 = Dryopteris spinulosa.
(Pl) **lanceolatum** L. sp. 2. 1082. 1753. HB. 356. Chr. 103 — Amer. trop. — Chile austr. Juan Fernandez. Africa occ. et austr. c. ins. Tristan d'Acunha, St. Helena. India. Ins. Hawaii. — Pleopeltis Klf. 1824; Phlebodium J. Sm. 1841; Niphobolus Keys. 1873; Lepicystis Diels 1899. NPfl. 323. Polypodium lepidotum Willd.; Schlecht. 1825. Mett. n. 155; Pleopeltis Pr. 1836; Drynaria Fée 1850-52. Polypodium macrocarpum Bory; Willd. 1810; Pleopeltis Klf. 1824; Phlebodium J. Sm. 1841; Drynaria Fée 1850-52 — Polypodium peltatum Cav. 1802 — P. marginale Willd. 1810; Pleopeltis Klf. 1824 — P. ensifolia Carm. 1818.
—* Grammitis elongata Sw. 1806; Synammia Pr. 1836; Phlebodium J. Sm. 1841; Mecosorus Kl. 1847; Drynaria Fée 1850-52; Loxogramma Moore 1857; Pleopeltis J. Sm. 1857; Polypodium Mett. n. 154. 1857 (non Ait. 1789); Gymnogramme Hk. 1864. HB. 387; Taenitis Swartzii Jenm. 1879; Grammitis lanceolata Schkuhr, Kr. Gew. 1. 9 t. 7. 1805 (non Sw. 1801). — Drynaria Prieurii Fée 1850-52; Polypodium Christ 1901 — Grammitis robusta Phil. 1896 — Drynaria crassinervata Fée 1857; Pleopeltis Moore 1862; Polypodium Kiærsk. 1874. Drynaria mexicana Fée 1857.
lanciforme Pr. Del. Prag. 1. 164. 1822 = P. laevigatum.
(P) **lanigerum** Desv. Berl. Mag. 5. 316. 1811 — Peru. Venezuela. ? P. lanigerum Eat. 1860. HB. 335; P. concinnum Mett.; Kuhn. Linn. 36. 132. 1869 (non Willd. 1810).
lanigerum Mett. Pol. 48. n. 49. 1857 = P. laxum.
lanigerum Eat. Mem. Amer. Acad. n. s. 8. 198. 1860; Hk. sp. 4. 189. 1862 = ? P. lanigerum Desv.
lanosum Poir. Enc. 5. 538. 1804 = Cheilanthes lanosa et Woodsia ilvensis.
lanosum Fée, Gen. 237. 1850-52 = P. lepidopteris var. rufulum.
(G) *lanuginosum* Schrad. Gött. gel. Anz. 1824. 867 — Brasilia. Goniophlebium Moore 1862. (an P. dasypleuron Kze. 1834?).

lanugisosum Vell. Fl. flum. **11** t. 70. 1827; Arch. Mus. Nac. Rio Jan. **5**. 448. 1881 = Dryopteris falciculata.
lanuginosum Nees, Linn. **19**. 683. 1847 = ? P. thyssanolepis.
lapathifolium Poir. Enc. **5**. 514. 1804 = P. laevigatum.
laserpitiifolium Scort.; Bedd. JoB. **1887**. 324 = Dryopteris.
lasiernos Spr. Syst. **4**. 61. 1827 = Dryopteris hirtula.
lasiolepis Mett.; Kuhn, Linn. **36**. 130. 1869 = P. tenuiculum.
(G) **lasiopus** Kl. Linn. **20**. 393. 1847. Mett. n. 109. HB. 343 — Mexico. Columbia. Venezuela. Guadeloupe. — Goniophlebium Moore 1862.
lasiosorum Hk. sp. **4**. 166. 1862 = P. hirtellum var.
(P) **lasiostipes** Mett. Ann. sc. nat. IV. **15**. 76. 1861. HB. 327 — N. Caledonia.
(Pl) **Lastii** Bak. JoB. **1891**. 5 — Madagascar.
latebrosum Salisb. Prod. 403. 1796 = Dryopteris phegopteris.
latebrosum Wall. List n. 318. 1828 = Alsophila.
laterepens Trotter; Hope, Journ. Bomb. Nat. Hist. Soc. **12**. 628 t. 14. 1899 [= Dryopteris.
(Pl) **lateritium** Bak. Ann. Bot. **5**. 476. 1891 — Cambodia.
latifolium Forst. Prod. 83. 1786 = Aspidium.
latifolium Poir. Enc. **5**. 512. 1804 = P. platyphyllum.
latifolium Vahl, Ecl. **3**. 50. 1807 = Aspidium plantagineum.
latifrons Witte; Salom. Nom. 304. 1883 (nomen) — Java.
(G) **latipes** Langsd. et Fisch. Ic. Fil. 10 t. 10. 1810. Mett. n. 116 — Amer. trop. Marginaria Pr. 1836; Goniophlebium J. Sm. 1846; Polypodium loriceum β HB. 343. P. harpeodes Link 1833; Marginaria Pr. 1836; Goniophlebium J. Sm. 1846 — Polypodium vacillans Link 1833; Goniophlebium Fée 1850-52 — G. pectinans Fée 1850-52 — G. grammatoides Fée 1869.
(C) **latum** (Moore) Sod. Cr. vasc. quit. 371. 1893 — Amer. trop. Campyloneurum Moore Ind. 225. 1861.
laxifolium J. Sm. Hist. Fil. 89. 1875 (err. typ.) = P. taxifolium.
laxifrons Liebm. Vid. Selsk. Skr.V. **1**. 204 (seors. 52). 1899 = P. suspensum.
(P) **laxum** Pr. Rel. Haenk. **1**. 23 t. 4 f. 1. 1825 — Peru. Brasilia. P. lanigerum Mett. n. 49. 1857 (non Desv. 1811).
Lebeufii Bak. Ann. Bot. **5**. 461. 1891 = Dryopteris.
Lecanopteris Mett. Ann. Lugd. Bat. **2**. 224. 1866 = Lecanopteris carnosa.
Lechleri Bak. Ann. Bot. **5**. 456. 1891 = Dryopteris laevigata.
legionarium Bak. Syn. 337. 1867 = P. macrodon.
(P) **Lehmanni** Mett. Pol. 117. n. 229 t. 3 f. 35. 1857. HB. 369. Chr. 114 — Birma. Sikkim. — Pleopeltis Bedd. 1868 — ? P. moulmeinensis Bedd. 1867.
Lehmannianum Hier. Engl. Jahrb. **34**. 513. 1904 = P. pastoense.
leiopteris Kze. Linn. **23**. 279, 319. 1850 = P. excavatum.
(My) **leiorhizum** Wall. List n. 303. 1828 (nomen); Mett. Fil. Lips. 37 t. 25 f. 7. 1856. Pol. n. 194. HB. 369. Chr. 115. NPfl. 321 — Ind. bor. et austr. China occ. — Phymatodes Pr. 1836 (nomen); Drynaria J. Sm. 1841 (nomen); Pleopeltis Moore 1862. Polypodium cuspidatum Don, Prod. Fl. Nepal. 2. 1825 (non Pr. 1825); Phymatodes J. Sm. 1857. ? Polypodium lucidum Roxb. 1844 (nom. opt.?).
(Pl) **Lenormandi** Bak. Syn. 514. 1874 — N. Caledonia.
lepidopodum Link, Hort. Berol. **2**. 94. 1833 = P. diversifolium.
(M) **lepidopteris** (Langsd. et Fisch.) Kze. Linn. **13**. 132. 1836. Mett. n. 103. HB. 346; Kuhn, Abh. Nat. Ges. Halle **11**. 23. Chr. 91 — Amer. trop. Acrostichum Langsd. et Fisch. Ic. Fil. 5 t. 2. 1810; Goniophlebium

Moore 1857; Lepicystis J. Sm. 1875. NPfl 323. Polypodium hirsutissimum Raddi 1819; Goniophlebium Brack. 1854 — Polypodium crassimargo Kze.; Mett. n. 103 b. 1857; Goniophlebium Moore 1862 — Polypodium rufulum Pr. 1822; Marginaria Pr. 1836. Polypodium lanosum Fée 1850-52 — P. sepultum Klf. 1824; Goniophlebium J. Sm. 1840; Lepicystis [J. Sm. 1842.

lepidopteris Sod. Cr. vasc. quit. 361. 1893 = P. baloonense.

lepidotum Willd; Schlecht. Adumbr. 17 t. 8. 1825 = P. lanceolatum.

lepigerum Schrad. Gött. Gel. Anz. **1824**. 868 = Dryopteris deflexa.

(Pl) **leptochiloides** Kuhn, Forschreise Gazelle **4** Farne 10. 1889 — N. Hannover.

leptophyllum L. sp. **2**. 1092. 1753 = Anogramma.

leptophyllum Moritz; Eat. Mem. Amer. Ac. n. s. **8**. 198. 1860 (nomen); Mett. Ann. sc. nat. V. **2**. 250. 1864 = P. apiculatum var.

leptophyllum Bak Ann. Bot. **5**. 458. 1891 = Dryopteris gracilifrons.

(Gr) **leptopodon** Wright, Tr. Linn. Soc. II Bot. **6**. 83. 1901 — Guiana.

leptosorum Kze. Linn. **23**. 320. 1850 = Dryopteris oligocarpa.

(P) **leptostomum** Fée, 7 mém. 58 t. 21 f. 2. 1857 — Mexico.

Leschenaultianum Wall. List n. 323. 1828 = Alsophila.

Leseblii Mérat, Fl. Paris. ed. II. 276. 1821 = Athyrium filix femina.

(G) *Le Sourdianum* Fourn. Mex. pl. **1**. 82. 1872 — Mexico.

leucatomos Poir. Enc. **5**. 516. 1804 = P. aureum var.

(P) **leucolepis** Gilbert, Bull. Torr. Cl. **24**. 260. 1897 — Jamaica.

(C) **leuconeuron** (Fée) Christ in Schwacke, Pl. Nov. Mineiras **2**. 22. 1900; Bull. Boiss. II. **2**. 374. 1902 — Brasilia. — Campyloneuron Fée, Cr. vasc. Br. **1**. 103 t. 35 f. 1. 1869 — C. fallax Fée 1869.

leuconeurum Diels, Engl. Jahrb. **29**. 203. 1900 = P. Dielseanum.

(Pl) **leucophorum** Bak. Journ. Linn. Soc. **22**. 229. 1886 — Borneo.

leucorhizon Kl. Linn. **20**. 400. 1847 = P. angustifolium var. amphostenon.

(P) **leucosorum** Boj. Hort. Maur. 417. 1837 (nomen); Hk. Ic. pl. t. 942, 943. 1854. HB. 324. Chr. 83. NPfl. 309 — Ins. Mascar. — (Conf. P. Cordemoyi).

(M?) **leucosporum** Kl. Linn. **20**. 404. 1847. Mett. n. 157. HB. 369 — Columbia. Pleopeltis Moore 1857; Lepicystis Diels 1897. NPfl. 324.

(P) **leucosticton** Kze.; Kl. Linn. **20**. 380. 1847; Hier. Engl. Jahrb. **34**. 521. 1904 — Columbia. — P. Karwinskyanum pt. Mett. n. 95. 1857.

leucosticton Fée, Gen. 240. 1850-52. 1857 = P. longiusculum.

Leuzeanum Gaud. Freyc. Voy. Bot. 361 t. 6. 1827 = Aspidium.

leveilleanum Desv. Prod. 232. 1827 = P. cultratum var. elasticum.

(Pl) **Lewisii** Bak. JoB **1875**. 201. Chr. 102 — China.

(P) **Leysii** Bak. JoB. **1879**. 66 — Sulu Archipel.

(P) **L'Herminieri** Fée, Gen. 238. 1850-52; 11 mém. 46 t. 12 f. 1; Lindm. Ark. f. Bot. **1**. 231 t. 11 f. 5 — Ind. occ. Venezuela.

(P) **Liebmanni** C Chr. Ind. 539. 1906 — Mexico. Guatemala.

P. griseum Liebm. Vid. Selsk. Skr. V. **1**. 198 (seors. 46). 1849. HB. 330. (non Schkuhr 1806).

lignosum Burm. Fl. Ind. Fl. Cap. Prod. 28. 1768 — Africa austr.

(P) **ligulatum** Bak. Syn. 320. 1867; Hier. Hedw. **44**. 78. 1905 — Fiji. Samoa.

ligustifolium Poir. Enc. **5**. 553. 1804 = Polystichum denticulatum.

ligustrifolium Desv. Prod. 225. 1827 = P. lycopodioides.

limbospermum Bellardi, Mém. Ac. Turin **5**. 253 (47). 1792 = Dryopteris oreopteris.

(Pl) **Lindbergii** Mett.; Kuhn, Linn. **36**. 136. 1869. HB. 358. NPfl. 316 — Brasilia. — Drynaria Fée 1872-73.

(P) **Lindenianum** Kze. Farnkr. **2**. 83 t. 134. 1851. Mett. n. 92 — Amer. centr. P. cancellatum Fée 1850-52.

(C) **Lindigii** Mett. Ann. sc. nat. V. **2**. 257. 1864. (HB. 348) — Columbia.

Lindleyanum Wall. List n. 304. 1828 = P. palmatum var. angustatum.

lineare Burm. Fl. Ind. 235 t. 67 f. 2. 1768 = Gleichenia.

lineare Houtt. Nat. Hist. **14** —. 1783 = P. ramosum.

(Pl) **lineare** Thbg. Fl. Jap. 335. 1784. HB. 354 (excl. β). Chr. 102. NPfl. 315 — Japonia. China. India bor. Ceylon. Malacca. Africa austr. c. ins. Ins. Hawaii. Pleopeltis Moore 1862; Niphobolus Keys. 1873; Pleopeltis Thunbergiana Klf. 1827 (nom. optim.). Pleopeltis nuda Hk. 1825; Polypodium Kze. 1850; Drynaria Fée 1850-52; Phymatodes J. Sm. 1857; Lepisorus J. Sm. 1857) — Pleopeltis elongata Klf. 1824; Phymatodes Pr. 1836; Polypodium Goldm. 1843; Drynaria Brack. 1854 (Hawaii) — Polypodium Wightianum Wall. 1829; Pleopeltis Bedd. 1863. Drynaria subspathulata Hk. 1857; Pleopeltis Moore 1862 — Polypodium leiopteris Kze. 1850 — P. atropunctatum Gaud. 1827. Mett. n 160 — P. Gueintzii Mett. n. 161. 1857 — Pleopeltis ussuriensis Regel et Maack 1861; Polypodium Regel 1881 — P. Alberti Regel 1881 — P. Onoei Franch. et Sav. 1879. Chr. 102. — P. oligolepidum Bak. 1880.

linearifolium Hk. sp. **5**. 53. 1863 = Cyclophorus.

lineatum Colebr.; Wall. List n. 300. 1828 = Dryopteris penangiana.

lingua Sw. Syn. 29. 1806 = Cyclophorus.

lingua Vahl, Ecl. **3**. 50. 1807 = P. Ryani.

(Pl) **linguaeforme** Mett. Ann. Lugd. Bat. **2**. 228. 1866. HB. 358. NPfl. 315 — Malesia orient. Melanesia.

lingulatum Sw. Syn. 30. 1806 = P. punctatum.

Linnei Bory, Ann. sc. nat. **5**. 464 t. 12. 1825 (Linnaei auctt.) = Drynaria sparsisora.

litigiosum Liebm. Vid. Selsk. Skr. V. **1**. 205 (seors. 53). 1849 = Dryopteris oligocarpa.

(G) **liukiuense** Christ, Bull. Boiss. II. **1**. 1014. 1901 — Ins. Liu-Kiu.

(P) **lividum** Mett. Ann. Lugd. Bat. **2**. 221. 1866. (HB. 328) — Java.

lobatum Huds Fl. angl. 459. 1762 = Polystichum.

lobatum Rich.; Willd. sp. **5**. 164. 1810 — Bahama. Sagenia Pr. 1836. (= Aspidium cicutarium?).

Lobbianum Hk. JoB. **5**. 309 t. 11. 1853 = Dipteris.

Lobbianum Hk. sp. **4**. 226 t. 278 B. 1862 = P. pediculatum.

(Gr) **locellatum** Bak. JoB. **1890**. 108. NPfl. 308 — N. Guinea.

(Gr) **Loherianum** Christ, Bull. Boiss. **6**. 197 t. 3 f. 2. 1898. NPfl. 308 — Luzon.

(P) **lomariiforme** Kze. Linn. **9**. 42. 1834. Mett. n. 76 — Venezuela. Peru.

(My) **lomarioides** (J. Sm.) Kze.; Mett. Pol. 102. n. 192 t. 2 f. 18-19. 1857. HB. 365. Chr. 113. NPfl. 321 — Malesia. Ins. Philipp. Formosa. Drynaria J. Sm. 1841 (nomen); Pleopeltis Moore 1857.

Lonchitis L. sp. **2**. 1088. 1753 = Polystichum.

(P) **longepilosum** C. Chr. Bot. Tidsskr. **25**. 78. 1902 — Brasilia. — P. villosum Fée, Cr. vasc. Br. **2**. 54 t. 97 f. 1. 1873 73 (non L. 1753 nec. Karst. 1869).

(P) **longepinnulatum** Fourn. Mex. pl. **1**. 77. 1872. HB. 510 — Mexico.

longicaudatum Liebm. Vid. Selsk. Skr. V. **1**. 209 (seors. 57) 1849 — Mexico.

(G) **longicaule** (Fée) C. Chr. Ind. 326. 1905; 541. 1906 — Columbia.
Goniophlebium Fée, 8 mém. 95. 1857.
longifolium Cav. Descr. 245. 1802 — Peru. — (prob. Cyclophorus sp.).
longifolium Pr. Del. Prag. **1**. 167. 1822 = P. brasiliense.
longifolium Roxb. Calc. Journ. **4**. 492. 1844 — Amboyna. — (Dryopteris sp.).
longifolium Mett. Pol. 87. n. 153 t. 1 f. 46. 1857 = P. revolutum.
longifrons Wall. List n. 274. 1828 = P. normale var.
longipes Wall. List n. 316. 1828 = Dryopteris brunnea.
longipes Link; Kze. Linn. **23**. 280. 1850 = P. phymatodes.
longipes Fée, Cr. vasc. Br. **2**. 53 t. 95 f. 3. 1872-73 = P. transiens.
(P) **longisetosum** Hk. sp. **4**. 225 t. 278 A. 1862. HB. 338. NPfl. 311 — Ecuador.
(S) **longisorum** (Bak.) C. Chr. Ind. 541. 1906 — Tonkin.
Gymnogramme Bak. JoB. **1890**. 267.
(Pl) **longissimum** Bl. Enum. 127. 1828; Fl. Jav. Fil. 159 t. 68. Mett. n. 191. HB. 366. Chr. 112 — Asia trop. Polynesia.
Phymatodes J. Sm. 1857; Pleopeltis Moore 1857; Colysis J. Sm. 1875 — (an Polypodium alternifolium Willd. 1810?).
longissimum Hk. sp. **5**. 7. 1863 = Dryopteris.
(P) **longiusculum** C. Chr. Ind. 188. 1905; 541. 1906 — Ecuador.
P. leucosticton Fée, Gen. 240. 1850-52; 7 mém. 58 t. 21 f. 3 (non Kl. 1847); Ctenopteris J. Sm. 1875.
(P) **longum** C. Chr. Ind. 541. 1906 — Ecuador. — P. alternifolium Hk. sp. **4**. 222 t. 277 A. 1862. HB. 335. Chr. 81 (non Willd. 1810).
(C) **Lorentzii** Hier. Engl. Jahrb. **22**. 406. 1896 — Argentina.
loreum Klf.; (Kze. Flora **1839**[1]. Beibl. 31. syn.) = P. angustifolium.
(G) **loriceum** L. sp. **2**. 1086. 1753. HB. 343 (pt). Chr. 92. NPfl. 312 — Amer. trop. — Marginaria Pr 1836: Goniophlebium J. Sm. 1840 — Polypodium laetum Raddi 1819; Marginaria Pr. 1836; Goniophlebium J. Sm. 1854 — G. pectinatum J. Sm. 1854.
loriforme Wall. List n. 271. 1828; Mett. Pol. 92. n. 164. 1857 = P. excavatum var.
loriforme Hk. Gard. Ferns t. 14. 1862 = P. lineare.
(G) **Lowei** C. Chr. Ind. 326. 1905; 541. 1906 — Guatemala.
Phlebodium inaequale Moore, Gard. Chr. **1855**. 660; Goniophlebium J. Sm. 1857; Polypodium Lowe 1858 (non Link 1833 nec Fée 1866); P. guatemalense Hk. 1863. HB. 345 (non Kl. 1855)
(L) **loxogramme** Mett. Pol. 112 n. 216 t. 3 f. 25. 1857. Chr. 107. NPfl. 316 — Japonia. Asia trop. Polynesia. Ins. Mascar. Natal. Guinea. Abyssinia.
Grammitis lanceolata Sw. Schrad. Journ. **1800**[2]. 18. 1801; Antrophyum Bl. 1828; Loxogramme Pr. 1836; Selliguea Fée 1850-52; Gymnogramme Hk. 1864. HB. 387 (non Polypodium L. 1753) — Grammitis coriacea Klf.; Spr. 1827; Loxogramme Pr. 1836; Selliguea Fée 1850-52; Polypodium Mett. n. 217. 1857 (non Raddi 1819) — Gymnogramme abyssinica Bak. 1874. HB. 517.
loxoscaphoides Bak. JoB. **1890**. 107 = Dryopteris.
lucens Schrad. Gött. gel. Anz. **1824**. 868 = P. brasiliense.
lucidum Rich. Act. Soc. Hist. nat. Paris **1**. 114. 1792 (nomen) — Cayenne.
lucidum Roxb. Calc. Journ. **4**. 486. 1844 = ? P. leiorhizum.
lucidum Beyrich; Hk. sp. **5**. 41. 1863 = ? P. angustifolium*.
(Gr) **ludens** Bak. Ann. Bot. **8**. 129. 1894 — Java.
(Pl) **Ludovicianum** Bak. Ann. Bot. **8**. 131. 1894 — Ins. Louisiade.

Lunanianum Hew. Mag. Nat. Hist. II. **2**. 460. 1838 = Dryopteris serrulata.
lunulatum Forst. Prod. 83. 1786 = Alsophila extensa.
lusitanicum L. sp. **2**. 1094. 1753 = Davallia canariensis.
luxurians Kze. Linn. **23**. 280. 1850 = Dryopteris prolifera.
luzonense Pr. Rel. Haenk. **1**. 26. 1825 = Goniopteris.
lyciaefolium Bory, Dup. Voy. Bot. **1**. 260. 1828 = P. accedens.
(Pl) **lycopodioides** L. sp. **2**. 1082. 1753. Mett. n. 174. HB. 357. Chr. 100. NPfl. 316 — Amer. trop. Africa trop. Ins. Mascar. ? Java. ? Ins. Hawaii.
Pleopeltis Pr. 1836; Phlebodium J. Sm. 1846; Drynaria Fée 1850-52; Anapeltis J. Sm. 1857; Niphobolus Keys. 1873; Phymatodes Millsp. 1903 — Polypodium lagopodioides Desv. 1811; Craspedaria Fée 1850-52 — Polypodium owariense Desv. 1811; Goniophlebium Lodd. 1849; Anapeltis J. Sm. 1866 — Polypodium myrtillifolium Klf. 1824; Goniophlebium Moore 1857 — Drynaria acuminata Fée 1869 (non Brack. 1854) — Craspedaria grandis Fée 1869.
—* Polypodium surinamense Jacq. 1789; Pleopeltis Pr. 1836; Craspedaria Fée 1850-52 — Polypodium salicifolium Willd. 1810. Mett. n. 172 (non Vahl 1807); Pleopeltis Pr. 1836; Phlebodium J. Sm. 1841; Polypodium salicinum Wikstr. 1826 — P. dictyophyllum Kze.; Mett. n. 173. 1857. (an potius species propria?).
lycopodioides Schkuhr, Kr. Gew. **1**. 187 t. 8 C. 1809 = P. persicariifolium.
(M) **Macbridense** Shimek, Bull. Lab. Nat. Hist. Univ. Iowa **4**. 199 t. 20 f. 6-9. 1897 — Nicaragua.
(P) **Macgregori** Bak. Ann. Bot. **8**. 130. 1894 — Ins. Louisiade.
(Pl) **Mackenii** Bak. Syn. 357. 1868 — Natal.
macrocarpum Bory; Willd. sp. **5**. 147. 1810 = P. lanceolatum.
macrocarpum Pr. Rel. Haenk. **1**. 23 t. 1 f. 4. 1825 = P. pycnocarpum.
(Pl) **macrochasmum** Bak. JoB. **1880**. 216 — Sumatra. Java. Perak.
— P. laciniatum Bl. Enum. 131. 1828; Fl. Jav. Fil. 149 t. 63. Mett. n. 204. HB. 365 (non Burm. 1768 nec Lam. 1778); Phymatodes Pr. 1836; Phymatopsis J. Sm. 1875; Pleopeltis Bedd. 1892.
(P) **macrodon** Hk. sp. **4**. 218. 1862 — Guatemala.
P. legionarium Bak. Syn. 337. 1867.
macrodon Reinw.; Bak. Syn. 318. 1867 = Aspidium.
(S) **macrophyllum** (Bl.) Reinw.; Mett. Fil. Lips. 37 t. 25 f. 22-23. 1856. Pol. n. 213. Chr. 108. NPfl. 316 — Malesia. N. Guinea. China austr.
Grammitis Bl. Enum. 119. 1828; Selliguea Bl. Fl. Jav. Fil. 127 t. 53. 1828; Colysis Pr. 1849; Gymnogramme Hk. 1864. HB. 388.
macrophyllum Hk. sp. **4**. 241. 1862 = Dryopteris pellucido-punctata.
macropodum Bak. Journ. Linn. Soc. **15**. 108. 1876 = Cyclophorus.
macropterum Klf. Enum. 111. 1824 = Dryopteris grandis*.
(P) **macrorhynchum** Bak. JoB. **1880**. 370 — Madagascar.
macrosorum Fée, Gen. 241. 1850-52; 6 mém. 11 t. 8 f. 1. 1853 = P. monosorum.
macrosorum Bak. JoB. **1885**. 106 = megasorum.
(Pl) **macrosphaerum** Bak. Kew Bull. **1895**. 55 — Yunnan.
(Pl) **macrourum** Bak. Gard. Chr. n. s. **25**. 136. 1886 — ? Queensland.
(P) **macrum** Copeland in Perkins, Fragm. 189. 1905 — Ins. Philipp. (Mindanao).
maculatum Spr. Nova Acta **10**. 228. 1821 = Alsophila quadripinnata.
(Pl) **maculosum** Christ, Bull. Boiss. **6**. 872. 1896 — Yunnan.
madagascariense Desv. Prod. 227. 1827 = P. phymatodes.

Madrense J. Sm. Bot. Voy. Herald 338 t. 73. 1854 = P. plebejum.
maenurum Link, Hort. Berol. **2**. 96. 1833 = P. recurvatum.
magnificum Hier. Engl. Jahrb. **34**. 535. 1904 = P. Fendleri.
(S) **Maingayi** (Bak.) Diels, Nat. Pfl. 1^4. 318. 1899 — Malacca. Formosa.
Gymnogramme Bak. Syn 517. 1874; Selliguea Bedd. 1876.
(L) **Makinoi** C. Chr. Ind. 339. 1905; 543. 1906 — Japan.
Gymnogramme salicifolia Mak. Phan. et Pter. Jap. Ic. t. 34. 1899 (non Polypodium Vahl 1807 nec Willd. 1810); Loxogramme Mak. 1905.
(P) **malaccanum** Bak. Ann. Bot. **8**. 129. 1894 — Malacca.
(Pl) **malacodon** Hk. sp. **5**. 87. 1863. HB. 363. NPfl. 318 — Himalaya. China merid
Pleopeltis Bedd. 1883 (non 1876).
Manabyanum Bak. JoB. **1877**. 165 = P. patentissimum.
manipurense Bedd. JoB. **1888**. 235 = Dryopteris.
(G) **manmeiense** Christ, Bull. Boiss. **6**. 870. 1898 — Thibet orient. China.
Mannianum Hk. sp. **4**. 253. 1862 = Dryopteris.
Marantae Hoffm. Deutschl. Fl. **2**. 5. 1759 = Woodsia ilvensis.
(P) **margaritiferum** Christ, Bull. Boiss. II. **5**. 2 c. fig. 1905 — Costa Rica.
marginale L. sp. **2**. 1091. 1753 = Dryopteris.
marginale Thbg. Fl. Jap. 337. 1784 = Microlepia marginata.
marginale Bory; Willd. sp. **5**. 149. 1810 = P. lanceolatum.
marginale Wall. List n. 322. 1828 = Dryopteris pulvinulifera, punctata,
marginatum Houtt. Pfl. Syst. 13^1. 199. 1786 = Microlepia. [scabrosa.
marginatum Moore, Ind. 297. 1861 = ? P. marginale Wall.
(Pl) **marginatum** Bak. Syn. 513. 1874 — Ins. Pinorum. N. Caledonia.
(Gr) **marginellum** Sw. Prod. 130. 1788. Mett. n. 10. HB. 321. Chr. 77. NPfl. 308 — Amer. trop. Ins. Caboverd. St. Helena. Samoa.
Grammitis Sw. 1801; Mecosorus Kl. 1847 — Grammitis limbata Fée 1850-52 — G. quaerenda Bolle 1863 — G. fluminensis Fée 1869.
(G) **maritimum** Hier. Engl. Jahrb. **34**. 527. 1904 — Columbia.
(P) **Martensii** Mett. Pol. 61 n. 83. 1857. HB. 333 — Mexico.
P. affine Mart. et Gal. Mém. Ac. Brux. **15**. 37 t. 8 f. 1. 1842 (non Bl. 1828). P. callolepis Fée 1857.
(M) **masafuerae** Phil. Linn. **29**. 107. 1857 — Juan Fernandez.
mascarense Bak. Syn. ed. I. 455. 1868 = Dryopteris.
(Pl) **Mathewii** Tutcher, Journ. Linn. Soc. **37**. 68. 1905 — China (Wei-hai-wei).
(G) **Mathewsii** Kze.; Mett. Pol. 74 n. 112. 1857. HB. 341. NPfl. 311 — Columbia-Peru. — Goniophlebium Villeminianum Fée, 7 mém. 63 t. 27 f. 3. 1857.
mauritianum Willd. sp. **5**. 150. 1810 = P. lycopodioides.
Maximowiczii Bak. Syn. 504. 1874 = Polystichum.
(Pl) **maximum** (Brack.) Hk. sp. **5**. 83. 1863. (HB. 364) — Tahiti.
Drynaria Brack. Expl. Exp. **16**. 51 t. 7. 1854; Pleopeltis Moore 1857.
(P) **Maxonii** C. Chr. Ind. 543. 1906 — Mexico. — P. firmulum Maxon, Contr. U. S. Nat. Herb. **8**. 275 t. 61 f. 4. 1903 (non Bak. 1893).
(Gr) **Maxwellii** Bak. Kew Bull. **1893**. 211 — Borneo.
Mayi ht. May, Gard. Chr. III. **36**. 44. 1904 = P. aureum var. hort. (vide
Mazei Bak. Ann. Bot. **5**. 457. 1891 = Dryopteris. [Phlebodium).
(Gr) **mediale** Bak. Syn. 507. 1874 — China merid. Ind. austr. Ceylon. Java. Celebes. — Grammitis attenuata Kze. Linn. **24**. 251. 1851 (non Polypodium H.B.Willd. 1810); Trichocalymma parasiticum Zenker msc. Polypodium

Mett. n. 16. 1857. HB. 319. NPfl. 308 (non L. 1753) — (an Grammitis pilosiuscula Bl. 1828?) — (an adhuc P. Christii Copeland?).
medullare Forst. Pl. escul. 74. 1786 — Cyathea.
megacuspe Bak. JoB. **1890**. 266 = Dryopteris.
megalocarpum Hk. sp. **5**. 102. 1864 = Aspidium.
megalodus Schkuhr, Kr. Gew. **1**. 24 t. 19 b. 1806 = Dryopteris.
(Pl) **megalophyllum** Desv. Prod. 227. 1827. NPfl. 316 — Guiana. Amazonas.
Phymatatodes Schomburgkianum J. Sm. 1842; Polypodium Kze. 1842. Mett. n. 177. HB. 355. Chr. 104; Drynaria Fée 1850-52; Pleopeltis Moore 1862; Mecosorus Schomburgkii Kl. 1847; Polypodium Hk. 1863.
(Pl) **megasorum** C. Chr. Ind. 544. 1906 — Formosa.
P. macrosorum Bak. JoB. **1885**. 106. NPfl. 316 (non Fée 1850-52).
(Gr) **melanoloma** (Boiv.) Cordem. Bull. Soc. sci. et arts Réun. **1890-91**. —. 1891; Fl. Réun. 83. 1895. Chr. 77 — Réunion. — Grammitis Boiv. msc.
melanopus R.Br.; Wall. List pag. 83 (n. 293). 1829 (nomen) = P. ebenipes.
(P) **melanopus** Grev. et Hk. Bot. Misc. **3**. 384 t. 111. 1833. HB. 329 — Costa Rica-Ecuador.
(P) **melanostictum** Kze. Linn. **9**. 44. 1834. Mett. n. 30 — Cuba. Columbia. Venezuela.
(P) **melanotrichum** Bak. Tr. Linn. Soc. II. Bot. **2**. 292. 1894 — Guiana.
(Pl) **Melleri** Bak. Syn. 359. 1868 — Madagascar.
(Pl) **membranaceum** Don, Prod. Fl. Nepal. 2. 1825. Mett. n. 232. HB 360. Chr. 105 — India bor. et austr. China austr. Ceylon. Celebes. Ins. Philipp.
Colysis J. Sm. 1857 (non Pr. 1849); Pleopeltis Moore 1860; Drynaria undulata J. Sm. 1841; ? Bathmium ? Fée 1850-52 — Polypodium grandifolium Wall. 1828; Phymatodes Pr. 1836; Pleopeltis Moore 1857.
membranaceum Hk. ap. Blakiston, Yang-tsze 365, 366. 1862 = Aspidium devexum.
membranifolium R.Br. Prod. Fl. N. Holl. 147. 1810 = ? P. pustulatum. (an P. nigrescens?).
(G) **mengtzeense** Christ, Bull. Boiss. **6**. 869. 1898. — Yunnan. Khasia.
P. aspersum Bak. Kew Bull. **1898**. 231 (non P. adspersum Schrad. 1818 nec Bl. 1828).
menisciicarpon Hk. sp. **5**. 86. 1863 = Aspidium irriguum et A. siifolium.
menisciifolium Langsd. et Fisch. Ic. Fil. 11 t. 12. 1810 = P. brasiliense var.
meniscioides Liebm. Vid. Selsk. Skr. V. **1**. 211 (seors. 59). 1849 = Dryopteris.
(P) **meridense** Kl. Linn. **20**. 380. 1847. HB. 333 (pt.) — Columbia. Brasilia.
P. radicale Bak. 1867 (non Moritz; Eat. 1860) — P. Spixianum Mart.;
meridionale Poir. Enc. **5**. 553. 1804 — Guadeloupe. [Mett. n. 71. 1857.
Lastrea Pr. 1836 — (Dryopteris sp.).
(Gr) **Merillii** Copeland in Perkins, Fragm. 188. 1905 — Ins. Philipp.
Metcalfei Bak. Ann. Bot. **5**. 461. 1891 = Dryopteris.
Mettenianum Ces. Att. Ac. Napoli 7^{8}. 27 (nota). 1876 = P. heterocarpum.
Mettenianum Ces. Rend. Ac. Napoli **16**. 27. 1877 (err.) = P. Meyenianum.
(S) **mexicanum** (Fée) Salom. Nom. 308. 1883 — Mexico. Guatemala. — Selliguea Fée, Gen. 177. 1850-52; 7 mém. 43 t. 10 f. 4. 1857; Gymnogramme Bak. 1868. HB. 387. Grammitis Salvinii Hk. 1861; Gymnogramme Hk. 1864.
(Pl) **Meyenianum** (Schott) Hk. sp. **5**. 94. 1863. HB. 366. Chr. 117. NPfl. 319 — Luzon. — Aglaomorpha Schott, Gen. Fil. t. 19. 1834; Psygmium elegans Pr. 1836; Drynaria Proustiana Gaud. 1846.

Michaelis Bak. JoB. 1877. 164 = Dryopteris.
microcarpum Willem. in Usteri Ann. 18. 63. 1796 — Mauritius.
Aspidium Kuhn 1868. — (Dryopteris sp.).
(P) **microchasmum** Bak. JoB. 1887. 44 — Jamaica.
microdontum Desv. Berl. Mag. 5. 319. 1811 = Alsophila.
(Pl) *microlepidum* Desv. Prod. 228. 1827 — Peru.
microlepis Fée, Gen. 238. 1850-52; 6 mém. 8 t. 6 f. 2 = P. polypodioides.
(P) **microphyllinum** Mett.; Bak. Syn. 339. 1867 — Peru.
(Gr) **microphyllum** Bak. Kew Bull. 1897. 299 — Madagascar.
(P) **micropteris** C. Chr. Ind. 545. 1906 — Amer. trop.
Xiphopteris setosa Klf. Enum. 275. 1824; Grammitis Pr. 1836 (non Bl. 1828); Polypodium Mett. n. 6. 1857; Hier. Hedw. 44. 91. 1905 (non Thbg. 1784 nec. Forst. 1786).
(G) **microrhizoma** Clarke; Bak. Syn. 511. 1874 — Himalaya. Yunnan.
Goniophlebium Bedd. 1876.
microsorum Mett. Cat. hort. Herrenb. 1855 = P. musifolium.
microsorum Lindm. Ark. för Bot. 1. 239 t. 11 f. 2. 1903 = P. pectinatiforme.
(P) **millefolium** Bl. Enum. 134. 1828; Fl. Jav. Fil. 190 t. 78 A. Mett. n. 63. HB. 339. NPfl. 311 — Java.
(Pl) **millisorum** Bak. Journ. Linn. Soc. 15. 109. 1876 — Kei Ins.
Milnei Hk. sp. 4. 254. 1862 = Aspidium viridans.
(G) **mindense** Sod. Cr. vasc. quit. 348. 1893 — Ecuador.
minimum Aublet, Hist. pl. Guiane 2. 963. 1775 — Guiana.
minimum Brack. Expl. Exp. 16. 5 t. 1 f. 3. 1854 = ? P. serrulatum var.
minimum Moore, Ind. 14. 1857 = Acrostichum serratum Lam.
minimum Bak. JoB. 1879. 41 = P. vittariifolium.
minus Fée, 1 mém. 14 t. 1 f. 12. 1844 (nomen) — ?
(P) **minutum** Bl. Enum. 130. 1828; Fl. Jav. Fil. 188 t. 87 D. Mett. n. 43. HB. 328. NPfl. 310 — Malesia. Ins. Philipp.
(My) **mirabile** C. Chr. Ind. 545. 1906 — Amboyna.
P. imbricatum Karst. Ann. Jard. Buit. 12. 168 t. 15 f. 29, 30, t. 16. f. 49, t. 19. 1895. Chr. 113. NPfl. 321 (non Liebm. 1849.)
miser Hew. Mag. Nat. Hist. II. 2. 460. 1838 = Dryopteris effusa vel Aspidium Christianae.
(P) **Mitchellae** Bak. in Hemsl. Biol. Centr. Amer. 3. 664. 1885 — Brit. Honduras.
(P) **mixtum** Sod. Cr. vasc. quit. 339. 1893 — Ecuador.
molestum Mett. Ann. sc. nat. V. 2. 254. 1864 = P. monosorum.
molle Schreb. Spic. Fl. Lips. 70. 1771 = Athyrium filix femina.
molle All. Fl. Pedem. 2. 287. 1785 = Athyrium alpestre.
molle Jacq. Coll. 3. 188. 1789 = Dryopteris parasitica.
molle HBK. Nov. Gen. et Sp. 1. 8. 1815 = P. pectinatum.
molle Roxb. in Beats. St. Helena 315. 1816 = Dryopteris Dianae.
molle Mett. Pol. 128. n. 261. 1857 = Cyclophorus.
(Gr) **Molleri** Bak. Bol. Soc. Brot. 4. 154 t. 2 f. B. 1886 — Ins. S. Thomé (Guinea).
(P) **mollicomum** Nees et Bl. Nova Acta 11. 121 t. 12 f. 2. 1823; Bl. Fl. Javae Fil. 184 t. 86 B. 1829. Chr. 81 — Java. Celebes.
Ctenopteris Kze. 1845. — (an adhuc P. fuscatum Bl. 1828. HB. 331?).
molliculum Kze.; Link. Fil. sp. 130. 1841 = Dryopteris.
molliculum Copeland, in Perkins, Fragm. 190. 1905 = P. tomentellum.
(Gr) **mollipilum** Bak. JoB. 1890. 107 — N. Guinea.

(P) **mollissimum** Fée, 8 mém. 47 t. 12 f. 2. 1866. (HB. 329) — Guadeloupe. Ecuador.
mollissimum Christ, Bull. Boiss. **7**. 5. 1899 = Cyclophorus malacophyllus.
molliusculum Wall. List n. 332. 1828 = Dryopteris.
(P) **moniliforme** Lagasca; Sw. Syn. 33. 1806. Mett. n. 29. HB. 325. Chr. 79. NPfl. 309 — Amer. trop. — Ctenopteris J. Sm. 1875 — Jamesonia adnata Kze. 1851 — Polypodium subcrenatum Hk. 1845 — P. angustissimum Fée 1869 — P. subdicarpon Fée 1869.
(P) **monosorum** Desv. Berl. Mag. 5. 319. 1811 — Columbia-Peru. P. onustum Hk. 1845. Mett. n. 97. HB. 339. Chr. 87; Lepicystis Diels 1899. NPfl. 323 — Polypodium macrosorum Fée 1850-52. P. sporadolepis γ [Mett. n. 96.
monosorum Pr. Tent. 181. 1836 = Dryopteris tristis var.
montanum Lam. Fl. franç. **1**. 23. 1778 = Cystopteris.
montanum Vogler, Dissert. 1781 = Dryopteris oreopteris.
montanum var. A Poir. Enc. **5**. 540. 1804 = Davallia canariensis.
montevidense Spr. Syst. **4**. 59. 1827 = Polystichum aculeatum var. 9.
monticola Kl. Linn. **20**. 377. 1847. Mett. n. 38 — Venezuela. Micropteris J. Sm. 1875.
montium Gmel. Syst. Nat. **2**[2]. 1314. 1791 = Cystopteris montana.
morbillosum Pr. Rel. Haenk. **1**. 22 t. 3 f. 3. 1825 = Drynaria quercifolia.
Morgani Zeiller, Bull. Soc. Fr. **32**. 77. 1885 = P. stenophyllum.
(Pl) **Moricandii** Mett. Pol. 87 n. 152. t. 1 f. 47-48. 1857 — Bahia. Drynaria Fée 1869.
(P) **Moritzianum** Link, Fil. sp. 126. 1841. HB. 333 — Venezuela.
Morlae Sod. Cr. vasc. quit. 641. 1893 = Dryopteris.
(Pl) **Moseleyi** Bak. Journ. Linn. Soc. **15**. 110. 1876. NPfl. 321 — Ins. Moluccae. [N. Guinea.
(G) **Mosenii** C. Chr. Bot. Tidsskr. **25**. 78. 1902 — Brasilia.
(G) **moupinense** Franch. Nouv. Arch. Mus. II. **10**. 121. 1887 — Tibet orient. Gonioplebium Bedd. 1892.
mucronatum Sw. Fl. Ind. occ. **3**. 1649. 1806 = Polystichum echinatum.
mucronatum Roxb. Calc. Journ. **4**. 490. 1844 — Sylhet. (= Polystichum auriculatum?).
multifidum Jacq. Coll. **3**. 187. 1789 = Dryopteris effusa.
multifidum Bory, Voy. **2**. 193 t. 20 f. 2. 3. 1804 = P. furcatum.
multifissum Goldm. Nova Acta **19**. Suppl. I. 453. 1843 = Athyrium scandicinum.
multiflorum Roth, Catal. bot. **1**. 135. 1797 = Dryopteris spinulosa *dilatata.
multiflorum Roxb. Calc. Journ. **4**. 493. 1844 — Amboyna.
multilineatum Wall.; Hk. sp. **5**. 11. 1863 = Dryopteris moulmeinensis.
(Pl) **Munchii** Christ, Bull. Boiss. II. **3**. 147. 1903 — Mexico.
munitum Klf.; Pr. Tent. 180. 1836 = Dryopteris rufescens.
muricatum L. sp. **2**. 1093. 1753 = Polystichum aculeatum var. 4.
muricatum Sw. Prod. 131. 1788 = Polystichum echinatum.
muricatum Powell; Bak. Syn. 506. 1874 = Dryopteris subspinosa.
(P) **murorum** Hk. Ic. pl. t. 70. 1837. HB. 339. Chr. 87 — Columbia-Ecuador. Lepicystis Diels 1899. NPfl. 323. Polypodium sporadolepis β Mett. n. 96.
(Pl) **musifolium** Bl. Enum. 134. 1828; Fl. Jav. Fil. 171 t. 79. Mett. n. 234. HB. 360. Chr. 118. NPfl. 315 — Malesia. N. Guinea. — Drynaria J. Sm. 1857; Pleopeltis Moore 1857 — Polypodium Schumannianum Diels 1901.
muscosum Vahl, Ecl. **3**. 54. 1807 = Dryopteris pubescens var.

muscosum Willd. **5**. 203. 1810 — Amer. trop.
(an P. muscosum Vahl? Dryopteris sp.?)
muscosum Christ in Schwacke, Pl. Nov. Mineiras **2**. 19. 1900; Bull. Boiss. II. **2**. 368. 1902 = P. Wittigianum.
(P) **musgravianum** Bak. JoB. **1890**. 108 — N. Guinea.
mutabile Kze. Linn. **9**. 46. 1834 = P. fraxinifolium.
(P) **myosuroides** Sw. Prod. 131. 1788. Mett. n. 5; Hier. Hedwigia **44**. 83. 1905 — Amer. trop. Afr. orient. — Grammitis Sw. 1801; Xiphopteris Klf. 1824.
myosuroides Jenm. Bull. Dep. Jam. **4**. 112. 1897 = P. delitescens.
myriocarpum Hk. Ic. pl. t. 84. 1837 = P. pellucidum var.
(Pl) **myriocarpum** (Pr.) Mett. Pol. 105 n. 198. t. 1 f. 3. 1857. HB. 358. NPfl. 315 — Cochinchina. Amboyna. Ins. Philipp.
Phymatodes Pr. Tent. 196 t. 8 f. 12. 1836. Pleopeltis Moore 1857. Drynaria longissima J. Sm. 1841; Microsorium Fée 1850-52.
myriolepis Christ, Bull. Boiss. **4**. 661. 1896 = P. Skinneri.
(P) **myriophyllum** Mett. Fil. Lechl. **1**. 6. 1856. HB. 338. NPfl. 311 — Peru.
P. tenuisectum Mett. l. c. t. 2 f. 1, 2. 1856 (non Bl. 1828).
myriotrichum Bak. JoB. **1885**. 217 = Dryopteris hirtula.
myrmecophilum Christ, Verh. Nat. Ges. Basel **11**. 24. 1894 = P. sarcopus.
myrrhidifolium Vill. Fl. Delph. 114. 1785 = Cystopteris montana.
myrtillifolium Klf. Enum. 89. 1824 = P. lycopodioides.
mysurense Heyne; Wall. List n. 269. 1828 = ? Cyclophorus porosus.
nanegalense Sod. Rec. 65. 1883 = P. loriceum.
(P) **nanum** Fée, Gen. 238. 1850-52; Hier. Hedwigia **44**. 102. 1905 — Guiana. [Brasilia.
nanum Sturm, Enum. pl. vasc. cr. chil. 13. 1858 = P. Billardieri.
nanum Vieill., Bak. Syn. 507. 1874 = P. pumilio.
nectariferum Becc.; Bak. Malesia **2**. 247 t. 65. 1886 = Drynaria.
(Pl) **neglectum** Bl. Enum. 121. 1828; Fl. Jav. Fil. 133 t. 54 f. 1. — Luzon. Bantam.
Microterus Pr. 1849; Niphobolus? Fée 1850-52; Pleopeltis Moore 1861. Craspedaria calva Fée 1850-52.
(Ph) **nematorhizon** Eat. Bot. Gaz. **3**. 89 (?). 1878; Jenm. JoB. **1887**. 100 — [Trinidad.
nemorale Salisb. Prod. 403. 1796 = Dryopteris filix mas.
nemorale Wall. List n. 317. 1828 = Dryopteris parasitica.
nemorale Brack. Expl. Exp. **16**. 16. 1854 = Dryopteris setigera.
nephrodioides Kl. Linn. **20**. 384. 1847 — Columbia.
neriifolium Schkuhr, Kr. Gew. **1**. 14 t. 15. 1806 = P. brasiliense.
nervosum Boj. Hort. Maur. 419. 1837 (nomen) = Dryopteris tomentosa.
nervosum Kl. Linn. **20**. 386. 1847 = Dryopteris.
(P) **nesioticum** Maxon, Smiths. Misc. Coll. **47**. 410 t. 57. 1905 — Jamaica.
(P) **Newtoni** Bak. Kew Bull. **1896**. 41 — Fernando Po.
nicotianaefolium Bak. Syn. ed. I. 455. 1868 = Aspidium.
(Pl) **nigrescens** Bl. Enum. 126. 1828; Fl. Jav. Fil. 101 t. 70. HB. 364. Chr. 111. NPfl. 318 — Ind. austr. Ceylon. Malesia-Polynesia-Austr. trop.
Phymatodes J. Sm. 1866; Pleopeltis Carr. 1873. — (an Polypodium alternifolium Willd. 1810? an P. membranifolium R.Br. 1810?).
nigrescentium Jenm. Gard. Chr. III. **17**. 100. 1895 = Dryopteris.
nigripes Hassk. Cat. pl. Bogor. 4. 1844; Zoll. et Moritz, Nat. Gen. Arch. Neerl. Ind. **1**. 397. 1844 — Java.
nigripes Hk. sp. **5**. 17. 1863 = P. plebejum var.
nigritianum Bak. Syn. 313. 1867 = Dryopteris.

nigrocarpum Bedd. Ferns S. Ind. t. 169. 1863-65 = Dryopteris scabrosa.
(Pl) **nigrocinctum** Christ, Bull. Boiss. 6. 874. 1896 — Yunnan.
(Gr) **nigrolimbatum** (Spruce) Jenm. Bull. Dep. Jam. 4. 69. 1897 — Jamaica.
Grammitis Spruce msc.
nigrovenium Christ, Verh. Nat. Ges. Basel 11. 443. 1896 = P. papillosum.
(P) **nimbatum** Jenm. JoB. 1886. 271. NPfl. 309 — Jamaica.
(Pl) **ningpoense** Bak. Ann. Bot. 5. 474. 1891 — China.
(G) **niponicum** Mett. Ann. Lugd. Bat. 2. 222. 1866. HB. 341. Chr. 93. NPfl. 311 — Japonia. China. Birma.
nitens Desv. Prod. 240. 1827 — Peru. — (Dryopteris sp.).
nitens Bak. Syn. ed. II. 353. 1874 = Cyclophorus splendens.
(C) **nitidissimum** Mett. Ann. sc. nat. V. 2. 258. 1864. (HB. 349) — Columbia.
nitidulum Kze. Linn. 23. 281. 1850 = P. angustifolium.
(C) **nitidum** Klf. Enum. 92. 1824. Mett. n. 138. (HB. 348) — Amer. trop.
Campyloneurum Pr. 1836; Cyrtophlebium J. Sm. 1841 — Campyloneurum polyanthum Pr. 1836.
nitidum Kl. Linn. 20. 398. 1847 = P. latum.
nitidum Hk. Fil. Exot. t. 12. 1857 = P. lucidum.
nivosum Fée, 8 mém. 89. 1857 = P. furfuraceum.
nodosum Kl. Linn. 20. 400. 1847 = P. angustifolium.
Pl) **normale** Don, Prod. Fl. Nepal. 1. 1825. Mett. n. 151. HB. 358. Chr. 104. NPfl. 315 — China. Asia trop. N. Guinea. Madagascar.
Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Pleopeltis Moore 1862; Colysis J. Sm. 1875 — Polypodium longifrons Wall. 1828; Hk. et Grev. Ic. Fil. t. 65; Drynaria J. Sm. 1841; Pleopeltis Moore 1862.
(Pl) **novae-zealandiae** Bak in Hk. Ic. pl. t. 1674. 1886 — N. Zealand.
P. sp. Cheesem. Tr. N. Zeal. Inst. 10. 356. 1878.
noveboracense L. sp. 2. 1091. 1753 = Dryopteris.
nudatum Roxb. Calc. Journ. 4. 491. 1844 — Amboyna.
nudiusculum Kze. Linn. 24. 253. 1851 = P. lineare et P. simplex.
nudum Forst. Prod. 82. 1786 = Microlepia speluncae.
nudum Kze. Linn. 23. 281. 1850 = P. lineare.
nudum Mett. Pol. 37 n. 18. 1857 = P. gramineum.
nummulariaefolium Mett. Pol. 123 n. 247 t. 3 f. 9, 10. 1857 = Cyclophorus.
nummularium Mett. Pol. 105 n. 199. 1857 = P. pyrolifolium.
(P) **nutans** Bl. Enum. 128. 1828; Fil. Jav. Fil. 182 t. 86 A — Malacca. Java. Sumatra. N. Caledonia. — Prosaptia Mett. 1870; Ctenopteris J. Sm. 1875.
nutans J. Sm. JoB. 3. 394. 1841; Mett. Pol. 41 n. 28. 1857 = P. decorum.
(P) **nutatum** Jenm. JoB. 1886. 272 — Jamaica.
nymphale Forst. Prod. 81. 1786 = Dryopteris parasitica.
(P) **oblanceolatum** Bak. Ann. Bot. 8. 128. 1894 — N. Guinea.
(P) **obliquatum** Bl. Enum. 128. 1828; Fl. Jav. Fil. 181 t. 85 B. Mett. n. 52. HB. 328. Chr. 85. NPfl. 310 — Ind. austr. Ceylon. Malesia. Ins. Philipp.
Grammitis Hassk. 1844; Prosaptia Mett. 1870; Cryptosorus J. Sm. 1875; C. Blumei Fée 1850-52. Ctenopteris rufescens Kze. 1846. (Coelopteris A.Br. msc. Cystidium J. Sm. msc. Leptostegia Zipp. msc. Thylacopteris Kze. msc.).
obliteratum Sw. Prod. 132. 1788 = Dryopteris.
obliteratum? Raddi. Opusc. sci. Bol. 3. 287. 1819 = Dryopteris vivipara.

(Pl) **oblongisorum** C. Chr. Ind. 549. 1906 — Yunnan.
P. subintegrum Bak. Kew Bull. **1898**. 231 (non 1877).
oblongo-dentatum Hoffm. in Roem. et Ust. Mag. Bot. **9**. 10. 1790 = Athyrium filix femina.
obovatum Mett. Pol. 124 n. 248. 1857 = Cyclophorus nummularifolius var.
obscurum Mett. Pol. 38 n. 23. 1857 = P. Reinwardtii var.
obscurum Hk. sp. **4**. 237. 1862 = Dryopteris sagenioides*.
obtusilobum Desv. Berl. Mag. **5**. 317. 1811 = Dryopteris.
(P) **obtusissimum** C. Chr. Ind. 549. 1906 — Luzon.
P. subobliquatum Christ, Bull. Boiss. **6**. 197. 1898 (non Hk. 1862).
obtusum Spr. Anleit. **3**. 92. 1804 = Woodsia.
obtusum Poir. Enc. **5**. 510. 1804 — S. Domingo.
obtusum Kuhn, Fil. Afr. 150. 1868 = ? P. furcatum.
occidentale Maxon, Fern Bull. **12**. 102. 1904 = P. glycyrrhiza.
(C) **occultum** Christ, Bull. Boiss. II. **5**. 7. 1905 — Costa Rica.
odoratum Poir. Enc. **5**. 541. 1804 = Dryopteris rigida.
odorum Vell. Fl. flum. **11** t. 78. 1827; Arch. Mus. Nac. Rio Jan. **5**. 450. 1881 = Dryopteris patens.
officinale Güldenst. Reise **1**. 421. 1787 = P. vulgare.
(P) **Okuboi** Yatabe, Bot. Mag. Tokyo **5**. 35 t. 21. 1891 — Japonia.
Oldhami Bak. Syn. 311. 1867 = Dryopteris.
(P) **oleandroides** Bak. Ann. Bot. **8**. 128. 1894 — N. Guinea.
oligocarpum H.B.Willd. sp. **5**. 201. 1810 = Dryopteris.
oligolepidum Bak. Gard. Chr. n. s. **14**. 494. 1880 = P. lineare var.
(Pl) **oligolepis** Bak. Kew Bull. **1898**. 231 — Yunnan.
oligophlebium Kze. Linn. **23**. 281, 320. 1850 = P. repens.
oligophlebium Bak. Syn. 506. 1874 = Dryopteris paucinervata.
oligophlebium Bak. JoB. **1891**. 6 = Dryopteris paucivenia.
oligosorum Kl. Linn. **20**. 387. 1847 = Dryopteris oligocarpa.
oligosorum Moritz, Bot. Zeit. **1854**. 855 (nomen); Mett.; Kuhn, Linn. **36**. 132. 1869 = P. flabelliforme var.
olivaceum Liebm. Vid. Selsk. Skr. V. **1**. 199 (seors. 47). 1849 = P. Ehrenbergianum.
omeiense Bak. JoB. **1888**. 229 = Dryopteris.
Onoei Franch. et Sav. Enum. pl. Jap. **2**. 246, 642. 1879 = P. lineare var.
ontolepis »Fée« (Hk. Bak. Syn. 330, err. script.) = P. oulolepis.
onustum Hk. Ic. pl. t. 749. 1845 = P. monosorum.
(Pl) **oodes** Kze. Bot. Zeit. **1846**. 421. Mett. n. 222. HB. 354 — Borneo. Luzon.
Drynaria Fée 1850-52; Pleopeltis Moore 1857; Phymatopsis J. Sm. 1875.
(P) **oosorum** Bak. Bol. Soc. Brot. **4**. 154 t. 2 f. A. 1886; Hier. Hedwigia **44**. 94. 1905 — Ins. S. Thomé. Kamerun.
(P) **opacum** Fée, Gen. 236. 1850-52 — Venezuela.
(C) **ophiocaulon** Kl. Linn. **20**. 401. 1847 — Peru. — Campylonevron Fée 1850-52.
oppositifolium Hk. sp. **5**. 8. 1863 = Dryopteris.
oppositum Vahl, Ecl. **3**. 53. 1807 = Dryopteris.
oppositum Wall. List n. 7080. 1829 = Dryopteris sparsa.
orbiculatum Poir. Enc. **5**. 525. 1804 = Stenochlaena pollicina var.
Oreopteris Ehrh.; Willd. Prod. 292. 1787 = Dryopteris.
(Gr) **organense** (Gardn.) Mett. Pol. 39. n. 39. 1857. HB. 324. NPfl. 309 — Brasilia.
Grammitis Gardn. in Hk. Ic. pl. t. 509. 1843; Fée, Cr. vasc. Br. **1**. 264 t. 78 f. 1. 1869.

orientale Gmel. Syst. Nat. 2^2. 1312. 1791 = Dryopteris.
ornatum Wall. List n. 327. 1828 = Dryopteris.
ornatum Kl. Linn. **20**. 396. 1847 = P. fraxinifolium.
(P) *Otites* L. sp. **2**. 1085. 1753. (Plum. t. 85) — Jamaica.
(an P. tenuifolium HBK. an P. pectinatum L.?).
Otites Willd. sp. **5**. 177. 1810; Mett. Pol. 59 n. 77. 1857 = P. pectinatum.
oulolepis Fée, 8 mém. 86. 1857 = P. plebejum.
ovalescens Fée, Cr. vasc. Br. **1**. 94 t. 27 f. 3. 1869 = P. ciliare.
ovato-crenatum Hoffm. in Roem. et Ust. Mag. Bot. **9**.[10]. 1790 = Athyrium [filix femina.
ovatum Burm. Fl. Ind. 233. 1768 — Java.
ovatum Wall.; Hk. et Grev. Ic. Fil. t. 41. 1827 = P. phyllomanes.
owariense Desv. Berl. Mag. **5**. 314. 1811 = P. lycopodioides.
(P) **oxylepis** C. Chr. Ind. 550. 1906 — Brasilia. — P. filipes Christ in Schwacke, Pl. Nov. Mineiras **2**. 20. 1900; Bull. Boiss. II. **2**. 370. 1902 (non Moore 1855).
oxylobum Wall. List n. 294. 1828 = P. hastatum var.
oxyodon Bak. JoB. **1879**. 66 = Dryopteris.
oxyphyllum Wall. List n. 324. 1828 = Athyrium drepanopterum.
oxyphyllum Kze. Bot. Zeit. **1848**. 116 = P. Zippelii.
oyamense Bak. JoB. **1877**. 366 = Dryopteris.
pachydermum Bak. Journ. Linn. Soc. **15**. 108. 1876 = Cyclophorus.
pachyphyllum Eat. Amer. Journ. Sci. II. **22**. 138. 1856 = P. Scouleri.
(Gr) **padangense** Bak. JoB. **1880**. 213 — Sumatra.
paleaceum Borckh. Röm. Arch. 1^3. 20. 1798 — Hort. — Dryopteris sp.
paleaceum Hk.fil. Tr. Linn. Soc. **20**. 166. 1847 = Dryopteris furcata var.
paleaceum Powell; Bak. Syn. 505. 1874 = Dryopteris samoensis.
pallens Bl. Fl. Jav. Fil. 178 t. 84 f. 1. 1829 = P. subauriculatum var.
pallidivenium Hk. sp. **5**. 8. 1863 = Dryopteris striata.
pallidum Brack. Expl. Exp. **16**. 18. 1854; ? Bail. Proc. Linn. N. S. Wales **5**. 31. 1881 = Dryopteris ornata.
palma Vell. Fl. flum. **11** t. 69. 1827; Arch. Mus. Nac. Rio Jan. **5**. 447. 1881 = P. brasiliense.
palmarum Goldm. Nova Acta **19**. Suppl. I. 454. 1843 = P. palmatum.
palmatopedatum Bak. Kew Bull. **1898**. 232 = Neocheiropteris.
palmatum Vell. Fl. flum. **11** t. 63. 1827 (sine descr.); Arch. Mus. Nac. Rio Jan. **5**. 446. 1881 = P. angustum.
(Pl) **palmatum** Bl. Enum. 131. 1828; Fl. Jav. Fil. 150 t. 64. Mett. n. 203. HB. 368. Chr. 114. NPfl. 319 — Malesia. Ins. Philipp.
Drynaria J. Sm. 1841; Pleuridium J. Sm. 1857; Pleopeltis Moore 1862; Phymatopsis J. Sm. 1875 — Polypodium angustatum Bl. 1828. Mett. n. 205 (non Sw. 1806); Pleopeltis Moore 1857; Pleuridium J. Sm. 1866.
palmatum Salom. Nom. 311. 1883 = P. christovalense.
palmatum Sessé et Mociño, Fl. Mex. 261. 1896 — Cuba.
palmoides Bory, Bél. Voy. Bot. **2**. 30. 1833 — Java. India.
paludosum Bl. Fl. Jav. Fil. 192 t. 90. 1829 = Dryopteris brunnea.
palustre Burm. Fl. Ind. 234. 1768 = Stenochlaena.
palustre Salisb. Prod. 403. 1796 = Dryopteris thelypteris.
palustre Lowe. Ferns **2** t. 45. 1858 = Dryopteris serrata.
Panangianum Hk. sp. **5**. 13. 1863 (err. typ.) = Dryopteris penangiana.
pannosum Mett.; Kuhn, Linn. **36**. 141. 1869 = Cyclophorus.

(P) **papillosum** Bl. Enum. 131. 1828; Fl. Jav. Fil. 191 t. 88. Mett. n. 69. HB. 332. Chr. 86. NPfl. 310 — Malesia. Perak. Ins. Philipp.
Thylacopteris Kze.; J. Sm. 1875.
papillosum Ces. Atti Ac. Napoli 7^8. 25. 1876 = P. Cesatianum.
(Pl) **Pappei** Mett; Kuhn, Fil. Afr. 150. 1868 — Africa austr.
P. normale HB. 358 quoad pl. afric.
(Pl) **papuanum** Bak. Malesia **3**. 48. 1886 — N. Guinea.
(P) **paradiseae** Langsd. et Fisch. Ic. Fil. 11 t. 11. 1810. Mett. n. 78. (Chr. 86) — Amer. trop. — P. pectinatum pt. HB. 333 — P. ptiloton Kze. 1834.
Paradisiastrum Fée, Cr. vasc. Br. **1**. 90 t. 29 f. 2. 1869 = P. recurvatum.
paradoxum Col. Tr. N. Zeal. Inst. **14**. 336. 1882 = P. Billardieri.
(P) **paraguayense** Bak. JoB. **1878**. 301 — Paraguay.
parallelinerve Desv. Prod. 230. 1827 — Ins. Africanae?
(L) **parallelum** (Copeland) C. Chr. Ind. 551. 1906 — Luzon.
Loxogramme Copeland in Perkins Fragm. 182. 1905.
parasiticum L. sp. **2**. 1090. 1753 = Dryopteris.
parasiticum Mett. Pol. 36 n. 16. 1857 = P. mediale var.
(P) **parietinum** Kl. Linn. **20**. 373. 1847 — Columbia. — Grammitis Fée 1850-52.
Parkeri Hk. et Grev. Ic. Fil. t. 232. 1831 = Alsophila blechnoides.
Parkinsoni Bak. Ann. Bot. **5**. 481. 1891 = Drynaria.
(P) **parvulum** Bory; Willd. sp. **5**. 182. 1810. Mett. n. 34. HB. 326 (pt.). NPfl. 309 — Africa austr. Ins. Mascar. — Ctenopteris J. Sm. 1875.
parvulum Bl. Fl. Jav. Fil. 187 t. 37 C. 1828 = P. inconspicuum.
parvulum Bedd. Ferns S. Ind. t. 166. 1863-65 = P. subfalcatum.
(P) **pastoense** C. Chr. Ind. 551. 1906 — Columbia. Ecuador. — P. Lehmannianum Hier. Engl. Jahrb. **34**. 513. 1904 (non P. Lehmanni Mett. 1857).
patelliferum Burck, Ann. Jard. Buit. **4**. 96 t. 7. 1884 = Lecanopteris carnosa.
patens Sw. Prod. 133. 1788 = Dryopteris.
(G) **patens** (J. Sm.) Hk. sp. **5**. 23. 1863. HB. 342 — Panama.
Goniophlebium J. Sm. Bot. Voy. Herald 230. 1854.
(P) **patentissimum** Mett.; Kuhn, Linn. **36**. 134. 1869. HB. 509 — Ecuador.
P. Manabyanum Bak. 1877.
pauciflorum Hk. sp. **4**. 242. 1862 = Hemitelia multiflora.
(Gr) **paucinervatum** (Fée) C. Chr. Ind. 332. 1905; 551. 1906 — Brasilia.
Grammitis Fée, Cr. vasc. Br. **2**. 51 t. 96 f. 1. 1872-73.
Pavonianum Kl. Linn. **20**. 386. 1847 = Dryopteris.
(P) **Pearcei** Bak. Syn. 508. 1874 — Columbia. Peru.
Pecten Fée, Gen. 240. 1850-52 = P. apiculatum.
(P) **pectinatiforme** Lindm. Hedw. **43**. 309. 1904 — Brasilia. — P. micrososum Lindm. Ark. för Bot. **1**. 239 t. 11 f. 2. 1903 (non Mett. 1855).
(P) **pectinatum** L. sp. **2**. 1085. 1753. HB. 333 pt. Chr. 85. NPfl. 310 — Amer. trop. — Goniophlebium J. Sm. 1841. Polypodium Otites Willd. 1810. Mett. n. 77 (an L. 1753?) — P. heteroclitum Fée 1869 (non Desv. 1811) — P. elasticum γ. Glaziovii Bak. 1870; P. Glaziovii Christ 1900 (non Bak. 1891; an ad P. truncorum Lindm. pertinens?) — conf. Lindm. Ark. för Bot. **1**. 236.
pectinatum Forsk. Fl. Aegypt.-Arab. 185. 1775 = Dryopteris orientalis.
pectinatum Schkuhr, Kr. Gew. **1**. 189 t. 17 b. 1809 = P. plumula.
pectinellum Fée, 7 mém. 61 t. 27 f. 2. 1857 = P. apiculatum.

pedatum Houtt. Nat. Hist. **14**. 174. 1783; Pfl. Syst. **13**[1]. 196. 1786 = Gleichenia linearis.

pedicularifolium Hoffm. Deutschl. Fl. **2**. 10. 1795 = Athyrium filix femina.

pedicularifolium Pr. Del. Prag. **1**. 168. 1822 = Dryopteris caudata.

pedicularis Willem. Usteri Ann. **18**. 62. 1796 — Mauritius. Aspidium Kuhn 1868 (Dryopteris sp).

(P) **pediculatum** Bak. Syn. ed. I. 455. 1868 — Borneo. — P. Lobbianum Hk. sp. **4**. 226 t. 278 B. 1862. HB. 331 (non Hk. 1853); Ctenopteris J. Sm. 1875.

(S) **pedunculatum** (Hk. et Grev.) Mett.; Salom. Nom. 312. 1883 — India bor. China austr. — Ceterach Hk. et Grev. Ic. Fil. t. 5. 1827; Grammitis Bl. 1828; Selliguea Pr. 1849; Grammitis Hamiltoniana Wall. 1828; Gymnogramme Hk. 1864. HB. 389; Selliguea Bedd. 1868; Polypodium Diels 1899. NPfl. 316; Selliguea Hamiltoni et Hookeri Pr. 1836.

(G) *pellitum* Willd.; Klf. Enum. 89. 1824 — Brasilia. Marginaria Pr. 1856; Goniophlebium Moore 1862.

(P) **pellucidum** Klf. Enum. 101. 1824. HB. 334. Chr. 83 — Ins. Hawaii. — P. myriocarpum Hk. 1837.

peltatum Cav. Descr. 244. 1802 = P. lanceolatum.

peltideum Link, Hort. Berol. **2**. 94. 1833 = P. phymatodes.

Penangianum Hk. sp. **5**. 52. 1863 = Cyclophorus.

(P) **pendulum** Sw. Prod. 131. 1788. Mett. n. 65. HB. 320. Chr. 81. NPfl. 310 — Amer. trop. — Ctenopteris J. Sm. 1875.

pendulum γ. Gardnerianum Bak. Fl. bras. **1**[2]. 513. 1870 = P. brevistipes.

Penna marina Poir. Enc. **5**. 520. 1804 = Blechnum.

pennatum Poir. Enc. **5**. 535. 1804 = Dryopteris megalodus vel D. tetragona.

pennigerum Forst. Prod. 82. 1786 = Dryopteris.

? pennigerum Pr. Rel. Haenk. **1**. 25. 1825 = Dryopteris gongylodes.

(Pl) **pentaphyllum** Bak. Ann. Bot. **5**. 478. 1891 — Ins. Philipp.

Percivalii Jenm.; Bak. Ann. Bot. **5**. 456. 1891 = Dryopteris.

(P) **percrassum** Bak. JoB. **1887**. 26. NPfl. 308 — Costa Rica.

(Pl) **percussum** Cav. Descr. 243. 1802. Mett. n. 159. HB. 356. Chr. 103 — Costa Rica-Peru-Brasilia. — Pleopeltis Hk. et Grev. 1828; Phlebodium J. Sm. 1841; Drynaria Fée 1850-52; Niphobolus Keys. 1873; Lepicystis Diels 1899. NPfl. 323.

(P) **perludens** Bak. Journ. Linn. Soc. **16**. 204. 1877 — Madagascar.

(Pl) **persicariifolium** Schrad. Gött. gel. Anz. **1824**. 867. Mett. n. 178. HB. 358. Chr. 104. NPfl. 316 — Ind. occ. Brasilia. Ecuador. — Microgramma Pr. 1836; Mecosorus Kl. 1847; Drynaria Fée 1850-52; Pleopeltis Moore 1857.

(G) **persicifolium** Desv. Berl. Mag. **5**. 316. 1811. HB. 344 — Java. Celebes. Goniophlebium Bedd. 1870; ? Phegopteris Christ 1900. Polypodium cuspidatum Bl. 1828; Fl. Jav. Fil. 175 t. 82 (non Pr. 1825 nec Don 1825); Goniophlebium Pr. 1836; Schellolepis J. Sm. 1866 — Polypodium grandidens Kze. 1846; Goniophlebium Fée 1850-52.

persimile Bak. JoB. **1876**. 344 = Dryopteris.

pertusum Roxb.; Hk. Exot. Flora **2**. t. 162. 1825 = Cyclophorus adnascens.

(P) **peruvianum** Desv. Prod. 231. 1827. HB. 326. NPfl. 310 — Venezuela-Peru-Argentina. — Ctenopteris J. Sm. 1875.

(Gr) **Pervillei** Mett.; Kuhn, Fil. Afr. 150. 1868. HB. ed. II. 321 — Ins. Sechellae.

(G) **petiolatum** Dav. Bot. Gaz. **19**. 394. 1894 — Mexico.

petiolosum Christ, Nu. Giorn. bot. It. n. s. **4**. 96 t. 1 f. 2. 1897 = Cyclophorus.

(Pl) **petraefolium** Jenm. Bull. Dep. Jam. **4**. 139. 1897 — Guadeloupe. Jamaica.
Drynaria elastica Fée, 11 mém. 72 t. 20 f. 2. 1866 (non Polypodium Bory 1810).

Petrarchae Guérin, Descr. font. Vaucl. 124. 1804 = Asplenium glandulosum.

petrophyum Bl. Fl. Jav. Fil. 163 t. 71. 1829 = Aspidium saxicola.

Phegopteris L. sp. **2**. 1089. 1753 = Dryopteris.

phegopteroides Desv. Journ. bot. appl. **2**. 264. 1814 = Dryopteris crinita.

Philippinum Hk. sp. **5**. 151. 1864 = ? Dryopteris sagenioides*.

phlebodes Kze.; Mett. Pol. 92. n. 163. 1857 = P. excavatum.

(G) **phlebodioides** Copeland, Polyp. Philipp. 123. 1905 — Ins. Philipp.

(P) **phlegmaria** J. Sm. Lond. JoB. **1**. 194. 1842 — Guiana-Ecuador.
— P. tovarense Kl. 1847. HB. ed. II. 324. NPfl. 309; P. subdimidiatum Bak. 1867. HB. ed. I. 324.

(C) **phyllitidis** L. sp. **2**. 1083. 1753. Mett. n. 141. HB. 348. Chr. 96. NPfl. 314 — Amer. trop. (Florida-Argentina. S. Thomé (Guinea)).
Campyloneurum Pr. 1836; Cyrtophlebium J. Sm. 1841. Polypodium repens (Willd. 1810?) Mett. 1856; Pol. n. 142 (non Aublet 1775) — Campylonevron macrosorum Fée 1857 — C. Moritzianum Fée 1850-52.

phyllitidis Thbg. Fl. Jap. 335. 1784 = P. ensatum.

phyllitidis Roxb. Calc. Journ. **4**. 483. 1844 = ? Aspidium singaporianum.

(Pl) **phyllomanes** Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 210 c. figg. — China. Japonia. India bor. — P. ovatum Wall.; Hk. et Grev. Ic. t. 41. 1827. HB. 359 (non Burm. 1768); Phymatodes Pr. 1836; Drynaria Fée 1850-52; Pleopeltis Moore 1857; Colysis J. Sm. 1875. — (an adhuc P. hemitomum Hance?).

(Pl) **phymatodes** L. Mant. 306. 1771. Mett. n. 195. HB. 364. Chr. 110. NPfl. 318 — Ceylon. Tsus-Sima. China merid.-Malesia-Austr. trop. Polynesia. Afr. trop. et austr. c. ins. — ? Tectaria Cav. 1802; Chrysopteris Link 1841; Drynaria Fée 1850-52; Pleopeltis Moore 1857; Phymatodes Maxon 1905; Ph. vulgaris Pr. 1836; Drynaria J. Sm. 1841. Polypodium grossum Langsd. et Fisch. 1810; Phymatodes Pr. 1836; Drynaria Fée 1850-52 — Chrysopteris sp. plur. Link 1841.

Pica L.fil. Suppl. 446. 1781 = Aspidium.

(P) **pichinchae** Sod. Cr. vasc. quit. 329. 1893 — Ecuador.

pichinchense Hier. Engl. Jahrb. **34**. 506. 1904 = P. ecuadorense.

(Pl) **Picoti** Carrière, Rév. Hort. **1886**. 206 f. 62 — Brasilia.

(Pl) **pictum** (Moore) Bak. Ann. Bot. **5**. 479. 1891 — »Polynesia«.
Pleopeltis Moore, Gard. Chr. n. s. **15**. 331. 1881.

piligerum Hk. Ic. pl. t. 321. 1841 = P. achilleifolium.

(P) **pilipes** Hk. Ic. pl. t. 221. 1840. HB. 330 — Peru. — Ctenopteris J. Sm. 1875.

pilipes Kl. Linn. **20**. 382. 1847; Mett. Pol. 51 n. 56. 1857 = P. capillare.

pilosella Vell. Fl. flum. **11** t. 57. 1827; Arch. Mus. Nac. Rio Jan. **5**. 445. 1881 — Brasilia.

(M) **piloselloides** L. sp. **2**. 1083. 1753. Mett. n. 167. HB. 340. (excl. varr.). Chr. 99 — Amer. trop. — Marginaria Pr. 1836; Goniophlebium J. Sm. 1840; Craspedaria Fée 1850-52; Lopholepis J. Sm. 1854; Niphobolus Keys. 1873; Lepicystis Diels 1899. NPfl. 323.

(P) **pilosissimum** Mart. et Gal. Mém. Ac. Brux. **15**. 39 t. 9 f. 2. 1842. Mett. n. 31. HB. 327. NPfl. 309 — Mexico-Ecuador-Brasilia.
— P. acrodontium Fée 1872-73.

(Gr) **pilosiusculum** (Bl.) Pr. Tent. 180. 1836 — Java.
Grammitis Bl. Enum. 115. 1828; Fl. Jav. Fil. 107 t. 46 f. 2.
pilosum Schkuhr, Kr. Gew. **1**. 15 t. 17. 1806 —?
Goniopteris Pr. 1836. (Dryopteris sp.).
pilosum Roxb. Calc. Journ. **4**. 492. 1844 — Chittagong.
pilosum Fée, Cr. vasc. Br. **2**. 58 t. 97 f. 3. 1872-73 = P. cultratum.
pilosum Salom. Nom. 314. 1883 = Dryostachyum splendens.
pinnatifidum Gilib. Exerc. Phytol. **2**. 577. 1792 = P. vulgare.
Pinwillii Bak. Ann. Bot. **5**. 460. 1891 = Dryopteris urophylla.
(C) **Pittieri** Christ, Prim. Fl. Costaric. **3**. 16. 1901 — Costa Rica.
planiusculum Mett. Kuhn, Fil. Afr. 151. 1868 (nomen) = P. ascensionense)
planiusculum Christ: Krug, Engl. Jahrb. **24**. 125. 1897 = P. Hartii.
plantagineum Jacq. Coll. **2**. 104 t. 3 f. 1. 1788 = Aspidium.
(Pl) *platanifolium* Goldm. Nova Acta **19**. Suppl. I. 455. 1843 — Ins. Hawaii.
(G) **platybasis** Bak. Syn. 511. 1874 — Guatemala. Salta.
(M) **platylepis** Mett.; Kuhn, Linn. **36**. 137. 1869. HB. 512 — Guatemala. Columbia.
platylobum Bak. Syn. 307. 1867 = Dryopteris.
(Pl) **platyphyllum** Sw. Syn. 27. 1806. HB. 359. NPfl. 315 — Java. Perak. Borneo.
Pleopeltis Bedd. 1892; Polypodium latifolium Poir. Enc. **5**. 512. 1804 (non Forst. 1786). P. crassinervium Bl. 1828; Fl. Jav. Fil. 145 t. 61. Mett. n. 209; Phymatodes Pr. 1836; Pleopeltis Moore 1857; Pleuridium J. Sm. 1866.
platyphyllum Hk. sp. **4**. 248. 1862 = Polystichum aculeatum var. 23.
(Pl) **Playfairii** Bak. Ann. Bot. **5**. 474. 1891. NPfl. 315 — Formosa.
(M) **plebejum** Schlecht. et Cham. Linn. **5**. 607. 1830. HB. 336. Kuhn, Abh. Nat. Ges. Halle **11**. 40. Chr. 84 — Mexico-Peru-Brasilia.
Lepicystis Diels 1899. NPfl. 322. Marginaria Karwinskyana Pr. 1836; Polypodium A.Br.; Kze. 1850. Mett. n. 95 (pt.) — P. Madrense J. Sm. 1854. HB. 330. P. oulolepis Fée 1857 — P. Pleopeltidis Fée 1869 — P. nigripes Hk. 1863. HB. 347 (non Zoll. 1844); Phlebodium J. Sm. 1877.
plebejum var. columbiensis Kuhn, Abh. Nat. Ges. Halle **11**. 40. 1869 = P. leucosticton.
(Pl) **plebiscopum** Bak. Journ. Linn. Soc. **15**. 110. 1876 — Ins. Ternate.
(G) **plectolepis** (Fée) Hk. sp. **5**. 30. 1863. HB. 344. NPfl. 312 — Mexico. Guatemala. Ecuador. — Goniophlebium Fée, 8 mém. 95. 1857. Polypodium insigne Liebm. Vid. Selsk. Skr. V. **1**. 193 (seors. 41). 1849 (non Bl. 1828).
(P) **pleiosoroides** Copeland in Perkins, Fragm. 189. 1905 — Ins. Philipp (Mindanao).
pleiosoros Hk.fil. Tr. Linn. Soc. **20**. 166. 1847 — Ins. Galapagos.
pleiosorum Mett.; Kuhn, Linn. **36**. 128. 1869 = P. tahitense.
Pleopeltidis Fée, Cr. vasc. Br. **1**. 86 t. 26 f. 1. 1869 = P. plebejum.
pleopeltifolium Raddi, Opusc. sci. Bol. **3**. 286. 1819; Pl. Bras. **1**. 16 t. 21 f. 2. 1825 = P. angustum.
(G) **plesiosorum** Kze. Linn. **18**. 313. 1844. Mett. n. 111. HB. 342 — Mexico-Venezuela. — Goniophlebium Fée 1857 — Polypodium gonatodes Kze. 1850 — Goniophlebium anisomeron Fée 1857; Polypodium Fourn. 1872 — ? Polypodium appendiculatum Kl. 1855 (non alior.); Goniophlebium Moore 1856 — Polypodium rhodopleuron Kze. 1844. Mett. n. 110; Goniophlebium Fée 1857; Polypodium Fourn. 1872 (non Mett. 1857) — conf. Fourn. Mex. pl. **1**. 81-82.
pleuridioides Mett. Ann. Lugd. Bat. **2**. 229. 1866 = Drynaria.

(Ph) **pleurosorum** Kze.; Mett. Pol. 101 n. 187 t. 2 f. 6, 7. 1857 — Mexico.
Plukenetii Loisel. Notice 146. 1810 = Polystichum lobatum.
(Pl) ? *Plukenetii* Pr. Rel. Haenk. **1**. 21. 1825 — Ins. Philipp. — Phymatodes Pr. 1836.
Plumieri Desv. Berl. Mag. **5**. 316. 1811 = Dryopteris opposita.
plumosum Pr. Tent. 178. 1836 (nomen) —?
(P) **plumula** H.B.Willd. sp. **5**. 178. 1810. Mett. n. 74 — Amer. trop. Florida. Ctenopteris J. Sm. 1875. Polypodium elasticum Rich. 1792 (nomen, non Bory 1810). HB. ed. II. 332. Chr. 86. NPfl. 310. P. elegans Poir. 1804 (non Cav.; Sw. 1806). P. Schkuhrii Raddi 1819.
(Pl) **podobasis** Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 215 — Yunnan.
(S) **podopterum** Christ, Journ. de Bot. **19**. 125. 1905 — Annam.
poecilophlebium Hk sp. **5**. 14. 1863 = Dryopteris.
Poeppigianum Mett. Pol. 37. n. 17. 1857 = P. Billardierii.
Poeppigii Kze. Linn. **9**. 50. 1834 = Dryopteris punctata.
Pohlianum Pr. Tent. 180. 1836 (nomen) —?
Poiretii Spr. Syst. **4**. 60. 1827 (Plum. t. 35) — Hispaniola. P. rostratum Poir. Enc. **5**. 555. 1804 (non Cav. 1802).
politum Poir. Enc. **5**. 553. 1804 = Polystichum adiantiforme.
polyanthum ht. Berol.; Mett. Pol. 84. n. 142 b. 1857 = P. nitidum.
polycarpon Cav.; Sw. Schrad. Journ. **1800**[2]. 21. 1801; Cav. Descr. 246. 1802 = P. punctatum.
polycarpon Hk. et Arn Bot. Beech. Voy. 104. 1832 = Dryopteris stegnogrammoides.
polycarpum Bak. Syn. 506. 1874 = Aspidium Kuhnii.
polycephalum Wall. List n. 273. 1828 = P. punctatum.
polydactylon Hance, JoB. **1883**. 269 = Cyclophorus.
(M) **polylepis** Roem.; Kze. Linn. **13**. 131. 1839. Mett. n. 156 — Mexico. Pleopeltis Moore 1862. Drynaria vestita Fée 1850-52.
polymorphum Vill. Hist. Pl. Dauph **1**. 282, 292. 1786; **3**. 846. 1789 = Cystopteris fragilis.
polyodon Wall. List n. 7079. 1829 = Polystichum Hookerianum.
(M) **polypodioides** (L.) Hitchcock, Rep. Mo. Bot. Gard. **4**. 156. 1893 — U. S. centr. et austr.-Argentina-Chile. Africa austr.
Acrostichum L. sp. **2**. 1068. 1753; Candollea Mirb. 1803; Polypodium incanum Sw. 1788. Mett. n. 100. HB. 346. Chr. 89; Pleopeltis Wall. 1828; Marginaria Pr. 1836; Goniophlebium J. Sm. 1841; Lepicystis J. Sm. 1842. NPfl. 323. Polypodium ceteraccinum Michx. 1803; Marginaria Bory 1826; Goniophlebium Fée 1869 — Marginaria minima Bory 1826 - Polypodium Eckloni Kze. 1836. Mett. n. 99 (Afr.) — P. microlepis Fée 1850-52; Goniophlebium Fée 1869 — Acrostichum ferrugineum L. 1753.
polystichoides Kl. Linn. **20**. 383. 1847 = Polystichum aculeatum var. 23.
polystichops O.Ktze. Rev. Gen. Pl. **2**. 819. 1891 = Dryopteris pubescens.
polystichum Link, Hort. Ber. **2**. 101. 1833 = P. fraxinifolium.
polyxiphion Bak. Bull. Linn. Soc. Paris **1**. 534. 1885 = Dryopteris biformis.
Pontederae All. Fl. Pedem. **2**. 286. 1785 = Cystopteris fragilis.
(P) **Poolii** Bak. Journ. Linn. Soc. **15**. 419. 1876 — Madagascar.
porosum Wall. List n. 266. 1828 = Cyclophorus.
porrectum Willd. sp. **5**. 161. 1810 = P. crassifolium.
portoricense Spr. Nova Acta **10**. 230. 1821 = Dryopteris pubescens var.
pothifolium Mett. Fil. Lips. 130 t. 25 f. 21. 1856 = P. ellipticum.

(Pl) **Powellii** Bak. Syn. 364. 1868. NPfl. 318 — Samoa. — Pleopeltis Carr. 1873.
(P) **pozuzoense** Bak. in Hk. Ic. pl. t. 1672. 1886. NPfl. 311 — Peru.
praelongum Poir. Enc. **5**. 521. 1804 = ? P. curvatum.
(G) **Prainii** (Bedd.) C. Chr. Ind. 327. 1905; 556. 1906 — Perak.
Goniophlebium Bedd. JoB. **1893**. 226.
prasinum Bak. Syn. 312. 1867 = Dryopteris.
Preslianum Spr. Syst. **4**. index 1827; Link, Hort. Ber. **2**. 102. 1833 = P. brasiliense.
Preslii Wikstr. Vet. Akad. Handl. **1825**. 442. 1826 — Brasilia.
P. auriculatum Pr. Del. Prag. **1**. 167. 1822 (non L. 1753 nec Raddi 1819).
Preslii Opiz, Flora **1834**. 591 = P. percussum (cuspidatum Pr.).
Prieurii Christ, Prim. Fl. Costaric. **3**. 17. 1901 = P. lanceolatum*.
princeps Mett. Ann. Lugd. Bat. **2**. 232. 1866 = Cyclophorus.
(G) **Pringlei** (Maxon) C. Chr. Ind. 327. 1905; 556. 1906 — Mexico.
Goniophlebium Maxon, Proc. U. S. Nat. Herb. **27**. 953 t. 48. 1904.
prionites Kze. Flora **1839**[1]. Beibl. 29 = Dryopteris.
procerum Willd. sp. **5**. 206. 1810 = Alsophila.
procerum Brack. Expl. Exp. **16**. 14 t. 3. 1854 = Dryopteris Keraudreniana.
procurrens Kze. Anal. 17. 1837 = Arthropteris altescandens.
proliferum Klf. Enum. 107. 1824 = Dryopteris vivipara.
proliferum Roxb.; Wall. List n. 312. 1828 = Dryopteris.
prolongatum Mak. —? = Polystichum flagellare.
propinquum Wall. List n. 293. 1828 = Drynaria.
protensum Poir. Enc. **5**. 553. 1804 = Dryopteris.
pruinatum Sw. Schrad. Journ. **1800**[2]. 29. 1801 = Alsophila quadripinnata.
(P) **pruinatum** Bak. Syn. 508. 1874 — Nicaragua.
(Gr) **pseudaustrale** Fourn. Ann. sc. nat. V. **18**. 282. 1873 — N. Caledonia.
Grammitis Fourn. Bull. Soc. Fr. **16**. 423. 1869.
pseudoaureum Cav. Descr. 247. 1802 = P. aureum var. sporadocarpum.
(G) **pseudo-dimidiatum** Christ, Bull. Soc. Fr. **52** Mém. I. 14. 1905 — Yunnan.
(Gr) **pseudogrammitis** Gaud. Freyc. Voy. Bot. **1**. 345. 1827. HB. 320 — Ins. Hawaii.
Grammitis tenella Klf. Enum. 84. 1824 (non Polypodium Forst. 1786); Polypodium Kaulfussii Pr. 1836. Mett. n. 11.
(G?) **pseudoserratum** Christ, Bull. Boiss. **6**. 871. 1898 — China merid. (an P. crenato-pinnatum?).
pteridioides Reich.; Linn. Syst. pl. ed. novis. **4**. 424. 1780 = Cheilanthes.
pterioides Lam. Fl. franç. **1**. 18. 1878 = Dryopteris thelypteris.
pteroides Retz. Obs. Bot. **6**. **39**. 1791 = Dryopteris.
pteroides Pr. Rel. Haenk. **1**. 25 t. 4 f. 4. 1825 = Aspidium Brongniartii.
pteroideum Kl. Linn. **20**. 389. 1847 = Dryopteris.
pteryoides Bory, Ess. isles fort. 311. 1803 = Cheilanthes pteridioides.
(Pl) **pteropus** Bl. Enum. addend. 1828; Fl. Jav. Fil. 168 t. 76. 1829. Mett. n. 197. HB. 362. Chr. 109. NPfl. 318 — Asia trop.
Pleopeltis Moore 1857. Polypodium tridactylon Wall. 1828; Hk. et Grev. t. 209; Phymatodes Pr. 1836; Drynaria Fée 1850-52; Pleopeltis Moore 1857; Colysis J. Sm. 1875.
pteropus Hk. sp. **4**. 192 t. 275 B. 1862 = P. subsessile.
(G) **ptilorhizon** Christ, Bull. Boiss. II. **5**. 6. 1905 — Costa Rica.
ptiloton Kze. Linn. **9**. 42. 1834 = P. paradiseae.

(P) **puberulum** Schlecht. et Cham. Linn. 5. 607. 1830. Mett. n. 85. HB. 336 — Mexico-Guatemala.

puberulum Bak. Syn. ed. I. 344. 1867 = P. Beddomei.

pubescens L. Syst. Nat. ed. X. 2. 1327. 1759 = Dryopteris.

pubescens Poir. Enc. 5. 546. 1804 = Nephrodium Poiretii.

pubescens Raddi, Pl. Bras. 1. 23 t. 34. 1825 = Dryopteris oligocarpa var.

pubescens Schum. Vid. Selsk. Afh. 4. 227 (seors. 453). 1829 = Dryopteris [protensa.

pubescens Gill.; Hk. et Grev. Ic. Fil. t. 182. 1830 = P. Gilliesii.

pubescens Fée, 8 mém. 87. 1857 = P. Fournieri.

pubigerum Wall. List n. 7078. 1829 = Aspidium Leuzeanum.

(Gr) **pubinerve** (Bl.) Christ, Ann. Jard. Buit. 15. 146. 1897 — Java. Celebes.
Grammitis Bl. Enum. 116. 1828; Fl. Jav. Fil. 117 t. 48 f. 4.

pulchellum Willem. Usteri Ann. 18. 62. 1796 — Mauritius.
Aspidium Kuhn 1868 (non Bl. 1828). — (Dryopteris? sp.).

pulchellum Salisb. Prod. 403. 1796 = Dryopteris Linnaeana.

pulchrum Mart. et Gal. Mém. Ac. Brux. 15. 41 t. 8 f. 2. 1842 — Mexico.

pulverulentum Poir. Enc. 5. 555. 1804. (Plum. t. 34) — Hispaniola.
Aspidium Desv. 1827; A. lutescens Willd. 1810; Lastrea? Moore 1858. (an Dryopteris ampla ?).

pulvinatum Link, Hort. Berol. 2. 99. 1833 = P. aureum*.

(Gr) **pumilio** Hier. Hedw. 44. 102 nota. 1905 — N. Caledonia.
P. nanum Vieill.; Bak. Syn. 507. 1874 (non Fée 1850-52).

punctatum Thbg. Fl. Jap. 337. 1784 = Dryopteris.

(Pl) **punctatum** (L.) Sw. Schrad. Journ. **1800**^{2}. 21. 1801. Chr. 106 — Asia trop.-Austr. trop. Polynesia. Afr. trop. Ins. Mascar.
Acrostichum L. sp. ed. II. 2. 1524. 1763; Pleopeltis Bedd. 1876. Polypodium irioides Poir. 1804. Mett. n. 235. HB. 360. NPfl. 315; Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Microsorium Fée 1850-52; Pleopeltis Moore 1857; Colysis J. Sm. 1875 — Polypodium polycarpon Cav. 1802; Niphobolus Spr. 1827; Phymatodes Pr. 1836; Drynaria Brack. 1854; Pleopeltis Moore 1857 — Polypodium sessile Klf. (Kze. 1848); Phymatodes Pr. 1836; Microsorium Fée 1850-52; Pleopeltis Moore 1857.

punctatum Hk. sp. 4. 172. 1862 = Monogramma seminuda.

punctatum Spruce; Hk. sp. 4. 262. 1862 = Dryopteris prasina.

punctatum var. rigescens Bak. Fl. bras. 1^{2}. 503 t. 65. 1870 = Hypolepis rigescens.

punctilobum Poir. Enc. 5. 549. 1804 = Dennstaedtia punctilobula.

punctulatum Poir. Enc. 5. 533. 1804; Vahl, Ecl. 3. 52. 1807 = Nephrolepis [biserrata*.

punctulatum Hk. Ic. pl. t. 720. 1845 = P. falcaria var.

pungens Willd. sp. 5. 206. 1810 = Alsophila.

purpurascens Nadeaud, Enum. 26. 1873 — Tahiti.

pusillum Christ, Ann. Jard. Buit. 15. 145. 1897 = P. hirtellum.

(Pl) **pustulatum** Forst. Prod. 81. 1786. Mett. n. 188. HB. 363. Chr. 111. NPfl. 318 — Australia. N. Caledonia. N. Zealand.
Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Pleopeltis Moore 1857 — Polypodium scandens Forst. 1786; Phymatodes Pr. 1836; Drynaria Fée 1850-52. (vide tamen Carr in Seem. Fl. Vit. 369. 1873).

(M) **pycnocarpum** C. Chr. Ind. 326. 1905; 557. 1906 — Mexico-Chile-Argentina.
P. macrocarpum Pr. Rel. Haenk. 1. 23 t. 1 f. 4. 1825. Mett. n. 94. HB. 330. Chr. 83 (non Bory 1810); Goniophlebium J. Sm. 1841; Lepi-

cystis Diels 1899. NPfl. 322. Pleopeltis pinnatifida Gill.; Hk. et Grev. 1829 (non Polypodium Gilib. 1792).
pycnolepis Hk. sp. **4**. 247. 1862 = Polystichum aculeatum var. 27.
pycnosorum Link, Fil. sp. 128. 1841 = P. latipes.
(Gr) **pygmaeum** Buchinger; Kuhn, Fil. Afr. 152. 1868. HB. 507 — Réunion. (an P. furcatum var.?).
pyramidale L. sp. **2**. 1093. 1753 = Adiantum.
(G?) **pyrolifolium** Goldm. Nova Acta **19**. Suppl. I. 453. 1843 — Luzon.
Marginaria nummularia Pr. 1836 (nomen); Crypsinus Pr. Epim. 123. 1849; Craspedaria Fée 1850-52; Polypodium Mett. 1857. n. 199. HB. 340; Goniophlebium Moore 1857; Pleopeltis Moore 1862.
pyrrhorhachis Kze. Linn. **24**. 257. 1851 = Dryopteris brunnea.
quadripartitum Poir. Enc. **5**. 543. 1804 = Gleichenia.
quadripinnatum Gmel. Syst. Nat. 2^2. 1314. 1791 = Alsophila.
quercicans Fée, 11 mém. 50. 1866 = P. Swartzii.
quercifolium L. sp. **2**. 1087. 1753 = Drynaria.
quercioides Desv. Prod. 235. 1827 = Drynaria quercifolia.
(Pl) **quinquefidum** Bak. JoB. **1880**. 216 — Sumatra.
quinquefurcatum Bak. Journ. Linn. Soc. **24**. 260. 1887 = Dipteris.
(P) **quitense** Bak. JoB. **1877**. 165 — Ecuador.
(S) **Raciborskii** C. Chr. Ind. 336. 1905; 558. 1906 — Java.
Gymnogramme grandis Racib. Pter. Buit. 72. 1898 (non Bak. 1868 nec Polypodium Pr. 1822).
raddii Desv. Prod. 232. 1827 = P. lepidopteris.
(P) **radicale** Moritz; Eat. Mem. Amer. Acad. n. s. **8**. 198. 1860 (nomen); Hier. Engl. Jahrb. **34**. 511. 1904 — Venezuela-Columbia.
radicale Bak. Syn. ed. I. 332. 1867 = ? P. meridense vel ? P. radicale.
radicans Burm. Fl. Ind. 233 t 66 f. 3. 1768 = Nephrolepis.
radicans Poir. Enc. **5**. 530. 1804 = Dryopteris reptans.
ramosum Burm. Fl. Ind. 235 t. 67 f. 2. 1868. (an Houtt. Pfl. Syst. 13^1. 197. 1786?) — Ceylon.
ramosum Lodd.; Link, Hort. Berol. **2**. 97. 1833 = P. loriceum.
(Pl) **rampans** Bak. Journ. Linn. Soc. **15**. 109. 1876 — Ins. Admiralitatis.
Rasamalae Racib. Pter. Buit. 99. 1898 = Cyclophorus.
recedens J. Sm. JoB. **3**. 394. 1841 = Dryopteris.
reclinatum Brack. Expl. Exp. **16**. 11. 1854 = P. cultratum var.
(Pl) **recreense** Hier. Engl. Jahrb. **34**. 537. 1904 — Ecuador.
rectangulare Zoll. Syst. Verz. 37, 48. 1854 — Java.
(P) **recurvatum** Klf. Enum. 106. 1824. Mett. n. 79. HB. 332. NPfl. 311 — Guadeloupe. Brasilia-Argentina. Ecuador.
P. Paradisiastrum Fée 1869 — P. extensum Fée 1869.
refractum Fisch. et Mey.; Kze. Linn. **23**. 283, 321. 1850 = Dryopteris.
refulgens Kl.; Bak. Syn. 307. 1857 = Dryopteris.
regium L. sp. **2**. 1091. 1753 = Cystopteris fragilis*.
(S) **regulare** Mett. Ann. Lugd. Bat. **2**. 225. 1866 — Borneo.
Gymnogramme Bak. 1868. HB. 388.
(Gr) **Reinwardtii** (Bl.) Pr. Tent. 180. 1836. Mett. n. 20. NPfl. 308 — Java.
Grammitis Bl. Enum. addend. 1828; Fl. Jav. Fil. 114 t. 48 f. 1; Polypodium hirtum pt. HB. 320 — Grammitis obscura Bl. 1828. Fl. Jav. Fil. 113 t. 50 f. 1; Polypodium Mett. 1857. n. 23.

Reinwardtii Kze. Linn. **23**. 283. 1850 = P. subauriculatum.
remissum Kl.; Salom. Nom. 317. 1883 (nomen) — Ceylon.
remotum Desv. Prod. 232. 1827 — Amer. calidior.
remotum Bak. Ann. Bot. **5**. 470. 1891 = P. uniseriale.
(P) **repandulum** (Kze.) Mett. Pol. 50. n. 54. 1857. HB. 328 — Ceylon.
Ctenopteris Kze.; Zoll. Syst. Verz. 37. 1854.
repandum Lour. Fl. Cochinch. **2**. 826. 1790 — China.
repandum Sw. Schrad. Journ. **1800**^{2}. 25. 1801 — Jamaica.
repandum Vahl, Ecl. **3**. 53. 1807 = Aspidium martinicense.
repandum Vell. Fl. flum. **11** t. 73. 1827; Arch. Mus. Nac. Rio Jan. **5**. 448. 1881 = Dryopteris grandis.
repandum Pr. Tent. 179. 1836 (nomen) — ?
repandum Fée, Cr. vasc. Br. **1**. 87 t. 29 f. 1. 1869 = P. suspensum.
(C) **repens** Aublet, Hist. pl. Guian. **2**. 962. 1775. Sw. Prod. 130. 1788. HB. 348. Chr. 96 — Amer. trop. — Campyloneurum Pr. 1836; Cyrtophlebium J. Sm. 1841 — Polypodium caespitosum Link 1833. Mett. n. 139; Campyloneuron Link 1841 — Polypodium brevifolium Link 1833. Mett. n. 140; Campyloneurum Link 1841.
repens Sw. Prod. 132 (non 130). 1788 = Dryopteris reptans.
repens (Willd. sp. **5**. 156. 1810?); Mett. Fil. Lips. 34 t. 24 f. 1-3. 1856 = [P. phyllitidis.
reptans Gmel. Syst. Nat. **2**^{2}. 1309. 1791 = Dryopteris.
resiniferum Desv. Berl. Mag. **5**. 317. 1811 — Amer. trop. — (Dryopteris sp.).
(M) **restingae** Christ in Schwacke, Pl. Nov. Mineiras **2**. 21. 1900; Bull. Boiss. II. **2**. 374. 1902 — Brasilia austr.
reticulatum L. Syst. Nat. ed. X. **2**. 1325. 1759 = Dryopteris.
reticulatum Boj. Hort. Maur. 419. 1837 = Aspidium subtriphyllum.
reticulatum Sessé et Mociño, Fl. Mex. 261. 1896 — Cuba.
retroflexum L. sp. **2**. 1089. 1753. (Plum t. 68) — Hispaniola. (an Dryopteris refracta?).
(G) *retrofractum* Desv. Prod. 237. 1827 — Amer. calidior.
Goniophlebium Moore 1862.
retusum Sw. Vet. Akad. Handl. **1817**. 61 = Dryopteris.
(G) **revertens** (Fée) Fourn. Mex. pl. **1**. 79. 1872 — Mexico.
Goniophlebium Fée, 8 mém. 94. 1857.
(Pl) **revolutum** (J. Sm.) C. Chr. Ind. 331. 1905; 559. 1906 — Malesia. Ins. Philipp. N. Caledonia. — Drynaria J. Sm. 1841 (nomen); Grammitis longifolia Bl. Enum. 119. 1828; Pleopeltis Bedd. 1828; Paragramma Moore 1857; Phymatodes J. Sm. 1857; Polypodium Mett. 1857. n. 153. HB. 355. Chr. 103. NPfl. 315 (non Cav. 1802 nec Pr. 1822); Niphobolus Keys. 1873 — Grammitis decurrens Bl. 1828; Pleopeltis Bl. 1828; Polypodium Kze. 1846 (non Raddi 1819); Paragramma Moore 1857; Polypodium contiguum Hk. 1854 (an Wall. 1828? nomen; non Brack. 1854).
rhachipterygium Liebm. Vid. Selsk. Skr. V. **1**. 191 (seors. 39). 1849 = P. synammia.
rhaeticum L. sp. **2**. 1091. 1753 = Athyrium filix femina et A. alpestre.
rhagadiolepis Fée, Gen. 237. 1850-52 = P. thyssanolepis.
rheosorum Bak. JoB. **1884**. 363 = Aspidium.
rheosorum Bak. Ann. Bot. **5**. 457. 1891 = Dryopteris lepidorachis.
rhizocaulon Willd. sp. **5**. 196. 1810 = P. fraxinifolium.
rhizophyllum Sw. Prod. 132. 1788 = Polystichum.

rhodopleuron Kze. Linn. **18**. 315. 1844 = P. plesiosorum var.
rhomboideum Bl. Enum. 124. 1828 = P. triquetrum.
(Pl) **rhynchophyllum** Hk. Ic. pl. t. 954. 1854. HB. 359. NPfl. 316 — India bor. Ins. Philipp. — Pleopeltis Moore 1857; Phymatopsis J. Sm. 1875.
Richardii Kl. Linn. **20**. 394. 1847 = P. adnatum var.
Riedelianum Fisch.; Kze. Linn. **23**. 284. 1850 = P. trichomanoides.
Riedleanum Mett. Pheg. u. Asp. 104. 1858 (err.) = Polystichum.
(P) **rigens** Maxon, Proc. U. S. Nat. Mus. **27**. 741. 1904; Bull. Dep. Agr. Jam. **2**. 176. 1904 — Jamaica.
(P) **rigescens** Bory; Willd. sp. **5**. 183. 1810. Mett. n. 29 b. HB. 331. NPfl. 309 — Amer. trop. Fernando Po. Réunion. — Ctenopteris J. Sm. 1875.
rigidulum Sw. Schrad. Journ. **1800**². 26. 1801 = Drynaria.
rigidum Aublet, Hist. pl. Guian. **2**. 963. 1775 — Guiana.
(f. Aubl. Plum. t. 135 = Elaphoglossum longifolium).
rigidum Hoffm. Deutschl. Fl. **2**. 6. 1795 = Dryopteris.
rigidum Hk. et Grev. Ic. Fil. t. 163. 1829 = Polystichum aculeatum var. 24.
rigidum Lowe, Ferns **2** t. 37 A. 1858 = P. angustifolium *lucidum.
(G) **Rimbachii** Sod. Cr. vasc. quit. 356. 1893 — Ecuador.
riograndense Lindm. Ark. för Bot. **1**. 230 t. 9 f. 6. 1903 = Dryopteris.
rivale Mett.; Bak. Syn. 368. 1867 = Drynaria mollis.
Rivoirei Bak. Ann. Bot. **5**. 460. 1891 = Dryopteris Poiteana var.
rivulare Vahl, Ecl. **3**. 51. 1857 = Nephrolepis.
rivulorum Raddi, Pl. bras. **1**. 23 t. 35. 1825 = Dryopteris opposita.
Robertianum Hoffm. Deutschl. Fl. **2**. 20. 1795 = Dryopteris.
robustum Fée, Cr. vasc. Br. **1**. 92 t. 28 f. 1. 1869 = P. pectinatum.
roraimense Bak. Tr. Linn. Soc. II. Bot. **2**. 291. 1887 = Dryopteris.
(Pl) **rosmarinifolium** Kunth, Syn. pl. aequin. **1**. 71. 1822; Hier. Engl. Jahrb. **34**. 536. 1904 — Ecuador.
(Pl) **Rosthornii** Diels, Engl. Jahrb. **29**. 205. 1900 — China.
rostratum Burm. Fl. Ind. 235. 1768 — India.
rostratum Cav. Descr. 244. 1802 — Quito.
rostratum Poir. Enc. **5**. 555. 1804 = P. Poireti.
rostratum H.B.Willd. sp. **5**. 193. 1810 = Alsophila blechnoides.
rostratum Hk. Ic. pl. t. 953. 1854 = P. subrostratum.
(P) **rosulatum** Christ, Bull. Boiss. **4**. 662. 1896 — Costa Rica.
rotundato-crenatum Sieb. Fl. Mart. exs. n. 367 = P. trifurcatum.
rotundatum Hk. sp. **4**. 238. 1862 = Dryopteris platyloba.
rotundatum Bak. Fl. bras. **1**². 499 t. 63. 1870 = Dryopteris.
rubidum Kze. Bot. Zeit. **1848**. 117 = P. longissimum.
rubidum Hk. sp. **5**. 12. 1863 = Dryopteris.
rubrinerve Bak. Syn. 315. ed. II. 1874 = Dryopteris.
rude Kze. Linn. **13**. 133. 1839 = Dryopteris.
(Pl) **rudimentum** Copeland in Perkins, Fragm. 190. 1905 — Luzon.
rufescens Bl. Fl. Jav. Fil. 194 t. 91. 1829 = Dryopteris.
ruffum Poir. Enc. **5**. 532. 1804 = Dryopteris rufa.
rufobarbatum Col. Tr. N. Zeal. Inst. **16**. 347. 1844 = Dryopteris punctata.
rufulum Pr. Del. Prag. **1**. 164. 1822 = P. lepidopteris var.
rugosulum Lab. N. Holl. pl. spec. **2**. 92 t. 241. 1806 = Dryopteris punctata*.
rugulosum auctt. = rugosulum.
Ruizianum Kl. Linn. **20**. 385. 1847 — Peru. — (Dryopteris sp.).

runcinatum Desv. Prod. 230. 1827 = P. Swartzii.
rupestre R.Br. Prod. Fl. N. Holl. 146. 1810 = Cyclophorus.
rupestre Bl. Enum. 124. 1828 = P. triquetrum var.
rupestre Roxb. Calc. Journ. 4. 488. 1844 — Penang. — (Nephrolepis sp.).
rupestre Hk. sp. 5. 46. 1863 = Cyclophorus serpens et C. confluens.
Russelianum Wall. List n. 7077. 1829 = Dryopteris setigera.
rusticum Bak. Syn. 306. 1867 = Dryopteris.
(Gr) **Rutenbergii** Luerss. Abh. Nat. Ver. Bremem 7. 48 t. 1 f. 1-2. 1882 — [Madagascar.
Ryani Klf. Enum. 92. 1824 — Montserrat.
P. Lingua Vahl, Ecl. 3. 50. 1807 (non Sw. 1806).
saccatum Fée, Gen. 239. 1850-52; 6 mém. 10 t. 7 f. 3. 1853 = P. jubiforme.
saccatum Lowe, Ferns 2 t. 59. 1858 = P. nigrescens.
sagitta Christ, Bull. Boiss. 6. 199. 1898 = Christopteris.
sagittatum Sw. Prod. 132. 1788 = Dryopteris.
salicifolium Vahl, Ecl. 3. 51. 1807 — Montserrat. (= P. brasiliense?).
salicifolium Willd. sp. 5. 149. 1810 = P. lycopodioides*.
salicifolium Hk. sp. 4. 242. 1862 = Dryopteris straminea.
salicinum Wikstr. Vet. Akad. Handl. 1825. 443. 1826 = P. lycopodioides*.
samarense Mett. Pol. 123. n. 246 t. 3 f. 1-4. 1857 = Cyclophorus.
samarense Christ, Bull. Boiss. 6. 199. 1898 = Cyclophorus lanuginosus.
(P) **samoense** Bak. Syn. 321. 1867. NPfl. 308 — Samoa. ? Ins. Hawaii.
samoense var. glabra Hill. Fl. Haw. 554. 1888 = P. Knudsenii.
(G) **Sampsoni** Bak. Ann. Bot. 5. 471. 1891 — Canton.
Sancti-Gabrieli Hk. sp. 4. 233. 1862 = Dryopteris.
sanctum Sw. Prod. 133. 1788 = Dryopteris.
sandwicense Hk. et Arn. Bot. Beech. Voy. 105. 1832 = Dryopteris.
sandwicense Hk. sp. 5. 5. 1863 = Dryopteris stegnogrammoides.
(Pl) **sarawakense** Bak. Journ. Linn. Soc. 22. 228. 1886 — Borneo.
(My) **sarcopus** de Vriese et Teysm.; Bak. Syn. 514. 1874. Chr. 112. NPfl. 321 — Celebes. China merid.
sarmentosum Boj. Hort. Maur. 419. 1837 (nomen) = Dryopteris orientalis.
(P) **sarmentosum** Brack. Expl. Exp. 16. 8 t. 2 f. 3. 1854. HB. 327 — Ins. Hawaii.
(Pl) **Sauvinieri** (rectius Sauvinerrei) Bak. Ann. Bot. 5. 480. 1891 — Celebes.
(Gr?) **savaiense** Powell; Bak. JoB. 1876. 344 — Samoa.
saxatile Kl. Linn. 20. 383. 1847 = Polystichum aculeatum var. 24?
saxatile Mett. Pol. 116. n. 227. 1857 = P. triquetrum var. rupestre.
saxicola Sw. Vet. Ak. Handl. 1817. 59 t. 3 f. 5 = Dryopteris aspidioides.
saxicolum Bak.; Jenm. JoB. 1877. 264 = P. induens.
scabridum Wall. List n. 302. 1828 = Dryopteris arida.
(Gr) **scabristipes** Bak. JoB. 1890. 108 — N. Guinea.
scabrum Pr. Del. Prag. 1. 169. 1822 — Brasilia.
Lastrea Pr. 1849. — (Dryopteris sp.).
scabrum Roxb.; Wall. List n. 2225. 1829; Roxb. Calc. Journ. 4. 491. 1844 = Dryopteris ferox.
(G) **scalare** Christ, Bull. Soc. Fr. 52 Mém. I. 14. 1905 — Yunnan.
scandens Forst. Prod. 81. 1786 = P. pustulatum.
scandens Lab. Nov. Holl. pl. spec. 2. 91 t. 204. 1806 = P. diversifolium.
scariosum Roxb. Calc. Journ. 4. 494. 1844 — Amboyna.
Schaffneri Fée, 9 mém. 22. 1857 = P. subpetiolatum.
(Gr) **Schenckii** Hier. Hedwigia 44. 87. 1905 — Brasilia.

(P) **Schenkii** Harr. Journ. Linn. Soc. **16.** 31. 1877 — Ins. Philipp.
Schimperianum Mett.; Kuhn, Fil. Afr. 152. 1868 = Cyclophorus.
Schkuhri Raddi, Opusc. sci. Bol. **3.** 287. 1819; Pl. Bras. **1.** 19 t. 27 f. 2. 1825 = P. plumula.
Schkuhrii Bory, Ann. sc. nat. **5.** 467. 1825 = Drynaria quercifolia.
Schneideri ht. Veitch; Gard. Chr. III. **15.** 665. 1894 — Hort.
(Pl) **Schneideri** Christ, Bull. Boiss. **6.** 835. 1898 — Sumatra.
(Pl) **Schnittspahnii** Christ, Bull. Boiss. **6.** 836. 1898 — »Andes«.
Schomburgkianum Kze. Farnkr. **1.** 88 t. 42. 1842 } = P. megalophyllum.
Schomburgkii Hk. sp. **5.** 68. 1863 }
(Pl) **Schraderi** Mett. Pol. 98 n. 179 t. 2 f. 11. 1857. HB. 354 — Africa austr. Niphobolus Keys. 1873; Polypodium elongatum Schrad. Gött. gel. Anz. **1818.** 915 (non Mett. 1857); Phymatodes Pappe et Raws. 1858.
Schraderi Milde, Fil. Eur. 16. 1867 = P. lineare.
Schroderi ht.; Gard. Chr. III. **24.** 241. 1898 — Hort. (P. aureum × vulgare).
(P) **Schwackei** Christ in Schwacke, Pl. Nov. Mineiras II. 20. 1900; Bull. Boiss. II. **2.** 370. 1902 — Brasilia.
Schumannianum Diels in Schum. et Laut. Fl. deut. Südsee 139 t. 3 C, D. 1901 = P. musifolium.
scolopendria Burm. Fl. Ind. 232. 1768 = P. phymatodes.
(L) **scolopendrinum** (Bory) C. Chr. Ind. 60. 1905; 562. 1906 — China. Asia trop. Melanesia. — Grammitis Bory, Dup. Voy. Bot. **1.** 257 t. 30 f. 1. 1828 (patria err.?); Loxogramme? Pr. 1836; Selliguea Fée 1850—52; Grammitis involuta Don, Prod. Fl. Nepal. 14. 1825; Antrophyum Bl. 1828; Loxogramme Pr. 1836; Selliguea Kze. 1851; Polypodium Mett. 1856. Pol. n. 218. Chr. 106 (non Desv. 1811); Gymnogramme Hk. 1864. HB. 387. Selliguea Wallichiana Hk. 1840; Colysis Pr. 1849; Polypodium Mett. n. 213 b. 1857 (non Spr. 1827) — ? Grammitis flavescens Wall. 1828; Loxogramme Pr. 1836; Selliguea J. Sm. 1841.
scolopendrioides L. sp. **2.** 1085. 1753 = Dryopteris.
scolopendrioides Lour. Fl. Cochinch. **2.** 827. 1790 — Cochinchina.
scolopendrioides Hk. et Grev. Ic. Fil. t. 42. 1828 = P. trifurcatum.
Scolopendrium Ham.; Don, Prod. Fl. Nepal. 1. 1825 = P. excavatum var.
(Pl) **Scortechinii** Bak. Ann. Bot. **5.** 477. 1891 — Perak. — Pleopeltis Bedd. 1892.
Scottii Bedd. Ferns br. Ind. t. 345. 1870 = Dryopteris hirtipes var.
(G) **Scouleri** Hk. et Grev. Ic. Fil. t. 56. 1828. HB. 342. Chr. 84. NPfl. 312 — Amer. bor. pacif. — Goniophlebium J. Sm. 1875. Polypodium carnosum Kellogg 1861 (non Mett. 1857).
(Pl) **sculpturatum** Bak. Syn. 514. 1874 — Java.
scutifrons Boj. Hort. Maur. 417. 1837 = P. phymatodes.
(P) **sechellarum** Bak. Syn. 508. 1874 — Ins. Sechellae. Mauritius.
sectifrons Kze.; Mett. Pol. 99. n. 181 t. 2 f. 3-4. 1857 = P. angustum var.
secundum Wall. List n. 301. 1828 = Dryopteris gongylodes.
(P) **segregatum** Bak. Syn. 510. 1874 — Ecuador.
(S) **selliguea** Mett. Pol. 111 n. 214. 1857. Chr. 108. NPfl. 316 — Malesia. Ins. Philipp. Queensland. — Grammitis membranacea Bl. Enum. 118. 1828 (non Polypodium Don 1825); Selliguea Bl. Fl. Jav. Fil. 123 t. 52 f. 2. 1828; Colysis Pr. 1849; Gymnogramme Hk. 1864. HB. 388 -- G. Sayeri F. Muell. et Bak. 1887; Grammitis F. Muell. 1889. (Queensland).

(Pl) **selligueoides** Bak. Syn. 513. 1874 — N. Caledonia.
(P) **semiadnatum** Hk. Ic. pl. t. 948. 1854. HB. 335 — Ecuador.
semibipinnatum Wall. List n. 2229. 1829 = Aspidium.
semicordatum Sw. Prod. 132. 1788 = Cyclopeltis.
(P) **semihirsutum** Kl. Linn. **20**. 379. 1847 — Jamaica. Guatemala-Peru.
semipinnatifidum Mett. Pol. 80 n. 130. 1857 = P. glaucophyllum var.
semipinnatum Roxb. Calc. Journ. **4**. 486. 1844 — »Malesia«.
semisagittatum Roxb. Calc. Journ. **4**. 491. 1844 = Dryopteris arida.
(Pl) **senanense** Maxim. Mél. biol. **12**. 571. 1886 — Japonia.
(P) **senile** Fée, 7 mém. 60 t. 25 f. 1. 1857 — Columbia-Ecuador.
sepultum Klf. Enum. 104. 1824 = P. lepidopteris.
(P) **sericeo-lanatum** Hk. sp. **4**. 221. 1862. HB. 335 — Columbia-Ecuador.
sericeum Hk. sp. **4**. 258. 1862 = Dryopteris pubescens var.
serpens Forst. Prod. 81. 1786 = Cyclophorus.
serpens Sw. Prod. 131. 1788 = P. Swartzii.
serra Sw. Prod. 132. 1788 = Dryopteris.
serra Wall. List n. 313. 1828 = P. decorum.
serraeforme J. Sm. JoB. **3**. 394. 1841 = Davallia alata.
serratifolium Diels, Nat. Pfl. **1**[4]. 312. 1899 = P. subauriculatum var.
serratum Aublet, Hist. Pl. Guian. **2**. 962. 1775 = Asplenium sp. vel Leptochilus guianensis.
serratum Vell. Fl. flum. **11**. t. 77. 1827; Arch. Mus. Nac. Rio Jan. **5**. 449. 1881 = Dryopteris serra.
serratum Blanco, Fl. Filip. 827. 1837 — Ins. Philipp.
serratum (Willd. sp. **5**. 173. 1810); Futó, Hedwigia **44**. 106-111 t. 3. 1905 = P. vulgare*.
serricula Fée, Gen. 238. 1850-52; 6 mém. 9 t. 7 f. 1. 1853 = P. trichomanoides.
serrulatum Sw. Schrad. Journ. **1800**[2]. 25. 1801 = Dryopteris.
(P) **serrulatum** (Sw.) Mett. Fil. Lips. 30. 1856. Pol. n. 4. HB. 323 (pt.). Chr. 79. NPfl. 308; Hier. Hedw. **44**. 80. 1905 — Amer. trop.-Juan Fernandez. Guinea. Madagascar. Ins. Mascar. Ins. Amsterdam. ? Ceylon. ? Ins. Hawaii. Acrostichum Sw. Prod. 128. 1788; Grammitis Sw. 1801; Asplenium Sw. 1806; Gymnopteris Bernh. 1806; Xiphopteris Klf. 1824; Micropteris Desv. 1827. Xiphopteris extensa Fée 1866 — Micropteris orientalis Desv. 1827; Xiphopteris Fourn. 1875 (Ins. Mascar. Amsterdam) — ? Polypodium minimum Brack. 1854; Hier. Hedw. **44**. 86. 1905 (non Aublet 1775, Ins. Hawaii; f. Hier. sp. bona).
sertularioides J. Sm. JoB. **3**. 394. 1841 = P. Hieronymusii.
sertularioides Bak. JoB. **1876**. 12 = P. tenuisectum.
sesquipedale Wall. List n. 275. 1828 = P. excavatum.
sessifolium Desv. Prod. 238. 1827 — Peru.
sessile Klf.; Sieb. exsicc.; Kze. Bot. Zeit. **1848**. 116 = P. punctatum.
G) **sessilifolium** Liebm. Vid. Selsk. Skr. V. **1**. 192 (seors. 40). 1849. HB. 512 — Mexico. — Goniophlebium serratum Fée 1857.
sessilifolium Hk. sp. **4**. 168 t. 272 A. 1862 = P. subevenosum var.
sessilifolium Hk. sp. **4**. 251. 1862 = Dryopteris cruciata.
setiferum Forsk. Fl. Aegypt. arab. 185. 1775 = Polystichum aculeatum.
(Gr) **setigerum** Bl. Enum. 123. 1828. HB. 320. Chr. 78. NPfl. 308 — Singapore. Java. Grammitis fasciculata Bl. Fl. Jav. Fil. 112 t. 47 f. 2; Polypodium Pr. 1836.
setigerum Hk. et Arn. Bot. Beech. Voy. 103 t. 21 A. 1832 = P. Hookeri

setosum Thbg. Fl. Jap. 337. 1784 = Polystichum varium.
setosum Forst. Prod. 82. 1786 = Polystichum hispidum.
setosum Pr. Tent. 180. 1836 = P. diplosorum var.
setosum Mett. Pol. 33. n. 6. 1857 = P. micropteris.
sharpianum Bak. JoB. **1880**. 369 = Dryopteris.
Sheareri Bak. JoB. **1875**. 201 = Cyclophorus.
(Pl) **shensiense** Christ, Nu. Giorn. bot. It. n. s. **4**. 99 t. 3 f. 2. 1897 — China. [Japonia.
(P) **Sherringii** Bak.; Jenm. JoB. **1882**. 326 — Jamaica.
(P) **sibarongae** Christ, Ann. Jard. Buit. II. **4**. 36. 1904 — Celebes.
(P) **siccum** Lindm. Ark. för Bot. **1**. 234 t. 11 f. 4. 1903; Hedwigia **43**. 309. 1904 — Brasilia.
signatum Blanco, Fl. Filip. ed. II. 572. 1845 — Ins. Philipp.
Pteris Blanco, Fl. Filip. ed. I. 830. 1837.
Sieberianum Klf.; Spr. Syst. **4**. 56. 1827 = Dryopteris cyatheifolia.
siifolium Willd. sp. **5**. 196. 1810 = Aspidium.
siifolium Goldm. Nova Acta **19**. Suppl. I. 454. 1843 = ?Drynaria quercifolia.
(P) **sikkimense** Hier. Hedwigia **44**. 97. 1905 — Sikkim.
P. trichomanoides Bedd. 1865 et auctt. quoad pl. sikkim.
silvaticum vide sylvaticum.
(P) **simile** L. Syst. Nat. ed. X. **2**. 1325. 1759; ?Jenm. Bull. Dep. Jam. **4**. 126. 1897 — Jamaica.
simile Lour. Fl. Cochinch. **2**. 828. 1790 — Cochinchina.
simplex Burm. Fl. Ind. 235. 1768 — Amboyna.
simplex Sw. Schrad. Journ. **1800**^{2}. 19. 1801 = P. excavatum.
simplex Lowe, Ferns **2** t. 41 B. 1858 = Dryopteris.
simplex Bak. JoB. **1876**. 12 = P. graminellum.
simplicifolium Salisb. Prod. 403. 1796 = P. phyllitidis.
simplicifolium Hk. sp. **5**. 2. 1863 = Dryopteris canescens*.
(Pl) **simplicissimum** F. Muell. Fragm. **7**. 120. 1870; 156. 1871. HB. 513 — Queensland. — P. lanceola F. Muell. Fragm. **7**. 120. 1870 (non Mett. 1861); Pleopeltis Bail. 1874.
sinense Christ, Bull. Soc. Fr. **52** Mém. I. 16. 1905 = P. chinense.
sinicum Christ, Bull. Boiss. **7**. 3. 1899 = P. subfalcatum.
(P) **Sintenisii** Hier. Hedwigia **44**. 101. 1905 — Ind. occ.
(Pl) ?*sinuatum* Pr. Rel. Haenk. **1**. 21. 1825 — Ins. Philipp. — Phymatodes Pr. 1836.
(My) **sinuosum** Wall. List n. 2231. 1829 (nomen); Hk. sp. **5**. 61 t. 284. 1863. HB. 355. Chr. 112. NPfl. 321; Yapp, Ann. Bot. **16**. 207. 1902 — Malacca. Malesia. Melanesia. — Pleopeltis Bedd. 1865; Phymatodes J. Sm. 1877.
(P) **Skinneri** Hk. sp. **4**. 214 t. 276 B. 1862. HB. 331 — Guatemala.
P. myriolepis Christ 1896.
Sloanei Poeppig; Spr. Syst. **4**. 59. 1827; Kze. Linn. **9**. 51. 1834 = Dryopteris ampla.
Sloanei Liebm. Vid. Selsk. Skr. V. **1**. 207 (seors. 55). 1849 = Dryopteris [Galeottii.
Sloani Desv. Prod. 238. 1827 = Dryopteris reptans.
Smithianum Hew. Mag. Nat. Hist. II. **2**. 459. 1838 — Jamaica. Trinidad.
(Pl) *Smithianum* Mett. Pol. 114. n. 220 b. 1857 — Ins. Philipp.
(P) **solidum** (Kze.) Mett. Pol. 53. n. 61 t. 1 f. 1-3. 1857. HB. 324. NPfl. 309 Java. Ins. Philipp. — Ctenopteris Kze.; Zoll. Syst. Verz. 37. 1854.
solutum Kl. Linn. **20**. 399. 1847 = P. angustifolium.

sophoroides Thbg. Tr. Linn. Soc. **2**. 341. 1794 = Dryopteris.
(Pl) **soridens** Hk. sp. **5**. 61 t. 283 B. 1863. HB. 355 — Borneo.
(P) **sororium** H.B.Willd. sp. **5**. 191. 1810. Mett. n. 84. HB. 337. Chr. 93. NPfl. 311 — Ind. occ. Mexico-Peru. — P. dissimile Gris. 1864 (an L. 1759?).
(Pl) **Soulieanum** Christ, Bull. Soc. Fr. **52** Mém. I. 15. 1905 — China.
sparsiflorum Hk. sp. **5**. 92. 1863 = Aspidium.
sparsisorum Desv. Berl. Mag. **5**. 315. 1811; Prod. 233. 1827 = Drynaria.
(Pl) *spathulaefolium* Bl. Enum. 134. 1828; Fl. Jav. Fil. 171 t. 78 A — Java.
(Gr) **spathulatum** Cordem. Bull. soc. sci. et arts Réunion **1890-91**; Fl. Réun. 84. 1895 — Réunion.
speciosum Bl. Enum. 132. 1828 = Drynaria rigidula.
speciosum Meyen, Reise um die Erde **1**. 108. 1834 = Alsophila.
speciosum Christ, Farnkr. 121. 1897 = Photinopteris.
spectabile Klf. Enum. 121. 1824 = Dryopteris subincisa.
(Pl) **spectrum** Klf. Enum. 94. 1824. HB. 361. Chr. 110. NPfl. 317 — Ins. Hawaii. Phymatodes Pr. 1836; Drynaria J. Sm. 1841; Pleopeltis Moore 1862; Colysis J. Sm. 1866; Polypodium Thouinianum Gaud. 1827; Drynaria [Fée 1850-52.
Speluncae L. sp. **2**. 1093. 1753 = Microlepia.
sphaerocephalum Wall. List n. 272. 1828 = Cyclophorus angustatus.
sphaeropteroides Bak. Kew Bull. **1895**. 55 = Dryopteris.
sphaerostichum Mett. Pol. 130. n. 265 t. 3 f. 13. 1857 = Cyclophorus.
(C) **sphenodes** Kze.; Kl. Linn. **20**. 402. 1847. Mett. n. 144. HB. 348. NPfl. 314 — Guatemala-Peru. — Campylonevron Fée 1850-52.
spinosum L. sp. **2**. 1092. 1753 = Pteris.
spinosum Schrank, Baier. Fl. **2**. 424. 1789 = Dryopteris spinulosa.
spinulosum Burm. Fl. Ind. 233 t. 67 f. 1. 1768 — Java. — (vix filix).
spinulosum Müll. Fl. dan. t. 707. 1777 = Dryopteris.
spinulosum Bak. Ann. Bot. **5**. 459. 1891 = Dryopteris acutidens.
spissum Bory; Willd. sp. **5**. 146. 1810 = Cyclophorus.
(P) *spissum* Fée, 8 mém. 85. 1857 — Columbia.
Spixianum Mart.; Mett. Pol. 57. n. 71. 1857 = P. meridense.
splendens Hk. sp. **5**. 52. 1863 = Cyclophorus.
splendens Hk. sp. **5**. 95. 1863 = Dryostachyum.
splendidum Klf. Enum. 112. 1824 = Dryopteris grandis.
sporadocarpum Willd. sp. **5**. 171. 1810 = P. aureum *areolatum var.
(P) **sporadolepis** (Kze.; Mett. Pol. 67. part. 1857.) Eat.; Hk. sp. **4**. 212. 1862. HB. 336 — Venezuela-Ecuador. — (nomen malum; verisim. non Mett.)
sporadolepis Mett. Pol. 67 n. 96. 1867. α. t. 1 f. 11-12 = P. Tweedianum.
— — — — β. t. 1 f. 13 = P. murorum.
— — — — γ. t. 1 f. 14 = ? P. monosorum.
Sprengelii Desv. Prod. 239. 1827 = Dryopteris megalodus.
(Gr) **Sprucei** Hk. 2 Cent. t. 10. 1860. HB. 319. NPfl. 308 — Guatemala-Peru. Grammitis J. Sm. 1875.
(S) **spurium** Mett. Ann. Lugd. Bat. **2**. 226. 1866 — Celebes. Gymnogramme Bak. 1868. HB. 388.
squalidum Vell. Fl. flum. **11** t. 76. 1827; Arch. Mus. Nac. Rio Jan. **5**. 449. 1881 = P. polypodioides.
(M) **squamatum** L. sp. **2**. 1086. 1753. Mett. n. 102. HB. 346; Kuhn, Abh. Nat. Ges. Halle **11**. 22. Chr. 90 — India occ. Mexico-Peru. Marginaria Pr. 1836; Lepicystis J. Sm. 1856; Goniophlebium Moore

1862; Drynaria Fée 1866; Pleopeltis J. Sm. 1875 — Goniophlebium lepidotrichum Fée 1857.

squamatum Phil. Linn. **29**. 107. 1857; Anal. Univ. chil. 94. 357. 1896 = P. atacamense.

(Pl) **squamulosum** Klf. Enum. 89. 1824. Mett. n. 170 — Mexico-Peru-Brasilia. Argentina. — Pleopeltis Pr. 1836; Phlebodium J. Sm. 1846; Anapeltis J. Sm. 1857. Polypodium lycopodioides pt. HB. 357.

squarrosum Roxb. Calc. Journ. **4**. 494. 1844 = Polystichum lobatum var. 4.

Gr) **Stanleyanum** Bak. JoB. **1890**. 107. NPfl. 308 — N. Guinea.

(Pl) **Steerei** Harr. Journ. Linn. Soc. **16**. 32. 1877 — Formosa.

stegnogrammoides Bak. Syn. 317. 1867 = Dryopteris.

stellatum Vahl, Symb. **3**. 104. 1794 = Cyclophorus serpens.

(P) **stenobasis** Bak. Ann. Bot. 8. 130. 1894 — Sumatra.

stenolepis Bak. Kew Bull. **1898**. 231 = Dryopteris.

(G) **stenoloma** Eat. Proc. Amer. Acad. **8**. 618. 1873 — Mexico.

(Pl) **stenophyllum** Bl. Enum. 124. 1828; Fl. Jav. Fil. 135 t. 55 f. 1. Mett. n. 180. HB. 354 — Penang. Perak. Malesia. Luzon. Fiji.
Drynaria J. Sm. 1841; Pleopeltis Moore 1857; Phymatodes J. Sm. 1861; Paragramma J. Sm. 1875 — Polypodium Morgani Zeiller 1885.

(P) **stenopteris** Bak. JoB. **1879**. 43; Christ, Ann. Jard. Buit. II. **5**. 122 t. 3 [f. 4. 1905 — Borneo.

(P) **stenopteron** Bak. JoB. **1886**. 183. NPfl. 309 — Fiji.

stenopteron Bak. JoB. **1888**. 229 = Dryopteris Dielsii.

(Pl) **Stewartii** (Bedd.) Clarke, Tr. Linn. Soc. II. Bot. **1**. 563. 1880. HB. 513 — Himalaya. — Pleopeltis Bedd. Ferns br. Ind. t. 204. 1866.

sticticum Mett. Pol. 128. n. 260. 1857 = Cyclophorus.

(Pl) **stigmaticum** Pr. Rel. Haenk. **1**. 20 t. 3 f. 2. 1825. Mett. n. 176. HB. 357 — Columbia-Brasilia. — Pleopeltis Pr. 1836; Anapeltis J. Sm. 1866. Polypodium venosum Desv. Berl. Mag. **5**. 314. 1811 (non Lour. 1790); Phlebodium Moore et Houlst. (1851); Anapeltis J. Sm. 1857; Polypodium venulosum Desv. 1827.

stigmosum Sw. Schrad. Journ. **1800**^{2}. 21. 1801 = Cyclophorus.

stigmosum auctt. quoad plant. ex Ind. bor. = Cyclophorus Beddomeanus.

stipitatum Hk. et Grev.; Hk. Bot. Misc. **2**. 239. 1831 = P. peruvianum.

stoloniferum Gmel. Syst. Nat. **2**^{2}. 1305. 1791 = Cyclophorus serpens.

stramineum Bak. Syn. 316. 1867 = Dryopteris.

(P) **streptophyllum** Bak. JoB. **1879**. 42. NPfl. 309 — Borneo. Singapore.

(P) **strictissimum** (Hk.) Hier. Engl. Jahrb. **34**. 501. 1904; Hedw. **44**. 84. 1905 Ind. occ. Columbia. Ecuador. — P. serrulatum β strictissimum Hk. sp. **4**. 175. 1862; Xiphopteris Jamesoni Hk. 2 Cent. t. 14. 1860. (HB. 323) (non Campylonevron Fée 1850-52).

struthionis L. sp. **2**. 1086. 1753. (Plum. t. 82) — S. Domingo.
Pteris Desv. 1827.

(G) **subalatum** Kl. Linn. **20**. 394. 1847 — Guiana.

(G) **subamoenum** Clarke, Tr. Linn. Soc. II. Bot. **1**. 550 t. 82 f. 2. 1880. Chr. 93 — China merid. Himalaya. — Goniophlebium Bedd. 1883.

(G) **subandinum** Sod. Cr. vasc. quit. 348. 1893 — Ecuador.

(Pl) **subaquatile** Christ, Ann. Jard. Buit. II. **5**. 124. 1905 — Borneo.

subarboreum Ham.; Wall. List n. 2226. 1829 (nomen) — ?

subarboreum Bak. Journ. Linn. Soc. **24**. 259. 1887 = Dryopteris.

(G) **subauriculatum** Bl. Enum. 133. 1828; Fl. Jav. Fil. 177 t. 83. HB. 344. Chr. 93. NPfl. 312 — Khasia?-Malesia-Samoa-Austr. trop. N. Caledonia. Goniophlebium Pr. 1836; Schellolepis J. Sm. 1866 — Polypodium pallens Bl. Fl. Jav. Fil. 178 t. 84 f. 1. 1829. Mett. n. 132; Goniophlebium Pr. 1836; Schellolepis J. Sm. 1875 — Goniophlebium serratifolium Brack. 1854; Polypodium Diels 1899. NPfl. 312.

(P) **subcapillare** Christ, Bull. Boiss. II. **5**. 1. 1905 — Costa Rica.

subcrenatum Hk. Ic. pl. t. 719. 1845 = P. moniliforme.

subdicarpon Fée, Cr. vasc. Br. **2**. 55 t. 96 f. 4. 1869 = P. moniliforme.

subdigitatum Bl. Fl. Jav. Fil. 196 t. 93. 1829 = Monachosorum.

subdimidiatum Bak. Syn. ed. I. 328. 1867 = P. phlegmaria.

(G) *subdimidiatum* Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 204 (nomen) — Yunnan.

(Pl) **subecostatum** Hk. sp. **5**. 59 t. 283 A. 1863. HB. 353 — Borneo.

(Gr) **subevenosum** Bak. Syn. 320. 1867 — Penang. Perak. Celebes. Ins. Philipp. — P. sessilifolium Hk. sp. **4**. 168 t. 272 A. 1863. HB. 322 (non Liebm. 1849); Grammitis J. Sm. 1875.

(P) **subfalcatum** Bl. Enum. 130. 1828; Fl. Jav. Fil. 186 t. 87 A, B. Mett. n. 57. HB. 328. NPfl. 310 — China merid. Ind. bor.-Malesia.
Ctenopteris Kze. 1848 — Polypodium sinicum Christ. 1897.

subfalcatum β glabrum Bedd. Ferns br. Ind. t. 189 B. 1866 = P. repandulum.

subfurfuraceum Hk. sp. **5**. 52. 1863 = Cyclophorus.

(Pl) **subgeminatum** Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 47. 1905 — N. Guinea.

(Pl) **subhastatum** Bak. JoB. **1889**. 177 — China.

(Pl) **subhemionitideum** Christ, Bull. Boiss. 7. 5. 1899 — Yunnan.

(Pl) **subimmersum** Bak. Kew Bull. **1895**. 55 — Yunnan.

subincisum Willd. sp. **5**. 202. 1810 = Dryopteris.

subintegrum Bak. JoB. **1877**. 164 = Dryopteris.

subintegrum Bak. Kew Bull. **1898**. 231 = P. oblongisorum.

sublanosum Hk. sp. **4**. 221. 1862 = P. chnoophorum.

submarginale Langsd. et Fisch. Ic. Fil. 12 t. 13. 1810 = Dryopteris.

(P) **subnudum** Mett.; Kuhn, Linn. **36**. 130. 1869. HB. 509 — Tahiti.

subobliquatum Hk. sp. **4**. 240. 1862 = Dryopteris.

subobliquatum Christ, Bull. Boiss. **6**. 197. 1898 = P. obtusissimum.

subpectinatum Wall. List n. 311. 1828 = Dryopteris otaria.

(P) **subpetiolatum** Hk. in Benth. Pl. Hartw. 54. 1840; Ic. pl. t. 391-92. Mett. n. 86. HB. 336. NPfl. 311; Maxon, Contr. U. S. Nat. Mus. **8**. 271 — Cuba. Mexico-Guatemala. — P. biserratum Mart. et Gal. 1842 — P. Cubense Fée 1850-52.

(P) **subpinnatifidum** Bl. Enum. 129. 1828. Mett. n. 24. HB. 324 — Java. Perak. Grammitis Bl. Fl. Jav. Fil. 118 t. 49 f. 2. 1828.

(P) **subpinnatum** Bak. Journ. Linn. Soc. **15**. 419. 1876 — Madagascar.

(Gr) **subpleiosorum** Racib. Pter. Buit. 89. 1898 — Java.

(P) **subrepandulum** Christ, Ann. Jard. Buit. II. **5**. 119. 1905 — Borneo.

(Pl) **subrostratum** C. Chr. Ind. 567. 1906 — Himalaya. China merid.
P. rostratum Hk. Ic. pl. t. 953. 1854. HB. 353 (non Burm. 1768 nec Cav. 1802 nec alior.); Pleopeltis Bedd. 1867; Phymatopsis J. Sm. 1875.

(P) **subscabrum** Kl. Linn. **20**. 377. 1847. Mett. n. 33 — Columbia-Ecuador. Ctenopteris J. Sm. 1875.

subscabrum Hk. sp. **4**. 183 t. 274 A. 1862 = P. ecuadorense.

(P) **subsecundo-dissectum** Zoll. Syst. Verz. 38, 48. 1854; Mett. Ann. Lugd. Bat. **2.** 220. 1866. HB. 328. NPfl. 310 — Java. N. Caledonia.

(Gr?) **subselligneum** Bak. JoB. **1890**. 107 — N. Guinea.

subserratum Hk. sp. **4**. 202. 1863 = Diplazium porphyrorachis.

subserratum Christ, Verh. Nat. Ges. Basel **11**. 211. 1895 = P. papillosum.

(P) **subsessile** Bak. Syn. 329. 1867. Chr. 82 — Amer. trop. — P. pteropus Hk. sp. **4**. 192 t. 275 B. 1862 (non Bl. 1828); Ctenopteris J. Sm. 1875. Polypodium euchlorum Kze.; Hier. 1904 (non Sod. 1883).

subsimile Col. Tr. N. Zeal. Inst. **20**. 238. 1888 = Dryopteris pennigera.

(Pl) **subsparsum** Bak. JoB. **1880**. 215 — Sumatra.

(Gr) **subspathulatum** Brack. Expl. Exp. **16**. 3 t. 1 f. 1. 1854. (HB. 320) — Tahiti.

subtetragonum Link, Hort. Berol. **2**. 105. 1883 = Dryopteris tetragona.

(P) **subtile** Kze.; Kl. Linn. **20**. 375. 1847. Mett. n. 42. HB. 326 — Jamaica. Venezuela-Peru. — Ctenopteris J. Sm. 1875 — Polypodium albopunctatum Bak. 1877. NPfl. 310 (non Raddi 1819).

subtriphyllum Hk. et Arn. Bot. Beech. Voy. 256 t. 50. 1836-40 = Aspidium.

subtripinnatum Clarke, Tr. Linn. Soc. II. Bot. **1**. 545 t. 80 f. 1. 1880 = Dryopteris Boryana.

subtruncatum Bory, Bél. Voy. Bot. **2**. 32. 1833 = Dryopteris truncata.

subulatum Vell. Fl. flum. **11**. t. 67. 1827; Arch. Mus. Nac. Rio Jan. **5**. 447. 1881 = ? Dryopteris opposita.

subvillosum Moore, Ind. 308. 1861 (nomen) = Dryopteris sp.

(Gr) **sucklingianum** Bak. Ann. Bot. **8**. 128. 1894 — N. Guinea.

(P) **sumatranum** Bak. JoB. **1880**. 214 — Sumatra.

sumatranum Bak. Ann. Bot. **8**. 131. 1894 = P. sundense.

(Pl) **sundense** C. Chr. Ind. 568. 1906 — Sumatra. P. sumatranum Bak. Ann. Bot. **8**. 131. 1894 (non 1880).

(Pl) **superficiale** Bl. Enum. 123. 1828; Fl. Jav. Fil. 136 t. 56 f. 1. HB. 355. Chr. 104. NPfl. 315 — Japonia. Korea. Tsus Sima. China. India bor.-Java. Queensland. — Pleopeltis Bedd. 1865; Colysis J. Sm. 1875; Polypodium hymenodes Kze. 1850. Mett. n. 223 — var. australiense Bail. Queensl. Bot. Bull. **4**. 21 t. 4. 1891 (Queensland).

(P) **suprasculptum** Christ, Bull. Boiss. II. **5**. 3. 1905 — Costa Rica.

surinamense Jacq. Coll. **3**. 285 t. 21 f. 4. 1789 = P. lycopodioides*.

(G) **surucuchense** Hk. Ic. pl. t. 69. 1837. HB. 343. NPfl. 312 — Ind. occ. Costa Rica-Ecuador. — Goniophlebium Moore 1857 — Polypodium andinum Karst. 1861 (non Hk. 1861).

(P) **suspensum** L. sp. **2**. 1084. 1753. Mett. n. 67. HB. 329. Chr. 80. NPfl. 310 Amer. trop.

Swartzii Wikstr. Vet. Ak. Handl. **1825**. 436. 1826 = Dryopteris vivipara.

(Pl) **Swartzii** Bak. Syn. 357. 1868. HB. 357. Chr. 104 — Florida. India occ. Guiana. Rio. — P. serpens Sw. Prod. 131. 1788. Mett. n. 171 (non Forst. 1786); Marginaria Pr. 1836; ? Pleopeltis Kze.; Pr. 1836; Phlebodium J. Sm. 1841; Craspedaria Fée 1850-52; Goniophlebium Fée 1850-52; Anapeltis J. Sm. 1857; Drynaria Fée 1872-73; Phymatodes Und. 1900 — Polypodium runcinatum Desv. 1827 — P. exiguum Hew. 1838; Phymatodes Und. 1903.

sylvaticum Schkuhr, Kr. Gew. **1**. 22 t. 8 b. 1806 = Drynaria quercifolia.

sylvaticum Col. Tasm. Journ. **2**. 163. 1844; Hk.fil. Fl. N. Zeal. **2**. 41. t. 81. 1854 = Polystichum vestitum*.

sylvaticum Mett. Novara Reise Bot. **1**. 215. 1870 = P. euryphyllum.
sylvicolum Bak. JoB. **1881**. 205 — Dryopteris.
(G) **synammia** (Fée) C. Chr. Ind. 328. 1905; 569. 1906 — Chile. Mexico. Goniophlebium Fée 1853; Polypodium trilobum Cav. Descr. 246. 1802. Mett. n. 105. HB. 341. Chr. 92. NPfl. 312 (non Houtt. 1783); Grammitis Desv. 1827; Synammia Pr. 1836; Mecosorus Kl. 1847; Goniophlebium Moore 1857 — Polypodium rachipterygium Liebm. 1849; Goniophlebium Moore 1862 (Mexico).
(Gr) **synsorum** Bak. Journ. Linn. Soc. **16**. 203. 1877 — Madagascar.
(G) **Tachiroanum** Luerss. Engl. Jahrb. **4**. 362. 1883 — Japonia.
taeniatum Sw. Schrad. Journ. **1800**[2]. 26. 1801 — Java.
(P) **taenifolium** Jenm. Bull. Dep. Jam. **4**. 114. 1897 — Jamaica.
taeniopsis Christ, Verh. Nat. Ges. Basel **11**. 443. 1896 = P. triquetrum.
taeniosum H.B.Willd. sp. **5**. 155. 1810 = P. angustifolium.
Taenitis Roth, Nov. pl. spec. 394. 1821 — Alsophila corcovadensis.
(P) **tahitense** C. Chr. Ind. 569. 1906 — Tahiti. Samoa. — P. pleiosorum Mett.; Kuhn, Linn. **36**. 128. 1869. HB. 507 (non Hk.fil. 1847).
taiwanense Christ in Warb. Monsunia **1**. 60. 1901 = Cyclophorus.
(G) **taliense** Christ, Bull. Soc. Fr. **52** Mém. I. 13. 1905 — Yunnan.
(P) **tamariscinum** Klf. Enum. 117. 1824. Mett. n. 2. HB. 338. Chr. 88. NPfl. 311 — Ins. Hawaii. Tahiti. Batjan. — Adenophorus bipinnatus Gaud. 1824. A. Tamarisci Hk. et Grev. 1830 — A. tripinnatifidus Gaud. 1824; Polypodium Pr. 1836; Amphoradenium Gaudichaudii Desv. 1827.
tanacetifolium Hoffm. Deutschl. Fl. **2**. 8. 1795 = Dryopteris spinulosa*.
(Gr) **tanalense** Bak. Kew Bull. **1897**. 300 — Madagascar.
tarapotense Bak. Syn. 505. 1874 = Dryopteris.
Tatei Bak. Syn. 506. 1874 = Aspidium.
(Pl) **tatsienense** Franch. et Bureau; Christ, Bull. Soc. Fr. **52** Mém. I. 19. 1905 — China.
(P) **taxifolium** L. sp. **2**. 1086. 1753. ? HB. 332. NPfl. 310 — India occ. Guatemala-Argentina. — P. gratum Fée 1869 — P. venustum Desv. 1811. Mett n. 50. P. Trichomanes St. Hil. 1833 — conf. Lindm. Ark. för Bot. **1**. 233.
(P) **taxodioides** Bak. JoB. **1879**. 42 — Borneo.
(M) **tectum** Klf. Enum. 87. 1824 — Brasilia. — Goniophlebium J. Sm. 1840. Polypodium aurisetum Raddi 1825. Mett. n. 168; Marginaria Pr. 1836; Craspedaria Fée 1850-52; Goniophlebium Brack. 1854; Polypodium piloselloides γ HB. 340 — Craspedaria Gestasiana Fée 1853; Polypodium [Mett.; Salom. 1883.
tenellum Forst. Prod. 81. 1786 = Arthropteris.
tenellum Don, Prod. Fl. Nepal. 2. 1825 = P. Donianum.
tenericaule Wall. List n. 335. 1828 = Dryopteris setigera.
tenerifrons Hk. sp. **5**. 104. 1864 = Aspidium tenerifrons et A. gaboonense.
tenerrimum Poir. Enc. **5**. 536. 1804 — ? (= Cystopteris fragilis?).
tenerum Roxb. Calc. Journ. **4**. 490. 1844 — Sylhet.
tenue Hoffm. Deutschl. Fl. **2**. 9. 1795 = Cystopteris fragilis.
tenuicauda Hk. sp. **5**. 90. 1863 = P. Wallichianum var.
tenuiculum Fée, Gen. 239. 1850-52. HB. 509 (pt.) — Ind. occ. Ecuador. P. lasiolepis Mett. 1869. HB. ed. II. 330.
tenuiculum Fée, Cr. vasc. Br. **1**. 86. 1869. HB. 509 (pt.) = P. brevistipes.
(P) **tenuifolium** H.B.Willd. sp. **5**. 185. 1810. Mett. n. 55. HB. 334. NPfl. 310 — Ind. occ. Columbia. Brasilia.

P. Otites Gris. 1864 (an L. 1753?) — P. camptonevron Fée 1850-52.

(Pl) **tenuilore** (J. Sm.) Kze.; Mett. Pol. 86 n. 150 t. 1 f. 28. 1857. HB. 354. NPfl. 315 — Ins. Philipp.
Drynaria J. Sm. JoB. **3**. 397. 1841 (nomen); Pleopeltis Moore 1857; Colysis J. Sm. 1875 — ? Diblemma samarensis J. Sm. 1841 (non Polypodium Mett. 1857); Taenitis Mett. 1856; Colysis J. Sm. 1875.

(P) **tenuisectum** Bl. Enum. 134. 1828; Fl. Jav. Fil. 189 t. 88 A. HB. 338. Chr. 88. NPfl. 311 — Perak. Java. Samoa. — Ctenopteris J. Sm. 1875 — Polypodium sertularioides Bak. 1876 (Samoa).

tenuisectum Mett. Fil. Lechl. **1**. 5 t. 2 f. 1-2. 1856; Hk. 2 Cent. t. 21. 1860 = P. myriophyllum.

(P) **Teresae** Maxon, Contr. U. S. Nat. Herb. **8**. 274 t. 61 f. 2. 1903 — Mexico.

terminale Spr.; Link, Hort. Berol. **2**. 93. 1833 = P. phymatodes.

tetragonum Sw. Prod. 132. 1788 = Dryopteris.

Thelypteris Weis, Pl. Cr. Fl. Gott. 307. 1770 = Dryopteris.

thelypteroides Desv. Berl. Mag. **5**. 317. 1811 = Dryopteris crinita.

Thomsonii Jenm. JoB. **1886**. 272 = Dryopteris.

Thouarsii Willd.; Mett. Pol. n. 238 b. 1857 = Drynaria Willdenowii.

Thouinianum Gaud. Freyc. Voy. Bot. 348 t. 5 f. 1. 1827 = P. spectrum.

(Pl) **Thurnii** Bak. Ann. Bat. **5**. 476. 1891. NPfl. 316 — Guiana.

(P) **Thwaitesii** Bedd. Ferns br. Ind. t. 188. 1866. HB. 508 — Ceylon.

(M) **thyssanolepis** A.Br.; Kl. Linn. **20**. 392. 1847. Mett. n. 101. HB. 512; Kuhn, Abh. Nat. Ges. Halle **11**. 21. Chr. 91 — Arizona-Peru. Cuba. Jamaica.
Goniophlebium Moore 1862; Lepicystis Diels 1899. NPfl. 323. Polypodium rhagadiolepis Fée 1850-52; Goniophlebium Fée 1857; Lepicystis J. Sm. 1866.

tijuccanum Raddi, Pl. Bras. **1**. 25 t. 37. 1825 = Dryopteris.

(G) **tomentellum** C. Chr. Ind. 570. 1906 — Luzon. — P. molliculum Copeland in Perkins, Fragm. 190. 1905 (non Kze.; Link 1841).

tomentosum Thouars, Fl. Trist. d. Ac. 32 t. 3. 1804 = Dryopteris.

tomentosum Bory, Bél. Voy. Bot. **2**. 31. 1833 = Dryopteris arthrothrix.

tomentosum Roxb. Calc. Journ. **4**. 483 t. 29. 1844 = Cyclophorus flocculosus.

(Pl) **tonkinense** Bak. JoB. **1890**. 266 — Tonkin.

(P) **torulosum** Bak. Journ. Linn. Soc. **16**. 204 1877. NPfl. 311 — Réunion.

torulosum Bak. JoB. **1880**. 215 = P. Bakeri.

tottum Thbg. Prod. Fl. Cap. 172. 1800 — Africa austr. — (Dryopteris sp. [dub.).

tottum Willd. sp. **5**. 201. 1810 = Dryopteris africana.

tovarense Kl. Linn. **20**. 374. 1847 = P. phlegmaria.

(P) **trachycarpum** Mett.; Kuhn, Linn. **36**. 127. 1869. HB. 506 — Tahiti.

(P) **transiens** Lindm. Ark. för Bot. **1**. 235. t. 11 f. 7. 1903 — Brasilia. Costa Rica.
P. longipes Fée, Cr. vasc. Br. **2**. 53 t. 95 f. 3. 1872-73 (non Link; Kze. 1850).

(G) **translucens** Kze. Anal. 16. 1837. HB. 342 — Juan Fernandez.
Goniophlebium Fée 1850-52; Polypodium intermedium Colla, Mem. Ac. Torino **39**. 51 t. 74. 1836 (non Willd. 1810). P. californicum Mett. 1857 n. 104 (non Klf. 1824).

(Pl) *transparens* (Pr.) Ettingsh. Denk. Ak. Wien **22**. 95 t. 18 f. 16. 1864 — Himalaya. — Pleuridium Pr. hb.

trapezoides Burm. Fl. Ind. 233 t. 66 f. 2. 1768 = Pteris longifolia v. vittata.

trapezoides Sw. Fl. Ind. occ. **3**. 1647. 1806 = ? Polystichum viviparum.

(Pl) **Treubii** Christ, Ann. Jard. Buit. II. **5**. 121. 1905 — Borneo.

triangulare Gmel. Syst. Nat. **2**^{2}. 1309. 1791 = Polystichum triangulum.

(P) **triangulare** Scort.; Bedd. JoB. **1887**. 324 t. 278 f. 1 — Perak.
triangulum L. sp. **2**. 1088. 1753 = Polystichum.
trichodes Reinw.; J. Sm. JoB. **3**. 394. 1841 = Dryopteris setigera.
tricholepis Schrad. Gött. gel. Anz. **1824**. 867 — P. lepidopteris.
tricholepis Mett.; Kuhn, Linn. **36**. 139. 1869 = Cyclophorus blepharolepis.
Trichomanes St. Hil. Voy. Distr. Diam. **2**. 378. 1833 = P. taxifolium var.
(P) **trichomanoides** Sw. Prod. 131. 1788. Mett. n. 26. HB. 326. Chr. 80. NPfl. 309; Hier. Hedwigia **44**. 99. 1905 — Amer. trop.
Ctenopteris J. Sm. 1875 — Polypodium serricula Fée 1850-52.
trichomanoides Bedd. Ferns br. Ind. t. 2. 1865 et auctt. quoad pl. ind. = P. sikkimense.
trichomanoides *β* jungermannioides Hk. sp. **4**. 178. 1862 = P. ascensionense.
(P) **trichopodum** F. Muell. Tr. Roy. Soc. Victoria **1**. 40. 1889; Bak. JoB. **1890**. 107 — N. Guinea.
(P) **trichosorum** Hk. 2 Cent. t. 12. 1860. HB. 323. NPfl. 308 — Ecuador.
Ctenopteris J. Sm. 1875.
trichosorum Nadeaud, En. pl. Tahiti »n. 167.« 1873 = P. tahitense.
tricuspe Sw. Schrad. Journ. **1800**^{2}. 21. 1801 = Cyclophorus hastatus.
tridactylon Wall. List n. 315. 1828; Hk. et Grev. Ic. Fil. t. 209. 1831 = P. pteropus.
tridactylon Mett. Pol. 104 n. 196 t. 1 f. 39, 40. 1857 = P. insigne.
(M) **tridens** Kze. Farnkr. **1**. 23 t. 13 f. 1. 1840. (HB. 346) — Ecuador. Galapagos.
Pleopeltis J. Sm. 1875. ? Marginaria Cumingiana Pr. 1836 (nomen).
tridentatum Roxb. Calc. Journ **4**. 495. 1844 — Banda.
trifidum Hoffm. Röm. et Ust. Mag. **9**. 10. 1790 = Athyrium filix femina.
trifidum Wither. Bot. Arrang. ed. III. **3**. 779. 1796 = Cystopteris fragilis.
trifidum Don, Prod. Fl. Nepal. 3. 1825 = P. hastatum var.
trifidum Franch. et Sav. Enum. pl. Jap. **2**. 247. 1876 = P. senanense.
trifoliatum L. sp. **2**. 1087. 1753 = Aspidium.
(P) **trifurcatum** L. sp. **2**. 1084. 1753. Mett. n. 64. HB. 323. Chr. 82. NPfl. 308 — Ind. occ.-Peru. — Goniopteris Moore 1857. Polypodium comptoniaefolium Desv. 1811. P. scolopendrioides Hk. et Grev. 1828 (non L. 1753).
(S) **triglossum** Bak. Kew Bull. **1898**. 232 — Yunnan.
Selliguea triphylla Christ 1898.
trilobum Houtt. Hist. nat. **14**. — t. —. 1783. ed. germ. Pfl. Syst. **13**^{1}. 166 t. 98 f. 1. 1786 — Batavia.
trilobum Cav. Descr. 246. 1802 = P. synammia.
trinidadensis Jenm. Gard. Chr. III. **18**. 235. 1895 = Dryopteris.
(Pl) *triphyllum* Jacq. Coll. **2**. 284 t. 22 f. 1. 1788 — Java. Ceylon.
triphyllum Desv. Berl. Mag. **5**. 315. 1811 = Aspidium puberulum.
tripinnatifidum Pr. Tent. 178. 1836 = P. tamariscinum.
(Pl) **triquetrum** Bl. Enum. 124. 1828; Fl. Jav. Fil. 141 t. 59. Mett. n. 226. HB. 359. Chr. 108. NPfl. 315 — Malesia-Samoa. — Pleuridium J. Sm. 1866 — Polypodium rupestre Bl. 1828; Fl. Jav. Fil. 142 t. 60 f. 1-3. HB. 359 (non R Br. 1810); Drynaria J. Sm. 1841; Pleuridium Fée 1850-52; Pleopeltis Moore 1857; Phymatodes Moore 1862; Polypodium saxatile Mett. n. 227. 1857 (non Kl. 1847) — P. taeniopsis Christ 1896.
(Pl) **trisectum** Bak. Kew Bull. **1898**. 232 — Yunnan.
triseriale Sw. Schrad. Journ. **1800**^{2}. 26. 1801 = P. fraxinifolium.
triste Kze. Linn. **9**. 47. 1834 = Dryopteris.

truncatum Poir. Enc. **5**. 534. 1804 = Dryopteris.
(P) **truncicola** Kl. Linn. **20**. 374. 1847. Mett. n. 27. HB. 508. NPfl. 309 — Columbia-Guatemala. — Ctenopteris J. Sm. 1875.
(P) **truncorum** Lindm. Hedwigia **43**. 309. 1904 — Brasilia.
P. Bakeri Lindm. Ark. för Bot. **1**. 240 t. 11 f. 9. 1903 (non Luerss. 1882). — conf. P. pectinatum.
tuberculatum Bak. Ann. Bot. **5**. 456. 1891 = Dryopteris.
(C) **tucumanense** Hier. Engl. Jahrb. 22. 405. 1896 — Argentina.
(M) **Türckheimii** Christ, Bull. Boiss. II. **5**. 254. 1905 — Guatemala.
(P) **turrialbae** Christ, Bull. Soc. bot. Belg. **35** 226. 1896 — Costa Rica.
(M) **Tweedianum** Hk. Ic. pl. t. 86. 1837 — Costa Rica-Bolivia.
Goniophlebium J. Sm. 1841. Polypodium sporadolepis α Mett. n. 96.
(P) **typicum** Fée, Cr. vasc. Br. **2**. 52 t. 96 f. 2. 1872-73 — Brasilia.
tyucanum Raddi, Opusc. sci. Bol. **3**. 288. 1819 = Dryopteris tijuccana.
(M) **Ulei** Hier. in Karsten et Schenck, Vegetationsbilder II Heft 1. t. 3. 1904; Christ, Hedwigia **44**. 363. 1905 — Peru-Brasilia.
umbilicatum Poir. Enc. **5**. 528. 1804 = Dryopteris filix mas var.
umbrosum Ait. Hort. Kew. **3**. 466. 1789 = Athyrium.
?umbrosum Wall. List n. 336. 1828 = Alsophila glabra.
(P) **undosum** Bak. JoB. **1890**. 108 — N. Guinea.
undulatum Willd. sp. **5**. 155. 1810 — Tranquebar.
undulatum Fourn. Mex. pl. **1**. 75. 1872 = P. gibbosum.
unidentatum Hk. et Arn. Bot. Beech. Voy. 105. 1832 = Dryopteris.
(G) **uniseriale** C. Chr. Ind. 572. 1906 — Guiana. Columbia.
P. remotum Bak. Ann. Bot. **5**. 470. 1891 (non Desv. 1827).
unisorum Bak. Syn. 307. 1867 = Sadleria polystichoides juv.
unitum L. Syst. Nat. ed. X. **2**. 1326. 1759 = Dryopteris.
unitum Thbg. Fl. Jap. 336. 1784 = Dryopteris sophoroides.
unitum Hk. sp. **5**. 5. 1863 = Dryopteris silvatica.
(Gr) **universe** Bak. Ann. Bot. **5**. 464. 1891 — Penang.
Urbani Sod. Cr. vasc. quit. 301. 1893 = Dryopteris.
urophyllum Wall. List n. 299. 1828 = Dryopteris.
(P) **ursipes** Moritz, Linn. **36**. 135. 1869 (syn.) — Venezuela.
P. ambiguum Mett.; Kuhn, Linn. **36**. 134. 1869. HB. 510 (non Desv. 1827).
ussuriense Regel, Acta Hort. Petr. **7**. 663. 1881 = P. lineare var.
(M) **vacciniifolium** Langsd. et Fisch. Ic. Fil. 8 t. 7. 1810. Mett. n. 169. HB. 340. Chr. 100 — Amer. trop. — Marginaria Pr. 1836; Goniophlebium J. Sm. 1840; Craspedaria Link 1841; Anapeltis J. Sm. 1857; Lopholepis J. Sm. 1866; Niphobolus Keys. 1873; Lepicystis Diels 1899. NPfl. 323 — Craspedaria cordifolia Fée 1869.
vacillans Link, Hort. Berol. **2**. 97. 1833 = P. latipes.
(G) **vagans** Mett. Ann. sc. nat. V. **2**. 255. 1864 — Columbia.
(G) **valdealatum** Christ, Bull. Boiss. **7**. 4. 1899 — Yunnan.
(Pl) **validum** Copeland in Perkins, Fragm. 191. 1905 — Ins. Philipp. (Mindanao).
(G) **van Heurckii** Fourn. Mex. pl. **1**. 82. 1872 — Mexico.
variabile Mett.; Kuhn, Linn. **36**. 133. 1869 = P. heteromorphum.
varians Bl. Enum. 132. 1828 = P. albidosquamatum.
variolatum Willd. sp. **5**. 192. 1810 = Aspidium martinicense.
varium L. sp. **2**. 1090. 1753 = Polystichum.
varium Mett. Pol. 126. n. 254. 1857 = Cyclophorus.

vastum Kze. Linn. **9**. 50. 1834 = Dryopteris subincisa*.
(G) **Veitchii** Bak. Gard. Chr. n. s. **14**. 494. 1880 — Japonia.
velatum Schkuhr, Kr. Gew. **1**. t. 11 b. 1809 = P. polypodioides.
(Pl) *velatum* (Pr.) Ettingsh. Farnkr. 49 t. 34 f. 2, 9. 1865 — »Asia«. Phymatodes Pr.
velutinum Sod. Rec. 59. 1883; Cr. vasc. quit. 292. 1893 = Dryopteris decussata var.
venosum Lour. Fl. Cochinch. **2**. 828. 1790 — Cochinchina.
venosum Desv. Berl. Mag. **5**. 314. 1811 = P. stigmaticum.
venosum Ham.; Wall. List n. 2228. 1829 (nomen) — India.
venulosum Desv. Prod. 226. 1827 = P. stigmaticum.
(P) **venulosum** Bl. Enum. 128. 1828; Fl. Jav. Fil. 180 t. 85 A. Mett. n. 53. HB. 334. NPfl. 310 — Malesia. — Ctenopteris Kze. 1846. Cryptosorus Dionaea Fée 1850-52. C. elasticus Fée 1850-52.
venustum Desv. Berl. Mag. **5**. 315. 1811 = P. taxifolium var.
venustum Wall. List n. 305. 1828; Clarke, Tr. Linn. Soc. II. Bot. **1**. 565. 1880 = P. himalayense.
(M) **verapax** Christ, Bull. Boiss. II. **5**. 253. 1905 — Guatemala.
(G) **verrucosum** Wall. List n. 296. 1828; Hk. Gard. Ferns t. 41. 1862. HB. 344. Chr. 93. NPfl. 312 — Malesia-Ins. Philipp. Austr. trop.
Marginaria Pr. 1836; Goniophlebium J. Sm. 1840; Schellolepis J. Sm. 1866 — Polypodium cuspidatum Mett. 1857. n. 134 (non Don 1825).
Vescoi Drake, Fl. Polyn. franç. 307. 1893 = Dryopteris Drakei.
vesiculiferum Rich. Act. Soc. Hort. nat. Paris **1**. 114. 1792 — Cayenne. (Cyatheacearum sp.).
vestitum Forst. Prod. 82. 1786 = Polystichum.
vestitum Raddi, Opusc. sci. Bol. **3**. 288. 1819; Pl. Bras. **1**. 24 t. 36. 1825 = Dryopteris deflexa.
vestitum Hk. sp. **4**. 271. 1862 = Dryopteris subincisa*.
(C) **vexatum** Eat. Mem. Amer. Acad. n. s. **8**. 199. 1860 — Cuba. Puerto Rico. — Campylonevron cubense Fée, Gen. 259. 1850-52; 6 mém. 14 t. 3 f. 2. 1853; Polypodium Christ 1897 (non Fée 1850-52).
(M) **vexillare** Christ in Schwacke, Pl. Nov. Mineiras **2**. 21. 1900; Bull. Boiss. II. **2**. 373. 1902 — Brasilia.
(Pl) **Vieillardii** Mett. Ann. sc. nat. IV. **15**. 77. 1861. (HB. 364) — N. Caledonia. Phymatodes Fourn. 1873.
Villarii Bellardi, Mém. Ac. Turin **5**. 255. (49). 1792 = Dryopteris rigida.
(P) **villosissimum** Hk. sp. **4**. 197. 1862. HB. 324. NPfl. 309 — Afr. occ. trop.
villoso-viscidum Thouars, Fl. Trist. d'Ac. 33. 1804 = Dryopteris punctata.
villosum L. sp. **2**. 1093. 1753 = Dryopteris.
(M) **villosum** Karst. Fl. Columb. **2**. 87 t. 144. 1865-69. HB. 512; Kuhn, Abh. Nat. Ges. Halle **11**. 23 — Venezuela. Peru.
villosum Fée, Cr. vasc. Br. **2**. 54 t. 97 f. 1. 1872-73 = P. longepilosum.
(Pl) **violascens** Mett. Ann. Lugd. Bat. **2**. 227. 1866. (HB. 365) — Java.
virginianum L. sp. **2**. 1085. 1753 = P. vulgare.
(P) **viride** Gilb. Bull. Torr. Cl. **26**. 316. 1899 — N. Zealand. (an P. vulgare var. Eatoni Bak?).
viscidum Roxb. in Beats. St. Helena 319. 1816 = Dryopteris punctata.
viscidum Col. Tasm. Journ. **2**. 164. 1844 — N. Zealand.
(Pl) **vitiense** Bak. JoB. **1879**. 298 — Fiji.
Vittaria Mett. Fil. Lechl. **1**. 8. 1856 = P. crassifolium.

(Gr) **vittariifolium** C. Chr. Ind. 574. 1906 — Borneo.
P. minimum Bak. JoB. **1879**. 41 (non Aublet 1775 nec Brack. 1854).
vittarioides Wall. List n. 270. 1828; Mett. Pol. 126. n. 256. 1857 = Cyclophorus spissus.
viviparum Raddi, Pl. Bras. **1**. 22 t. 32. 1825 = Dryopteris.
Vogelii Hk. sp. **4**. 271. 1862 = Dryopteris subcoriacea.
vulcanicum Bl. Enum. 122. 1828 = P. Feei.
(P) **vulgare** L. sp. **2**. 1085. 1753. HB. 334. Chr. 83. NPfl. 311 — Europa. Ins. Atlant. Mediterr. Thibet. China. Japonia. Africa. austr. Amer. bor. Ins. Kerguelen. ?N. Zealand. — Ctenopteris Newm. 1846 — Polypodium cambricum L. 1753 — P. australe Fée 1850-52 — var. Eatoni Bak. Phil. Trans. Roy. Soc. Lond. **168**. 23. 1879 (Kerguelen). conf. P. viride Gilb. —* P. serratum (Willd. 1810) Futó, Hedwigia **44**. 106-111 t. 3. 1905 (non Aublet 1775).
vulgare var. occidentale Hk. Fl. bor. Am. **2**. 258. 1840 = P. glycyrrhiza.
vulgare var. japonicum Franch. et Sav. Enum. pl. Jap. **2**. 244. 1876 = P. Fauriei.
(C) **vulpinum** Lindm. Ark. för Bot. **1**. 245 t. 11 f. 12. 1903 — Brasilia.
(G) **Wagneri** Mett. Ann. sc. nat. V. **2**. 255. 1864 (HB. 333) — Mexico-Brasilia.
Walkerae Hk. sp. **4**. 233. 1862 = Dryopteris.
(Pl) **Wallichianum** Spr. Syst. **4**. 53. 1827 — India bor.
P. juglandifolium Don, Prod. Fl. Nepal. **3**. 1825. HB. 368. Chr. 114. NPfl. 321 (non H.B.Willd. 1810); Pleuridium J. Sm. 1857; Pleopeltis Moore 1857; Arthromeris J. Sm. 1875. Polypodium capitellatum Wall. 1828. Mett. n. 208; Phymatodes Pr. 1836; Drynaria J. Sm. 1846; Pleopeltis Bedd. 1865 — Polypodium tenuicauda Hk. 1863.
Wallichianum Mett. Pol. 111 n. 213 b. 1857 = P. scolopendrinum.
Wallichii R.Br.; Wall. List n. 287. 1828 = Dipteris.
(P) **Wallii** Bedd. Ferns br. Ind. Suppl. 20 t. 380. 1876 — Ceylon.
(Gr) **Warburgii** Christ in Warb. Monsunia **1**. 59. 1900 — Batjan.
(Pl) **Wardii** Clarke, Journ. Linn. Soc. **25**. 99 t. 43. 1889 — Bhotan.
Pleopeltis Bedd. 1892. (f. Bedd. = P. himalayense).
P) **Warmingii** C. Chr. Bot. Tidsskr. **25**. 77. 1902 — Brasilia.
(Pl) **Weinlandii** Christ, Bull. Boiss. II. **1**. 448 c. fig. 1901 — N. Guinea.
(M) **Wendlandii** Hier. Hedwigia **44**. 180. 1905 — Costa Rica. — P. costaricanum Hier. Engl. Jahrb. **34**. 530. 1904 (non P. costaricence Christ 1896).
(Pl?) **Wercklei** Christ, Bull. Boiss. II. **5**. 8. 1905 — Costa Rica.
(Gr) **Whitmeei** Bak. JoB. **1876**. 12 — Samoa.
Wiesbaueri Sod. Rec. 65. 1883 = P. falcaria.
Wightianum Wall. List n. 2222. 1829 = P. lineare.
(Pl) **Wilkesii** C. Chr. Ind. 247. 1905; 574. 1906 — Fiji. Samoa.
Drynaria alata Brack. Expl. Exp. **16**. 48 t. 6 f. 1. 1854; Phymatodes J. Sm. 1861; Pleopeltis Moore 1862; Polypodium Hk. 1863. HB. 365 (non L. 1753).
Willdenowii Bory, Ann. sc. nat. **5**. 468 t. 13. 1825 = Drynaria.
Willdenowii Bl. Fl. Jav. Fil. 156 t. 66. 1829 = Drynaria pleuridioides.
Willsii Bak. Ann. Bot. **5**. 458. 1891 = Dryopteris.
(P) **Wittigianum** (Fée) (? Christ in Schwacke, Pl. Nov. Mineiras **2**. 19. 1900); Hier. Hedw. **44**. 88. 1905 — Brasilia. — Grammitis Fée, Cr. vasc. Br. **2**. 50 t. 95 f. 1. 1872-73 — G. muscosa Fée 1872-73; Polypodium Christ 1900 (non Vahl 1807).

(Pl) **Wrayi** Bak. JoB. **1887**. 206 — Perak. — Pleopeltis Bedd. 1892.

(S) **Wrightii** (Hk.) Mett.; Diels, Nat. Pfl. **1**^{4}. 316. 1899 — Liu Kiu. Formosa. Gymnogramme Hk. sp. **5**. 160 t. 303. 1864. HB. 388; Selliguea J. Sm. 1875.

Wrightii Bak. Syn. 304. 1867 = Dryopteris Sauvallei.

(C) **xalapense** (Fée) Christ, Bull. Soc. bot. Belg. **35**. 231. 1896 — Mexico-Costa Rica. Camylonevron Fée, Gen. 258. 1850-52.

(G) **xantholepis** Harr. Journ. Linn. Soc. **16**. 36. 1877 — Peru.

(P) **xanthotrichium** Kl. Linn. **20**. 376. 1847. Mett. n. 47 — Guiana. Venezuela. P. ellipticosorum Fée 1850-52.

(Pl) **xiphias** (Moore) Bak. Ann. Bot. **5**. 478. 1891 — »Polynesia austr.«. Pleopeltis Moore, Gard. Chr. n. s. **15**. 331. 1881.

xiphophoron Kze.; Mett. Pol. 73. n. 107. 1857 = P. attenuatum var.

xiphopteroides Liebm. Vid. Selsk. Skr. V. **1**. 196 (seors. 44). 1849 = P. pilosissimum.

(P) **xiphopteroidifolium** Jenm. Gard. Chr. III. **18**. 612. 1895 — Cuba.

(L) **Yakushimae** Christ, Bull. Boiss. II. **1**. 1014. 1901 — Liu Kiu. Gymnogramme lanceolata var. minor Bak.; Mak. Bot. Mag. Tokyo **10**. 178. 1896; Loxogramme minor Mak. 1905.

(P) **yarumalense** Hier. Engl. Jahrb. **34**. 499. 1904 — Columbia.

(G) **yunnanense** Franch. Bull. Soc. Fr. **32**. 29. 1885 — Yunnan. Goniophlebium Bedd. 1892. — (an P. amoenum var.?).

Zeyheri Sieb. Fl. Mart. exs. n. 135 = P. lancolatum.

(Gr) **zeylanicum** (Fée) Mett. Pol. 38 n. 21. 1857. HB. 321. NPfl. 308 — Ceylon. Grammitis Fée, Gen. 234. 1850-52.

(Pl) **Zippelii** Bl. Fl. Jav. Fil. 172 t. 80. 1829. Mett. n. 224 — Himalaya. Java. Ins. Philipp. — Pleopeltis Moore 1862; Colysis J. Sm. 1875; Polypodium heterocarpum β HB. 360.

(Pl) **Zollingerianum** Kze. Bot. Zeit. **1846**. 422. Mett. n. 233 — Java. Celebes. P. heterocarpum Bl. Fl. Jav. Fil. 167 t. 75. 1829. HB. 360 (excl. β). Chr. 105 (non supra); Pleopeltis Moore 1857.

(Pl) **zosteriforme** Wall. List n. 280. 1828; Mett. Pol. 86 n. 149 t. 1 f. 26-27. 1857. HB. 357 — Tenasserim. — Phymatodes Pr. 1836; Pleopeltis Bedd. 1866; Colysis J. Sm. 1875. — (f. Bedd. = P. pteropus var.).

Polystichopsis J. Smith, Hist. Fil. 217. 1875 *(Lastrea §)* = Polystichum et Dryopteris.

POLYSTICHUM Roth, Röm. Mag. **2**^{1}. 106. 1799; Bernh. Schrad. Journ. **1799**^{1}. 298; Roth, Tent. Fl. Germ. **3**. 69. 1800 — (Polypodiaceae Gen. **17**).

abbreviatum Lam. et DC. Fl. franç. **2**. 560. 1805 = Dryopteris filix mas.

abbreviatum (Schrad.) (? J. Sm. Lond. JoB. **1**. 199. 1842); Pr. Epim. 58. 1849. NPfl. 193 — Brasilia. Guiana. Ecuador. — Aspidium Schrad. Gött. gel. Anz. **1824**. 869. Mett. n. 72. HB. 257. Chr. 233; Cyclodium Pr. 1849; Nephrodium Fée 1850-52; Cyrtomium J. Sm. 1877; Dryopteris O.Ktze. 1891. —* Aspidium guianense Kl. 1847. Mett. n. 83; Polystichum Pr. 1849. Peltochlaena nephrodioides Fée 1850-52.

acanthophyllum (Franch.) Christ, Bull. Soc. Fr. **52** Mém. I. 30. 1905 — India bor. China merid. — Aspidium Franch. Bull. Soc. Fr. **32**. 28. 1885.

acrostichoides (Michx.) Schott, Gen. Fil. ad t. 9. 1834. NPfl. 190 — Amer. bor. Nephrodium Michx. Fl. bor. Am. **2**. 267. 1803; Polypodium Poir. 1804; Aspidium Sw. 1806. Mett. n. 95. HB. 250. Chr. 236; Dryopteris O.Ktze. 1891.

aculeatum (L.) Schott, Gen. Fil. ad t. 9. 1834. NPfl. 191 — sp. collect. fere ubique inter reg. temp. et trop. — Polypodium L. sp. **2**. 1090. 1753 (pt.);

Aspidium Sw. 1801. HB. 250 (pt.). Chr. 237; Tectaria Lag. Garcia et Clem. 1802; Hypopeltis Todaro 1866 (β); Dryopteris O.Ktze. 1891. Aspidium angulare Kit.; Willd. 1810; Polystichum Pr. 1836.

I. *Formae Africanae.*

— 1. Polypodium ammifolium Poir. Enc. **5**. 554. 1804; Aspidium Desv. 1811; Phegopteris Fée 1850-52. Aspidium stramineum Klf.; Spr. 1827. Mett. n. 116; Hypopeltis Bory 1833; Polystichum Pr. 1836 — Hypopeltis Mauritiana Bory 1833; Polystichum Fée 1850 52 — Ins. Mascarenae.

— 2. Polystichum Schizolobium Fée, 8 mém. 99. 1857; Aspidium Kuhn 1868. Ins. Mascarenae.

II. *Formae Americanae.*

1. *Indusiatae.*

— 3. Aspidium aculeatum var. scopulinum Eat. Ferns. N. Am. **2**. 125 t. 62 f. 8. 1880; Polystichum scopulinum Maxon 1900 — Amer. bor. pacif.

— 4. Polypodium muricatum L. sp. **2**. 1093. 1753; Aspidium Willd. 1810; Kuhn, Abh. Nat. Ges. Halle **11**. 33. 1869; Polystichum Fée 1850-52. Aspidium ordinatum Kze. 1844. Mett. n. 106; Polystichum Liebm. 1849 — Aspidium Moritzianum Kl. 1847. Mett. n. 105. A. crenatum Mett. 1856. n. 114. A. Hartwegii Kl. 1847 — Polystichum grande Fée 1857 — P. rachichlaena Fée 1857 — Ind. occ.; Mexico-Venezuela.

— 5. Polystichum Mülleri (Mett.) Fourn. Mex. pl. **1**. 91. 1872 (non Schum. 1803); Aspidium Mett. hb. — Mexico.

— 6. Polystichum pallidum Fourn. Mex. pl. **1**. 91. 1872 — Mexico.

— 7. Polystichum distans Fourn. Mex. pl. **1**. 91. 1872 — Mexico.

— 8. Polystichum confluens Fourn. Mex. pl. **1**. 92. 1872 — Mexico.

— 9. Polypodium montevidense Spr. Syst. **4**. 59. 1827; Phegopteris Mett.; Salom. 1883; Aspidium Hier. 1896; Polystichum Sellowianum Pr. 1836 — Amer. austr.

—10. Aspidium subintegerrimum Hk. et Arn. Bot. Beech. Voy. 52. 1832 — Chile.

—11. Polystichum Bridgesi Schott, Gen. Fil. ad t. 9. 1834 (nomen); Aspidium Sturm 1858.

—12. Polystichum Brongniartianum Remy; Gay, Fl. Chil. **6**. 518. 1853; Aspidium Sturm 1858.

—13. Aspidium orbiculatum Desv. Berl. Mag. **5**. 321. 1811; Polystichum Gay 1853; P. orbiculare Christ 1905 — Chile-Amer. antarct.

—14. Aspidium aculeatum var. chilense Christ. Ber. Schw. bot. Ges. **3**. —. 1893; Polystichum chilense Diels 1899. NPfl. 192. (an P. vestitum var.?)

—15. Polystichum aculeolatum Fée, Cr. vasc. Br. **1**. 128 t. 38 f. 2. 1869 — Brasilia.

—16. Polystichum giganteum Fée, Cr. vasc. Br. **1**. 127 t. 39 f. 2. 1869 — Brasilia.

—17. Polystichum lanosum Fée, Cr. vasc. Br. **1**. 126 t. 40 f. 2. 1869 — Brasilia.

—18. Polystichum longecuspis Fée, Cr. vasc. Br. **1**. 129 t. 41 f. 2. 1869 — Brasilia.

—19. Polystichum microsorium Fée, Cr. vasc. Br. **1**. 125 t. 40 f. 1. 1869 — Brasilia.

—20. Polystichum platylepis Fée, Cr. vasc. Br. **1**. 129 t. 41 f. 1. 1869 — Brasilia.

(—21. Rochaleanum supra pag. 91 vide P. Rochaleanum).

—22. Tijucense Fée, Cr. vasc. Br. **1**. 127 t. 38 f. 1. 1869 — Brasilia.

2. *Exindusiatae.*

—23. Aspidium platyphyllum Willd. sp. **5**. 255. 1810; Nephrodium Desv. 1827; Polystichum Pr. 1836; Phegopteris Mett. n. 9. 1858; Polypodium Hk. 1862. HB. 310 (non Sw. 1806) — P. polystichoides Kl. 1847. Phegopteris polysti-

chiformis Fée 1850-52 — Polystichum laxum Wawra 1863; Phegopteris polystichops Wawra 1866 — Amer. trop.

—24. Polypodium rigidum Hk. et Grev. Ic. Fil. t. 163. 1829. HB. 310 (non Hoffm. 1795); Phegopteris Mett. n. 5. 1858. Nephrodium polyphyllum Pr. 1825; Polystichum Pr. 1836; Aspidium Mett. 1856 — Andes.

—25. Aspidium caudatum Sw. Vet. Ak. Handl. **1817**. 621 (non 1806); Polystichum Fée 1850-52; Aspidium Freyreissii Wikstr. 1826 — Brasilia.

—26. Polypodium cochleatum Kl. Linn. **20**. 388. 1847; Phegopteris Mett. n. 6. 1858; Polystichum Hier. 1904. Aspidium robustum Kze.: Mett. 1856 — Peru. Venezuela. — (Affine est P. Lehmannii Hier.).

—27. Jamesonia paleacea Kze. Bot. Zeit. **1844**. 739; Aspidium pycnolepis Kze.: Kl. Linn. **20**. 365. 1847; Phegopteris Mett. n. 7. 1858; Polystichum? Moore 1858; Polypodium Hk. 1862 — Aspidium gelidum Kze.: Kl. 1847; Polystichum Fée 1850-52 — Andes.

III. *Formae Australienses.*

—28. Aspidium aculeatum var. samoensis Luerss. Fil. Graeff. 174. 1871 — Samoa.

—29. Aspidium proliferum R.Br. Prod. Fl. N. Holl. 147. 1810. Mett. n. 113; Hypopeltis Bory 1833; Polystichum Pr. 1836. Aspidium radicans Sieb. exsicc.; Polystichum Pr. 1836 — Australia.

IV. *Formae Asiaticae.*

30. Aspidium biaristatum Bl. Enum. 164. 1828; Polystichum Moore 1858. — ? Phegopteris eximia Mett.: Kuhn 1869 — India. Malesia.

—31. Aspidium vulcanicum Bl. Enum. 163. 1828; Polystichum Moore 1858 — Java.

—32. Aspidium mucronifolium Bl. Enum. 164. 1828; Polystichum Pr. 1849. Aspidium congener Bl. 1828 — A. amblyotus Kze. 1848; Polystichum Pr. 1849 — Aspidium microphyllum Bl. 1828. Mett. n. 111; Polystichum Pr. 1849 — India. Malesia.

—33. Aspidium moluccense Bl. Enum. 168. 1828; Polystichum Moore 1858; P. horizontale Pr. 1849. Aspidium acutifolium Reinw. hb.; Polystichum Pr. 1849 — Ins. Moluccae.

—34. Aspidium aculeatum var. Batjanense Christ, Farnkr. 238. 1897 et in Warb. Monsunia **1**. 77. 1900 — Malesia orient. Ins. Philipp.

—35. Aspidium tacticopterum Kze. Linn. **24**. 290. 1851. Mett. n. 112; Polystichum Moore 1858 — India austr.

—36. Aspidium subinerme Kze. Linn. **24**. 290. 1851. Mett. n. 117 — India austr.

—37. Polypodium anomalum Hk. et Arn. JoB. 8. 360 t. 11. 1856; Aspidium Hk. 1862. HB. 253. (Chr. 239); Polystichum J. Sm. 1866; Dryopteris O.Ktze. 1891 — Ceylon.

—38. Polystichum aculeatum var. travancoricum Bedd. Hdb. 209. 1883 — India austr.

—39. Polystichum aculeatum var. nigropaleaceum Christ. Ber. Schweiz. bot. Ges. **3**. —. 1893; Polystichum Diels 1899. NPfl. 191 — India.

—40. Polystichum aculeatum var. japonicum Christ. Ber. Schweiz. bot. Ges. **3**. —. 1893; Polystichum Diels 1899. NPfl. 191 — Japonia.

V. *Formae Europaeae.*

—41. Aspidium hastulatum Tenore. Atti Ist. Incorr. Nap. **5**. 149 t. 4 f. 7 A. b. 1830; Hypopeltis Todaro 1866 — Europa austr.

Obs. Conf. P. lobatum. — Vide Mett. Pheg. u. Aspid. 1858 et Ann. Lugd. Bat. **1**. 226. 1864. Hk. sp. **4**. 18, 246. 1862. Kuhn, Abh. Nat. Ges. Halle **11**. 11. 1869. Clarke, Tr. Linn. Soc. II. Bot. **1**. 508. 1880. Christ, Ber. Schweiz. bot. Ges. **3**. 1893. Burck, Receuil Trav. Bot. Néerl. **1**. 33. 1904.

aculeatum Roth, Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 79. 1800 = P. lobatum.

aculeatum × lobatum C. Chr. Ind. 66. 1905; 578. 1906 — Europa centr. Aspidium Bicknellii Christ ap. Burnat, Matér p. fl. Alp. marit. — 1900; A. lobatum × aculeatum Christ 1900.

aculeatum × lonchitis (Halácsy) C. Chr. Ind. 63. 1905; 578. 1906 — Graecia. Aspidium Halácsy, Verh. zool. bot. Ges. **54**. 129 c. fig. 1904; A. lonchitiforme Hal. l. c.

aculeolatum Fée, Cr. vasc. Br. **1**. 128 t. 38 f. 2. 1869 = P. aculeatum var. 15.

acutidens Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 259 — China.

acutifolium Pr. Epim. 259. 1849 = P. aculeatum var. 33.

acutum Pr. Tent. 83. 1836 = Nephrolepis biserrata.

adiantiforme (Forst.) J. Sm. Hist. Fil. 220. 1875 — Australia. Polynesia. N. Zealand. Afr. austr. c. ins. Tristan d'Acunha. Cuba. Amer. austr.-Chile. Juan Fernandez. — Polypodium Forst. Prod. 82. 1786; Dryopteris O.Ktze. 1898. Polypodium coriaceum Sw. 1788; Aspidium Sw. 1801. Mett. n. 121; Tectaria Link 1833; Hypopeltis Bory 1833; Polystichum Schott 1834. Aspidium capense Willd. 1810 (non Polypodium L.fil. 1781). HB. 254. Chr. 242; Polystichum J. Sm. 1846. NPfl. 193 — Tectaria Calahuala Cav. 1802 — Rumohra aspidioides Raddi 1819 — Aspidium discolor Langsd. et Fisch. 1810; Polystichum Brack. 1854.

adscendens auctt. vide ascendens.

aemulum Pr. Tent. 83. 1836 = Dryopteris.

affine Pr. Tent. 83. 1836 = P. speciosum.

affine Ledeb. Fl. ross. **4**. 515. 1853 = Dryopteris filix mas.

alcicorne (Bak.) Diels, Engl. Jahrb. **29**. 194. 1900 — China. Polypodium Bak. JoB. **1888**. 229; Aspidium Christ 1901.

amabile (Bl.) J. Sm. Ferns br. and. for. 152. 1866. NPfl. 193 — Japonia. China austr. Ind. bor.-Malesia. — Aspidium Bl. Enum. 165. 1828. HB. 254. Chr. 240; Lastrea Moore 1858; Dryopteris O.Ktze. 1891. Aspidium rhomboideum Wall. 1828. Mett. n. 155; Polystichum Schott 1834.

amblyotum Pr. Epim. 55. 1849 = P. aculeatum var. 32.

amplissimum Pr. Tent. 84. 1836; Epim. 58. 1849 = Dryopteris.

andinum Phil. Linn. **29**. 108. 1857; Christ, Ark. för Bot. **4**[12]. 2. 1905 — Chile-Amer. antarct. Ins. Falkland. South Georgia.

angulare Pr. Tent. 83. 1836 = P. aculeatum.

angustatum Schum. En. pl. Saell. **2**. 22. 1803 = Dryopteris spinulosa.

anomalum J. Sm. Ferns br. and for. 151. 1866 = P. aculeatum var. 37.

apiifolium (Sw.) C. Chr. Ind. 64. 1905; 578. 1906 — Cuba. Jamaica. Guatemala. Dicksonia Sw. Schrad. Journ. **1800**[2]. 91. 1801 (non auctt.); Aspidium Mett.; Kuhn 1869; Dryopteris O.Ktze. 1891. Aspidium ascendens Hew. 1838. HB. 255. Chr. 242; Lastrea J. Sm. 1841; Polystichum Moore 1858. NPfl. 193. Nephrodium duale Donn. Smith 1890.

aquifolium Und. et Maxon, Bull. Torr. Cl. **29**. 584. 1902 = P. triangulum var.

aristatum (Forst.) Pr. Tent. 83. 1836. NPfl. 193 — Japonia. China. India-Malesia-Polynesia. Natal. — Polypodium Forst. Prod. 82. 1786; Aspidium Sw. 1801. Mett. n. 107. HB. 255. Chr. 240. Nephrodium Pr. 1825; Lastrea Moore 1858; Dryopteris O.Ktze. 1891.

— Aspidium cornu-cervi Don 1825; A. Hameltoni Spr. 1827; Hypopeltis Bory 1833; Polystichum Moore 1858.

— Aspidium assamicum Kuhn 1869. (HB. 493).

— Aspidium platypus Kze. 1848. Mett. n. 161; Lastrea Moore 1858; Nephrodium Hk. 1862. HB. 286; Dryopteris O.Ktze. 1891; Polystichum Diels 1899. NPfl. 193 — ? Arachniodes aspidioides Bl. 1828.

aristatum Hk.f. Fl. N. Zeal. **2**. t. 78. 1854 = P. lobatum var. 5.

artemisioides Goldm Nova Acta **19**. Suppl. 1. 463. 1843 — Oahu.

ascendens Moore, Ind. 86. 1858 = P. apiifolium.

Atkinsoni Bedd. Ferns br. Ind. Suppl. 14 t. 362. 1876 — Ind. bor. China austr. Aspidium Clarke 1880.

auriculatum (L.) Pr. Tent. 83. 1836. NPfl. 189 — India. Formosa. ? Ins. Philipp. Polypodium L. sp. **2**. 1088. 1753; Aspidium Sw. 1801. Mett. n. 90. HB. 251 (pt.). Chr. 237; Nephrodium Rich. 1834; Dryopteris O.Ktze. 1891. — Conf. Polypodium harpophyllum Zenk. — ? Phegopteris nervosa Fée 1850-52 (Ins. Philipp.) — (Conf. Dryopteris philippina).

Bakerianum (Atkins.) Diels, Nat. Pfl. **1**[4]. 191. 1899 — Ind. bor. Japonia. Aspidium Atkins. msc.; Bak. Hk. Ic. pl. t. 1656. 1886; A. Prescottianum var. Bakeriana Clarke, Tr. Linn. Soc. II. Bot. **2**. 510 t. 66. 1880 — A. microchlamys Christ 1899.

basipinnatum (Bak.) Diels, Nat. Pfl. **1**[4]. 189. 1899 — China austr. Aspidium Bak. JoB. **1889**. 176.

Benoitianum Gaud. Freyc. Voy. Bot. 331 t. 11. 1827 — Ins. Moluccae. Aspidium Gaud. l. c. sub t. 11; Nephrodium Moore 1858. — (Dryopteris sp. dub.).

Berterianum (Colla) C. Chr. Ind. 66. 1905; 579. 1906 — Juan Fernandez. Aspidium Colla, Mem. Ac. Torino **39**. 42 t. 70. 1836. Mett. n. 122; A. flexum Kze. 1837. HB. 255; Polystichum Gay 1853. NPfl. 193; Dryopteris O.Ktze. 1891.

biaristatum Moore, Ind. 86. 1858 = P. aculeatum var. 30.

Boottii Gaudog. Fl. Eur. **27**. 168. 1891 = Dryopteris cristata × spinulosa.

Braunii (Spenn.) Fée, Gen. 278. 1850-52. NPfl. 192 — Europa. Caucasus. Amur. China. Amer. bor. temp. Ins. Hawaii. — Aspidium Spenn. Fl. Friburg. **1**. 9 t. 2. 1825. Mett. n. 109. Chr. 238; Dryopteris Und. 1896 — Polystichum Haleakalense Brack. 1854 (non Aspidium Mann 1868) — var. Clarkei Christ, Bull. Boiss. **7**. 15. 1899 (Yunnan).

Braunii × lobatum C. Chr. Ind. 80. 1905; 579. 1906 — Europa centr. Aspidium lobatum × Braunii Luerss. Rabenh. Krypt. Fl. ed. II. **3**. 357 f. 143. 1884-89; A. Luerssenii Dörfler 1890 (non Christ 1898). — A. lobatiforme Waisb. 1899.

Bridgesi Schott. Gen. Fil. ad t. 9. 1834 = P. aculeatum var. 11.

Brongniartianum Remy; Gay. Fl. chil. **6**. 518. 1853 = P. aculeatum var. 12.

caducum Diels, Nat. Pfl. **1**[4]. 194. 1899 = P. Hookerianum.

caespitosum Schott. Gen. Fil. ad t. 9. 1834 = P. obliquum.

californicum (Eat.) Diels. Nat. Pfl. **1**[4]. 191. 1899 — Amer. bor. pacif. Aspidium Eat. Pr. Amer. Ac. **6**. 555. 1865. HB. 253; Dryopteris O.Ktze. 1891.

Callipteris Bernh. Schrad. Journ. **1799**[1]. 307 = Dryopteris filix mas, cristata, rigida, spinulosa.

Canariense Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris filix mas *oligodonta.

capense J. Sm. Bot. Mag. **72** Comp. 35. 1846 = P. adiantiforme.

capillipes (Bak.) Diels, Nat. Pfl. 1⁴. 191. 1899 — China occ.
Aspidium Bak. JoB. **1888**. 228.

caripense Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris submarginalis.

carthusianum Sanio, Verh. Bot. Ver. Brand. **25**. 84. 1884 = Dryopteris filix mas × spinulosa.

carvifolium (Kze.) C. Chr. Ind. 70. 1905; 580 1906 — Japonia. China austr. Ind. bor.-Malesia-Polynesia ad Ins. Hawaii. Australia. Natal.
Aspidium curvifolium Kze. Bot. Zeit. **1848**. 283 (err. typ.); A. carvifolium Kze. 1851 (non Bak. 1888); A. coniifolium Wall. 1828 (nomen, non Pr. 1822). Mett. n. 157. Chr. 240; Polystichum Pr. 1836 (nomen, non Schum. 1803). NPfl. 193; Lastrea Moore 1857; Dryopteris Und. 1897; Nephrodium Hope 1903; Aspidium aristatum β coniifolium. HB. 255.

carvifolium Diels, Engl. Jahrb. **29**. 194. 1900 = P. omeiense.

caryotideum Diels, Nat. Pfl. **1⁴**. 194. 1899 = P. falcatum*.

caudatum Fée, Gen. 278. 1850-52 = P. aculeatum var. 25

chaerophylloides Pr. Tent. 84. 1836 = Dryopteris pubescens var.

chilense Diels, Nat. Pfl. **1⁴**. 192. 1899 = P. aculeatum var. 14.

chlaenosticta Fée, Gen. 280. 1850-52 — Australia.

cochleatum Hier. Engl. Jahrb. **34**. 452. 1904 = P. aculeatum var. 26.

concavum Moore, Proc. Roy. Hort Soc. **2**. 377. 1862 (nomen) = P. Standishii.

confluens Fourn. Mex. pl. **1**. 92. 1872 = P. aculeatum var. 8.

conifolium Schum. En. pl. Saell. **2**. 22. 1803 = Dryopteris spinulosa.

coniifolium Pr. Tent. 84. 1836; Epim. 55. 1849 = P. carvifolium.

consanguineum Gaud. Freyc. Voy. Bot. 331. 1827 — Ins. Mariannae. (Dryopteris sp. dub.).

constrictum Marshall; Gard. Chr. III. **18**. 588. 1895 — Hort.

conterminum Gaud. Freyc. Voy. Bot. 329. 1827 = Dryopteris opposita.

coriaceum Schott, Gen. Fil. ad t. 9. 1834 = P. adiantiforme.

craspedosorum (Maxim.) Diels, Nat. Pfl. **1⁴**. 189. 1899 — Japonia. China.
Aspidium Maxim. Bull. Ac. St. Pétersb. **15**. 231. 1871. HB. 492. Chr. 235; Dryopteris O.Ktze. 1891.

cristatum Roth, Röm. Arch. **2¹**. 106. 1799; Tent Fl. Germ. **3**. 84. 1800 = Dryopteris.

cristatum × spinulosum Sanio, Verh. Bot. Ver. Brand. **25**. 82. 1884 = Dryopteris.

Cumingianum Pr. Epim. 54. 1849 — Chile.

cuspidatum Pr. Tent. 82. 1836 = Leptochilus.

cyphochlamys Fée, Gen. 279. 1850-52; 6 mém. 20 t. 3 f. 4 = P. triangulum.

cystopteroides Nees, Linn. **19**. 685 1847 = Dryopteris patula.

cystostegia (Hk.) Armstr. Tr. N. Zeal. Inst. **13**. 364. 1881 (cystostigium); NPfl. 193 — N. Zealand. — Aspidium Hk. sp. **4**. 26 t. 227. 1862. HB. 253. Chr. 241; Dryopteris O.Ktze. 1891.

davallioides Diels, Nat. Pfl. **1⁴**. 193. 1899 = Dryopteris.

decompositum Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris.

deltodon (Bak.) Diels, Nat. Pfl. **1⁴**. 191. 1899 — China centr.
Aspidium Bak. Gard. Chr. n. s. **14**. 494 1880.

denticulatum (Sw.) J. Sm. JoB. **4**. 195. 1841. NPfl. 193 — Amer. trop.
Polypodim Sw. Prod. 134; Aspidium Sw. 1801. Mett. n. 159. Chr. 240; Nephrodium Fée 1850-52. HB. 287; Lastrea Moore 1858; Dryopteris O.Ktze. 1891. —* Aspidium laetum Sw. 1817. Mett. n. 158; Polystichum Pr. 1836; Lastrea Moore 1857. Aspidium Klotzschii Hk. 1854.

diaphanum (Zoll. et Moritz) Moore, Ind. 90. 1858 — Java. — Aspidium Zoll. et Moritz, Nat. en Geneesk. Arch. **2**. 203. 1845; Flora **1847**. 319. Mett. n. 96.
dictyophyllum (Mett.) C. Chr. Ind. 581. 1906 — Ecuador.
Phegopteris Mett.; Kuhn, Linn. **36**. 106 1869.
dilatatum Schum. En. pl. Saell. **2**. 24. 1803 = Dryopteris spinulosa*.
diplazioides Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 260 — China centr.
discolor Brack. Expl. Exp. **16**. 207. 1854 = P. adiantiforme.
discretum J. Sm. JoB. **3**. 413. 1841 = P. aculeatum var. 33.
discretum Diels. Nat. Pfl. **1**[4]. 199. 1899 = P. lobatum var. 3.
dissectum Bernh. Schrad. Journ. **1801**[1]. 16. 1802 = Dryopteris.
distans Fourn. Mex. pl. **1**. 91. 1872 = P. aculeatum var. 7.
drepanoides Fourn. Mex. pl. **1**. 92. 1872 — Mexico.
drepanum (Sw.) Pr Tent. 84. 1836. NPfl. 192 — Madeira.
Aspidium Sw. Schrad. Journ. **1800**[2]. 38. 1801; Polypodium Lowe 1838. HB. 310; Phegopteris J. Sm. 1857. Mett. n. 11. Chr. 276.
Dryopteris Roth, Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 80. 1800; Bernh. Schrad. Journ. **1799**[1]. 306 = Dryopteris Linnaeana.
dubium (Karst.) Diels, Nat. Pfl. **1**[4]. 194. 1899 — Columbia-Ecuador.
Phegopteris Karst. Fl. Col. **1**. 169 t. 84. 1861; Polypodium Hk. 1863 (excl. β). HB. 313. (Chr. 233). (§ Cyrtomiphlebium Hk.).
Dubrueillianum Gaud. Freyc. Voy. Bot. 333 t. 9. 1827 = Dryopteris cyatheoides.
? *duriusculum* Pr. Tent. 84. 1836 — Chile.
Nephrodium? Pr. Rel. Haenk. **1**. 38. 1825. Aspidium Sturm 1858.
Duthiei (Hope) C. Chr. Ind. 72. 1905; 581. 1906 — India bor.
Aspidium Hope, Journ. Bomb. Nat. Hist. Soc. **12** 532 t. 6. 1899.
echinatum (Gmel.) C. Chr. Ind. 83. 1905; 581. 1906 — Ind. occ.
Polypodium Gmel. Syst. Nat. **2**[2]. 1309. 1791; P. muricatum Sw. Prod. 131. 1788 (non L. 1753); Aspidium mucronatum Sw 1801. Mett. n. 91. HB. 250. Chr. 236; Polypodium Sw. 1806; Polystichum Pr. 1836. NPfl. 191; Dryopteris O.Ktze. 1891.
effusum Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris.
elegans Remy; Fée, Gen. 278. 1850-52 (nomen); Gay, Fl. chil. **6**. 514. 1853 = P. mohrioides.
emarginatum Moore. Ind. 91. 1858 = Aspidium.
falcatum Fée, Gen. 279. 1850-52 = ? P. echinatum.
falcatum (L.fil.) Diels, Nat. Pfl. **1**[4]. 194. 1899 — Japonia. China-Ind. bor. Celebes. Ins. Hawaii. Afr. austr. Madagascar. — Polypodium L.fil. Suppl. 446. 1781; Aspidium Sw. 1801. Mett. n. 74. HB. 257. Chr. 232; Cyrtomium Pr. 1836; Dryopteris O.Ktze. 1891 — Cyrtomium Fortunei J. Sm. 1866. —* Aspidium caryotideum Wall. 1828: Hk. et Grev. Ic. Fil. t. 69. 1828. Mett. n. 76; Cyrtomium Pr. 1836; Dryopteris Und. 1897; Polystichum Diels 1899. NPfl. 194. Aspidium anomophyllum Zenker 1835. Mett. n. 75.
falciculatum Gaud. Freyc. Voy. Bot. 328. 1827 = Dryopteris.
falcinellum (Sw.) Pr. Tent. 83. 1836. NPfl. 189 — Madeira.
Aspidium Sw. Syn. 46, 243. 1806. Mett. n. 94. HB. 249. Chr. 236; Dryopteris O.Ktze. 1891 — Polystichum Maderense Johns. 1866.
Filix mas Roth. Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 82. 1800 = Dryopteris.
fimbriatum Pr. Epim. 58. 1849 = Athyrium.
flagellare (Maxim.) C. Chr. Ind. 581. 1906 — Japonia. — Polypodium Maxim. msc.; Phegopteris Mak. Bot. Mag. Tokyo **9**. - . 1895; Ptilopteris Mak. 1899.
flexum Gay, Fl. chil. **6**. 517. 1853 = P. Berterianum.

foeniculaceum (Hk.) J. Sm. Hist. Fil. 220. 1875 — Sikkim.
Aspidium Hk. sp. **4**. 36 t. 237. 1862. HB. 256. Chr. 241: Lastrea Bedd. 1865; Diacalpe Clarke 1880. NPfl. 159; Dryopteris O.Ktze. 1891.
Obs. vide Bedd. JoB. **1888**. 234.
formosissimum Goldm. Nova Acta **19** Suppl. I. 463. 1843 = P. aristatum.
fragrans Ledeb. Fl. Ross. **4**. 514. 1853 = Dryopteris.
Franchetii Christ, Bull. Soc. Fr. **52** Mém. I. 28. 1905 — China.
fraxinellum (Christ) Diels, Nat. Pfl. **1**. 494. 1899 — China austr.
Aspidium Christ. Bull. Boiss. **7**. 15. 1899: Cyrtomium Christ 1902.
frondosum J. Sm. Cat. cult. Ferns 60. 1857 = P. Webbianum.
fuscipes Pr. Tent. 83. 1836 = Aspidium (Wall.).
gelidum Fée, Gen. 278. 1850-52 = P. aculeatum var. 27.
giganteum Fée, Cr. vasc. Br. **1**. 127 t. 39 f. 2. 1869 = P. aculeatum var. 16.
glaciale Christ, Bull. Soc. Fr. **52** Mém. I. 28. 1905 — China.
? glandulosum Pr. Tent. 82. 1836 = Adenoderris viscidula.
goggilodus Gaud. Freyc. Voy. Bot. 326. 1827 = Dryopteris gongylodes.
Goldieanum Keys. Pol. Cyath. Hb. Bung. 44. 1873 = Dryopteris.
grande Fée, 8 mém. 98. 1857 = P. aculeatum var. 4.
? Grevilleanum Pr. Tent. 82. 1836 = Fadyenia Fadyenii.
guadalupense Fée, 11 mém. 74 t. 19 f. 2. 1866 — Guadeloupe.
guianense Pr. Epim. 58. 1849 = P. abbreviatum*.
gymnogrammoides Gaud. Freyc. Voy. Bot. 324. 1827 = Dryopteris polypo-[dioides.
Haenkeanum Pr. Tent. 83. 1836 = P. mohrioides.
Haleakalense Brack. Expl. Exp. **16**. 204 t. 28. 1854 = P. Braunii.
Hamiltoni Moore, Ind. 89. 1858 = P. aristatum var.
Hancockii (Hance) Diels, Nat. Pfl. 1^4. 191. 1899 — Formosa. — Ptilopteris Hance, JoB. **1884**. 139; Aspidium Bak. 1891; A. reductum Bak. 1888.
hecatopterum Diels, Engl. Jahrb. **29**. 193. 1900 — China.
heterodon Pr. Epim. 58 1849 = Cyclodium meniscioides.
heterolepis Fée, Gen. 279. 1850-52 = P. viviparum.
Hillebrandii Carr. in Seem. Fl. Vit. 358. 1873 = P. lobatum var. 6.
hirsutulum Bernh. Schrad. Journ. **1801**^{1}. 16. 1802 = Nephrolepis.
hispidum (Sw.) J. Sm. JoB. **4**. 195. 1841. NPfl. 193 — N Zealand. Australia.
Aspidium Sw. 1801. Mett. n. 165. Chr. 266; Lastrea Houlst. (1851 ?): Moore 1857: Nephrodium Hk. 1862. HB. 286; Dryopteris O.Ktze. 1891. Polypodium setosum Forst. Prod. 82. 1786 (non Thbg. 1784).
Hookerianum (Pr.) C. Chr. Ind. 67. 1905; 582. 1906 — Ind. bor. China austr.
Lastrea Pr. 1836; Aspidium caducum Wall. 1828; Hk. et Grev. Ic. Fil. t. 171. 1829. Mett. n. 78. HB. 257. Chr. 233 (non HBK. 1815); Cyrtomium Moore 1861; Dryopteris O.Ktze. 1891; Polystichum Diels 1899. [NPfl. 194.
horizontale Pr. Epim. 57. 1849 = P. aculeatum var. 33.
ichangense Christ, Bull. Soc. Fr. **52** Mém. I. 28. 1905 — China.
ilicifolium Fée. Gen. 279. 1850-52: 6 mém. 21 t. 6 f. 4 = P. triangulum var.
ilicifolium (Don) Moore, Ind. 94. 1858 — Himalaya. Yunnan.
Aspidium Don. Prod. Fl. Nepal. 3. 1825. HB. 251. Chr. 238; Dryopteris O.Ktze. 1891; Polystichum stimulans Pr. 1836; Aspidium Kze.; Mett. n. [97. 1858.
inaequale Keys. Pol. Cyath. Hb. Bung. 44. 1873 = Dryopteris.
incisum Fée, Gen. 278. 1850-52 — Ins. Philipp.
induratum Schur. En. pl. Trans. 834. 1866 = Dryopteris filix mas.
? *inerme* Fée, Gen. 281. 1850-52 — Guiana.

intermedium Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris rhodolepis.
japonicum Diels, Nat. Pfl. 1⁴. 191. 1899 = P. aculeatum var. 40.
juglandifolium Diels, Nat. Pfl. 1⁴. 193. 1899 = Phanerophlebia.
Krugii Maxon, Proc. Biol. Soc. Wash. **18**. 215. 1905 — Puerto Rico. Cuba.
lacerum Pr. Epim. 56. 1849 = Dryopteris.
lachenense (Hk.) Bedd. Ferns br. Ind. t. 32. 1865. NPfl. 191 — Himalaya. China (Shen-si). — Aspidium Hk. sp. **4**. 8 t. 212. 1862. HB. 250; Dryopteris O.Ktze. 1891.
lactum Pr. Tent. 84. 1836 = P. denticulatum*.
lanceolatum (Bak.) Diels, Engl. Jahrb. **29**. 193. 1900 — China centr. Aspidium Bak. Gard. Chr. n. s. **14**. 494. 1880.
lanosum Fée, Cr. vasc. Br. **1**. 126 t. 40 f. 2. 1869 = P. aculeatum var. 17.
lanuginosum Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris.
laserpitiifolium J. Sm. Ferns br. and for. ed. II. 304. 1877 = P. Standishii.
laxum Wawra. Oest. Bot. Zeit. **13** 143. 1863 = P. aculeatum var. 23.
Lehmannii Hier. Engl. Jahrb. **34**. 452. 1904 — Columbia.
Lemmoni Und. Our Nat. Ferns ed. VI. 116. 1900 — California-Alaska. Aspidium mohrioides auctt. quoad. pl. amer. bor.
lentum (Don) Moore, Ind. 86. 1858 — India bor. — Aspidium Don, Prod. Fl. Nepal. 4. 1825; A. ocellatum Wall. 1828. Mett. n. 98; Polystichum Schott 1834; Àspidium auriculatum γ HB. 251.
lepidocaulon (Hk.) J. Sm. Ferns br. and for. 286. 1866. NPfl. 190 — Japonia. Aspidium Hk. sp. **4**. 12 t. 217. 1862. HB. 250; Dryopteris O.Ktze. 1891.
lepidomanes Fée, 8 mém. 99. 1857 — Java.
Levingei Hope; Christ, Bull. Ac. Géogr. Bot Mans **1902**. 260 (nomen) — Sikkim.
lobatum (Huds.) Pr. Tent. 83. 1836. NPfl. 191 — Europa. Asia minor. Caucasus. Persia. Japonia. China. India. Ins. Hawaii. Africa austr.
Polypodium Huds. Fl. angl. 459. 1762; Aspidium Sw. 1801. Mett. n. 108. Chr. 238. A. aculeatum part. auctt. HB. 352; Polystichum Roth 1799.

I. *Formae Africanae.*

— 1. Aspidium luctuosum Kze. Linn. **10**. 548. 1836. Mett. n. 119; Polystichum Moore 1858. NPfl. 191 — Africa austr.

(*Formae Americanae* vide P. californicum).

II. *Formae Asiaticae.*

— 2. Aspidium Tsus-Simense Hk. sp. **4**. 16 t. 220. 1862; Polystichum J. Sm. 1875. NPfl. 191 — Japonia. China bor.

— 3. Aspidium discretum Don, Prod. Fl. Nepal. 4. 1825; Polystichum Diels 1899. NPfl. 191 — Aspidium setosum Wall. 1828 (non Sw. 1801); Polystichum Schott 1834. NPfl. 191; P. Wallichianum Pr. 1836; Aspidium Kze. 1851 — A. polyblepharon Röm.: Kze. 1848: Polystichum Pr. 1849 — Japonia. India. ? Malesia.

— 4. Aspidium squarrosum Don, Prod. Fl. Nepal. 4. 1825. Mett. n. 103; Hypopeltis Bory 1833; Polypodium Roxb. 1844; Polystichum Fée 1850-52. Aspidium rufobarbatum Wall. 1828; Polystichum Schott 1834. NPfl. 191 — India.

III. *Formae Polynesiae.*

— 5. Polystichum aristatum Hk.fil. Fl. N. Zeal. **2**. t. 78. 1854 (non Pr. 1836) — N. Zealand.

— 6. Polystichum Hillebrandii Carr in Seem. Fl. Vit. 358. 1873. NPfl. 191; Aspidium Hill. 1888 — Ins. Hawaii.

Obs. Vide et P. Richardi. Conf. P. aculeatum.

lobatum × lonchitis (Murbeck) — Europa alpina. — Aspidium Murbeck, Lunds Univ. Årsskr. 27⁵. 16. 1892. ? A. illyricum Borbás 1891.

lonchitis (L.) Roth. Röm. Mag. 2¹. 106. 1799; Tent. Fl. Germ. 3. 71. 1800. Bernh. Schrad. Journ. 1799¹. 307. NPfl. 190 — Europa. Asia et Amer. arct. et temp. Himalaya. — Polypodium L. sp. 2. 1088. 1753; Aspidium Sw. 1801. Mett. n. 92. HB. 250. Chr. 236; Hypopeltis Todaro 1866; Dryopteris O.Ktze. 1891.

lonchitoides (Christ) Diels. Nat. Pfl. 1⁴. 194. 1899 — China.
Aspidium Christ. Bull. Boiss. 7. 16. 1899; Cyrtomium Christ 1902.

longecuspis Fée. Cr vasc. Br. 1. 129 t. 41 f. 2. 1869 = P. aculeatum var. 18.

luctuosum Moore, Ind. 95. 1858 = P. lobatum var. 1.

Macleaii (Bak.) Diels, Nat. Pfl. 1⁴. 190. 1899 (Macleanii) — Africa austr.
Aspidium Bak. in Hk. Ic. pl. t. 1654. 1886 Chr. 237.

macroporum J. Sm. Hist. Fil. 220. 1875 = P. adiantiforme.

Maderense Johns. Ann. Nat. Hist. III. 17. 287. 1866 = P. falcinellum.

Marantae Roth, Röm. Arch. 2¹. 106. 1799; Tent. Fl. Germ. 3. 92. 1800 = Woodsia ilvensis.

marginale Keys. Pol. Cyath. Hb. Bung. 44. 1873 = Dryopteris.

marginatum Schott, Gen. Fil. ad t. 9. 1834 = P. nepalense.

Martini (Christ) Bull. Ac. Géogr. Bot. Mans 1902. 263 c. tab. — China.

Mauritianum Fée, Gen. 278. 1850-52 = P. aculeatum var. 1.

Maximowiczii (Bak.) Diels, Nat. Pfl. 1⁴. 189. 1899 — Japonia. Liu Kiu.
Polypodium Bak. Syn. 504. 1874; Ptilopteris Hance 1884; Phegopteris Christ 1897. 271.

melanochlamys (Fée) Diels. Nat. Pfl. 1⁴. 193. 1899 — Cuba. — Aspidium Fée. Gen. 294. 1850 52. HB. 256; Lastrea Moore 1858; Dryopteris O.Ktze. 1891.

melanostictum (Kze.) Liebm. Vid. Selsk. Skr. V. 1. 276 (seors. 124). 1849. NPfl. 193 — Mexico. — Aspidium Kze. Linn. 13. 148. 1839. Mett. n. 173. HB. 256; Lastrea Moore 1858; Dryopteris O.Ktze. 1891.

microphyllum Pr. Epim. 55. 1849 = P. aculeatum var. 32.

microsorium Fée. Cr. vasc. Br. 1. 125 t. 40 f. 1. 1869 = P. aculeatum var. 19.

minusculum Christ. Bull. Soc. Fr. 52 Mém. I. 31. 1905 — Yunnan.

mohrioides (Bory) Pr. Tent. 83. 1836. NPfl. 193 — Quito-Peru. Amer. austr. temp. et antarct. Ins. Falklandicae. Ins. Marion. Ins. Amsterdam.
Aspidium Bory, Mém. Soc. Linn. Paris 4. 597; Dup. Voy. Bot. 1. 267 t. 35 f. 1. 1828. Mett. n. 100. HB. 252; Nephrodium Desv. 1827; Dryopteris O.Ktze. 1891. Aspidium plicatum Poepp.; Kze. 1834 — ? A. fuscatum Willd. 1810 (f. Kuhn nom. opt.) — Polystichum elegans Remy; Gay 1853; Aspidium Sturm 1858 (non Mett. 1858).

molle Gaud. Freyc. Voy. Bot. 326. 1827 = Dryopteris parasitica.

moluccense Moore. Ind. 84. 1858 = P. aculeatum var. 33.

monotis (Christ) C. Chr. Ind. 83. 1905; 584. 1906 — China.
Aspidium Christ, Bull. Soc. bot. It. 1901. 294.

montanum Roth, Röm. Arch. 2¹. 106. 1799; Tent. Fl. Germ. 3. 74. 1800 = Dryopteris oreopteris.

Moorei Christ, Arkiv för Bot. 4¹². 3. 1905 — Ins. Lord Howe.

moupinense (Franch.) Bedd. Handb. Suppl. 42. 1892. NPfl. 191 — China occ. Tibet orient. — Aspidium Franch. Nouv. Arch. Mus. II. **10**. 115. 1887.
mucronatum Pr. Tent. 83. 1836 = P. echinatum.
mucronifolium Pr. Epim. 55. 1849 = P. aculeatum var. 32.
Mülleri Schum. En. pl. Saell. **2**. 20. 1803 = Dryopteris spinulosa.
Mülleri Fourn. Mex. pl. **1**. 91. 1872 = P. aculeatum var. 5.
Münchii (Christ) C. Chr. Ind. 83. 1905; 585. 1906 — Mexico.
Aspidium Christ, Bull. Boiss. II. **1**. 1120. 1901.
multifidum (Mett.) Moore, Ind. LXXXIV. 1857. NPfl. 193 — Chile austr.
Aspidium Mett. Fil. Lechl. **1**. 20. t. 3 f. 14-17. 1856. Mett. n. 102. HB. 256. Chr. 241; Dryopteris Metteniana O.Ktze. 1891. Polystichum Pearcei Phil. 1865.
multiflorum Roth, Röm. Arch. **2**[1]. 106. 1799; Tent. Fl. Germ. **3**. 87. 1800 = Dryopteris spinulosa *dilatata.
munitum (Klf.) Pr. Tent. 83. 1836. NPfl. 189 — Alaska-California.
Aspidium Klf. Enum. 236. 1824. Mett. n. 93. HB. 249. Chr. 236; Dryopteris O.Ktze. 1891. Nephrodium plumula Pr. 1825; Polystichum Pr. 1836.
muricatum Fée, Gen. 278. 1850-52 = P. aculeatum var. 4.
Napoleonis Keys. Pol. Cyath. Hb. Bung. 44. 1873 = Dryopteris.
Neo-Zelandicum Fée, 8 mém. 99. 1857 — N. Zealand.
nepalense (Spr.) C. Chr. Ind. 84. 1905; 585. 1906 — India bor. China.
Aspidium Spr. Syst. **4**. 97. 1827. A. marginatum Wall. 1828. Mett. n. 88; Polystichum Schott 1834; Aspidium auriculatum β HB. 251. A. Manmeiense Christ 1898.
nephrolepioides Christ, Bull. Ac. Géogr. Bot. Mans **1902**. 258 c. fig. — China.
nigropaleaceum Diels, Nat. Pfl. **1**[4]. 191. 1899 = P. aculeatum var. 39.
nitens Fée, 8 mém. 101. 1857 — Ceylon. — Phegopteris Fée, Gen. 246. 1850-52.
nymphale Gaud. Freyc. Voy. Bot. 330. 1827 = Dryopteris parasitica.
obliquum (Don) Moore, Ind. 87. 1858. NPfl. 191 — India bor.
Aspidium Don. Prod. Fl. Nepal. 3. 1825; A. caespitosum Wall. 1828. Mett. n. 87; Polystichum Schott 1834; Aspidium auriculatum δ HB. 493.
obtusum J. Sm. JoB. **3**. 412. 1841 (nomen); Pr. Epim. 53. 1849 — Luzon.
Aspidium Kze.; Mett. n. 120. 1858. HB. 252; Dryopteris O.Ktze. 1891.
ocellatum Schott, Gen. Fil. ad t. 9. 1834 = P. lentum.
oculatum Armstr. Tr. N. Zeal. Inst. **13**. 364. 1881 = P. Richardi*.
omeiense C. Chr. Ind. 67. 1905; 585. 1906 — China occ. — Aspidium caruifolium Bak. JoB. **1888**. 228 (non Kze. 1851); Polystichum Diels 1900.
orbiculare Christ, Arkiv för Bot. **4**[12]. 3. 1905 } = P. aculeatum var. 13.
orbiculatum Gay, Fl. chil. **6**. 515. 1853 } = P. aculeatum var. 13.
ordinatum Liebm. Vid. Selsk. Skr. V. **1**. 276 (seors. 123). 1849 = P. aculeatum [var. 4.
Oreopteris Bernh. Schrad. Journ. **1799**[1]. 305 = Dryopteris.
otophorum (Franch.) Bedd. Handb. Suppl. 42. 1892. NPfl. 190 — Tibet orient.
Aspidium Franch. Nouv. Arch. Mus. II. **10**. 116. 1887.
pallidum Gardn. Lond. JoB. **1**. 547. 1842 — Rio Janeiro.
pallidum Todaro, Syn. pl. acot. vasc. Sic. 35. 1866 = Dryopteris rigida *australis.
pallidum Fourn. Mex. pl. **1**. 91. 1872 = P. aculeatum var. 6.
parvulum Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 114 c. fig. — China.
patens (Bernh. Schrad. Journn. **1799**[1]. 298); Gaud. Freyc. Voy. Bot. 329. 1827 [= Dryopteris.
Pearcei Phil. Linn. **33**. 305. 1865 = P. multifidum.
pennigerum Gaud. Freyc. Voy. Bot. 328. 1827 = Dryopteris.

perelegans (Col.) C. Chr. Ind. 87. 1905: 586. 1906 — N. Zealand.
Aspidium Col. Tr. N. Zeal. Inst. **29**. 416. 1897.

Phegopteris Roth, Röm. Arch. 2[1]. 106. 1799; Tent. Fl. Germ. **3**. 72. 1800; Bernh. Schrad. Journ. **1799**[1]. 305 = Dryopteris.

Plaschnickianum (Kze.) Moore, Ind. 100. 1858. NPfl. 189 — Jamaica.
Aspidium Kze. Linn. **23**. 229, 302. 1850. Mett. n. 84. HB. 249; Dryopteris O.Ktze. 1891.

platylepis Fée, Cr. vasc. Br. **1**. 129 t. 41 f. 1. 1869 = P. aculeatum var. 20.

platyphyllum Pr. Tent. 84. 1836 = P. aculeatum var. 23.

platypus Diels, Nat. Pfl. 1[4]. 193. 1899 = P. aristatum var.

Plukenetii DC. Fl. franç. **5**. 241. 1815 = P. lobatum.

plumosissimum ht. Birkenh.; Gard. Chr. III. **24**. 241. 1898 — Hort.

plumula Pr. Tent. 83. 1836 = P. munitum.

polyblepharum Pr. Epim. 56. 1849 = P. lobatum var. 3.

polyphyllum Pr. Tent. 83. 1836 = P. aculeatum var. 24.

praelongum Christ. Bull. Ac Géogr. Bot. Mans **1902**. 260 — China.

Prescottianum (Wall.) Moore, Ind. 101. 1858. NPfl. 191 — Himalaya.
Aspidium Wall. 1828 (nomen); Mett. Pheg. u. Asp. 48. n. 110. 1858. HB. 253; Dryopteris O.Ktze. 1891.

Prescottianum var. sinense Christ, Bull. Soc. bot. It. **1901**. 289 = P. sinense.

Preslianum Moore, Ind. LXXXIV. 1857 = Cyclopeltis.

proliferum Pr. Tent. 83. 1836 = P. aculeatum var. 29.

propinquum Gaud. Freyc. Voy. Bot. 330. 1827 = Dryopteris gongylodes*.

pseudo-cristatum Schur, »Sert. Pl. Transs. 94. 1852;« En. pl. Transs. 836. 1866 = Dryopteris spinulosa.

pulchellum Moore, Ind. 101. 1858 = P. aristatum.

pungens (Klf.) Pr. Tent. 83. 1836. NPfl. 192 — Africa austr. Réunion.
Aspidium Klf. Enum. 242. 1824. Mett. n. 115. HB. 252. Chr. 239; Dryopteris O.Ktze. 1891.

?pycnolepis Moore, Ind. 92. 1858 = P. aculeatum var. 27.

quadrangulare Fée, Cr. vasc. Br. **2**. 68 t. 99 f. 1. 1872-73 — Brasilia.

rachichlaena Fée, 8 mém. 100. 1857 = P. aculeatum var. 4.

radicans Pr. Tent. 83. 1836 = P. aculeatum var. 29.

recurvum Gandog. Fl. Eur. **27**. 174. 1891 = Dryopteris aemula.

remotum Fée, Cr. vasc. Br. **1**. 125 t. 39 f. 1. 1869 — Brasilia. Columbia.
Aspidium Christ 1900 (non A.Br. 1850).

rhizophyllum (Sw.) Pr. Tent. 82. 1836. NPfl. 189 — Ind. occ.
Polypodium Sw. Prod. 132. 1788; Aspidium Sw. 1801. Mett. n. 86. HB. 249. (Chr. 239): Nephrodium Pr. 1825; Dryopteris O.Ktze. 1891.

rhomboideum Schott, Gen. Fil. ad t. 9. 1834 = P. amabile.

Richardi (Hk.) J. Sm. Hist. Fil. 220. 1875. NPfl. 191 — N. Zealand. Fiji.
Aspidium Hk. sp. **4**. 23 t. 222. 1862. HB. 253. (Chr. 238); Dryopteris O.Ktze. 1891.
—* Aspidium oculatum Hk. sp. **4**. 24 t. 228. 1862. HB. 253; Polystichum Armstr. 1881. NPfl. 192; Dryopteris O.Ktze. 1891.

Riedleanum Gaud. Freyc. Voy. Bot. 327. 1827 — Ins. Moluccae.
Aspidium Gaud. l. c.: Nephrodium Moore 1858. — (Dryopteris sp. dub.).

rigidum Lam. et DC. Fl. franç. **2**. 560. 1805 = Dryopteris.

riparium Pr. Tent. 83. 1836 = ? Dryopteris Napoleonis.

Rochaleanum Glaz.; Fée, Cr. vasc. Br. **2**. 69 t. 99 f. 2. 1872-73 — Brasilia austr.
Aspidium Christ 1902.
?rufescens Moore, Ind. 102. 1858 = Dryopteris
rufobarbatum Schott, Gen. Fil. ad t. 9. 1834 = P. lobatum var. 4.
Schimperianum Keys. Pol. Cyath. Hb. Bung. 44. 1873 = Dryopteris.
Schizolobium Fée, 8 mém. 99. 1857 = P. aculeatum var. 2.
Schkuhrii Pr. Tent. 84. 1836 = P. hispidum.
scopulinum Maxon, Fern Bull. **8**. 29. 1900 = P. aculeatum var. 3.
Seemanni (Hk.) J. Sm. Hist. Fil. 220. 1875 — Panama.
Aspidium Hk. sp. **4**. 34 t. 230. 1862. HB. 255; Dryopteris O.Ktze. 1891.
Sellowianum Pr. Tent. 83. 1836; Epim. 53. 1849 = P. aculeatum var. 9.
semicordatum Moore, Ind. LXXXIV. 1857 = Cyclopeltis.
serratum Gaud. Freyc. Voy. Bot. 325. 1827 = Dryopteris gongylodes.
setigerum Pr. Tent. 83. 1836 = P. aculeatum.
setosum Schott, Gen. Fil. ad t. 9. 1834 = P. lobatum var. 3.
setosum Pr. Tent. 84. 1836 = P. varium.
Sieberianum Pr. Tent. 83. 1836; Epim. 53. 1849 = P. aculeatum var. 1.
Sieboldii Keys. Pol. Cyath. Hb. Bung. 41. 1873 = Dryopteris.
sikkimense Bedd. Ferns br. Ind. t. 127. 1866 = Dryopteris.
silvaticum Diels, Nat. Pfl. 1^4. 192. 1899 = P. vestitum var.
sinense Christ, Bull. Soc. Fr. **52** Mém. I. 30. 1905 — China.
P. Prescottianum var. sinense Christ, Bull. Soc. bot. It. **1901**. 289.
Sloanei Pr. Tent. 83. 1836 = P. triangulum.
Sodiroi Christ, Arkiv för Bot. 4^{12}. 4. 1905 — Ecuador.
sparsum Keys. Pol. Cyath. Hb. Bung. 43. 1873 = Dryopteris.
speciosum (Don) J. Sm. JoB. **4**. 195. 1841 — Ind. bor.
Aspidium Don, Pr. Fl. Nepal. 5. 1825. A. affine Wall. 1828. Mett. n. 104; Polystichum Pr. 1836; Lastrea Bedd. 1883.
spinosum Roth, Röm. Arch. 2^1. 106. 1799; Tent. Fl. Germ. **3**. 91. 1800 } = Dryopteris spinulosa.
spinulosum Lam. et DC. Fl. franç. **2**. 561. 1805 } = Dryopteris spinulosa.
spinulosum × cristatum Sanio, Verh. Bot. Ver. Brand. **25**. 82. 1884 = Dryopteris cristata × spinulosa.
spinulosum × dilatatum Sanio, Verh. Bot. Ver. Brand. **25**. 83. 1884 = Dryopteris spinulosa.
squamatum Lodd. Cat. 1849 — Hort. — Aspidium Kze. 1850.
squarrosum Fée, Gen. 278. 1850-52 = P. lobatum var. 4.
Standishii (Moore) C. Chr. Ind. 79. 1905: 587. 1906 — Japonia. Canton. Tonkin.
Lastrea Moore, Gard. Chr. **1863**. 292. Aspidium laserpitiifolium Mett. Ann. Lugd. Bat. **1**. 227 t. 6 f. 3. 1864. HB. 254. Chr. 240; Polystichum J. Sm. 1877. NPfl. 193; Dryopteris O.Ktze. 1891 — Aspidium festinum Hance 1883.
stenophyllum Christ, Bull. Soc. Fr. **52** Mém. I. 27. 1905 — Tibet orient. China.
stenopteris Moore, Ind. LXXXIV. 1857 = Dryopteris incisa.
stimulans Pr. Tent. 83. 1836 = P. ilicifolium.
stipitatum Chauvin, Mém. Soc. Linn. Calvados **1825**. 502 = P. lobatum.
stramineum Pr. Tent. 83. 1836 = P. aculeatum var. 1.
strigosum Roth, Röm. Arch. 2^1. 106. 1799; Tent. Fl. Germ. **3**. 86. 1800 = Dryopteris rigida.
submite (Christ) Diels, Engl. Jahrb. **29**. 192. 1900 — China centr.
Aspidium Christ, Nu. Giorn. bot. It. n. s. **4**. 93. 1897.
subquinquefidum Keys. Pol. Cyath. Hb. Bung. 45. 1873 = Dryopteris protensa.

tacticopterum Moore, Ind. 105. 1858 = P. aculeatum var. 35.
tanacetifolium Lam. et DC. Fl. franç. **2**. 562. 1805 = Dryopteris spinulosa [*dilatata.
tenerum Fée, Gen. 280. 1850-52 = P. denticulatum.
tenue Gilb. Fern Bull. **8**. 63. 1900 — Jamaica.
tetragonum Fée, 8 mém. 99. 1857 — Chile. — Aspidium Sturm 1858.
Thelypteris Roth, Röm. Arch. **2**^{1}. 106. 1799; Tent. Fl. Germ. **3**. 77. 1800: Bernh. Schrad. Journ. **1799**^{1}. 306 = Dryopteris.
Thomsoni (Hk.fil.) Bedd. Ferns br. Ind. t. 126. 1866. NPfl. 191 — Ind. bor. Aspidium Hk.fil. in Hk. 2 Cent. t. 25. 1860. HB. 251; Dryopteris O.Ktze. 1891 — Lastrea gracilis Moore; Bedd. 1866; Nephrodium sparsum var. gracilis HB. 498.
Tijucense Fée, Cr. vasc. Br. **1**. 127 t. 38 f. 1. 1869 = P. aculeatum var. 22.
Torresianum Gaud. Freyc. Voy. Bot. 333. 1827 — Ins. Mariannae. Aspidium Gaud. l. c.; Lastrea Moore 1858. — (Dryopteris sp. dub.).
trapezoides Pr. Tent. 83. 1836 = ? P. viviparum.
Trejoi (Christ) Bull. Boiss. II. **5**. 728. 1905 — Mexico. Costa Rica. Aspidium Christ, Bull. Boiss. II. **5**. 255. 1905.
triangulum (L.) Fée, Gen. 279. 1850-52. NPfl. 191 — Ind. occ. Polypodium L. sp **2**. 1088. 1753; Aspidium Sw. 1801. Mett. n. 89. HB. 250. Chr. 239; Dryopteris O.Ktze. 1891 — Polystichum cyphochlamys Fée 1850-52. —* Polystichum ilicifolium Fée 1850-52 (non supra); P. aquifolium Und. [et Maxon 1902.
tridens (Hk.) Fée, 11 mém. 74. 1866 — Jamaica. Aspidium Hk. sp. **4**. 15 t. 215. 1862. HB. 251; Dryopteris O.Ktze. 1891.
tripinnatum Goldm. Nova Acta **19** Suppl. I. 463. 1843 = P. carvifolium.
tripteron (Kze.) Pr. Epim. 55. NPfl. 191 — Japonia. Ins. Tsus Sima. Aspidium Kze. Bot. Zeit. **1848**. 569. Mett. n. 118. HB. 254. Chr. 237; Dryopteris O.Ktze. 1891; Aspidium tripteris Eat. 1856.
truncatum Gaud. Freyc. Voy. Bot. 332. 1827 = Dryopteris.
Tsus-Simense J. Sm. Hist. Fil. 219. 1875 = P. lobatum var. 2.
unitum Gaud. Freyc. Voy. Bot. 325. 1827 = Dryopteris.
varium (L.) Pr. Epim. 57. 1849. NPfl. 193 — Japonia. Korea. China. Assam. Polypodium L. sp. **2**. 1090. 1753; Aspidium Sw. 1801. Mett. n. 147. HB. 254. Chr. 259; Nephrodium Desv. 1827; Lastrea Moore 1858; Dryopteris O.Ktze. 1891. Polypodium setosum Thbg. 1784; Aspidium Sw. 1801; Hypopeltis Bory 1833; Polystichum Pr. 1849; Lastrea Moore 1858 — Lastrea opaca Hk. 1857; Aspidium Benth. 1861 — Nephrodium coriaceum Hope 1890.
venustum Hombr. et Jacq. Voy. au Pôle Sud t. 5. 1852 = P. vestitum.
vestitum (Forst.) Pr. Tent. 83. 1836. NPfl. 192 — N. Zealand. Insulae antarcticae. Amer. antarct. Australia. — Polypodium Forst. Prod. 82. 1786; Aspidium Sw. 1801. Mett. n. 101. (HB. 252). Chr. 239 — A. Tasmaniae Mett. 1856.
—* Polypodium silvaticum Col. Tasm. Journ. **2**. 163. 1844; Hk.fil. Fl. N. Zeal. **2**. 41 t. 81. 1854; Polystichum Diels 1899. NPfl. 192.
viscidulum C. Chr. Ind. 75. 1905 = Adenoderris.
vittatum (Christ) C. Chr. Ind. 202. 1905; 588. 1906 — Yunnan. Cyrtomium Christ, Bull. Soc. Fr. **52** Mém. I. 33. 1905.
viviparum Fée. Gen. 280. 1850-52; 6 mém. 21 t. 3 f. 3 — India occ. Aspidium Mett. n. 99. 1858. HB. 251. (Chr. 239); Dryopteris O.Ktze. 1891

— Aspidium caudatum Jenm. 1879 (non Sw. 1806) — ? A. trapezoides Sw. 1801; Polypodium Sw. 1806; Polystichum Pr. 1836.

Volkensii (Hier.) C. Chr. Ind. 97. 1905; 589. 1906 — Africa orient.
Aspidium Hier. in Engl. Pflw. Ostafr. C. 86. 1895.

vulcanicum Moore, Ind. 108. 1858 = P. aculeatum var. 31.

Wallichianum Pr. Tent. 83. 1836 = P. lobatum var. 3.

Wattii (Bedd.) C. Chr. Ind. 98. 1905; 589. 1906 — Munipore.
Aspidium Bedd. JoB. **1888**. 234; Lastrea Bedd. 1892.

Webbianum (A.Br.) C. Chr. Ind. 74. 1905; 589. 1906 — Madeira.
Aspidium A.Br. Flora **1841**. 711; A. frondosum Lowe, Tr. Cambr. Phil. Soc. **4**. 6. 1830. Mett. n. 156. HB. 255 (non Wikstr. 1826); Polystichum J. Sm. 1857. NPfl. 193; Lastrea Moore 1858; Dryopteris O.Ktze. 1891.

xiphophyllum (Bak.) Diels, Nat. Pfl. 1^4. 189. 1899 — China occ.
Aspidium Bak. JoB. **1888**. 227.

zerophyllum (Col.) C. Chr. Ind. 98. 1905; 589. 1906 — N. Zealand.
Aspidium Col. Tr. N. Zeal. Inst. **29**. 418. 1897.

POLYTAENIUM Desvaux Prodrome 174 (218: Polyaetnium). 1827 = **Antrophyum.**

Grevillii Moore, Ind. LVII. 1857.

lanceolatum Desv. Prod. 218. 1827 = A. lineatum.

lineatum J. Sm. JoB. **4**. 68. 1841.

Poronema J. Smith, Hist. Fil. 108. 1875 *(Drynaria §)* = Drynaria.

PORPAEA Presl msc. (Ettingsh. Farnkr. 174. 1865) = **Polystichum.**

PRESLIA Opiz (sec. Milde, Fil. Eur. 164. 1867) = **Woodsia.**

paleacea Opiz = W. ilvensis.

PRIONOPTERIS Wallich, List n. 184. 1828 = **Matonia.**

Farquhariana Wall. List n. 184. 1828 = M. pectinata.

PROFEREA Presl, Epim. Bot. 259. 1849 = **Aspidium.**

excellens Pr. Epim. 259. 1849.

PRONEPHRIUM Presl, Epim. 258. 1849 = **Dryopteris.**

acrostichoides Pr. Epim. 259. 1849 = D. canescens*.

affine Pr. Epim. 259. 1849 = D. lineata.

lastreoides Pr. Epim. 259. 1849 — Java. — Aspidium Mett; Salom. 1883.

lineatum Pr. Epim. 259. 1849.

PROSAPTIA Presl. Tent. Pterid. 165. 1836 = **Davallia.**

alata Christ, Ann. Jard. Buit. II. **5**. 127. 1905.

bipinnata Pr. Tent. 166 t. 6 f. 19. 1836 = Odontosoria uncinella.

contigua Pr. Tent. 166. 1836.

? Emersoni Pr. Tent. 166. 1836 = D. alata.

Friederici et Pauli Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 41. 1905.

nutans Mett. Novara Reise. Bot. **1**. 214. 1870 = Polypopium n.

obliquata Mett. Novara Reise. Bot. **1**. 214. 1870 = Polypodium o.

pectinata Moore, Ind. 298. 1861 = D. contigua.

pinnatifida Pr. Tent. 166 t. 6 f. 25. 1836 = D. contigua.

Preslii Fée, Gen. 324. 1850-52 = D. contigua.

Reineckei Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 41. 1905.

Schlechteri Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 41 t. 1 B. 1905.

serraeformis Christ. Bull. Boiss. **6**. 143. 1898 = D. alata.

Prosechium Moore. Index Fil. LXXI. 1857 *(Polypodium §)* = Polypodium.

Protogleichenia Copeland in Perkins. Fragmenta 194. 1905 *(Dicranopteris §)* = Gleichenia.

Pseudachomanes Presl, Epim. Bot. 16. 1849 *(Trichomanes §)* = Trichomanes.
Pseudallantodia Clarke, Tr. Linn. Soc. II. Bot. **1**. 495. 1880 *(Asplenium §)* = Athyrium.
Pseudangiopteris Presl, Suppl. Tent. Pterid. 23. 1845 *(Angiopteris §)* = Angiopteris.
PSEUDATHYRIUM Newman, The Phytologist **4**. 370. 1851 = **Athyrium.**
alpestre Newm. Phytol. 4. app. XIV. 1851.
flexile Newm. Phytol. 4. 394. 1851 = A. alpestre.
PSIDOPODIUM Necker, Element. Bot. **3**. 315. 1790 = **Dryopteris.**
PSILODOCHEA Presl, Suppl. Tent. Pterid. 28. 1845 = **Angiopteris.**
salicifolia Pr. Suppl. 28. 1845.
PSILOGRAMME Kuhn, Chaetopt. 332 (12). 1882.
aureo-nitens Kuhn, l. c. 341 — Biardii Kuhn, l. c. 336 — cheilanthoides Kuhn, l. c. 335 — dubia Kuhn, l. c. 337 — elongata Kuhn, l. c. 335 — flabellata Kuhn, l. c. 336 — flexuosa Kuhn, l. c. 339 — hirsutula Kuhn, l. c. 340 — hirta Kuhn, l. c. 338 — hispidula Kuhn, l. c. 341 — incisa Kuhn, l. c. 336 — insignis Kuhn, l. c. 337 — Karstenii Kuhn, l. c. 340 — Lechleri Kuhn, l. c. 339 — Lindigii Kuhn, l. c. 340 — Mathewsii Kuhn, l. c. 337 — myriophylla Kuhn, l. c. 339 — Orbignyana Kuhn, l. c. 338 — Ottonis Kuhn, l. c. 340 — rufescens Kuhn, l. c. 336 — Schomburgkiana Kuhn, l. c. 341 — Sellowiana Kuhn, l. c. 337 — Warscewiczii Kuhn, l. c. 337 = **Gymnogramma** sp. homonym.
canescens Kuhn, l. c. 333 — cinnamomea Kuhn, l. c. 334 — glutinosa Kuhn, l. c. 333 — imbricata Kuhn, l. c. 334 — laxa Kuhn, l. c. 333 — nivea Kuhn, l. c. 333 — rotundifolia Kuhn, l. c. 334 — scalaris Kuhn, l. c. 334 — verticalis Kuhn, l. c. 332 = **Jamesonia** sp. homonym.
domingensis Und. Bull. Torr. Cl. **29**. 628. 1902 = Gymnogramma.
glandulosa Kuhn, Chaetopt. 338. 1882 = Gymnogramma caracasana.
schizophylla Und. Bull. Torr. Cl. **29**. 629. 1902 = Anogramma.
PSOMIOCARPA Presl, Epim. Bot. 161. 1849 = **Polybotrya.**
acuminata Pr. Epim. 162. 1849 = P. scandens.
apiifolia Pr. Epim. 161. 1849.
caudata Pr. Epim. 162. 1849.
incisa Pr. Epim. 162. 1849 = P. osmundacea var.
PSYGMIUM Presl, Tent. Pterid. 199. 1836 = **Polypodium.**
elegans Pr. Tent. 200 t. 8 f. 17. 1836 = P. Meyenianum.
PTERIDELLA Mettenius; Kuhn, v. Deck. Reis. **3**[3]. Bot. 13. 1879 = **Pellaea.**
adiantoides Kuhn, l. c. 14 = P. Boivini.
angulosa Mett.; Kuhn, l. c. 15.
Belangeri Mett.; Kuhn, l. c. 16 = Cheilanthes B.
Doniana Mett.; Kuhn, l. c. 13.
dura Mett.; Kuhn, l. c. 14.
hastata Mett.; Kuhn, l. c. 14.
Holstii Hier. in Engl. Pflw. Ostafr. C. 78. 1895.
involuta Mett.; Kuhn, v. Deck. Reis. 3[3]. Bot. 15. 1879.
leucomelas Mett.; Kuhn, l. c. 14.
pectiniformis Mett.; Kuhn, l. c. 14 = P. Goudotii.
quadripinnata Mett.; Kuhn, l. c. 16.
Schweinfurthii Hier. in Engl. Pflw. Ostafr. C. 78. 1895.
viridis Mett.; Kuhn, v. Deck. Reis. 3[3]. Bot. 16. 1879.
Pteridellastrum Prantl, Engl. Jahrb. **8**. 421. 1882 *(Pellaea §)* = Pellaea.

PTERIDIUM Gleditsch; Scopoli, Flora carniolica 169. 1760 — (Polypodiaceae Gen. **86**).

aquilinum (L.) Kuhn, v. Deck. Reisen **3**[3]. Bot. 11. 1879. NPfl. 296 — Reg. temp. et trop. — Pteris L. sp. **2**. 1075. 1753. HB. 162. Chr. 164; Asplenium Bernh. 1799; Allosorus Pr. 1836; Eupteris Newm. 1845; Paesia Keys. 1873; Cincinalis Trev. 1874; Ornithopteris J. Sm. 1875 — Pteris arachnoidea Klf. 1824; Allosorus Pr. 1836 — Pteris psittacina Pr. 1822; Allosorus Pr. 1836.

—* Pteris caudata L. 1753; Allosorus Pr. 1836; Ornithopteris J. Sm. 1875; Pteridium Maxon 1901.

—* Pteris esculenta Forst. 1786; Allosorus Pr. 1836; Ornithopteris J. Sm. 1875. (Reg. austral.).

—* Pteris capensis Thbg. 1800; Allosorus Pappe et Raws. 1858 (non Pr. 1836); Pteridium Krasser 1900 — Pteris lanuginosa Bory; Willd. 1810; Allosorus Pr. 1836 — Pteris coriifolia Kze. 1844; Allosorus Pappe et Raws. 1858. (Africa austr.).

capense Krasser, Ann. Hofmus. Wien **15**. 6. 1900 = P. aquilinum*.

caudatum Maxon, Proc. U. S. Nat. Mus. **23**. 631. 1901 = P. aquilinum*.

lanuginosum Clute, Fern Bull. **8**. 38. 1900 = P. aquilinum.

Pteridopsis Link, Fil. Spec. 49. 1841 *(Pteris §)* = Pteris.

Pteridopsis J. Smith, Hist. Fil. 283. 1875 *(Cheilanthes §)* = Cheilanthes.

PTERIGLYPHIS Fée, X[e] Congr. scient. de France **1**. 178. 1843; Gen. Fil. 219. 1850-52 = **Diplazium.**

elegans Fée, Gen. 220 t. 18 B. 1850-52 = D. Cumingii.

PTERINODES Siegesb.; O.Kuntze, Rev. Gen. Pl. **2**. 819. 1891.

orientale O.Ktze. l. c. 820 = Matteuccia.

sensibile O.Ktze. l. c. 820 = Onoclea.

Struthiopteris O.Ktze. l. c. 820 = Matteuccia.

PTERIS Linné, (Hort. Cliff. 443. 1737) Spec. plant. **2**. 1073. 1753 — (Polypodiaceae Gen. **83**).

Subgenera: C = **Campteris.**
L = **Litobrochia.**
P = **Eupteris.**

acclivis Mett. Fil. Lips. 59. 1856 = Paesia.

acinaciformis Sessé et Mociño. Fl. Mex. 262. 1896 — Cuba.

acrosticha Balbis, Elenco del Piante 98. 1801 = Cheilanthes pteridioides.

(P) **actiniopteroides** Christ, Bull. Boiss. **7**. 6. 1899. NPfl. 292 — China merid.

aculeata Sw. Prod. 129. 1788 = P. spinosa.

aculeata Nadeaud, Enum. pl. Tahiti »n. 129«. 1873 = P. Nadeaudii.

acuminata Desv. Berl. Mag. **5**. 324. 1811 = P. pungens.

acuminata Bak. JoB. **1891**. 5 = P. Lastii.

acuminatissima Bl. Enum. 208. 1828 = P. longifolia.

acutangula Nees, Linn. **19**. 684. 1847 = Doryopteris rigida.

adiantoides Burm. Fl. Ind. Cap. Prod. 28. 1768 (nomen) — Afr. austr.

adiantoides Bory; Willd. sp. **5**. 391. 1810 = Pellaea viridis.

adiantoides Vell. Fl. flum. **11** t. 88. 1827; Arch. Mus Nac. Rio Jan. **5**. 452. 1881 = Lindsaya lancea.

Adscensionis vide Ascensionis.

aequalis Pr. Rel. Haenk. **1**. 54. 1825 = P. longifolia.

affinis A. Rich. Fl. N. Zeal. 81. 1832 = P. tremula.

alabamensis Buckl. Amer. Journ. Sci. **45**. 177. 1843 = Cheilanthes.
alata Lam. Illustr. t. 869; Poir. Enc. **5**. 716. 1804 = P. semipinnata.
(L) ?*alata* Pr. Rel. Haenk. **1**. 56. 1825 — Ecuador. Luzon.
Litobrochia Pr. 1836.
alata Desv. Prod. 297. 1827 = P. woodwardioides.
alata Gaud. Freyc. Voy. Bot. 391 t. 19. 1827 = P. irregularis.
Alcyonis Lind.; Salom. Nom. 332. 1883 — Brasilia.
(Litobrochia »Pr.«; Doryopteris »J. Sm.«).
Alexandrae ht. May: Gard. Chr. III. **31**. 279. 1902 — Hort
allosora Link, Hort. Berol. **2**. 31. 1833 = P. biaurita.
alpina Field. Ferns N. Zeal. 97 t. 28 f. 2. 1890 = Cheilanthes tenuifolia.
Alpinii Desv. Prod. 295. 1827; Spr. Syst. **4**. 71. 1827 = P. longifolia.
(L) **altissima** Poir. Enc. **5**. 722. 1804 — Ind. occ. Guatemala-Ecuador.
?Litobrochia Fée 1850-52. Pteris elata Ag. 1839. HB. 173. Chr. 173; Litobrochia Fée 1850-52.
alternifolia Wall. List n. 2182. 1829 = Pellaea falcata.
Amazonica Christ, Hedwigia **44**. 364. 1905 = Paesia.
ambigua Desv. Prod. 297. 1827 — Timor. India.
(P) **amoena** Bl. Enum. 210. 1828. HB. 479 — Java.
ampla Kze. Linn. **9**. 74. 1834 = P. Haenkeana.
amplectens Wall. List n. 112. 1828 = P. longifolia.
amplexicaulis Roxb. Calc. Journ. **4**. 505. 1844 = P. longifolia.
(L) **Andreana** Sod. Cr. vasc. quit. 102, 635. 1893 — Ecuador.
andromedaefolia Klf. Enum. 188. 1824 = Pellaea.
anfractuosa Christ. Bull. Boiss. II. **5**. 257. 1905 = Paesia.
angulata Pr. Rel. Haenk. **1**. 54. 1825 = Schizoloma ensifolium.
angulosa Bory; Willd. sp. **5**. 377. 1810 = Pellaea.
angusta Bory; Willd. sp. **5**. 388. 1810 = P. scabra
angustata Wall. List n. 93. 1828 = Schizoloma ensifolium.
angustifolia Sw. Prod. 129. 1788 = Vittaria.
angustifolia F. Muell. Fragm. **7**. 121 (nota). 1870 = Cheilanthes.
angustisora Remy, Ann. sc nat. III. **6**. 345. 1846 — Bolivia. — (Pellaea sp.?).
apicalis Liebm. Vid. Selsk. Skr. V. **1**. 230 (seors. 78). 1849 = P. podophylla.
appendiculata Klf. Enum. 187. 1824 = P. decurrens.
appendiculata Bak. JoB. **1881**. 366 = P. Curtisii.
aquilina L. sp. **2**. 1075. 1753 = Pteridium.
arachnoidea Klf. Enum. 190. 1824 = Pteridium aquilinum var.
arborea L. sp. **2**. 1073. 1753 = ? P. spinosa.
arborescens Mart. et Gal. Mém. Ac. Brux. **15**. 54. 1842 = P. Kunzeana.
argentea Gmel. Nova Comm. Petr. **12**. 519 t. 12 f. 2. 1768 = Cheilanthes.
argentea Bory. Voy. **1**. 327. 1804 = Cheilanthes farinosa.
argentea ht. Sander: Gard. Chr. III. **27**. 206. 1900 — Hort. = P. tremula var.
(P) **arguta** Ait. Hort. Kew. **3**. 458. 1789. HB. 160. Chr. 169. NPfl. 293 — Marocco. Ins Atlant. Portugal. ? Graecia. — P. palustris Poir. 1804. (an Pteris serrulata Forsk. 1775?).
arguta Ham.; Wall. List n. 2179. 1829 = P. scabra.
argyraea Moore, Gard. Chr. **1859**. 671 = P. biaurita* var.
argyrophylla Sw. Syn. 105. 1806 = Cheilanthes farinosa.
armata Pr. Rel. Haenk. **1**. 56. 1825; Zoll. et Moritz. Nat. Gen. Arch. Neerl. Ind. **1**. 398. 1844 = P. biaurita.

articulata Klf.; Spr. Syst. 4. 76. 1827 = Pellaea angulosa.
Ascensionis Sw. Schrad. Journ. **1800**^{2}. 67. 1801 = P. dentata var.
aspera Poir. Enc. **5**. 713. 1804 = Adiantum lucidum.
aspera Fée, Gen. 126. 1850-52 = P. longifolium.
aspericaulis Wall. List n. 107. 1828 = P. biaurita*.
asperiuscula Willd.; Pr. Tent. 145. 1836 (nomen) — ?
asperula J. Sm. JoB. **3**. 405. 1841 = P. biaurita*.
atropurpurea L. sp. **2**. 1076. 1753 = Pellaea.
atropurpurea Goldm. Nova Acta **19** Suppl. I. 457. 1843 = P. excelsa.
(L) **atrovirens** Willd. sp. **5**. 385. 1810. HB. 170. Chr. 168 — Africa trop.
P. spinulifera Schum. 1827; Litobrochia J. Sm. 1866.
attenuata Sw. Schrad. Journ. **1800**^{2}. 66. 1801 = P. tripartita.
attenuata Pr. Rel. Haenk. **1**. 55. 1825 = P. biaurita*.
attenuata Bl. Enum. 210. 1828 = P. ensiformis.
aurantiaca Cav. Descr. 266. 1802 = Cheilanthes.
aurata Mett. Fil. Lips. 54. 1856 = Onychium siliculosum.
aurea Poir. Enc. **5**. 710. 1804 = Notholaena.
auriculata Thbg. Prod. Fl. Cap. 172. 1800 = Pellaea viridis.
auriculata Sw. Schrad. Journ. **1800**^{2}. 69. 1801 = Pellaea.
auriculata Goldm. Nova Acta **19** Suppl. I. 458. 1843 = ? Pteridium aquilinum.
aurita Bl. Enum. 213. 1828 = Histiopteris incisa var.
australasica Desv. Prod. 302. 1827 = Histiopteris incisa.
australis Thouars, Obs. sur l'Isl. d'Afr. 11. —; Hk. et Grev. Ic. Fil. t. 8. 1827 = Actiniopteris.
Backeri ht. Bat Backer, Gard. Chr. III. **35**. 317. 1904 — Hort.
Bahamensis Fée, Gen. 125. 1850-52 = P. longifolia.
(P) **Bakeri** C. Chr. Ind. 593. 1906 — Peru.
P. decomposita Bak. Syn. 479. 1874. NPfl. 293 (non Gaud. 1827).
(P) **Balansae** Fourn. Ann. sc. nat. V. **18**. 320. 1873 — N. Caledonia.
Barkleyae Mett.; Kuhn, Fil. Afr. 77. 1868 = Pellaea Barklyae.
Barteri Bak. Syn. 168. 1867 (var.) = Lonchitis Currori var.
Baueriana Dies; Endl. Prod. Fl. Norf. 16. 1833 — Norfolk.
(P) **Beccariana** C. Chr. Ind. 593. 1906 — N. Guinea.
P. concinna Bak. Malesia **3**. 37. 1886 (non Hew. 1838).
(L) **Beecheyana** Ag. Rec. 68. 1839 — Polynesia.
Litobrochia Moore 1857; Pteris aculeata auctt. quoad pl. Polynesiae.
Belangeri Bory, Bél. Voy. Bot. **2**. 44. 1833 = Cheilanthes.
Berteroana Ag. Rec. 66. 1839 = P. comans.
(C) **biaurita** L. sp. **2**. 1076. 1753. HB. 164. NPfl. 292 — Trop. et Subtrop.
Campteria Hk. 1840; Litobrochia J. Sm. 1857 — Pteris nemoralis Willd. 1809; Campteria J. Sm. 1846.
(P) —* P. quadriaurita Retz. 1791. HB. 158. Chr. 166. NPfl. 292 — P. Swartziana Ag. 1839 — P. deltea Ag. 1839 — P. Blumeana Ag. 1839 — P. pyrophylla Bl. 1828 — P. felosma J. Sm. 1846 — P. subindivisa Clarke 1880 — P. argyraea Moore 1859 — P. asperula J. Sm. 1841 — P. reducta Bak. 1880 — P. subquinata Wall. 1828; Hope 1901.
bicolor Roxb. Calc. Journ. **4**. 507. 1844 = Cheilanthes farinosa.
biformis Splitgb. Tijdschr. Nat. Gesch. **7**. 422. 1840 = P. propinqua.
Binoti ht. Hill.; Gard. Mag. **1904**. 376 — Hort. (Brasilia). — (Doryopteris sp.).
Blanchetiana Pr.; Ettingsh. Farnkr. 103 t. 36 f. 7. 1865 = P. propinqua.

blechnoides Willd. Phyt. 13 t. 9 f. 3. 1794 = Taenitis.

Blumeana Ag. Rec. 22. 1839 = P. biaurita* var.

Boivini Moore; Bedd. Ferns S. Ind. t. 36. 1863-65 = Pellaea.

(L) **borbonica** (Fée) Hk. sp. **2**. 228. 1858. (HB. 170) — Réunion. Litobrochia Fée, Gen. 136. 1850-52.

borealis Salisb. Prod. —. 1796 = Pteridium aquilinum.

Boultoni ht. Boult.; Gard. Chr. III. **19**. 652 1896 — Hort.

brasiliensis Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 47 t. 68. 1825 = P. denticulata.

Brazzaiana ht. Lambeau, Rev. Hort. Belge **1903**. 298 f. 66 — (Hort.) Congo. (Eadem ac Compteris Brazzaiana?).

(L) **brevinervis** (Fée) Jenm. W. Ind. and Guiana Ferns 125. 1900 — Guadeloupe. Guiana. — Litobrochia Fée, 11 mém. 27 t. 8 f. 2. 1866.

brevipes Tausch, Flora **1836**^{2}. 427 = Pteridium aquilinum.

(P) **brevisora** Bak. Syn. 162. 1867. Chr. 167. NPfl. 293 — Africa occ. trop. Hypolepis pteridioides Hk. 2 Cent. t. 59. 1861 (non Mett. 1856).

Brownii Desv. Prod. 294. 1827 = Doryopteris concolor.

Brunoniana Endl. Prod. Fl. Norf. 12. 1833 = Histiopteris incisa.

(?) **Buchanani** Bak. ap. Sim, Ferns S. Afr. 111, 112 t. 46. 1892 — Afr. orient. et austr.

(L) **bulbifera** Jenm. Bull. Dep. Jam. **1**. n. 42. 6. 1893 — Jamaica.

Burkeana Hk. sp. **2**. 213 t. 126 B. 1858 = Pellaea dura.

(L) **Burtoni** Bak. Ann. Bot. **5**. 218. 1891 — Guinea.

(P) **Cadieri** Christ, Journ. de Bot. **19**. 72. 1905 — Annam.

caespitosa Wall. List n. 90. 1828 = Cheilanthes Belangeri.

calcarata Bory, Bél. Voy. Bot. **2**. 42. 1833 = P. biaurita*.

calomelanos Sw. Schrad. Journ. **1800**^{2}. 70. 1801 = Pellaea hastata.

(P) **camerooniana** Kuhn, Fil. Afr. 207. 1868 — Africa occ. trop. P. Manniana Mett.; Kuhn, Fil. Afr. 84. 1868 (non P. Mannii Bak. 1867). P. pellucida pt. HB. 154.

campestris Schrad. Gött. gel. Anz. **1824**. 871 = Pteridium aquilinum.

camptocarpa Hk. sp. **2**. 229. 1858 = P. podophylla.

canariensis Pr. Tent. 145. 1836 = Notholaena Marantae.

candifera Zoll. Syst. Verz. 39. 1854 = P. biaurita.

capensis Thbg. Prod. Fl. Cap. 172. 1800 = Pteridium aquilinum*.

capensis Hk. sp. **2**. 145. 1858 = Adiantopsis.

cartilaginea Pr. Rel. Haenk. **1**. 57 t. 9 f. 3. 1825 = Doryopteris rigida.

catoptera Kze. Linn. **18**. 119. 1844 = P. biaurita.

caudata L. sp. **2**. 1075. 1753 = Pteridium aquilinum*.

caudata Lour. Fl. Cochinch. **2**. 835. 1790 = P. ensiformis.

caudata Zoll. et Moritz, Nat. Gen. Arch. Neerl. Ind. **1**. 398. 1844 = P. biaurita.

caudiformis Moritz, Syst. Verz. 118. 1844-45 = P. biaurita.

ceilanica Wikstr. Vet. Akad. Handl. **1825**. 443. 1826 = ? Drymoglossum heterophyllum.

Cervantesii Loud.: Kze. Linn. **23**. 287. 1850 — Mexico.

chamaedryfolia Goldm. Nova Acta **19**. Suppl. I. 458. 1843 = Doryopteris pedata.

Childsii ht. Childs; Gard. Chr. III. **20**. 470. 1896 — Hort.

(P) **chilensis** Desv. Berl. Mag. **5**. 325. 1811. HB. 161. Chr. 169. NPfl. 293 — Chile. Juan Fernandez. — P. tenera Klf. 1824 — P. Fernandeziana Phil. 1857.

chrysocarpa Hk. et Grev. Ic. Fil. t. 107. 1829 = Onychium siliculosum.

chrysocarpa Link, Hort. Berol. **2**. 33. 1833 = P. tremula.

(L) **chrysodioides** (Fée) Hk. sp. **2**. 211. 1858. HB. 168 — »Amer. austr.«
Litobrochia Fée, Gen. 135. 1850-52.
chrysosperma Hk. et Grev. Ic. Fil. t. 107. 1829 = Onychium siliculosum.
(L) **ciliaris** Eat. Mem. Amer. Acad. n. s. **8**. 203. 1860. HB. 170 — Cuba.
ciliata Pr. Tent. 145. 1836 (nomen) — ?
collina Raddi, Opusc. sci. Bol. **3**. 292. 1819 = Doryopteris.
(L) **comans** Forst. Prod. 79. 1786. HB. 171, 480. Chr. 172. NPfl. 294 — Polynesia. Ins. Norfolk. Austr. Tasmania. N. Zealand. Juan Fernandez.
Litobrochia Pr. 1836 — Pteris Endlicheriana Ag. 1839; Litobrochia Fée 1850-52 — Pteris Berteroana Ag. 1839; Litobrochia Fée 1850-52 — Pteris microptera Mett.; Kuhn 1869. — (vide Litobrochia areolata Moore).
(L ?) **commutata** Kuhn, v. Deck. Reis **3**[3]. Bot. 20. 1879 — Africa centr.
concinna Hew. Mag. Nat. Hist. II. **2**. 461. 1838 = P. mutilata var.
concinna Bak. Malesia **3**. 37. 1886 = P. Beccariana.
concolor Langsd. et Fisch. Ic. Fil. 19 t. 21. 1810 = Doryopteris.
confluens Thbg. Prod. Fl. Cap. 171. 1800 = Pellaea auriculata.
connexa J. Sm. JoB. **3**. 405. 1841 = P. tripartita et P. Wallichiana.
(L) **consanguinea** Mett.; Kuhn, Linn. **36**. 89. 1869. (HB. 480) — Venezuela-Ecuador.
consobrina Kze. Linn. **10**. 526. 1836 = Pellaea quadripinnata.
contracta Link, Fil. sp. 53. 1841 = P. cretica.
contracta Mett. Cheil. 5. 1859 = Pellaea quadripinnata.
cordata Cav. Descr. 267. 1802 = Pellaea.
cordata Sieb. Fl. mixt. n. 269 = Pellaea paradoxa.
(P) **Cordemoyi** C. Chr. Ind. 595. 1906 — Réunion.
P. straminea Cord. 1890-91; Fl. Réun. 52. 1895 (non Mett. 1874).
cordifolia Bak. JoB. **1891**. 4 = Doryopteris.
(P) **coriacea** Desv. Prod. 300. 1827. HB. 162. Chr. 169. NPfl. 293 — Costa Rica-Venezuela-Peru. — P. Jamesoni Hk. 1858 — P. muricata Hk. 1858.
coriifolia Kze. Linn. **18**. 120. 1844 = Pteridium aquilinum *capense.
cornuta Pal. Beauv. Fl. d'Ow. **2**. 63 t. 38. (? 1807) = Ceratopteris thalictroides.
costata Bory; Willd. sp. **5**. 367. 1810 = P. longifolia.
crassipes Ag. Rec. 59. 1839 = P. gigantea.
crenata Sw. Schrad. Journ. **1800**[2]. 65. 1801 = P. ensiformis.
crenata Vell. Fl. flum. **11** t. 86. 1827; Arch. Mus. Nac. Rio Jan. **5**. 452. 1881 = Dryopteris gongylodes.
(P) **cretica** L. Mant. 130. 1767. HB. 154 (excl. β). Chr. 165. NPfl. 292 — Mediterr. Ins. Atlant. Abyssinia. Afr. austr. Ins. Mascar. Asia occ. Japonia. China. Ind. bor. et austr. Ceylon. Ins. Philipp. Fiji. Hawaii. N. Zealand-U. S. merid.-Ind. occ. Bermuda-Brasilia. — P. nervosa Thbg. 1784 — P. serraria Sw. 1801 — P. pentaphylla Willd. 1810 — P. triphylla Mart. et Gal. 1842 (non Ag. 1839) — P. lomarioides Col. 1888 — P. melanocaulon Fée 1850-52. P. Treacheriana Bak. 1879. ? P. scabripes Wall. 1828; Ag. 1839 (an = P. pellucida?).
crispa All. Fl. pedem. **2**. 284. 1785 = Cryptogramma.
cristata Hort; Gard. Chr. III. **12**. 249. 1892 — Hort.
croesoides Fée, 10 mém. 17. 1865 (nomen) — Réunion.
(P) **croesus** Bory. Voy. **2**. 192. 1804. HB. 479. Chr. 167. NPfl. 293 — Réunion.
cruciata Klf.; Sieb. Syn. Fil. exs. n. 79 = Histiopteris incisa.
Currori Hk. sp. **2**. 232 t. 140. 1858 = Lonchitis.

(P) **Curtisii** C. Chr. Ind. 596. 1906 — Madagascar.
P. appendiculata Bak. JoB. **1881**. 366 (non Klf. 1824).
cuspidata Thbg. Prod. Fl. Cap. 171. 1800 = Angiopteris sp. fol. ster.
(P) **dactylina** Hk. sp. **2**. 160 t. 130 A. 1858. HB. 155. NPfl. 292 — Ind. bor. China.
(P) **Dalhousiae** Hk. sp. **2**. 170 t. 121 A. 1858. HB. 157 — Penang. Java.
P. rangiferina Pr.; Miq. 1868-69.
daucifolia Roxb. Calc. Journ. **4**. 508. 1844 = ? Cheilanthes tenuifolia.
decipiens Hk. sp. **2**. 209. 1858 = Doryopteris.
decomposita Gaud. Freyc. Voy. Bot. 393 1827 = Pteridium aquilinum.
decomposita Bak. Syn. 479. 1874 = P. Bakeri.
decora Hk. sp. **2**. 210. 1858 = Doryopteris.
(L) **decurrens** Pr. Del. Prag. **1**. 183. 1822. HB. 171. Chr. 171. NPfl. 294 — Brasilia. — Litobrochia Pr. 1836 — L. horizontalis Fée 1869.
decurrens Vell. Fl. flum. **11**. t. 85. 1827; Arch. Mus. Nac. Rio Jan. **5**. 452. 1881 = P. leptophylla.
decursiva Forsk. Fl. Aeg.-Arab. 186. 1775 = Cheilanthes farinosa.
(P) **decussata** J. Sm. JoB. **3**. 405. 1841 (nomen) — Ceylon. Borneo. Celebes. Ins. Phil. Samoa. Ins. Societ. — P. patens Hk. sp. **2**. 177 t. 137. 1858. HB. 165. Chr. 168. NPfl. 293 (non Klf. 1837). — (Conf. P. Gardneri).
(P) **deflexa** Link, Hort. Berol. **2**. 30. 1833. HB. 162. Chr 169. NPfl. 293 — Amer. trop. — P. stridens Ag. 1839 — P. platychlamys Fée 1872-73 — P. lata Klf; Link 1833.
deltea Ag. Rec. 33. 1839 = P. biaurita* var.
(P) **deltodon** Bak. JoB. **1888**. 226 — China.
densa Wall. List n. 99. 1828 = Pteridium aquilinum.
(P) **dentata** Forsk. Fl. Aeg.-Arab. 186. 1775 — Afr. trop. et austr.; var. Ascension. P. flabellata Thbg. 1800. HB. 161. Chr. 168. NPfl. 293 — Lonchitis Adscensionis Forst. 1789; Pteris Sw. 1801; Asplenium Bernh. 1802.
(L) **denticulata** Sw. Prod. 129. 1788. HB. 169. Chr. 171. NPfl. 294 — Ind. occ.-Brasilia. — Litobrochia Pr. 1836. Pteris tristicula Raddi 1819; Litobrochia Pr. 1836 — Pteris brasiliensis Raddi 1819; Litobrochia Pr. 1836 — L. dissimilis Fée 1872-73 — Pteris undulata Christ 1902 (non Litobrochia Moore 1867).
Deplanchei Fourn. Ann. sc. nat. V. **18**. 322. 1873 = P. comans.
diaphana Tausch, Flora **1839**². 479 — Hort.
dichotoma L. sp. **2**. 1076. 1753 = Gleichenia furcata.
dichotoma Cav. (Sw. Syn. 335. 1806) = Adiantopsis.
dichotoma Kuhn, Fil. Deck. 18. 1867 = Actiniopteris australis.
difformis Poir. Enc. **5**. 713. 1804 — Ind. orient.
digitata Wall. List n. 91. 1828 = P. stenophylla.
dimidiata Willd. sp. **5**. 381. 1810 = P. semipinnata.
diplazioides Moritz. Bot. Zeit. **1854**. 855 (nomen) — ? Venezuela.
discolor Kze.; Steud. Nom. **2**. 356. 1824; Spr. Syst. **4**. 76. 1827 = Histiopteris incisa.
dispar Kze. Bot. Zeit. **1848**. 539 = P. semipinnata var.
(P) **dissitifolia** Bak. JoB. **1890**. 263 — Tonkin.
(P) **distans** J Sm. JoB. **3**. 405. 1841 (nomen); Hk. sp. **2**. 169 t. 124 B. 1858. HB. 156. NPfl. 292 — Luzon.
divaricatissima Bak. Syn. ed. II. 163. 1874 = ? Paesia rugosula.
diversifolia Sw. Syn. 96, 288. 1806 = P. longifolia.
dolabriformis Poir. Enc. **5**. 722. 1810 = ? Adiantum pulverulentum.

(L) **dominicensis** Bak. in Hk. Ic. pl. t. 1642. 1886 — Dominica.
Doniana Kuhn, Fil. Afr. 80. 1868 = Pellaea.
Drinkwateri ht.; Gard. Chr. III. **19**. 592. 1896 — Hort.
(?) *Droogmantiana* L. Linden, La Sem. Hort. **4**. 435, 487 f. 162. 1900 — Congo.
(P) **dubia** Kuhn, v. Deck. Reis. **3**^{3}. Bot. 71. 1879 — Ins. Comor.
dura Willd. sp. **5**. 376. 1810 = Pellaea.
echinulata Kze.; Mett. Fil. Lechl. **1**. 12 in obs. 1856 — ?
edentula Kze. Linn. **9**. 75. 1834 = P. biaurita.
edulis Link, Fil. sp. 55. 1841 — Hort.
elastica Tausch in Sieb. Syn. Fil. exs. n. 15 = P. scabra.
elata Ag. Rec. 63. 1839 = P. altissima.
elata var. ? Karsteniana Kze. Linn. **23**. 287. 1850 = P. Kunzeana.
elegans Poir. Enc. **5**. 718. 1804 — ? Ind. orient.
elegans Jacq. Fragm. 74 t. 116. 1809 = P. dentata.
elegans Sw. Vet. Ak. Handl. **1817**. 70 = Histiopteris incisa.
elegans Vell. Fl. flum. **11**. t. 81. 1827; Arch. Mus. Nac. Rio Jan. **5**. 451. 1881 = Doryopteris nobilis.
elliptica Poir. Enc. **5**. 714. 1804 — Ins. Sechellae.
elliptica Willd. sp. **5**. 356. 1810 = Drymoglossum heterophyllum.
elongata Pr. Tent. 145. 1836 (nomen) — ?
emergens Link, Fil. sp. 52. 1841 = P. cretica.
Enderi Regel; Salom. Nom. 336. 1883 — Hort. (P. leptophylla × macroptera).
Endlicheriana Ag. Rec. 66. 1839 = P. comans var.
ensifolia Poir. Enc. **5**. 711. 1804 = P. longifolia.
(P) **ensiformis** Burm. Fl. Ind. 230. 1768. HB. 155. Chr. 166. NPfl. 292 — China. Ind. bor.-Malesia-Polynesia-Austr. trop.
P. crenata Sw. 1801. Phorolobus chinensis Desv. 1827. — var. victoriae Bak. Gard. Chr. III. **7**. 576. 1890; Pteris victoriae ht.
ensiformis Goldm. Nova Acta **19** Suppl. I. 457. 1843 — China.
esculenta Forst. pl. escul. 74. 1786 = Pteridium aquilinum*.
(P) **excelsa** Gaud. Freyc. Voy. Bot. 388. 1827. HB. 159. Chr. 167. NPfl. 292 — Ind. bor.-Malesia-Fiji-Hawaii. — P. terminalis Wall. 1828; Ag. 1839.
excelsa Bl. Enum. 213. 1828 = Pteridium aquilinum.
falcata R.Br. Prod. Fl. Nov. Holl. 154. 1810 = Pellaea.
fallax Mart. et Gal. Mém. Ac. Brux **15**. 53 t. 14 f. 2. 1842 = Pellaea intra- [marginalis var.
farinosa Forsk. Fl. Aeg.-Arab. 187. 1775 = Cheilanthes.
Feei Schaffn.; Fée, 8 mém. 73. 1857 = Pteridium aquilinum.
Feliciennae F. Muell. Fragm. **5**. 124. 1866 = P. tripartita juv.
felosma J. Sm. Comp. Bot. Mag. **72** Comp. 24. 1846 = P. biaurita*.
Fernandeziana Phil. Linn. **29**. 105. 1857 = P. chilensis.
(L) **Finoti** Christ. Journ. de Bot. **19**. 72. 1905 — Annam.
firma Wall. List n. 100. 1828 = Pteridium aquilinum.
flabellata Thbg. Prod. Fl. Cap. 172. 1800 = P. dentata.
flabellata Schkuhr, Kr. Gew. **1**. t. 93. 1809 = P. semipinnata.
flaccida Fée. Gen. 126. 1850-52 = Lonchitis hirsuta.
flava Goldm. Nova Acta **19** Suppl. I. 457. 1843 = P. biaurita*.
flavescens Colla. Mem. Ac. Torino **39**. 37 t. 67. 1836 = Histiopteris incisa.
flexuosa Klf.; Schlecht. et Cham. Linn. **5**. 614. 1830 = Pellaea.
flexuosa Mett. Fil. Lechl. **1**. 12. 1856 = P. semiadnata.
foemina Gray. Nat. Arr. br. Pl. **2**. 16. 1821 = Pteridium aquilinum.

(P) **formosana** Bak. JoB. **1885**. 103 — Formosa.
(P) **Fournieri** C. Chr. Ind. 598. 1906 — N. Caledonia.
P. polymorpha Fourn. Ann. sc. nat. V. **18**. 320. 1873 (non Poir. 1804).
fragrans Lag., Garcia et Clem. Anal. cienc. **5**. 158. 1802 = Cheilanthes pteridioides.
(L) **Fraseri** Mett.; Kuhn. Bot. Zeit. **1868**. 41. HB. 480. Chr. 173. NPfl. 294 — [Ecuador.
(P) **furcans** Bak. JoB. **1888**. 324 — Borneo.
furcata L. sp. **2**. 1073. 1753 = Eschatogramme.
furcata Pr. Tent. 145. 1836 (nomen) — ?
Galeotti Hk. sp. **2**. 204. 1858 = P. biaurita.
(C) *Gardneri* (Fée) Hk. sp. **2**. 206. 1858 — Ceylon. — Litobrochia Fée, Gen. 139. 1850-52; Campteria Moore 1857. (an = P. decussata?).
Gardneri Ettingsh. Denkschr. Ak. Wien **23**. 42. 1864 = Pteridium aquilinum.
(P) **Gaudichaudii** Ag. Rec. 42. 1839 — Brasilia. — P. palustris Gaud. Freyc. Voy. Bot. 391. 1827 (non Poir. 1804). (an P. deflexa?).
(C) **geminata** Wall. List n. 2180. 1829; Ag. Rec. 31. 1839 — Japonia. China. Ind. austr. Ins. Comor. — Campteria J. Sm. 1875; C. Kleiniana Pr. 1836. (Tent. t. 5 f. 19 sine descr.); Litobrochia Fée 1850-52; Pteris Christ 1896; Campteria anamallayensis Bedd. 1863 — Pteris maxima Bak. 1874. HB. 480 (Comor.)
geraniifolia Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 46 t. 67. 1825 = Doryopteris concolor.
Ghiesbreghtii J. Sm. Ferns br. and for. 191. 1866 (Gheisbeghtii) = Lonchitis hirsuta.
(L) **gigantea** Willd. sp. **5**. 381. 1810. HB. 173 — Ind. occ. Columbia-Peru. Litobrochia Pr. 1836 — Pteris crassipes Ag. 1839; Litobrochia Fée 1850-52.
glaberrima Kuhn. Fil. Afr. 81. 1868 = Lonchitis occidentalis.
glabra Mett. Fil. Lips. 59 t. 25 f. 29. 1856 = Lonchitis.
glandulosa Mett. Ann. sc. nat. IV. **15**. 68. 1861 = Gymnogramma.
glauca Cav. Descr. 269. 1802 = Cheilanthes.
glauca Moritz, Bot. Zeit. **1854**. 855 (nomen); Ettingsh. Farnkr. 101 f. 55 t. 65 f. 8. 1865 — Venezuela.
glaucescens Bory in Willd. sp. **5**. 396. 1810 (syn.) = Histiopteris incisa.
glaucovirens Goldm. Nova Acta **19**. Suppl. I. 457. 1843 = P. biaurita.
glaucovirens Linden, Cat. 1858; J. Sm. Ferns br. and for. 190. 1866 — (Hort.)
(L) *Goeldiana* Christ, Hedwigia **44**. 364. 1905 (nomen) — Brasilia austr. Paraguay. Amazonas.
gracilis Michx. Fl. bor. Am. **2**. 262. 1803 = Cryptogramma Stelleri.
gracilis Roxb. Calc. Journ. **4**. 508. 1844 = Cheilanthes tenuifolia.
gracilis Fée, Gen. 128. 1850-52; 7 mém. 32 t. 19 f. 2 = P. leptophylla var.
graminea Poir. Enc. **5**. 708. 1804 = Monogramma.
graminifolia Roxb. Calc. Journ. **4**. 502 t. 33. 1844 = Vittaria elongata.
grandifolia L. sp. **2**. 1073. 1753. HB. 164. Chr. 170. NPfl. 293 — Ind. occ. Mexico-Peru. — Litobrochia J. Sm. 1841; Heterophlebium Fée 1850-52. Pteris socorrensis Karst. 1860.
grandifolia Blanco, Fl Filip. 829. 1837 = ? P. opaca.
grandis Hk. sp. **2**. 229. 1858 = P. Kunzeana.
(P) **Grevilleana** Wall. List n. 2680. 1829; Ag. Rec. 23. 1839; Clarke Tr. Linn. Soc. II. Bot. **1**. 466 t. 54. 1880 — Ind. bor. Siam. Borneo. Luzon.
(P) **Griffithii** Hk. sp. **2**. 170 t. 123 A. 1858. HB. 156 — Mishmee.

Guichenotiana Gaud. Freyc. Voy. Bot. 387. 1827 } = P. longifolia.
Guichenotii Dcne. Nouv. Ann. Mus. **3**. 348. 1834 }
Guilleminei Ag.; Guill. Ann. sc. nat. II. **6**. 314. 1836 = P. tripartita.
guineensis Desv. Prod. 293. 1827 = Vittaria.
(L) **Haenkeana** Pr. Rel. Haenk. **1**. 55. 1825. HB. 169. Chr. 171 — Guatemala. Peru. Litobrochia Pr. 1836. Pteris ampla Kze. 1834; ? Litobrochia Pr. 1836.
Harrisonae Jenm. Gard. Chr. III. **24**. 414. 1898 = Doryopteris.
(L) **Hartiana** Jenm. W. Ind. and Guiana Ferns 130. 1900 — Trinidad.
hastaefolia Schrad. Gött. gel. Anz. **1818**. 917 = Pellaea viridis.
hastata Thbg. Prod. Fl. Cap. 172. 1800 = Pellaea.
hastata Sw. Schrad. Journ. **1800**². 69. 1801 = Pellaea viridis.
hastata Raddi, Pl. bras. **1**. 43 t. 63 f. 2. 1825 = Doryopteris sagittifolia.
hastata Goldm. Nova Acta **19**. Suppl. I. 458. 1843 — Manila.
hederacea Pr. Del. Prag. **1**. 181. 1822 = Doryopteris varians.
hemipteris Hk. sp. **2**. 248. 1858 = P. Orizabae.
(P) **Henryi** Christ, Bull. Boiss. **6**. 957. 1898 — Yunnan.
heptaphyllos Poir. Enc. **5**. 714. 1804 = P. cretica.
Heredia Colmeiro, Enum. Cr. Esp. y Port. **1**. 16. 1867 — Hispania.
(C) **heteroclita** Desv. Prod. 299. 1827 — Mozambique. Ins. Comor. Madagascar. P. triplicata Ag. 1839. HB. 165; Litobrochia Fée 1850-52; Campteria Moore 1861 — Pteris Melleri Bak. Syn. 165. 1867.
heteroclita Bory, Bél. Voy. Bot. **2**. 41. 1833 — Pegu.
heterodactyla Reinw.; J. Sm. JoB. **3**. 405. 1841 = P. ensiformis.
(P) **heteromorpha** Fée, Gen. 127. 1850-52. HB. 156 — Luzon. Birma. Celebes. (f. Bedd. = P. cretica var.).
(C) *heterophlebia* Kze. Bot. Zeit. **1848**. 196. (HB. 172) — Java. Campteria Moore 1857.
heterophylla L. Syst. Nat. ed. X. **2**. 1322. 1759 = Anopteris.
Hillii ht. Hill; Gard. Mag. **1904**. 164 — Hort. (Brasilia).
hirsuta Poir. Enc. **5**. 719. 1804 = Notholaena.
hirsuta J. Sm. JoB. **4**. 165. 1841 = Lonchitis.
hispida Pr. Tent. 145. 1836 = P. biaurita*.
(L) **hondurensis** Jenm. W. Ind. and Guiana Ferns 128. 1900 — Honduras.
(P) **Hookeriana** Ag. Rec. 12. 1839. HB. 155 — Ceylon.
Hostmanniana Ettingsh. Denk. Ak. Wien **23**. 55. 1864 = P. propinqua.
humilis Forst. Prod. 79. 1786 = Cheilanthes tenuifolia.
(L) **hymenophylla** Sod. Cr. vasc. quit. 101. 1893 — Ecuador.
imbricata Cav. Hort. Matr. tab. —; Sw. Syn. 102. 1806 = Jamesonia.
inaequalis Pr. Tent. 145. 1836 (nomen) — ?
inaequalis Fée. 8 mém. 114. 1857 nomen) = P. heteromorpha.
(L **inaequalis** (Fée) Jenm. W. Ind. and Guiana Ferns 124. 1900 — Trinidad. Guiana. — Litobrochia Fée, 11 mém. 26 t. 8 f. 1. 1866.
inaequalis Bak. JoB. **1875**. 199 = P. semipinnata var.
incisa Thbg. Prod. Fl. Cap. 171. 1800 = Histiopteris.
incompleta Cav. Anal. Hist. Nat. **4**. 107. 1801 = P. arguta.
indica Poir. Enc. **5**. 712. 1804 = P. longifolia vel P. moluccana.
Indica Christ, Farnkr. 165. 1897 = P. moluccana.
inframarginalis Link. Hort. Berol. **2**. 34. 1833 = Pellaea intramarginalis.
(P) **insignis** Mett.; Kuhn. JoB. **1868**. 269 — China.
intermedia Bl. Enum. 211. 1828 = P. tripartita.

intermedia Klf.; Ettingsh. Farnkr. 101 t. 66 f. 6. 1865 = P. macroptera.
interrupta Willd. Phyt. 13 t. 10 f. 1. 1794 = Dryopteris pteroides.
intramarginalis Klf.; Schlecht. et Cham. Linn. **5**. 613. 1830 = Pellaea.
involuta Sw. Schrad. Journn. **1800**[2]. 69. 1801 = Pellaea.
(P) **irregularis** Klf. Enum. 189. 1824. HB. 158. NPfl. 292 — Ins. Hawaii.
P. alata Gaud. 1827.
itatiaiensis Szysz. in Beck, It. Princ. Cob. **2**. 119. 1888 = Doryopteris coriacea var.
Jamesoni Hk. sp. **2**. 193 t. 133 A. 1858 = P. coriacea var.
japonica Mett. Fil. Lips. 54. 1856 = Onychium.
(L) **Johnstoni** Bak. Ann. Bot. **5**. 218. 1891 — Sierra Leone.
(L) **Junghuhnii** (Reinw.) Bak. Syn. 170. 1867 — Java.
Campteria Reinw. msc.
Kingiana Endl. Prod. Fl. Norf. 13. 1833 = P. tremula.
Kingiana Lowe, Ferns **3** t. 46. 1857 = P. arguta.
Kleiniana Christ, Bull. Boiss. **4**. 666. 1896 = P. geminata.
(L) **Kunzeana** Ag. Rec. 62. 1839. HB. 173 — Ind. occ. Mexico-Peru-Brasilia.
Litobrochia Fée 1850-52 — L. grandis Fée 1857; Pteris Hk. 1858 — P. arborescens Mart. et Gal. 1842; Litobrochia Fée 1850-52 — Pteris protea Liebm. 1849.
laciniata Willd. sp. **5**. 397. 1810 = Lonchitis hirsuta.
laciniata Vell. Fl. flum. **11** t. 89. 1827; Arch. Mus. Nac. Rio Jan. **5**. 453. 1881 = Doryopteris concolor.
(L) **lactuca** Poir. Enc. **5**. 717. 1804 — Ind. occ. — Litobrochia Pr. 1836.
laeta Wall. List n. 95. 1828 = P. cretica.
laeta Klf.; Pr. Tent. 145. 1836 = P. deflexa.
(P) **laevis** Mett. Ann. sc. nat. IV. **15**. 294. 1861. (HB. 171). NPfl. 294 — N. Caledonia.
(L) **lanceaefolia** Ag. Rec. 53. 1839. HB. 168. NPfl. 294 — Madagascar.
Litobrochia J. Sm. 1875.
lanceolata L. sp. **2**. 1073. 1753 = Paltonium.
lanceolata Lour. Fl. Cochinch. **2**. 834. 1790 — Cochinchina.
lanceolata Desf. Fl. Atlant. **2**. 401. 1800 = P. longifolia.
lanigera Bl. Enum. 214. 1828 = Pteridium aquilinum.
lanuginosa Bory; Willd. sp. **5**. 403. 1810 = Pteridium aquilinum *capense.
lanuginosa Spr. Nova Acta **10**. 231 t. 16 f. 1-2. 1821 = Pteridium aquilinum.
(L) **Lastii** C. Chr. Ind. 600. 1906 — Madagascar.
P. acuminata Bak. JoB. **1891**. 5 (non Desv. 1811).
lata Klf.; Link, Hort. Berol. **2**. 28. 1833 = P. deflexa.
latifolia H.B.Willd. sp. **5**. 370. 1810 = Amphiblestra.
latiuscula Desv. Prod. 303. 1827 = Pteridium aquilinum.
(L) **laurea** Desv. Prod. 299. 1827. HB. 168 — Madagascar.
(C?) Campteria Moore 1861. Pteris triphylla Ag. 1839.
(L) **Lechleri** Mett. Fil. Lechl. **2**. 13. 1859 — Peru.
lendigera Cav. Descr. 268. 1802 = Cheilanthes.
(L?) **leptophylla** Sw. Vet. Ak. Handl. **1817**. 70. HB. 171. Chr. 170. NPfl. 294 — Brasilia. Columbia. — Litobrochia J. Sm. 1846 — Pteris spinulosa Raddi 1819; Cheilanthes Link 1833; Litobrochia J. Sm. 1841 — Pteris gracilis Fée 1850-52. HB. 160 (non Michx. 1803) — P. semidentata Fée 1857.
Lessonii Bory, Dup. Voy. Bot. **1**. 276. 1828 = Histiopteris incisa.
leucomelas Mett.; Kuhn, Fil. Afr. 83. 1868 = Pellaea.
(P) **ligulata** Gaud. Freyc. Voy. Bot. 385. 1827 — Moluccae. Borneo. N. Guinea.

Lindeniana Christ, Prim. Fl. Costaric. **3**. 22. 1901 = Lonchitis aurita.
linearis Poir. Enc. **5**. 723. 1804 = ? P. tripartita.
linearis Wall. List n. 105. 1828 = P. tripartita.
linearis Roxb. Calc. Journ. **4**. 505. 1844 (an Wall.?) — Amboyna.
lineata L. sp. **2**. 1073. 1753 = Vittaria.
lineata (Pr.) Ettingsh. Farnkr. 96 t. 56 f. 3, t. 57 f. 7. 1865 — Luzon. Heteropteris Pr. hb.
(P) **litobrochioides** Kl. Linn. **20**. 341. 1847. HB. 157 — Guiana. Para. Ecuador.
(L) **livida** Mett. Ann. sc. nat. V. **2**. 222. 1864. HB. 481 — Columbia.
lobata Roxb. Calc. Journ. **4**. 504. 1844 — Moluccae.
lobata Goldm. Nova Acta **19** Suppl. I. 457. 1843 — Manila.
lomariacea Kze. Linn. **22**. 576. 1849 = Doryopteris.
lomariacea γ digitato-palmata Hk. sp. **2**. 133. 1858 = Doryopteris ornithopus.
lomarioides Col. Tr. N. Zeal. Inst. **13**. 380. 1881 = P. cretica.
lonchitoides Desv. Prod. 301. 1827 — Peru.
lonchophora Mett. Cheil. 4 t. 3 f. 1-3. 1859 = Doryopteris.
(L) **longibrachiata** Ag. Rec. 75. 1839; Bak. Ann. Bot. **5**. 165. 1891 — St. Vincent. Martinique. — Litobrochia Fée 1866.
longibracteata Bak. Syn. 173. 1867 (err. scr.) = P. longibrachiata.
(P) **longicauda** Christ, Prim. Fl. Costar. **3**. 21. 1901 — Costa Rica.
(P) **longifolia** L. sp. **2**. 1074. 1753. HB. 153. Chr. 165. NPfl. 292 — Mediterr. Ins. Atlant.-Afr. austr.-China. Japonia. Asia trop. Austr. Polynesia. N. Zealand. Ind. occ. Amer. centr.-Venezuela. — P. vittata L. 1753 — P. ensifolia Poir. 1804 — P. costata Bory; Willd. 1810 — P. diversifolia Sw. 1806.
longifolia Cav. Descr. 269. 1802 — S. Domingo.
(P) **longipes** Don, Prod. Fl. Nepal. 15. 1825. HB. 161. NPfl. 293 — Asia trop. P. pellucens Ag. 1839 — P. Zollingeri Mett.; Miq. 1868-69.
longipes Bl. Enum. 212. 1828 = P. tripartita.
(P) **longipinnula** Wall. List n. 108. 1828; Ag. Rec. 19. 1839. HB. 158 — Japonia. China. Ind. bor. Birma. Malesia. Samoa. ? Jamaica (Jenman). — P. umbraculifera Mett.; Miq. 1868-69.
longula Mett.; Kuhn, Linn. **36**. 88. 1869 = ? Doryopteris subsimplex.
lorigera Wall. List n. 103. 1828 = Pteridium aquilinum.
lucida Cav. Descr. 266. 1802 = Adiantum.
ludens Wall. List n. 88. 1828 = Doryopteris.
lunulata Retz. Obs. **2**. 28 t. 4. 1781 = Adiantum.
(L) **Luschnathiana** (Kl.) Bak. Syn. 172. 1867 — Brasilia. Campteria Kl. 1840 (nomen).
lutea Cav. Descr. 267. 1802 = Cheilanthes.
luxuriosa Kze. Linn. **23**. 289, 321. 1850 — Hort.
(P) **Lydgatei** (Hill.) Christ. Farnkr. 167. 1897. NPfl. 293 — Ins. Hawaii. Schizopteris Hill msc.; Cheilanthes Lidgatii Bak. Syn. 475. 1874; Schizostege Lydgatei Hill. Fl. Haw. 632. 1888.
(L) **macilenta** A. Rich. Fl. N. Zél. 82 t. 12. 1832. HB. 171. Chr. 171. NPfl. 294. N. Zealand. — Litobrochia J. Sm. 1841 — Pteris pendula Col. 1888.
macrocarpa Ettingsh. Farnkr. 100 t. 70. f. 5. 1865 = P. macroptera.
(P) **macrodon** Bak. Journ. Linn. Soc. **15**. 414. 1876 — Madagascar.
(L) **macroptera** Link. Hort. Berol. **2**. 32. 1833. HB. 169. NPfl. 294 — Brasilia. Litobrochia J. Sm. 1841. L. angustata Fée 1869.
macroura Willd. sp. **5**. 380. 1810 = P. pungens.

(P) **madagascarica** Ag. Rec. 17. 1839. HB. 156 — Madagascar.
madagascariensis Kuhn, Fil. Afr. 84. 1868 = Lonchitis.
Maissonieri ht. Sander, Gard. Chr. III. **33**. 266. 1903 — Hort.
(P. multifida × tremula).
Manniana Mett.; Kuhn, Fil. Afr. 84. 1868 = P. camerooniana.
(L) **Mannii** Bak. Syn. 168. 1867. NPfl. 294 — Fernando Po.
Litobrochia J. Sm. 1875.
marginata Bory, Voy. **2**. 192. 1804 = P. tripartita.
marattiaefolia Hk. sp. **2**. 177 t. 122 B. 1858 = P. semiadnata.
mascarenensis Spr. Syst. **4**. 75. 1827 = P. scabra.
mauritiana Willem. Usteri Ann. **18**. 61. 1796 — Mauritius.
maxima Bak. Syn. 480. 1874 = P. geminata.
megaphylla Mett.; Miq. Ann. Lugd. Bat. **4**. 96. 1868-69 = P. biaurita*.
melanocaulon Fée, Gen. 127. 1850-52; 7 mém. 31 t. 19 f. 1 = P. cretica var.
melanolepis Ettingsh. Farnkr. 88 f. 35, 39. 1865 = Onychium.
Melleri Bak. Syn. 165. 1867 = P. heteroclita var.
(?) **mertensioides** Willd. sp. **5**. 394. 1810 — Amboyna. Java. Borneo.
metallica ht. May; Gard. Chr. III. **34**. 64. 1903 — Hort.
(C) **Mettenii** Kuhn, Fil. Afr. 85. 1868 — Ins. Comor. Madagascar.
(L) **mexicana** (Fée) Fourn. Mex. pl. 117. 1872 — Mexico.
Litobrochia Fée, Gen. 136. 1850-52 (an = P. Haenkeana?). Pteris pulchra Bak. Syn. 169. 1867 (pt.?). NPfl. 294.
(L) *microdictyon* (Fée) Hk. sp. **2**. 228. 1858 — Ins. Philipp.
Litobrochia Fée, Gen. 138. 1850-52.
microdonta Gaud. Freyc. Voy. Bot. 387. 1827 = P. longifolia.
microphylla Cav. (Sw. Syn. 324. 1806) = Cheilanthes micropteris.
microphylla Colla, Mem. Ac. Torino **39**. 38. 1836 — Chile.
microphylla A. Cunn. in Hk. Comp. Bot. Mag. **2**. 366. 1836 = Paesia scaberula.
microptera Mett.; Kuhn, Linn. **36**. 92. 1869 = P. comans.
(L) **Milneana** (Hk.) Bak. Syn. 170. 1867. Chr. 168. NPfl. 294 — Polynesia. Austr. trop. — P. tripartita γ Milneana Hk. sp. **2**. 226 t. 138 B. 1858; Litobrochia Milneana Carr. 1873.
minuta Turcz. Bull. Soc. Nat. Moscou **1838**. 105 = Cryptogramma Stelleri.
misera Lodd. Cat. 1849 — Hort.
(P) **mixta** Christ in Schum. et Laut. Nachtr. Fl. deutsch. Südsee 45. 1905 — N. Guinea.
(L) **mollis** Christ, Bull. Boiss. **4**. 658. 1896. NPfl. 293 — Costa Rica.
(P) **moluccana** Bl. Enum. 208. 1828. HB. 154. NPfl. 292 — Malesia. Ins. Salomon.
P. indica Christ 1897. 165 (an Poir. 1804?).
montana Col. Tasm. Journ. **2**. 172. 1844 = Histiopteris incisa.
(L) **Montbrisonis** (Fée) Hk. sp. **2**. 228. 1858 — Réunion.
Litobrochia Fée, Gen. 137. 1850-52.
Moerenhoutii Ag.; Guill. Ann. sc. nat. II. **6**. 314. 1836 } = Histiopteris incisa.
Morrenhouttiana Ag. Rec. 79. 1839 }
multiaurita Ag. Rec. 12. 1839 = P. cretica.
multidentata Wall. List n. 2681. 1829 = P. ensiformis.
(P) **multifida** Poir. Enc. **5**. 714. 1804 — China. Japonia. — P. serrulata L.fil. Suppl. 445. 1781. HB. 155. Chr. 166. NPfl. 292 (non Forsk. 1775).
multifida Roxb. Calc. Journ. **4**. 507. 1844 = Doryopteris ludens.
multifida × tremula = P. Maissonieri.
(L) **multiserialis** Jenm. W. Ind. and Guiana Ferns 129. 1900 — Trinidad.

muricata Hk. sp. **2**. 193 t. 123 B. 1858 = P. coriacea var.

(P) **muricella** Fée, 8 mém. 73. 1857 — Mexico.

(P) **mutilata** L. sp. **2**. 1076. 1753. HB. 156. NPfl. 292 — Ind. occ. — P. concinna Hew. 1838.

mysorensis Fée, 10 mém. 16 t. 32 f. 2. 1865 = P. cretica.

mysurensis Heyne; Wall. List n. 87. 1828 = Doryopteris concolor.

(L) **Nadeaudii** Drake, Fl. Polyn. franç. 287. 1893 — Tahiti.
P. aculeata Nadeaud, Enum. pl. Tahiti »n. 129«. 1873 (non Sw. 1788; an = P. Beecheyana?).

(P) **nana** Christ in C. Chr. Ind. 603. 1906 - Yunnan.
P. trifoliata Christ, Bull. Boiss. **7**. 7. 1899. NPfl. 292 (non Fée 1857).

Natalensis Kuhn, Fil. Afr. 85. 1868 = Lonchitis glabra.

nemoralis Willd. Enum. 1073. 1809; sp. **5**. 386. 1810 = P. biaurita.

nervosa Thbg. Fl. Jap. 332. 1784 = P. cretica.

nervosa Wall. List n. 96. 1828 = P. pellucida.

(L) **Nevillei** Bak. Ann. Bot. **5**. 219. 1891 — Réunion.

nigra Retz. Obs. **6**. 38. 1791 = Cheilanthes tenuifolia.

nigrescens Poir. Enc. Suppl. **4**. 607. 1816 = P. croesus.

nigricans Willd. sp. **5**. 396. 1810 = Notholaena.

nitida R.Br. Prod. Fl. N. Holl. 155. 1810 = Pellaea.

nitida Mett.; Kuhn, Fil. Afr. 86. 1868 = P. repens.

nitidula Wall. List n. 89. 1828 = Pellaea.

nivea Poir. Enc. **5**. 718. 1804 = Pellaea.

nivea Bl. Enum. 213. 1828 = Histiopteris incisa.

normalis Don, Prod. Fl. Nepal. 15. 1825 = P. biaurita*.

notholaenoides Desv. Prod. 299. 1827 = Cheilanthes microphylla.

(P) **novae caledoniae** Hk. sp. **2**. 189. 1858. HB. 159. Chr. 168. NPfl. 293 — N. Caledonia.

nudicaulis Güldenst. Reise **1**. 421. 1787 = Pteridium aquilinum.

nudiuscula R.Br. Prod. Fl. N. Holl. 155. 1810 α = Cheilanthes tenuifolia.
β = Notholaena hirsuta.

obliqua Forsk. Fl. Aeg. Arab. 185. 1775 = P. longifolia.

(L) **obscura** Mett.; Kuhn, Linn. **36**. 90. 1869. HB. 480 — Venezuela.

occidentalis Kuhn, Fil. Afr. 207. 1868 = Lonchitis.

(C) **oligodictyon** Bak. JoB. **1880**. 328. NPfl. 293 — Madagascar.

oligophylla Viv. Ann. Bot. **1**^{2}. 189. 1804 = P. cretica.

(P) **opaca** J. Sm. JoB. **3**. 405. 1841 (nomen). HB. 154. Chr. 165. NPfl. 292 — Ins. Philipp. Celebes. — Pycnodoria Pr. Epim. 101. 1849.

ophioderma Fée, 10 mém. 15. 1865 = P. longifolia.

ophioglossoides Vell. Fl. flum. **11**. t. 84. 1827; Arch. Mus. Nac. Rio Jan. **5**. 451. 1881 = Polybotrya cervina.

oppositi-pinnata Fée, 10 mém. 17 t. 33 f. 3. 1865 = P. biaurita.

orbiculata Houtt. Nat. Hist. **14**. 108. 1783; Pfl. Syst. **13**^{1}. 120 t. 96 f. 3. 1786 = Pellaea pteroides.

orbiculata Poir. Enc. **5**. 710. 1804 = Jamesonia imbricata.

(L) **orizabae** Mart. et Gal. Mém. Ac. Brux. **15** 53 t. 13. 1842 — Mexico-Columbia. — Litobrochia Fée 1850-52.

ornithopus Mett.: Bak. Syn. 166. 1867 = Doryopteris.

osmundoides Bory. Voy. **2**. 194 t. 32. 1804 = Blechnum tabulare.

Otaria Bedd. Ferns S. Ind. t. 41. 1863-65 = P. biaurita*.

ovata Desv. Prod. 301. 1827 — Peru.
owahiensis Pr. Tent. 145. 1836 (nomen) — ?
(P) **paleacea** Roxb. in Beats. St. Helena 319. 1816; Ac. Rec. 33. 1839. HB. 159. NPfl. 293 — St. Helena.
pallens Mett. Fil. Lips. 54 t. 15 f. 7-10. 1856 = Ochropteris.
pallida Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 49 t. 71. 1825 = Histiopteris incisa.
palmaeformis Thouars, Fl. Trist. d'Ac. 30. 1804 = Blechnum tabulare.
palmata Willd. sp. **5**. 357. 1810 = Doryopteris pedata var.
palmata Vell. Fl. flum. **11**. t. 87. 1827; Arch. Mus. Nac. Rio Jan. **5**. 452. 1881 = P. dendiculata.
palustris Poir. Enc. **5**. 722. 1804 = P. arguta.
palustris Raddi, Opusc. sci. Bol. **3**. 293. 1819 = P. pilosiuscula.
palustris Gaud. Freyc. Voy. Bot. 391. 1827 = P. Gaudichaudii.
(P) **Pancheri** Bak. Syn. 479. 1874 — N. Caledonia.
(P) **papuana** Ces. Rend. Ac. Napoli **16**. 25, 29. 1877 — N. Guinea.
papyracea Hk. sp. **2**. 212. 1858 = P. splendens var.
paradoxa Schrad. Gött. gel. Anz. **1824**. 871 — Brasilia. (Doryopteris sp. prob. = D. lomariacea).
paradoxa Desv. Prod. 294. 1827 = Pellaea.
paradoxa Kuhn, Hedwigia **35**. 160. 1896 = Doryopteris.
patens Kze. Anal. 28. 1837 = Histiopteris incisa.
patens Hk. sp. **2**. 177 t. 137. 1858 = P. decussata.
(P) **paucinervata** Fée, 8 mém. 73. 1857. HB. 157 — Mexico.
(L) **Pearcei** Bak. Ann. Bot. **5**. 218. 1891 — »Amer. austr.« (Brasilia?).
pectinata Cav. Descr. 266. 1802 — Ins. Mascarenae.
pectinata Desv. Berl. Mag. **5**. 324. 1811 = P. plumula.
pectinata Don, Prod. Fl. Nepal. 15. 1825 = P. biaurita*.
pectinata Roxb. Calc. Journ. **4**. 507. 1844 — Moluccae.
pectiniformis Goldm. Nova Acta **19** Suppl. I. 458. 1843 — Brasilia.
pectiniformis Godet; Kuhn, Fil. Afr. 87. 1868 = Pellaea Goudotii.
pedata L. sp. **2**. 1075. 1753 = Doryopteris.
pedata R.Br. Prod. Fl. N. Holl. 155. 1810 = Doryopteris concolor.
pedata Hk. et Arn. Bot Beech. Voy. 107. 1832 = Doryopteris decipiens.
pedata Kuhn, Hedwigia **35**. 161. 1896 = P podophylla.
pedata var. β et γ Raddi, Opusc. sci. Bol. **3**. 293. 1819 = Doryopteris [Raddiana.
pedatifida Roxb. Calc. Journ. **4**. 508. 1844 — Malesia.
pedatoides Desv. Prod. 293 t. 7 f. 2. 1827 = Doryopteris.
pellucens Ag. Rec. 43. 1839 = P. longipes.
(P) **pellucida** Pr. Rel. Haenk. **1**. 55. 1825. HB. 154 — Ind. bor.-Java-Ins. Philipp. P. venulosa Bl. 1828. P. venusta Kze. 1848.
pellucida Bak. Syn. 154. 1867 quoad pl. afric. = P. camerooniana.
pellucida Klf.; Spr. Syst. **4**. 73. 1827 = P. woodwardioides.
pendula Col. Tr. N. Zeal. Inst. **20**. 218. 1888 = P. macilenta var.
pentaphylla Willd. sp. **5**. 362. 1810 = P. cretica.
peruviana Poir. Enc. **5**. 718. 1804 = Pellaea ternifolia.
peucedanifolia Schrad. Gött. gel. Anz. **1824**. 871 = P. leptophylla.
phanerophlebia Bak. JoB. **1881**. 367 = Doryopteris.
(P) **philippinensis** Fée, 8 mém. 114. 1857; 10 mém. 17 t. 33 f. 2. 1865 — [Ins. Philipp.
pilosa Poir. Enc. **5**. 717. 1804 = Doryopteris.

piloselloides L. sp. ed. II. **2**. 1530. 1763 = Drymoglossum heterophyllum.
piloselloides Thbg. Fl. Jap. 331. 1784 = Drymoglossum microphyllum.
piloselloides Blanco, Fl. Filip. 830. 1837 = ? Nephrolepis biserrata.
pilosiuscula Desv. Prod. 296. 1827 — Brasilia.
P. palustris Raddi, Opusc. sci. Bol. **3**. 293. 1819 (non Poir. 1804).
pinnata Mett. Fil. Lips. 55. 1856 = Pellaea.
? platychlamys Fée, Cr. vasc. Br. **2**. 25. 1872-73 = P. deflexa.
(L) **platyodon** Bak. Journ. Linn. Soc. **15**. 415. 1876 — Madagascar.
(L) **platypteris** Sod. Rec. 28. 1883; Cr. vasc. quit. 109. 1893 — Ecuador.
(P) **platysora** Bak. JoB. **1880**. 211. NPfl. 292 — Sumatra.
Plumeria »H. S.«; Ettingsh. Farnkr. 95 t. 61 f. 3. 1865 = P. biaurita*.
Plumierii Link, Enum. alt. **2**. 462. 1822 = P. biaurita*.
plumula Desv. Prod. 297. 1827 — Ind. occ.
P. pectinata Desv. Berl. Mag. **5**. 324. 1811 (non Cav. 1802).
(L) **podophylla** Sw. Schrad. Journ. **1800**[2]. 67. 1801. HB. 172. Chr. 172. NPfl 294 — Amer. trop. — Litobrochia Pr. 1836; Lonchitis pedata L. sp. ed. II. 1536. 1763; Pteris Kuhn 1896 (non L. 1753) — P. pulchra Schlecht. et Cham. 1830. ? HB. 169 (pt.?). NPfl. 294; Litobrochia Pr. 1836 — Pteris apicalis Liebm. 1849; Litobrochia setifera Fée 1850-52; Pteris Hk. 1858 — Litobrochia camptocarpa Fée 1850-52; Pteris Hk. 1858.
podophylla Kze. Linn. **9**. 75. 1834 = P. Kunzeana.
Pohliana Pr. Del. Prag. **1**. 181. 1822 = Doryopteris concolor.
polita Wall. List n. 92. 1828 = Schizoloma ensifolium.
(L) *polita* Link, Hort. Berol. **2**. 30. 1833. (HB. 171) — Brasilia.
Litobrochia J. Sm. 1841. (an = P. Kunzeana an P. propinqua?).
polymorpha Poir. Enc. **5**. 719. 1804 = Pellaea viridis.
polymorpha Fourn. Ann. sc. nat. V. **18**. 320. 1873 = P. Fournieri.
(L) *polyphylla* (Pr.) Ettingsh. Denkschr. Ak. Wien **23**. 53 t. 5 f. 1. 1864 — Brasilia. — Litobrochia Pr. hb.
polypodioides Poir. Enc. **5**. 716. 1804 = Dryopteris unita.
polytoma Kze. Linn. **23**. 289, 322. 1850 = Doryopteris pedata.
Presliana Ag. Rec. Index 83. 1839 = P. biaurita*.
Prionitis Fée, Gen. 126. 1850-52 = P. ensiformis.
(L) **propinqua** Ag. Rec. 65. 1839. (HB. 171) — Amer. trop.
Litobrochia J. Sm. 1854. (an = P. polita Link 1833?).
propinqua J. Sm. JoB. **3**. 405. 1841 = P. heteromorpha.
protea Liebm. Vid. Selsk. Skr. V. **1**. 228 (seors. 76). 1849 = P. Kunzeana
Pseudo-Lonchitis Bory; Willd. sp. **5**. 389. 1810 = P. tripartita.
Pseudo-Lonchitis Ettingsh. Farnkr. 97 t. 61 f. 6. 7. 1865 = P. Mettenii.
psittacina Pr. Del. Prag. **1**. 185. 1822 = Pteridium aquilinum.
pubescens Kuhn, Fil. Afr. 88. 1868 = Lonchitis.
pulchra Vell. Fl. flum. **11**. t. 83. 1827; Arch. Mus. Nac. Rio Jan. **5**. 451. 1881 = Dryopteris opposita.
pulchra Schlecht. et Cham. Linn. **5**. 614. 1830 = P. podophylla.
pulchra Bak. Syn. 169. 1867 = P. mexicana et ? P. podophylla.
punctata Fée, 10 mém. 18 t. 32 f. 4. 1865 = P. scabra.
(P) **pungens** Willd. sp. **5**. 387. 1810. HB. 159. (Chr. 167) — Ind. occ.
pyrophylla Bl. Enum. 212. 1828 = P. biaurita*.
quadrialata Willd. sp. **5**. 385. 1810 } = P. biaurita*.
quadriaurita Retz. Obs. **6**. 38. 1791 }

quadriaurita var. digitata Bak. JoB. **1879**. 40 = P. Grevilleana.
quadrifida Pr. Del. Prag. **1**. 182. 1822 = P. denticulata.
quadrifoliata L. Syst. Nat. ed. X. **2**. 1321. 1759 = Marsilea quadrifolia.
quadripinnata Forsk. Fl. Aeg.-Arab. 186. 1775 = Pellaea.
quinquelobata Bak. Ann. Bot. **5**. 215. 1891 = Doryopteris.
radiata Boj. Hort. Maur. 399. 1837 = Actiniopteris australis.
(P) **radicans** Christ. Verh. Nat. Ges. Basel **11**. 431. 1896; Ann. Jard. Buit. **15**. 110. 1897. NPfl. 293 — Celebes.
radula Bak. JoB. **1880**. 211 = Paesia.
rangiferina Pr.; Miq. Ann. Lugd. Bat. **4**. 95. 1868-69 = P. Dalhousiae.
recurvata Wall. List n. 113. 1828 = Pteridium aquilinum.
reducta Bak. JoB. **1880**. 211 = P. biaurita*.
(L) **regia** Jenm. Gard. Chr. III. **17**. 39. 1895 — Jamaica.
reginae ht.; Gard. Chr. III. **12**. 249. 1892 — Hort.
regularis Forsk. Fl. Aeg.-Arab. 186. 1775 — Arabia.
regularis Bail. Haw. Ferns 26. 1883 — Ins. Hawaii.
(P) **remotifolia** Bak. Journ. Linn. Soc. **16**. 199. 1877 — Madagascar.
repandula Link, Fil. sp. 56. 1841 = P. biaurita*.
(P) **repens** C. Chr. Ind. 606. 1906 — Africa occ. trop.
P. nitida Mett.; Kuhn, Fil. Afr. 86. 1868. HB. 479 (non R.Br. 1810).
resistens Mett. Fil. Lips. 59. 1856 = Paesia viscosa.
reticulata Desv. Berl. Mag. **5**. 324. 1811 = P. denticulata.
reticulata Mett.; Kuhn, Linn. **36**. 91. 1869 = P. spinosa var.
revoluta Bl. Enum. 214. 1828 = Pteridium aquilinum.
revolvens Ac. Rec. 74. 1839 = P. tripartita.
rigida Sw. Syn. 104, 299. 1806 = Doryopteris.
Rochfordi ht. Rochf.; Gard. Chr. III. **24**. 241. 1898 — Hort.
rostrata Fée, 10 mém. 16 t. 33 f. 1. 1865 = P. Gaudichaudii.
Rottleriana Link, Fil. sp. 48. 1841 = P. biaurita.
rotundifolia Forst. Prod. 79. 1786 = Pellaea.
ruffa L. sp. **2**. 1074. 1753 = Gymnopteris rufa.
rugosula Lab. Sert. Austr. Cal. 6 t. 8. 1824 = Paesia.
rugulosa auctt. = rugosula.
sagittata Cav. Descr. 267. 1802 = Pellaea cordata var.
sagittifolia Raddi, Opusc. sci. Bol. **3**. 292. 1819 = Doryopteris.
salicifolia Schrad. Gött. gel. Anz. **1824**. 871 = ? P. denticulata.
scaberula Rich. Fl. N. Zél. 82 t. 11. 1832 = Paesia.
(P) **scabra** Bory; Willd. sp. **5**. 386. 1810. HB. 159. Chr. 167 — Ins. Mascar. Madagascar. — P. angustata Bory; Willd. 1810 — P. sexalata Desv. 1827 — P. punctata Fée 1865.
scabripes Wall. List n. 94. 1828 = P. cretica vel P. pellucida.
scalaris Moritz, Bot. Zeit. **1854**. 856; Mett. Pteris 9. n. 2. t. 5. (16). 1858 = Paesia viscosa.
scandens Roxb. Calc. Journ. **4**. 505. 1844 = Stenochlaena palustris.
Schiedeana Ettingsh. Farnkr. 99 t. 66 f. 1. 1865 = P. Kunzeana.
(L) **Schwackeana** Christ in Schwacke, Pl. Nov. Mineiras **2**. 97. 1900; Bull. Boiss. II. **2**. 551. 1902 — Brasilia.
(L) **sclerophylla** Sod. Cr. vasc. quit. 101. 1893 — Ecuador.
scolopendrina Bory, Voy. **2**. 323. 1804 = Vittaria.

(P) **semiadnata** Phil. Linn. **29**. 106. 1857 — Chile. Chiloe. — P. flexuosa Mett. 1856 (non Klf. 1830); P. marattiaefolia Hk. 1858. HB. 157. NPfl. 292.
semidentata Fée, 7 mém. 32 t. 18 f. 3. 1857 = P. leptophylla var.
semihastata Wall. List n. 102. 1828 = Pteridium aquilinum.
semihirta Link, Fil. sp. 50. 1841 = P. longifolia.
semiovata Poir. Enc. **5**. 723. 1804 = P. tripartita.
(P) **semipinnata** L. sp. **2**. 1076. 1753. HB. 157. Chr. 166. NPfl. 292 — Japonia. China. Asia trop. — P. dispar Kze. 1848 — P. inaequalis Bak. 1875 (non Jenm. 1900).
semisagittata ht.; Wien. Ill. Gartenzeit. **1902**. 203 — Hort. (Brasilia austr.).
semiserrata Forsk. Fl. Aeg.-Arab. 186. 1775 = P. cretica.
semiserrata Roxb. in Beats St. Helena 319. 1816 = P. dentata.
(L) **sericea** (Fée) Christ, Bull. Boiss. II. **2**. 551. 1902 — Brasilia. Litobrochia Fée, Cr. vasc. Br. **1**. 48 t. 11 f. 3. 1869.
serraria Sw. Schrad. Journ. **1800**^{2}. 65. 1801 = P. cretica.
serrata Gmel. Syst. Nat. **2**^{2}. 1299. 1791 = P. serrulata Forsk.
serrata Vell. Fl. flum. **11**. t. 82. 1827; Arch. Mus. Nac. Rio Jan. **5**. 451. 1881 = P. denticulata.
serrulata Forsk. Fl. Aeg.-Arab. 187. 1775 = P. arguta vel P. dentata.
serrulata L.fil. Suppl. 445. 1781 = P. multifida.
seticaulis Hk. Ic. pl. t. 209. 1840 = Pellaea falcata.
setifera Hk. sp. **2**. 224. 1858 = P. podophylla var.
sexalata Desv. Prod. 298. 1827 = P. scabra.
Sieberi Tausch, Flora **1834**. 526 — Hort.
signata Blanco, Fl. Filip. 830. 1837 = Polypodium.
siliculosa Desv. Berl. Mag. **5**. 324. 1811 = Onychium.
(L) **similis** Kuhn. v. Deck. Reis. **3**^{8}. Bot. 21. 1879 — Africa centr.
sinuata Thbg. Fl. Jap. 332. 1784 — Japonia. — (Cheilanthes sp.).
sinuata Wall. List n. 84. 1828; Hk. sp. **2**. 232. 1858 = Histiopteris incisa.
sinuata Blanco, Fl. Filip. 830. 1837 = ? Nephrolepis acuminata.
Sloani Raddi, Pl. Bras. **1**. 49 t. 71 bis. 1825 = ? P. gigantea.
Smithiana Pr.: Ettingsh. Farnkr. 95 t. 57 f. 2. 1865 = P. biaurita*.
socorrensis Karst. Fl. Col **1**. 118 t. 58. 1860 = P. grandifolia.
(L) **speciosa** Mett.: Kuhn, Linn. **36**. 91. 1869. HB. 480 — Ecuador-Peru.
spiculata Schkuhr. Kr. Gew. **1**. 92 t. 100. 1809 = Pellaea atropurpurea.
spinescens Pr. Rel. Haenk. **1**. 56. 1825 = P. biaurita*.
(L) **spinosa** (L.) Desv. Prod. 301. 1827 — Amer. trop. Polypodium L. sp. **2**. 1092. 1753; Pteris aculeata Sw. 1788 (f. Sw. Obs. bot. 403). HB. 171. Chr. 172. NPfl. 294; Litobrochia Pr. 1836 — Pteris reticulata Mett. 1869 (non Desv. 1811).
spinulifera Schum. Vid. Selsk. Afh. **4** 233 (seors. 459). 1827 = P. atrovirens.
spinulosa Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 47 t. 70. 1825 = P. leptophylla.
(L) **splendens** Klf. Enum. 186. 1824. HB. 168. Chr. 170. NPfl. 293 — Brasilia. Litobrochia Pr. 1836 — L. papyracea Fée 1850-52; Pteris Hk. 1858 — Litobrochia praealta Fée 1869.
Sprengelii Steud. Nom. **2**. 358. 1824 = Pteridium aquilinum.
Stelleri Gmel. Nov. Comm. Petr. **12**. 519 t. 12 f. 1. 1768 = Cryptogramma.
(P) **stenophylla** Wall.; Hk. et Grev. Ic. Fil. t. 130. 1829 — India bor. P. cretica β HB. 154. P. digitata Wall. 1828 (nomen).

stenophylla Link. Hort. Berol. **2**. 23. 1833 = P. denticulata.
stenophylla Pr. Tent. 145. 1836 (nomen) — ?
sterilis Pr. Del. Prag. **1**. 184. 1822 = ? P. spinosa.
stipulacea Hk. sp. **2**. 233. 1858 = Histiopteris incisa.
stipularis L. sp. **2**. 1074. 1753 = P. longifolia.
(P) **straminea** Mett.; Bak. Syn. 479. 1874 — Chile.
straminea Cord. Bull. Soc. sci. et arts Réunion **1890-91**; Fl. Réun. 52.
striata Desv. Prod. 300. 1827 — Réunion. [1895 = P. Cordemoyi.
stricta Poir. Enc. **5**. 713. 1804 = Schizoloma ensifolium.
stridens Ag. Rec. 45. 1839 = P. deflexa.
struthionis Desv. Prod. 294. 1827 = Polypodium.
subciliata Forsk. Fl. Aeg.-Arab. 185. 1775 — Arabia.
subindivisa Clarke, Tr. Linn. Soc. II. Bot. **1**. 467 t. 56 f. 1. 1880 = P. bi-
subpedata Wall. List n. 110. 1828 = P. tripartita. [aurita*.
subquinata Wall. List n. 104. 1828; Hope, Journ. Bomb. Nat. Hist. Soc. **13**. 452 t. 17. 1901 = P. biaurita*.
subverticillata Sw. Syn. 103. 1806 = Pellaea ternifolia.
succulenta Roxb. Calc. Journ. **4**. 508. 1844 = Ceratopteris thalictroides.
sulcata Meyen; J. Sm. Bot. Mag. **72** Comp. 24. 1846; Kze. Linn. **23**. 290.
sulfurea Cav. Descr. 269. 1802 = Notholaena. [1850 = P. biaurita*.
(P) **sumatrana** Bak. JoB. **1881**. 367 — Sumatra.
Swartziana Ag. Rec. **34**. 1839 = P. biaurita var.
tabularis Thbg. Prod. Fl. Cap. 171. 1800 = Blechnum.
taeniosa J. Sm. JoB. **3**. 405. 1841 = P. cretica.
tenera Klf. Enum. 191. 1824 = P. chilensis.
tenera Moritz, Bot. Zeit. **1854**. 856 (nomen) — Venezuela.
tenuifolia Lam. Fl. franç. **1**. 13. 1778 = Cryptogramma crispa.
tenuifolia Brack. Expl. Exp. **16**. 112. 1854 = P. longifolia.
tenuis A. Cunn. in Hk. Comp. Bot. Mag. **2**. 365. 1836 = P. tremula.
terminalis Wall. List n. 101. 1828; Ag. Rec. 20. 1839 = P. excelsa.
ternifolia Cav. Descr. 266. 1802 = Pellaea.
thalictroides Sw. Schrad. Journ. **1800**^{2}. 65. 1801 = Ceratopteris.
thalictroides Mühlenb. Trans. Amer. Phil. Soc. **3**. (157-184). 1793 — Amer. bor.
(P) **torricelliana** Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 45. 1905
trachyphylla Kze. Bot. Zeit. **1848**. 196 = P. biaurita*. [— N. Guinea.
(L) **transparens** Mett. Ann. sc. nat. V. **2**. 221. 1864. (HB. 480) — Columbia.
Trattinickiana Endl. Prod. Fl. Norf. 14. 1833 — Ins. Norfolk.
Treacheriana Bak. JoB. **1879**. 65 = P. cretica var. melanocaulon.
(P) **tremula** R.Br. Prod. Fl. N. Holl. 154. 1810. HB. 161. Chr. 169. NPfl. 293 — Austr. Tasmania. N. Zealand. Ins. Norfolk. ? Afr. austr.
— P. Kingiana Endl. 1833.
(L) **trialata** Sod. Cr. vasc. quit. 107. 1893 — Ecuador.
trichomanoides L. sp. **2**. 1074. 1753 = Notholaena.
trichomanoides Blanco, Fl. Filip. 830. 1837 — Ins. Philipp.
tricolor Linden, Gard. Chr. **1860**. 123 = P. biaurita var.
tricuspidata L. sp. **2**. 1073. 1753 = Eschatogramme furcata var.
trifoliata Fée, 8 mém. 114. 1857 = P. cretica.
trifoliata Christ, Bull. Boiss. **7**. 7. 1899 = P. nana.
(L) **tripartita** Sw. Schrad. Journ. **1800**^{2}. 67. 1801 — Africa, Asia, Austr., Polynesia trop. — Litobrochia Pr. 1836. Pteris marginata Bory 1804. HB. 172.

Chr. 172. NPfl. 294; Litobrochia Pr. 1836 — Pteris Pseudo-Lonchitis Bory; Willd. 1810; Campteria Pr. 1836; Litobrochia Fée 1850-52.
tripartita γ Milneana Hk. sp. **2**. 226 t. 238 B. 1858 = P. Milneana.
triphylla Bertero in »Mercator, Chil. n. 16. 748« = Pellaea ternifolia.
triphylla Ag. Rec. 16. 1839 = P. laurea.
triphylla Mart. et Gal. Mém. Ac. Brux. **15**. 51 t. 14 f. 1. 1842 = P. cretica.
triphylla Mett. Fil. Lips. 55. 1856 = Doryopteris.
triphylla Bak. Fl. bras. 1^2. 596. 1870 = Doryopteris subsimplex.
tripinnatifida Roxb. Calc. Journ. **4**. 509. 1844 — Moluccae.
triplicata Ag. Rec. 29. 1839 = P. heteroclita.
tristicula Raddi, Opusc. sci. Bol. **3**. 293. 1819; Pl. Bras. **1**. 46 t. 69. 1825 [= P. denticulata.
(L) **Tussaci** (Fée) Hk. sp. **2**. 229. 1858 — San Domingo. Litobrochia Fée, Gen. 137. 1850-52.
umbraculifera Mett.; Miq. Ann. Lugd. Bat. **4**. 97. 1868-69; Kuhn, Linn. **36**. 89. 1869 = P. longipinnula.
(P) **umbrosa** R.Br. Prod. Fl. N. Holl. 154. 1810. HB. 155. Chr. 166. NPfl. 292 [— Australia.
umbrosa Wall. List n. 109. 1828 = P. Wallichiana.
undulata Christ. Bull. Boiss. II. **2**. 550. 1902 = P. denticulata var.
uniseriata Poir. Enc. Suppl. **4**. 608. 1816 = P. tripartita.
(P) **usambarensis** Hier. in Engl. Pflwlt. Ostafr. C. 80. 1895 — Afr. orient. trop.
varia Sw. Schrad. Journ. **1800**^{2}. 69. 1801 — Ind. orient.
varians Raddi, Opusc. sci. Bol. **3**. 292. 1819 = Doryopteris.
varians Wall. List n. 86. 1828 = Cheilanthes Belangeri.
venulosa Bl. Enum. 209. 1828. } = P. pellucida.
venusta Kze. Bot. Zeit. **1848**. 195 } = P. pellucida.
Vespertilionis Lab. Nov. Holl. pl. spec. **2**. 96 t. 245. 1806 = Histiopteris [incisa.
(L) **vestita** Bak. Syn. 169. 1867 — Peru.
Victoriae ht. Bull; Gard. Chr. III. **9**. 676. 1891 = P. ensiformis var.
(L) **Vieillardii** Mett. Ann. sc. nat. IV. **15**. 66. 1861. HB. 167. NPfl. 293 — N. Caledonia. — Litobrochia Carr. 1873.
villosa Sw. Schrad. Journ. **1800**^{2}. 67. 1801 = ? Lonchitis hirsuta.
villosa Fée, Gen. 126. 1850-52 = Pteridium aquilinum.
(L) *villosa* Linden; Bak. Ann. Bot. **5**. 218. 1891 — Hort. (Assam?).
viridis Forsk. Fl. Aeg.-Arab. 186. 1775 = Pellaea.
viscosa Moore, Gard. Chr. **1858**. 878 = Paesia.
(P) **vitiensis** Bak. JoB. **1879**. 295 — Fiji.
vittarioides Thouars, Fl. Trist. d'Ac. 31 t. 1. 1804 = Vittaria.
vittata L. sp. **2**. 1074. 1753 = P. longifolia var.
vittata Schkuhr, Kr. Gew. **1**. t. 89. 1809 = P. grandifolia.
vulcanica Bertol. Misc. bot. **18**. 21. 1857; (Rend. Sess. Acc. Sci. Bol. **1856-57**. 54. 1857) = P. longifolia.
(P) **Walkeri** Bak. JoB. **1888**. 324 — Borneo.
Wallichiana Pr. Tent. 290. 1836 (nomen) = P. cretica.
(C) **Wallichiana** Ag. Rec. 69. 1839. HB. 165 — India bor. Java. Celebes. Ins. Philipp. Samoa. — Litobrochia Fée 1850-52; Campteria Moore 1861 C. pectinata Pr. 1836 (nomen). (an = P. tripartita forma?).
(L) **Warburgii** Christ in Warb. Monsunia **1**. 70. 1900 — Moluccae. N. Guinea.
Wavrini ht.: Rev. Hort. Belg. **1901**. 192 — Hort. (hybr.).
Wightiana Wall. List n. 2178. 1828 = Pteridium aquilinum.
Wimsettii ht. = P. cretica var. hort.

(L) **woodwardioides** Bory; Willd. sp. 5. 360. 1810. HB. 170 — Ins. Mascar. Litobrochia Pr. 1836 — Pteris alata Desv. 1827.

(L) **yunnanensis** Christ, Bull. Boiss. 6. 957. 1898. NPfl. 294 — Yunnan.

Zahlbruckneriana Endl. Prod. Fl. Norf. 13. 1833 = P. comans.

Zollingeri Mett.; Miq. Ann. Lugd. Bat. 4. 97. 1868-69 = P. longipes.

PTERONEVRON Fée, Gen. Fil. 320. 1850-52 = **Humata.**

parallelum Fée, Gen. 320. t. 25 B. f. 1. 1850-52 = H. Gaimardiana.

PTEROPSIS Desvaux, Prodrome 218. 1827.

angustifolia Desv. Prod. 218. 1827 = Vittaria.

angustifolia Pappe et Raws. Syn. Fil. Afr. austr. 43. 1857 = Vittaria isoetifolia.

angustifolia Fourn. Pl. mex. 1. 86. 1872 = Polypodium angustum.

Blumei Fée, 3 mém. 25. 1851-52 = Antrophyum callifolium.

elliptica Desv. Prod. 218. 1827 = Drymoglossum heterophyllum.

elongata Desv. Prod. 219. 1827 — Guiana. Grenada.

furcata Desv. Prod. 219. 1827 = Eschatogramme.

lanceolata Desv. Prod. 218. 1827 = Paltonium.

marginalis Pr. Tent. 226. 1836 = Antrophyum callifolium.

nummularia Desv. Prod. 218. 1827 = Drymoglossum heterophyllum.

piloselloides Desv. Prod. 218. 1827 = Drymoglossum heterophyllum.

scolopendrina Desv. Prod. 218. 1827 = Vittaria.

tricuspidata Desv. Prod. 219. 1827 = Eschatogramme furcata.

vittarioides Desv. Prod. 219. 1827 — Peru.

PTEROZONIUM Fée, Gen. Fil. 178. 1850-52 — (Polypodiaceae Gen. 57).

cyclophyllum (Bak.) Diels, Nat. Pfl. 1⁴. 257. 1899 — Guiana. Gymnogramme Bak. Tr. Linn. Soc. II. Bot. 2. 293 t. 53. 1887. Chr. 68.

reniforme (Mart.) Fée, Gen. 178 t. 16 A. 1850-52. NPfl. 257 — Guiana. Brasilia. Peru. — Gymnogramme Mart. Ic. Cr. Bras. 88 t. 26. 1834. HB 378. Chr. 68; Trichiogramme Kuhn 1882.

Ptiloneura Prantl, Ber. deutsch. bot. Ges. 1. 351. 1883 *(Ophioglossum §)* = Ophioglossum.

PTILOPHYLLUM van den Bosch. Versl. Akad. Wet. Amsterdam 11. 321. 1861 = **Trichomanes.**

accedens Prantl. Hym. 47. 1875 — bicorne Prantl, l. c. 48 — botryoides Prantl, l. c. 49 — crispum Prantl, l. c. 47 — cristatum Prantl, l. c. 47 — heterophyllum Prantl, l. c. 48 — holopterum Prantl. l. c. 47 — Hostmannianum Prantl, l. c. 49 — Lambertianum Prantl, l. c. 48 — pellucens Prantl, l. c. 48 — pennatum Prantl, l. c. 49 — pilosum Prantl, l. c. 48 — pinnatum Prantl, l. c. 49 — plumosum Prantl. l. c. 47 — Schomburgkianum Prantl, l. c. 49 — Vittaria Prantl, l. c. 49 = Trichomanes sp. homonym.

attenuatum Prantl, Hym. 48. 1875 = T. alatum var.

auratum Prantl, Hym. 48. 1875 = T. lucens.

Bancroftii Prantl, Hym. 48. 1875 = T. arbuscula.

elegans Prantl, Hym. 49. 1875 = T. diversifrons.

Kaulfussii Prantl, Hym. 48. 1875 = T. trigonum.

Martiusii Prantl, Hym. 48. 1875 = T. pilosum var.

ptilodes Prantl, Hym. 48. 1875 = T. alatum var.

sinuosum Prantl, Hym. 47. 1875 = T. polypodioides.

spicatum Prantl, Hym. 49. 1875 = T. osmundoides.

splendidum Prantl, Hym. 48. 1875 = T. lucens.

Spruceanum Prantl, Hym. 48. 1875 = T. heterophyllum var.

PTILOPTERIS Hance, Journ. Bot. **1884**. 138 = **Polystichum.**
flagellaris Mak. Bot. Mag. Tokyo **13**. 56. 1899.
Hancockii Hance JoB. **1884**. 139.
Maximowiczii Hance JoB. **1884**. 139.
PTYCHOMANES Hedwig, Fil. Gen. et Spec. fasc. II. 1800 = **Hymenophyllum.**
PTYCHOPHYLLUM Presl, Hymenoph. 28. 1843 = **Hymenophyllum.**
dichotomum Pr. Abh. böhm. Ges. V. **5**. 337. 1848.
magellanicum Kl. Linn. **18**. 533. 1844.
plicatum Pr. Hym. 29 t. 11 f. E. 1843 = H. dichotomum.
PYCNODORIA Presl, Epim. Bot. 100. 1849 = **Pteris.**
opaca Pr. Epim. 101. 1849.
PYCNOPTERIS Moore, Gard. Chr. **1855**. 468 = **Dryopteris.**
Sieboldii Moore, Gard. Chr. **1855**. 468 c. fig.
PYRROSIA Mirbel, Hist. nat. des Vég. **5**. 91. 1803; Hist. nat. pl. **4**. 70. 1803.
chinensis Mirb. Hist. nat. Vég. **5**. 92. 1803 = ? Cyclophorus Beddomeanus.
PYXIDARIA Gleditsch, Syst. plant. 291. 1764 = **Trichomanes.**
RAGATELUS Presl, Hymenoph. 16. 1843 = **Trichomanes.**
crinitus Pr. Hym. 16. 1843.
eriophorus Pr. Epim. 18 t. 9. 1849.
RAGIOPTERIS Presl, Tent. Pterid. 96. 1836 = **Onoclea.**
obtusiloba Pr. Tent. 96. 1836. } = O. sensibilis.
onocleoides Pr. Tent. 96 t. 3 f. 9, 10. 1836 } (conf. Milde, Bot. Zet. **1867**. 57).
RAMONDIA Mirbel, Bull. Soc. Philom. **2**. 179. 1801 = **Lygodium.**
flexuosa Mirb. l. c. t. 12 f. 3.
scandens Mirb. l. c.
REGNELLIDIUM Lindman, Arkiv för Botanik 3^6. 2. 1904 — (Marsileaceae
diphyllum Lindm. l. c. 14. f. 1-10. 1904 — Brasilia austr. (Gen. 2).
RHIPIDOPTERIS Schott, Gen. Fil. ad t. 14. 1834 = **Elaphoglossum.**
flabellata Fée, Acrost. 78. 1845.
foeniculacea Schott, Gen. ad t. 14. 1834.
Humboldtiana Schott, Gen. ad t. 14. 1834 = E. flabellatum.
peltata Schott, Gen. ad t. 14. 1834.
Rusbyi Christ, Farnkr. 46. 1897 = E. Moorei.
sphenophylla Fée. Acrost. 79. 1845 = E. flabellatum.
tripartita Schott. Gen. ad t. 14. 1834 = E. flabellatum.
RHIZOGLOSSUM Presl, Suppl. Tent. Pterid. 48. 1845 = **Ophioglossum.**
Bergianum Pr. l. c.
RHIZOSPERMA Meyen, Reise um die Erde **1**. 337. 1834; Nova Acta $\mathbf{18}^1$. 523. 1836 = **Azolla.**
RIEDLEA Mirbel, Hist. nat. des Vég. **5**. 71. 1803; Hist. nat. plant. **4**. 65. 1803
sensibilis Mirbel, Hist. nat. des Vég. **5**. 74. 1803. [= **Onoclea.**
RIPIDIUM Bernhardi, Schrad. Journ. $\mathbf{1800}^2$. 127. 1801 = **Schizaea.**
dichotomum Bernh. l. c. 127 t. 2 f. 3. 1801.
RUMHORA Desvaux, Prodrome 262. 1827 = Rumohra.
RUMOHRA Raddi, Opusc. scient. Bologn. **3**. 290. 1819 = **Polystichum.**
aspidioides Raddi l. c. t. 12 f. 1. 1819; Pl. Bras. **1**. 28 t. 43. 1825 = P. adiantiforme.
Runcinaria K. Müller, Bot. Zeit. **1854**. 545 (*Vittaria §*) = Vittaria.
SACCOLOMA Kaulfuss, Berl. Jahrb. f. d. Pharmacie **1820**. 51; Enum. Fil. 224. 1824 — (Polypodiaceae Gen. **29**).

adiantoides Pr. Tent. 126. 1836 = Lindsaya.

adiantoides Mett. Ann. sc. nat. IV. **15**. 80. 1861 = S. domingense.

alatum Mett. Ann. sc. nat. IV. **15**. 80. 1861 = S. inaequale.

amblylobum Prantl, Arb. bot. Gart. Breslau **1**. 21. 1892 (nomen) — ?

Boryanum Pr. Tent. 126. 1836 = Lindsaya repens var.

brasiliense Mett. Ann. sc. nat. IV. **15**. 80. 1861 = S. inaequale.

campylurum Mett. Ann. sc. nat. IV. **15**. **80**. 1861 = S. moluccanum var.

cuneifolium Pr. Tent. 126. 1836 = Lindsaya adiantoides.

domingense (Spr.) Prantl, Arb. bot. Gart. Breslau **1**. 21. 1892 — Amer. trop. Davallia Spr. Anleit. **3**. 149 t. 4 f. 33. 1804; Davallia adiantoides Sw. 1806; Cibotium Pr. 1836; Microlepia J. Sm. 1842; Patania Pr. 1849; Saccoloma Mett. 1861. NPfl. 212. Dicksonia Plumieri Hk. 1844. HB. 52; Microlepia Fée 1850-52; Sitolobium J. Sm. 1876; Saccoloma Christ 1897. 309 — Dicksonia Lindeni Hk. 1844; Patania Pr. 1849; Microlepia Fée 1850-52; Saccoloma Prantl 1892 — Dicksonia Eggersii Prantl 1897.

elegans Klf. Berl. Jahrb. Pharm. **1820**. 51; Enum. 224 t. 1. 1824. Chr. 309. NPfl. 211 — Amer. trop. — Neuropteris Desv. 1827; Microlepia Mett. 1856; Davallia Saccoloma Spr. 1827. HB. 97.

? hemipterum Pr. Tent. 126. 1836 = Lindsaya repens var.

Hookerianum Fée, Gen. 321. 1850-52 = Microlepia.

Imrayanum Hk. Gen. t. 58 B. 1839 (nomen); Kze. Farnkr. **1**. 86. 1842. NPfl. 211 — India occ. Guiana. — Davallia Hk. sp. **1**. 171 t. 49 A. 1846. HB. 90; Humata? Moore 1857; Acrophorus Moore 1861; Schizoloma Kuhn 1882.

inaequale (Kze.) Mett. Ann. sc. nat. IV. **15**. 80. 1861. Chr. 310. NPfl. 211 — Amer. trop. — Davallia Kze. Linn. **9**. 87. 1834. HB. 99. Microlepia Pr. 1836 — M. brasiliensis Pr. 1836; Davallia Hk. 1846; Saccoloma Mett. 1861 — Davallia alata Hew. 1838 (non Bl. 1828); Microlepia J. Sm. 1842; Saccoloma Mett. 1861 — Microlepia Galeottii Fée 1850-52; M. Boryana Fourn. 1872 — Davallia nigrescens Kze. 1850; Microlepia nigricans Pr. 1849. — (an Davallia distans Klf. Enum. 222. 1824; Microlepia J. Sm. 1842?).

Lindeni Prantl, Arb. bot. Gart. Breslau **1**. 20. 1892 = S. domingense var.

moluccanum (Bl.) Mett.; Kuhn, Verh. zool. bot. Ges. **19**. 581. 1869. Chr. 310. NPfl. 211 — Malesia. Polynesia. — Davallia Bl. Enum. 237. 1828. HB. 99; Microlepia Blumei Moore 1861; Davallia campylura Kze. 1850; Microlepia Moore 1861; Saccoloma Mett. 1861 — Microlepia tenuis Brack. 1854; Saccoloma Mett. 1861 — Microlepia papillosa Brack. 1854; Saccoloma Mett. 1861.

papillosum Mett. Ann. sc. nat. IV. **15**. 80. 1861 = S. moluccanum var.

pinnatum Pr. Tent. 126. 1836 = Tapeinidium.

Plumieri Christ, Farnkr. 309. 1897 = S. domingense.

Sandwicense A. Brongn. Voy. Venus Bot. t. 6. 1846 = Lindsaya repens.

Sloanei (Jenm.) C. Chr. Ind. 214. 1905; 612. 1906 — Jamaica. Davallia Jenm. JoB. **1886**. 37.

sorbifolium (Sm.) Christ, Farnkr. 309. 1897. NPfl. 211 — Malesia. N. Guinea. Dicksonia Sm. in Rees Cycl. 7. — (?). HB. 52; Cystodium J. Sm. 1842; Hk. Gen. t. 96. 1842. Dicksonia Papuana F. Muell. 1876.

tenue Mett. Ann. sc. nat. IV. **15**. 80. 1861 = S. moluccanum.

Wercklei Christ, Bull. Boiss. II. **4**. 1100. 1904 — Costa Rica.

SADLERIA Kaulfuss, Enum. Fil. 161. 1824 — (Polypodiaceae Gen. **52**).

cyatheoides Klf. Enum. 162. 1824. HB. 187. Chr. 184. NPfl. 251 — Ins. Hawaii. Sumatra. Perak. — Woodwardia Mett. 1856; Blechnum Fontanesianum Gaud. 1827.

pallida (Hk. et Arn. Bot. Beech. Voy. 75, 107. 1832); Hill. Fl. Haw. 581. 1888. NPfl. 251 — Ins. Hawaii. — Blechnum Brack. 1854. (S. pallida Hk. et Arn. est f. Bak. HB. 187 = S. cyatheoides).

polystichoides (Brack.) Heller, Minn. bot. St. **1**. 788. 1897 — Ins. Hawaii. Blechnum Brack. Expl. Exp. **16**. 134. 1854; B. squarrosum Gaud. Voy. Bonite, Bot. t. 2. 1846 (sine descr.); Sadleria Moore 1857. HB. 187. Chr. 184. NPfl. 251 — Polypodium unisorum Bak. 1867. HB. 307. Gymnogramme sadlerioides Und. 1897.

Souleytiana (Gaud.) Moore, Ind. XXVI. 1857; Hill. Fl. Haw. 581. 1888. Chr. 184. NPfl. 251 — Ins. Hawaii. — Blechnum Gaud. Voy. Bonite, Bot. t. 2, 134. 1846 (sine descr.).

squarrosa Moore, Ind. XXVI. 1857 = S. polystichoides.

SAGENIA Presl, Tent. Pter. 86. 1836 = **Aspidium.**

alata Bedd. Ferns br. Ind. t. 169. 1866 = A. vastum.
ampla Moore, Ind. 228. 1861 = Cardiochlaena a.
angulata Moore, Ind. 96. 1858 = A. Plumieri.
apiifolia J. Sm. JoB. **4**. 184. 1841 = A. cicutarium*.
calcarea J. Sm. JoB. **3**. 410. 1841.
cicutaria Moore, Ind. LXXXVI. 1857.
coadunata J. Sm. JoB. **4**. 184. 1841 = A. cicutarium*.
confluens Moore, Ind. 228. 1861 = Cardiochlaena c.
decurrens Houlst.; Moore, Ind. LXXXVI. 1857.
difformis Christ, Ann. Jard. Buit. II. **5**. 109. 1905 = A. irregulare.
dilacerata Moore, Ind. LXXXVI. 1857 = A. cicutarium.
gemmifera Fée, Gen. 313. 1850-52 = A. cicutarium.
gigantea Moore, Ind. 92. 1858.
gigantea var. minor Bedd. Ferns S. Ind. t. 243. 1863-65 = A. devexum.
grandis Moore, Ind. LXXXVI. 1857 = A. grandifolium.
Griffithii Bedd. Ferns br. Ind. t. 337. 1870 = A. multicaudatum.
heterocarpa Bedd. Ferns br. Ind. t. 47. 1865 = A. heterosorum.
Hippocrepis Pr. Tent. 87 t. 2. f. 24, 25. 1836.
Hippocrepis Brack. Expl. Exp. **16**. 181. 1854 = A. latifolium.
intermedia J. Sm. JoB. **3**. 410. 1841 = A. cicutarium.
Labrusca Christ, Ann. Jard. Buit. II. **5**. 108. 1905.
latifolia Pr. Tent. 87 t. 2 f. 23. 1836 = A. cicutarium*.
latifolia Carr. in Seem. Fl. Vit. 364. 1873.
Lawrenceana Moore, Gard. Chr. n. s. **15**. 9. 1881.
lobata Pr. Tent. 87. 1836 = Polypodium l.
Lobbii Christ, Ann. Jard. Buit. II. **5**. 100. 1905.
macrodonta Fée, Gen. 313 t. 24 A f. 1. 1850-52 = A. cicutarium.
macrophylla Moore, Ind. LXXXVI. 1857 = A. martinicense.
mamillosa Moore, L'Ill. hort. IV. **6**. t. 598; Gard. Chr. n. s. **26**. 38. 1886.
melanocaulis Moore, Ind. 90. 1858.
melanocaulis Bail. Hdb. Ferns Queensl. 47. 1874 = A. Mülleri.
membranacea Bedd. Ferns br Ind. Correct. II. 1870 = A. devexum.
membranifolia Christ, Bull. Acad. Géogr. Bot. Mans **1902**. 257.

? menisciicarpa Moore, Ind. 96. 1858 = Dryopteris lineata.
Menyanthidis Moore, Ind. 96. 1858.
Mexicana Fée, Gen. 313. 1850-52 = A. cicutarium.
microsora Moore, Ind. LXXXVI. 1857 = A. melanocaulon.
pachyphylla Moore, Ind. LXXXVI. 1857.
palmata Christ, Ann. Jard. Buit. II. 5. 108. 1905.
Pica Moore, Ind. 90. 1858.
platyphylla J. Sm. JoB. **3**. 410. 1841 = A. cicutarium.
polymorpha Christ, Ann. Jard. Buit. II. **5**. 108. 1905.
pteropoda Christ, Ann. Jard. Buit. II. **5**. 108. 1905.
pteropus Moore, Ind. 89. 1858 = A. decurrens.
puberula Moore, Ind. LXXXVI. 1857.
repanda Pr. Epim. 59. 1849 = A. cicutarium.
repanda Moore, Ind. 102. 1858.
rufescens Pr. Tent. 87. 1836 = A. cicutarium.
semibipinnata Moore, Ind. 103. 1858.
siifolia Moore, Ind. 93. 1858 = A. polymorphum.
sinuata Moore, Ind. 104. 1858 } = A. sinuatum.
sinuosa Moore, Ind. LXXXVI. 1857 (err.) } = A. sinuatum.
sorbifolia Pr. Tent. 87. 1836 = ? Polybotrya polyphylla.
subdigitata Christ, Ann. Jard. Buit. II. **5**. 100. 1905 = A. psilopodum.
subtriphylla Bedd. Ferns S. Ind. t. 242. 1863-65.
Thwaitesii Bedd. Ferns S. Ind. t. 244. 1863-65.
varia Pr. Tent. 87. 1836; Epim. 60. 1849 — Luzon.
Nephrodium Pr. Rel. Haenk. **1**. 36. 1825.
varia J. Sm. JoB. **4**. 184. 1841; Moore, Ind. 107. 1858 = Aspidium (Willd.).
variolata Moore, Ind. 107. 1858 = A. variolosum.
variolosa Bedd. Ferns br. Ind. t. 170. 1866 = A. cicutarium.
vasta Moore, Ind. LXXXVI. 1857.
Vitis Christ, Ann. Jard. Buit. II. **15**. 108 t. 3 f. 2. 1905.

SALPICHLAENA J. Smith in Hook. Gen. Fil. t. 93. 1842 = **Blechnum.**
Salpiglaena Kl. 1844; Salpinchlaena Pr. 1849; Trev. 1869.
adnata Trev. Atti Ist. Veneto III. **14**. 576. 1869 = B. orientale.
brasiliensis Trev. l. c. 574. 1869.
cartilaginea Trev. l. c. 575. 1869.
Cumingiana Fée, Gen. 78. 1850-52 = B. orientale.
denticulata Trev. l. c. 575. 1869 = B. serrulatum.
elongata Trev. l. c. 575. 1869 = B. orientale.
Finlaysoniana Fée, Gen. 78. 1850-52.
imbricata Trev. l. c. 575. 1869 = B. orientale.
latifolia Trev. l. c. 575. 1869 = B. orientale.
longifolia Trev. l. c. 575. 1869 = B. orientale.
malaccensis Trev. l. c. 576. 1869 = B. serrulatum.
nitida Trev. l. c. 574. 1869 = B. elongatum.
orientalis Fée, Gen. 78. 1859-52.
Patersonii Fée, Gen. 78. 1850-52.
pectinata Trev. l. c. 575. 1869 = B. orientale.
pyrophylla Trev. l. c. 575. 1869 = B. orientale.
scandens Pr. Epim. 122. 1849 = B. volubile.
serrulata Trev. l. c. 576. 1869.

stenophylla Trev. l. c. 575. 1869 = B. orientale.
striata Trev. l. c. 576. 1869 = B. serrulatum.
volubilis J. Sm. in Hk. Gen. t. 93. 1841.

SALPIGLAENA Klotzsch, Linnaea **20**. 348. 1847 = Salpichlaena.

SALPINCHLAENA Presl, Epim. Bot. 122. 1849; Trevisan, Atti Ist. Veneto III. **14**. 574. 1869 = Salpichlaena.

SALVINIA (Micheli, Nov. Gen. 107 t. 58); Adanson, Fam. d. plant. **2**. 15. 1763 — (Salviniaceae Gen. **2**).

adnata Desv. Prod. 177. 1827. Bak. Hdb. 137 — Africa occ.
affinis Desv. Prod. 177. 1827 = S. auriculata.
auriculata Aublet, Hist. Pl. Guian. **2**. 969 t. 367. 1775. Bak. Hdb. 136. NPfl. 402 — Amer. trop. — S. rotundifolia Willd. 1810. S. hispida HBK. 1815. S. biloba Raddi 1825 — S. affinis Desv. 1827.
Azolla Raddi, Pl. Bras. **1**. 2 t. 1 f. 3. 1825 = Azolla caroliniana.
biloba Raddi, Pl. Bras. **1**. 1 t. 1 f. 4. 1825 = S. auriculata.
cucullata Roxb.; Wall. List n. 399. 1828; Bory, Bél. Voy. Bot. **2**. 6. 1833; Roxb. Calc. Journ. **4**. 470. 1844. Bak. Hdb. 186. NPfl. 402 — India. W. Austr.
europaea Desv. Prod. 177. 1827 = S. natans.
hastata Desv. Prod. 177. 1827. Bak. Journ. Linn. Soc. **24**. 98. 1887. Hdb. 135. NPfl. 402 — Madagascar.
Hildebrandtii Bak. JoB. **1886**. 98. Hdb. 135. NPfl. 402 — Madagascar.
hispida HBK. Nov. Gen. et Sp. **1**. 44. 1815 = S. auriculata.
imbricata Roxb. Calc. Journ. **4**. 470. 1844 = Azolla pinnata.
laevigata H.B.Willd. sp. **5**. 537. 1810 = Trianaea bogotensis (Hydrocharidaceae).
minima Bak. JoB. **1886**. 98. Hdb. 136. NPfl. 402 — Brasilia austr.
mollis Mett.; Kuhn, Fil. Afr. 201. 1868. Bak. Hdb. 135. NPfl. 402 — Madagascar.
natans (L.) All. Fl. Pedem. **2**. 289. 1785. Bak. Hdb. 135. NPfl. 402 — Eur. centr.-China bor.-India. — Marsilea L. sp. **2**. 1099. 1753; Salvinia europaea Desv. 1827; S. vulgaris Rupr. 1845 — S. Sprengelii Corda 1829 — S. verticillata Roxb. 1844.
nigropunctata A.Br. in Kuhn, Fil. Afr. 201. 1868. Bak. Hdb. 135. NPfl. 402 — Africa occ.
nymphellula Desv. Prod. 177. 1827. Bak. Hdb. 136 — Africa occ.
oblongifolia Mart. Ic. Cr. Bras. 128 t. 75 f. 2. t. 76. 1834. Bak. Hdb. 135. NPfl. 402 — Brasilia.
Olfersiana Kl. (Bak. Hdb. 136) = S. auriculata var.
radula Bak. JoB. **1886**. 98. Hdb. 136. NPfl. 402 — Guiana. Brasilia.
rotundifolia Willd. sp. **5**. 537. 1810 = S. auriculata.
Sprengelii Corda. Monogr. Rhizosp. 10 t. 2 f. 12-23. 1829 = S. natans.
Sprucei Kuhn, Fl. Bras. 1^2. 655 t. 81 f. 11-13. 1884. Bak. Hdb. 136. NPfl. 402 [— Amazonas.
verticillata Roxb. Calc. Journ. **4**. 469. 1844 = S. natans.
vulgaris Rupr. Distr. Cr. vasc. Ross. 27. 1845 = S. natans.

SCEPTRIDIUM Lyon, Bot Gazette **40**. 457. 1905 = **Botrychium.**

australe, biforme, biternatum, californicum, Coulteri, daucifolium, decompositum, dissectum, japonicum, Jenmani Lyon, Bot. Gaz. **40**. 457. 1905 — matricariae, obliquum, pusillum, robustum, Schaffneri, silaifolium, subbifoliatum, tenuifolium, ternatum, Underwoodianum Lyon l. c. 458 = B. sp. homonym.

SCHAFFNERIA Fée, 7 mémoire 56. 1857 = **Phyllitis.**

nigripes Fée, 7 mém. 56 t. 17 f. 1. 1857.

SCHELLOLEPIS J. Smith, Ferns brit. and foreign 82. 1866 = **Polypodium.**
Goniophlebium § Schellolepis J. Sm. in Hk. Gen. ad t. 51. 1840.
amoena J. Sm. Hist. Fil. 93. 1875.
arguta J. Sm. Hist. Fil. 93. 1875.
cuspidata J. Sm. Ferns br. and for. 82. 1866 = P. persicifolium.
lachnopus J. Sm. Hist. Fil. 93. 1875.
pallens J. Sm. Hist. Fil. 93. 1875 = P. subauriculatum var.
subauriculata J. Sm. Ferns br. and for. 82. 1866.
verrucosa J. Sm. Ferns br. and for. 83. 1866.

SCHIZAEA Smith, Mém. Acad. Turin **5**. 419. 1793 — (Schizaeaceae Gen. **1**).
attenuata Beyrich; Sturm, Fl. bras. **1**². 184. 1859 = S. elegans.
australis Gaud. Ann. sc. nat. **5**. 98. 1825 = S. fistulosa var.
Balansae Fourn. Ann. sc. nat. V. **18**. 353. 1873 — N. Caledonia.
bifida Willd. Schrift. Akad. Erf. **1802**. 30 t. 3 f. 3. HB. 429. Prantl. Schiz. 136. Chr. 345. NPfl. 363 — Australia. N. Zealand.
bifida auctt. quoad pl. amer. = S. incurvata.
chilensis Phil. Linn. **30**. 207. 1859-60 = S. fistulosa.
cristata Willd. sp. **5**. 88. 1810 = S. dichotoma.
dichotoma (L.) Sm. Mém. Ac. Turin **5**. 422 t. 9 f. 9. 1793. HB. 430 (pt). Prantl, Schiz. 138. Chr. 345. NPfl. 363 — Asia, Austr., Polynesia trop. Madagascar. Ins. Mascarenae. — Acrostichum L. sp. **2**. 1068. 1753; Osmunda Spr. 1799; Ripidium Bernh. 1801 — Schizaea Forsteri Spr. 1804 — S. cristata Willd. 1810.
dichotoma Kze. Linn. **9**. 19. 1834 et auctt. quoad pl. americ. = S. Poeppigiana.
digitata (L.) Sw. Syn. 150, 380 t. 4 f. 1. 1806. HB. 430. Prantl, Schiz. 133. Chr. 344. NPfl. 363 — Asia, Polynesia trop. ? Madagascar. — Acrostichum L. sp. **2**. 1068. 1753; Belvisia Mirb. 1803; Actinostachys Wall. 1828.
digitata var. orbicularis Bak. Journ. Bot. **1881**. 208 = S. orbicularis.
elegans (Vahl) Sw. Schrad. Journ. **1800**². 103. 1801. HB. 430. Prantl, Schiz. 140. Chr. 346. NPfl. 363 — Amer. trop. — Acrostichum Vahl, Symb. **2**. 104 t. 50. 1791; Lophidium Pr. 1845. L. latifolium Rich. 1792 — Schizaea Flabellum Mart. 1834; Lophidium Pr. 1845.
filifolia La Pylaie, Ann. Soc. Linn. Paris **4**. 450. 1825 = S. pusilla.
fistulosa Labill. Nov. Holl. Pl. Spec. **2**. 103 t. 250 f. 3. 1806. HB. 429. Prantl, Schiz. 135 (pt.). Chr. 345. NPfl. 363 — Tasmania. N. Zealand. Amer. antarctica. Madagascar. Borneo. — Acrostichum Poir. 1810. Schizaea propinqua A. Cunn. 1836 — S. chilensis Phil. 1859-60 — S. australis Gaud. 1825. HB. 428. (Ins. Falklandicae).
Flabellum Mart. Ic. Cr. Bras. 115 t. 55 f. 2. 1834 = S. elegans var.
fluminensis Miers; Sturm, Fl. bras. **1**². 184 t. 15 f. 2. 1859. HB. 429. Prantl, Schiz. 140. Chr. 346. NPfl. 363 — Brasilia. Grenada.
Forsteri Spr. Anleit. **3**. 157. 1804 = S. dichotoma.
Germani (Fée) Prantl, Schiz. 132. 1881 — Guadeloupe. Florida.
Actinostachys Fée, 11 mém. 123 t. 29 f. 3. 1866.
incurvata Schkuhr. Kr. Gew. **1**. 138 t. 137. 1809. HB. 429. Prantl, Schiz. 137. NPfl. 363 — Brasilia. Guiana. — S. bifida auctt. quoad pl. americ.
intermedia Mett. Ann. sc. nat. IV. **15**. 86. 1861. Prantl, Schiz. 132 — N. Caledonia.
laevigata Mett. Ann. sc. nat. IV. **15**. 85. 1861. Prantl. Schiz. 133 — N. Caledonia.
malaccana Bak. Syn. 428. 1868 — Birma. Malesia. Ins. Philipp.
S. fistulosa δ Prantl, Schiz. 136.

occidentalis Gris. Cat. Pl. Cub. 273. 1866 = S. Poeppigiana.

orbicularis (Bak.) C. Chr. Ind. 617. 1906 — Columbia.
S. digitata var. orbicularis Bak. JoB. **1881**. 208.

pacificans Mart. Ic. Cr. Bras. 116 t. 56 f. 1. 1834. Prantl, Schiz. 141 — Brasilia.
Lophidium Pr. 1845 — Schizaea spectabilis Mart. 1834; Lophidium Pr. 1845.

palmata Hombr. et Jacq. Voy. au Pôle sud t 4. 1852 = S. fistulosa var. australis.

pectinata (L.) Sw. Schrad. Journ. **1800²**. 102. 1801. HB. 429. Prantl, Schiz. 134. Chr. 345. NPfl. 363 — Africa austr. St. Helena.
Acrostichum L. sp. **2**. 1068. 1753.

penicillata H.B.Willd. sp. **5**. 86. 1810. Prantl, Schiz. 132. NPfl. 363 — Amer. austr. trop. Trinidad. — S. subtrijuga Mart. 1834; Actinostachys Pr. 1845.
S. Pennula β HB. 430.

penicillata Bory, Dup. Voy. Bot. **1**. 250 t. 27. 1828 = S. pennula.

pennula Sw. Syn. 150, 379. 1806. HB. 430. (excl. β). Prantl, Schiz. 131. Chr. 344. NPfl. 363 — Amer. trop. — Acrostichum Poir. 1810; Actinostachys Hk. 1842. Schizaea trilateralis Schkuhr 1809; Actinostachys J. Sm. 1842.

plana Fourn. Ann. sc. nat. V. **18**. 353. 1873 — N. Caledonia.

Poeppigiana Sturm, Fl. bras. **1²**. 181. 1859. Prantl, Schiz. 139. NPfl. 363 — Amer. trop. — S. dichotoma auctt. quoad pl. amer. HB. 430. — S. occidentalis Gris. 1866.

propinqua A. Cunn. in Hk. Comp. Bot. Mag. **2**. 362. 1836 = S. fistulosa.

pusilla Pursh, Fl. Am. sept. **2**. 657. 1814. HB. 428. Prantl, Schiz. 134. Chr. 344. NPfl. 363 — N. Jersey. Nova Scotia. N. Foundland.

robusta Bak. Syn. 429. 1868 — Ins. Hawaii. Ins. Societatis.
S. fistulosa γ Prantl, Schiz. 136.

rupestris R.Br. Prod. Fl. N. Holl. 162. 1810. HB. 429. Prantl, Schiz. 135. NPfl. 363 — Australia.

serrata Fée, 2 mém. 112. 1845 = Acrostichum (Lam.).

spectabilis Mart. Ic. Cr. Bras. 117. 1834 = S. pacificans.

?spicata Sm. Mém. Ac. Turin **5**. 43. 1793 = Hymenolepis.

Sprucei Hk.; Bak. in Hk. Ic. pl. t. 1016. 1867. HB. 430. Prantl, Schiz. 140. NPfl. 363 — Amazonas.

subtrijuga Mart. Ic. Cr. Bras. 117. 1834 = S. penicillata.

tenella Klf. Enum. 50 t. 1 f 7. 1824. HB. 428. Prantl, Schiz. 135. Chr. 345. NPfl. 363 — Africa austr.

tenuis Fourn. Ann. sc. nat. V. **18**. 353. 1873 — N. Caledonia.

tortuosa Mühl. Cat. 102. 1813 (?1818 nomen?) = S. pusilla.

trilateralis Schkuhr. Kr. Gew. **1**. 137 t. 136. 1809 = S. pennula.

valdiviana Phil. (Hk. & Bak. Syn. 429. 1868 syn.) = S. fistulosa.

SCHIZOGRAMMA Link. Fil. spec. 136. 1841 = ?**Hemionitis**.

SCHIZOCAENA J. Smith in Hook. Gen. Fil. t. 2. 1838 = **Cyathea**.

alternans J. Sm. Hist. Fil. 244. 1875.

Brunonis J. Sm. in Hk. Gen. Fil. t. 2. 1838 = C. moluccana.

Gaudichaudii Fée, Gen. 354. 1850-52 = C. moluccana.

laevigata J. Sm. Lond. JoB. **1**. 661. 1842 = C. borbonica var.

sinuata J. Sm. Lond. JoB. **1**. 661. 1842.

SCHIZOLEPTON Fée, 3 mémoire 10, 27. 1851-52.

cordatum Fée, 3 mém. 27. 1851-52 = Schizoloma.

rigidum Moore. Ind. XXX. 1857 = Drymoglossum.

SCHIZOLOMA Gaudichaud, Ann. sc. nat. **3**. 507. 1824 — (Polypodiaceae Gen. **38**).
Agatii Brack. Expl. Exp. **16**. 216 t. 30 f. 1. 1854 = S. ensifolium.
alutaceum Kuhn, Chaetopt. 346. 1882 = Lindsaya.
Billardieri Gaud. Ann. sc. nat. **3**. 508. 1824; Freyc. Voy. Bot. 380. t. 17 = S. ensifolium var.
Boivini (Mett.) Kuhn, Chaetopt. 346. 1882 — Ins. Comorae.
Lindsaya Mett.; Kuhn, Fil. Afr. 67, 206. 1868.
clavatum Kuhn, Chaetopt. 346. 1882 = Odontosoria.
cordatum Gaud. Ann. sc. nat. **3**. 507. 1824; Freyc. Voy. Bot. 379 t. 16. NPfl. 219 — Malesia. — Lindsaea Gaud.; Desv. 1827. HB. 111; Schizolepton Fée 1851-52.
cuneatum (Willd.) Kuhn, Chaetopt. 346. 1882 — Ins. Mascarenae.
Lindsaea Willd. sp. **5**. 423. 1810. (HB. 112).
davallioides Moore, Ind. XXXV. 1857 = Lindsaya.
divergens (Roxb.) Kuhn, Chaetopt. 346. 1882. NPfl. 219 — Malacca. Borneo.
Vittaria Roxb.; Wall. 1828 (nomen); Lindsaea Hk. et Grev. Ic. Fil. t. 226. 1831. HB. 109. Chr. 295; Isoloma J. Sm. 1841.
elongatum Kuhn, Chaetopt. 346. 1882 = Lindsaya deltoidea.
ensifolium (Sw.) J. Sm. JoB. **3**. 414. 1841. NPfl. 219 — Asia trop. Austr. trop. Polynesia. Afr. occ. et austr. c. ins. — Lindsaea Sw. Schrad. Journn. **1800**[2]. 77. 1801. HB. 112. Chr. 297; Adiantum Poir. 1810 — Lindsaea pentaphylla Hk. 1844; Schizoloma Fée 1850-52 — Lindsaea Griffithiana Hk. 1844; Schizoloma Fée 1850-52 — Lindsaea lanceolata Lab. 1806. Adiantum Poir. 1810; Schizoloma? Pr. 1836; S. Billardieri Gaud. 1824; Lindsaya Carr. 1873. Schizoloma Agatii Brack. 1854 — S. Javae Fée 1850-52.
erectum Moore, Ind. XXXV. 1857 = Diellia.
falcatum Moore, Ind. XXXV. 1857 = Diellia.
ferrugineum Kuhn, Chaetopt. 346. 1882 = Odontosoria Melleri.
ferulaceum Kuhn, Chaetopt. 346. 1882 = Davallia.
Fraseri (Hk.) Fée, Gen. 108. 1850-52, NPfl. 219 — Queensland.
Lindsaea Hk. sp. **1**. 221 t. 70 B. 1846. HB. 112.
Griffithianum Fée, Gen. 108. 1850-52 = S. ensifolium.
Guerinianum Gaud. Ann. sc. nat. **3**. 508. 1824; Freyc. Voy. Bot. 380. t. 18. NPfl. 219 — Malesia. — Lindsaea Gaud.; Desv. 1827. HB. 111; Isoloma Fée 1850-52; Guerinia articulata J. Sm. 1875 — Lindsaea ovata J. Sm.; Hk. 1846. HB. 105; Isoloma Pr. 1849.
heterophyllum (Dry.) J. Sm. JoB. **3**. 414. 1841. NPfl. 219 — China merid. India austr. Asia trop. — Lindsaea Dry. Tr. Linn. Soc. **3**. 41 t. 8 f. 1. 1797. HB. 112. Chr. 298; Adiantum Poir. 1810. Lindsaea variabilis Hk. [et Arn. 1836-40.
Imrayanum Kuhn. Chaetopt. 346. 1882 = Saccoloma.
induratum (Bak.) C. Chr. Ind. 618. 1906 — Borneo. — Lindsaea Bak. JoB. [**1888**. 324.
Javae Fée, Gen. 109 t. 29 f. 1. 1850-52 = S ensifolium.
?lanceolatum Pr. Tent. 132. 1836 = S. ensifolium var.
lobatum Bedd. Ferns br. Ind. Suppl. 6. 1876 = Lindsaya decomposita.
macrophyllum (Klf.) Pr. Tent. 132. 1836. NPfl. 219 — Guiana. Brasilia.
Lindsaya Klf. Enum. 218. 1824. HB. 112. Chr. 298.
madagascariense Kuhn, Chaetopt. 346. 1882 = Lindsaya.
malabaricum Bedd. Ferns br. Ind. t. 268. 1868 = Lindsaya decomposita var.
medium Kuhn, Chaetopt. 346. 1882 = Lindsaya.
microphyllum Kuhn. Chaetopt. 346. 1882 = Lindsaya.
nervosum Kuhn, Chaetopt. 346. 1882 = Lindsaya.

nitens Bedd. Ferns brit. Ind. Correct. II. 1870 = Lindsaya decomposita.
orbiculatum Kuhn, Chaetopt. 346. 1882 = Lindsaya.
pellaeiforme (Christ) C. Chr. Ind. 619. 1906 — Celebes. — Lindsaya Christ, Verh. Nat. Ges. Basel **11**. 430. 1896; Ann. Jard. Buit. **15**. 102 t. 14 f. 14. [1897.
pentaphyllum Fée, Gen. 108. 1850-52 = S. ensifolium.
Pickeringii Moore, Ind. XXXV. 1857 = Lindsaya.
prolongatum Kuhn, Chaetopt. 346. 1882 = Lindsaya.
propinquum Moore, Ind. XXXV. 1857 = Lindsaya decomposita var.
pumilum Moore, Ind. 320. 1861 = Diellia.
recurvatum Moore, Ind. XXXV. 1857 = Lindsaya decomposita var.
reniforme (Dry.) Diels, Nat. Pfl. **1**[4]. 218. 1899 — Guiana. Amazonas. Lindsaea Dry. Tr. Linn. Soc. **3**. 40 t. 7 f. 1. 1797. HB. 109. Chr. 294.
retusum Kuhn, Chaetopt. 346. 1882 = Odontosoria.
sagittatum (Aubl.) Diels, Nat. Pfl. **1**[4]. 218. 1899 — Guiana. Guadeloupe. Adiantum Aubl. Hist. pl. Guian. **2**. 964. t. 366. 1875; Lindsaea Dry. 1797. HB. 109. Chr. 295.
scoparium Kuhn, Chaetopt. 346 t. 1 f. 1. 1882 = Odontosoria.
trichomanoides Kuhn, Chaetopt. 346. 1882 = Lindsaya cuneata.
Vieillardii (Mett.) Kuhn, Chaetopt. 346. 1882 — N. Caledonia. Lindsaea Mett. Ann. sc. nat. IV. **15**. 65. 1861. HB. 471.
Walkerae (Hk.) Kuhn, Chaetopt. 346. 1882. NPfl. 218 — Ceylon. Bangka. Lindsaea Hk. sp. **1**. 209 t. 69 A. 1846. HB. 109; Isoloma Pr. 1849.

SCHIZOPTERIS Hillebrand msc. (Hk. & Bak. Syn. 475. 1874) = Schizostege.

SCHIZOSTEGE Hillebrand, Fl. Hawaii. 631. 1888 = **Pteris**.
Lydgatii Hill. Fl. Haw. 632. 1888.

SCOLIOSORUS Moore, Index Fil. XXIX. 1857 = **Antrophyum**.
ensiformis Moore, Ind. XXIX. 1857.

SCOLOPENDRIUM Adanson, Fam. des plant. **2**. 20. 1763; Smith, Mém. Acad. Turin **5**. 410. 1793 = **Phyllitis**.
alternifolium Roth, Röm. Mag. **2**[1]. 105. 1799; Tent. Fl. Germ. **3**. 53. 1800 = Asplenium germanicum.
ambiguum Raddi, Opusc. sci. Bol. **3**. 292. 1819; Pl. Bras. **1**. 40 t. 57 f. 1. [1825 = P. brasiliensis.
Balansae Bak. in Hk. Ic. pl. t. 1653. 1886.
brasiliense Fisch.; Kze. Linn. **23**. 291. 1850.
breve Bertol. Misc. bot. **18** 20 t. 5. 1857; (Rend. Sess. Acc. Sci. Bol. **1856-57**. 54. 1857) = P. hemionitis.
Ceterach Symons, Syn. pl. 193. 1798 = Ceterach officinarum.
cordatum Fée, Gen. 209. 1850-52 = P. hemionitis.
Delavayi Franch. Bull. Soc. Fr. **32**. 29. 1885.
Douglasii Fisch.; Hk. sp. **4**. 3. 1862 = P. plantaginea.
dubium Don, Prod. Fl. Nepal. 9. 1825 = Diplazium lanceum.
Durvillaei Bory, Dup. Voy. Bot. **1**. 273 t. 37 f. 1. 1828 = Stenochlaena sorbifolia [var 16.
hemionitis Lag., Garcia et Clem. Anal. Hist. nat. **5**. 150 t. 41. f. 2. 1802.
hybridum Milde, Verh. zool. bot. Ges. **14**. 235 t. 18. 1864.
Ikenoi Mak. Bot. Mag. Tokyo **13**. 130. 1899.
integrifolium Christ. Farnkr. 212. 1897 = Diplora i.
Krebsii Kze. Linn. **18**. 118 1844 = Blechnum punctulatum var.
lanceolatum Roxb. Calc. Journ. **4**. 501. 1844 — Chittagong.
Lindeni Hk. Ic. pl. t. 488. 1842 = P. scolopendrium var.
Lingua Cav. Descr. 253. 1802 = P. scolopendrium.

lobatum Rouy, Ill. pl. Eur. rar. t. 100. 1896 — Gallia.
(S. vulgare × Asplenium marinum).
longifolium Pr. Rel. Haenk. **1**. 48 t. 9 f. 1. 1825 = Triphlebia l.
mambare Bailey, Queensl. Agric. Journ. $\mathbf{3}^2$. 9 (seors.) 1898 — N. Guinea.
(an Triphlebia sp.?)
minus Fée, Gen. 209 t. 17 D f. 3. 1850-52. = P. hemionitis.
nigripes Hk. sp. **4**. 3. 1862.
oblongatum Schrad. Gött. gel. Anz. **1824**. 870 = P. brasiliensis.
officinale Lam. et DC. Fl. franç. **2**. 552. 1805 } = P. scolopendrium.
officinarum Sw. Schrad. Journ. $\mathbf{1800}^2$. 61. 1801 } = P. scolopendrium.
Phyllitis Roth, Röm. Mag. $\mathbf{2}^1$. 105. 1799; Tent. Fl. Germ. **3**. 47. 1800 } = P. scolopendrium.
pinnatifidum Diels, Nat. Pfl. $\mathbf{1}^4$. 231. 1799 = Asplenium p.
pinnatum J. Sm. JoB. **3**. 406. 1841 = Triphlebia p.
plantagineum Schrad. Gött. gel. Anz. **1824**. 870.
repandum Pr. Del. Prag. **1**. 180. 1822 = P. brasiliensis.
rhizophyllum Endl. Gen. pl. Suppl. I. 1348. 1840 = Camptosorus r.
Ruta muraria Roth, Röm. Mag. $\mathbf{2}^1$. 105. 1799; Tent. Fl. Germ. **3**. 52. 1900 = Asplenium.
sagittatum DC. Fl. franç. **5**. 238. 1815 = P. hemionitis.
Scolopendrium Karst. Deutschl. Flora 278. 1880-83.
septentrionale Roth, Röm. Mag. $\mathbf{2}^1$. 105. 1799; Tent. Fl. Germ. **3**. 49. 1900 = [Asplenium s.
sibiricum Hk. 2 Cent. t. 35. 1860 = Camptosorus s.
subsessile Diels, Nat. Pfl. $\mathbf{1}^4$. 233. 1899 — P. brasiliensis.
vulgare Sm. Mém. Ac. Turin **5**. 421 t. 9 f. 2. 1793 = P. scolopendrium.

Scutigera Fée, Gen. Fil. 62. 1850-52 (*Neuroplatyceros §*) = Platycerium.

SCYPHOFILIX Thouars, Nov. Gen. Madag. **1** —; Römer, Collect. bot. 195. 1809 = ? **Microlepia**.

SCYPHOLEPIA J. Smith, Hist. Fil. 261. 1875 = **Microlepia**.
Hookeriana J. Sm. Hist. Fil. 262. 1875.

SCYPHULARIA Fée, Gen. Fil. 324. 1850-52 = **Davallia**.
pentaphylla Fée, Gen. 325. t. 26 B. f. 1. 1850-52.
triphylla Fée, Gen. 325. 1850-52.

SCYTOPTERIS Presl, Epim. Bot. 133. 1849 = **Cyclophorus**.
Niphobolus § Scytopteris Pr. Tent. 202. 1836.
acrostichoides Pr. Epim. 134. 1849 = C. confluens.

Selenidium Kunze, Anal. pterid. 38. 1837 (*Dicksonia? §*) = Microlepia.

Selenodesmium Prantl, Hymenoph. 53. 1875 (*Trichomanes §*) = Trichomanes.

SELLIGUEA Bory, Dict. class. d'hist. nat. **6**. 587. 1824; **15**. 344. 1829 = [**Polypodium**.
aliena Brack. Expl. Exp. **16**. 58. 1854 = ? P. selliguea.
anceps Christ, Bull. Boiss. **6**. 879. 1898.
avenia Kze. Bot. Zeit. **1846**. 420 = P. Blumeanum.
Blumeana Fée, Gen. 177. 1850-52
Blumei Kze. Bot. Zeit. **1846**. 420 = P. Blumeanum.
campyloneuroides Bedd. Handb. Suppl. 101. 1892 = P. Hosei.
caudiformis J. Sm. Ferns br. and for. 97. 1866 = P. Feei var.
coriacea Fée, Gen. 177. 1850-52 = P. loxogramme.
decurrens Pr. Tent. 216. 1836 = P. ellipticum.
elliptica Bedd. Ferns br. Ind. Index 1870.
elliptica var. flagellaris Christ, Bull. Boiss. **7**. 6. 1899 = P. flexilobum.
Feei Bory, Dict. class. **6**. 588. 1824; **17** t. 41.

Finlaysoniana Moore, Ind. LXVI. 1857 = P. digitatum.
flavescens J. Sm. JoB. **3**. 399. 1841 = ? P. scolopendrinum.
grammitoides Christ, Bull. Boiss. II. **3**. 510. 1903.
Hamiltoni Pr. Tent. 216. 1836 } = P. pedunculatum.
Hamiltoniana Bedd. Ferns br. Ind. t. 239. 1867 }
hemionitidea Pr. Tent. 216. 1836.
Henryi Christ, Bull. Boiss. **6**. 879. 1898.
heterocarpa Bl. Enum. addend. 1828.
Hookeri Pr. Tent. 216. 1836 = P. pedunculatum.
involuta Kze. Linn. **24**. 252. 1851 = P. scolopendrinum.
interrupta Fée, 1 mém. 14 t. 2 f. 25. **1844**. —?
Lanceola Fourn. Ann. sc. nat. V. **18**. 280. 1873.
lanceolata Fée, Gen. 177. 1850-52 = P. loxogramme.
macrophylla Bl. Enum. addend. 1828.
Maingayi Bedd. Ferns br. Ind. Suppl. 24. 1876.
marginata Meyen; Pr. Tent. 216. 1836; Goldm. Nova Acta **19**. Suppl. I. 452. 1843 = P. selliguea.
membranacea Bl. Enum. addend. 1828; Fl. Jav. Fil. 123 t. 52 f. 2. 1828 = [P. selliguea.
Mexicana Fée, Gen. 177. 1850-52.
minor Fée, Gen. 177. 1850-52 (nomen) — ?
pedunculata Pr. Epim. 146. 1849.
peltatisquama Fourn. Bull. Soc. Fr. **16**. 432. 1869 = P. lanceola.
plantaginea Brack. Expl. Exp. **16**. 58. 1854 = P. Feei var. caudiforme.
pothifolia J. Sm. JoB. **3**. 399. 1841 = P. ellipticum.
scolopendrina Fée, Gen. 177. 1850-52.
triphylla Christ, Bull. Boiss. **6**. 878. 1898 = P. triglossum.
Wallichiana Hk. Ic. pl. t. 204. 1840 = P. scolopendrinum.
Wrightii J. Sm. Hist. Fil. 102. 1875.

Sericonitis J. Smith, Hist. Fil. 150. 1875 *(Hemionitis §)* = Gymnopteris et Trachypteris.

SERPYLLOPSIS van den Bosch, Versl. Akad. Wet. Amsterdam **11**. 318. 1861 = **Trichomanes**.
antarctica v. d. B. Ned. Kr. Arch. **4**. 377. 1859 (syn.) = T. caespitosum.

SITOBOLIUM Desvaux, Prodrome 262. 1827 = **Dennstaedtia**.
punctilobum Desv. Prod. 263. 1827 = D. punctilobula.

SITOLOBIUM J. Smith, Journ. Bot. **3**. 418. 1841; Lond. Journ. Bot. **1**. 434. [1842 = **Dennstaedtia**.
adiantoides J. Sm. Lond. JoB. **1**. 434 1842.
anthriscifolium J. Sm. Ferns br. and for. 237. 1866.
apiifolium J. Sm. Lond. JoB. **1**. 434. 1842 = D. adiantoides.
cicutarium J. Sm. Lond. JoB. **1**. 434. 1842.
cuneatum J. Sm. JoB. **3**. 418. 1841.
davallioides J. Sm. Lond. JoB. **1**. 434. 1842.
dissectum J. Sm. Lond. JoB. **1**. 434. 1842.
dubium Brack. Expl. Exp. **16**. 273. 1854 = Davallia ? d.
flaccidum J. Sm. JoB. **3**. 418. 1841 = D. Smithii.
glutinosum J. Sm. Lond. JoB. **1**. 434. 1842 = D. appendiculata.
moluccanum J. Sm. Cat. cult. Ferns 70. 1857.
obtusifolium J. Sm. Lond. JoB. **1**. 434. 1842.
Pavoni J. Sm. Ferns br. and for. 237. 1866.
pilosiusculum J. Sm. Lond. JoB. **1**. 434. 1842 = D. punctilobula.

Plumieri J. Sm. Ferns br. and for. ed. II. 319. 1877 = Saccoloma domingense.
punctilobum J. Sm. Bot. Mag. **72**. Comp. 36. 1846 = D. punctilobula.
rubiginosum J. Sm. Lond. JoB. **1**. 434. 1842.
samoense Brack. Expl. Exp. **16**. 274 t. 38 f. 1. 1854.
? scandens Brack. Expl. Exp. **16**. 275. 1854.
Smithii J. Sm. Ferns br. and for. ed. II. 319. 1877.
solidum Lodd. Cat. 1849.
stramineum Brack. Expl. Exp. **16**. 273. 1854 = Balantium s.
strigosum J. Sm. Lond. JoB. **1**. 434. 1842 = D. scabra.
tenerum Brack. Expl. Exp. **16**. 275. 1854.

SOLENOPTERIS Wallich, Tidsskr. f. Naturvid. **1**. 259. 1822; Flora **1823**[1]. 65 (nomen) = **Drymoglossum et Antrophyum**

SOLENOPTERIS Zenker msc. (Kze. Linn. **24**. 267. 1851) = **Athyrium**.

SOROMANES Fée, 2 mém. Hist. d'Acrost. 16. 1845 = **Polybotrya**.
Caenopteris Christ, Bull. Boiss. II. **3**. 613. 1903 } = P. serratifolia.
dentata Fée, Acrost. t. 43. 1845 } = P. serratifolia.
integrifolia Fée, Acrost. 82 t. 42. 1845 } = P. serratifolia.
serratifolia Fée, Acrost. 82. 1845 } = P. serratifolia.

SPATHEPTERIS Presl, Suppl. Tent. Pterid. 95. 1845 = **Aneimia**.
verticillata Pr. Suppl. 95. 1845.

SPHAEROCIONIUM Presl, Hymenoph. 33. 1843 = **Hymenophyllum**.
abietinum Pr. Hym. 35. 1843 = H. polyanthos.
asterothrix Fée, 9 mém. app. 1865 = H. hirsutum.
aureum Pr. Hym. 34, 57. 1843 = H. plumosum.
australe Pr. Hym. 35. 1843.
axillare Pr. Hym. 35. 1843.
badium Pr. Hym. 35. 1843 = H. dilatatum*.
bivalve Pr. Hym. 34. 1843.
Boryanum Pr. Hym. 34. 1843 = H. ciliatum.
Breutelii Pr. Abh. böhm. Ges. V. **5**. 337. 1848 = H. polyanthos var.
Bridgesii Kl. Linn. **18**. 536. 1844 = H. dentatum.
Bulnesianum Schaffn.; Fée, 9 mém. 39. 1857 = H. undulatum.
caudiculatum Pr. Hym. 35. 1843.
ciliatum Pr. Hym. 34. 1843.
commutatum Pr. Hym. 34. 1843 = H. ciliatum.
crispatum Pr. Hym. 35. 1843 = H. australe.
crispum Kl. Linn. **18**. 537. 1844.
cristatum Pr. Hym. 34. 1843 = H. fucoides.
demissum Pr. Hym. 35. 1843.
diffusum Hassk. Obs. Bot. Fil. **2**. 26. 1857-58 = Trichomanes d.
dilatatum Pr. Hym. 35. 1843.
diversilobum Pr. Hym. 34, 59. 1843.
elasticum Pr. Hym. 34 t. 4 f. B. 1843 = H. hygrometricum.
Gaudichaudii Hassk. Obs. Bot. Fil. **2**. 29. 1857-58 = ? H. recurvum.
glanduliferum Pr. Epim. 22 t. 12. 1849 = H. scabrum.
gracile Pr. Hym. 35. 1843 = H. inaequale.
Grevilleanum Pr. Hym. 34. 1843 = H. ciliatum.
hirsutum Pr. Hym. 34. 1843.
hirtellum Pr. Hym. 34. 1843.
infortunatum Pr. Hym. 35. 1843 = H. capillaceum.

interruptum Pr. Hym. 34 t. 10 f. C. 1843.
lineare Pr. Hym. 34. 1843.
macrocarpum Pr. Hym. 35, 61. 1843 = H. dilatatum*.
myriocarpum Kl. Linn. **18**. 536. 1844.
nigricans Pr.; Kl. Linn. **18**. 536. 1844 = H. nigrescens.
pendulum Pr. Hym. 34. 1843 = H. capillare.
Plumieri Pr. Hym. 34. 1843 = H. ciliatum.
productum Pr. Hym. 35, 61. 1843 = H. caudiculatum var.
pulchellum Pr. Hym. 34. 1843 = H. sericeum var.
ricciaefolium Pr. Hym. 35. 1843 = H. tenellum.
Ruizianum Kl. Linn. **18**. 535. 1844.
rupestre Pr. Hym. 35. 1853.
sanguinolentum Pr. Hym. 35. 1843.
scabrum Pr. Hym. 34. 1843.
Schiedeanum Pr. Hym. 34, 60. 1843 = H. ciliatum.
sericeum Pr. Hym. 34. 1843.
Sieberi Pr. Hym. 34, 58. 1843 = H. ciliatum.
Smithii Hassk. Obs. Bot. Fil. **2**. 27. 1857-58 = H. australe.
tomentosum Pr. Hym. 34. 1843.
trifidum Pr. Hym. 34. 1843 = H. lineare.
undulatum Pr. Hym. 35. 1843.
vestitum Pr. Hym. 34, 58. 1843 = H. ciliatum.

Sphaerodium Presl, Hymenoph. 31. 1843 *(Hymenophyllum §)* = Hymenophyllum.

SPHAEROPTERIS Bernhardi, Schrad. Journ. **1800²**. 122. 1801 = **Cyathea.**
medullaris Bernh. l. c.

SPHAEROPTERIS Wallich, List n. 183. 1828; Pl. Asiat. Rar. **1**. 42. t. 52. 1830 [= **Peranema.**
barbata Wall. l. c. = P. cyatheoides.
Hookeriana Wall. List. pag. 248 (n. 775). 1829 = Diacalpe aspidioides.

SPHAEROSTEPHANOS J. Smith in Hook. Gen. Fil. t. 24. 1839 = **Mesochlaena.**
asplenioides J. Sm. l. c. = M. polycarpa.

SPHAEROSTICHUM Presl, Epim. Bot. 134. 1849 = **Cyclophorus.**
abbreviatum Pr. Epim. 135. 1849.
acrostichoides Pr. Epim. 135. 1849 = C. sphaerostichus.

Sphenopteris Mettenius, Fil. Lechl. **1**. 15. 1856 *(Asplenium §)* = Asplenium.

SPICANTA Presl, Epim. Bot. 114. 1849; O.Ktze. Rev. Gen. Pl. **2**. 820. 1891 = **Blechnum.**

Obs. Omnes species Blechni et Lomariae a cl. O.KUNTZE ad Spicantam translatae sunt. Homonyma sub Blechno citata sunt.

acuminata O.Ktze. Rev. **2**. 821. 1891 = B. norfolkianum.
auriculata O.Ktze. Rev. **2**. 821. 1891 = B. Fullageri.
blechnodes O.Ktze. Rev. **2**. 821. 1891 = B. valdiviense.
borealis Pr. Epim. 114. 1849 = B. spicant.
caudata O.Ktze. Rev. **2**. 821. 1891 = B. Sprucei.
ciliata O.Ktze. Rev. **2**. 821. 1891 = B. Moorei.
deflexa O.Ktze. Rev. **2**. 822. 1891 = B. Faberi.
hastata O.Ktze. Rev. **2**. 821. 1891 = B. australe*.
Hookeriana O.Ktze. Rev. **2**. 821. 1891 = B. volubile var.
onocleoides Pr. Epim. 114. 1849 = B. polypodioides.
osmundodes O.Ktze. Rev. **2**. 821. 1891 = B. tabulare.
pubescens O.Ktze. Rev. **2**. 822. 1891 = B. Bakeri.

pumila O.Ktze. Rev. **2**. 822. 1891 = B. parvifolium.
stenophylla O.Ktze. Rev. **2**. 822. 1891 = B. Humblotii.
unilateralis O.Ktze. Rev. **2**. 822. 1891 = B. blechnoides.

STEGANIA R. Brown, Prod. Fl. Nov. Holl. 152. 1810 = **Blechnum**.
alpina R.Br. Prod. 152. 1810 = B. penna marina.
discolor A.Rich. Fl. N. Zél. 87. 1832.
falcata R.Br. Prod. 153. 1810 = B. discolor.
fluviatilis R.Br. Prod. 152. 1810.
lanceolata R.Br. Prod. 152. 1810.
minor R.Br. Prod. 153. 1810 = B. capense.
nuda R.Br. Prod. 153. 1810 = B. discolor var.
onocleoides Gray, Nat. Arr. Br. Pl. **2**. 16. 1821 = Cryptogramma crispa.
Patersoni R.Br. Prod. 152. 1810.
procera R.Br. Prod. 153. 1810 = B. capense.

STEGNOGRAMMA Blume, Enum. pl. Javae 172. 1828.
aspidioides Bl. Enum. 173. 1828 = Dryopteris stegnogramme.
Ehrenbergiana Fourn. Mex. pl. **1**. 71. 1872 = Gymnopteris.
Javanica Fée, Gen. 204. 1850-52 } = Mesochlaena polycarpa.
Mesochlaena Fée, Gen. 204. 1850-52 } = Mesochlaena polycarpa.
Moluccana Fée, Gen. 204. 1850-52 } = Mesochlaena polycarpa.
Sandwicensis Brack. Expl. Exp. **16**. 26 t. 4 f. 2. 1854 = Dryopteris stenogrammoides.

STENOCHLAENA J. Smith, Journ. Bot. **3**. 401. 1841; **4**. 149. 1841 — (Polypodiaceae Gen. **54**).
Conf. Underwood, Bull. Torr. Cl. **33**. 35-50. 1906.

aculeata (Bl.) Kze. Bot. Zeit. **1848**. 142 — Asia trop. — Lomaria Bl. Enum. 205. 1828; Olfersia Bl. Fl. Jav. Fil. Suppl. t. 96 —; Teratophyllum Mett.; Kuhn 1869; Acrostichum Racib. 1898. Lomariopsis spinescens Fée 1845. Davallia achilleaefolia Wall. 1828. Lomaria polymorpha Reinw.; Zoll. 1845. *Obs.* Species anomala, t. auctt. plur. = S. sorbifolia vel S. palustris forma.
Blumeana Pr. Epim. 163. 1849 = S. palustris.
decrescens (Bak.) Und. Bull. Torr. Cl. **33**. 46 f. 5. 1906 — Afr. occ. trop. Acrostichum Bak. 1868. HB. 412; Lomariopsis Kuhn 1869; Acrostichum polyphyllum Hk. sp. **5**. 243. 1864 (non pag. 269); Lomariopsis Kuhn 1868; Chrysodium Salom. 1883.
feejeensis Brack. Expl. Exp. **16**. 78 t. 11 f. 1. 1854 = Blechnum filiforme.
fraxinifolia Pr. Epim. 164. 1849 = S. palustris.
gracilis (Bl.) Kze. Bot. Zeit. **1848**. 142 — Java.... Lomaria Bl. Enum. 205. 1828; Olfersia Bl. Fl. Jav. Fil. Suppl. t. 95 —; Acrostichum Racib. 1898. — vide Obs. sub S. aculeata.
Haenkeana Pr. Epim. 165. 1849 = S. palustris.
heteromorpha J. Sm. JoB. **4**. 149. 1841 = Blechnum filiforme.
japurensis Gris. Fl. br. W. Ind. 676. 1864 = S. marginata.
juglandifolia Pr. Epim. 164. 1849 = S. palustris.
laurifolia Pr. Epim. 164. 1849. NPfl. 252 — Celebes. Ins. Philipp. Ins. Salomonis. — Acrostichum Hk. 1864. HB. 412.
limonifolia J. Sm. JoB. **4**. 149. 1841 = S. sorbifolia var. 20.
longifolia J. Sm. JoB. **3**. 402. 1841 = S. sorbifolia var. 7.
marginata (Schrad.) C. Chr. Ind. 624. 1906 — Amer. trop. Afr. occ. et orient. trop. — Lomaria Schrad. Gött gel. Anz. **1824**. 871; Lomariopsis Kuhn 1879. Lomaria scandens Raddi 1819 (non Willd. 1810); L. fraxinifolia Raddi 1825.

Acrostichum japurense Mart. 1834; Olfersia Pr. 1836; Stenochlaena Gris. 1864; Lomariopsis J. Sm. 1875 — Acrostichum erythrodes Kze. 1839; Lomariopsis Fée 1845 — Acrostichum phlebodes Kze. 1834; Lomariopsis Fée 1845 — L. Prieuriana Fée 1845.

Meyeriana Pr. Epim. 166. 1849 = S. tenuifolia.

oleandraefolia Brack. Expl. Exp. **16**. 75. 1854 = S. sorbifolia var. 9.

palustris (Burm.) Bedd. Ferns br. Ind. Suppl. 26. 1876. NPfl. 251 — Asia, Austr., Polynesia trop. — Polypodium Burm. Fl. Ind. 234. 1768; Lomariopsis Kuhn 1869. Chr. 41; Chrysodium Luerss. 1871; Acrostichum Clarke 1880; Onoclea scandens Sw. 1806; Lomaria Willd. 1810; Olfersia Pr. 1836; Stenochlaena J. Sm. 1841; Pteris Roxb. 1844; Lomariopsis Mett. 1856; Acrostichum Hk. 1864. HB. 412 (non Raddi 1819 nec Bory; Fée 1845). — Conf. S. aculeata et S. gracilis.

Pittieri (Christ) Diels, Nat. Pfl. 1⁴. 251. 1899 — Costa Rica. — Acrostichum Christ, Bull. Soc. bot. Belg. **35**. 243. 1896; Lomariopsis Christ 1901.

pollicina (Willem.) C. Chr. Ind. 7. 1905; Und. Bull. Torr. Cl. **33**. 49. f. 8. 1906 — Madagascar. Ins. Mascarenae. Afr. occ. — Osmunda Willem. Usteri Annal. **18**. 61. 1796; Lomariopsis Mett.; Kuhn 1868. Osmunda trifrons Bory 1804; Acrostichum Commers. 1816. Lomaria fraxinea Willd. 1810; Lomariopsis J. Sm 1866; Lomariopsis integrifolia Klf.; Spr. 1827; Olfersia Pr. 1836 — Acrostichum lomarioides Bory 1833; Olfersia Pr. 1836; Lomariopsis Boryana Fée 1845 — Polypodium orbiculatum Poir. 1804; Lomaria Desv. 1827. Acrostichum buxifolium Kze. 1845; Lomariopsis Fée 1845 — ? Polypodium? binerve Hk. 1862.

? pycnophylla Pr. Epim. 165. 1849 = Plagiogyria.

Raciborskii C. Chr. Ind. 18. 1905; 625. 1906 — Ins. Moluccae. Acrostichum Smithii Racib. Bull. int. Ac. Cracovie **1902**. 59 (non Bak. 1891).

scandens J. Sm. JoB. **3**. 401. 1841 = S. palustris.

sessilifolia Pr. Epim. 163. 1849 = S. palustris.

sorbifolia (L.) J. Sm JoB. **4**. 149. 1841. NPfl. 251 — Trop. (Typ. Amer. trop.) Acrostichum L. sp. **2**. 1069. 1753. HB. 412; Onoclea Sw. 1806; Lomaria Klf. 1824; Olfersia Pr. 1836; Lomariopsis Fée 1845. Chr. 39; Chrysodium Luerss. 1871; Polybotrya Keys. 1873.

Subspecies (vel species?).

— 1. Lomaria variabilis Willd. sp. **5**. 294. 1810; Onoclea Poir. 1816; Olfersia Pr. 1836; Lomariopsis Fée. Acrost. 70 t. 31. 1845; Lomaria marattiaefolia Desv. 1827. L. acrostichoides Klf.; Spr. 1827 — Lomariopsis cuspidata Fée, Acrost. 68 t. 27. 1845 — Mauritius.

— 2. Lomaria longifolia Klf. Enum. 153. 1824 — Amer. trop.

— 3. Acrostichum alatum Roxb. Calc. Journ. **4**. 480. 1844. Lomaria? spondiaefolia Wall. 1828; Stenochlaena J. Sm. 1841 (spondicifolia, nomen); Pr. 1849; Lomariopsis Moore 1857. L. cochinchinensis Fée, Acrost. 66 t. 26. 1845 — Malesia. Cochinchina.

— 4. Lomariopsis elongata Fée. Acrost. 67. 1845 — Brasilia.

— 5. Lomariopsis leptocarpa Fée. Acrost. 69 t. 29. 1845. Stenochlaena variabilis Brack. 1854; Lomariopsis Brackenridgei Carr. 1873 — Ins. Philipp. Polynesia.

— 6. Lomariopsis recurvata Fée, Acrost. 68 t. 28. 1845 —

— 7. Lomariopsis Smithii Fée. Acrost. 71 t. 33 f. 2. t. 53. 1845: Stenochlaena longifolia J. Sm. 1841 (nomen): Lomariopsis J. Sm. 1857 — Luzon.

— 8. Lomaria spectabilis Kze. Bot. Zeit. **1848**. 144: Lomariopsis Mett. 1856; Acrostichum Racib. 1898 — Malesia.

— 9. Stenochlaena oleandrifolia Brack. Expl. Exp. **16**. 75. 1854; Lomariopsis Mett., Kuhn 1869 — Polynesia.

—10. Lomariopsis Fendleri Eat. Mem. Amer. Ac. n. s. **8**. 195. 1860 — Venezuela.

—11. Lomariopsis Wrightii Mett.: Eat. Mem. Amer. Ac. n. s. **8**. 195. 1860: Stenochlaena Gris. 1866 — Cuba.

—12. Lomariopsis Novae-Caledoniae Mett. Ann. sc. nat. IV. **15**. 58. 1861 — N. Caledonia.

—13. Acrostichum Brightiae F. Muell. Fragm. **7**. 119. 1870; Lomariopsis Bail. 1874 — Queensland.

—14. Lomariopsis vestita Fourn. Bull. Soc. Fr. **19**. 250. 1872 — Nicaragua.

—15. Lomariopsis Balansae Fourn. Ann. sc. nat. V. **18**. 271. 1873 — N. Caledonia. — Vide S. marginata et S. pollicina. Conf. Lomariopsis.

Formae monstrosae: (an ad S. palustrem pertinentes?).

—16. Scolopendrium D'Urvillaei Bory. Dup. Voy. Bot. **1**. 273 t. 37 f. 1. 1828. HB. 247: Micropodium Mett. 1866; Phyllitis O.Ktze. 1891 — Melanesia.

—17. Gymnogramme subtrifoliata Hk. sp. **5**. 152 t. 298. 1864; Toxopteris Trevis 1877 — Fiji.

—18. Lindsaea Parishii Bak. Syn. ed. I. 109. 1867 — Tenasserim.

—19. Triphlebia dimorphophylla Bak. Malesia **3**. 42 t. 4. 1886. NPfl. 230 — N. Guinea.

—20. ? Lomariopsis ludens Fée, Acrost. 70 t. 30. 1845 — Lomaria? limonifolia Wall. 1828: Stenochlaena J. Sm. 1841. Teratophyllum aculeatum var. inermis Mett. Ann. Lugd. Bat. **4**. 297. 1869 — Malesia. Polynesia. — (Bedd. Ferns br. Ind. t. 209. 210. 1866). — Vide S. aculeata et S. gracilis.

spondiaefolia Pr. Epim. 165. 1849 } = S. sorbifolia var. 3.
spondicifolia J. Sm. JoB. **4**. 149. 1841 }

tenuifolia (Desv.) Moore, Gard. Chr. **1856**. 193 c. fig. NPfl. 253 — Afr. trop. et austr. cum ins. adjac. — Lomaria Desv. Berl. Mag. **5**. 326. 1811; Lomariobotrys Fée 1850-52; Polybotrya Kuhn 1868; Acrostichum Bak. 1868. HB. 412; Lomariopsis Christ 1897. 42; Lomaria Meyeriana Kze. 1836: Stenochlaena Pr. 1849; Lomariobotrys Fée 1850-52; Polybotrya Mett. 1856; Acrostichum Hk. 1862.

triquetra J. Sm. JoB. **4**. 149. 1841 = Plagiogyria euphlebia.

variabilis Brack. Expl. Exp. **16**. 76. 1854 = S. sorbifolia var. 5.

Wrightii Gris. Cat. Pl. Cub. 277. 1866 = S. sorbifolia var. 11.

yapurensis vide japurensis.

?STENOCHLAMYS Griffith. Notulae **2**. 610. 1849.

Edgeworthii Griff. Notulae **2**. 611. 1849: Ic. pl. as. **2**. t. 138 f. 1 — India bor. (Davallia sp. dub.).

Stenogramme Klotzsch, Linnaea **20**. 411. 1847 *(Gymnogramme §)* = Gymnopteris (Bommeria).

Stegnogramme Fourn. Mex. pl. **1**. 71. 1872; vide Stegnogramme.

STENOLOBUS Presl. Tent. Pterid. 130. 1836 = **Davallia.**

Fejeensis Pr. Epim. 99. 1849 = D. solida var.

Kunzeanus Pr. Tent. 130. t. 4 f. 30. 1836 = D. solida.

mauritianus Pr. Epim. 99. 1849.
ornatus Pr. Tent. 130. 1836 = D. solida.
pentaphyllus Pr. Epim. 99. 1849.
solidus Pr. Tent 130 1836.
triphyllus Pr. Epim. 99. 1849.

STENOLOMA Fée, Gen. Fil. 330. 1850-52 = **Odontosoria.**
aculeatum Fée, Gen. 330. t. 27 bis A. f. 4. 1850-52.
Blumeanum Fée, Gen. 330. t. 27 bis A. f. 2. 1850-52 = Lindsaya B.
?capillaceum Fée, 11 mém. 93. 1866 = Trichomanes c.
chinensis Bedd. Handb. 70. 1883.
clavatum Fée, Gen. 330. t. 27 bis A. f. 3. 1850-52.
dumosum Fée, Gen. 330. 1850-52 = O. aculeata.
fumarioides Fée, Gen. 330. t. 27 bis A. f. 6. 1850-52.
Glaziovii Fée, Cr. vasc. Br. **1**. 153 t. 52 f. 2. 1869 = O. bifida.
?Goudotianum Fée, Gen. 330. 1850-52 = Lindsaya G.
gratissima Fée, Cr. vasc. Br. **1**. 153 t. 52 f. 1. 1869 = O. meifolia.
?lindsayoides Fée, Gen. 330. t. 27 bis A. f. 5. 1850-52 = Lindsaya microphylla.
retusum Fée, Gen. 330. 1850-52.
Schlechtendalii Fée, Gen. 330. t. 27 bis A. f. 1. 1850-52.
?tegulare Fée, 11 mém. 93. 1866 = Davallia t.
tenuifolium Fée, Gen. 330. 1850-52 = O. chinensis.
?trifoliatum Fée, 11 mém. 93. 1866 = Davallia t.

Stenoneura Christ, Monogr. Elaph. 20. 1899 (*Elaphoglossum* §) = Elaphoglossum.

STENOSEMIA Presl, Tent. Pterid. 237. 1836 — (Polypodiaceae Gen. **22**).
aurita (Sw.) Pr. Tent. 237. t. 10 f. 24. 1836. NPfl. 198 — Malesia. Ins. Salomonis. — Acrostichum Sw. Schrad. Journ. **1800**². 12. 1801. HB. 416; Polybotrya Bl. 1828. Gymnopteris Keys. 1873. Chr. 52 — Polybotrya cicutaria Bl. 1828; Stenosemia? Pr. 1836; Acrostichum Hk. 1864.
aurita J. Sm. in Hk. Gen. t. 94 f. 1-4. 1842 = Aspidium ambiguum.
?cicutaria Pr. Tent. 237. 1836 = S. aurita.
Teysmanniana (Bak.) Diels. Nat. Pfl. **1**⁴. 198. 1899 — Ins. Sumba (Malesia). Acrostichum Bak. Malesia **3**. 56. 1886.

STIBASIA Presl, Suppl. Tent. Pterid. 15. 1845 = **Marattia.**
Douglasii Presl, Suppl. 16. 1845.

STICHERUS Presl, Tent. Pterid. 51. 1836 = **Gleichenia.**
laevigatus Pr. Tent. 52. 1836.
laniger Pr. Tent. 52. 1836.

STROMATOPTERIS Mettenius. Ann. sc. nat. IV. **15**. 84. 1861 — (Gleicheniaceae Gen. **1**).
moniliformis Mett. Ann. sc. nat. IV. **15**. 84 t. 3. 1861. NPfl. 352 — N. Caledonia. Gleichenia Moore 1862. HB. 11. Chr. 339.

STRUTHIOPTERIS Weis. Plant. Crypt. Fl. Gott. 286. 1770 = **Blechnum.**
appendiculata Trev. Atti Ist. Veneto III. **14**. 572. 1869.
australis Trev. l. c. 572. 1869.
doodioides Trev. l. c. 571. 1869 = B. spicant var.
Dregeana Trev. l. c. 572. 1869 = B. punctulatum.
hastata Trev. l. c. 572. 1869 = B. auriculatum.
japanensis Trev. l. c. 571. 1869 = B. spicant var.
javanica Trev. l. c. 573. 1869 = B. Blumei.
mucronulata Trev. l. c. 572. 1869 = Lomaria m.

plantaginea Trev. l. c. 573. 1869 = B. lanceola.
polypodioides Trev. l. c. 571. 1869.
punctulata Trev l. c. 573. 1869.
remota Trev. l. c. 572. 1869 = B. auriculatum var.
rigida Trev. l. c. 572. 1869 = B. australe.
Spicant Weis, Pl. Cr. Gott. 287. 1770.
triloba Trev. l. c. 572. 1770 = B. auriculatum var.

STRUTHIOPTERIS Bernhardi. Schrad. Journ. **1800**[2]. 126. 1801 = **Osmunda.**
cinnamomea Bernh. Schrad. Journ. **1800**[2]. 126. 1801.
claytoniana Bernh. Schrad. Journ. **1800**[2]. 126. 1801.
regalis Bernh. Schrad. Journ. **1800**[2]. 126. 1801.

STRUTHIOPTERIS Willdenow, Mag. Ges. Nat. Fr. Berlin **3**. 160. 1809 = **Matteuccia.**
Cavaleriana Christ, Bull. Ac. Géogr. Bot. Mans **1904**. 118 c. figg.
crispa Wallr. Fl. Cr. Germ. **1**. 27. 1831 = Cryptogramma c.
europaea Hornem. Hort. reg. bot. Hafn. **2**. 945. 1815 = M. struthiopteris.
filicastrum All. Fl. Pedem. **2**. 283. 1785 = M. struthiopteris.
germanica Willd. Enum. 1071. 1809 = M. struthiopteris.
nodulosa Desv. Prod. 287. 1827 = Woodwardia areolata.
orientalis Hk. 2 Cent. t. 4. 1860.
pensylvanica Willd. sp. **5**. 289. 1810 = M. struthiopteris.

SYMPHLEBIUM Fée, Gen. Fil. 386. 1850-52 = Synaphlebium.

Symplecium Kunze, Bot. Zeit. **1846**. 422 *(Polypodium §)* = Polypodium.

SYNAMMIA Presl, Tent. Pterid. 212. 1836 = **Polypodium.**
elongata Pr. Tent. 212. 1836 = P. lanceolatum*.
triloba Pr. Tent. 212. t. 9 f. 11. 1836 = P. synammia.

SYNAPHLEBIUM J. Smith, JoB. **3**. 415. 1841 et in Hook. Gen. Fil. t. 101. 1842 [= **Lindsaya.**
davallioides J. Sm. Lond. JoB. **1**. 424. 1842.
nitens J. Sm. Hist. Fil. 268. 1875 = L. decomposita.
obtusum J. Sm. JoB. **3**. 415. 1841 = L. decomposita var.
Pickeringii Brack. Expl. Exp. **16**. 223 t. 30 f. 2. 1854.
propinquum Fée, Gen. 109. 1850-52 = L. decomposita var.
pulchrum Brack. Expl. Exp. **16**. 223. 1854 = L. decomposita.
recurvatum J. Sm. JoB. **3**. 415. 1841; Hk. Gen. t. 101. 1842 = L. decomposita var.
tetraphyllum Fée, Gen. 109. 1850-52 = Adiantum t.
Urvillei Fée, Gen. 110. 1850-52 = L. decomposita.

Synechia Fée, Gen. Fil. 113. 1850-52 *(Adiantum §)* = Adiantum.

SYNEURON J. Smith msc. (Hk. Ic. plant. t. 950. 1854) = **Dryopteris** (stegnogramme).

SYNGRAMMA J. Smith, Lond. Journ. Bot. **4**. 166. 1845 — (Polypodiaceae Gen. 58).

alismifolia (Pr.) J. Sm. Lond. JoB. **4**. 168 t. 7-8 f. B. 1845. NPfl. 257 — Malesia. Ins. Philipp. — Diplazium Pr. Rel. Haenk. **1**. 49 t. 8. f. 3. 1825; ? Oxygonium Pr. 1836 (non J. Sm. 1842); Gymnogramme Hk. 1864. HB. 386; Hemionitis Trev. 1877; Trichiogramme Kuhn 1882; Neurogramme Christ 1897. 64. Callogramme Caeciliae Fée 1850-52; Gymnogramme Mett. 1856 — G. Wallichii Hk. 1864. HB. 386; Syngramma Bedd. 1866; Dictyogramme Trev. 1877.

borneensis (Hk.) J. Sm. Hist. Fil. 152. 1875. NPfl. 257 — Borneo. Celebes. Fiji. Gymnogramme Hk. sp. **5**. 154 t. 301. 1864. HB. 386; Toxopteris Trev. 1877; Trichiogramme Kuhn 1882.

cartilagidens (Bak.) Diels, Nat. Pfl. 1^4. 257. 1899 — Borneo.
Gymnogramme Bak. Syn. 386. 1868; Toxopteris Trev. 1877; Trichiogramme Kuhn 1882; Neurogramme Chr. 1897. 63.

Dayi Bedd. Handb. Suppl. 100. 1892 — Perak.
Gymnogramme Bedd. JoB. **1888**. 5. t. 279 B.

?**elaphoglossoides** (Bak.) Diels, Nat. Pfl. 1^4. 257. 1899 — Guiana.
Gymnogramme Bak. Tr. Linn. Soc. II. Bot. **2**. 293 t. 54. 1887; Neurogramme Chr. 1897. 63.

fraxinea Bedd. Handb. 386. 1883 = Coniogramme.

Hookeri C. Chr. Ind. 346. 1905; 629. 1906 — N. Guinea. Fiji.
Hemionitis lanceolata Hk. 2 Cent. t. 55. 1861. HB. 398. Chr. 59 (non L. 1753); Gymnogramme Luerss. 1871; Trichiogramme Kuhn 1882; Syngramme Diels 1899. NPfl. 257.

Hosei (Bak.) Diels, Nat. Pfl. 1^4. 257. 1899 — Borneo.
Hemionitis Bak. JoB. **1891**. 108.

lanceolata Diels, Nat. Pfl. 1^4. 257. 1899 = S. Hookeri.

Lobbiana (Hk.) J. Sm. Hist. Fil. 152. 1875. NPfl. 257 — Borneo.
Gymnogramme Hk. sp. **5**. 153 t. 300. 1864. HB. 386; Toxopteris Trev. 1877; Trichiogramme Kuhn 1882.

marginata (Mett.) Diels, Nat. Pfl. 1^4. 257. 1899 — N. Caledonia.
Gymnogramme Mett. Ann. sc. nat. IV. **15**. 59. 1861. HB. 378; Anstrogramme Fourn. 1873; Trichiogramme Kuhn 1882; Neurogramme Chr. 1897. 63.

obtusifolia (Hk.) J. Sm. Hist. Fil. 152. 1875. NPfl. 257 — Java. Sumatra.
Gymnogramme Hk. sp. **5**. 153 t. 299. 1864. HB. 386; Dictyogramme Trev. 1877; Trichiogramme Kuhn 1882.

pinnata J. Sm. Lond. JoB. **4**. 168 t. 7-8 f. C. 1845. NPfl. 257 — Austr. trop. Polynesia occ. — Dictyogramme Moore 1861; Gymnogramme Hk. 1864. HB. 390; Grammitis F. Muell. 1868; Trichiogramme Kuhn 1882 — Hemionitis elongata Brack. 1854 (non Cav. 1802); Dictyogramme Moore 1861.

quinata (Hk.) Carr. in Seem. Fl. Vit. 372. 1873. NPfl. 257 — Malesia. N. Guinea. Ins. Salomon. — Gymnogramme Hk. sp. **5**. 152 t. 297. 1864. HB. 387; Dictyogramme Trev. 1877; Trichiogramme Kuhn 1882; Neurogramme Chr. 1897. 64.

scolopendrioides (Bak.) C. Chr. Ind. 340. 1905; 629. 1906 — Fiji.
Gymnogramme Bak. JoB. **1879**. 299.

valleculata (Bak.) C. Chr. Ind. 340. 1905; 629. 1906 — Borneo.
Gymnogramme Bak. JoB. **1888**. 325.

vestita Moore, Ind. LX 1857 = Gymnopteris.

vittaeformis J. Sm. Lond. JoB. **4**. 168 t. 7-8 f. A. 1845. NPfl. 257 — Ins. Philipp. — Callipteris J. Sm. 1841 (nomen); Oxygonium J. Sm. 1841 (nomen); Gymnogramme Mett. 1856. HB. 386; Hemionitis Trev. 1877; Trichiogramme Kuhn 1882.

Wallichii Bedd. Ferns br. Ind. t. 153. 1866 = S. alismifolia var.

Zollingeri (Kurz) Diels, Nat. Pfl. 1^4. 257 — Malesia. Ins. Philipp.
Hemionitis Kurz, Journ. As. Soc. Bengal 39^2. 90 t. 5 1870. HB. 518. Chr. 59; Gymnogramme Salom. 1883.

SYNOCHLAMYS Fée, 7 mémoire, 35 t. 20 f. 4. 1857 = **Pellaea.**
ambigua Fée, l. c. 1857.

TAENIOPSIS J. Smith, Journ. Bot. **4**. 67. 1841 = **Vittaria.**
amboynensis Bedd. Ferns br. Ind. t. 117. 1866.

falcata Bedd. Ferns br. Ind. t. 175. 1866 = V. sulcata.
falcata Bedd. Ferns br. Ind. Suppl. 25. 1876 = V. pusilla.
Forbesii Moore, Ind. XXXIII. 1857 = V. scolopendrina.
furcata Moore, Ind. XXXIII. 1857 = Eschatogramme f.
graminifolia J. Sm. JoB. **4**. 67. 1841.
lineata J. Sm. JoB. **4**. 67. 1841.
mauruensis Nadeaud, Journ. de Bot. **13**. 8. 1899.
revoluta J. Sm. Job. **4**. 67. 1841.
Richiana Brack. Expl. Exp. **16**. 60. 1856 = V. scolopendrina.
scolopendrina J. Sm. JoB. **4**. 67. 1841.
stipitata Moore, Ind. XXXIII. 1857.
sulcata Bedd. Ferns br. Ind. Suppl. 25. 1876.
tricuspidata Moore. Ind. XXXIII. 1857 = Eschatogramme furcata.
zeylanica Moore, Ind. XXXIII. 1857 = V. scolopendrina.

TAENIOPTERIS Hook. Gen. Fil. t. 76 B. 1842 = **Vittaria**.
Forbesii Hk. l. c. = V. scolopendrina.

TAENITIS Willdenow; Sprengel, Anleit. **3**. 374. 1804; Schkuhr, Kr. Gew. **1**. 21. 1804; Sw. Syn. 3. 1806 — (Polypodiaceae Gen. **96**).
augustifolia Spr. Syst. **4**. 42. 1827 = Vittaria.
blechnoides (Willd.) Sw. Syn. 24. 220. 1806. HB. 397. Chr. 130. NPfl. 305 — Asia trop.-Fiji. — Pteris Willd. Phyt. 13 t. 9 f. 3. 1794; Oetosis O.Ktze. 1891: — Taenitis pteroides Schkuhr 1804 — T. chinensis Desv. 1811 — T. obtusa Hk. 1854. HB. 396: Oetosis Ktze. 1891 — T. interrupta Hk. et Grev. 1828.
Blumei Hk. sp. **5**. 187. 1864 = Antrophyum callifolium.
carnosa Mett. Fil. Lips. 28. 1856 = Drymoglossum.
chinensis Desv. Berl. Mag. **5**. 308. 1811 = T. blechnoides.
Desvauxii Kl. Linn. **20**. 431. 1847 = Eschatogramme furcata var.
furcata Willd. sp. **5**. 136. 1810 = Eschatogramme.
graminifolia Hk. Exot. Fl. **2**. t. 77. 1825 = Monogramma seminuda.
graminoides Mett. Fil. Lips. 27. 1856 = Monogramma.
interrupta Hk. et Grev. Ic. Fil. t. 62. 1828 = T. blechnoides var.
lanceolata Klf. Enum. 130. 1824 = Paltonium.
linearis Klf. Enum. 131. 1824 = Monogramma seminuda.
linearis Mart. et Gal. Mém. Ac. Brux. **15**. 44. 1842 = Vittaria filifolia.
macrorhynchos Mett. Fil. Lips. 28. 1856 = Hymenolepis platyrhynchos.
marginalis Moore. Ind. XXX. 1857 = Antrophyum callifolium.
microphylla Mett.: Kuhn. Fil. Afr. 58. 1868 = Drymoglossum.
Miyoshiana Mak. Bot. Mag. Tokyo **12**. 26. 1898 = Drymotaenium.
niphoboloides Moore. Ind. XXX. 1857 = Cyclophorus flocciger.
niphoboloides Luerss. Abh. Nat. Ver. Bremen **7**. 49 t. 1 f. 3-5. 1882 = Drymoglossum.
obtusa Hk. Ic. pl. t. 994. 1854 = T. blechnoides var.
ophioglossoides Mett. Fil. Lips. 28. 1856 = Hymenolepis spicata.
piloselloides Mett. Fil. Lips. 28 t. 10 f. 6-8. 1856 (R.Br. Prod. 154. 1810) = Drymoglossum heterophyllum.
piloselloides Wall. List pag. 62. 1829 = Drymoglossum carnosum.
platyrhyncha Mett. Ann. Lugd. Bat. **4**. 173. 1868-69 = Hymenolepis.
pteroides Schkuhr. Kr. Gew. **1**. 21 t. 6 b. 1804 = T. blechnoides.
pumila Klf. Enum. 132. 1824 = Monogramma seminuda.
pusilla Mett. Ann. Lugd. Bat. **4**. 172. 1868-69 = Vittaria.
revoluta Mett. Fil. Lips. 28. 1856 = Hymenolepis spicata var.

sagittaefera Bory, Dup. Voy. Bot. 1. 258 t. 30 f. 2. 1828 = Blechnum auriculatum var.
samarensis Mett. Fil. Lips. 27. 1856 = Polypodium tenuilore.
seminuda Kuhn, Fil. Afr. 59. 1868 = Monogramma.
simplicivenia Ces. Rend. Ac. Napoli 16. 28. 31. 1877 = Vittaria sulcata.
spicata Mett. Ann. Lugd. Bat. 4. 173. 1868-69 = Hymenolepis.
stenophylla Christ, Ann. Jard. Buit. II. 5. 129. 1905 — Borneo.
Swartzii Jenm. JoB. 1879. 263 = Polypodium lanceolatum*.
tricuspidata Spr. Syst. 4. 42. 1827 = Eschatogramme furcata var.
validinervis Mett. Ann. Lugd. Bat. 4. 173. t. 7. 1868-69 = Hymenolepis spicata var.

TAPEINIDIUM (Presl) C. Chr. Index 631. 1906 — (Polypodiaceae Gen. 35).
Microlepia § Tapeinidium Presl, Epim. 96. 1849.

amboynense (Hk.) C. Chr. Ind. 207. 1905; 631. 1906 — Amboyna. Moluccae. Samoa. — Davallia Hk. sp. 1. 178 t. 56 C. 1846. HB. 470: Lindsaya Mett.: Kuhn 1869; Wibelia Kuhn 1882. NPfl. 216. Davallia moluccana Roxb.: Wall. 1829 (non Bl. 1828); Microlepia Pr. 1849.

Denhami (Hk.) C. Chr. Ind. 209. 1905; 631. 1906 — Polynesia. Davallia Hk. 2 Cent. t. 47. 1860. HB. 100; Microlepia Moore 1861: Lindsaya Mett.; Kuhn 1869; Wibelia Kuhn 1882. NPfl. 217.

oligophlebium (Bak.) C. Chr. Ind. 631. 1906 — Borneo. Davallia Bak. JoB. 1888. 323; Wibelia Christ 1905.

pinnatum (Cav.) C. Chr. Ind. 213. 1905; 631. 1906 — Malesia. Polynesia. Davallia Cav. Descr. 277. 1802. HB. 98. Chr. 305; Saccoloma Pr. 1836: Microlepia J. Sm. 1841; Wibelia Fée 1850-52. NPfl. 216; Lindsaya Mett.: Kuhn 1869. Davallia flagellifera Wall. 1828 — D. biserrata Bl. 1828: Microlepia Pr. 1849 — Wibelia Javae Fée 1850-52 — Dicksonia linearis Cav. 1802; Davallia gracilis Bl. 1828; Microlepia J. Sm. 1842; Wibelia Christ 1905; Davallia luzonica Hk. 1846. Wibelia bipinnata Fée 1850-52 — Davallia intramarginalis Ces. 1877 — D. longipinnula Ces. 1877.

TARACHIA Presl. Epim. Bot. 74. 1849 = **Asplenium.**

Adiantum nigrum Pr. l. c. 82 — attenuata Pr. l. c. 75 — caudata Pr. l. c. 79 — Chamissoniana Pr. l. c. 77 — contigua Pr. l. c. 78 — cuneata Pr. l. c. 81 — dimidiata Pr. l. c. 76 — erosa Pr. l. c. 76 — fragrans Pr. l. c. 80 — germanica Pr. l. c. 79 — lacera Pr. l. c. 81 — lanceolata Pr. l. c. 82 — laserpitiifolia Pr. l. c. 83 — lepida Pr. l. c. 81 — macrophylla Pr. l. c. 78 — nitida Pr. l. c. 83 — paradoxa Pr. l. c. 260 — pumila Pr. l. c. 75 — pusilla Pr. l. c. 260 — Ruta-muraria Pr. l. c. 81 — setisecta Pr. l. c. 260 — solida Pr. l. c. 80 — splendens Pr. l. c. 83 — stereophylla Pr. l. c. 80 = A. sp. homonym.

acuta Pr. l. c. 82 = A. adiantum nigrum *onopteris.
angustata Pr. l. c. 260 = A. laserpitiifolium.
arguta Pr. l. c. 82 = A. adiantum nigrum.
Browniana Pr. l. c. 260 = A. praemorsum.
caespitosa Pr. l. c. 81 = A. laciniatum.
canaliculata Pr. l. c. 77 = A. macrophyllum.
caryotoides Pr. l. c. 76 = ? A. dimidiatum.
falcata Pr. l. c. 77 = A. adiantoides.
Finlaysoniana Pr. l. c. 76 = A. macrophyllum.
furcata Pr. l. c. 80 = A. praemorsum.
geminaria Pr. l. c. 79 = A. praemorsum.

Haenkeana Pr. l. c. 76 = A. adiantoides.
insignis Pr. l. c. 260 = A. nitidum.
multicaulis Pr. l. c. 81 = A. ruta muraria.
nigricans Pr. l. c. 79 = A. praemorsum.
obtusa Pr. l. c. 81 = A. adiantum nigrum.
oxyphylla Pr. l. c. 77 = A. macrophyllum.
palmata Pr. l. c. 75 = A. hemionitis.
polymorpha Pr. l. c. 260 = A. abscissum.
polyphylla Pr. l. c. 83 = A. acuminatum.
polyodon Pr. l. c. 76 = A. adiantoides.
polystichoides Pr. l. c. 260 = A. nitidum.
Ruiziana Pr. l. c. 76 = A. abscissum.
similis Pr. l. c. 260 = A. macrophyllum.
trapezoides Pr. l. c. 75 = A. trilobum.
truncata Pr. l. c. 78 = A. laciniatum var.
truncatiloba Pr. l. c. 77 = A. caudatum.
vacillans Pr. l. c. 78 = Diplazium v.
zamiaefolia Pr. l. c. 76 = ? A. dimidiatum.

TASCHNERIA Presl, Epim. Bot. 258. 1849 = **Trichomanes.**
Filicula Pr. l. c. = T. bipunctatum.

TECTARIA Cavanilles, Anales de Historia Natural **1**. 115. 1799; Descripcion d. l. plantas 249. 1802.
aculeata Lag., Garcia et Clem. Anal. Cienc. **5**. 145. 1802 = Polystichum.
Calahuala Cav. Descr. 252. 1802 = Polystichum adiantiforme.
caudata Cav. Anal. Cienc. **4**. 100. 1801 = Athyrium umbrosum.
cinnamomea Cav. Descr. 252. 1802 = ? Polystichum denticulatum.
coriacea Link, Hort. Berol. **2**. 127. 1833 = Polystichum adiantiforme.
crenata Cav. Descr. 250. 1802 = ? Aspidium repandum.
cristata Lag., Garcia et Clem. Anal. Cienc. **5**. 147. 1802 = Dryopteris.
elongata Cav. Anal. Cienc. **4**. 101. 1801 = Dryopteris filix mas*.
falcata Cav. Descr. 250. 1802 = Nephrolepis biserrata.
ferruginea Cav. Descr. 252. 1802 = Polystichum adiantiforme.
filix femina Cav. Anal. Cienc. **4**. 100. 1801 = Athyrium.
Filix mas Cav. Descr. 251. 1802 = Dryopteris.
fragilis Lag., Garcia et Clem. Anal. Cienc. **5**. 147. 1802 = Cystopteris.
fraxinea Cav. Descr. 250. 1802 — Peru.
incisa Cav. Descr. 249. 1802 = Dryopteris serra var.
marginalis Cav. ? = Dryopteris pteroides.
phymatodes Cav. Descr. 249. 1802 = ? Polypodium.
pterioides Lag., Garcia et Clem. Anal. Cienc. **5**. 144. 1802 = Dryopteris thelypteris.
regia Lag., Garcia et Clem. Anal. Cienc. **5**. 148. 1802 = Cystopteris fragilis*.
serrata Cav. Descr. 251. 1802 = Dryopteris gongylodes.
trifoliata Cav. Descr. 249. 1802 = Aspidium.

TEGULARIA Reinw. Sylloge Pl. **2**. 3. 1824 = **Didymochlaena.**
adiantifolia Reinw. l. c. = D. truncatula.

TELEOZOMA R. Brown ap. Richards. in Frankl. Narr. of a journey 767. 1823 = **Ceratopteris.**

TERATOPHYLLUM Mettenius: Kuhn. Annal. Mus. Lugd. Bat. **4**. 296. 1869.
aculeatum Mett.: Kuhn, Ann. Lugd. Bat. **4**. 296. 1869 = Stenochlaena.
— var. inermis Mett. l. c. 297 = Stenochlaena sorbifolia var. 20.
articulatum Mett.: Kuhn. Ann. Lugd. Bat. **4**. 297. 1869 = Polybotrya.

THAMNOPTERIS Presl, Epim. Bot. 68. 1849 = **Asplenium.**
Asplenium § Thamnopteris Pr. Tent. 105. 1836.
australasica Moore, Ind. 115. 1859 = A. nidus var.
Grevillei Moore, Ind. L. 1857.
mauritiana Pr. Epim. 68. 1849 = A. nidus.
musaefolia Pr. Epim. 68. 1849 = A. nidus var.
Nidus Pr. Epim. 68. 1849
orientalis Pr. Epim. 69. 1849 = A. nidus.
pachyphylla Pr. Epim. 69. 1849.
Phyllitidis Pr. Epim. 68. 1849 = A. nidus var.
Simonsiana Moore, Ind. L. 1857.
simplex Pr. Epim. 69. 1849 = A. nidus.
squamulata Pr. Epim. 260. 1849.
stipitata Pr. Epim. 68. 1849 = A. squamulatum.
taeniosa Pr Epim. 69. 1849 = A. amboinense.
THELYPTERIS Adanson, Fam. des plantes **2**. 20. 1763 = **Pteris.**
THELYPTERIS Schott, Gen. Fil. ad tab. 10. 1834 = **Dryopteris.**
palustris Schott, Gen. ad t. 10. 1834 = D. thelypteris.
Themelium Moore, Ind. Fil. LXXI. 1857 *(Polypodium §)* = Polypodium.
THYLACOPTERIS Kunze msc. (Mett. Pol. 50. 1857); J. Sm. Hist. Fil. 87 1875
papillosa Kze.; J. Sm. Hist. Fil. 88. 1875. [= **Polypodium.**
THYRSOPTERIS Kunze, Linnaea **9**. 507. 1834 — (Cyatheaceae Gen. **4**).
elegans Kze. Linn. **9**. 507. 1834. HB. 16. Chr. 331. NPfl. 123 — Juan Fernandez.
Dicksonia Mett. 1861; Panicularia Berteri Colla 1836.
TODEA Willdenow, Schrift. Akad. Erfurt **1802**. 14 — (Osmundaceae Gen. 1).
africana Willd. Schrift. Ak. Erf. **1802**. 14 t. 3 f. 1 = T. barbara.
alpina Bak. Kew Bull. **1899**. 121 = Leptopteris.
arborea ht. (Gard. Chr. III. **19**. 652. 1896) = T. barbara.
barbara (L.) Moore, Ind. CXIX. 1857. HB. 427. Chr. 333. NPfl. 378 — Africa austr. *Austr. temp. Tasmania. N. Zealand. — Acrostichum L. sp. **2**. 1072. 1753; Osmunda Thbg. 1800; O. totta Sw. 1801; Todea africana Willd. 1802 — *T. rivularis Sieb.; Kze. 1837.
Fraseri Hk. et Grev. Ic. Fil. t. 101. 1829 = Leptopteris.
grandipinnula Moore, Gard. Chr. n. s. **25**. 752. 1886 = ? Leptopteris hymenophylloides.
hymenophylloides A. Rich. Fl. Nouv. Zél. 97 t. 16. 1832 = Leptopteris.
intermedia ht. Angl.; André, L'Ill. hort. **24**. 141 t. 290. 1877 — Hort. (e. N. Zealand. — (Leptopteris sp.).
marginata Col. Tr. N. Zeal. Inst. **29**. 419. 1897 = Leptopteris.
Moorei Bak. JoB. **1873**. 16 = Leptopteris.
pellucida Carm.; Grev. et Hk. in Hk. Bot. Misc. **3**. 232. 1833 = Leptopteris hymenophylloides.
rivularis Sieb. Fl. mixt. n. 232; Kze. Anal. 7 t. 4. 1837 = T. barbara*.
superba Col. Tasm. Journ. **2**. 188. 1844; Moore, Gard. Chr. **1861**. 697 = Leptopteris.
Wilkesiana Brack. Expl. Exp. **16**. 309 t. 43. 1854 = Leptopteris.
Tomophyllum Fournier, Ann. sc. nat. V. **18**. 283. 1873 *(Polypodium §)* = Polypodium.
TOXOPTERIS Trevisan, Atti Ist. Veneto V. **3**. 591. 1877 = **Syngramma.**
borneensis Trev. l. c. 592.
cartilagidens Trev. l. c. 592.
Lobbiana Trev. l. c. 592.
?subtrifoliata Trev. l. c. 592 = Stenochlaena sorbifolia var. 17.

TRACHYPTERIS André. (Rev. hort. 1899): Christ, Monogr. Elaph. 150. 1899 — (Polypodiaceae Gen. **108**).

aureo-nitens André; Christ. Mon. Elaph. 150. 1899 = T. pinnata.

pinnata (Hk.fil.) C. Chr. Ind. 5. 1905; 634. 1906 — Galapagos. Ecuador. Brasilia centr. — Hemionitis Hk.fil. Tr. Linn. Soc. **20**. 167. 1847; Acrostichum aureo-nitens Hk. 1854. HB. 421; Neurocallis Moore 1857; Hemionitis J. Sm. 1875; Chrysodium Christ 1897. 47; Trachypteris André; Christ 1899; Elaphoglossum Diels 1899. NPfl. 334 — Acrostichum gillianum Bak. 1882; Elaphoglossum Diels 1899. NPfl. 334.

Triblemma J. Smith. Ferns br. and for. 209. 1866 (*Asplenium §*) = Diplazium.

TRICHIOCARPA (Hook.) J. Smith, Cat. Kew Ferns 7. 1856 = **Aspidium.**
Deparia § Trichiocarpa Hook. JoB. 4. 55. 1852.

Moorei J. Sm. Cat. Kew Ferns 7. 1856.

TRICHIOGRAMME Kuhn. Chaetopterides 325. (5). 1882 = **Syngramma.**

alismaefolia Kuhn, l. c. 327 — Borneensis Kuhn, l. c 326 — cartilagidens Kuhn, l. c. 326 — Lobbiana Kuhn. l. c. 327 — marginata Kuhn, l. c. 326 obtusifolia Kuhn, l. c. 326 — pinnata Kuhn. l. c. 328 — quinata Kuhn. l. c. 328 — vittaeformis Kuhn, l. c. 327 = S. sp. homonym.

lanceolata Kuhn, l. c. 328 = S. Hookeri.

reniformis Kuhn, l. c. 326 = Pterozonium r.

TRICHIPTERIS Presl. Del. Prag. 1. 172 1822 = Trichopteris.

TRICHOCALYMMA Zenker msc. (Kze. Linn. **24**. 251. 1851) = **Polypodium.**

TRICHOMANES Linné. Spec. plant. 2. 1097. 1753 — Hymenophyllaceae Gen. **1**).

Subgenera (Genera in Prantl: Die Hymenophyllaceen).

C = **Cardiomanes.**
F = **Feea.**
G = **Gonocormus.**
H = **Hemiphlebium.**
L = **Lacostea.**
P = **Ptilophyllum.**
T = **Eutrichomanes.**

(T) **abrotanifolium** v.d.B. Ned. kr. Arch. **4**. 367. 1859 — Brasilia.

abruptum Christ. Hedwigia **44**. 361. 1905 = T. Hostmannianum.

(P) **accedens** Pr. Epim. 14. 1849 — Amer. trop. — Ptilophyllum Prantl 1875; Trichomanes crispum Hk. et Grev. Ic. Fil. t. 12. 1827. HB. 82 (pt.). NPfl. 107 (pt.) (non L. 1753).

achilleaefolium Bory; Willd. sp. 5 512 1810 = T. stylosum.

acrocarpon Fée. Cr. vasc. Br. **1**. 188 t. 70 f. 1. 1869 = T. Kraussii.

aculeatum Sw. Prod. 137. 1788 = Odontosoria fumarioides.

aculeatum Poir. Enc. 8. 81. 1808 = Odontosoria.

aculeatum J. Sm. JoB. **3**. 417. 1841 = Hymenophyllum.

acutum Pr. Hym. 16, 42. 1843 = T. pallidum var.

acutum Mak. Bot. Mag. Tokyo **6**. 45. 1892 (nomen); Christ, Bull. Boiss. **4**. 665. 1896 = T. Makinoi.

adianthinum Bory. Bél. Voy. Bot. **2**. 78. 1833 = T. cuspidatum.

adiantoides L. sp. **2**. 1098. 1753 = Asplenium.

(P) **adscendens** Kze. Farnkr. **1**. 189. 1845; Bot. Zeit. 1847. 320 — Martinique.

aerugineum v.d.B. Ned. kr. Arch. **5**[3]. 201. 1863 = T. erosum.

aeruginosum Poir. Enc. **8**. 76. 1808 = Hymenophyllum.
aethiopicum Burm. Fl. Ind. Fl. Cap. Prod. 28. 1768 — Africa austr.
(P) **alatum** Sw. Schrad. Journ. **1800**². 97. HB. 79. Chr. 27. NPfl. 105 — Amer. trop. T. delicatum v.d.B. 1861 — T. lineare v.d.B. 1861 (non Sw.) — T. attenuatum Hk. 1844; Ptilophyllum Prantl 1875 — Trichomanes ptilodes v.d.B. 1859 (= alatum Hk. et Gr. t. 11); Ptilophyllum Prantl 1875.
alatum Schkuhr. Kr. Gew. **1**. 133 t. 135. 1809 = T. bipunctatum.
alatum Hk. in Curtis. Fl. Londin. ed. nov. t. 53. 1821 = T. radicans.
alatum Bory. Dup. Voy. Bot. **1**. 282 t. 38 f. 2. 1828 = T. javanicum.
album Bl. Enum. 226. 1828 = T. pallidum var.
alchemillaefolium Telfair; Wall. List n. 159. 1828 = T. stylosum.
(T) **alternans** Carr. in Seem. Fl. Vit. 343. 1873. HB. 465 — Fiji.
(F) **amazonicum** Christ. Hedvigia **44**. 359. 1905 — Amazonas.
ambiguum Mart.; Kl. Linn. **14**. 287. 1840 (an Sieb.; Pr. Hym. 16. 1843?) = T. pyxidiferum var. diaphanum.
ampliatum Mett. (Sadeb. NPfl. **1**⁴. 105. 1899) = T. digitatum.
anceps Wall. List n. 166. 1828 = T. radicans var.
anceps α Hk. sp. **1**. 135 t. 40 C. 1846 = T. elegans.
anceps β Hk. sp. **1**. 135 t. 40 C. f. 3. 1846 = T. maximum.
Andrewsii Newm. Brit. Ferns ed. II. 14. 1844 = T. radicans.
(T) **angustatum** Carm. Tr. Linn. Soc. **12**. 513. 1818. NPfl. 107 (pt.) — Tristan d'Acunha. — (an adhuc T. tenerum Spr. 1827?).
angustatum J. Sm. JoB. **3**. 417. 1841 = T. Cumingii.
angustissimum Pr. Abh. böhm. Ges. V. **5**. 333. 1848; Epim. 18 t. 8 A. 1849 = T. capillaceum.
(L) **Ankersii** Parker; Hk. et Grev. Ic. Fil. t. 201. 1831. NPfl. 105 — Amer. trop. Lacostea Prantl 1875. Trichomanes brachypus pt. HB. 80. Chr. 30 — T. commutatum Sturm 1859.
Antillarum v.d.B. Ned. kr. Arch. **5**². 164. 1861 = T. radicans var.
(T) **aphlebioides** Christ in C. Chr. Ind. 635. 1906 — N. Guinea. — T. tenuissimum Christ in Schum. et Laut. Fl. deut. Südsee 106. 1901 (non v.d.B. 1861).
(T) **apicilare** Fourn. Ann. sc. nat. V. **18**. 257. 1873 — N. Caledonia.
apiifolium Pr. Hym. 16, 44. 1843 = T. Bauerianum.
apodum Hk. et Grev. Ic. Fil. t. 117. 1829 = T. hymenoides.
(P) **arbuscula** Desv. Prod. 326. 1827. NPfl. 105 — Amer. trop. T. Bancroftii Hk. et Grev. 1831. HB. 78; Ptilophyllum Prantl 1875. Trichomanes coriaceum Kze. 1834.
Armstrongii Bak. Syn. ed. I. 452. 1868 = Hymenophyllum.
Asae Grayi v.d.B. Ned. kr. Arch. **5**². 180. 1861 = T. leptophyllum.
(G) **Asnykii** Racib. Nat. Tijdschr. Ned. Ind. **59**. 238 t. 2 f. 6. 1900 — Java.
asplenioides Sw. Prod. 136. 1788 = Hymenophyllum.
asplenioides Pr. Hym. 15, 37. 1843 = T. javanicum var.
(G) **assimile** Mett.; Kuhn. Linn. **35**. 386. 1868. HB. 464. NPfl. 105 — N. Hebridae.
astylum Klf. in Sieb. Fl. mixt. n. 340 = T. trigonum.
atrovirens Kze. Bot. Zeit. **1847**. 371 = T. javanicum var.
attenuatum Hk. sp. **1**. 122 t. 39 C. 1844 = T. alatum var.
auratum Fée. Cr. vasc. Br. **1**. 186 t. 67 f. 1. 1869 = T. lucens.
aureum v.d.B. Ned. kr. Arch. **5**³. 208. 1863 = T. humile.
(L) **auriculatum** Bl. Enum. 225. 1828. HB. 82. Chr. 29. NPfl. 107 — Japonia. China. Asia trop. — Cephalomanes v.d.B. 1859; Hym. Jav. 34 t. 25;

Lacostea Prantl 1875. Trichomanes Belangeri Bory 1833 — T. dissectum J. Sm. 1841; Cephalomanes v.d B. 1859 — Trichomanes dimidiatum [Pr. 1843.
axillare Poir. Enc 8. 76. 1808 = Hymenophyllum.
(T) **axillare** Sod. Cr. vasc. quit. 11. 1893 — Ecuador.
(P) **badium** Fourn. Bull. Soc. Fr. **15**. 144. 147. 1868 — Ind. occ. Venezuela.
Bancroftii Hk. et Grev. Ic. Fil. t. 204. 1831 = T. arbuscula.
(H) **Barklianum** Bak. Journ. Linn. Soc. **9**. 338 t. 8 f. F. 1866. HB. 74 — Mauritius. Réunion.
Barnardianum Bailey, Queensl. Flora Suppl. III. 89 c. fig. 1890 = T. pyxidiferum.
(T) **Bauerianum** Endl. Prod. Fl. Norf. 17. 1833. NPfl. 107 — Malesia. Polynesia. Austr. — T. meifolium Bl 1828. Hk. et auctt. (non Bory 1810); T. apiifolium Pr. 1843; v.d.B. Hym. Jav. 26 t. 19. HB. 86. Chr. 31 — T. eminens Pr. 1843 — T. myrioplasium Kze. 1847 — Hymenophyllum polyanthos Hk. 1835 (non Sw. 1801); Trichomanes Hk. 1844. T. exaltatum Brack. 1854 — ? Hymenophyllum puellum Ces. 1877.
Beccarianum Ces. Atti Ac. Napoli 7^8. 8 t. 1 f. 2. 1876 = T. Motleyi.
(T) **Beckeri** Krause; Phil. Linn. **33**. 305. 1865. NPfl. 107 — Chile.
Belangeri Bory, Bél. Voy. Bot. **2**. 79 t. 8 f. 1. 1833 = T. auriculatum.
(P) **bicorne** Hk. Ic. pl. t. 982. 1854. HB. 79 — Brasilia. Ecuador. Ptilophyllum Prantl 1875.
bifidum Vent.; Willd. sp. **5**. 511. 1810 = T. rigidum.
bifolium Bl. Enum. 224. 1828 = T. minutum.
(T) **bilabiatum** Nees et Bl. Nova Acta **11**. 123 t. 13 f. 2. 1823 — Malesia. Melanesia. — Didymoglossum v.d.B. 1856; D. laxum v.d.B. 1861; Hym. Jav. 37 t. 27. D. brevipes Pr. 1843; Trichomanes Bak. 1867. HB. 84. NPfl. 107; T. melanorhizon Hk. 1846.
bilingue Hk.; J. Sm. JoB. **3**. 417. 1841 = T. bilabiatum.
(H) **bimarginatum** v.d.B. Ned. kr. Arch. 5^2. 143. 1863 — Formosa. Ind. austr. Ceylon. Malesia-Polynesia. Austr. — Microgonium v.d.B. Hym. Jav. 7. 1861; Ann. sc. nat. IV. **15**. 91. 1861; Hemiphlebium Luerss. 1882 — Trichomanes Neilgherrense Bedd. 1863. HB. 74. T. muscoides Brack. 1854 et auctt. quoad pl. males. (non Sw. 1801).
(T) **bipunctatum** Poir. Enc. 8. 69. 1808. Chr. 32. NPfl. 105 — Japonia. Formosa. Asia et Austr. trop. Polynesia. Afr. occ. et austr. c. ins. ? Mexico. Didymoglossum Fourn. 1873. Hymenophyllum Filicula Bory; Willd. 1810; Didymoglossum Desv. 1827; v.d.B. Hym. Jav. 35 t. 26; Trichomanes Bory 1828. HB. 81; Taschneria Pr. 1849; Hymenophyllum alatum Schkuhr 1809 (non Sw. 1809); Didymoglossum Pr. 1849 — D. decipiens Desv. 1827 — D. anomalum v.d.B. 1863 — D. insigne v.d.B. 1863; Trichomanes Bedd. 1868 — D. Griffithii v.d.B. 1863 — D. euphlebium v.d.B. 1863 — D. racemulosum v.d.B. 1863 — D. dilatatum v.d.B. 1863; Trichomanes Kuhn 1868 — Didymoglossum latealatum v.d.B. 1863; Trichomanes Christ 1896 — Didymoglossum plicatum v.d.B. 1863; Trichomanes Bedd. 1868 — T. capillatum Taschn. 1843; Didymoglossum Pr. 1843; v.d.B. Hym. Jav. 38 t. 28 — Trichomanes latifrons v.d.B. 1863 — T. striatum Mett. NPfl. 105 (an Don 1825?).
birmanicum Bedd. Ferns br. Ind. Suppl. 3 t. 349. 1876 = T. radicans var.
bivalve Forst. Prod. 84. 1786 = Hymenophyllum.
Blumei Hassk. Obs. Bot. Fil. **2**. 4. 1857-58 = T. digitatum var.
Boivini (Bovini) v.d.B. Ned. kr. Arch. **4**. 373. 1859 = T. rigidum.

Bojeri Hk. et Grev. Ic. Fil. t. 155. 1829 = T. cuspidatum.
Borbonicum v.d.B. Ned. kr. Arch. 5^2. 158. 1861 = T. pyxidiferum var.
Boryanum Kze. Farnkr. **1**. 237 t. 97. 1847 = T. javanicum.
Boschianum Sturm; v.d.B. Ned. kr. Arch. 5^2. 160. 1861 = T. radicans.
(F) **botryoides** Klf. Enum. 263. 1824. HB. 72. NPfl. 108 — Guiana. Panama. Peru. — Feea v.d.B. 1859; Ptilophyllum Prantl 1875; Feea nana Bory 1824; Trichomanes Hk. 1844.
(T) **brachyblastos** Mett.; Kuhn, Linn. **35**. 388. 1865. HB. 466. NPfl. 107 — Peru.
brachypus Kze. Linn. **9**. 105. 1834 = T. pedicellatum.
brasiliense Desv. Prod. 328 t. 7 f. 4. 1827 = T. pyxidiferum var.
Braunii v.d.B. Pl. Jungh. **1**. 550. 1856 = T. pallidum.
brevipes Bak. Syn. 84. 1867 = T. bilabiatum.
brevisetum R.Br. in Ait. Hort. Kew. ed. II. **5**. 529. 1813 = T. radicans.
bryoides Goldm. Nova Acta **19** Suppl. 1. 467. 1843 — Brasilia.
(?) **caespitosum** (Gaud.) Hk. sp. **1**. 132 t. 40 B. 1846. HB. 83. Chr. 23. NPfl. 107 — Amer. antarct. Ins. Falkland. — Hymenophyllum Gaud. Ann. sc. nat. **5**. 99. 1825; Leptocionium Pr. 1848; Serpyllopsis v.d.B. 1861. Hymenophyllum densifolium Phil. 1857.
(G) **calvescens** v.d.B. Ned. kr. Arch. 5^3. 199. 1863. HB. 77. NPfl. 105 — N. S. Wales.
campanulatum Roxb.; Wall. List n. 2199. 1829; Roxb. Calc. Journ. **4**. 518. [1844 — Chittagong.
canariense L. sp. **2**. 1099. 1753 = Davallia.
(T) **capillaceum** L. sp. **2**. 1099. 1753. Chr. 30. NPfl. 107 — Amer. trop. Davallia Willd. 1810; Microlepia? Fée 1866. Trichomanes trichoideum Sw. 1801. HB. 85. T. tenellum Hedw. 1799 — T. angustissimum Pr. 1848 — T. Schiedeanum K. Müll. 1854. — (conf. Hier. Engl. Jahrb. **34**. 424).
capillatum Taschner. Dissert. 34 t. 1 f. 2. 1843 = T. bipunctatum var.
(T) **cartilagineum** Vieill. et Pansch.; v.d.B. Ned. kr. Arch. 5^2. 183. 1861 — Polynesia.
caruifolium Roxb. Calc. Journ. **4**. 519. 1844 — Prince of Wales Ins..
(T) **caudatum** Brack. Expl. Exp. **16**. 256 t. 36 f. 5. 1854. HB. 85. NPfl. 107 — N. S. Wales. Queensland. Polynesia. — T. flavofuscum v.d.B. 1861.
cavifolium K. Müll. Bot. Zeit. **1854**. 753 = T. pyxidiferum.
(T) **cellulosum** Kl. Linn. **18**. 531. 1844 — Brasilia. T. filiforme Sturm 1859; T. gemmatum pt. HB. 87.
chaerophylloides Poir. Enc. **8**. 80. 1808 = Davallia.
chinense L. sp. **2**. 1099. 1753 = Odontosoria.
ciliatum Sw. Prod. 136. 1788 = Hymenophyllum.
clavatum Sw. Prod. 137. 1788 = Hymenophyllum polyanthos*.
cochinchinense Poir. Enc. **8**. 83. 1808 = Lygodium scandens.
(T) **cocos** Christ, Bull. Boiss. II. **4**. 943. 1904 — Costa Rica.
cognatum Pr. Hym. 16. 41. 1843 = T. polypodioides var. incisum.
cognatum Ces. Rend. Ac. Napoli **16**. 24. 28. 1877 = T. Motleyi var.
Colensoi Hk.fil. in Hk. Ic. pl. t. 979. 1854. HB. 85. NPfl. 107 — N. Zealand.
collariatum v.d.B. Ned. kr. Arch. **4**. 368. 1859 = T. radicans.
commutatum Sturm. Fl. bras. 1^2. 261. 1859 = T. Ankersii.
compressum Desv. Berl. Mag. **5**. 329. 1811 = T. rigidum.
(G) **concinnum** Mett.; Kuhn, Linn. **35**. 385. 1868. HB. 465. NPfl. 105 — Tahiti.
contiguum Forst. Prod. 84. 1786 = Davallia.
(T) **corcovadense** v.d.B. Ned. kr. Arch. 5^2. 162. 1859 — Brasilia.
coriaceum Kze. Linn. **9**. 105. 1834 = T. arbuscula.
? cormophyllum Klf. Enum. 266. 1824 = Hemitelia capensis fol. adv.

corticola Bedd. Ferns S. Ind. t. 264. 1863-65 = T. digitatum var.
crenatum Gilib. Exerc. Phytol. **2**. 556. 1792 = Asplenium trichomanes.
(P) *crenatum* v.d.B. Ned. kr. Arch. 5^3. 205. 1863 — Afr. occ. trop.
(P) **crinitum** Sw. Prod. 136. 1788. HB. 80. Chr. 28. NPfl. 107 — Ind. occ. Costa Rica-Ecuador. — Ragatelus Pr. 1843 – Trichomanes L'Herminieri [Fée 1866.
crispulum v.d.B. Ned. kr. Arch. 5^3. 200. 1863 = T. erosum.
(P) **crispum** L. sp. **2**. 1097. 1753. HB. 82 (pt.). Chr. 28. NPfl. 107 (pt.) — Amer. trop. Ptilophyllum Prantl 1875 — Trichomanes fastigiatum Sieb.: Pr. 1843 — T. Haenkeanum Pr. 1843.
crispum Hk. et Grev. Ic. Fil. t. 12. 1827 = T. accedens.
crispum HB. 82 quoad pl. Afr. occ. = ? T. crenatum v.d.B.
(P) **cristatum** Klf. Enum. 265. 1824 — Amer. trop. Ptilophyllum Prantl 1875: Trichomanes crispum pt. HB. 82. NPfl. 107 — T. Schomburgkii v.d.B. 1861.
(H) **cultratum** Bak. JoB. **1879**. 293 — Fiji.
(T) **Cumingii** (Pr.) C. Chr. Ind. 3. 1905; 638. 1906 — Ins. Philipp. Ins. Moluccae. Abrodictyum Pr. Hym. 21 t. 7. 1843; Habrodictyon v.d.B. 1861; Hym. Jav. 17 t. 12; Trichomanes Smithii Hk. 1844. HB. 84. NPfl. 107; Habrodictyon v.d.B. 1861.
cuneiforme Forst. Prod. 85. 1786 = Odontosoria chinensis.
cuneiforme Schkuhr. Kr. Gew. **1**. 131. 1809 = T. capillaceum.
(T) **cupressoides** Desv. Prod. 330. 1827 — Ins. Comorae. Madagascar. Ins. Mascar. Ins. Sechellae. Asia trop. — T. obscurum Bl. 1828; v.d.B. Hym. Jav. 23 t. 17; Didymoglossum Hassk. 1857-58. D. longisetum Pr. 1843 – Trichomanes setaceum v.d.B. 1861. T. rigidum pt. auctt. HB. 86.
curvatum J. Sm. JoB. **3**. 417. 1841 = T. javanicum var.
(H) **cuspidatum** Willd. sp. **5**. 499. 1810. HB. 73. Chr. 24. NPfl. 104 — Afr. trop. Madagascar. — Microgonium Pr. 1843; Hemiphlebium Prantl 1875. Trichomanes Bojeri Hk. et Grev. 1829 — T. adianthinum Bory 1833.
(T) **cyrtotheca** Hill. Fl. Haw. 636. 1888. (NPfl. 107) — Ins. Hawaii.
(P) **dactylites** Sod. Cr. vasc. quit. 12. 1893 — Ecuador. T. digitatum Sod. Rec. 4. 1883 (non Sw. 1806).
daucoides Pr. Epim. 12 t. 7. 1849 = T. rigidum.
(T) **davallioides** Gaud. Freyc. Voy. Bot. 378. 1827. (NPfl. 107) — Ins. Hawaii. Fiji. — T. Sandwicense v.d.B. 1861.
(T) **debile** v.d.B. Ned. kr. Arch. 5^2. 154. 1861 — Venezuela.
decurrens Poir. Enc. Suppl. 8. 71. 1808 = Hymenophyllum.
(T) **delicatum** v.d.B. Ned. kr. Arch. 5^2. 145. 1861 — Ecuador. (an T. Poeppigii var.?).
demissum Forst. Prod. 85. 1786 = Hymenophyllum.
(T) **dentatum** v.d.B. Ned. kr. Arch. 5^2. 182. 1861. (NPfl. 107) — Polynesia.
denticulatum Houtt. Nat. Hist. **14**. 212 t. 100 f. 2. 1783 = Davallia.
denticulatum Poir. Enc. 8. 75. 1808 = Hymenophyllum.
denticulatum Bak. Syn. 82. 1867 = Hymenophyllum praetervisum.
depauperatum Bory, Dup. Voy. Bot. **1**. 283. 1828 = ? T. humile.
diaphanum H.B.K. Nov. Gen. et Sp. **1**. 25. 1825 = T. pyxidiferum var.
diaphanum Korth.: Salom. Nomenkl. 367. 1883 (nomen) — Java.
(G) **dichotomum** Kze. Bot. Zeit. **1847**. 302 (nomen); **1848**. 285; v.d.B. Hym. Jav. 22 t. 16. 1861 — Java. Ins. Societatis. — Gonocormus Prantl 1875.
dichotomum Phil. Bot. Zeit. **1856**. 650 = T. Philippianum.

(G) **diffusum** Bl. Enum. 225. 1828 — Java. — Gonocormus v.d.B. Hym. Jav. 9 t. 4. 1861; Sphaerocionium Hassk. 1857-58.

(G) **digitatum** Sw. Syn. 370, 422. 1806; Poir. Enc. 8. 70. 1808. HB. 239 (pt.). Chr. 27. NPfl. 105 — Ins. Mascar. Asia trop. N. S. Wales.
Gonocormus Prantl 1875. Trichomanes lanceum Bory; Willd. 1810 (Ins. Mascar.) — T. flabellatum v.d.B. 1859; Hym. Jav. 19 t. 13 (non Bory 1827): Gonocormus Prantl 1875: Trichomanes Blumei Hassk. 1857-58 (Java) — Hymenophyllum corticola Thwait. 1864; Trichomanes Bedd. 1863-65 (Ceylon. Austr.).

digitatum Bedd. Ferns br. Ind. t. 301. 1869 = T. pyxidiferum.
digitatum Sod. Rec. 4. 1883 = T. dactylites.
dilatatum Forst. Prod. 85. 1786 = Hymenophyllum.
dilatatum Kuhn, Fil. Afr. 33. 1868 = T. bipunctatum var.
dimidiatum Pr. Hym 15, 38. 1843 = T. auriculatum.
dimorphum Mett. (Sadeb. Nat. Pfl. 1^4. 108. 1899) = T. diversifrons.
dissectum J. Sm. JoB. 3. 417. 1841; Hk. sp. 1. 140. 1846 = T. auriculatum.
divaricatum Poir. Enc. 8. 78. 1808 — Persia. — (Hymenophyllum sp.).

(F) **diversifrons** (Bory) Mett.; Sadeb. Nat. Pfl. 1^4. 108. 1899 — Amer. trop.
Hymenostachys Bory. Dict. class. 8. 462. 1825; 17. t. 42 f. 2; Trichomanes elegans Rudge 1805 (non Rich. 1792). HB. 72. Chr. 29; Ptilophyllum Prantl 1875; Feea Boryi v.d.B. 1859.

(T) **Draytonianum** Brack. Expl. Exp. 16. 252 t. 36 f. 3. 1854. (NPfl. 105) —
Dregei v.d.B. Ned. kr. Arch. 4. 372. 1859 = T. rigidum. Ins. Hawaii.
elatum Forst. Prod. 85. 1786 = Davallia denticulata.
elatum Desv. Prod. 327. 1827 — Amer. trop.
elatum v.d.B. Ned. kr. Arch. 5^2. 177. 1861 = T. giganteum.

(T) **elegans** Rich. Act. Soc. Hist. Nat. Paris 1. 114. 1792. NPfl. 108 — Amer. trop.
Adiantopsis v.d.B. 1861: Trichomanes Prieurii Kze. 1837. HB. 86. Chr. 31: (Davalliopsis v.d.B. 1861). Trichomanes anceps α Hk. 1846 — T. Weddellii v.d.B. 1861.

elegans Rudge. Ic. pl. Guian. 24 t. 35. 1805 = T. diversifrons.
elegans Poir. Enc. 8. 79. 1808 = Davallia denticulata.

(T) **elongatum** A. Cunn. in Hk. Comp. Bot. Mag. 2. 368. 1836 — N. Zealand. N. Caledonia. Fiji. — T. rigidum β HB. 86. NPfl. 107.

emarginatum Poir. Enc. 8. 76. 1808 = Hymenophyllum dilatatum*.
emarginatum Pr. Abh. böhm. Ges. V. 5. 332. 1848; Epim. 11 t. 5 f. B. 1849
eminens Pr. Hym. 16. 44. 1843 = T. meifolium. [= T. pyxidiferum var.
Endlicherianum Pr. Abh. böhm. Ges. V. 5. 333. 1848; Epim. 10 t. 5 A. 1849 = T. humile.
epiphyllum Forst. Prod. 85. 1786 = Davallia denticulata var.
erectum Brack. Expl. Exp. 16. 250 t. 56 f. 1. 1854 = T. humile.
ericoides Hedw. in Weber et Mohr. Beitr. 1. 116. 1805 = T. meifolium.

(P) **eriophorum** (Pr.) v.d.B. Ned. kr. Arch. 4. 360. 1859 — Guiana. Brasilia.
Ragatelus Pr. Epim. 18 t. 9. 1849. (= T. crispum var.?).

(H) **erosum** Willd. sp. 5. 501. 1810 — Africa occ. trop.
Microgonium Pr. 1848 — Trichomanes aerugineum v.d.B. 1863 — T. crispulum v.d.B. 1863 — T. muscoides pt. HB. 75. NPfl. 105.

europaeum Sm. in Rees. Cycl. 36. —. 1817 = T. radicans.
exaltatum Brack. Expl. Exp. 16. 259. 1854 = T. Bauerianum var.

(H) **exiguum** (Bedd.) Bak. Syn. 464. 1874. NPfl. 105 — India austr. Ceylon. Hymenophyllum Bedd. Ferns br. Ind. t. 275. 1868.
eximium Kze. Bot. Zeit. **1847**. 350 = T. pyxidiferum var.
(H) **exsectum** Kze. Anal. 47 t. 29 f. 2. 1837. HB. 85. NPfl. 107 — Chile austr. Chiloe. Juan Fernandez.
falcatum Poir. Enc. **8**. 78. 1808 = Davallia (Sw.). (Nephrolepis sp.).
(T) **Fargesii** Christ, Bull. Soc. Fr. **52** Mém. I. 10. 1905 — China.
fastigiatum Sieb. Syn. Fil. n. 144; Pr. Hym. 15. 1843 = T. pellucens.
(T) **ferrugineum** Fourn. Bull. Soc. Fr. **16**. 392. 1869 — N. Caledonia.
filicula Bory. Dup. Voy. Bot. **1**. 283. 1828 = T. bipunctatum.
(T) **filiculoides** Christ in Schum. et Laut. Fl. deut. Südsee 108. 1901 — N. Guinea.
filiforme Sturm, Fl. bras. **1**^{2}. 269 t. 18 f. 4. 1859 = T. cellulosum.
(P) **fimbriatum** Backh. Cat. 12. 1861; Moore, Gard. Chr. **1862**. 44. — Trinidad. Guiana. — T. superbum v.d.B. 1863. HB. 78.
firmulum Pr. Hym. 16, 46. 1843 = T. rigidum.
flabellatum Bory; Desv. Prod. 325. 1827 (nomen); Bory, Dup. Voy. Bot. **1**. 281. 1828 = T. sibthorpioides.
flabellatum v.d.B. Ned. kr. Arch. **4**. 353. 1859 = T. digitatum var.
flabelliforme Hassk. Obs. Bot. Fil. **2**. 5. 1857-58 — Java.
flabellula D'Urv. Mém. Soc. Linn. Paris **4**. 597. (1824?) = T. sibthorpioides.
flaccidum Forst. Prod. 85. 1786 = Dennstaedtia.
flavofuscum v.d.B. Ann. sc. nat. IV. **15**. 88. 1861 = T. caudatum.
floribundum H.B.Willd. sp. **5**. 505. 1810 = T. pinnatum.
floribundum Hk. et Grev. Ic. Fil. t. 9. 1827 = T. pinnatum.
foeniculaceum Bory; Willd. sp. **5**. 511. 1810 = T. parviflorum.
(H) **fontanum** Lindm. Ark. för Bot. **1**. 44 c. figg. 1903 — Brasilia austr.
(H) **formosanum** Yabe, Bot Mag. Tokyo **16**. 45 (pars japonica) 1902; **19**. 31 f. 1-4. 1905 — Ins. Botol Tobago (prope Formosa).
formosum Trev. Atti Ist. Veneto II. **2**. 163. 1851 (nomen) — Jamaica.
fragile Hedw. Gen. et sp. fil. sub t. 17. 1802 = Hymenophyllum hirsutum.
(T) **Frappieri** Cord. Bull. soc. sci. et arts Réunion **1890-91** —; Fl. Réunion 37. 1895 — Réunion.
(H) **Fraseri** Jenm. Gard. Chr. III. **20**. 266. 1896 — Grenada.
Friedrichsthalii Trev. Atti Ist. Veneto II. **2**. 163. 1851 (nomen) — Guatemala.
frondosum Fée. Cr. vasc. Br. **1**. 190 t. 68 f. 1. 1869 = T. rupestre.
(H) **fruticulosum** Jenm. Gard. Chr. III. **15**. 71. 1894 — Guiana.
fucoides Sw. Prod. 136. 1788 = Hymenophyllum.
fulvum Kl.; Sturm, Fl. bras. **1**^{2}. 272 t. 18 f. 6. 1859 = T. tenerum.
fumarioides Poir. Enc. **8**. 82. 1808 = Odontosoria.
furcatum Klf.; Spr. Anleit. ed. II. **2**. 95. 1818 = T. meifolium.
(L) **furcatum** v.d.B. Ned. kr. Arch. **5**^{2}. 168. 1861 — Guiana.
fusco glaucescens Hk.; J. Sm JoB. **3**. 417. 1841 = T. pallidum.
fuscum Bl. Enum. 225. 1828 = Hymenophyllum.
(P) **Galeottii** Fourn. Bull. Soc. Fr. **15**. 144, 147. 1868 — Mexico. Cuba.
(P) **Gardneri** v.d.B. Ned. kr. Arch. **5**^{2}. 149. 1861 — Amer. trop.
(T) **gemmatum** J. Sm. JoB. **3**. 417. 1841 (nomen); Bak. Syn. 87. 1867 (pt.). NPfl. 108. — Malacca. Java. Ins. Philipp. N. Guinea.
gemmatum Hk. Bak. Syn. 87. 1867 et. auctt. quoad pl. brasil. = T. cellulosum.
gibberosum Forst. Prod. 85. 1786 = Asplenium.

(H) **Giesenhagenii** C. Chr. Ind. 641. 1906 — Ins. Johanna (Comorae).
T. microphyllum Gies. Flora **1890**. 439 t. 14 f. 2-3 (non Kuhn 1868).
(T) **giganteum** Bory; Willd. sp. **5**. 514. 1810. HB. 87. NPfl. 107 — Ins. Comor. et Mascar. Malesia. Fiji. — T. elatum v.d.B. 1861 (non Forst. 1786 nec [Desv. 1827).
glaucescens v.d.B. Pl. Jungh. **1**. 551. 1856 = T. pallidum.
glaucofuscum Hk. in Night. Ocean. sketch. app. 131. 1835; sp. **1**. 128 t. 40 A. 1844 = T. pallidum.
(H) **Godmani** Hk.; Bak. Journ. Linn. Soc. **9**. 337 t. 8 f. A. 1866. HB. 74. NPfl. 104 — Guatemala-Panama.
(H) **Goebelianum** Gies. Flora **1892**. Ergänzbd. 179 f. 3. Chr. 25. NPfl. 104 — [Venezuela.
(T) **Goetzii** Hier. Engl. Jahrb. **28**. 339. 1900 — Africa centr.
(H) *Gourlianum* Grev.; J. Sm. Bot. Voy. Herald 240. 1854 — Panama.
(T) **gracile** v.d.B. Ned. kr. Arch. **5**². 157. 1861 — Java.
(L) **guianense** Sturm. Fl. bras. 1². 262. 1859 — Guiana. — Lacostea Prantl 1875.
guineense Afz.; Sw. Schrad. Journ. **1800**². 96. 1801 — Africa occ. trop.
Haenkeanum Pr. Hym. 15. 36, 65. 1843 = T. crispum var.
(T) **Hartii** Bak. Gard. Chr. n. s. **18**. 680. 1882 — Sierra Leone.
Harveyi Carr. in Seem. Fl. Vit. 344. 1873 = T. rigidum var.
(H) **Henzaianum** Parish; Hk. 2 Cent. t. 1. 1860. HB. 75. NPfl. 105 — Moulmein. Pegu. — Microgonium Henzariense v.d.B. 1861.
Henzaiense Bedd. Ferns br. Ind. t. 183. 1866 = T. Motleyi.
heterophyllum Poir. Enc. **8**. 78. 1808 = Humata.
(F) **heterophyllum** H.B.Willd. sp. **5**. 503. 1810. HB. 72. NPfl. 108 — Essiquibo. Brasilia bor. — Homoeotes Pr. 1848; Ptilophyllum Prantl 1875; Feea Humboldtii v.d.B. 1859 — Trichomanes Spruceanum Hk. 1854; Feea v.d.B. 1859; Ptilophyllum Prantl 1875.
hibernicum Spr. Syst. **4**. 130. 1827 = T. radicans.
(H) **Hildebrandtii** Kuhn. v. Deck. Reis. **3**³. Bot. 70. 1879. Chr. 25. NPfl. 104 — Ins. Johanna (Comorae).
hirsutum L. sp. **2**. 1098. 1753 = Hymenophyllum.
hirsutum Thbg. Fl. Jap. 339. 1784 = Microlepia pilosella.
hirsutum Lour. Fl. Cochinch. **2**. 837. 1790 — Cochinchina.
hirsutum Thouars Fl. Trist. d'Ac. 34. 1804 = Hymenophyllum capillare.
hirtellum Poir. Enc. **8**. 76. 1808 = Hymenophyllum.
(T) **hispidulum** Mett.; Kuhn. Linn. **35**. 389. 1868. HB. 466. NPfl. 107 — Borneo. Perak.
hispidum Poir. Enc. **8**. 71. 1808 = Hymenophyllum hirsutum.
(P) **holopterum** Kze. Farnkr. **1**. 185 t. 77 f. 2. 1845 — Amer. trop. Ptilophyllum Prantl 1875.
(H **Hookeri** Pr. Hym. 16. 1843. NPfl. 105 — Amer. trop.
Microgonium Pr. 1848; Hemiphlebium Prantl 1875; Trichomanes muscoides Hk. et Grev. Ic. Fil. t. 179. 1830 (non Sw. 1801); HB. 75 (pt.).
Hookeri Kze. Bot. Zeit. **1847**. 300 = T. sphenoides et T. sublimbatum.
(T) **Hosei** Bak. Journ. Linn. Soc. **22**. 223 t. 12. 1886 — Borneo.
(P) **Hostmannianum** (Kl.) Kze. Bot. Zeit. **1847**. 352 — Guiana. Amazonas. Neurophyllum Kl. Linn. **18**. 532. 1844; Odontomanes Pr. 1849; Ptilophyllum Prantl 1875; Neurophyllum abruptum Fée 1844-45 (nomen); Neuromanes v.d.B. 1859; Trichomanes Christ 1905; T. pinnatum pt. HB. 84. NPfl. 107.
(P) **Huberi** Christ. Bull. Boiss. **6**. 992. 1898 — Amazonas.

(T) **humile** Forst. Prod. 84. 1786. HB. 80. NPfl. 105 — Java. Formosa. N. Guinea. Polynesia. Austr. N. Zealand. — Didymoglossum Pr. 1843; Crepidomanes v.d.B. Hym. Jav. 16 t. 11. 1861 — Trichomanes minutulum Gaud. 1827; Didymoglossum Pr. 1843 — Trichomanes Endlicherianum Pr. 1848. T. erectum Brack. 1854 — T. aureum v.d.B. 1863.

hygrometricum Poir. Enc. 8. 79. 1808 = Hymenophyllum.

(H) **hymenoides** Hedw. Fil. Gen. et Sp. t. 4 f. 3. 1799; Lindm. Ark. för Bot. 1. 12. 1903 — Amer. trop. — T. muscoides Sw. 1801 (non HB. et auctt.): Didymoglossum Desv. 1827; Hemiphlebium Prantl 1875. Trichomanes apodum Hk. et Grev. 1829. HB. 75; Didymoglossum v.d.B. 1859; Hemiphlebium Prantl 1875. Trichomones Schaffneri Schlecht. 1853; Microgonium Fée 1857; Didymoglossum v.d.B. 1859.

hymenophylloides v.d.B. Ned. kr. Arch. 5^3. 209. 1863 = T. pyxidiferum.

(T) **hypnoides** Christ, Bull. Boiss. II. 5. 725. 1905 — Mexico.

(T) **ignobile** Ces. Atti Ac. Napoli 7^8. 9. 1876 — Borneo.

(P) **imbricatum** Sod. Cr. vasc. quit. 13. 1893 — Ecuador.

inaequale Poir. Enc. 8. 74. 1808 = Hymenophyllum.

incisum Thbg. Prod. Fl. Cap. 173. 1800 = Hemitelia capensis fol. adv.

incisum Klf. Enum. 261. 1824 = T. polypodioides.

insigne Bedd. Ferns br. Ind. t. 284. 1868 = T. bipunctatum var.

intermedium Klf.; Pr. Hym. 16. 1824 = T. tenerum.

intermedium v.d.B. Ned. kr. Arch. 5^2. 179. 1861 = T. maximum.

(T) **intramarginale** Hk. et Grev. Ic. Fil. t. 211. 1831. HB. 79. NPfl. 105 — Ceylon. Didymoglossum Pr. 1848; Crepidomanes Pr. 1849.

japonicum Thbg. Fl. Jap. 340. 1784 = Onychium.

japonicum Poir. Enc. 8. 79. 1808 = Microlepia pilosella.

japonicum Franch. et Sav. Enum. pl. Jap. 2. 207, 618. 1879 = T. orientale.

(L) **javanicum** Bl. Enum. 224. 1828. HB. 83. Chr. 29. NPfl. 107 — Liu Kiu. Asia, Polynesia, Austr. trop. — Cephalomanes Pr. 1848; v.d.B. Hym. Jav. 30 t. 22; Lacostea Prantl 1875 — Trichomanes curvatum J. Sm. 1841; Cephalomanes v.d.B. 1859. Trichomanes Zollingeri v.d.B. 1856; Cephalomanes v.d.B. 1859; Hym. Jav. 31 t. 23; Lacostea Prantl 1875 — Cephalomanes atrovirens Pr. 1843; Trichomanes Kze. 1847. T. rhomboideum J. Sm. 1841 (nomen); Cephalomanes v.d.B. 1859; Hym. Jav. 33 t. 24; Lacostea Prantl 1875 — Cephalomanes Singaporianum v.d.B. 1859 (= Trichomanes javanicum Hk. et Gr. t. 240); Lacostea Prantl 1875 — Trichomanes alatum Bory 1828 (non Sw. 1801); Cephalomanes Pr. 1848; Trichomanes Boryanum Kze. 1847; Cephalomanes v.d.B. 1859; Lacostea Prantl 1875 — Cephalomanes Wilkesii v.d.B. 1861 — C. australicum v.d.B. 1861.

(T) **Johnstonense** Bail. Proc. Roy. Soc. Queensl. 1. 14 t. 1. 1884 — Queensland.

(T) **junceum** Christ, Bull. Boiss. II. 4. 944. 1904 — Amer. trop.

(T) **jungermannioides** Fourn. Ann. sc. nat. V. 18. 258. 1873 — N. Caledonia.

(P) **Kalbreyeri** Bak. JoB. 1881. 203 — Columbia.

(H) **Kapplerianum** Sturm, Fl. bras. 1^2. 276. 1859 — Brasilia. Hemiphlebium Prantl 1875.

Kaulfussii Hk. et Grev. Ic. Fil. addend. 1831 = T. trigonum.

Kirkii Hk. Syn. 78. 1867 = T. Lenormandi.

(H) **Kraussii** Hk. et Grev. Ic. Fil. t. 149. 1829. HB. 77. NPfl. 105; Lindm. Ark. för Bot. 1. 24. 1903 — Amer. trop. Florida. — Didymoglossum Pr. 1843;

Hemiphlebium Prantl 1875 — Didymoglossum fructuosum Fée 1866 — Trichomanes acrocarpon Fée 1869.

Kraussii var. crispatum Hk. Bak. Syn. ed. I. 77. 1867 = T. melanopus.

(T) **Krugii** Christ, Engl. Jahrb. **24**. 90. 1897 — Ind. occ.

Kunzeanum Hk. sp. **1**. 127 t. 39 D. 1844 = T. radicans var.

(G) **Kurzii** Bedd. Ferns br. Ind. t. 286. 1868 — Asia trop. Formosa. T. nanum v.d.B. Ned. kr. Arch. **5**[3]. 206. 1863. HB. 77. NPfl. 105 (non [Hk. 1844).

(H) **labiatum** Jenm. Gard. Chr. n. s. **24**. 7. 1885 — Guiana.

labiatum Sadeb. Nat. Pfl. **1**[4]. 105. 1899 = T. Giesenhagenii.

laceratum Desv. Journ. de Bot. appl. **3**. —. 1813 = T. pedicellatum.

laciniatum Roxb. Calc. Journ. **4**. 518. 1844 — Ins. Moluccae.

laetevirens Fée, Cr. vasc. Br. **1**. 191 t. 67 f. 2. 1869 = T. radicans.

laetum v.d.B. Ann. sc. nat. IV. **15**. 90. 1861 = T. pluma vel T. trichophyllum.

(P) **Lambertianum** Hk. sp. **1**. 139 t. 41 B. 1846. HB. 83. Chr. 28. NPfl. 107 — Peru. — Ptilophyllum Prantl 1875.

lanceolatum Poir. Enc. **8**. 83. 1808 — Madagascar. — Davallia Steud. 1824.

lanceum Bory; Willd. sp. **5**. 500. 1810 = T. digitatum.

Langsdorffii v.d.B. Ned. kr. Arch. **5**[2]. 150. 1861 = T. laxum.

lastreoides Pr. Hym. 15. 1843 = T. elegans.

latealatum Christ, Verh. Nat. Ges. Basel **11**. 424. 1896 = T. bipunctatum var.

(T) **latemarginale** Eat. Proc. Amer. Acad. **4**. 111. 1858. HB. 79. NPfl. 105 — Hongkong.

latifrons v.d.B. Ned. kr. Arch. **5**[3]. 209. 1863 = T. bipunctatum var.

(T) **Lauterbachii** Christ in Schum. et Laut. Fl. deut. Südsee 108. 1901 — N. Guinea.

(P) **laxum** Kl. Linn. **18**. 530. 1844 — Amer. trop. — T. Langsdorffii v.d.B. 1861.

Lechleri v.d.B. Ned. kr. Arch. **4**. 363. 1859 = T. pyxidiferum var.

(H) **Lehmannii** Hier. Engl. Jahrb. **34**. 420. 1904 — Columbia.

(H) **Lenormandi** v.d.B. Ned. kr. Arch. **5**. 144. 1861 — Ins. Comorae. T. Kirkii Hk. 1867. HB. 78.

lentum Poir. Enc. **8**. 80. 1808 = Davallia.

Lepervanchii Cord. Bull. soc. sci. et arts Réunion **1890-91**. —. 1891 = T. [parvulum.

Leprieurii Hk. Gard. Ferns t. 11. 1862 = T. elegans.

(T) **leptophyllum** A. Cunn. in Hk. Comp. Bot. Mag. **2**. 368. 1836 — N. Zealand. N. Caledonia. Fiji. — T. Asae Grayi v.d.B. 1861.

leptophyllum v.d.B. Ned. kr. Arch. **4**. 363. 1859 = T. pyxidiferum var.

(?) **levissimum** Fée, 11 mém. 110. 1866 — Jamaica.

L'Herminieri Fée, 11 mém. 107 t. 29 f. 1. 1866 = T. crinitum var.

Lindbergii Mett. (Sadeb. Nat. Pfl. **1**[4]. 105. 1899) = T. polypodioides.

Lindeni Pr. Epim. 11 t. 6. 1849 = T. scandens.

(P) **Lindigii** Fourn. Bull. Soc. Fr. **15**. 144. 147. 1868 — Columbia.

lineare Sw. Prod. 137. 1788 = Hymenophyllum.

lineare v.d.B. Ned. kr. Arch. **5**[2]. 159. 1861 = T. alatum var.

(H) **lineolatum** (v.d.B.) Hk. Syn. 73. 1867; Lindm. Ark. f. Bot. **1**. 54. 1903 — Cuba. Jamaica. Merida. Costa Rica. — Didymoglossum v.d.B. Ned. kr. Arch. **5**[3]. 136. 1863; Hemiphlebium v.d.B.; Salom. 1883. Didymoglossum laceratum Fée 1866.

(T) **liukiuense** Yabe, Bot. Mag. Tokyo **19**. 35. 1905 — Liu Kiu.

(T) **longicollum** v.d.B. Ann. sc. nat. IV. **15**. 90. 1861 — N. Caledonia.

longifolium Desv. Berl. Mag. **5**. 328. 1811 = T. crispum.

longisetum Bory; Willd. sp. **5**. 510. 1810 = T. meifolium.

longisetum Wall. List n. 164. 1828 = T. bipunctatum.
loreum Bory, Bél. Voy. Bot. **2**. 79. 1833 = T. digitatum.
(P) **lucens** Sw. Prod. 136. 1788. HB. 83. Chr. 28. NPfl. 107 — Amer. trop.
T. splendidum v.d.B. 1859; Ptilophyllum Prantl 1875 — Trichomanes auratum Fée 1869; Ptilophyllum Prantl 1875.
lucens Hk. et Grev. Ic. Fil. t. 10. 1827 = T. trigonum.
lucidum Roxb. Calc. Journ. **4**. 519. 1844 = Davallia denticulata.
(T) **Luerssenii** F. Muell.; Luerss. Bot. Centralbl. **9**. 440. 1882 — Aneiteum.
Luschnatianum Pr. Hym. 16, 45. 1843 = T. rupestre.
luzonicum Pr. Hym. 16, 42. 1843 = T. humile.
(G) **Lyallii** (Hk.fil.) Hk. Syn. 77. 1867. NPfl. 105 — N. Zealand. N. Caledonia.
Hymenophyllum Hk.fil. Fl. N. Zeal. **2**. 16. 1854.
(T) **Macgillivrayi** Bak. Ann. Bot. **5**. 195. 1891 — Fiji.
(P) **macilentum** v.d.B. Ned. kr. Arch. $\mathbf{5^2}$. 146. 1861. HB. 78 — Amer. trop.
macroclados Kze. Farnkr. **2**. 72 t. 130. 1850 = T. scandens.
(L) **madagascariense** (v.d.B.) Moore, Ind. 230. 1861 — Madagascar.
Cephalomanes v.d.B. Ned. kr. Arch. **4**. 351. 1859. Trichomanes javanicum HB. 83 quoad pl. madag.
(T) **Makinoi** C. Chr. Ind. 644. 1906 — Japonia. China.
T. acutum Mak. 1892; Christ. Bull. Boiss. **4**. 665. 1896 (non Pr. 1843)
malayanum Roxb. Calc. Journ. **4**. 519. 1844 = ? Odontosoria chinensis.
Malingii Hk. Gard. Ferns t. 64. 1862 = Hymenophyllum.
mandiocanum Raddi, Pl. Bras. **1**. 64 t. 79 f. 2. 1825 = T. rigidum.
(G) **Mannii** Hk. Syn. 75. 1867. NPfl. 105 — Fernando Po.
T. Mannianum Mett.; Kuhn 1868.
Mannianum Mett.; Kuhn, Fil. Afr. 34. 1868 = T. Mannii.
marchantioides Zippel; Moritz, Verz. 107. 1845-46; Hassk. Obs. **2**. 1. 1857-58 = T. sublimbatum.
marginatum Mett. (Sadeb. Nat. Pfl. $\mathbf{1^4}$. 107. 1899) = T. rigidum.
Martiusii Pr. Hym. 15, 36. 1843 = T. pilosum var.
(T) **maximum** Bl. Enum. 228. 1828.; v.d.B. Hym. Jav. 25 t. 18. HB. 86. Chr. 31. NPfl. 108 — Malesia. Polynesia. Queensland. — T. anceps β Hk. 1846 — T. intermedium v.d.B. 1861 (non Pr. 1843). T. Millefolium Pr. 1843 v.d.B. Hym. Jav. 27 t. 20.
maximum Pohl; Kl. Linn. **21**. 240. 1848 = T. Sellowianum.
(T) **meifolium** Bory; Willd. sp. **5**. 509. 1810 — Réunion. Malesia. Polynesia.
T. ericoides Hedw. 1805 (nomen). HB. 87. Chr. 31. NPfl. 108 — T. longisetum Bory; Willd. 1810; v.d.B. Hym. Jav. 28 t. 21. HB. ed. I. 87.
meifolium Bl. Enum. 227. 1828; HB. ed. I. 86 = T. Bauerianum.
(H) **melanopus** Bak. Syn. 465. 1874; Lindm. Ark. f. Bot. **1**. 28. 1903 — Ecuador.
melanorhizon Hk. sp. **1**. 140. 1846 = T. bilabiatum.
(T) **melanotrichum** Schlecht. Adumbr. 56; Linn. **10**. 553. 1836 — Afr. trop. et austr. c. ins. ? Asia trop.
(H) **membranaceum** L. sp. **2**. 1097. 1753. HB. 73. Chr. 25. NPfl. 104 — Amer. trop.
Lecanium Pr. 1843; Hemiphlebium Prantl 1875.
membranaceum Hk. et Arn. Bot. Beech. Voy. 76. 1832 = ? T. parvulum.
(H) **Mettenii** C. Chr. Ind. 644. 1906 — Afr. occ. trop. — T. subsessile Mett.; Kuhn. Fil. Afr. 37. 1868. HB. 465. NPfl. 105 (non Splitgb. 1840).
Mexicanum v.d.B. Ned. kr. Arch. **5**. 164. 1861 = T. radicans.

(?) *Meyenianum* (Pr.) v.d.B. Ned. kr. Arch. **4**. 379. 1859 — Luzon.
Meringium Pr. Hym. 24 t. 8 f. B. 1843.

(P) **micayense** Hier. Engl. Jahrb. **34**. 422. 1904 — Columbia.

(T) **microchilum** Bak. Tr. Linn. Soc. II. Bot. **4**. 250. 1894 — Borneo.

microphyllum Hk.; Kuhn, Linn. **35**. 389. 1868 = Hymenophyllum parvifolium.

microphyllum Gies. Flora **1890**. 439 t. 14 f. 2-3 = T. Giesenhagenii.

millefolium Desv. Prod. 329. 1827 = T. rigidum.

Millefolium Pr. Hym. 16, 43. 1843 = T. maximum.

(T) **Milnei** v.d.B. Ann. sc. nat. IV. **15**. 89. 1861 — N. Caledonia.

minutulum Gaud. Freyc. Voy. Bot. 377 t. 12 f. 2. 1827 = T. humile.

(G) **minutum** Bl. Enum. 223. 1828 — Java. Borneo. N. Guinea.
Gonocormus v.d.B. Hym. Jav. 7 t. 3. 1861 — Trichomanes bifolium et T. parvulum Bl. 1828. T. saxifragoides pt. HB. 75. NPfl. 105.

(T) **Miyakei** Yabe, Bot. Mag. Tokyo **19**. 34. 1905 — Formosa.

molle Fourn., Bull. Soc. Fr. **15**. 147. 1868 = T. crispum.

montanum Salisb. Prod. 404. 1796 = Davallia canariensis.

(H) **montanum** Hk. Ic. pl. t. 187. 1837 — Amer. trop. Natal.
T. quercifolium Hk. et Grev. Ic. Fil. t. 115. 1829; Lindm. Ark. f. Bot. **1**. 20. 1903 (non Desv. 1811); Didymoglossum Pr. 1843 — Trichomanes Robinsoni Bak. 1867. T. pusillum pt. HB. 77.

Moritzii v.d.B. Ned. kr. Arch. **4**. 366. 1859 = T. rupestre.

(H) **Mosenii** Lindm. Ark. f. Bot. **1**. 46 f. 25 D. E, 27. 1903 — Brasilia.

(H) **Motleyi** v.d.B. Ned. kr. Arch. **5**². 145. 1861. HB. 73. Chr. 24. NPfl. 104 — Formosa. Asia trop. Melanesia. Queensland. — Microgonium v.d.B. Hym. Jav. 5 t. 1. 1861; Hemiphlebium Prantl 1875 — Trichomanes Beccarianum Ces. 1876 — T. cognatum Ces. 1877 (non Pr. 1843) — T. pannosum Ces. 1877.

Mougeoti v.d.B. Ned. kr. Arch. **4**. 378. 1859 = T. osmundoides var.

multifidum Forst. Prod. 85. 1786 = Hymenophyllum.

muscoides Sw. Schrad. Journ. **1800**². 95. 1801 = T. hymenoides.

muscoides Hk. et Grev. Ic. Fil. t. 179. 1830 = T. Hookeri.

muscoides Bak. Syn. 75. 1867 = spec. plur.: T. Hookeri (pl. amer.), erosum (pl. afr.). bimarginatum et sublimbatum (pl. asiat.).

(H) **myrioneuron** Lindm. Ark. f. Bot. **1**. 48 f. 25 F-I, 28. 29. 1903 — Guiana. Brasilia. Costa Rica.

myriophyllum Desv. Prod. 329. 1827 = T. parviflorum.

myrioplasium Kze. Bot. Zeit. **1846**. 477 = T. Bauerianum.

nanum Hk. sp. **1**. 115. 1844 = T. botryoides.

nanum v.d.B. Ned. kr. Arch. **5**³. 206. 1863 = T. Kurzii.

(T) **Naseanum** Christ, Bull. Soc. Fr. **52** Mém. I. 11. 1905 — China. Liu Kiu.

(T) **Naumanni** Kuhn et Luerss.: Christ, Engl. Jahrb. **23**. 336. 1896 — Samoa.

Neesii Bl. Enum. 226. 1828 = Hymenophyllum denticulatum.

neilgherrense Bedd. Ferns S. Ind. t. 6. 1863 = T. bimarginatum.

(G) **nitidulum** v.d.B. Pl. Jungh. **1**. 547. 1856; Hym. Jav. 21 t. 15. 1861 — Java.
Gonocormus Prantl 1875.

niveum Burm. Fl. Ind. 238. 1768 — India.

nudum Poir. Enc. **8**. 73. 1808 = Hymenophyllum.

(H) **nummularium** (v.d.B.) C. Chr. Ind. 226. 1905; 645. 1906 — Brasilia septentr.
Didymoglossum v.d.B. Ned. kr. Arch. **5**³. 135. 1863.

(?G) **Nymani** Christ in Schum. et Laut. Nachtr. Fl. deut. Südsee 36. 1905 — [N. Guinea.
obscurum Bl. Enum. 227. 1828 = T. cupressiforme.
olivaceum Kze.; Kl. Linn. **20**. 437. 1847 = T. pyxidiferum.
(H) **omphalodes** (Vieill.) C. Chr. Ind. 646. 1906 — Liu Kiu. Formosa. Asia trop. Polynesia. Austr. trop. — Microgonium Vieill. apud Fourn. Ann. sc. nat. V. **18**. 255. 1873 (syn.); Trichomanes peltatum Bak. Journ. Linn. Soc. **9**. 336 t. 8 f. C. 1866. HB. 73. Chr. 24. NPfl. 104 (non Poir. 1808); Hemiphlebium Luerss. 1882 — ? Trichomanes tahitense Nadeaud 1873.
(T) **opacum** v.d.B. Ned. kr. Arch. **5**[2]. 175. 1861 — Peru.
(H) **orbiculare** Christ, Engl. Jahrb. **19**. Beibl. 26. 1894; Farnkr. 25. 1897 (pl. ster.); in Schwacke, Pl. Nov. Mineiras **2**. 16. 1900 (fert.); Bull. Boiss. II. **2**. 326. 1902. NPfl. 104 — Brasilia.
(T) **orientalis** C. Chr. Ind. 646. 1906 — China. Japonia. Himalaya. T. japonicum Franch. et Sav. Enum. pl. Jap. **2**. 207. 618. 1879 (non Thbg. 1784 nec Poir. 1808).
(P) **ornatulum** v.d.B. Ned. kr. Arch. **5**[2]. 146. 1861 — Brasilia.
(F) **osmundoides** DC.; Poir. Enc. **8**. 65. 1808 — Amer. trop. Hymenostachys Pr. 1843. Trichomanes spicatum Hedw. 1805 (nomen). HB. 72, Chr. 29. NPfl. 108; Feea Pr. 1848; Ptilophyllum Prantl 1875. Feea polypodina Bory 1824. Trichomanes spicisorum Desv. 1811 — T. Mougeoti v.d.B. 1859; Maschalosorus v.d.B. 1861.
(H) **Pabstianum** K. Müll. Bot. Zeit. **1854**. 738 — Brasilia. Didymoglossum v.d.B. 1859.
?pachycarpum Fée, 9 mém. 39. 1857 = Hymenophyllum lineare.
pacificum Hedw. Fil. Gen. et Sp. 1803 = Hymenophyllum bivalve.
(T) **pallidum** Bl. Enum. 225. 1828. HB. 80. Chr. 32. NPfl. 105 — Asia trop. Polynesia. — Pleuromanes Pr. 1849; Craspedoneuron v.d.B. Hym. Jav. 14 t. 8. 1861 — Trichomanes glauco-fuscum Hk. 1835; Craspedoneuron v.d.B. 1861 — Trichomanes album Bl. 1828; Leucomanes Pr. 1849; Craspedoneuron v.d.B. Hym. Jav. 12 t. 7. 1861 — Trichomanes Braunii v.d.B. 1856; Craspedoneuron v.d.B. Hym. Jav. 15 t. 9. 1861 — Trichomanes acutum Pr. 1843; Pleuromanes Pr. 1849.
(G) **palmatifidum** K. Müll. Bot. Zeit. **1854**. 732; v.d.B. Hym. Jav. 20 t. 14 — Java. — Gonocormus Prantl 1875; Trichomanes digitatum β HB. 76.
palmatum Pr. Hym. 16, 39. 1843 = T. proliferum.
pannosum Ces. Rend. Ac. Napoli **16**. 24, 28. 1877 = T. Motleyi var.
papillatum K. Müll. Bot. Zeit. **1854**. 751 = T. rigidum var.
Pappei Kze. Bot. Zeit. **1847**. 438 = T. tamarisciforme.
(T) **parviflorum** Poir. Enc. **8**. 83. 1808. NPfl. 108 — Madagascar. Ins. Mascar. Perak. Singapore. Borneo. Australia. — T. foeniculaceum Bory; Willd. 1810. HB. 88. T. myriophyllum Desv. 1827.
(G) **parvulum** Poir. Enc. **8**. 64. 1808. HB. 75. Chr. 27. NPfl. 105 — Ins. Comorae. Madagascar. Réunion. Japonia. China. Asia trop. Polynesia. N. S. Wales. Queensland. — T. Thouarsianum Pr. 1843.
parvulum Klf. Enum. 206. 1824 = T. sibthorpioides.
parvulum Bl. Enum. 223. 1828 = T. minutum.
pectinatum Poir. Enc. **8**. 78. 1808 = Humata.
(L) **pedicellatum** Desv. Berl. Mag. **5**. 328. 1811. NPfl. 105 — Amer. trop. T. brachypus Kze. 1834. HB. 80 (excl. β). Chr. 30; Lacostea Prantl

1875. T. radicans Hk. et Grev. 1831 (non Sw. 1801) — T. Sonderi v.d.B. 1861 — T. subsessile Splitgb. 1840 — T. volubile Arrab.; v.d.B. 1861.

(P) **pellucens** Kze. Linn. **9**. 104. 1834 — Amer. trop. Ptilophyllum Prantl 1875; T. crispum pt. HB. 82. NPfl. 107.

pellucidum Goldm. Nova Acta **19** Suppl. I. 466. 1843 = T. rupestre.

pellucidum Pr. Hym. 15. 1843 = T. pellucens.

peltatum Poir. Enc. **8**. 76. 1808 = Hymenophyllum.

peltatum Bak. Journ. Linn. Soc. **9**. 336 t. 8 f. C. 1866 = T. omphalodes.

(P) **pennatum** Klf. Enum. 264. 1824 — Amer. trop. — Neurophyllum Pr. 1843; Ptilophyllum Prantl 1875; Neuromanes Kaulfussii v.d.B. 1859. Trichomanes floribundum Hk. et Grev. 1827 (non H.B.Willd. 1810). T. pinnatum pt. HB. 84.

(H) **Petersii** A. Gray. Amer. Journ. Sci. II. **15**. 326. 1853. HB. 74. Chr. 26. NPfl. 105 — Alabama. Georgia. Mississippi. — Microgonium v.d.B. 1861; Hemiphlebium Prantl 1875.

(G?) **Philippianum** Sturm, Enum. pl. cr. vasc. chil. 38. 1858 — Chile. Juan Fernandez. — T. dichotomum Phil. Bot. Zeit. **1856**. 650 (non Kze. 1848).

(P) **pilosum** Raddi, Opusc. sci. Bol. **3**. 296. 1819; Pl. Bras. **1**. 63 t. 79 f. 1. 1825 — Guiana. Brasilia. — Ptilophyllum Prantl 1875 — Trichomanes Martiusii Pr. 1843 (= T. pilosum Mart. Ic. Cr. Br. 104 t. 68. 1834); Ptilophyllum Prantl 1875 — Trichomanes plumula Pr. 1843.

(P) **pinnatifidum** v.d.B. Ned. kr. Arch. **5**³. 202. 1861. HB. 78 — Jamaica.

(H) **pinnatinervium** Jenm. Gard. Chr. n. s. **25**. 787. 1886 — Guiana.

(P) **pinnatum** Hedw. Fil. Gen. et Sp. t. 4 f. 1. 1799. HB. 84 (pt.). Chr. 28. NPfl. 107 — Amer. trop. — Neurophyllum Pr. 1843; Ptilophyllum Prantl 1875; Neuromanes Hedwigii v.d.B. 1859; Neurophyllum Fée 1866. Trichomanes floribundum H.B.Willd. 1810 — ? T. rhizophyllum Cav. 1802 (an = T. pennatum?); Neurophyllum Fée 1866 — N. thecaphyllum Fée 1866 — Neuromanes immersum v.d.B. 1859.

pinnatum pt. Hk. & Bak. Syn. 84 et auctt. = T. pennatum.

pinnatum (a) Vell. Fl. flum. **11** t. 110. 1827; Arch. Mus. Nac. Rio Jan. **5**. 459. 1881 — Brasilia.

platyderon Fourn. Bull. Soc. Fr. **16**. 321. 1869 = T. dentatum.

plicatum Bedd. Ferns br. Ind. t. 285. 1868 = T. bipunctatum var.

(T) **pluma** Hk. Ic. pl. t. 997. 1854. HB. 466. Chr. 31. NPfl. 108 — Perak. Malesia. Samoa. N. Caledonia. — T. elatum v.d.B. 1861 (an = T. trichophyllum?).

(T) **plumosum** Kze. Linn. **9**. 104. 1834 — Peru. Ptilophyllum Prantl 1875. 47. Trichomanes crispum pt. HB. 82.

plumula Pr. Hym. 15, 36. 1843 = T. pilosum var.

(T) **Poeppigii** Pr. Hym. 16. 41. 1843 — Columbia-Peru.

polyanthos Sw. Prod. 137. 1788 = Hymenophyllum.

polyanthos Hk. Ic. pl. t. 703. 1844 = T. Bauerianum var.

(T) **polyodon** Col. Tr. N. Zeal. Inst. **28**. 618. 1896 — N. Zealand.

(P) **polypodioides** L. sp. **2**. 1098. 1753. NPfl. 105 — Amer. trop. T. sinuosum Rich.; Willd. 1810. HB. 78. Chr. 27; Ptilophyllum Prantl 1875 — T. quercifolium Desv. 1811 — P. incisum Klf. 1824 (non Thbg. 1800).

polysperma Poir. Enc. **8**. 82. 1808 = Odontosoria chinensis.

(G) **Powellii** Bak. Syn. 76. 1867 — Samoa.

Prieurii Kze. Anal. 48. 1837 = T. elegans.

(P) **procerum** Fée. 11 mém. 106 t. 28 f. 2. 1866 — Peru. Ind. occ.
T. undulatum v.d.B. 1861 (non Sw. 1788).

(G) **proliferum** Bl. Enum. 224. 1828. HB. 76. Chr. 27. NPfl. 105 — Asia trop. Gonocormus Prantl 1875; Trichomanes palmatum Pr. 1843; Gonocormus v.d.B. Hym. Jav. 11 t. 6. 1861.

ptilodes v.d.B. Ned. kr. Arch. **4**. 357. 1859 = T. alatum var.

Puigari Glaziou (Sadeb. Nat. Pfl. **1**[4]. 105. 1899) = T. polypodioides var.

pulchellum Salisb. Prod. —. 1796 = Hymenophyllum tunbridgense.

(T) **pumilum** v.d.B. Ann. sc. nat. IV. **15**. 89. 1861 — N. Caledonia.

(H) **punctatum** Poir. Enc. 8. 64. 1808. HB. 74. Chr. 26 (pt.). NPfl. 104; Lindm. Ark. f. Bot. **1**. 51. 1903 — Amer. trop. — Didymoglossum Desv. 1827; Hemiphlebium v.d.B. 1861.

(H) **pusillum** Sw. Prod. 136. 1788. (HB. 77 (pt.). Chr. 26 (?); NPfl. 105 (?); Lindm. Ark. f. Bot. **1**. 33. 1903 — Amer. trop. — Didymoglossum Desv. 1827; Hemiphlebium Prantl 1875 (pt.). Didymoglossum angustifrons Fée 1866.

pusillum Bak. Syn. 77. 1867 = T. montanum (et pusillum?).

pusillum Bedd. Ferns br. Ind. t. 302. 1869 = T. viridans.

pyramidale Wall. List n. 162. 1828 = T. cupressoides.

(T) **pyxidiferum** L. sp. **2**. 1098. 1753. HB. 81. Chr. 32. NPfl. 105 — Trop. — T. diaphanum H.B.K. 1815. T. eximium Kze. 1847 — T. brasiliense Desv. 1827. T. emarginatum Pr. 1848 (non Poir. 1808). T. cavifolium K. Müll. 1854 — T. leptophyllum v.d.B. 1859 (non A. Cunn. 1836); T. hymenophylloides v.d.B. 1863 — T. olivaceum Kze.; Kl. 1847 — T. Borbonicum v.d.B. 1861 — T. Schmidianum Zenker; Taschner 1843 — T. Lechleri v.d.B. 1859 — T. tenellum Mart.; v.d.B. 1861 — T. tranninense Fée 1869 — T. Barnardianum Bailey 1890.

pyxidiferum Huds. Fl. Angl. 392. 1762 = Hymenophyllum tunbridgense.

pyxidiferum Schkuhr, Kr. Gew. **1**. t. 134. 1809 = T. capillaceum.

quercifolium Desv. Berl. Mag. **5**. 328. 1811 = T. polypodioides.

quercifolium Hk. et Grev. Ic. Fil. t. 115. 1829 = T. montanum.

(T) **racemulosum** v.d.B. Ned. kr. Arch. **5**[3]. 210. 1863 — Borneo.

(T) **radicans** Sw. Schrad. Journn. **1800**[2]. 97. 1801. HB. 81 (excl. γ). NPfl. 107 — Eur. atlant. Ins. atlant. Afr. trop. occ. Japonia. China. Himalaya. Birma. Alabama-Brasilia. — T. scandens Hedw. 1799 (non L. 1753). T. Kunzeanum Hk. 1844 — T. speciosum Willd. 1810. Chr. 30. T. hibernicum Spr. 1827 – T. Antillarum v.d.B. 1861 — T. Mexicanum v.d.B. 1861 — T. Boschianum Sturm; v.d.B. 1861 — T. repens Schott; Sturm 1859 — T. collariatum v.d.B. 1859 — T. birmanicum Bedd. 1876.

radicans Hk. et Grev. Ic. Fil. t. 218. 1831 = T. pedicellatum.

(C) **reniforme** Forst. Prod. 84. 1786. HB. 73. Chr. 33. NPfl. 105 — N. Zealand. Cardiomanes Pr. 1843.

repens Schott; Sturm, Fl. bras. **1**[2]. 264. 1859 = T. radicans var.

(H) **reptans** Sw. Prod. 136. 1788. HB. 74 (pt.); Lindm. Ark. f. Bot. **1**. 30. 1903 — Amer. trop. — Didymoglossum Pr. 1843; Hemiphlebium Pr. 1848.

reptans Vell. Fl. flum. **11** t. 99. 1827; Arch. Mus. Nac. Rio Jan. **5**. 459. 1881 — Brasilia. — (Hymenophyllum sp.).

reptans Hk. et Grev. Ic. Fil. t. 32. 1828; Hk. Bak. Syn. 74. 1867 (part.?) = T. sphenoides.

rhizophyllum Cav. Descr. 279. 1802 = T. pinnatum (vel pennatum?).

rhomboideum J. Sm. JoB. **3**. 417. 1841 = T. javanicum var.

(T) **rigidum** Sw. Prod. 137. 1788. HB. 86 (pt.). Chr. 30. NPfl. 107 (pt.) — Trop. (an Asia et Polynesia?). — T. bifidum Vent.; Willd. 1810 — T. mandiocanum Raddi 1825 — T. daucoides Pr. 1849 — T. compressum Desv. 1811 — T. Dregei v.d.B. 1859 — T. papillatum K. Müll. 1854 — T. Harveyi et T. Seemanni Carr. 1873.

rigidum Hk. & Bak. Syn. 86. 1867 quoad pl. males. = T. cupressoides.

rigidum Hedw. Fil. Gen. et Sp. 1802 = Hymenophyllum hirsutum.

rigidum Wall. List n. 161. 1828 = T. javanicum.

Robinsoni Hk.; Bak. Journ. Linn. Soc. **9**. 339 t. 8 f. B. 1867 = T. montanum.

(P) **robustum** Fourn. Bull. Soc. Fr. **15**. 147. 1868 — Guadeloupe.

(?) **roraimense** Jenm. Gard. Chr. III. **20**. 716. 1896 — Guiana.

(T) **rupestre** (Raddi) v.d.B. Ned. kr. Arch. **4**. 370. 1859 — Amer. trop. Hymenophyllum Raddi. Pl. Bras. **1**. 67 t. 80. 1825 — Trichomanes venustum Desv. 1827. T. Luschnathianum Pr. 1843; T. radicans γ. HB. 81. NPfl. 107. — T. Moritzii v.d.B. 1859 — T. frondosum Fée 1869.

(H) **rupicolum** Racib. Pterid. Buit. 24. 1898 — Java.

Sandvicense v.d.B. Ned. kr. Arch. 5^2. 165. 1861 = T. davallioides.

sanguinolentum Forst. Prod. 84. 1786 = Hymenophyllum.

(T) **saxatile** Moore, Gard. Chr. **1862**. 45 — Borneo.

(G) **saxifragoides** Pr. Hym. 16. 39. 1843. HB. 75. NPfl. 105 — Java. Ins. Philipp. Melanesia. Fiji. Ins. Amsterdam. — Gonocormus v.d.B. 1861.

(H) **Sayeri** F. Muell. et Bak.; Bak. Ann. Bot. **5**. 195. 1891 — Queensland.

(T) **scandens** L. sp. **2**. 1098. 1753. HB. 85. NPfl. 108 — Mexico. Cuba. Jamaica. T. Lindeni Pr. 1849 — T. macroclados Kze. 1850.

scandens Hedw. Fil. Gen. et Sp. t. 6. 1799 = T. radicans.

Schaffneri Schlecht. Linn. **26**. 368. 1853 = T. hymenoides.

Schiedeanum K. Müll. Bot. Zeit. **1854**. 716 = T. capillaceum.

Schiedeanum Fée, Cr. vasc. Br. **1**. 188. 1869 = T. tenerum.

Schmidianum Zenker; Taschner, Dissert. 34 t. 1 f. 1. 1843 = T. pyxidiferum var.

(P) **Schomburgkianum** Sturm, Fl. bras. 1^2. 249. 1859 — Guiana. Neuromanes v.d.B. 1861; Ptilophyllum Prantl 1875.

Schomburgkii v.d.B. Ned. Kr. Arch. 5^2. 148. 1861 = T. cristatum.

Seemanni Carr. in Seem. Fl. Vit. 345. 1873 = T. rigidum.

(P) **Sellowianum** Pr. Hym. 15, 37. 1843 — Amer. trop. P. maximum Pohl; Kl. 1848 (non Bl. 1828). T. crispum pt. HB. 82.

sericeum Sw. Prod. 136. 1788 = Hymenophyllum.

(T) **serratulum** Bak. Syn. 80. 1867 — Borneo.

serricula Fée, Cr. vasc. Br. **1**. 189 t. 68 f. 3. 1869 = T. polypodioides.

setaceum v.d.B. Ned. kr. Arch. 5^2. 176. 1861 = T. cupressoides.

(H) **setiferum** Bak.; Jenm. JoB. **1881**. 52 — Jamaica.

setigerum Wall. List n. 158. 1828 (nomen) = T. javanicum.

(T) *setigerum* Backh. Cat. 14. 1861; Moore Gard. Chr. **1862**. 45 — Borneo. (an = T. gemmatum?)

setilobum F. Muell. (Hk. & Bak. Syn. 88. 1867) = T. parviflorum.

(T) **siamense** Christ, Bot. Tidsskr. **24**. 103. 1901 — Koh Chang (Siam).

(G) **sibthorpioides** Bory; Willd. sp. **5**. 498. 1810 — Ins. Mascar. — Hymenophyllum Mett.; Kuhn 1868. Trichomanes flabellatum Bory 1828. HB. 76. (non v.d.B. 1859). Hymenophyllum palmatum v.d.B. 1859.

sinuosum Rich.; Willd. sp. **5**. 502. 1810 = T. polypodioides.

Smithii Hk. Ic. pl. t. 704. 1844 = T. Cumingii.

(H) **sociale** (Fée) Lindm. Ark. för Bot. **1**. 17. 1903 — Brasilia.
Didymoglossum Fée. Cr. vasc. Br. **2**. 85 t. 85 f. 3. 1872-73.

solidum Forst. Prod. 86. 1786 = Davallia.

(H) **solitarium** Jenm. Gard. Chr. III. **16**. 592. 1894 — Jamaica. Grenada.

Sonderi v.d.B. Ned. kr. Arch. 5^2. 169. 1861 = T. pedicellatum.

speciosum Willd. sp. **5**. 514. 1810 = T. radicans.

spectabile Kl. Linn. **14**. 287. 1840 = T. pennatum.

(H) **sphenoides** Kze. Linn. **9**. 102. 1834 — Amer. trop. Florida.
Didymoglossum Pr. 1843; Hemiphlebium Prantl 1875. Trichomanes reptans Hk. et Grev. 1828. HB. 74 (pt.?). NPfl. 104 (non Sw. 1788). Didymoglossum Hookeri Pr. 1843; Trichomanes Kze. 1847; Hemiphlebium Pr. 1848. H. punctatum pt. Prantl 1875, Trichomanes Christ 26 (pt.). 1897 (non Poir. 1808).

spicatum Hedw. in Weber et Mohr. Beitr. **1**. 116. 1805 = T. osmundoides.

spicisorum Desv. Berl. Mag. **5**. 329 t. 7 f. 7. 1811 = T. osmundoides.

(T) **spinulosum** Phil. Linn. **30**. 208. 1860 — Chile.

splendidum v.d.B. Ned. kr. Arch. **4**. 360. 1859 = T. lucens.

Spruceanum Hk. Ic. pl. t. 981. 1854 = T. heterophyllum var.

Sprucei Bak. Syn. 87. 1867 = T. subdeltoideum.

squarrosum Forst. Prod. 86. 1786 = Dicksonia.

striatum Don, Prod. Fl. Nepal. 11. 1825 — Nepal.
(an = T. bipunctatum?)

(T) **strictum** Menzies; Hk. et Grev. Ic. Fil. t. 122. 1829. HB. 466 — N. Zealand.

strigosum Thbg. Fl. Jap. 339. 1784 = Microlepia.

(T) **stylosum** Poir. Enc. **8**. 82. 1808 — Madagascar. Ins. Mascar.
Davallia Steud. 1824. Trichomanes achilleaefolium Bory; Willd. 1810. T. thuioides Desv. 1827. T. rigidum pt. HB. 86. NPfl. 107.

(T) **subdeltoideum** C. Chr. Ind. 650. 1906 — Columbia. Brasilia sept.
T. Sprucei Bak. Syn. 87. 1867 (non T. Spruceanum Hk. 1854).

subexsertum v.d B. Ned. kr. Arch. 5^2. 155. 1861 = T. tenerum.

(P) **sublabiatum** v.d.B. Ned. kr. Arch. 5^2. 152. 1861 — Ecuador.

(H) **sublimbatum** K. Müll. Bot. Zeit. **1854**. 737 — Khasia. Java. Borneo.
Microgonium v.d.B. Hym. Jav. 6 t. 2. 1861; Hemiphlebium Prantl 1875. Trichomanes muscoides auctt. quoad pl. asiat. HB. 75. (pt.). T. marchantoides Zippel 1845-46.

(G) **subpinnatifidum** v.d.B. Ned. kr. Arch. 5^2. 141. 1861 — Ceylon.

subsessile Splitg. Tijdschr. Nat. Gesch. **7**. 437. 1840; v.d.B. Ned. Kr. Arch. 5^2. 167. 1861 = T. pedicellatum.

subsessile Mett.; Kuhn. Fil. Afr. 37. 1868 = T. Mettenii.

superbum v.d.B. Ned. kr. Arch. 5^3. 203. 1863 = T. fimbriatum.

tahitense Nadeaud. Enum. pl. Tahiti 18. 1873 = ? T. omphalodes.

(T) **tamarisciforme** Jacq. Coll. **3**. 285 t. 21 f. 2. 1789 — Madagascar. Ins. Mascar. — T. rigidum pt. HB. 86. NPfl. 107.

tamariscinum Bory. Bél. Voy. Bot. **2**. 81. 1833 = T. tamarisciforme.

(L) **tanaicum** Hk.; Sturm. Fl. bras. 1^2. 260. 1859 — Brasilia.
Lacostea Prantl 1875. Trichomanes brachypus β HB. 80.

tenellum Hedw. Fil. Gen. et Sp. t. 3 f. 1. 1799 = T. capillaceum.

tenellum Mart.; v.d.B. Ned. kr. Arch. 5^2. 154. 1861 = T. pyxidiferum.

(T) **tenerum** Spr. Syst. **4**. 129. 1827. HB. 85. Chr. 30 — Amer. trop. — T. fulvum Kl.; Sturm. 1859; T. angustatum NPfl. 107 (an Carm. 1818?)
(T) **tenue** Brack. Expl. Exp. **16**. 251 t. 36 f. 2. 1854. HB. 84 — Tahiti. Samoa.
tenuifolium Burm. Fl. Ind. 237. 1768 = Cheilanthes.
tenuifolium Cav.; Sw. Syn. 144. 1806 = T. uniflorum.
(T) **tenuissimum** v.d.B. Ned. kr. Arch. **5**^{2}. 156. 1861 — Ecuador.
tenuissimum Christ in Schum. et Laut. Fl. deut. Südsee 106. 1901 = T. aphlebioides.
(G) **Teysmanni** v.d.B. Ned. kr. Arch. **5**^{2}. 142. 1861 — Sumatra.
Gonocormus v.d.B. Hym. Jav. 10 t. 5. 1861.
Thouarsianum Pr. Hym. 16, 40. 1843 = T. parvulum.
thuioides Desv. Prod. 328. 1827 = T. stylosum.
(T) **thysanostomum** Mak. Bot. Mag. Tokyo **12**. 193. 1898 — Japonia. Liu Kiu. Formosa.
tranninense Fée. Cr. vasc. Br. **1**. 187 t. 69 f. 1. 1869 = T. pyxidiferum.
translucens Kze. Bot. Zeit. **1847**. 302 = T. humile.
trichodes Sw. Fl. Ind. occ. **3**. 1741. 1806 } = T. capillaceum.
trichoideum Sw. Schrad. Journ. **1800**^{2}. 98. 1801 } = T. capillaceum.
(T) **trichophyllum** Moore, Gard. Chr. **1862**. 45. HB. 466. NPfl. 108 — Borneo. [N. Caledonia.
(an = T. laetum v.d.B. 1861?)
(P) **trigonum** Desv. Berl. Mag. **5**. 328. 1811. NPfl. 105 — Amer. trop.
T. lucens Hk. et Grev. Ic. Fil. t. 10. 1827 (non Sw. 1788); T. Kaulfussii Hk. et Grev. 1831. HB. 79; Ptilophyllum Prantl 1875.
(G) **trinerve** Bak. Fl. Maurit. 465. 1877 — Mauritius.
tunbrigense L. sp. **2**. 1098. 1753 = Hymenophyllum.
(L) **Türckheimii** Christ, Hedwigia **44**. 361. 1905 — Guatemala. Peru.
(T) **Ulei** Christ in Schwacke, Pl. Nov. Mineiras **2**. 16. 1900; Bull. Boiss. II. **2**. 326. 1902 — Brasilia.
umbrosum Wall. List n. 165. 1828 = T. radicans.
undulatum Sw. Prod. 137. 1788 = Hymenophyllum.
undulatum Wall. List n. 160. 1828 = T. cuspidatum.
undulatum v.d.B. Ned. kr. Arch. **5**^{2}. 147. 1861 = T. procerum.
uniflorum Cav. Descr. 280. 1802 — Chiloe.
T. *tenuifolium* Cav.; Sw. 1806 (non Burm. 1768).
(T) **venosum** R.Br. Prod. Fl. N. Holl. 159. 1810. HB. 82. Chr. 32. NPfl. 107 — Austr. Tasmania. N. Zealand.
Phlebiophyllum v.d.B. 1861 — Trichomanes venustulum Col. 1880.
venustulum Col. Tr. N. Zeal. Inst. **12**. 366. 1880 = T. venosum.
venustum Desv. Prod. 328. 1827 = T. rupestre.
(T) **vestitum** Bak. Kew Bull. **1894**. 7 — Borneo.
(T) **Vieillardii** v.d.B. Ann. sc. nat. IV. **15**. 90. 1861 — N. Caledonia.
villosulum Wall. List n. 163. 1828 = ? Salvinia natans.
(T) **virgatulum** v.d.B. Ned. kr. Arch. **5**^{2}. 163. 1861 — Brasilia.
(an T. radicans var.?)
(H) **viridans** Mett.; Kuhn, Linn. **35**. 389. 1868. HB. 465. NPfl. 105 — Moulmein.
T. pusillum Bedd. 1869 (non Sw. 1788).
(H) **vitiense** Bak. Journ. Linn. Soc. **9**. 338 t. 8 f. D. 1866. HB. 74 — Fiji. N. S. Wales. Queensland.
(P) **vittaria** DC.; Poir. Enc. **8**. 65. 1808 — Guiana. Amazonas.
Neurophyllum Pr. 1843; Neuromanes v.d.B. 1859; Ptilophyllum Prantl 1875; Trichomanes pinnatum β HB. 84.

volubile Vell. Fl. flum. **11**. t. 108. 1827: Arch. Mus. Nac. Rio Jan. **5**. 459. 1881; v.d.B. Ned. kr. Arch. 5^2. 170. 1861 = T. pedicellatum.

(H) **Wallii** Thwait.; Ferguson, Ceylon Ferns preface 1880; Trimen, JoB. **1885**. 274; Bak. Ann. Bot. **5**. 194. 1891 — Ceylon.

(T) **Warburgii** Christ in Warb. Monsunia **1**. 55. 1900 — Mindanao.

Weddellii v.d.B. Ned. kr. Arch. 5^2. 174. 1861 = T. elegans var.

(?) **Wildii** Bail. Queensl. Bot. Bull. **4**. 19 t. 5 a. 1891 — Queensland.

Yandinense Bail. Proc. Linn. Soc. N. S. Wales. **5**. 30. 1881 = T. bimarginatum.

Zollingeri v.d.B. Pl. Jungh. **1**. 552. 1856 = T. javanicum.

TRICHOPTERIS Presl (Del. Prag. **1**. 172. 1822 — Trichipteris): Schott, Gen. Fil. tab. 5. 1834 = **Alsophila**.

crenata Pohl: Fée, Cr. vasc. Br. **1**. 175. 1869 = A. elegans var.

denticulata Pr. Tent. 59. 1836 = A. corcovadensis.

elegans Pr. Tent. 59. 1836.

elegans Fée, Cr. vasc. Br. **2**. 83. 1872-73 = A. Feeana.

excelsa Pr. Del. Prag. **1**. 172. 1822 = A. corcovadensis.

TRICHOSORUS Liebmann, Vid. Selsk. Skr. V. **1**. 281 (seors. 129). 1849.

densus Liebm. Vid. Selsk. Skr. V. **1**. 284 (seors. 132). 1849 } = **Alsophila** quadripinnata varr.

frigidus Liebm. Vid. Selsk. Skr. V. **1**. 284 (seors. 132). 1849 } = **Alsophila** quadripinnata varr.

glaucescens Liebm. Vid. Selsk. Skr. V. **1**. 283 (seors. 131). 1849 } = **Alsophila** quadripinnata varr.

Trichostegia J. Smith, Lond. Journ. Bot. **1**. 666. 1842 (*Alsophila* §) = Alsophila.

Trichotemeleum Kunze, Linnaea **24**. 251. 1851 (*Grammitis* §) = Polypodium (mediale).

Trigonophyllum Prantl, Hymenoph. 48. 1875 (*Ptilophyllum* §) = Trichomanes.

TRIPHLEBIA Baker, Malesia **3**. 41. 1886 — (Polypodiaceae Gen. **44**).

dimorphophylla Bak. Malesia **3**. 42 t. 4. 1886 = Stenochlaena sorbifolia var. 19.

linza (Ces.) Bak. Malesia **3**. 42 t. 5. 1886. NPfl. 230 — N. Guinea.
Asplenium Ces. Rend. Ac. Napoli **16**. 26, 29. 1877.

longifolia (Pr.) Bak. Malesia **3**. 41. 1886. NPfl. 230 — Luzon. Moluccae.
Scolopendrium Pr. Rel. Haenk. **1**. 48 t. 9 f. 1. 1825. HB. 247; Micropodium Mett. 1866; Phyllitis O.Ktze. 1891.

pinnata (J. Sm.) Bak. Malesia **3**. 42. 1886. NPfl. 230 — Ins. Philipp.
Scolopendrium J. Sm. 1841 (nomen); Kze. Farnkr. **1**. 124 t. 56. 1843. HB. 247. Chr. 212; Micropodium Mett. 1866; Phyllitis O.Ktze. 1891.

TRISMERIA Fée, Gen. Fil. 164. 1850-52 — (Polypodiaceae Gen. **66**).

argentea Fée, Gen. 165. t. 14 A. f. 1. 1850-52 = T. trifoliata.

aurea Fée, Gen. 165. t. 14 A. f. 2. 1850-52 = T. trifoliata.

longipes (Bak.) Diels, Nat. Pfl. 1^4. 265. 1899 — Paraguay.
Gymnogramme Bak. JoB. **1878**. 301.

microphylla Fée, Gen. 165. 1850-52 = T. trifoliata.

trifoliata (L.) Diels, Nat. Pfl. 1^4. 265. 1899 — Amer. trop.
Acrostichum L. sp. **2**. 1070. 1753; Gymnogramma Desv. 1811. HB. 384. Chr. 69; Ceropteris Kuhn 1896; Trismeria aurea Fée 1850-52.

TROCHOPTERIS Gardner, London Journ. Bot. **1**. 74. 1842 = **Aneimia**.

elegans Gardn. Lond. JoB. **1**. 74 t. 4. 1842.

UGENA Cavanilles, Icon. Descr. Plant. **6**. 73. 1801; Anal. Cienc. **4**. 249. 1801 [= **Lygodium**.

alba Blanco, Fl. Filip. 823. 1837 = ? L. scandens.

dichotoma Cav. Ic. Descr. Pl. **6**. 74 t. 594 f. 3. 1801 = L. circinnatum.

macrostachya Cav. l. c. 75 t. 594 f. 2 = L. circinnatum.

microphylla Cav. l. c. 76 t. 595 f. 2 = L. scandens.

polymorpha Cav. l. c. 75 t. 595 f. 1 = L. venustum.

semihastata Cav. l. c. 74 t. 594 f. 1 = L. flexuosa.
semihastata Blanco, Fl. Filip. 822. 1837 = L. circinnatum.

VAGINULARIA Fée, Dix. sess. d. congr. sci. France 1. 178. 1843; 3 mémoire 50. 1851-52 = **Monogramma**.
angustissima Mett. Ann. Lugd. Bat. 4. 174. 1868-69 } = M. paradoxa.
Junghuhnii Mett. Fil. Lips. 25 t. 27 f. 24-28. 1856 } = M. paradoxa.
paradoxa Mett. Ann. Lugd. Bat. 4. 174. 1868-69.
trichoidea Fée, 3 mém. 54. 1851-52.

VALLIFILIX Thouars, Nov. Gen. Madagascar. 1. —; in Römer, Collect. bot. 195. 1809 = **Lygodium**.

VITTARIA J. Smith, Mém. Acad. Turin 5. 413 t. 9. 1793 — (Polypodiaceae Gen. 89).
acrostichoides Hk. et Grev. Ic. Fil. t. 186. 1830 = Elaphoglossum conforme.
amboinensis Fée, 3 mém. 14 t. 1 f. 1. 1851-52 — Amboyna. Martaban. Taeniopsis Bedd. 1866.
angustifolia Bl. Enum. 199. 1828 = V. elongata. (an sp. propr.?).
angustifolia (Sw.) Bak. Fl. bras. 1^2. 544. 1870. NPfl. 299 — Amer. trop. Mauritius. — Pteris Sw. Prod. 129. 1788; Taenitis Spr. 1827. HB. 396; Pteropsis Desv. 1827. Chr. 55; Oetosis O.Ktze. 1891.
angustifrons Michx. Fl. bor. Am. 2. 261. 1803 = V. lineata.
angustifrons Bory, Voy. 2. 324. 1804 = V. zosterifolia.
anodontolepis Fée, 3 mém. 23 t. 4 f. 3. 1851-52 = V. elongata var.
bisulcata Kze. Bot. Zeit. 1846. 445 = V. zosterifolia.
Bommeri Christ, Bull. Boiss. II. 5. 11. 1905 — Costa Rica.
boninensis Christ in Warb. Monsunia 1. 57. 1900 — Bonin.
caricina Christ, Bull. Ac. Géogr. Bot. Mans 1904. 109 — China.
caudiformis Bl. Enum. 200. 1828 — Java.
Ceterach, Bernh. Schrad. Journ. 1799^1. 315 } α = Ceterach officinarum.
} β = Notholaena Marantae.
coccygocarpa Trev. Atti. Ist. Veneto II. 2. 167. 1851 (nomen) — Afr. austr.
confusa Fourn. Ann. sc. nat. V. 18. 276. 1873 — N. Caledonia.
congoensis Christ, Ann. Mus. Congo Bot. 5. 1. 9 t. 1. 1903 — Congo.
costata Kze. Linn. 9. 77. 1834; Anal. 29 t. 18 f. 2. 1837 — Brasilia. Ecuador. ? Yunnan. — V. Gardneriana Fée 1851-52. HB. 395. NPfl. 299.
crassifolia Bak. Kew Bull. 1898. 212 — Borneo.
curvidentata K. Müll. Bot. Zeit. 1854. 546 = V. lineata.
debilis (Mett.) Kuhn, Linn. 36. 67. 1869. HB. 518 — Borneo. Pleurogramme Mett. msc.
Deppeana K. Müll. Bot. Zeit. 1854. 547 = V. filifolia.
dimorpha K. Müll. Bot. Zeit. 1854. 547 — Mexico.
divergens Roxb.; Wall. List n. 2191. 1829; Roxb. Calc. Journ. 4. 510. 1844 = Schizoloma.
elongata Sw. Syn. 109. 302. 1806. HB. 395 pt. Chr. 53. NPfl. 299 — Asia, Polynesia trop. N. S. Wales. — Oetosis Greene 1900. — Vittaria anodontolepis Fée 1851-52. — V. loricea Fée 1851-52.
ensifolia Bl. Enum. 198. 1828 = V. ensiformis.
ensiformis Sw. Ges. Nat. Fr. Berl. Neu. Schr. 2. 134 t. 7 f. 1. 1799 — Ins. Mascar. Java. — Oetosis Greene 1900. Vittaria incurvata Cav. 1802. V. planta-
falcata Kze. Bot. Zeit. 1848. 198 = V. pusilla. ginea Bory 1804.
filifolia Fée, 3 mém. 20 t. 3 f. 6. 1851-52. NPfl. 299 — Ind. occ. Mexico-Venezuela. — V. Deppeana K. Müll. 1854.

filiformis Cav. Descr. 270. 1802 = V. lineata.
flexuosa Fée, 3 mém. 16. 1851-52 — India.
Forbesii Fée, 3 mém. 14. 1851-52 = V. scolopendrina.
Fudinoi Mak. Bot. Mag. Tokyo **6**. 47. 1892 (nomen); **12**. 28. 1898 — Japonia.
furcata Bak. Fl. bras. 1². 545. 1870 = Eschatogramme.
Gardneriana Fée, 3 mém. 15 t. 3 f. 1. 1851-52 = V. costata.
gracilis Moritz; Kuhn, Linn. **36**. 67. 1869. (HB. 518) — Columbia.
graminifolia Klf. Enum. 192. 1824 — Amer. trop. — Taeniopsis J. Sm. 1841.
Gueinzii Trev. Atti Ist. Veneto II. **2**. 167. 1851 = V. isoetifolia.
guineensis Desv. Berl. Mag. **5**. 325. 1811 — Africa trop. — Pteris Desv. 1827.
hirta Fée, 10 mém. 12 t. 31 f. 4. 1865 — Borneo.
incurvata Cav. Descr. 270. 1802 = V. ensiformis.
intermedia Bl. Enum. 199. 1828 = V. elongata.
interrupta Roxb. Calc. Journ. **4**. 511. 1844 = Lindsaya orbiculata.
intramarginalis Bak.; Jenm. JoB. **1877**. 266 — Jamaica.
isoetifolia Bory, Voy. **2**. 325. 1804. NPfl. 299 — Africa austr. Réunion. Oetosis Greene 1900. Vittaria sarmentosa Ruiz; Fée 1851-52 — V. tenera [Fée 1851-52.
japonica Miq. Ann. Lugd. Bat. **3**. 169. 1867 — Japonia.
Karsteniana Mett. Ann. sc. nat. V. **2**. 207. 1864. (HB. 396) — Columbia.
lanceola Christ, Bull. Boiss. **4**. 674. 1896 — Japonia. China. (= V. japonica?)
lanceolata Sw. Ges. Nat. Fr. Berl. Neu. Schr. **2**. 133 t. 7 f. 2. 1799 = Antrophyum lineatum.
lanceolata Bak. Fl. bras. 1². 544. 1870 = Paltonium.
latipes Fée, 10 mém. 11 t. 31 f. 3. 1865 = V. scolopendrina.
lineata (L.) Sm. Mém. Ac. Turin. **5**. 421 t. 9 f. 5. 1793. HB. 396 (pt.). Chr. 53. NPfl. 299 — Florida. Amer. trop. — Pteris L. sp. **2**. 1073. 1753; Taeniopsis J. Sm. 1841; Haplopteris J. Sm. 1866; Oetosis Greene 1900 — Vittaria filiformis Cav. 1802; Oetosis Greene 1900.
lloydiifolia Racib. Pter. Buit. 59. 1898 — Java. Celebes.
longicoma Christ, Ann. Jard. Buit. II. **5**. 129. 1905 — Borneo.
longidentata K. Müll. Bot. Zeit. **1854**. 546 = V. isoetifolia.
longipes Sod. Cr. vasc. quit. 417. 1893 — Ecuador.
loricea Fée, 3 mém. 14 t. 1 f. 2. 1851-52 = V. elongata.
lunulata Roxb. Calc. Journ. **4**. 510. 1844 — Prince of Wales Ins. (Lindsaya sp.)
maurnensis (Nadeaud) C. Chr. Ind. 654. 1906 — Ins. Societatis. Taeniopsis Nadeaud, Journ. de Bot. **13**. 8. 1899.
Mertensii Trev. Atti Ist. Veneto II. **2**. 167. 1851 (nomen) = Ins. Mariannae.
minor Fée, 3 mém. 23 t. 4 f. 2. 1851-52 = V. pusilla.
minor Bedd. Ferns br. Ind. t. 56. 1865 = V. sikkimensis.
Moritziana Mett. Ann. sc. nat. V. **2**. 207. 1864. (HB. 395) — Columbia.
onusta Trev. Atti Ist. Veneto II. **2**. 167. 1851 = V. costata.
Orbignyana Mett.; Kuhn, Linn. **36**. 66. 1869. HB. 518 — Bolivia.
owariensis Fée, 3 mém. 21. t. 3 f. 2. 1851-52 — Africa occ. trop.
pachydictyon K. Müll. Bot. Zeit. **1854**. 547 = V. lineata.
parasitica Roxb. Calc. Journ. **4**. 510. 1844 = Lindsaya lancea.
parvula Bory, Bél. Voy. Bot. **2**. 35. 1833 — Java.
planipes Kze. Bot. Zeit. **1848**. 197 = V. elongata.
plantaginea Bory, Voy. **2**. 325. 1804 = V. ensiformis.

plantaginea Hk. et Grev. Ic. Fil. t. 187. 1830 = V. rigida.

pumila Mett.; Kuhn, Linn. **36**. 65. 1869. HB. 517 — Borneo. (an = V. hirta?)

pusilla Bl. Enum. 199. 1828 — Ceylon. Java. Malacca. Queensland. Taenitis Mett. 1868-69. Vittaria minor Fée 1851-52 — V. falcata Kze. 1848. HB. 395; Taeniopsis Bedd. 1876.

remota Fée, 7 mém. 26 t. 20 f. 1. 1857. HB. 396. Chr. 54. NPfl. 299 — Ind.-occ.-Ecuador.

resecta Roxb. Calc. Journ. **4**. 510. 1844 — Chittagong. (Lindsaya sp.)

revoluta Willd.; Klf. Enum. 193. 1824 = V. vittarioides.

revoluta Don, Prod. Fl. Nepal. 15. 1825 — Nepal. — Taeniopsis J. Sm. 1840.

rigida Klf. Enum. 193. 1824 — Malesia. Polynesia.

Ruiziana Fée, 3 mém. 16 t. 3 f. 3. 1851-52 = V. stipitata.

sarmentosa Ruiz; Fée, 3 mém. 17. 1851-52 = V. isoetifolia.

scabrida Kl.; Fée, 3 mém. 20. 1851-52. HB. 395 — Mexico-Ecuador. Brasilia.

Schkuhrii Raddi, Pl. Bras. **1**. 51. 1825 = V. lineata.

scolopendrina (Bory) Thwait. Enum. pl. Zeyl. 381. 1864. HB. 396. Chr. 54. NPfl. 299 — Africa austr. c. ins. India. Malesia-Samoa. Pteris Bory, Voy. **2**. 323. 1804; Pteropsis Desv. 1827; Haplopteris Pr. 1836; Taeniopsis J. Sm. 1841 — Taeniopteris Forbesii Hk. 1842; Vittaria Fée 1851-52; Taeniopsis Moore 1857 — Vittaria zeylanica Fée 1851-52; Taeniopsis Moore 1857 — Vittaria latipes Fée 1865.

scolopendrioides Mett. Fil. Lips. 25. 1856 (err.) = V. scolopendrina.

sikkimensis Kuhn, Linn. **36**. 66, 1869. Chr. 53. NPfl. 299 — Himalaya. Birma. — V. minor Bedd. 1865 (non Fée 1851-52).

spartimorpha Trev. Atti Ist. Veneto II. **2**. 167. 1851 = V. filifolia.

stipitata Kze. Linn. **9**. 77. 1834; Anal. 28 t. 18 f. 1. HB. 395 — Cuba. Columbia-Peru. Brasilia. — Taeniopsis Moore 1857 — Vittaria Ruiziana Fée 1851-52.

stricta Carm. Tr. Linn. Soc. **12**. 513. 1818 = V. vittarioides.

suberosa Christ, Bull. Soc. Fr. **52**. Mém. I. 12. 1905 — China.

sulcata (Mett.) Kuhn, Linn. **36**. 68. 1869. HB. 518. Chr. 53 — Ceylon. Perak. N. Guinea. Polynesia. — Pleurogramme Mett. msc. Taenitis simplicivenia Ces. 1877.

tenera Fée, 3 mém. 17 t. 2 f. 1. 1851-52 = V. isoetifolia.

vittarioides (Thouars) C. Chr. Ind. 655. 1907 — Tristan d'Acunha. Pteris Thouars. Fl. Trist. d'Ac. 31 t. 1. 1804 (?); Vittaria stricta Carm. 1818. HB. 396; V. revoluta Willd.; Klf. 1824.

wooroonooran Bail. Rep. Govt. Sci. Exp. Bellenden-Ker 77. 1889; Syn. Queensl. Fl. Suppl. III. 91. 1890 — Queensland.

xiphophylla Trev. Atti Ist. Venoto II. **2** 167. 1851 (nomen) — Ins. Philipp.

zeylanica Fée, 3 mém. 15 t. 1 f. 3. 1851-52 = V. scolopendrina.

zosterifolia Willd. sp. **5**. 406. 1810 — Ins. Mascar. Comorae. Malesia. Polynesia. Oetosis Greene 1900. Vittaria angustifrons Bory, Voy. **2**. 324. 1804 (non Michx. 1803).

WIBELIA Bernhardi, Schrad. Journ. **1800**[2]. 122 t. 1 f. 2. 1801 = **Davallia**.

elata Bernh. l. c. } = D. denticulata.
multifida Bernh. l. c. }

WIBELIA Fée, Gen. Fil. 331. 1850-52 = **Tapeinidium**.

amboynensis Kuhn, Chaetopt. 346. 1882.

bipinnata Fée. Gen. 331. 1850-52 = T. pinnatum var.

Denhami Kuhn, Chaetopt. 346. 1882.

gracilis Christ, Ann. Jard. Buit. II. **5**. 134. 1905 = T. pinnatum var.
Javae Fée, Gen. 331. 1850-52 = T. pinnatum.
oligophlebia Christ, Ann. Jard. Buit. II. **5**. 134. 1905.
pectinalis Fée, Gen. 331. 1850-52 = Lindsaya dubia.
pinnata Fée, Gen. 331. t. 27 bis B. 1850-52.
Wilfordia Christ, Farnkr. 305. 1897 *(Davallia §)* = Microlepia.

WOODSIA R.Brown, Prod. Fl. N. Holl. 158. 1810; Tr. Linn. Soc. **11**. 170. 1815 — (Polypodiaceae Gen. **8**).

alpina (Bolton) Gray. Nat. arr. br. pl. **2**. 17. 1821. NPfl. 161 — Reg. arct.-U. S. bor.-Eur. centr.-Altai. — Acrostichum Bolt. Fil. Brit. 76 t. 42. 1790; Ceterach Lam. et DC. 1805; Acrostichum hyperboreum Liljebl. 1793; Polypodium Sw. 1801; Woodsia R.Br. 1810. HB. 46. Chr. 282; Polypodium arvonicum Wither. 1796. Woodsia hyperborea A arvonica Milde, Fil. Eur. 161. 1867.
alpina Tausch. Flora **1839**². 480 = W. alpina.
Andersoni Christ, Bull. Soc. Fr. **52** Mém. I. 45. 1905 = W. lanosa.
asplenioides Rupr. Distr. Cr. vasc. Ross. 55. 1845 = W. alpina.
Brandtii Franch. et Sav. Enum. pl. Jap. **2**. 205, 616. 1879 — Japonia.
Brownii Mett. Fil. Lips. 98. 1856 = Hypoderris.
Burgessiana Gerr.: Hk. & Bak. Syn. 48. 1866. NPfl. 161 — Natal. Transvaal. Diacalpe Trev. 1875.
canescens Mett. Ann. sc. nat. V. **2**. 249. 1864 = W. crenata.
caucasica J. Sm. JoB. **4**. 191. 1841 = W. fragilis.
crenata (Kze.) Hier. Engl. Jahrb. **34**. 440. 1904 — Andes. — Cheilanthes Kze. Linn. **9**. 84. 1834; Physematium Cumingianum Kze. 1837; Woodsia Hk. 1844 — W. Peruviana Hk. 1844; Diacalpe Trev. 1875. Polypodium flabellatum Kl. 1847. Woodsia canescens Mett. 1864 (non Cheilanthes Kze. 1839); Physematium Trev. 1875; Woodsia obtusa β et γ. HB. 48.
Cumingiana Hk. sp. **1**. 61. 1844 = W. crenata.
Delavayi Christ. Bull. Soc. Fr. **52** Mém. I. 45. 105 — Yunnan.
?dubia Desv. Prod. 319. 1827 = Davallia.
elongata Hk. sp. **1**. 62 t. 21 C. 1844. HB. 47. NPfl. 161 — Himalaya. Yunnan. Physematium Trev. 1875.
europolepis Trevis. Rend. Ist. Lombardo II. **9**. 812. 1876 — Brasilia. Physematium Trevis. Nu. Giorn. bot. It. **7**. 155. 1875. (an = Polypodium lepidopteris?)
fragilis Liebm. Vid. Selsk. Skr. V. **1**. 280 (seors 128). 1849 = W. mollis.
fragilis (Trevir.) Moore, Ind. Cl. 1857. Chr. 283. NPfl. 161 — Caucasus. Dicksonia Trevir. Berl. Mag. **7**. 155. t. 3 f. 18, 19. 1816; Physematium Kze. 1837; Hymenocystis caucasica C. A. Mey. 1831; Woodsia J. Sm. 1841. HB. 47; Cystopteris Fée 1850-52.
frigida Gandoger, Oest. Bot. Zeit. **31**. 18. 1881 = W. ilvensis.
glabella R.Br. ap. Richards. in Franklin, Narr. of a journ. 754. 1823. HB. 47. Chr. 282. NPfl. 161 — Actica-U. S. bor.-Eur. bor. Asia bor. Eur. centr. mont.
Guatemalensis Hk. sp. **1**. 60 t. 21 A. 1844 = W. mollis.
Hancockii Bak. Ann. Bot. **5**. 196. 1891 — China.
hyperborea R.Br. Prod. 158. 1810; Tr. Linn. Soc. **11**. 173 t. 11. 1815 = W. alpina.
ilvensis (L.) R.Br. Prod. 158. 1810; Tr. Linn. Soc. **11**. 173. 1815. HB. 46. Chr. 282. NPfl. 161 — Arctica-U. S. bor.-Eur. mont.-Altai. — Acrostichum L. sp. **2**. 1071. 1753; Polypodium Vill. 1789; Nephrodium rufidulum Michx. 1803; Aspidium Sw. 1806; Notholaena Desv. 1827; Woodsia Beck 1833; W. hyperborea B rufidula Milde, Fil. Eur. 164. 1867.

incisa Gill.; Hk. et Grev. Ic. Fil. t. 191. 1830 = W. montevidensis.
insularis Hance, Ann. sc. nat. IV. **15**. 228. 1861. HB. 47. NPfl. 161 — Sibiria orient. China bor. orient. — (f. Bak. = W. ilvensis et manchuriensis).
japonica Mak. Bot. Mag. Tokyo **18**. 134. 1904 — Japonia.
W. sinuata Mak. Bot Mag. Tokyo **11**. 64. 1897 (non Christ 1902).
laetevirens Prentice, Queensl. Dept. Agric. Bull. **7**. 37. 1891 = Cystopteris fragilis [var.
lanosa Hk. Syn. 47. 1866. NPfl. 161 — Himalaya. China.
Gymnogramme Andersoni Bedd. 1866. HB. 380; Woodsia Christ 1905.
lapponica Ångstr. in Hartm. Handb. ed. IX. 267. 1864 = W. glabella.
macrochlaena Mett.; Kuhn, JoB. **1868**. 270; Linn. **36**. 126. 1869 — China orient.
manchuriensis Hk. 2 Cent. t. 98. 1861. HB. 48. Chr. 283. NPfl. 161 — China bor. Korea. Japonia. — Diacalpe Trev. 1875.
mexicana R.Br. in Wall. Pl. Asiat. rar. **1**. 41. 1830 (nomen) = W. mollis.
mexicana Fée, 7 mém. 66 t. 26 f. 3. 1857 — Mexico. New Mexico. Arizona.
mollis (Klf.) J. Sm. JoB. **4**. 191. 1841. HB. 47. NPfl. 161 — Mexico. Peru.
Physematium Klf. Flora **1829**[1]. 341 — Woodsia Guatemalensis Hk. 1844; Diacalpe Trev. 1875 — Woodsia fragilis Liebm. 1849 (non Moore 1857) — Cheilanthes canescens Kze. 1839; Physematium Trev. 1875.
montevidensis (Spr.) Hier. Engl. Jahrb. **22**. 363. 1896. NPfl. 161 — Peru-Argentina-Brasilia austr. — Dicksonia Spr. Syst. **4**. 122. 1827. Woodsia incisa Gill.; Hk. et Grev. 1830; Physematium Pr. 1836; W. obtusa pt. HB. 48.
obtusa (Spr.) Torrey, Cat. Pl. in Geol. Rep. N. Y. 195. 1840. HB. 48 (excl. varr.). Chr. 282. NPfl. 161 — Amer. bor. (Alaska-Texas-Nova Scotia)-Andes-Peru.
Polypodium Spr. Anleit. **3**. 92. 1804; Aspidium Sw. 1806; Cystopteris Pr. 1836; Physematium Hk. 1840. Alsophila Perriniana Spr. 1821; Woodsia Hk. et Grev. 1828; Cystopteris Link 1833; Physematium Pr. 1836 — Woodsia Plummerae Lemmon 1882.
obtusa var. Lyallii Hk. Syn. ed. I. 48. 1866 = W. scopulina.
oregana Eat. Canad. Naturalist II. **2**. 90. 1865. HB. 48. NPfl. 161 — Br. Columbia-Wisconsin-Arizona — Physematium Trev. 1875.
paleacea Opiz, Böh. phan. u. crypt. Gew. —. 1823 = W. ilvensis.
Perriniana Hk. et Grev. Ic. Fil. t. 68. 1828 = W. obtusa.
Peruviana Hk. sp. **1**. 61 t. 21 B. 1844 = W. crenata.
pilosella Rupr. Distr. Cr. vasc. Ross. 54. 1845 = W. alpina.
Plummerae Lemmon. Bot. Gaz. **7**. 6. 1882 = W. obtusa.
polystichoides Eat. Proc. Amer. Acad. **4**. 110. 1858. HB. 48 (excl. γ). Chr. 283. NPfl. 161 — Japonia. China orient. et bor. — Physematium Trevis. 1875.
polystichoides var. sinuata Hk. Gard. Ferns t. 32 f. 3. 1862 = W. sinuata.
pubescens Spr. Nova Acta **10**. 233 t. 16 f. 5-7. 1821 = ? Dryopteris reptans.
pubescens Opiz, Böh. phan. et crypt. Gew. —. 1833 = W. alpina.
pulchella Bertol. Fl. Ital. Cr. **1**. 111. 1858 = W. glabella.
pusilla Fourn. Bull. Soc. Fr. **27**. 329. 1880 — Mexico.
Raiana Newm. Hist. br. Ferns ed. II. 140. 1844 = W. ilvensis.
Rosthorniana Diels. Engl. Jahrb. **29**. 187. 1900 — China.
rufidula Beck, Bot. North. St. 458. 1833 = W. ilvensis.
scopulina Eat. Canad. Naturalist II. **2**. 91. 1865. HB. 48. NPfl. 161 — Alaska-Arizona-Michigan. — Physematium Trev. 1875.
sinuata (Hk.) Christ. Bull. Boiss. II. **2**. 830. 1902; Mak. Bot. Mag. Tok. **18**. 135. 1904 (non 1897) — Korea. China bor. — W. polystichoides var. sinuata Hk. Gard. Ferns t. 32 f. 3. 1862. HB. 48.

sinuata Mak. Bot. Mag. Tokyo **11**. 64. 1897 = W. japonica.

subcordata Turcz. Bull. Soc. Nat. Moscou **5**. 206. 1832; Rupr. Distr. Cr. vasc. Ross. 52. 1845 = W. alpina.

uralensis Gandoger, Oest. Bot. Zeit. **31**. 18. 1881 = W. ilvensis.

vestita Spr. Nov. prov. hort. 44. 1819 = W. ilvensis.

Yazawai Mak. Bot. Mag. Tokyo **18**. 132. 1904 — Japonia.

WOODWARDIA Smith, Mém. Acad. Turin **5**. 411. 1793 – (Polypodiaceae Gen. **55**).

angustifolia Sm. Mém. Ac. Turin **5**. 411. 1793 = W. areolata.

angustiloba Hance, JoB. **1868**. 176 = W. radicans var.

areolata (L.) Moore, Ind. XLV. 1857. HB. 189. Chr. 187. NPfl. 253 — U. S. atlant.-Michigan-Louisiana. — Acrostichum L. sp. **2**. 1069. 1753; Lorinseria Pr. 1849; Woodwardia angustifolia Sm. 1793; W. onocleoides Willd. 1810.

aspera Fée, Gen. 207. 1850-52 = Doodia.

auriculata Bl. Enum. 196. 1828 = W. radicans var.

Bannisteriana Michx. Fl. bor. Am. **2**. 263. 1803 = W. virginica.

biserrata Pr. Rel. Haenk. **1**. 53. 1825 = W. radicans.

blechnoides Fée, Gen. 207. 1850-52 = Doodia maxima.

caudata Cav. Descr. 264. 1802 = Doodia.

Chamissoi Brack. Expl. Exp. **16**. 138. 1854 = W. spinulosa.

connexa Fée, Gen. 207. 1850-52 = Doodia media var.

cyatheoides Mett. Fil. Lips. 63. 1856 = Sadleria.

Desvauxii Moore, Ind. 342. 1862 = Doodia blechnoides.

dispar Willd. sp. **5**. 419. 1810. (Plum. t. 16) — Martinique.

dives Fée, Gen. 207. 1850-52 = Doodia.

duriuscula Moore, Gard. Chr. **1868**. 1114 (syn.) = Doodia media.

floridana Schkuhr, Kr. Gew. **1**. 103 t. 111. 1809 = W. areolata.

Harlandii Hk. Fil. Exot. t. 7. 1857. HB. 189. NPfl. 253 — Hongkong. Lorinseria J. Sm. 1875.

intermedia Christ, Bull. Boiss. II. **4**. 618. 1904 — Japonia. W. japonica × radicans.

japonica (L.fil.) Sm. Mém. Ac. Turin **5**. 411. 1793. HB. 188. NPfl. 253 — Japonia. China orient. — Blechnum L.fil. Suppl. 445. 1781; Woodwardia virginica pt. Chr. 186.

Kunthiana Fée, Gen. 207. 1850 52 = Doodia media var.

lunulata Fée, Gen. 207. 1850-52 = Doodia media.

media Fée, Gen. 207. 1850-52 = Doodia.

onocleoides Willd. sp. **5**. 416. 1810 = W. areolata.

orientalis Sw. Schrad. Journ. **1800**². 76. 1801 = W. radicans var.

prolifera Hk. et Arn. Bot. Beech. Voy. 275 t. 56. 1836-40 = W. radicans var.

radicans (L.) Sm. Mém. Ac. Turin **5**. 412. 1793. HB. 188. Chr. 186. NPfl. 253 — Mediterr. occ. Ins Atlant. China austr. Ind. bor. Java. — var. Japonia. Formosa. China orient. — Blechnum L. Mant. 307. 1771 — Woodwardia auriculata Bl. 1828; W. angustiloba Hance 1868 — W. orientalis Sw. 1801. HB. 188. W. prolifera Hk. et Arn. 1836-40.

radicans var. americana Hk. sp. **3**. 67. 1860 = W. spinulosa.

spinulosa Mart. et Gal. Mém. Ac. Brux. **15**. 64. 1842 — Washington-Guatemala. W. Chamissoi Brack. 1854; W. radicans auctt. quoad pl. amer.

Stans Cav. Hort. Matrit. I. Tab. —.; Sw. Syn. 117. 1806 = W. radicans.

thelypteroides Pursh. Fl. Am. sept. **2**. 670. 1814 = W. virginica.

virginica (L.) Sm. Mém. Ac. Brux. **5**. 412. 1793. HB. 188. Chr. 186 (pt.). NPfl. 253 — U. S. atlant. Bermuda. — Blechnum L. Mant. 307. 1771; Doodia Pr. 1836; Anchistea Pr. 1849.

XIPHOPTERIS Kaulfuss, Jahrb. f. d. Pharmacie, Berlin **1820**. 35; Enum. Fil. [85. 1824 = **Polypodium.**

cucullata Spr. Syst. **4**. 43. 1827.

extensa Fée, 11 mém. 14 t. 19 f. 3. 1866 = P. serrulatum.

heterophylla Spr. Syst. **4**. 44. 1827 = P. grammitidis.

Jamesoni Hk. 2 Cent. t. 14. 1860 = P. strictissimum.

myosuroides Klf. Enum. 85, 275. 1824.

orientalis Fourn. Compt. rendus **81**. 1140. 1875 = P. serrulatum.

serrulata Klf. Enum. 85. 1824.

setosa Klf. Enum. 275. 1824 = P. micropteris.

ZALUSIANSKYA Necker, Acta Theod. palat. phys. **3**. 303. 1775 = **Marsilea.**

Obs. Omnes species Marsileae a cl. O.KUNTZE (Rev. Gen. Pl. **2**. 823. 1891) ad Zalusianskyam translatae sunt. Vide sub Marsilea.

gymnocarpa O.Ktze. Rev. Gen. Pl. **2**. 823. 1891 = M. pygmaea.

macropus O.Ktze. l. c. = M. macropoda.

marsilioides Neck. Acta Theod. pal. phys. **3**. 303. 1775 = M. quadrifolia.

senegalensis O.Ktze. Rev. Gen. Pl. **2**. 823. 1891 = M. crenulata var.

ADDENDA

NOMINA ANNO 1905 PUBLICATA ET NOMINA IN INDICE OMISSA

ACROPHORUS Macraeanus Carr. in Seem. Fl. Vit. 336. 1873 = Lindsaya repens.

ACROSTICHUM

actinolepis Sod. Sert. Fl. Ecuad. 11. 1905 = Elaphoglossum.
alcicorne auctt. quoad pl. austral. = Platycerium bifurcatum.
Angamarcanum Sod. Sert. Fl. Ecuad. 5. 1905 = Elaphoglossum.
Christii Sod. Sert. Fl. Ecuad. 4 t. 1. 1905 = Elaphoglossum.
cladotrichium Sod. Sert. Fl. Ecuad. 9 1905 = Elaphoglossum.
excelsum Maxon, Proc. Biol. Soc. Wash. **18**. 224. 1905 — Florida-India occ.-Guiana. Mexico-Guatemala. Brasilia. — Chrysodium lomarioides Jenm. Timehri **4**. 314. 1885; Acrostichum Jenm. Bull. Dep. Jam. **5**. 154. 1898 (non Bory 1833).
Hieronymi Sod. Sert. Fl. Ecuad. 11. 1905 = Elaphoglossum.
intermedium Mett; Rosenstock, Hedwigia **43**. 232. 1904 (nomen) — Brasilia austr. (Elaphoglossum sp. aff. E. stricto).
Labrusca Christ, Ann. Mus. Congo Bot. 5. **1**. 10. 1903 = Leptochilus.
Litanum Sod. Sert. Fl. Ecuad. 10. 1905 = Elaphoglossum.
longissimum Sod. Sert. Fl. Ecuad. 3. 1905 = Elaphoglossum.
Oleandropsis Sod. Sert. Fl. Ecuad. 7. 1905 = Elaphoglossum.
Pangoanum Sod. Sert. Fl. Ecuad. 6. 1905 = Elaphoglossum.
pellucidum Sod. Sert. Fl. Ecuad. 7. 1905 = Elaphoglossum.
pteropodum Sod. Sert. Fl. Ecuad. 8. 1905 = Elaphoglossum.
subsessile Sod. Sert. Fl. Ecuad. 6. 1905 = Elaphoglossum oblanceolatum.
trichophorum Sod. Sert. Fl. Ecuad. 10. 1905 = Elaphoglossum.

ADENODERRIS J. Smith. Hist. Fil. 222. 1875 — (Polypodiaceae Gen. **16 a**).
glandulosa J. Sm. Hist. Fil 223. 1875 = A. viscidula.
sororia Maxon, Bot. Gaz. **39**. 368 f. 2. 1905 — Guatemala.
viscidula (Mett.) Maxon, Bot. Gaz. **39**. 367 f. 1. 1905 — Cuba. Jamaica. Aspidium Mett. Pheg. u. Asp. 38 n. 85. 1858; Dryopteris O.Ktze. 1891; Polystichum C. Chr. 1905; Aspidium glandulosum Hk. et Grev. Ic. Fil. t. 140. 1829. HB. 249 (non Bl. 1828); Polystichum Pr. 1836. NPfl. 189.

ADIANTUM

alatum Copeland in Perkins, Fragm. 192. 1905 = A. Mettenii.

blumenavense Rosenst. Festschr. Alb. v. Bamberg 57. 1905 — Brasilia austr.

conglomeratum ht.; Gard. Mag. **1903**. 269 — Hort.

Croweanum ht. Crowe; Gard. Mag. **1904**. 211 — Hort.

Delavayi Christ, Bull. Soc. Fr. **52** Mém. I. 63. 1905 — China.

edentulum Christ, Bull. Soc. Fr. **52** Mém. I. 63. 1905 — China.

fimbriatum Christ, Bull. Soc. Fr. **52** Mém. I. 62. 1905 — China.

lancea Blanco, Fl. Filip. 833. 1837 = ? Schizoloma ensifolium.

Martinii Pirotta, Ann. Ist. bot. Roma **8**. 15 t. 12 A. 1903 — Eritrea. Abyssinia.

monochlamys var. latedeltoidea Christ, Nu. Giorn. bot. It. n. s. **4**. 88. 1897 = A. Davidi.

ALCICORNIUM andinum, angolense Und. Bull. Torr. Cl. **32**. 593. 1905 — bifurcatum, coronarium, Ellisii, grande Und. l. c. 594 — Hillii, madagascariense, Stemaria Und. l. c. 595 — sumbawense, Veitchii, Wallichii, Willinckii Und. l. c. 596 = Platycerium sp. homonym.

ALLANTODIA marginata Racib. Bull. int. Ac. Cracovie **1905**. 349 = Diplazium m.

ALSOPHILA

congoensis ht.; Garden **63**. 288. 1903; Le Jardin **1903**. 133 — Congo. (= A. Sanderi ht.?).

cyclodonta Christ, Ann. Jard. Buit. II. **5**. 137. 1905 — Borneo.

lepidoclada Christ in Schum. et Laut. Nachtr. Fl. deutsch. Südsee 37. 1905 — N. Guinea.

Margarethae Schroeter; Christ, Ann. Jard. Buit. II. **5**. 136. 1905 — Borneo.

Munchii Christ, Bull. Boiss. II. **5**. 734. 1905 — Mexico.

Sanderi ht. Sander; Gard. Chr. III. **33**. 259 f. 103. 1903 — Hort. (= A. congoensis ht.?).

Ulei Christ, Hedwigia **44**. 367. 1905 — Amazonas.

ANEIMIA

Brandegeea Dav. Fern Bull. **13**. 20 c. tab. 1905 — Mexico.

flexuosa × phyllitidis Rosenst. Festschr. Alb. v. Bamberg 66 t. 2. 1905 — Brasilia austr. — A. Ulbrichtii Rosenst. l. c.

Ulbrichtii Rosenst. l. c. = A. flexuosa × phyllitidis.

ARTHROPTERIS

glabra Copeland in Perkins, Fragm. 178. 1905 — Ins. Philipp.

ASPIDIUM

(Ar) **Bryanti** Copeland in Perkins, Fragm. 175. 1905 — Ins. Philipp.

Chiapasense Christ, Bull. Boiss. II. **5**. 727. 1905 = Dryopteris.

(S) **Copelandii** C. Chr. Ind. 661. 1906 — Ins. Philipp. — A. heterodon Copeland in Perkins, Fragm. 177. 1905 (non Schrad. 1824 nec Bl. 1828).

Cunninghamii Col. Tasm. Journ. **1**. 376. 1841 = Polystichum adiantiforme.

elegans Sturm, Enum. pl. vasc. cr. chil. 32. 1858 = Polystichum mohrioides.

eumundi Bail. Queensl. Bot. Bull. **5**. 27. 1892 = Arthropteris obliterata var.

Hallieri Christ, Ann. Jard. Buit. II. **5**. 106. 1905 = Dryopteris.

heterodon Copeland in Perkins, Fragm. 177. 1905 = A. Copelandii.

incanum Christ, Hedwigia **44**. 367. 1905 = Dryopteris.

(Ar) **lamaoense** Copeland in Perkins, Fragm. 176. 1905 — Luzon.

lobatiforme Waisb. Oest. Bot. Zeit. **49**. 65. 1899 = Polystichum Braunii × lobatum.

Motleyanum Christ, Ann. Jard. Buit. II. **5**. 105. 1905 = Dryopteris.

(S) **persoriferum** Copeland in Perkins, Fragm. 177. 1905 — Mindanao.

(S) **pteropodum** (Bak.) Diels, Nat. Pfl. 1^4. 183. 1899 — Borneo.
Nephrodium Bak. JoB. **1888**. 325; Sagenia Christ 1905.

sphaerocarpum Christ, Bull. Boiss. II. **5**. 726. 1905 = Dryopteris athyrioides.

tablanum Christ, Bull. Boiss. II. **5**. 727. 1905 = Dryopteris.

tosaense Mak. Bot. Mag. Tokyo **6**. 46. 1892 (nomen) — Japonia.

(Ar) **Whitfordi** Copeland in Perkins, Fragm. 176. 1905 — Luzon.

Yoshinagae Mak. Bot. Mag. Tokyo **6**. 46. 1892 (nomen) — Japonia.

ASPLENIUM

Andrewsii Nelson, Proc. Biol. Soc. Wash. **17**. 174. 1904 — Colorado.

aneitense Carr. in Seem. Fl. Vit. 353. 1873 — Aneiteum.

angustatum × mucronatum Rosenst. Festschr. Alb. v. Bamberg 61 t. 1. 1905 = A. Martianum × mucronatum.

apoense Copeland in Perkins, Fragm. 185. 1905 — Mindanao.

biscaynianum (Eat.) A. A. Eaton, Fern Bull. **12**. 45. 1904 — Florida.
A. rhizophyllum var. biscaynianum Eat. Bull. Torr. Cl. **14**. 97 t. 68. 1887.

Clermontae Syme, Engl. Bot. ed. III. **12**. 132 t. 1879. 1886 — Ireland.
(= A. ruta muraria × trichomanes?).

epiphyticum Copeland in Perkins, Fragm. 184. 1905 — Mindanao.

escaleroense Christ, Hedwigia **44**. 366. 1905 — Peru.

germanicum × ruta muraria Waisb. Oest. Bot. Zeit. **49**. 63. 1899 (syn.) — Ungarn. — A. murariaeforme Waisb. l. c.

germanicum × septentrionale Waisb. Oest. Bot. Zeit. **49**. 62. 1899 (syn.) — Ungarn. — A. intercedens Waisb. l. c. — A. Luersseni Waisb. 1903; A. septentrionale × germanicum Waisb. 1903.

griseum Copeland, Polyp. Philipp. 82. 1905 — Mindanao. — A. insigne Copeland in Perkins, Fragm. 185. 1905 (non Bl. 1828 nec Liebm. 1849).

insigne Copeland in Perkins, Fragm. 185. 1905 = A. griseum.

intercedens Waisb. Oest Bot. Zeit. **49**. 62. 1899 = A. germanicum × septentrionale.

Luersseni Waisb. Mag. bot. lapok **2**. 63. 1903 = A. germanicum × septentrionale.

Martianum × mucronatum — Brasilia austr. — A. angustatum × mucronatum Rosenst. Festschr. Alb. v. Bamberg 61 t. 1. 1905.

Miyoshii Mak. Bot. Mag. Tokyo **6**. 45. 1892 (nomen) — Japonia.

monotis Christ in Schum. et Laut. Nachtr. Fl. deutsch. Südsee 43. 1905 — N. Guinea.

Moritzianum Kl.; Hemsl. Biol. Centr.-Amer. **3**. 636. 1885 = Athyrium.

murariaeforme Waisb. Oest. Bot. Zeit. **49**. 63. 1899 = A. germanicum × ruta muraria.

rhizophyllum var. biscaynianum Eat. Bull. Torr. Cl. **14**. 97 t. 68. 1887 = A. biscaynianum.

Sarasinorum Copeland, Polyp. Philipp. 87. 1905 = Athyrium.

septentrionale × germanicum Waisb. Mag. bot. lapok **2**. 63. 1903 (syn.) = A. germanicum × septentrionale.

subnormale Copeland in Perkins, Fragm. 183 t. 4 f. B. 1905 — Luzon.

Toppingianum Copeland in Perkins, Fragm. 184. 1905 — Luzon.

Ulbrichtii Rosenst. Hedwigia **43**. 220. 1904 — Brasilia austr.

ATHYRIUM

dentatum Gray, Nat. Arr. br. Pl. **2**. 11. 1821 = A. filix femina.

BLECHNUM

(B) **egregium** Copeland in Perkins, Fragm. 187. 1905 — Ins. Philipp.

(B) **Juergensii** Rosenst. Festschr. Alb. v. Bamberg 59. 1905 — Brasilia austr.

(B) **Stierii** Rosenst. Festschr. Alb. v. Bamberg 60. 1905 — Brasilia austr.

BOTRYCHIUM

californicum Und. Torreya **5**. 107. 1905 — California.

CALLIPTERIS

cordifolia Copeland, Polyp. Philipp. 70. 1905 = Diplazium.

pariens Copeland in Perkins, Fragm. 186. 1905 = Diplazium.

CHEILANTHES

Boltoni Copeland in Perkins, Fragm. 192. 1905 — Ins. Philipp.

undulata Hope et Wright, Gard. Chr. III. **34**. 397. 1903 — Yunnan.

CHRISTENSENIA Maxon, Proc. Biol. Soc. Washington **18**. 239. 1905 — (Marattiaceae Gen. **4**).

aesculifolia (Bl.) Maxon, Proc. Biol. Soc. Wash. **18**. 240. 1905 — Assam. Malesia. — Aspidium Bl. Enum. 143. 1828; Kaulfussia Bl. 1828. HB. 444. Chr. 362. NPfl. 442 — K. assamica Griff. 1836; (Macrostoma Griff. 1836) — Kaulfussia Korthalsii de Vriese 1852 — K. Lobbiana de Vriese 1853.

CHRISTOPTERIS Copeland in Perkins, Fragmenta 188. 1905 — (Polypodiaceae Gen. **99 a**).

sagitta (Christ) Copeland in Perkins, Fragm. 188. 1905 — Ins. Philipp. Polypodium Christ, Bull. Boiss. **6**. 199. 1898. NPfl. 318.

CYATHEA

Laurentiorum Christ, Mission E. Laurent 14. 1905 — Congo.

leucotricha Christ, Ann. Jard. Buit. II. **5**. 135. 1905 — Borneo.

CYRTOMIUM Butterfieldii ht.; Gard. Chr. III. **35**. 142. 1904 = Polystichum falcatum var. hort.

DAVALLIA

(P) **exaltata** Copeland in Perkins, Fragm. 180. 1905 — Mindanao.

(P) **Schlechteri** (Christ) C. Chr. Ind. 663. 1905 — N. Guinea. — Prosaptia Christ in Schum. et Laut. Nachtr. Fl. deutsch. Südsee 41 t. 1 B. 1905.

(D) **Wagneriana** Copeland in Perkins, Fragm. 180. 1905 — Mindanao.

Davallodes Copeland, Polyp. Philipp. 55. 1905 *(Microlepia §)* = Microlepia.

DICRANOPTERIS dolosa Copeland in Perkins, Fragm. 193 t. 4 f. C. 1905 = Gleichenia.

DIPLAZIUM

Griffithii Diels, Nat. Pfl. **1**^{4}. 227. 1899 = Athyrium spectabile.

palauanense Copeland in Perkins, Fragm. 186. 1905 — Ins. Philipp.

pariens (Copeland) C. Chr. Ind. 663. 1905 — Ins. Philipp. Callipteris Copeland in Perkins, Fragm. 186. 1905.

DIPTERIS

Nieuwenhuisii Christ, Ann. Jard. Buit. II. **5**. 124 t. 3 f. 3. 1905 — Borneo.

DORYOPTERIS

Juergensii Rosenst. Festschr. Alb. v. Bamberg 58. 1905 — Brasilia austr.

DRYMOGLOSSUM confertum Copeland in Perkins, Fragm. 187. 1905 = Polypodium Copelandii.

Drynariopsis Copeland, Polyp. Philipp. 116, 133. 1905 *(Polypodium §)* = Polypodium.

DRYOPTERIS

(D) **Juergensii** (Rosenst.) C. Chr. Ind. 663. 1906 — Brasilia austr. Nephrodium Rosenstock, Festschr. Alb. v. Bamberg 63. 1905.

(C) **Laurentiorum** (Christ) C. Chr. Ind. 664. 1906 — Congo.
Nephrodium Christ, Mission E. Laurent 5. 1905.

(L) **Stierii** (Rosenst.) C. Chr. Ind. 664. 1906 — Brasilia austr.
Gymnogramme Rosenstock, Festschr. Alb. v. Bamberg 64. 1905.

(D) **Tonduzii** (Christ) C. Chr. Ind. 664. 1906 — Costa Rica.
Aspidium Christ, Prim. Fl. costar. **3**. 34. 1901.

GLEICHENIA

dolosa (Copeland) C. Chr. Ind. 664. 1906 — Ins. Philipp.
Dicranopteris Copeland in Perkins, Fragm. 193 t. 4 f. C. 1905.

GYMNOGRAMMA

longisora Bak. JoB. **1890**. 267 = Polypodium.

LOXOGRAMME

minor Mak. Bot. Mag. Tokyo **19**. 139. 1905 = P. Yakushimae.
salicifolia Mak. Bot. Mag. Tokyo **19**. 138. 1905 = P. Makinoi.

PLATYCERIUM

coronarium (König) Desv. Prod. 213. 1827 — Siam-Malesia. Ins. Philipp.
Osmunda König; Müll. Naturforsch. Halle **21**. 107 t. 3. 1785; Alcicornium Und. 1905; Acrostichum biforme Sw. 1801; Platycerium Bl. 1828. HB. 425. Chr. 127. NPfl. 339; Neuroplatyceros Fée 1845.

CORRIGENDA

Pag. 5 lin. 13 lege
alcicorne Willem. Usteri Ann. **18**. 61. 1796 = ? Platycerium stemaria.
— 6 lin. 3-4 dele C. lomarioides Jenm. . . . . 1833).
— » — 5 dele lineam totam. (aureum Cav.).
— » — 29 dele alcicorne.
— 8 — 1 lege 1858 loco 1868.
— 9 — 11 ab infra lege Quito loco Peru.
— 11 — 8 lege 78 loco 80.
— 12 — 7 lege = A. aureum loco — Ins. Philipp.
— 13 — 4 lege excelsum loco aureum var.
— 15 — 20 ab infra post Willd. pone sp.
— » — 13 » — lege 102 loco 103.
— 16 — 1 lege Thouars loco Thours.
— 17 — 27 lege 1886 loco 1885.
— 19 — 28 lege 372 loco 373.
— 20 — 11 adde = ? Polybotrya apiifolia.
— 21 — 15 lege 1803 loco 1802.
— 26 — 28 adde = ? Odontosoria retusa.
— 28 — 25 lege 1856 loco 1854.
— 29 — 16 ab infra lege = A. caudatum loco — Ins. Philipp.
— 30 — 22 adde Argentina.
— 32 — 21 adde Argentina.
— 34 — 19 adde Florida.
— » — 26 adde = ? A. lunulatum.
— 36 — 13 ab infra adde = ? Diplazium bulbiferum.
— 41 — 8 » — lege 1866 loco 1865.
— 43 — 21 » — lege **goyazensis** loco goyazensis.
— » — 10 » — lege 1866 loco 1886.
— 45 — 10 dele Cyathea ferox Pr. 1822; lege Alsophila ferox Pr. 1836.
— 46 — 15 lege **36** loco 36.
— 48 — 14 ab infra lege 1845-46 loco 1844.
— 50 — 2 » — lege Swartzii loco runcinatum.

Pag. 55 lin. 16 lege 1753 loco 1763.
— 58 — 9 ab infra adde China (Shen-si).
— 59 — 11 lege **hexagona** loco hexagona.
— » — 14 ab infra lege 1827 loco 1877.
— 60 — 7 dele var.
— » — 17 lege Pl. loco Fl.
— 61 — 3-4 ab infra lege
APHYLLOCALPA Lagasca, Garcia et Clemente, Anal. Cienc. **5**. 164. 1802; Cavanilles, Descr. 556. 1803 = **Osmunda.**
regalis Lag. Garcia et Clem. Anal. Cienc. **5**. 165. 1802; Cav. l. c.
— 63 lin. 18 loco Haláczy lege Halácsy et adde Verh. zool. bot. Ges. **54**. 129 c. fig. 1904.
— » — 19 ab infra lege Christensenia loco Kaulfussia.
— » — 16 » — post Soc. adde **6**.
— 65 — 22 » — pone 6 loco —.
— » — 16 » — lege 123 loco 133.
— 66 — 2 adde Assam.
— » — 3 post 265 adde Aspidium Christ 1905.
— » — 10 lege N. Mecklenburg loco Fiji.
— 67 — 10 et 11 pone 31 loco —.
— 68 — 24 adde Dryopteris O.Ktze. 1891.
— 70 — 2 lege 1842 loco 1843.
— » — 19 dele ? et hirta var.
— » — 26 pone 32 loco —.
— » — 20 ab infra lege **2**. 166. 1844.
— » — 12 » — lege humida loco nudata.
— 71 — 4 lege pteropus loco decurrens.
— » — 10 lege 1880 loco 1810.
— 72 — 16 lege Linnaeana loco d.
— » — 22 pone 32 loco —.
— 75 — 26 lege Adenoderris viscidula loco Polystichum viscidulum.
— 77 — 2 lege 1789 loco 1879.
— 78 — 11 post 1903 adde Sagenia Christ 1905.
— » — 2 ab infra post 1902 adde Sagenia Christ 1905.
— 80 — 22 adde Sagenia Christ 1905.
— » — 28 adde Verh. zool. bot. Ges. **54**. 129 c. fig. 1904.
— 81 — 9 lege vivipara loco Swartzii.
— » — 10 ab infra pone **6**. 46 loco —.
— 84 — 12 pone 7 loco —.
— 85 — 22 ab infra pone **6**. 47 loco —.
— 86 — 17 adde Sagenia Christ 1905.
— » — 12 ab infra pone 33 loco —.
— 87 — 7 dele lineam totam. (pentangularum Col.).
— 88 — 20 post 1891 adde Sagenia Christ 1905.
— » — 24 pone **6**. 46 loco —.
— » — 16 ab infra pone 34 loco —.
— 89 — 12 adde Sagenia Christ 1905.
— » — 18 ab infra lege 167. 1844 loco —. 1845.
— 91 — 6 dele aculeatum var. 21.
— 92 — 24 pone **6**. 46 loco —.

Pag. 95 lin. 16 dele lineam totam. (tenue Retz. . . .).
— » — 24 ab infra pone 34 loco —.
— » — 11 » — dele Leprieurii.
— 97 — 13 post 1876 adde Phegopteris Salom. 1883.
— » — 14 ab infra lege Adenoderris loco Polystichum.
— » — 11 » — adde Sagenia Christ 1905.
— » — 6 lege 168. 1844 loco —. 1845.
— 98 — 2 post Dryopteris adde debilis.
— » — 5 lege
Wawraeanum Szysz. in Beck, It. Princ. Cob. **2**. 126 t. 15. 1888 — N. Zealand. (Polystichum sp.).
— 100 lin. 8 adde — A. venosum Hk. 1860.
— » — 20 lege fraxinifolium loco lineolatum.
— 102 — 1 dele (Anal. Cienc. **4**. 100. 1801).
— 103 — 19 ab infra post Diplazium adde grammitoides.
— 105 — 2 adde post Ceterach officinarum.
— 111 — 21 lege 1859 loco 1857.
— 114 — 21 lege Athyrium spectabile loco Diplazium.
— 121 — 22 ab infra pone **6**. 46 loco —.
— 122 — 12 adde Paraguay.
— » — 24 ab infra lege murale Salisb. Prod. 403. 1796.
— » — 8 » — lege **6**. 46 loco —.
— 125 — 21 » — lege Pechuēlii loco Pechuēllii.
— 131 — 8 » — lege 1803 loco 1802.
— 132 — 25 lege
shikokianum Mak. Bot. Mag. Tokyo **6**. 45. 1892 (nomen); **13**. 1899.
— » lin. 29 dele Flora **1834**[1]. Beibl. 28 (nomen).
— 135 — 21 ab infra pone 125 loco —.
— » — 13 » — lege
Toramanum Mak. Bot. Mag. Tokyo **6**. 45. 1892 (nomen); **13**. 1899.
— 137 lin. 6 ab infra pone **6**. 46 loco —.
— » — 2 » — dele Diplazium Kze. 1848.
— 138 — 15 » — lege
woodsioides Christ, Bull. Soc. Bot. It. **1900**. 261 — China.
— 142 lin. 11 lege 1801 loco 1802.
— » — 19 ab infra lege formis loco formibus.
— 143 — 11 lege Athyrium spectabile loco Diplazium.
— 144 — 18 pone **6**. 46 loco — et adde (nomen).
— » — 25 adde post hb. Hemsl. 1885.
— 145 — 24 ab infra dele ? Davallia F. Muell. 1865.
— 146 — 3 adde Asplenium Copeland 1905.
— » — 8 ab infra adde — Asplenium Griffithii Bak. 1867. HB. 239. Athyrium Milde 1870; Bedd. 1876; Diplazium Diels 1899. NPfl. 227 (non Moore 1861).
— 147 lin. 15 ab infra pone **6**. 46 loco — et adde (nomen).
— 149 — 24-18 ab infra lege
BELVISIA Mirbel, Hist. nat. vég. **5**. 111. 1803; Hist. nat. des pl. **4**. 65. 1803.
australis Mirb. Hist. nat. vég. **5**. 115. 1803 — digitata Mirb. l. c. 116 — septentrionalis Mirb. l. c. 115 — siliquosa Mirb. l. c. 116 — spicata Mirb. l. c. 114.

Pag. 150 lin. 2 ab infra lege Copeland, Polyp. Philipp. 90. 1905 loco C. Chr. Ind. 150. 1905.
— 152 — 18 ab infra lege = Oleandra loco Ins. Philipp.
— » — 3 » — pone 23 loco —.
— 154 — 12 » — pone 24 loco —.
— 155 — 11 lege **1883** loco 1883.
— 156 — 5 pone 25 loco —.
— 158 — 23 pone 26 loco —.
— 161 — 9 ab infra adde 20 Species Botrychii ad novum genus Sceptridium a Lyon 1905 translatae sunt. Vide Sceptridium.
— 163 lin. 8 lege
occidentale Und. Bull. Torr. Cl. **25**. 538. 1898 = B. silaifolium.
— 164 lin. 7 ab infra adde = ? *Onychium siliculosum.*
— 165 — 4 lege homonym. loco hymonym.
— 168 — 13 lege 13 loco 12.
— » — 17 lege 176 loco 376.
— 170 — 23 ab infra dele (Acrostichum Cav. 1801 non L. 1753).
— » — 22 » — post Cav. adde Anal. Cienc. **4**. 100. 1801.
— 171 — 18 lege (Polypodiaceae Gen. **72**).
— 173 — 6 ab infra lege Calc. loco Galc.
— 175 — 13 » — lege n. 15 loco n. 24.
— 178 — 16 adde Bolivia. Argentina.
— 181 — 12 adde Borneo.
— 182 — 21 lege excelsum loco aureum var.
— 186 — 20 ab infra lege Swartzii loco runcinatum.
— 187 — 18 » — lege **12** loco 12.
— 190 — 13 lege
Boylei F. Muell. apud C. French in South. Science Record **1**. 33. 1881 = C. Cunninghami.
— 192 lin. 9 ab infra adde an Myriotheca arborescens Blanco 1837?
— 196 — 12 lege 1900 loco 1899.
— » — 1 ab infra dele bor.
— 198 — 18 adde Borneo.
— 200 — 9 ab infra lege 1857 loco 1587.
— 201 — 10 post 350 adde Niphobolus Copeland 1905.
— 202 — 11 lege 1841 loco 1842.
— 203 — 24 adde Amer. bor. orient. temp. Alaska.
— » — 4 ab infra adde Tectaria Lag., Garcia et Clem. 1802.
— 204 — 1 adde Tectaria Lag., Garcia et Clem. 1802.
— 207 — 18 post 1849 adde Prosaptia Christ 1905.
— 208 — 2 ab infra lege 248 loco 258.
— 210 — 2 adde Humata Copeland 1905.
— » — 22 adde Prosaptia Christ 1905.
— » — 3 ab infra post 1876 adde Humata Copeland 1905.
— 211 — 4 adde Ins. Philipp.
— 212 — 6 ab infra lege oligophlebia Bak. JoB. **1888**. 323 = Tapeinidium.
— 214 — 11 ad finem adde Prosaptia Christ 1905.
— 217 — 19 adde N. Guinea.
— 218 — 10 ab infra lege 1861 loco 1857.

Pag. 228 lin. 6 ab infra lege
brachypodum Copeland, Polyp. Philipp. 75. 1905 = D. grammitoides.
— 229 lin. 15 lege Copeland, Polyp. Philipp. 73. 1905 loco C. Chr. Ind. 229. 1905.
— 230 — 16 post 1875 adde Callipteris Copeland 1905.
— 233 — 9 adde Asplenium brachypodum Bak. 1874. HB. 490; Diplazium Copeland 1905.
— » — 14 dele NPfl. 227.
— » — 15 dele lineana totam.
— 235 — 20 ab infra post 1870 adde Allantodia Racib. 1905.
— 236 — 18 » — lege 7. 66 t. 48. 1804 loco (fasc.? **19**. 67 t. 48).
— 250 — 4 » — lege t. 271 loco t. 171.
— 252 — 23 adde Nephrodium Copeland 1905.
— 253 — 20 adde Nephrodium Copeland 1905.
— 257 — 18 ab infra adde Nephrodium (pallidum) Hemsl. 1885.
— » — 16 » — lege Hemsl. 1885 loco Bak. 1891.
— 259 — 23 — post 258 adde Tectaria Lag., Garcia et Clem. 1802.
— 260 — 11 » — adde N. Guinea.
— » — 9 » — adde Aspidium Warburgii Kuhn et Christ 1900: Nephrodium Diels 1901.
— 264 — 2 ab infra lege 1801 loco 1802.
— 267 — 15 » — lege (D) loco (?) et adde Nephrodium Clute 1902.
— 268 — 6 » — adde Nephrodium Copeland 1905.
— 274 — 10 » — dele Aspidium Tonduzii Christ 1901.
— 275 — 16 » — adde Nephrodium Copeland 1905.
— 282 — 10 adde Nephrodium Copeland 1905.
— 287 — 12 ab infra adde Nephrodium Copeland 1905.
— 292 — 15 post 27 adde 1825.
— 293 — 23 ab infra dele Lastrea Bedd. 1876.
— 300 — 1 adde Jamaica.
— 302 — 23 ab infra post Platycerium adde bifurcatum.
— 304 — 15 » — lege **cinnamomeum** loco **cinnamorneum**.
— 315 — 4 lege 1886 loco 1885.
— 318 — 4 ab infra lege Fawcettii loco Fawzettii.
— 346 — 1 ab infra lege **39**^{2} loco **30**^{2}.
— 367 — 11 » — lege semibivalve loco semibivaloe.
— 374 — 1 » — lege D. Balbisii loco D. patens *stipularis.
— 377 — 18 lege Aspidium loco Asyndium.
— » — 13 ab infra lege ochthodes loco prolixa.
— 399 — 27 lege 1877 loco 1876.
— 401 nota lege 1906 loco 1905.
— 406 lin. 13 ab infra lege Photinopteris loco Dryostachyum.
— 409 — 3 lege 1877 loco 1876.
— 414 — 21 lege 442 loco 142.
— 423 — 3 ab infra lege 1869 loco 1899.
— 459 — 21 lege 373 loco 273.
— » — 13 ab infra lege Notholaena loco Nothochlaena.
— 470 — 5 » — lege 1901 loco 1801.
— 507 — 5 ante 288 adde **3**.
— 508 — 8 ante 287 adde **3**.
— 537 — 10 lege Willd. loco Wild.

Pag. 539 lin. 6 lege balaonense loco baloonense.
— 543 — 16 lege 1795 loco 1759.
— » — 7 ab infra lege 274 t. 61 f. 3 loco 275 t. 61 f. 4.
— 544 — 10 » — lege **Merrillii** loco **Merillii**.
— » — 16 lege 384 loco 884.
— 556 — 13 ab infra lege 1778 loco 1878.
— 576 — 19, 17, 13 ab infra adde Chile.
— » — 5 ab infra ante Tijucense adde Polystichum.
— 581 — 9 lege 191 loco 199.
— 604 — 3 lege Ag. loco Ac.
— » — 11 lege denticulata loco dendiculata.
— 624 — 21 lege stegnogrammoides loco stenogrammoides.
— 637 — 24 ab infra lege Panch. loco Pansch.
— 647 — 22 lege 1863 loco 1861.
— » — 17 ab infra lege 391 loco 321.

CATALOGUS LITERATURAE

ENUMERATIO ALPHABETICA ET SYSTEMATICA
OMNIUM OPERUM IN QUIBUS NOVA
GENERA ET SPECIES FILICUM ET HYDROPTERIDUM
DESCRIPTA SUNT

CUM

ABBREVIATIONIBUS OMNIBUS NOMINUM AUCTORUM
ET LIBRORUM IN INDICE CITATORUM

I.

ENUMERATIO ALPHABETICA

ABBREVIATIONES NONNULLAE SCRIPTORUM PERIODICORUM

Amer. Journ. Sci. = American Journal of Science, ed. Silliman etc.

Ann. Bot. = Annals of Botany. Oxford. 8⁰.

Ann. Jard. Buit. = Annales du Jardin Botanique de Buitenzorg. Leide. 8⁰.

Ann. sc. nat. = Annales du sciences naturelles. Botanique. Paris. 8⁰.

Ark. för Bot. = Arkiv för Botanik utgifvet af K. svenska Vetenskapsakademien. Stockholm. 8⁰.

Berl. Mag. = Der Gesellschaft Naturforschender Freunde zu Berlin Magazin für die neuesten Entdeckungen in der gesammten Naturkunde. Berlin 1807-1818. 4⁰.

Bot. Gaz. = The Botanical Gazette. Chicago. 8⁰.

Bot. Mag. Tokyo = The Botanical Magazine. Edited by the Tōkyō Botanical Society. Tōkyō. 8⁰.

Bot. Tidsskr. = Botanisk Tidsskrift. København. 8⁰.

Bot. Zeit. = Botanische Zeitung. Berlin. 4⁰.

Bull. Acad. Géogr. Bot. Mans = Bulletin de l'Académie internationale de Géographie Botanique. Le Mans. 8⁰.

Bull. Boiss. = Bulletin de l'herbier Boissier. Genève. 8⁰.

Bull. Soc. bot. it. = Bulletino della Società botanica italiana. Firenze. 8⁰.

Bull. Soc. Fr. = Bulletin de la Société botanique de France. Paris. 8⁰.

Bull. Torr. Cl. = Bulletin of the Torrey Botanical Club. New York. 8⁰.

Engl. Jahrb. = Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie ed. A. Engler. Leipzig. 8⁰.

Gard. Chr. = The Gardener's Chronicle. London. 4⁰.

JoB. = Journal of Botany ed. W. J. Hooker (vol. 1-4 et 1-9), Seemann, Trimen, Britten. London. 8⁰.

Journ. Linn. Soc. = Journal of the Linnean Society. Botany. London. 8⁰.

Kew Bull. = Bulletin of Miscellaneous Information. Royal Gardens Kew. London. 8⁰.

Linn. = Linnaea. Ein Journal für die Botanik in ihrem ganzen Umfange. Berlin. 8⁰.

Lond. JoB. = W. J. Hooker: The London Journal of Botany. London. 8⁰.

Ned. kr. Arch. = Nederlandsch kruidkundig Archief. Leyden. 8⁰.

Nova Acta = Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae curiosorum. 4⁰.

Nu. Giorn. bot. It. = Nuovo Giornale botanico Italiano (n. s. = Nuova Serie). Firenze. 8⁰.

Proc. Amer. Ac. = Proceedings of the American Academy of Arts and Science. Boston. 8⁰.

Schrad. Journ. = Journal für die Botanik, herausgegeben von Professor Schrader. Göttingen 1799-1803. 8⁰.

Schrad. Neu. Journ. = Neues Journal für die Botanik, herausgegeben von Professor Schrader. Erfurt 1806-1810. 8⁰.

Tr. Linn. Soc. = Transactions of the Linnean Society. London. 4⁰.

Tr. N. Zeal. Inst. = Transactions and Proceedings of the New Zealand Institute. Wellington. 8⁰.

Verh. zool. bot. Ges. = Verhandlungen der K. K. zoologischen-botanischen Gesellschaft in Wien. Wien. 8⁰.

Vet. Ak. Handl. = Kongliga Svenska Vetenskaps-Akademiens Handlingar. Stockholm. 8⁰. Ny Följd. 4⁰.

Vid. Selsk. Afh. (Skr.) = Kgl. Danske Videnskabernes Selskabs Afhandlinger (Skrifter). Naturvidenskabelige og mathematiske Afdeling. Kobenhavn. 4⁰.

Series scriptorum periodicorum numero Romano (II) indicatur.
Volumen libri numero crassiusculo (**2**) indicatur.
Signum * ante titulum abbreviatum opus a me non visum indicat.

Cl. A. GEPP nomina nova ex libris nonnullis a me non visis mecum communicavit.

Tituli librorum a PRITZEL in »Thesauro Literaturae botanicae« enumerati abbreviati sunt.

Ach. Vet. Ak. Handl. **1809**.	**E. Acharius:** Ophioglossum microstichum, en ny Svensk Wäxt beskrefven af — pag. 59 t. 3 A.
Adans. Fam. d. pl. **2**. 1763.	**M. Adanson:** Familles des plantes . . . Paris. 8⁰.
Ag. Rec. 1839.	**J. G. Agardh:** Recensio specierum generis Pteridis. Lundae. 8⁰.
Ait. Hort. Kew. **3**. 1789. ed. II. **5**. 1813.	**W. Aiton:** Hortus Kewensis; or, a Catalogue of the Plants cultivated in the Royal Botanic Garden at Kew. London. 8⁰. — ed. II. Filices a R.Brown.
All. Fl. pedem. **2**. 1785.	**C. Allione:** Flora pedemontana . . . . . Augustae Taurinorum. Fol.
Amann: Nat. Tijds. Ned. Ind. **23**. 1861.	**J. Amann** (pseudonymus = Sulpice Kurz): Opsomming der Vaatkrijptogamen van het Eiland Bangka. — pag. 399.
Anders. Vet. Ak. Handl. **1853**. 1855.	**N. J. Andersson:** *Om Galapagos-öarnes vegetation.* — pag. 61. (etiam in kgl. Svenska Fregatten Eugenies Resa. Vetensk. Iakttagelser. Botanik. Stockholm 1857. 4⁰).
Ångstr. Bot. Notiser **1854**.	**J. Ångström:** Anmärkningar om de skandinaviska arterna af slägtet Botrychium, med en Planch. — pag. 65.
Armstr. Tr. N. Zeal. Inst. **13**. 1881.	**J. B. Armstrong:** 1. Descriptions of new and rare New Zealand Plants. — pag. 335.
——— — —	— 2. A natural Arrangement of the New Zealand Ferns founded on the System of Smith's »Historia Filicum«, with critical Notes on certain species. — pag. 359.
— ——— **14**. 1882.	— 3. Descriptions of new plants. — pag. 359.
Aschers. Fl. Brandenb. (1859-) 1864.	**P. Ascherson:** 1. Flora der Provinz Brandenburg. Berlin. 8⁰.
— Verh. bot. Ver. Brand. **37**. 1896.	— 2. Asplenium trichomanes × ruta muraria (A. Preismanni Aschers. et Luerss.). — pag. XLVI.
— **et Graebn.** Syn. **1**. 1896.	- **und P. Graebner:** Synopsis der mitteleuropäischen Flora. Leipzig 1898. 8⁰. (Filices Ascherson elaboravit).
Aublet: Hist. pl. Guian. **2**. 1775.	**F. Aublet:** Histoire des plantes de la Guiane françoise . . . . Londres et Paris. 8⁰.
Bab. Trans. Bot. Soc. Edinb. **2**. 1846.	**C. C. Babington:** On the correct Nomenclature of the Lastraea spinosa and L. multiflora of Newman. — pag. 189.

Bail *Hdb. Ferns Queensl. 1874.	**F. M. Bailey:** 1. Handbook of the Ferns of Queensland. Brisbane. 8^0.
— *Proc. Linn. Soc. N. S. Wales **3**. 1879.	— 2. On the ferns of Queensland. — pag. 118.
— * ——— **4**. 1880.	— 3. On a new species of fern, Asplenium Prenticei. — pag. 36.
— * ——— **5**. 1881.	— 4. On Queensland ferns, with a description of two (four) new species. — pag. 29.
— *Fern World 1881.	— 5. The Fern World of Australia. Brisbane. 8^0.
— *Syn. Queensl. Flora 1883. *Suppl. I. 1886. * — II. 1888. * — III. 1890.	— 6. Synopsis of the Queensland Flora. Brisbane. 8^0. First-Third Supplement. — (t. A. Gepp).
— *Proc. Roy. Soc. Queensl. **1**. 1884.	— 7. Contributions to the Queensland Flora. — pag. 9, 148. Tab. — (t. A. Gepp).
— *Rep. Govt. Sci. Exp. Bellenden-Ker. 1889.	— 8. Botany in Queensland Department of Agriculture: Report Government scientific expedition to Bellenden-Ker Range. Brisbane 1899. (Species novae etiam in nr. 6 Suppl. III. descriptae). — (t. A. Gepp).
— *Queensl. Dept. Agric. Bull. **7**. 1891. **13**. 1891. — *Queensl. Bot. Bull. **4**. 1891. **5**. 1892. **13**. 1896. **14**. 1896.	— 9. Contributions to the Queensland Flora. Queensland Department of Agriculture. Brisbane. Bulletin nr. **7** (March 1891); nr. **13** (Dec. 1891) = Queensland Botany Bulletin no. **4**; Queensl. Bot. Bull. nr. **5**; nr. **13** (April 1896); nr. **14** (Oct. 1896). Tab. — (t. A. Gepp).
— *Lithogr. 1892.	—10. Lithograms of the Ferns of Queensland. Brisbane. 8^0. 191 Tab. — (t. A. Gepp).
— *Queensl. Agric. Journ. **1**. 1897. **3**. 1898. **9**. 1901.	—11. Contributions to the Flora of Queensland. **1**. pag. 368 (?); **3**. pag. ?; **9**. pag. 140. Tab. — (t. A. Gepp).
— *Proc. R. Soc. Queensl. **4**. 1899.	—12. Notes on the vegetation of New Guinea. — pag. 14.
Bail. *Haw. Ferns, 1883.	**Bailey:** Hawaiian Ferns. ?
Bak. Syn. 1866-1868. ed. II. 1874.	**J. G. Baker:** 1. vide Hooker and Baker.
— Journ. Linn. Soc. **9**. 1866.	— 2. Descriptions of six new Species of simple-fronded Hymenophyllaceae. — pag. 335 t. 8.
— in Hk. Ic. pl. 1867-1892.	— 3. Filices in Hooker: Icones plantarum. vol. **11**. t. 1015, 1016. 1867. **12**. t. 1132. 1873. **17**. t. 1601-1675. 1886; t. 1676-1700. 1887 (saepe ab auctt. cit. III. century of Ferns). **23**. t. 2229. 1892.
— Fl. bras. 1^2. 1870. $^1/_5$.	— 4. Cyatheaceae et Polypodiaceae in Martius: Flora brasiliensis vol. **1**. pars 2. — pag. 305. t. 20-70.
— Gard. Chr. **1872**.	— 5. Ferns of Lord Howe's Island. — pag. 252.
— JoB. **1872**.	— 6. On a new Ceylonese Acrostichum. — pag. 146.
— — —	— 7. On a new Asplenium from Cape Colony. — pag. 362.
— — 1873.	— 8. New Ferns from Lord Howe's Island. — pag. 16.

Bak. JoB. **1873**.	**J. G. Baker:** 9. A new Genus of Ferns of the Tribe Asplenieae from the Solomon Islands. — pag. 235 cum fig.
— Journ. Linn. Soc. **14**. 1873.	—10. On the Recent Synonyms of Brazilian Ferns. — pag. 12.
— JoB. **1874**.	—11. On two New Species of Pellaea from Namaqualand. — pag. 199.
— — —	—12. Tree Ferns from Lord Howe's Island. — pag. 279.
— — **1875**.	—13. Diplora integrifolia. — pag. 78.
— — —	—14. On Lindsaea viridis of Colenso: an undescribed New Zealand Fern. — pag. 108.
— — —	—15. On a Collection of Ferns gathered in Central China by Dr. Shearer. — pag. 199.
— — —	—16. On a Collection of Chinese Ferns gathered by Mr. J. F. Quekett. — pag. 291.
— Journ. Linn. Soc. **14**. 1875.	—17. Filices apud J. D. Hooker: On the Flora of the Island of St. Paul. — pag. 479.
— Tr. Irish Acad. **25**. 1875.	—18. On the Seychelles Fern flora. — pag. 509. t. 28-31.
— JoB. **1876**.	—19. On a Collection of Ferns made in Samoa by the Rev. S. J. Whitmee. — pag. 9.
— — —	—20. On a second Collection of Ferns made in Samoa by the Rev. S. J. Whitmee. — pag. 342.
— Journ. Linn. Soc. **15**. 1876.	—21. On the Polynesian Ferns of the »Challenger« Expedition. — pag. 104.
— ———— —	—22. On a Collection of Ferns made by Mr. William Pool in the interior of Madagascar. — pag. 411.
— Fl. Maur. 1877.	—23. Flora of Mauritius and the Seychelles etc. London. 8⁰.
— JoB. **1877**.	—24. Descriptive Notes on a few of Hildebrandt's East African Plants. — pag. 65.
— — —	—25. New Ferns from the Andes of Quito. — pag. 161.
— — —	—26. Two new Ferns from Japan. — pag. 366.
— Journ. Linn. Soc. **16**. 1877.	—27. On a Collection of Ferns made by Miss Helen Gilpin in the interior of Madagascar. — pag. 197.
— JoB. **1878**.	—28. List of Balansa's Ferns of Paraguay, with Descriptions of the new Species. — pag. 299.
— — **1879**.	—29. Report on a Collection of Ferns made in the North of Borneo by Mr. F. W. Burbidge. — pag. 37.
— — —	—30. Report on Burbidge's Ferns of the Sulu Archipelago. — pag. 65.
— — —	—31. On a Collection of Ferns gathered in the Fiji Islands by Mr. John Horne, F. L. S. — pag. 292.
— — —	—32. Four new Ferns from South China. — pag. 304.
— — **1880**.	—33. On a Collection of Ferns made by Dr. Beccari in Western Sumatra. — pag. 209.
— — —	—34. On a Collection of Ferns made by Langley Kitching Esq. in Madagascar. — pag. 336. 369.

Bak. Gard. Chr. n. s. **14**. 1880.	**J. G. Baker:** 35. New Ferns of China and Japan. — pag. 494.
— JoB. **1881**.	—36. On a Collection of Ferns made by Mr. W. Kalbreyer in New Granada. — pag. 202.
— — —	—37. On a Collection of Ferns made by Mr. Curtis in the Malay Islands and Madagascar. — pag. 366.
— Journ. Linn. Soc. **18**. 1881.	—38. Filices apud G. Watt: Notes on the Vegetation etc. of Chumba State and British Labone etc. t. 14 A.
— *in Oates Matabelel. App. **5**. 1881.	—39. Filices in Oates: Matabele Land and the Victoria Falls etc. Appendix V. Botany by M. D. Oliver. 8⁰.
— JoB. **1882**.	—40. New Ferns from Southern Brazil. pag. 309.
— Journ. Linn. Soc. **19**. 1882.	—41. On a Collection of Ferns made by the Rev. R. B. Comins in the Solomon Islands. — pag. 293.
— JoB. **1883**.	—42. Ferns collected by the Rev. J. Hannington in E. Tropical Africa. — pag. 245.
— Journ. Linn. Soc. **20**. 1883.	—43. Contributions to the Flora of Madagascar. part III. — pag. 237.
— JoB. **1884**.	—44. New Plants from the Zambesi Country. — pag. 52.
— — —	—45. Ferns collected in Madagascar by M. Humblot. — pag. 139.
— — —	—46. Ferns collected in Costa Rica by Mr. P. G. Harrison. — pag. 362.
— Bull. Soc. Linn. Paris **1**. 1885. (n. 67.)	—47. Liste des Fougères des Comores, rapportées par M. Humblot. — pag. 532.
— JoB. **1885**.	—48. Ferns collected in North Formosa by Mr. William Hancock. — pag. 102.
— — —	—49. New Ferns from Brazil collected by Dr. Glaziou. — pag. 217.
— Journ. Linn. Soc. **21**. 1885.	—50. On the Flora of Madagascar. — pag. 455.
— Malesia **2**. **3**. 1886.	—51. Filices in Beccari: Malesia, vide Beccari.
— JoB. **1886**.	—52. A Synopsis of the Rhizocarpeae. — pag. 97, 274, 381.
— — —	—53. New Ferns collected by J. B. Thurston, Esq., in Fiji. — pag. 182.
— — —	—54. A New Tree Fern from Central America. — pag. 243.
— Journ. Linn. Soc. **22**. 1886.	—55. On a Collection of Ferns made in North Borneo by the Bishop of Singapore and Sarawak. — pag. 222 t. 11-12.
— Bol. Soc. Brot. **4**. 1886.	—56. Filices apud Henriques: Flora de San Thomé. — pag. 149 t. 1-2.
— in Hk. Ic. pl. t. 1601-1700. 1886-87.	— vide n. 3.
— Handb. 1887.	—57. Handbook of the Fern-allies. A Synopsis of the Genera and Species of the Natural Orders Equisetaceae, Lycopodiaceae, Selaginellaceae, Rhizocarpeae. London. 8⁰.

Bak. JoB. **1887**.	**J. G. Baker:** 58. Mr. J. J. Cooper's Costa Rica Ferns. — pag. 24.
— — —	—59. A new Polypodium from Jamaica. — pag. 44.
— — —	—60. On a collection of Ferns made in West Central China by Dr. A. Henry. — pag. 170.
— Journ. Linn. Soc. **22**. 1887.	—61. Further Contributions to the Flora of Madagascar. — pag. 441.
— ——— **24**. 1887.	—62. On a further collection of Ferns from West-Borneo made by the Bishop of Singapore and Sarawak. — pag. 256.
— Tr. Linn. Soc. II. Bot. **2**. 1887.	—63. Filices in The Botany of the Roraima Expedition of 1884. — pag. 288 t. 53-55.
— ——— —	—64. Filices apud Oliver: Enumeration of the Plants collected by Mr. H. H. Johnston on the Kilimanjaro Expedition 1884. — pag. 357.
— JoB. **1888**.	—65. On a Collection of Ferns made by Baron Eggers in St. Domingo. — pag. 33.
— — —	—66. On two recent Collections of Ferns from Western China. — pag. 225.
— — —	—67. On a third Collection of Ferns made in West Borneo by the Bishop of Singapore and Sarawak. — pag. 323.
— — —	—68. On a new Acrostichum from Trinidad. — pag. 371.
— — **1889**.	—69. New Ferns from Western China. — pag. 176.
— — —	—70. On a new Species of Polypodium from Jamaica. — pag. 270.
— Journ. Linn. Soc. **25**. 1889.	—71. Further Contributions to the Flora of Madagascar. — pag. 294.
— ——— —	—72. Filices in Report on the Botanical Collections from Christmas Island, Indian Ocean. — pag. 351.
— JoB. **1890**.	73. Vascular Cryptogamia of New Guinea, collected by Mr. W. Macgregor. — pag. 103.
— — —	—74. Tonquin Ferns. — pag. 262.
— Ann. Bot. **5**. 1891.	—75. On the Vascular Cryptogamia of the Island of St. Vincent. — pag. 163 t. 10-11.
— ——— —	—76. A Summary of new Ferns discovered since 1874. — pag. 181, 301, 455. t. 14.
— JoB. **1891**.	—77. Ferns of North West Madagascar. — pag. 3.
— — —	—78. New Ferns from West Borneo. — pag. 107.
— Ann. Bot. **6**. 1892.	—79. On the Vascular Cryptogamia of the Island of Grenada. — pag. 95.
— Journ. Linn. Soc. **29**. 1892.	—80. Filices apud Hemsley: Observations on an Botanical Collection made by Mr. A. E. Pratt in Western China. — pag. 298.
— Kew Bull. **1892**. **1893**. **1894**. **1895**. **1896**. **1898**. **1901**.	—81. Filices in Decades Kewenses I. **1892**. pag. 86. VI. **1893**. pag. 209. VII. **1894**. pag. 7. XIV. **1895**. pag. 53. XXVII. **1896**. pag. 40. XXXI-XXXIII. **1898**. pag. 224 (Filices Henryanae). XXXIV-XXXV. **1901**. pag. 144.

Bak. Engl. Jahrb. **17**. 1893.	**J. G. Baker:** 82. Filices apud P. Taubert: Plantae Glaziovianae novae vel minus cognitae, pars IV. — pag. 522.
— Ann. Bot. **8**. 1894.	—83. New Ferns of 1892-3. — pag. 121.
— Tr. Linn. Soc. II. Bot. **4**. 1894.	—84. Filices apud O. Stapf: On the Flora of Mount Kinabalu in North Borneo. — pag. 69.
— Kew Bull. **1895**. **1897**. **1901**.	—85. Filices in Diagnoses Africanae. VII. **1895**. pag. 229. X. **1897**. pag. 299. XIII. **1901**. pag. 137.
——— **1899**.	—86. Filices in Flora of british New Guinea. — pag. 95.
— Gard. Chr.	—87. [Diagnoses of single Species in several of the volumes].
Balbis: Elenco del. Piante 1801.	**G. B. Balbis:** Elenco delle Piante crescenti ne'contorni di Torino. Torino. 8⁰.
Balfour: Phil. Tr. R. Soc. London **168**. 1879.	**Is. Baylay Balfour:** Botany of Rodriguez. — pag. 302.
Baumg. Enum. stirp. Trans. **4**. 1846.	**J. C. G. Baumgarten:** Enumeratio stirpium Magno Transsilvaniae principatui praeprimis indigenarum . . . . Vindobonae. 8⁰.
Beccari: Malesia **2**. 1886. **3**. 1886.	**O. Beccari:** Malesia. Firenze. 4⁰.
	— 1. — **2**. pag. 243 t. 65. (Species novae a Baker descr.).
	— 2. Rivista delle Felci e Licopodiacee di Borneo e della Nueva Guinea, enumerate o descritte dal Barone V. Cesati etc. **3**. pag. 16 t. 4-5. — (Species novae a Baker descr.).
	— 3. Nota sopra alcune Felci raccolti dal Sign. J. E. Teysmann all' Isola di Sumba o Sandal-Wood Ed in Timor. **3**. pag. 56. (Species novae a Baker descr.).
Beck: *Bot. North. St. 1833.	**L. C. Beck:** Botany of the Northern and Midland States . . . Albany. 8⁰.
Bedd. *Ferns S. Ind. 1863-1864 (1865?).	**R. H. Beddome:** 1. The Ferns of Southern India; being descriptions and plates of the Ferns of the Madras presidency. Madras. 4⁰. 271 Tab. (fasc. 1-10) = tab. 1-123 ed. 1863. (— 11-ult.) = tab. 124-271 ed. 1864 (an 1865 pro parte ?) — (t. B. Daydon Jackson in litt.).
— Ferns br. Ind. 1865-1870.	— 2. The Ferns of British India; being descriptions and figures of ferns from all parts of British India (excluded of those figured in the Ferns of Southern India and Ceylon). Madras. 4⁰. 345 Tab.

(fasc. 1- 8)	=	tab. 1-120:	1865.
(— 9-14)	=	» 121-210:	1866.
(— 15-17)	=	» 221-255:	1867.
(— 18-20)	=	» 256-300:	1868.
(— 21-22)	=	» 301-330:	1869.
(— 23)	=	» 331-345:	1870.

— (t. B. Daydon Jackson in litt.).

Bedd. Fernsbr. Ind. Suppl. 1876.	**R. H. Beddome:** 3. Supplement to the Ferns of Southern India and British India etc. Madras. 4⁰. t. 346-390.
— Handb. 1883.	— 4. Handbook to the Ferns of British India, Ceylon and the Malay Peninsula. Calcutta. 8⁰. 300 Illustr.
— JoB. **1887.**	— 5. Ferns collected in Perak by Father Scortechini. — pag. 321 t. 278.
— — **1888.**	— 6. Ferns collected in Perak and Penang by Mr. J. Day. — pag. 1 t. 279.
— — —	— 7. New Manipur Ferns collected by Dr. Watt. — pag. 234.
— — **1889.**	— 8. Two new Athyriums from the N. W. Himalayas. — pag. 72.
— Handb. Suppl. 1892.	— 9. Supplement to the Ferns of British India, Ceylon and the Malay Peninsula, containing ferns which have been discovered since the publication of the Handbook to the Ferns of British India. Calcutta. 8⁰.
— JoB. **1893.**	—10. Scortechini's Malayan Ferns. — pag. 225.
— — —	—11. Notes on Indian Ferns. — pag. 227.
Bellardi: Mém. Ac. Turin **5.** 1792.	**L. Bellardi:** Appendix ad Floram Pedemontanam. — pag. 209.
Benth. Fl. Hongk. 1861.	**G. Bentham:** 1. Flora Hongkongensis . . . London. 8⁰. Supplementum vide Hance Nr. 8.
— Fl. austr. **7.** 1878.	— 2. Flora australiensis. London. 8⁰.
Berg. Acta Ac. Petr. **1782**[2]. 1786.	**P. J. Bergius:** Caenopteris, novum e filicibus genus. — pag. 248 t. 5-7. (Acta Acad. scient. imp. Petropol. pro anno 1782 pars posterior. 1786. 4⁰).
Bernh. Schrad. Journ. **1799**[1].	**J. J. Bernhardi:** 1. Tentamen novae generum filicum et specierum earum Germaniae indigenarum dispositionis. — pag. 291.
— ——— **1800**[2]. 1801.	— 2. Tentamen alterum filices in genera redigenda. — pag. 121. t. 1-2.
— Schrift. Akad. Erfurt **1802.**	— 3. Ueber Asplenium und einige ihm verwandte Gattungen der Farrenkräuter. t. 1. (seorsim impr. v. Willd. n. 3).
— Schrad. Journ. **1801**[1]. 1802.	— 4. Wahrscheinliche Entdeckung der Befruchtungsart der Farrenkräuter. — pag. 1. tab. 1.
— Schrad. neu. Journ. **1**[2]. 1806.	— 5. Dritter Versuch einer Anordnung der Farrnkräuter. — pag. 1. tab. 2-3.
Bertol. Opusc. sci. Bol. **1.** 1817.	**A. Bertoloni:** 1. Osservazioni Botaniche. — pag. 229. tab. 8.
— Amoen. Ital. 1819.	— 2. Amoenitates Italicae. Bononiae. 4⁰.
— Nov. Comment. Bonon. **4.** 1840.	— 3. Florula guatemalensis. — pag. 403.
— *Misc. bot. **18.** 1857. **21.** 1860.	— 4. Miscellanea botanica. Bononiae. 4⁰. — (Diagnoses breves et in Rendic. dell' Accadem. delle Scienze. Bologna. misc. **18** in **1856-57**. p. 53. 1857; misc. **21** in **1859-60**. p. 62. 1860).

Bertol. Fl. Ital. Cr. **1**. 1858.	**A. Bertoloni:** 5. Flora Italica Cryptogama. Bononiae. 8^0.
Bitter: Nat. Pfl. 1^4. 1900.	**G. Bitter:** Marattiaceae, Ophioglossaceae in Engler und Prantl: Die natürlichen Pflanzenfamilien. 1 Teil 4 Abt. Leipzig. 8^0.
Blanco: Fl. Filip. 1837. ed. II. 1845.	**M. Blanco:** Flora de Filipinas. Manila 8^0. Conf. Merrill.
Blanf. *Simla Nat. Hist. Soc. **1886**.	**H. F. Blanford:** 1. Silver Ferns of Simla. —
— Journ. As. Soc. Bengal 57^2. 1888.	— 2. A. List of the Ferns of Simla in the N. W. Himalaya between Levels of 4,500 and 10,500 Feet. — pag. 294. tab. 16-21.
Bl. Enum. 1828.	**C. L. Blume:** 1. Enumeratio Plantarum Javae et Insularum adjacentium. fasc. II. Filices. Lugduni Batavorum. 8^0. ed. nov. 1830.
— Fl Jav. Fil. 1828-1829.	— 2. Florae Javae nec non Insularum adjacentium. — Filices. Bruxellis. Fol. t. 1-94. (t. 1-65. 1828; t. 66-94. 1829). Supplementum t. 95-96. Anno?
Boiss. *Elench. pl. nov. 1838.	**E. Boissier:** Elenchus plantarum novarum minusque cognitarum, quas in itinere hispanico legit. Genevae. 8^0.
Boj. *Hort. Maur. 1837.	**W. Bojer:** Hortus Mauritianus, ou énumération des plantes exotiques et indigènes qui croissent à l'île Maurice. Maurice. 8^0. Nov. Sp. sine descript.
Bolle: Bonpl. **3**. 1855.	**C. Bolle:** 1. Novitiae florae caboverdicae ... Filices. — pag. 121.
— — **7**. 1859.	— 2. Asplenium Newmani und Cheilanthes guanchica, zwei neue Farnspezies. — pag. 106.
— Zeitschr. f. Erdk. n. F. **14**. 1863. **17**. 1864. Zeitschr. Ges. f. Erdk. **1**. 1866.	— 3. Die Standorte der Farrn auf den canarischen Inseln, pflanzen-topographisch geschildert. Berlin. **14**. pag. 289. **17**. pag. 249. **1**. pag. 209, 273.
Bolt. Fil. Brit. 1785. 1790.	**J. Bolton:** Filices Brittanicae: an History of the British proper Ferns etc. 4^0. part I. Leeds t. 1-31 col. part II. Huddersfield t. 32-46 col.
Bomm. Mon. Foug. 1867.	**J.-E. Bommer:** 1. Monographie de la classe des fougères. Classification. Bull. Soc. Bot. Belg. **5**.
— Bull. Soc. Fr. **20**. 1873.	— 2. Revue et Classification des Cyathéacées. — pag. XVI.
— et Christ: Bull. Boiss. **4**. 1896.	**— et H. Christ:** 1. Filices novae. — pag. 656.
— ——— Bul. Soc. bot. Belg. **35**. 1896.	— ——— 2. Filices in Th. Durand et H. Pittier: Primitiae Florae costaricensis; troisième fascicule. — pag. 167. (vide Christ n. 37).
Borck. Röm. Arch. 1^3. 1798.	**M. B. Borckhausen:** Monographie der in der oberen Graffschaft Catzenellebogen und der benachbarten Gegend einheimischen, auch einiger anderer kryptogamischer Gewächse aus Linné's erster Ordnung der 24ten Classe. — pag. 1.

Bory: Ess. Isles fort. 1803.	**Bory de Saint Vincent:** 1. Essais sur les isles fortunées et l'antique Atlantide etc. Paris. 4^0. (an XI).
— Voy. **1-3**. 1804.	— 2. Voyage dans les quatre principales iles des mers d'Afrique. Paris. 8^0. Tabb. in 4^0.
— Dict. class. 1824-1831.	— 3. [Plusieurs genres de fougères dans] »Dictionnaire classique d'histoire naturelle.« Paris. 8^0. **6**. 1824. **8**. 1825. **9**. 1826. **10**. 1826. **15**. 1829. **17**. 1831.
— Ann. sc. nat. **5**. 1825.	— 4. Sur un sousgeure à former parmi les Polypodes, sous le nom Drynaire, Drynaria. — pag. 462 t. 12-14.
— Dup. Voy. Bot. **1**. 1827-29.	— 5. in L. J. Duperrey: Voyage antour du Monde etc. ... sur la corvette de Sa Majesté, La Coquille. Botanique. **1**. partie. Cryptogamie. Paris. 4^0. Atlas: Fol. t. 27-38. p. 1-96. 1827; p. 97-136. 1828; p. 137-300. 1829 (t. JoB. **1901**. 206).
— Exp. sci. Morée 3^2. 1832.	— 6. in Expédition scientifique de Morée. Section des sciences physiques. Tome **3**. 2. partie. Botanique. Paris. 4^0. Atlas in Fol.
— Bél. Voy. Bot. **2**. 1833.	— 7. in C. Bélanger: Voyage aux Indes orientales etc. Botanique. **2**. partie. Cryptogamie. Paris. 8^0. Atlas in 4^0. t. 2-8. — (Sine anno; teste Kuhn: Bot. Zeit. **1868**. 233 hoc vol. anno 1833 editum est. Vide et JoB. **1901**. 205.
v.d.B. *Pl. Jungh. **1**. 1856.	**R. van den Bosch:** 1. Hymenophyllaceae in Plantae Junghuhnianae. Enumeratio plantarum quas in insulis Java et Sumatra detexit Fr. Junghuhn. Lugduni-Batavorum. 8^0. **fasc. 5. inedita.**
— Ned. kr. Arch. **4**. 1859.	— 2. Synopsis Hymenophyllacearum. — pag. 341.
— ——— 5^2. 1861.	— 3. Hymenophyllaceas novas, cum ab aliis, tum a semet ipso distinctas ceu Synopsis Supplementum, exposuit. vol. **5** fasc. 2. — pag. 135.
— Hym. Jav. 1861.	— 4. Hymenophyllaceae javanicae sive Descriptio Hymenophyllacearum Archipelagi indici, iconibus illustrata. Amstelaedami. 4^0. 52 Tab. (Verhand. d. koningk. Akad. van Wetensch. Amsterdam. 9^6. 1861).
— Versl. Akad. Wet. Amsterd. **11**. 1861.	— 5. Eerste Bijdrage tot de Kennis der Hymenophyllaceae. — pag. 300. (Verslagen en Mededeelingen der koninkl. Akad. van Wetenschappen. Afdeeling Natuurkunde).
— Ann. sc. nat. IV. **15**. 1861.	— 6. Hymenophyllaceae (Novae Caledoniae). — pag. 88.
— Ned. kr. Arch. 5^3. 1863.	— 7. Hymenophyllaceas novas, exposuit. — pag. 135. — vol. **5** fasc. 3.
Bower: Ann. Bot. **18**. 1904.	**F. Bower:** Ophioglossum simplex Ridley. — pag. 205 t. 15.

Brack. Expl. Exp. **16**. 1854.	**Brackenridge:** Filices in Charles Wilkes: U. S. Exploring Expedition. vol. **16**. Philadelphia. 4⁰. Atlas in Folio. 46 Tab.
A. Br. Flora **1839**.	**A. Braun:** 1. Ueber Marsilea. — pag. 297.
— — **1841**.	— 2. Bemerkungen über die Flora von Abyssinien. — pag. 705.
— Amer. Journ. Sci. II. **3**. 1847.	— 3. On the North American Species of Isoetes und Marsilea. — pag. 52.
— Linn. **25**. 1852.	— 4. Marsiliaceae in »Plantae Muellerianae« — pag. 721.
— App. Ind. Sem. Berol. 1854.	— 5. Filicum hybridarum formae novae in horto Berolinensis enatae. — pag. 17.
— Monatster. Berl. Ak. **1863**. 1864.	— 6. Über die Marsiliaceen-Gattungen Marsilia und Pilularia. — pag. 413.
— in Schweinf. Beitr. **1**. 1867.	— 7. Pteridophyta in G. Schweinfurth: Beitrag zur Flora Aethiopiens. Berlin. 4⁰.
— App. Ind. Sem. Berol. 1867. (Ann. sc. nat. V. **9**. 1868).	— 8. Species novae vel minus cognitae, quae in horto regio botanico Berolinensis 1867 coluntur. pag. 3. — etiam in Ann. sc. nat. pag. 379.
— ap. Kuhn, Fil. Afr. 1868.	— 9. Ophioglossaceae, Marsilia ap. Kuhn: Fil. Afr. — pag. 176, 197. vide Kuhn.
— Monatschr. Berl. Akad. **1870**. 1871.	—10. Neuere Untersuchungen über die Gattungen Marsilia und Pilularia. — pag. 653.
——— **1872**. 1873.	—11. Nachträgliche Mittheilungen über die Gattungen Marsilia und Pilularia. — pag. 635.
E. Britt. Bull. Torr. Cl. **15**. 1888.	**E. G. Britton:** 1. Pteridophyta in »An Enumeration of the plants collected by Dr. H. H. Rusby in South America. 1885-1886. III.« — pag. 247.
— *Mem. Torr. Cl. **4**. 1895.	— 2. apud Rusby: An enumeration of the plants collected in Bolivia by Miguel Bang. II. — pag. 271.
— Bull. Torr. Cl. **24**. 1897.	— 3. A. Revision of the North American Species of Ophioglossum. — pag. 545 t. 318-319.
Brockm. Arch. Ver. Fr. Naturg. Meckl. **1863**.	**H. Brockmüller:** 4. Beiträge zur Kryptogamen-Flora Mecklenburgs. — pag. 162.
A. Brongn. Bull. Soc. Philom. **1821**.	**A. Brongniart:** 1. Description d'un nouveau genre de Fougères, nommé Ceratopteris. — pag. 184. cum tab.
— Dict. class. 1824-1831.	— 2. [Plusieurs genres de fougères. vide Bory nr. 3].
— Voy. Vénus 1846.	— (A.Br.) 3. in Abel du Petit Thouars: Voyage autour du Monde sur la Frégate la Vénus. Atlas de Botanique. Paris. Fol. t. 5-6 (sine descr.?).
R. Br. Prod. Fl. N. Holl. 1810.	**R. Brown:** 1. Prodromus Florae Novae-Hollandiae et Insulae Van Diemen. London. 8⁰.
— Hort. Kew. ed. II. **5**. 1813.	— 2. vide Aiton.
— Tr. Linn. Soc. **11**. 1815.	— 3. On Woodsia, a new Genus of Ferns. — pag. 170 t. 11.

L. v. Buch: Abh. Akad. Berlin **1816-17**. 1819.	**L. v. Buch:** 1. Allgemeine Uebersicht der Flora auf den canarischen Inseln. — pag. 337.
— Beschr. Canar. Ins. 1825.	— 2. Physicalische Beschreibung der canarischen Inseln. Berlin. 4^0.
Buckley: Amer. Journ. Sci. **45**. 1843.	**S. B. Buckley:** Description of some New Species of Plants. — pag. 170.
Burck: Ann. Jard. Buit. **4**. 1884.	**W. Burck:** 1. Contributions of the Fern-Flora of Borneo. — pag. 88 t. 7.
— Recueil Trav. Bot. Néerl. **1**. 1904.	— 2. Sur quelques formes du Polystichum aculeatum de l'Archipel Malais et sur un caractère spécial et peu connu de cette espèce. — pag. 33.
Burm. Fl. Ind. 1768.	**N. L. Burmann:** Flora Indica . . . nec non Prodromus Florae capensis. Lugduni Batavorum. 4^0.
Carm. Tr. Linn. Soc. **12**. 1818.	**D. Carmichael:** Some Account of the Island of Tristan da Cunha and of its Natural Production. — pag. 483.
Carrière: Rev. Hort. **1886**.	**Carrière:** Polypodium Picoti. — pag. 206 f. 62.
Carr. in *Seem. Fl. Vit. 1873.	**W. Carruthers:** 1. An Enumeration of the Ferns collected by Dr. Seemann in Fiji. — in Seemann: Flora vitiensis. London. 4^0. — (t. A. Gepp).
— JoB. **1900**.	— 2. The Nomenclature of Platycerium. — pag. 123.
— Cat. Welw. Pl. 2^2. 1901.	— 3. Pteridophyta in Catalogue of the African Plants collected by Dr. Friedrich Welwitsch in 1853-61. vol. **2** part 2. Cryptogamia. London. 8^0.
Cav. *Anal. Hist. Nat. **1**. 1799.	**A. J. Cavanilles:** 1. Diez especies nuevas del género Acrostichum. — pag. 101. — (t. A. Gepp). (Anales de Historia Natural. Madrid).
— * ——— —	— 2. Nuevos caractéras genéricos de los Helechos por Smith (traducidos y aumentados con los géneros Tectaria y Oleandra. — pag. 108. — (t. A. Gepp).
— *Anal. Cienc. **4**. 1801.	— 3. De las Plantas que el ciudadano Augusto Broussonet colectó en las costas septentrionales de la Africa y en las Islas Canarias. — pag. 54. (Anales de ciencias naturales. Madrid). — (t. A. Gepp).
— * ——— —	— 4. Del género Ugena. — pag. 249. — (t. A. Gepp).
— Icon. **6**. 1801.	— 5. Icones et descriptiones plantarum, quae aut sponte in Hispania crescunt, aut in hortis hospitantur. Matriti. Fol. t. 592-595.
— Descr. 1802.	— 6. Descripcion de las plantas que D. Antonio Josef Cavanilles demostró en las lecciones publicas del año 1801 . . . Madrid. 8^0.
— Descr. 1803.	— 7. (Continuatio operis nr. 6) Géneros y Especies de plantas demostradas en las lecciones publicas del año de 1802. Madrid. 8^0.
— *Anal. Cienc. **7**. 1804.	— 8. Del Macrocnemo y de algunas plantas descubiertas por los Españoles. — pag. 55 t. 48. — (t. A. Gepp).
— *Hort. Madrit. ?	— 9. Elenchus horti Madritensis — ?

Ces. Atti Acad. Napoli 7^{8}. 1876.	**V. Cesati:** 1. Felci e specie nei Gruppi affini raccolte a Borneo dal Signor Odoardo Beccari. — Atti Acad. sci. fis. e mat. Napoli. vol. 7 nr. 8 t. 1-4.
— Rend. Acad. Napoli **16**. 1877.	— 2. Prospetto delle Felci raccolte dal Signor O. Beccari nella Polinesia durante il suo secondo viaggio di esplorazione in que' mari — pag. 23. — (conf. Beccari no. 2).
Chauvin: Mém. Soc. Linn. Calvados **1825**.	**M. Chauvin:** Essai sur les Fougères du Calvados. — Mémoires de la Société Linnéenne du Calvados. Année 1825. — pag. 384.
Cheesem. Tr. N. Zeal. Inst. **10**. 1878.	**T. F. Cheeseman:** 1. Description of a new Species of Polypodium. — pag. 356.
——— **23**. 1891.	— 2. Further Notes on the Three Kings Islands. — pag. 408.
— Tr. Linn. Soc. II. Bot. **6**. 1903.	— 3. The flora of Rarotonga, the chief island of the Cook Group. — pag. 261.
Chiovenda: Ann. di Bot. **1**. 1903.	**E. Chiovenda:** Sul nome di alcune Felci nostrali. — Annali di Botanica. Roma. 8⁰. — pag. 208.
Christ: Journ. de Bot. **4**. 1890.	**H. Christ:** 1. Une nouvelle Fougère du Tonkin français: Cyathea Bonii Christ. — pag. 410.
— *Ber. Schweiz. Bot. Ges. **3**. 1893.	— 2. Les différentes formes de Polystichum aculeatum (L. sub Polypodio), leur groupement et leur dispersion, y compris les variétés exotiques. — pag. 26.
— Engl. Jahrb. **19** Beibl. 1894.	— 3. Trichomanes orbiculare n. sp. — pag. 26.
— Journ. de Bot. **8**. 1894.	— 4. Une liste de Fougères du Tonkin français. — pag. 149.
— Verhdl. Nat. Ges. Basel **11**. 1894-97.	— 5. Filices Sarasinianae. I. pag. 1. 1894. — II. pag. 198 t. 2. 1895. — III. pag. 221 t. 3. 1895. — IV. pag. 421. 1896.
— Ann. Jard. Buit. **12**. 1895.	— 6. Ueber einige javanische Arten von Diplazium. — pag. 217.
— Bot. Gaz. **20**. 1895.	vide Donnell Smith.
— Bull. Soc. bot. Belg. **33**². 1895.	— 7. Fougères nouvelles ou peu connues. — pag. 92.
— Ann. Jard. Buit. **13**. 1896.	— 8. Zur Farn-Flora der Sunda-Inseln. — pag. 90.
— Bull. Boiss. **4**. 1896.	vide Bommer et Christ.
— ——— —	— 9. Filices Faurieanae. Fougères recueillies par le Père Urban Faurie . . . dans les différentes Iles de l'Archipel Japonais. — pag. 664. (Continuationes vide nr. 23, 34, 41, 52).
— Bull. Soc. bot. Belg. **35**. 1896.	vide Bommer et Christ.
— Engl. Jahrb. **23**. 1896.	—10. Pteridophyten apud Reinecke: Die Flora der Samoa-Inseln. — pag. 333 t. 4-5.
— Hedwigia **35**. 1896.	vide Schenck.

Christ: Farnkr. 1897.	**H. Christ:** 11. Die Farnkräuter der Erde. Beschreibende Darstellung der Geschlechter und wichtigeren Arten der Farnpflanzen mit besonderer Berücksichtigung der exotischen. Mit 292 Abbildungen. Jena. 8⁰.
— Ann. Jard. Buit. **15.** 1897.	—12. Die Farnflora von Celebes. — pag. 73 t. 13-17. (Revisio operis n. 5).
— Bull. Soc. bot. Belg. **36²**. 1897.	—13. Filices in Durand et de Wildeman: Matériaux pour la flore du Congo. — pag. 92.
— Engl. Jahrb. **24**. 1897.	vide Krug.
— Nu. Giorn. Bot. It. n. s. **4**. 1897.	—14. in Baroni et Christ: Filices plantaeque filicibus affines in Shen-si septentrionali Prov. Imperii sinensis a Rev. patre Josepho Giraldi collectae. — pag. 86 t. 1-3.
— Bull. Soc. bot. It. **1898**. **1900**. **1901**.	Idem titulus. Manipulus alter. **1898**. pag. 27. tertius. **1898**. pag. 182. quartus. **1900**. pag. 260. quintus. **1901**. pag. 288.
— Bull. Boiss. **6**. 1898.	—15. Filices insularum Philippinarum. Collections de M. A. Loher. — pag. 127, 189 t. 2-4.
— ——————— —	—16. Filices novae. — pag. 835.
— ——————— — **7**. 1899.	—17. Fougères de Mengtze, Yunnan méridional (Chine). **6**. pag. 860, 956; **7**. pag. 1 t. 1.
——— **6**. 1898.	—18. Fougères recueillies dans le bassin inférieur de l'Amazone par Dr. J. Huber à Para. — pag. 991. Etiam in Boletim do Museu Paraense (Museu Goeldi). **3**. 1900-1902. pag. 60 sub titulo: Fetos do Amazonas inferior e de algumas regiões limitrophes, colleccionados pelo Dr. J. Huber).
— ——— **6** App. I. 1898. II. **1**. 1901. II. **3**. 1903.	—19. Filices ap. Chodat: Plantae Hasslerianae, soit énumération des plantes récoltées au Paraguay par le Dr. Émile Hassler, d'Aarau (Suisse). **6**. Appendix I. pag. 4. II. **1**. pag. 425. II. **3**. pag. 612.
— Bull. sci. Françe et Belg. **28**. 1898.	—20. apud Billet: Note sur la Flore du Haut-Tonkin (Région de Cao-Bang). Cryptogames vasculaires. — pag. 255 t. 12.
— Mon. (Elaph.) 1899.	—21. Monographie des Genus Elaphoglossum in Neue Denkschr. d. allg. schweiz. Ges. f. d. ges. Naturwiss. **36**. 84 Figg. 4 Tab.
— Ann. Cons. Jard. bot. Genève **3**. 1899.	—22. Énumération de quelques Fougères de l'herbier Delessert. — pag. 29.
— Bull. Boiss. **7**. 1899.	—23. Filices Faurieanae. — pag. 817.
— apud Dusén, Sv. Exp. Magell. 1899.	vide Dusén.
— in Warb. Monsunia **1**. 1900.	—24. Filicineae in O. Warburg: Monsunia Beiträge zur Kenntniss der Vegetation des Süd- und ostasiatischen Monsungebietes. Leipzig. Fol.
— Farnkr. d. Schweiz. 1900.	—25. Die Farnkräuter der Schweiz. Beiträge zur Kryptogamenflora der Schweiz. **1**. Heft 2. Bern. 8⁰.

Christ: *in Burnat: Mat. p. fl. Alp. Marit. 1900.	**H. Christ:** 26. Les Fougères des Alpes Maritimes in E. Burnat: Matériaux pour servir à l'histoire de la flore des Alpes Maritimes. Genève.
— in Schwacke: Pl. Nov. Mineiras **2**. 1900.	—27. Specilegium pteridologicum austro-brasiliense in Schwacke: Plantas Novas Mineiras. fasc. **2**. p. 10. Cidade de Minas. 8⁰. *Obs.* Vide Revisionem hujus operis nr. 39.
— Ann. Cons. Jard. bot. Genève **4**. 1900.	—28. Sur quelques Fougères de l'herbier Delessert. — pag. 207.
— Bol. Mus. Para **3**. 1900-1902.	vide nr. 18.
— Bot. Tidsskr. **24**. 1901.	—29. Pteridophyta in J. Schmidt: Flora of Koh Chang. — pag. 102.
— Bull. Boiss. II. **1**. 1901.	—30. Fougères collectées par M. Le Dr. J. Huber au Bas-Ucayali et au Bas-Huallaga (Alto-Amazonas) en octobre-décembre 1898. — pag. 65. Etiam in Boletim do Museu Goeldi **4**. 1905. pag. 512 (apud Huber: Materiaes para a Flora Amazonica).
— ——— —	—31. Reliquiae Weinlandianae. — Eine Pteridophyten-Sammlung aus Deutsch Neu-Guinea leg. † Dr. C. A. F. Weinland. — pag. 445. c. fig.
— ——— —	—32. Filices in De Candolle: Plantae Madagascarienses ab Alberto Mocquerysio lectae. — pag. 552.
— ——— —	—33. Elaphoglossum (Microstaphyla) Bangii. Une Fougère ancestrale. — pag. 588. c. fig.
— ——— —	—34. Filices Faurieanae. — pag. 1013.
— ——— —	—35. Une Fougère nouvelle. Aspidium (Polystichum) Münchii n. sp. — pag. 1120.
— Bull. Soc. bot. It. **1901**.	—36. E. Baroni et Christ: Filices Setciouenses a Rev. patre U. Scallan collectae in Se-Tciouen, provincia imperii sinensis, in monte Uo-Mi-San prope Tcen-To-Sen, anno 1899. — pag. 293.
— Prim. Fl. Costaric. **3**. 1901.	—37. Filices (2 d. Mémoire) in H. Pittier: Primitiae Florae Costaricensis. San José de Costa Rica. 8⁰. (vide Bomm. et Christ nr. 2).
— Bull. Ac. Géogr. Bot. Mans (9). **1902**.	—38. Filices Bodinierianae, déterminées et décrites. — pag. 189. figg.
— Bull. Boiss. II. **2**. 1902.	—39. Specilegium pteridologicum austro-brasiliense. — pag. 313, 361, 545, 633, 689. (Revisio op. n. 27).
— ——— —	—40. Filices novae. — pag. 561.
— ——— —	—41. Filices Faurieanae. IV. — pag. 825.
— ——— II. **3**. 1903.	—42. Fougères de Madagascar, récoltées en 1894 par le Docteur C.-J. Forsyth-Major. — pag. 31.
— ——— —	—43. Filices novae. — pag. 147.
— ——— —	—44. Filices Chinae centralis leg. Wilson. — pag. 508.
— Hedwigia **42**. 1903.	—45. Die Varietäten und Verwandten des Asplenium Ruta muraria L. — pag. 153 t. 5-8.

Christ: *Ann. Mus. Congo Bot. 5. **1**. 1903.	**H. Christ:** 46. Filices in de Wildeman: Études de systématique et de géographie botanique sur la flore du Bas- et du Moyen-Congo. — pag. 1. tab. 1. (t. H. Christ).
— Ann. Cons. Jard. bot. Genève **7-8**. 1904.	— 47. Quelques remarques concernant une collection de Fougères du Bhotan, recoltées par W. Griffith et acquisé par l'Herbier Delessert en 1856. — pag. 330.
— Ann. Jard. Buit. II. **4**. 1904.	—48. Zur Farnflora von Celebes. — pag. 33.
— Bull. Ac. Géogr. Bot. Mans 1904 (**11**).	—49. Filices Cavalerianae. — pag. 105. figg.
— ——— —	—50. Les fougères de la Galicie Espagnole. — pag. 76.
— Bull. Boiss. II. **4**. 1904.	—51. Loxomopsis costaricensis nov. gen. et spec. — pag. 393 t. 1.
— ——— —	—52. Filices Faurieanae. — pag. 610.
— ——— — II. **5**. 1905.	—53. Primitiae Florae Costaricensis. Filices et Lycopodiaceae III. **4**. pag. 936, 957, 1089. **5**. pag. 1, 248. — Conf. n. 37 et Bomm. et Christ n. 2.
— Ann. Jard. Buit. II. **5**. 1905.	—54. Filices borneenses. Fougères receuillies par les expéditions de Messieurs Nieuwenhuis et Hallier dans la partie équatoriale de Bornéo. — pag. 62 t. 3.
— Ark. för Bot. 4^{12}. 1905.	—55. Über die australen Polystichum-Arten.
— Bol. Mus. Goeldi **4**. 1905.	vide nr. 30.
— Bull. Boiss. II. **5**. 1905.	—56. Filices Mexicanae. I. German Munch. — pag. 725.
— Bull. Soc. Fr. **52** Mém. I. 1905.	—57. Les collections de Fougères de la Chine au Muséum d'histoire naturelle de Paris.
— Hedwigia **44**. 1905.	—58. Filices Uleanae Amazonicae. — pag. 359.
— Journ. de Bot. **19**. 1905.	—59. Filices Cadierianae. — pag. 58.
— *Mission E. Laurent 1905.	—60. Filices in de Wildeman: Mission E. Laurent 1903-1904. (fasc. I). Bruxelles 1905. — pag. 1 t. 4.
— in Schum. et Laut. Nachtr. Fl. deut. Südsee 1905.	—61. Filices in K. Schumann et K. Lauterbach: Nachträge zur Flora der deutschen Schutzgebiete in der Südsee. — pag. 34. — Conf. Schum. et Laut.
Christ et Gies. Flora **1899**.	**H. Christ et K. Giesenhagen:** Pteridographische Notizen. — pag. 72. 7 figg.
C. Chr. Bot. Tidsskr. **25**. 1902.	**C. Christensen:** 1. New Ferns from Brazil. — pag. 77.
— ——— **26**. 1904.	— 2. On the American Species of Leptochilus Sect. Bolbitis. — pag. 283 figg.
— ——— —	— 3. A new Elaphoglossum from Brazil. — pag. 299.
Clarke: Journ. Linn. Soc. **15**. 1876.	**C. B. Clarke:** 1. Botanical Notes from Darjeeling to Tonglo. — pag. 116.

Clarke: Tr. Linn. Soc. II. Bot. **1**. 1880.	**C. B. Clarke:** 2. A Review of the Ferns of Northern India, with an Index of the species and 36 Plates. — pag. 425 t. 49-84.
— Journ. Linn. Soc. **19**. 1882.	— 3. Note on two Himalayan Ferns erroneously treated in the »Ferns of Northern India.« — pag. 289.
— ——— **25**. 1888.	— 4. On the Plants of Kohima and Muneypore. — pag. 1 t. 41-44.
— **and Bak.** Journ. Linn. Soc. **24**. 1888.	— **and J. G. Baker:** Notes on the Ferns of Northern India. — pag. 408.
Clute: Fern Bull. 8. 1900.	**W. N. Clute:** 1. Pteris aquilina and its allies in North-America. – pag. 37.
— ——— 8. 1900.	— 2. A new Dryopteris from Jamaica. — pag. 67.
— ——— 8. 1900. **9**. 1901. **10**. 1902. **11**. 1903.	— 3. A list of the fernworts collected in Jamaica. **8**. pag. 64, 89; **9**. 16, 43, 67; **10**. 26, 88; **11**. 54.
Col. *Tasm. Journ. **1**. 1841.	**W. Colenso:** 1. Description of some new Ferns lately discovered in New Zealand. — pag. 375. — Tasmanian Journal of Natural Science. Hobart. 8⁰. — (t. A. Gepp).
— ——— 2. 1844.	— 2. A Classification and description of some newly discovered ferns, collected in the northern Island of New Zealand in the Summer of 1841-42. — pag. 161. — (t. A. Gepp).
— Tr. N. Zeal. Inst. **11**. 1879.	— 3. A Description of two New Zealand Ferns, believed to be new to Science. — pag. 429.
— ——— **12**. 1880. **13**. 1881. **14**. 1882.	— 4. A Description of a few new Plants from our New Zealand Forests. **12**. pag. 359. **13**. pag. 376. **14**. pag. 329.
— ——— **13**. 1881.	— 5. The Ferns of Scinde Island (Napier). — pag. 370.
— ——— **15**. 1883.	— 6. A Description of four new Ferns from our New Zealand Forests. — pag. 304.
— ——— —	— 7. On the large Number of Species of Ferns noticed in a small Area in the New Zealand Forests. — pag. 311.
— ——— **16**. 1884.	— 8. A further Contribution towards making known the Botany of New Zealand. — pag. 325.
— ——— **17**. 1885. **18**. 1886. **19**. 1888. **23**. 1891.	— 9. A Description of some newly-discovered and rare Indigenous*) Plants; being a further Contribution towards the making known the Botany of New Zealand. **17**. pag. 237. **18**. pag. 219. **19**. pag. 271. **23**. pag 381. *) in vol. 18, 19. »Cryptogamic« loco »Indigenous«.
— ——— **19**. 1888.	—10. A few Observations on the Tree-Ferns of New Zealand, with particular Reference to their peculiar Epiphytes, their Habit, and their manner of Growth. — pag. 253 t. 19.
— *——— **20**. 1888.	—11. On newly discovered and imperfectly-known Ferns of New Zealand, with critical observations. — pag. 212. — (t. A. Gepp).

Col. Tr. N. Zeal. Inst. **22**. 1890. — **W. Colenso:** 12. A Description of some Newly-discovered Indigenous Cryptogamic Plants. — pag. 452.

— — **24**. 1892. — —13. A Description of Three Species of newly-discovered New Zealand Ferns. — pag. 394.

— — **25**. 1893. — —14. A Description of a few lately-discovered Rare Indigenous Ferns. — pag. 319.

— **27**. 1895. — —15. A Description of Two New Ferns and One New Lycopodium, lately detected in our New Zealand Forests. — pag. 399.

— — **28**. 1896. — —16. A Description of Two New Ferns. — pag. 615.

— — — —17. A Description of Three Ferns believed to be undescribed, discovered more than Fifty Years ago in the Northern District of New Zealand. — pag. 618.

— — **29**. 1897. — —18. A Description of some New Indigenous New Zealand Forest Ferns. — pag. 414.

— — **31**. 1899. — —19. A Description of some Newly Discovered Indigenous New Zealand Ferns. — pag. 263.

Colla: Mem. Ac. Torino **39**. 1836. — **L. Colla:** Plantae rariores in regionibus chilensibus a clarissimis M. D. Bertero nuper detectae. t. 61-75.

Colm. Enum. crypt. Esp. y Port. **1**. 1867. — **M. Colmeiro:** Enumeracion de las cryptogamas de España y Portugal. Parte I. Acrogenas. Madrid. 8⁰.

Colomb: Compt. rendus **107**. 1888. — **G. Colomb:** Sur la place de quelques Fougères dans la classification. — pag. 1012. — (Comptes rendus des séances de l'Acad. des Sci. Paris).

Copeland: Copeland in Perkins. Fragm. 1905. — **E. B. Copeland:** 1. Ferns in J. Perkins: Fragmenta Florae Philippinae. Contributions to the Flora to the Philippine Islands. fasc. III. Febr. 20. 1905. Leipzig. 8⁰. — pag. 175 t. 4.

— Polyp. Philipp. 1905. — — 2. The Polypodiaceae of the Philippine Islands. — Bureau of Government Laboratories nr. 28. Manila. July 1905.

Corda: Mon. Rhizosp. 1829. — **A. J. Corda:** Monographia Rhizospermarum et Hepaticarum. Heft 1. Prag. 4⁰. Tabb.

Cord. *Bull. Soc. sci. et arts Réunion **1890-91**. — **J. D. Cordemoy:** 1. Flore de l'île de la Réunion. — pag. 129. 5 Tab. St. Denis 1891.

— Fl. Réun. 1895. — — 2. Flore de l'île de la Réunion. Paris. 8⁰.

— Rev. gen. Bot. **8**. 1895. — — 3. Sur la Polyphormisme de l'Asplenium lineatum Sw. — pag. 81 t. 5-6.

Cosentini: Atti Acad. Gioen. **2**. 1827. — **F. Cosentini:** Memoria sull' Acrostichum catanense, pianta ultimamente scoperta ne'contorni di Catania. — pag. 207. c. tab. — (Atti dell' Acad. Gioenia di Sci. Nat. di Catania. 4⁰).

Coss. et Germ. *Fl. Paris 1845. — **E. Cosson et E. Germain:** Flore descriptive et analytique des environs de Paris. Paris. 8⁰.

Coville: Bull. Torr. Cl. **28**. 1901. — **F. V. Coville:** The Home of Botrychium pumicola. — pag. 109 t. 7.

A. Cunn. in Hk. Comp. Bot. Mag. **2**. 1836.	**Allan Cunningham:** Florae Insularum Novae Zelandiae Precursor: or a Specimen of the Botany of the Islands of New Zealand. — pag. 361 t. 31-32.
Currey: JoB. **1863**.	**F. Currey:** On the Nardoo Plant of Australia. — pag. 161 t. 6.
Dav. *Cat. Dav. Herb. 1876. 1879. Suppl. 1883.	**G. E. Davenport:** 1. Catalogue of the »Davenport Herbarium« 1876. Supplement 1883. — Catalogue of the »Davenport Herbarium« of North American Ferns . . . 1879. (Mass. Hort. Soc.). Salem. 8⁰.
— Bull. Torr. Cl. **6**. 1877.	— 2. A new Cheilanthes. — pag. 190.
— *Amer. Natur. **12**. 1878.	— 3. Aspidium spinulosum Swartz, and its varieties. — pag. 707. (American Naturalist).
— ——— 8. 1881.	— 4. A new American Fern. — pag. 61 t. 8.
— ——— **10**. 1883.	— 5. A new Fern (Cheilanthes Pringlei). — pag. 61 t. 34.
— ——— **12**. 1885. **13**. 1886. **15**. 1888.	— 6. Fern Notes. **12**. pag. 21. **13**. pag. 129 t. 58. **15**. pag. 225.
— *Gard. and Forest. **4**. 1891.	— 7. Filices mexicanae. I-IV. — pag. 448, 483, 519. 555 f. 71, 75.
— Bot. Gaz. **19**. 1894. **21**. 1896.	— 8. Filices mexicanae. V. VI. **19**. pag. 390. **21**. pag. 253 t. 18.
— ——— **19**. 1894.	— 9. Two new ferns from New England. — pag. 492.
— Bull. Torr. Cl. **26**. 1899.	— 10. Acrostichum lomarioides Jenm. — pag. 318.
— Fern Bull. **13**. 1905.	— 11. A new type of Aneimia. — pag. 18 c. tab.
Deakin: *Florig. brit. **4**. 1841-48.	**Deakin:** Florigraphia britannica . . . London. 8⁰.
DC. Fl. franç. **5**. 1815.	vide Lam. et DC.
Dcne. Nouv. Ann. Mus. **3**. 1834.	**M. J. Decaisne:** 1. Herbarii Timorensis descriptio. pag. 345. (Nouv. Ann. de Mus. d'hist. nat. Paris. 4⁰).
— Arch. Mus. **2**. 1841.	— 2. Plantes de l'Arabie Heureuse, recueillies par M. P.-E. Botta. — pag. 89. (Arch. de Mus. d'hist. nat. Paris. 4⁰).
— *Jacq. Voy. **4**. 1844.	— 3. J. V. Jacquemont: Voyage dans l'Inde pendant les années 1828-1832. **4**. Botanique par M. J. Decaisne Paris. 4⁰. tabb. — (t. A. Gepp).
Desf. Fl. atlant. **2**. 1800.	**R. L. Desfontaines:** Flora atlantica . . . Parisiis. 4⁰.
Desv. Berl. Mag. **5**. 1811.	**A. N. Desvaux:** 1. Observations sur quelques nouveaux genres de fougères et sur plusieurs espèces nouvelles de la même famille. — pag. 297 t. 7 f. 4-7. — Hoc opus emendatum (?) etiam in:
— *Journ. de bot. appl. **1**. 1813 (sive Journ. de bot. **3**. 1813).	— 2. . . . . . Journal de botanique appliquée à l'agriculture, à la pharmacie, à la médicine et aux arts. 8⁰. (vol. 1-4 = Journal de botanique, rédigé par une société de Botanistes (vol. 1-2. 1808-1809) vol. 3-6).
— *——— **4**. 1814 (sive Journ. de bot. **6**. 1814).	— 3. Sur la Belvisia. — pag. 128.

Desv. *Journ. de bot. appl. **4**. 1814 (sive Journ. de bot. **6**. 1814).	**A. N. Desvaux:** 4. Description de plusieurs espèces du genre Polypode. — pag. 257.
— Prod. 1827.	— 5. Prodrome de la famille des Fougères. In Mém. de la Soc. Linnéenne. Paris. **6**. — pag. 171 t. 7-8.
Dicks. *Fasc. pl. cr. Britt. 1785-1801.	**J. Dickson:** Fasciculi (IV) plantarum cryptogamicarum Britanniae. London. 4^0.
Diels: Nat. Pfl. 1^4. 1899-1900.	**L. Diels:** 1. Loxsoma, Cyatheaceae, Polypodiaceae. — pag. 112-336. 1899. Parkeriaceae, Matoniaceae, Gleicheniaceae, Schizaeaceae, Osmundaceae. — pag. 337-380. 1900; in Engler und Prantl: Die natürlichen Pflanzenfamilien 1 Teil, Abt. 4.
— Engl. Jarhrb. **29**. 1900.	— 2. Die Flora von Central-China. — pag. 169.
— in Schum. et Laut. Fl. deut. Südsee 1900.	vide Schum. et Laut.
— Hedwigia **44**. 1905.	— 3. Die primitivste Form von Lygodium. — pag. 133. Fig.
Döll: Rhein. Fl. 1843.	**J. C. Döll:** Rheinische Flora . . . Frankfurt a. M. 8^0.
Dörfler: Oest. Bot. Zeit. **40**. 1890.	**J. Dörfler:** 1. Beiträge und Berichtigungen zur Gefässkryptogamenflora der Bukowina. — pag. 196, 226, 271.
— ——— **45**. 1895.	— 2. Asplenium Baumgartneri mihi; die intermediäre Form der Hybriden Asplenium septentrionale (L.) Hoffm. × Trichomanes Huds. — pag. 169 t. 9.
Don: Prod. Fl. Nepal. 1825.	**D. Don:** 1. Prodromus Florae Nepalensis . . . . . Londini. 8^0. NB. t. Opiz in Flora **1834**. 591 hoc opus X/1824 editum est.
— Tr. Linn. Soc. **17**. 1837.	— 2. Remarks on some British Ferns. — pag. 435.
Drake: Fl. Polyn. fr. 1893.	**E. Drake del Castillo:** Flore de la Polynésie française. Description des Plantes vasculaires qui croissent spontanément ou qui sont généralement cultivées aux îles de la Société, Marquise, Pomotou, Gambier et Wallis. Paris. 8^0.
Dry. Tr. Linn. Soc. **3**. 1797.	**J. Dryander:** Lindsaea, a new Genus of Ferns. Read XI/1794. — pag. 39 t. 7-11.
Dusén: Sv. Exp. Magell. **3**. 1899.	**P. Dusén:** Die Gefässpflanzen der Magellansländer . . . . in Svenska Expeditionen til Magellansländerna. Wissenschaftliche Ergebnisse. **3**. — pag. 77. — Species novas Christ descripsit.
A. A. Eat. Fern Bull. **7**. 1899.	**A. A. Eaton:** 1. A new species of Botrychium. — pag. 7.
——— **12**. 1904.	— 2. A preliminary list of Pteridophyta collected in Dade County, Florida, during November and December, 1903. — pag. 45.
Eat. in Perry: Narr. Exp. to China **2**. 1856.	**D. C. Eaton:** 1. Filices apud A. Gray: Account of the Botanical Specimens. List of Dried Plants collected in Japan by S. Wells Williams Esq.,

	and Dr. James Morrow. — In Narrative of Expedition of an American Squadron to the China Seas and Japan . . . 1852-54 under the command of Commodore Perry. Washington. 4^0. **2**. pag. 303.
Eat. Amer. Journ. Sci. II. **22**. 1856.	**D. C. Eaton:** 2. On three new Filices from California and Oregon. — pag. 138.
— Proc. Amer. Ac. **4**. 1858.	— 3. Characters on some New Filices from Japan and Adjacent Regions, collected by Charles Wright in the North Pacific Exploring Expedition under Captain John Rodgers. — pag. 110.
— *U. S. and Mex. Bound. Surv. 1859.	— 4. Equisetaceae, Filices etc. in J. Torrey: Botany of the U. S. and Mexican Boundary Survey. — pag. 233.
— Amer. Journ. Sci. II. **27**. 1859.	— 5. Enumeration of Ferns collected by Mr. Charles Wright in Eastern Cuba in 1856-7. — pag. 197.
— Mem. Amer. Acad. n. s. **8**. 1860.	— 6. Filices Wrightianae et Fendlerianae, — nempe in Insula Cuba a Carolo Wright et in Venezuela ab Aug. Fendler, ann. 1854-60, (nonnullis Panamensibus etc. interjectis). — pag. 193.
— *Canad. Naturalist. II. **2**. 1865.	— 7. On the Genus Woodsia. — pag. 89.
— Proc. Amer. Ac. **6**. 1865.	— 8. Filices ap. A. Gray: Characters of some New Plants of California and Nevada, chiefly from the Collections of Professor William H. Brewer . . . ., and of Dr. Charles L. Anderson, with revisions of certain Genera or Groups. — pag. 519.
— Bot. Zeit. **1869**.	— 9. Ein neues Adiantum von Cuba. — pag. 361.
— Proc. Amer. Ac. **8**. 1873.	—10. Characters of New Ferns from Mexico. — pag. 618.
— Bull. Torr. Cl. **4-10**. 1873-1883.	—11. New or little known Ferns of the United States. **4**. 1873. pag. 11, 18. **6**. 1875. pag. 33. **6**. 1876. pag. 71. **6**. 1878. pag. 263. **6**. 1879. pag. 306, 360. **7**. 1880. pag. 62. **8**. 1881. pag. 4, 99, 111. **9**. 1882. pag. 49. **10**. 1883. pag. 26, 101.
— Rep. U. S. Geogr. Surv. West of 100 Merid. **6**. 1879.	—12. Ferns of the Southwest etc. in Report of U. S. Geogr. Survey West of the One Hundredth Meridian. vol. **6**. Botany. 1878. — pag. 299 t. 30. (Prodiit V/1879).
— *Bot. Gaz. **3**. 1878. **5**. 1880.	—13. Fendler's Ferns of Trinidad. **3**. pag. 89; **5**. pag. 109.
— Ferns N. Am. **1**. 1877. **2**. 1880.	—14. The Ferns of North America. Colored Figures and Descriptions of the Ferns (including the Ophioglossaceae) of the United States of America and the British North American Posessions. 4^0. **1**. Salem. **2**. Boston. 81 Tab. col.

Eat. Bull. Torr. Cl. **6**. 1879.	**D. C. Eaton:** 15. A new Hawaiian fern, Hymenophyllum Baldwinii. — pag. 293.
— — —	—16. A new Hawaiian fern, Aspidium Boydiae. — pag. 361.
— Proc. Amer. Ac. **18**. 1883.	—17. Filices ap. S. Watson: List of Plants from Southwestern Texas and Northern Mexico, collected by Dr. E. Palmer in 1879-80. II. — pag. 183.
— Bull. Torr. Cl. **14**. 1887.	—18. Asplenium rhizophyllum, Kunze, var. Biscaynianum, N. Var. — pag. 97 t. 68.
— Proc. Amer. Ac. **22**. 1887.	—19. Filices ap. S. Watson: List of Plants collected by Dr. Edward Palmer in the State of Jalisco, Mexico in 1886. — pag. 461.
— Bull. Torr. Cl. **17**. 1890.	—20. A new Fern. — pag. 215 t. 104.
— Zoë **1**. 1890.	—21. A new Fern from Lower California. — pag. 197 t. 7.
Eat. et Dav. *Contr. U. S. Nat. Herb. **5**. 1897.	**(A. A.?) Eaton et G. Davenport** ap. J. N. Rose: Studies on Mexican and Central American plants. — pag. 138 t. 16.
Ehrh. Beitr. **4**. 1789.	**F. Ehrhart:** Kennzeichen seltener und unbestimmter Pflanzen in Beiträge zur Naturkunde **4**. pag. 42.
Emys: Tr. N. Zeal. Inst. **16**. 1884.	**J. D. Emys:** On the Occurrence of the Fern Botrychium lunaria Sw. (Moonwort) in New Zealand. — pag. 363.
Endl. Prod. Fl. Norf. 1833.	**S. L. Endlicher:** 1. Prodromus Florae Norfolkicae etc. Vindobonae. 8⁰.
— Gen. pl. 1836-1843.	— 2. Genera plantarum secundum ordines naturalis disposita. Accedit suppl. primum. Vindobonae. 4⁰. 1836-40. Suppl. sec. 1842. Suppl. tertium 1843.
Ettingsh. Denkschr. Ak. Wien **22**. 1864. **23**. 1864.	**C. v. Ettingshausen:** 1. Beiträge zur Kenntniss der Flächen-Skelete der Farnkräuter. Mit Darstellungen der Flächen-Skelete auf 24 (und 18) Tafeln in Naturselbstdruck, in Denkschriften d. Akad. d. Wiss. Wien. Mat. et Naturw. Classe. **22**. pag. 37. 24 Tab. **23**. pag. 39. 18 Tab. — (etiam in opere seq.).
— Farnkr. 1865.	— 2. Die Farnkräuter der Jetzwelt zur Untersuchung und Bestimmung der in der Formationen der Erdrinde eingeschlossenen Ueberreste von vorweltlichen Arten dieser Ordnung nach dem Flächen-Skelet bearbeitet. Mit zahlreichen in den Text gedruckten Abbildungen und 180 Tafeln im Naturselbstdruck. Wien. 4⁰.
Fée: Mém.	**A. L. A. Fée:** Mémoires sur la famille des fougères:
— 1 mém. 1844.	— 1. Premier mémoire: Examen des bases adoptées dans la classification des fougères et en particulier de la nervation. 2 Tab.

Fée: Acrost. (2 mém.) 1845.	**A. L. A. Fée:** 2. Deuxième mémoire: Histoire des Acrostichées. 64 Tab. — (1 et 2 mém. in 1 vol. Strasbourg. Fol. 1844-45).
— 3 mém. 1851-52.	— 3. Troisième mémoire: Histoire des Vittariées et Pleurogrammées . . . et
— 4 mém. 1851-52.	— 4. Quatrième mémoire: Histoire des Antrophyées. 3 et 4 mém. in 1 vol. 5 Tab. Paris. Fol.
— Gen. (5 mém.) 1850-52.	— 5. Cinquième mémoire: Genera Filicum. Exposition de la Famille des Polypodiacées (Classe des Fougères). Paris et Strasbourg. 4⁰. 32 Tab. — (prob. anno 1853 prodiit).
— 6 mém. 1853.	— 6. Sixième mémoire: Description de Fougères exotiques rares ou nouvelles, 1me Partie; in Mémoires de la Soc. du Mus. d'Hist. nat. de Strasbourg. **4**. Livr. 2-3. t. 1-8. 4⁰.
— 7 mém. 1857.	— 7. Septième mémoire: Description de Fougères exotiques rares ou nouvelles, 2me partie et
— 8 mém. 1857.	— 8. Huitième mémoire: Description de Fougères exotiques rares ou nouvelles, 3me partie. — 7 et 8 mém. in Mémoires de la Soc. des sci. nat. de Strasbourg. **5**. Livr. 1. 7 mém. pag. 1. t. 9-27. 8 mém. pag. 67. 4⁰.
— *9 mém. 1857.	— 9. Catalogue méthodique des Fougères et des Lycopodiacées du Mexique. 4⁰.
— 10 mém. 1865.	—10. Dixième mémoire: Description de Fougères exotiques rares ou nouvelles. Mém. Soc. sci. nat. Strasbourg. **6**. Livr. 1. pag. 1. t. 28-44. 4⁰.
— 11 mém. 1866.	—11. Onzième et dernier mémoire: Histoire des Fougéres et des Lycopodiacées des Antilles. Paris. 4⁰. 34 Tab.
— Cr. vasc. Br. **1**. 1869. **2**. 1872-73.	—12. Cryptogames vasculaires du Brésil. Paris. 4⁰. 108 Tab.
Field: *Ferns N. Zeal. 1890.	**H. C. Field:** 1. The Ferns of New Zealand and its immediate Dependencies, with directions for their Collection and Cultivation. London. 4⁰. 29 Tab. — (t. A. Gepp).
— Tr. N. Zeal. Inst. **27**. 1895.	— 2. On the Growth of Ferns, and on a New Fern now first reported. — pag. 401.
Field. et Gardn. Sert. Pl. 1844.	**H. B. Fielding et G. Gardner:** Sertum Plantarum, or Drawings and Descriptions of rare and undescribed Plants from the Author's Herbarium. London. 8⁰. tabb.
Fisch. et Mey. *Ind. Sem. ht. Petr. 1833. Ann. sc. nat. II. **5**. 1836. Linn. **10**. 1836.	**F. E. L. Fischer et C. A. Meyer:** 1. Animadversiones botanicae nonnullae novarumque aut non recte cognitarum plantarum diagnoses. (etiam impr. in Ann. sc. nat. pag. 120 et Linn. Litteratur. pag. 83).
— Bull. Soc. Nat. Moscou 1838.	— 2. in Hohenacker: Enumeratio plantarum Provinciae Talysch et regionum adjacentium. — pag. 239.

Forbes: JoB. **1888.**	**H. O. Forbes:** A new Fern from New Guinea (Polypodium Annabellae). — pag. 33 t. 280.
Formánek: Oest. Bot. Zeit. **38.** 1888.	**E. Formánek:** Beitrag zur Flora von Bosnien und Hercegovina. — pag. 240.
Forsk. Fl. Aeg.-arab. 1775.	**P. Forskål:** Flora aegyptiaco-arpbica, sive descriptiones plantarum quas per Aegyptum inferiorem et Arabiam Felicem detexit, illustravit. Hauniae. 4°.
Forst. Pl. escul. 1786.	**G. Forster:** 1. De plantis esculentis insularum oceani australis. Commentatio botanica. Berolini. 4°.
— Prod. 1786.	— 2. Florulae Insularum australicum Prodromus. Gottingae. 8°.
— Comm. Soc. Reg. Gott. 9. 1789.	— 3. Plantae Atlanticae ex insulis Madeira, Sti Jacobi, Adscensionis, Stæ Helenae et Fayal reportatae. — pag. 46.
Fourn. Bull. Soc. Fr. **14.** 1867.	**E. Fournier:** 1. Sur la valeur du genre Aconiopteris. — pag. 261.
— ——— **15.** 1868.	— 2. Sur les Hyménophyllées recueillies dans l'Amérique centrale par MM. Ch. Wright, Fendler et Th. Husnot. — pag. 143.
— ——— **16.** 1869.	— 3. Sur les fougères de la Nouvelle-Calédonie. — pag. 389, 422. (Conf. nr. 9).
— ———	— 4. Sur deux Pellaea nouveaux. — pag. LXVIII.
— ——— **17.** 1870.	— 5. Sur deux fougères nouvelles du Nicaragua. — pag. 236.
— Mex. pl. **1.** 1872.	— 6. Mexicanas plantas nuper a collectoribus expeditionis scientificae allatas aut longis ab annis in herbario Musei Parisiensis depositas praeside J. Decaisne, enumerandas curavit E. Fournier. Pars Prima: Cryptogamia. Parisiis. Fol. 6 Tab.
— Bull. Soc. Fr. **19.** 1872.	— 7. Sur les Hymenophyllées recueillies dans l'Amérique centrale par MM. Ch. Wright, Fendler et Th. Husnot. — pag. 239.
— ———	— 8. Sertum Nicaraguense. — pag. 247.
— Ann. sc. nat. V. **18.** 1873.	— 9. Filices Novae-Caledoniae. Enumeratio monographica. — pag. 253.
— Bot. Zeit. **1873.**	—10. Einige neue Farnpflanzen aus Nicaragua. — pag. 6.
— Compt. rendus **81.** 1875.	—11. Sur les Fougères et les Lycopodiacées des îles Saint-Paul et Amsterdam. — pag. 1139.
— L'Ill. hort. **23.** 1876.	—12. Fougères nouvelles introduites par M. J. Linden. — pag. 99. (L'Illustration horticole).
— Bull. Soc. Fr. **27.** 1880.	—13. Sur le Genre Bommeria. — pag. 327.
Franch. Nouv. Arch. Mus. II. **7.** 1883. **10.** 1887.	**A. Franchet:** 1. Plantae Davidianae ex sinarum imperio. **7.** pag. 55. **10.** pag. 33.
— Bull. Soc. Fr. **32.** 1885.	— 2. Plantes de Yunnan récoltées par M. L'abbé Delavay. — pag. 28.
Franch. et Sav. Enum. pl. Jap. **2.** 1876-1879.	**A. Franchet et L. Savatier:** Enumeratio plantarum in Japonia sponte crescentium. Parisiis. 8°. (Enumeratio filicum 1876 (vel 1877) prodiit, diagnoses spec. nov. 1879).

Fr. Novit. ed. II. 1828.	**E. Fries:** 1. Novitiae Florae Suecicae. Londini Gothorum. 8⁰.
— Novit. Mant. **1-3.** 1832-1842.	— 2. Continuatio sistens Mantissam I. Lundae 1832. II. 1839. III. 1842. Upsalae. 8⁰.
— Summ. Veg. **1.** 1845.	— 3. Summa Vegetabilium Scandinaviae etc. Holmiae et Lipsiae. 8⁰.
French: *South. Sci. Record **1-2.** 1881-82.	**C. French:** Victorian Ferns and their Habitats; in »The Southern Science Record«. Melbourne. **1.** 1881. pag. 2-4, 17-21, 33-36, 45-52, 65-68, 189-192, 207 208. **2.** 1882. pag. 31-34, 56-57, 173-775, 205-208. — (t. A. Gepp).
Fritsch: Bull. Boiss. II. **1.** 1901.	**K. Fritsch:** Beitrag zur Flora von Angola. Bearbeitung einer von E. Dekindt aus Huilla an das botanische Museum der k. k. Universität in Wien eingesendeten Pflanzenkollection. — pag. 1082.
Futó: Hedwigia **44.** 1905.	**Mihály Futó:** Polypodium vulgare L. und Polypodium vulgare *γ* serratum Willd. — pag. 106 t. 3.
Gammie: Journ. Linn. Soc. **29.** 1893.	**G. Gammie:** On Sikkim Tree Ferns. — pag. 482.
Gandog. Oest. Bot. Zeit. **31.** 1881.	**M. Gandoger:** 1. Pugillus plantarum novarum vel minus recte cognitarum. — pag. 18.
— Fl. Eur. **27.** 1891.	— 2. Flora Europae. — *Obs.* Numerosissimas species in indice non enumeravi.
Gardn. Lond. JoB. **1.** 1842.	**G. Gardner:** 1. Description of Trochopteris, a new Genus of Ferns. — pag. 73 t. 4.
— ———— —	— 2. Observations on the Genus Hemitelia of Mr. R. Brown. — pag. 438 t. 12.
— ———— —	— 3. List of Ferns from the neighbourhood of Rio de Janeiro, collected in 1836. — pag. 545.
— *Calc. Journ. **7.** 1847.	— 4. Contributions to a Flora of Ceylon. — pag. 1.
Gaspar. *Progr. Sci. Napoli **4.** 1833.	**G. Gasparrini:** 1. Notizie interno ad alcune piante della Lucania. — pag. 161. (Il Progresso dell. Scienze, Litt. ed. Arti. Napoli. 8⁰.
— *Rend. Acad. Napoli **4.** 1845.	— 2. Descrizione di una nuova spezie di Felce (A. brachyphyllum) appartenenti al genere Asplenium. — pag. 107. (Rendic. del. Real. Acad. Sci. Napoli).
Gaud. Ann. sc. nat. **3.** 1824.	**C. Gaudichaud-Beaupré:** 1. Description de quelques nouveaux genres de Plantes recueillies dans le voyage autour du monde, sous les ordre du capitaine Freycinet. — pag. 507.
— ———— **5.** 1825.	— 2. Rapport sur la Flore des îles Malouines. — pag. 89.
— Freyc. Voy. Bot. 1826 29.	— 3. Botanique in Freycinet: Voyage autour du Monde . . . sur l'Uranie et la Physicienne. Paris. 4⁰. Atlas in Fol. t. 3-21. *Obs.* In Indice ubique annum 1827 scripsi, sed sec. JoB. **1901.** 206 hoc opus prodiit: pag. 1-88: 1826; pag. 89-216: 1827; pag. 217-360: 1828; pag. 361-464: 1829; pag. 465-522: 1830.

Gaud. Voy. Bonite Bot. 1846 (-49?). — **C. Gaudichaud-Beaupré:** 4. Voyage autour du monde . . . sur la corvette la Bonite. Botanique. Atlas. Paris. Fol. — Conf. JoB. **1901**. 206. (—* Explication et description des planches de l'Atlas par C. d'Alleizette. 1866).

Gay: Fl. Chil. **6**. 1853. — **C. Gay:** Historia fisica y politica de Chile. Botanica. Flora Chilena. Paris. 8⁰. — (Filices Fée et Remy verisimiliter elaboraverunt).

Gies. Flora **1890**. — **K. Giesenhagen:** 1. Die Hymenophyllaceen. — pag. 411. t. 14-17.

— — **1892**. Ergänzbd. — 2. Ueber hygrophile Farne. — pag. 157. Fig.

— Niph. 1901. — 3. Die Farngattung Niphobolus. Eine Monographie. Jena. 8⁰. Fig.

Gilb. Bull. Torr. Cl. **23**. 1896. — **B. D. Gilbert:** 1. A new Gymnogramme with Remarks on some other Venezuelan Ferns. — pag. 448.

— ——— **24**. 1897. — 2. Two new Bolivian Ferns of the Miguel Bang Collection. — pag. 258.

— ——— — — 3. Three new Ferns from Jamaica. — pag. 259.

— ——— **25**. 1898. — 4. Revision of the Bermuda Ferns. — pag. 593.

— ——— **26**. 1899. — 5. Two new Polypodia from New Zealand. — pag. 316.

— Fern Bull. **8**. 1900. — 6. Two new Ferns from Jamaica. — pag. 62. Fig.

— ——— **9**. 1901. — 7. A new species of Asplenium. — pag. 53.

— *Amer. Botanist **4**. 1903. — 8. A New Fern from Bermuda. — pag. 86. — Vide etiam Fern Bull. **11**. 77. 1903 (»Asplenium muticum«).

Gilib. *Exerc. phyt. 1792. — **J. E. Gilibert:** Exercitia phytologica . . . Lugduni Gallorum. 8⁰.

Gleditsch: Syst. Pl. 1764. — **J. G. Gleditsch:** Systema Plantarum a staminum situ. Berolini. 8⁰.

Gmel. Syst. nat. **2**². 1791. — **J. F. Gmelin:** Linné: Systema naturae ed. XIII. aucta, reformata. Lipsiae. 8⁰.

Gmel. Nov. Comm. Ac. Petr. **12**. 1768. — **S. G. Gmelin:** Observationes et descriptiones botanicae. — pag. 508 t. 8-13.

Goebel: Flora **1896**. — **E. Goebel:** Archegoniatenstudien 8. Hecistopteris, eine verkannte Farngattung. — pag. 67.

Goldie: Edinb. Phil. Journ. **6**. 1822. — **J. Goldie:** Description of some new and rare Plants discovered in Canada in the year 1819. — pag. 319.

Goldm. Nova Acta **19**. Suppl. I. 1843. — **J. G. Goldmann:** Filices in Meyen: Beiträge zur Botanik. — pag. 451.

Goodd. Bot. Gaz. **33**. 1902. — **L. N. Goodding:** Rocky Mountain plant studies. — pag. 66.

Gouan: Ill. obs. bot. 1773. — **A. Gouan:** Illustrationes et observationes botanicae . . . Tiguri. Fol.

A. Gray: Manual 1848. — **Asa Gray:** 1. Manual to the Botany of the Northern United States. Boston & Cambridge. 8⁰.

— Amer. Journ. Sci. II. **15**. 1853. — 2. On the Discovery of two species of Trichomanes in the State of Alabama, one of which is new. — pag. 324.

Gray: *Nat. Arr. br. pl. **2**. 1821. — **S. F. Gray:** A natural arrangement of british plants . . . London. 8⁰. — (t. A. Gepp).

Greene: *Pittonia **4**. 1900.	**E. L. Greene:** Necker's genera of ferns I. — pag. 103.
Grev. Ann. Mag. Nat. Hist. II. **1**. 1848.	**R. K. Greville:** 1. On two new species of Ferns. — pag. 320 t. 17. (etiam in Tr. Bot. Soc. Edinb. **3**).
— ——— II. **2**. 1848.	— 2. Notice of a new species of Antrophyum. — pag. 10 t. 1. (etiam in Tr. Bot. Soc. Edinb. **3**).
Grev. et Hk. in Hk. Bot. Misc. **3**. 1833.	— **et W. J. Hooker:** 1. Enumeratio filicum. — pag. 216.
— ——— —	— 2. Descriptions of two new Species of Peruvian Ferns. — pag. 384 t. 111-112. vide Hk. et Grev.
Griff. Asiat. Research. **19**. 1836. } JoB. **3**. 1840. }	**W. Griffith:** 1. Description of one species of Kaulfussia. As. Res. pag. 94 t. 18; JoB. pag. 375 t. 11-12.
— Calc. Journ. **4**. 1844.	vide Roxburgh.
— ——— **5**. 1845.	— 2. On Azolla and Salvinia. — pag. 227 t. 15-20. (Calcutta Journal of Natural History).
— Notulae **2**. 1849.	— 3. Notulae ad plantas Asiaticas part II. On the higher Cryptogamous Plants. Calcutta. 8^0.
— Ic. pl. as. **2**. 1849.	— 4. Icones plantarum Asiaticarum. part II. On the higher Cryptogamous Plants. Calcutta. 4^0. t. 113-138.
Gris. Abh. Ges. Wiss. Gött. **7**. 1857.	**A. Grisebach:** 1. Plantae caribaeae. — pag. 161.
— Fl. br. W. Ind. 1864.	— 2. Flora of the British West Indian Islands. London 8^0.
— Cat. Pl. Cub. 1866.	— 3. Catalogus Plantarum Cubensium exhibens collectionem Wrightianam aliasque minores ex insula Cuba missas. Lipsiae. 8^0.
— Abh. Ges. Wiss. Gött. **19**. 1874.	— 4. Plantae Lorentzianae. Bearbeitung des ersten und zweiten Sammlung argentinischer Pflanzen des Professor Lorentz zu Cordoba. — pag. 49.
— ——— **24**. 1879.	— 5. Symbolae ad Floram argentinam. Zweite Bearbeitung argentinische Pflanzen. Abh. d. phys. Classe. — pag. 3.
Guérin: *Descr. font. Vaucl. 1804; ed. II. 1813.	**J. Guérin:** Description de la fontaine de Vaucluse. Avignon. 12^0.
Guill. Ann. sc. nat. II. **6**. 1836.	**J. B. A. Guillemin:** Zephyritis Taitensis. Enumération des plantes découvertes par les voyageurs, dans les Iles de la Société, principalement dans celle de Taïti. — pag. 297.
Güldenst. Reisen **1**. 1787. **2**. 1791.	**J. A. Güldenstaedt:** Reisen durch Russland . . . St. Petersburg. 4^0.
Guss. *Pl. rar. 1826.	**G. Gussone:** 1. Plantae rariores, quas in itinere per oras Jonii et Adriatici maris . . . collegit. Neapoli. 4^0.
— Fl. Sic. Syn. 2^2. 1844.	— 2. Florae Siculae Synopsis etc. . . . Napoli. 8^0. vol. 2 pars 2.
Halácsy: Verh. zool. bot. Ges. **54**. 1904.	**Halácsy,** E. von: Aspidium lonchitiforme. — pag. 129. 2 figg.

Hall: Nieuwe Verhdl. v. d. Nederl. Inst. **5**. 1836.	**H. C. van Hall:** Polypodium decursive-pinnatum, nova species descripta. (Nieuwe Verhandelingen de eerste Klasse van de konink. Nederland. Instituut van Wetenschapen. Amsterdam. 4⁰). — pag. 204 c. tab.
Hance: Ann. sc. nat. IV. **15**. 1861.	**H. F. Hance:** 1. Symbolae ad floram sinicam. — pag. 220.
— ——— IV. **18**. 1862.	— 2. Manipulus plantarum novarum, potissime chinensium — pag. 217.
— JoB. **1866**.	— 3. Some remarks on the classification of Ferns. — pag. 253.
— Ann. sc. nat. V. **5**. 1866	— 4. Adversaria in stirpes imprimis Asiae orientalis criticas minusve notas interjectis novarum plurimarum diagnosibus scripsit. — pag. 202.
— JoB. **1868**.	— 5. On two new chinese ferns; with some remarks on the genus Woodwardia. — pag. 175.
— — **1869**.	— 6. Notes on the Fern Flora of China. — pag. 234.
— Journ. Linn. Soc. **13**. 1873.	— 7. Notes on some Plants from Northern China. — pag. 74.
— ——— —	— 8. Florae Hongkongensis Supplementum. — pag. 95. — vide Benth. nr. 1.
— JoB. **1874**.	— 9. De nova Asplenii specie. — pag. 142.
— — **1875**.	—10. Diagnosis of two new Chinese Ferns. — pag. 197.
— — **1883**.	—11. Heptadem Filicum Novarum sinicarum. — pag. 267.
— — **1884**.	—12. Ptilopteris, Novum Polypodiacearum Genus. — pag. 138.
— — **1887**.	—13. Spicilegia Florae Sinensis. — pag. 12.
Hanstein: Monatsber. Akad. Berl. **1862-1863**.	**Hanstein:** Über das Nardoo genannten Nahrungsmittels der Urbewohner Australiens, einer Marsilea Frucht, nebst Bemerkungen zur Entwicklung dieser Gattung. — pag. 103.
Haračić: Verh. Zool. bot. Ges. **43**. 1903.	**A. Haračić:** Ueber das Vorkommen einiger Farne auf der Insel Lussin. — pag. 207 t. 3.
Harr. Journ. Linn. Soc. **16**. 1877.	**M. V. Harrington:** The Tropical Ferns collected by Professor Steere in the years 1870-75. — pag. 25.
Hartm. Handb. 1820: ed. IX. 1864.	**C. J. Hartman:** Handbok i Skandinaviens Flora . . . Stockholm. 8⁰.
Hassk. Tijdschr. Nat. Geschr. **10**. 1843.	**J. K. Hasskarl:** 1. Adnotationes de Plantis quibusdam Javanicis nonnullisque Japonicis, haud rite cognitis, e Catalogo Horti Bogoriensis excerptae. Accedunt nonnullae novae species. — pag. 115. — (in Hoev. en de Vriese: Tijdschr. Nat. Geschied.).
— Cat. pl. Bogor. 1844.	— 2. Catalogus plantarum in horto botanico Bogoriensi cultarum alter. Batavia. 8⁰.
— Flora **1847**.	— 3. Zollinger: Observationes phytographicae. — pag. 317. — vide Zoll. nr. 1.

Hassk. JoB. **7**. 1855.	**J. K. Hasskarl:** 4. Observations on the Gleicheniaceae and Cyatheaceae of Java. — pag. 321.
— Obs. Bot. Fil. **1**. 1856. **2**. 1857-58.	— 5. Observationes Botanicae, quas de Filicibus horti Bogoriensis nec non ad montem Gedeh aliisque locis sua sponte crescentibus annis 1855 et 1856. Batavia. 4⁰. — (in *Verhandelingen der Natuurkundige Vereenigung in Nederlandsch Indië: **1**. in vol. **1**. **2**. in vol. **3**.)
Hedw. Fil. Gen. et. Sp. 1799-1803.	**J. Hedwig:** Filicum Genera et Species recentiori Methodo accomodatae analytice descriptae. Lipsiae Fol. 24 tab. col.
Heller: Minn. Bot. St. **1**. 1897	**A. A. Heller:** Observations on the ferns and flowering plants of the Hawaiian Islands. — pag. 760. t. 42-43. — (Species novas Underwood descripsit).
Hemsl. *Biol. Centr. Amer. **3**. 1885-86. **4**. 1887.	**W. B. Hemsley:** 1. Biologia Centrali-Americana, or, Contributions to the Knowledge of the Fauna and Flora of Mexico and Central America. Botany. London. 4⁰. **3**. pag. 569-664. 1885. pag. 665-711. 1886. **4**. pag. 115 (Costa Rica Ferns T. T. Cooper leg.). — (t. A. Gepp).
— Chall. Voy. Bot. **1**. 1885.	— 2. Voyage of H. M. S. Challenger. Botany. London. Fol. 9 Tab. Fil. — Insular Floras. part I. Bermuda. part II. Ascension. St. Helena. Tristan d'Acunha. Prince Edward Group. The Crozets. Kerguelen Island. Amsterdam and St. Paul Islands. part III. Juan Fernandez. Arrou and Ki Islands.
— Journ. Linn. Soc. **30**. 1895.	— 3. The Flora of the Tonga or Friendly Islands. — pag. 158.
Hériband *1880.	**Hériband:** Note sur une nouvelle espèce de Fougère. Riom 1880. (conf. Le Grand).
Heufl. Verh. zool. bot. Ges. **6**. 1856.	**L. v. Heufler:** Asplenii species Europaeae. Untersuchungen über die Milzfarne Europas. — pag. 235. 4 Tab.
Hew. Mag. Nat. Hist. II. **2**. 1838.	**R. Heward:** 1. Some Observations on a Collection of Ferns from the Island of Jamaica. — pag. 453. — (Diagnoses nov. spec. etiam in Linn. **13**. Litteraturber. 66. 1839).
— Lond. JoB. **1**. 1842.	— 2. On A. Cunningham's plants from Norfolk Island. — pag. 121 nota.
Hier. in Engl. Pflw. Ostafr. C. 1895	**G. Hieronymus:** 1. Pteridophyta in A. Engler: Die Pflanzenwelt Ost-Afrikas und der Nachbargebiete. Theil C. Verzeichniss der bis jetzt aus Ost-Afrika bekannt gewordene Pflanzen. Berlin. 8⁰.
— Engl. Jahrb. **22**. 1896.	— 2. Beiträge zur Kenntniss der Pteridophyten-Flora der Argentina und einiger Teile von Uruguay, Paraguay und Bolivien. — pag. 359.
— Notizbl. Bot. Gart. Berl. **2**. 1898.	— 3. Pteridophyta apud K. Schumann: Die Flora von Neu-Pommern. — pag. 82.

Hier. Engl. Jahrb. **28**. 1900.	**G. Hieronymus:** 4. Filicales apud Engler: Die von W. Goetze und Dr. Stuhlmann im Uluguru-gebirge, sowie die von W. Goetze in den Kisaki- und Khutu-Steppe und in Uhehe gesammelten Pflanzen. — pag. 332.
— ——— **34**. 1904-1905.	— 5. Plantae Lehmannianae in Guatemala, Columbia et Ecuador regionibus finitimis collectae, additis quibusdam ab aliis collectoribus ex iisdem regionibus allatis determinatae et descriptae. Pteridophyta. — pag. 418-560: 1904; pag. 561-582: 1905.
— Hedwigia **44**. 1905.	— 6. Polypodiorum species novae et non satis notae. Beschreibungen von neuen Arten und Bemerkungen zu älteren Arten der Gattung Polypodium. I. Mitteilung. — pag. 78.
— —	— 7. Einige Berichtungen zu der Abhandlung: Plantae Lehmannianae etc. (v. n. 5). — pag. 179.
— ——— —	— 8. Aspleniorum species novae et non satis notae. Beschreibungen von neuen Arten und Bemerkungen zu älteren Arten der Gattung Asplenium. I. Mitteilung. — pag. 193 t. 6.
Hill. Fl. Haw. 1888.	**W. F. Hillebrand:** Flora of the Hawaiian Islands. — Heidelberg. 8⁰.
Hitchc. Amer. Journ. Sci. **6**. 1823.	**E. Hitchcock:** Description of a New Species of Botrychium, with a drawing. — pag. 103 t. 8.
Hoffm. Röm. et Usteri, Mag. Bot. **9**. 1790.	**G. F. Hoffmann:** 1. Tabula in qua Συνοπτικῶς Filices explicantur prima. — pag. 1.
— Deutschl. Fl. **2**. 1795.	— 2. Deutschlands Flora oder Botanisches Taschenbuch. Zweyter Theil für das Jahr 1795. Cryptogamie. Erlangen. 12⁰.
— Comm. Soc. reg. Gotting. **12**. 1796.	— 3. Descriptiones et Icones Plantarum. — pag. 22. t. 5. (Classis physicae, pro anno 1793).
Holl: Flora **1830**[1].	**F. Holl:** Verzeichniss der auf der Insel Madeira beobachteten Pflanzen. — pag. 369.
Hombr. et Jacq. Voy. Pôle sud 1852.	**Hombron et Jacquinot:** Cryptogames vasculaires in Voyage au Pôle sud et dans l'oceanie sur les corvettes l'Astrolabe et Zélée ..... Botanique. Paris. Atlas Fol. t. 1-5. (sine descr.).
Hk.fil.	**J. D. Hooker:** The Botany of the Antarctic Voyage of H. M. S. »Erebus« and Terror, in the years 1839-43, under the command of Captain James Clark Ross. London. 4⁰. Partes I-III:
— Fl. antarct. **1**. 1844. **2**. 1847.	— 1. I. Flora antarctica. t. 149-150.
— Fl. N. Zel. **2**. 1854.	— 2. II. Flora Novae-Zelandiae. t. 71-82.
— Fl. Tasm. **2**. 1859.	— 3. III. Flora Tasmaniae. t. 166-169.
— Tr. Linn. Soc. **20**. 1847.	— 4. An Enumeration of the Plants of the Galapagos Archipelagos, with Description of those, which are new. — pag. 163.

Hk.fil. Hdb. Fl. N. Zeal. 1864. 1867. — **J. D. Hooker:** 5. Handbook of the New Zealand Flora. London. 8^0. — pag. 1-392: 1864. pag. 393-end. 1867.

Hk. Exot. Fl. **1-3**. 1823-1827. — **W. J. Hooker:** 1. Exotic Flora. Edinburgh. gr. 8^0. **1**. 1823. **2**. 1825. **3**. 1827. Tabb. col.

— Brit. Fl. **1**. 1830. — 2. The British Flora. Volumen I. comprising the phaenogamous or flowering plants and the ferns. London. 8^0.

— Bot. Misc. **2**. 1831. — 3. Notice of some of the Plants collected during the above excursion (i. e. Cruckshanks: from Lima to Pasco). — pag. 205. — (Hooker: Botanical Miscellany. London. 8^0).

— *in Night. Ocean. sketch. 1835. — 4. List of the ferns in the collection made by Mr. Nightingale in the pacific isles; in Nightingale: Oceanic sketches. London. gr. 12^0.

— Ic. pl. **1-10**. 1837-1854. — 5. Icones plantarum ... London. 8^0.
1. t. 1-100: 1837. **2**. t. 101-200: 1837. **3**. t. 201-300: 1840. **4**. t. 301-400: 1841. **5**. t. 401-500: 1842. **6**. t. 501-600: 1843. **7**. t. 601-700: 1844. **8**. t. 701-750: 1844; t. 751-800: 1848. **9**. t. 801-900: 1852. **10**. t. 901-1000: 1854. vol. **10**. separatim edit. sub titulo: A century of Ferns. t. 1-100.

— Gen. 1838-1842. — 6. Genera Filicum, or illustrations of the Ferns and other allied genera; from the original coloured drawings of Francis Bauer Esq., with descriptive letterpress. London. gr. 8^0. 120 tab. col.
t. 1-20: 1838. t. 21-30: 1839. t. 31-69: 1840. t. 70-120: 1842. (an partim 1841?).

— *Fl. bor. am. **2**. 1840. — 7. Flora boreali-americana ... London. 8^0.

— in Benth. Pl. Hartw. 1840-41. — 8. Filices in G. Bentham: Plantas Hartwegianas imprimis Mexicanas adjectis nonnullis Grahamianis enumerat novasque describit. London. 8^0. — pag. 54-55: 1840. pag. 73: 1841.

— Lond. JoB. **1**. 1842. — 9. On Trichomanes Vittaria, De Cand. — pag. 137 t. 5.

— ——— — —10. On a new Species of Meniscium, from China. — pag. 294 t. 11.

— sp. **1-5**. 1844-1864. —11. Species Filicum; being descriptions of the known ferns, particularly of such as exist in the author's herbarium, or are with sufficient accuracy described in works to which he has had access; accompanied with numerous figures. London. 8^0. 304 Tab.

1. pag. 1-128, t. 1-40: 1844. — pag. 129-245, t. 41-70: 1846.

2. pag. 1-60, t. 71-90: 1851. — pag. 61-124, t. 91-110: 1852. — pag. 125-238, t. 111-140: 1858.

	3. t. 141-210: 1860. (an partim 1859?)
	4. t. 211-280: 1862. (an partim 1861?)
	5. pag. 1-96, t. 281-296: 1863. — pag. 97-314, t. 297-304: 1864.
Hk. Lond. JoB. **5.** 1846.	**W. J. Hooker:** 12. On a new Fern from Java, collected by. Mr. Th. Lobb (Gymnopteris Vespertilio). — pag. 193 t. 7-8.
— ———— **7.** 1848.	—13. Note on Anemia Seemanni. — pag. 564 t. 16.
— JoB. **4.** 1852.	—14. Notices of a new species of Deparia, discovered by Mr. Charles Moore in New Caledonia. — pag. 54 t. 3.
— — **5.** 1853.	—15. Notice of Two New Ferns from China. — pag. 236. t. 1-2.
— — —	—16. Description and Figure of a new Fern from Malacca. — pag. 309 t. 11.
— Cent. 1854.	vide n. 5. Ic. pl. **10.**
— JoB. **8.** 1856.	—17. On Polypodium anomalum, a new Species of Ferns, bearing its Sori on the upper side of the Frond. — pag. 360 t. 11.
— — **9.** 1857.	—18. On Asplenium (§ Schaffneria) nigripes, a Mexican, and on Davallia nodosa, an Indian Fern. — pag. 268. t. 9-10.
— — —	—19. Filices in G. Bentham: Florula Hongkongensis. — pag. 333, 353.
— *Fil. Exot. 1857-59.	—20. Filices exoticae, or, figures and descriptions of exotic ferns; chiefly of such as are cultivated in the Royal gardens of Kew. London. 4⁰. 100 Tab. col. t. 1-32: 1857. t. 33-88: 1858. t. 89-100: 1859. — (t. A. Gepp).
— 2 Cent. (1860) 1861.	—21. A second Century of Ferns. London. gr. 8⁰. 100 Tab.
— *Brit. Ferns 1861.	—22. The British Ferns, or, coloured figures and descriptions ... of the ferns of Great Britain and Ireland, systematically arranged. London. 8⁰. 66 Tab. col.
— *Gard. Ferns 1862.	—23. Garden Ferns. London. 8⁰. 64 Tab. col.
— *in Blakiston: Yangt-tsze 1862.	—24. Filices in Th. W. Blakiston: Five months on the Yangt-tsze. Appendix. London. 8⁰. (t. A. Gepp).
— Syn. 1865-1866.	vide Hk. et Bak.
— **et Arn.** Bot. Beech. Voy. 1830-1841.	— **et G. A. Walker-Arnott:** The Botany of Captain Beechey's Voyage ... London. 4⁰. 8 Tab. Fil. pag. 1-48: 1830. — pag. 49-96: 1832. — pag. 97-144: 1832. — pag. 145-192: 1833. — pag. 193-240: 1836. — pag. 241-288: ? — pag. 289-336: 1840. — pag. 337-384: 1840. — pag. 385-432: 1841? — pag. 433-486: 1841 (?) — (f. Jackson: JoB. **1893.** 298).
— **et Bak.** Syn. 1865-1868. ed. II. 1874.	— **and J. G. Baker:** Synopsis Filicum, or, a Synopsis of all known Ferns, including the Osmundaceae, Schizaeaceae, Marattiaceae and

Ophioglossaceae (chiefly derieved from the Kew Herbarium). Accompanied by Figures representing the essential Characters of each Genus. London. 8⁰. 9 Tab. — pag. 1-48 Hooker, pag. 49 ad finem et ed. II. Baker elaboravit.
pag. 1-32: 1865. — pag. 33-64: 1866. — pag. 65-352: 1867. — pag. 353-482: 1868. ed. II cum appendice: 1874.

Hk. et Grev. Ic. Fil. 1827-1831. — **W. J. Hooker et R. K. Greville:** 1. Icones Filicum: ad eas potissimum species illustrandas destinatae, quae hactenus vel in herbariis delituerunt prorsus incognitae, vel saltem nondum per icones botanicis innotuerunt. London. Fol. 2 voll. 240 Tab. col.
tab. t. 1-60: 1827. — t. 61-120: 1828 (?) — t. 121-180: 1829. — t. 181-240: 1830-31.

— — JoB. **1.** 1834. — — — 2. Description of two new Species of Gymnogramma from Peru. — pag. 61. t. 119-120.
vide Grev. et Hk.

Hope: JoB. **1890.** — **C. W. Hope:** 1. A new Lastrea from Assam. — pag. 145.

— — — — 2. Three new Lastreas from Assam. — pag. 326.

— — — — 3. The Ferns of the Chitral Relief Expedition. — pag. 122.

— Bull. Torr. Cl. **26.** 1899. — 4. Note on Asplenium Glenniei Baker in Synopsis Filicum 2 Ed. 488. — pag. 58.

— Journ. Bomb. Nat. Hist. Soc. **12-15.** 1899-1904. — 5. The Ferns of North-Western India. Including Afghanistan, the Trans-Indus Protected States, and Kashmir: arranged and named on the basis of Hooker and Baker's Synopsis Filicum and other works, with new Species added. 35 Tab. (Journal of the Bombay Natural History Society).
12. pag. 315, 527, 621: 1899.
13. pag. 25, 236: 1900; pag. 443, 657: 1901.
14. pag. 118, 252, 458: 1902; pag. 720: 1903.
15. pag. 78: 1903; pag. 415: 1904.

— et **Wright:** Journ. Linn. Soc. **35.** 1903. — et **C. N. Wright:** Filices in Descriptions of New Chinese Plants chiefly by Stephen Troyle Dunn F. L. S. . . . pag. 483.

Hoppe: Neu. Tachenb. **1805.** — **D. H. Hoppe:** Verzeichniss der in Deutschland wild wachsenden Farrenkräuter. -- Neues Botanisches Taschenbuch auf das Jahr 1805.

Hornem. Hort. reg. bot. Hafn. **2.** 1815. Suppl. 1819. — **J. W. Hornemann:** Hortus regius botanicus hafniensis in usum Tyronum et Botanophilorum. — Hafniae. 8⁰.

Horsf. Pl. Jav. Rar. 1838. — **Th. Horsfield:** Plantae Javanicae Rariores . . . Londini. Fol. t. 1-2.

Houtt. *Nat. Hist. **14**. 1783. — **M. Houttuyn:** 1. Natuurlijke historie oft uitvoerige beschryving der Dieren, Planten en Mineraalen, volgens het samenstel van Linnaeus. Deel II. Planten. Amsterdam. 8⁰. Filices in 14 Stuck. — Edit. germanica:

— Pfl. Syst. **13**[1]. 1786. — — 2. Des Ritters Carl von Linné vollständiges Pflanzensystem nach der 13 lateinischen Ausgabe und nach Anleitung des holländischen Houttuyn'schen Werkes . . . Nürnberg. 8⁰. vol. 13. erster Band.

Huds. Fl. angl. 1762. — **W. Hudson:** Flora anglica . . . Londini. 8⁰.

Humb. et Bonpl. Pl. aequin. **1-2**. 1805-1818. — **A. v. Humboldt et A. Bonpland:** 1. Plantae aequinoctiales . . . Parisiis. Fol.

HBK. Nov. Gen. et Sp. **1**. 1815. **7**. 1825. — — — 2. Nova Genera et Species plantarum, quas in peregrinatione orbis novi collegerunt . . . Ex schedis autographis A. Bonpland in ordinem digessit **C. S. Kunth** . . . Lutetiae Parisiorum. 4⁰. **1**. pag. 1. t. 1-2. **7**. t. 661-670. vide Kunth.

Husnot: *Cat. Crypt. Antill. franç. 1870. — **C. Husnot:** Catalogue des cryptogames recueillis aux Antilles françaises en 1868 et Essai sur leur distribution géographique dans ces îles. Caen 1870.

Jacq. Obs. bot. **1**. 1764. — **N. J. v. Jacquin:** 1. Observationum botanicarum iconibus ab auctore delineata illustratarum. Vindobonae. Fol.

— Misc. austr. **2**. 1781. — — 2. Miscellanea austriaca ad botanicam . . . spectantia. Vindobonae. 4⁰.

— Collect. **1-5**. 1786-1796. — — 3. Collectanea ad Botanicam . . . spectantia, cum figuris. Vindobonae. 4⁰. **1**. 1786. **2**. 1788. **3**. 1789. **4**. 1790. **5**. (Suppl.) 1796.

— Ic. pl. rar. **1**. 1786. **3**. 1793. — — 4. Icones plantarum rariorum. Vindobonae. Fol. **1**. t. 199-200. **3**. t. 637-646.

— Hort. Schönnbr. **4**. 1804. — — 5. Plantarum rariorum horti caesarei Schönbrunnensis descriptiones et icones. Viennae. Fol.

— Fragm. 1809. — — 6. Fragmenta botanica figuris coloratis illustrata. Viennae. Fol. t. 116.

Jan: Flora **1835**[1]. — **Jan:** Asplenium Tretteneriannm, ein neues Farrnkraut. — pag. 32.

Jaub. et Spach: Ill. pl. orient. **3**. 1847. — **Jaubert et Spach:** Illustrationes plantarum orientalium. Paris. Fol. t. 201.

Jenm. JoB. **1877**. — **G. S. Jenman:** 1. Supplement to the Jamaican Ferns recorded in Grisebach's »Flora of the British West Indies«. — pag. 263.

— — **1879**. — — 2. Second Supplement to the Jamaican Ferns etc. (vide n. 1). — pag. 257.

— — **1881**. — — 3. Third Supplement to the Ferns etc. (vide n. 1.) — pag. 51.

— — — — — 4. A New Tree-Fern from Jamaica. — pag. 275.

Jenm. JoB. **1882.**	**G. S. Jenman:** 5. Jamaica Ferns. — pag. 323.
— — **1886.**	— 6. On the Jamaica Ferns of Sloane's Herbarium. — pag. 14, 33.
— — —	— 7. Some additional Jamaica Ferns. — pag. 265.
— — **1887.**	— 8. The Ferns of Trinidad. — pag. 97.
— Bull. Dep. Jam. **1.** (n. s.) **1-5.** 1890-1898.	— 9. Synoptical List, with descriptions of the Ferns and Fern-Allies of Jamaica. — in Bull. of the Bot. Department of Jamaica. (old series **1.**) nr. 18. 20: 1890. nr. 21. 22. 23. 25. 26: 1891. nr. 29. 32. 33. 34. 35. 36. 38: 1892. nr. 40-49: 1893. (new series) **1.** 1894. pag. 28, 60, 75, 85. **2.** 1895. pag. 195, 266, 285. **3.** 1896. pag. 20, 45, 66, 110, 141, 162, 188, 211, 260. **4.** 1897. pag. 17, 42, 68, 112, 197, 251. **5.** 1898. pag. 44, 88, 153, 187, 208, 261, 272.
— Bull. Dep. Agr. Jam. **1.** 1903.	—10. Additions [to Synoptical List] in Bull. of the Departm. of Agriculture of Jamaica. — pag. 42, 116.
— Gard. Chr. 1894-1900.	—11. [Diagnoses specierum novarum].
— W. Ind. and. Guiana Ferns 1898. 1899. 1900.	—12. West Indian and Guiana Ferns. Appendix to Bull. of misc. inform. Bot. Department of Trinidad. **3.** 1898. **4.** 1899. **5.** 1900.
Johns. Ann. Nat. Hist. III. **17.** 1866.	**J. Y. Johnson:** Some Account of a new Species of Fern (Polystichum Maderense) recently discovered in the Island of Madeira. — pag. 287.
Johow: Hel. Juan Fern. 1893.	**F. Johow:** Los helechos de Juan Fernandez. Santiago de Chile. 8⁰. 1 Tab.
Jungh. Hoev. Tijdschr. **8.** 1841.	**F. Junghuhn:** Uitstapje naar de bosschen van de gebergten Malabar, Wayang en Tilu op Java. — pag. 349 (Hoeven en de Vriese: Tijdschrift voor Natuurlijke Geschiedenis en Physiologie. Amsterdam. 8⁰).
Juss. Gen. pl. 1789.	**A. L. de Jussieu:** Genera plantarum secundum ordines naturales disposita . . . Parisiis. 8⁰.
G. Karst. Ann. Jard. Buit. **12.** 1895.	**G. Karsten:** Morphologische und biologische Untersuchungen über einige Epiphytenformen der Molukken. — pag. 117. Tabb.
Karst. Linn. **28.** 1856.	**H. Karsten:** 1. Plantae columbianae — pag. 387.
— Fl. Col. **1. 2.** 1858-1869.	— 2. Florae Columbiae terrarumque adjacentium specimina selecta . . . Berolini. Fol. **1.** 1858-1861. **2.** 1862-1869. Tabb. col.
Klf. Flora 1822[1].	**G. F. Kaulfuss:** 1. Ueber die Entwickelung der Wedel der Ophioglosseen. — pag. 97. t. 2.
— — 1823[1].	— 2. Bemerkungen über die Farrnkränter welche in der, durch Hrn. Sieber veranstalteten Pflanzensammlung, von Martinique enthalten sind. — pag. 360.

Klf. Enum. 1824.	**G. F. Kaulfuss:** 3. Enumeratio Filicum, quas in itinere circa terram legit cl. Adalbertus von Chamisso, adjectis in omnia harum plantarum genera permultasque species non satis cognitas vel novas animadversiones. — Lipsiae. 8⁰. 2 Tab.
— Wes. d. Farrnkr. 1827.	— 4. Das Wesen der Farrnkränter, besonders ihrer Fruchttheile, zugleich mit Rücksicht auf systematische Anordnung betrachtet ... Erste Hälfte. Leipzig. 4⁰.
— Flora **1829**[1].	— 5. Antwortschreiben an Prof. Dr. Hoppe in Regensburg. — pag. 321, 337.
— Linn. **6**. 1831.	— 6. Lycopodiaceae et Filices in »Plantae Ecklonianae«. — pag. 181.
Kellogg: *Proc. Calif. Acad. **1**. 1854.	**A. Kellogg:** 1. Description of a new fern from Shoalwater Bay: Polypodium falcatum. — pag. 19.
— *——— **2**. 1861.	— 2. Description of a species of Polypodium found in the vicinity of San Francisco. — pag. 88.
Keys. Pol. Cyath. Hb. Bung. 1873.	**A. Keyserling:** 1. Polypodiacea et Cyatheacea Herbarii Bungeani. Lipsiae. gr. 8⁰.
— Mém. Ac. St. Petersb. VII. **22**[2]. 1875.	— 2. Gen. Adiantum L. 1 Tab.
Kiærsk. Cat. Fil. Hb. ht. Haun. 1874.	**H. Kiærskou:** Catalogus Filicum herbariorum horti botanici Hauniensis. In Beretning om Universitetets botaniske Have for Aarene 1871-73. Kjøbenhavn. 8⁰.
Kirk: Tr. N. Zeal. Inst. **5**. 1873.	**T. Kirk:** 1. On the Specific Characters of Dicksonia antarctica Br., and Dicksonia lanata Col. — pag. 345.
— ——— **6**. 1874.	— 2. On the New Zealand Forms of Cheilanthes. — pag. 247.
— ——— **9**. 1877.	— 3. Description of a New Species of Pilularia. — pag. 547 t. 29.
— ——— **10**. 1878.	— 4. Description of a new Species of Hymenophyllum. — pag. 394 t. 21 A. B.
— ——— —	— 5. On Hymenophyllum villosum Col. — pag. 395.
— ——— —	— 6. On Lindsaya viridis Col. — pag. 396.
— ——— —	— 7. On Nephrodium decompositum Br. and N. glabellum A. Cunn. — pag. 398.
— ——— **11**. 1879.	— 8. Description of a new Species of Hymenophyllum. — pag. 457 t. 19 A.
— ——— **17**. 1885.	— 9. On the Ferns and Ferns Allies of Stewart Island. — pag. 228.
— ——— **23**. 1891.	—10. On a remarkable Variety of Asplenium umbrosum J. Sm. — pag. 424.
Klinsm. Bot. Zeit. **1852**.	**Klinsmann:** Ueber die Botrychien der deutschen Flora und über Botrychium Kannenbergii als eine neue Species und deren Stellung zu den übrigen. — pag. 377 t. 6.

Kl. Linn. **14**. 1840.	**J. F. Klotzsch:** 1. Filices in Collectio Plantarum Bahiensium a Luschnathio decerptarum, exsiccatarum et veno positarum. — pag. 284.
— — **18**. 1844. **20**. 1847.	— 2. Beiträge einer Flora der Aequinoctial-Gegenden der neuen Welt. **18**. pag. 515. **20**. pag. 337. 433.
— Allg. Gartenzeit. **1852**.	— 3. Einige neue Baumfarrn, beschrieben von — pag. 41, 49. — etiam in Bot. Zeit. **1854**. 439.
— ———— **1855**.	— 4. Ein neues Polypodium aus Central-Amerika. — pag. 33.
— ————	— 5. Zwei neue Farrnkräuter. — pag. 65.
— ———— —	— 6. Nothochlaena (Cincinalis) chrysophylla. — pag. 265.
— ———— —	— 7. Ein neues Polypodium. — pag. 377.
— ———— —	— 8. Adiantopsis elongata Kl. — pag. 392.
— ———— **1856**.	— 9. Ein neues Farrn: Aspidium (Bathmium) nigripes Kl. — pag. 189.
Koch: Wochenschr. Gärtn. u. Pflzk. **1858**.	**Karl Koch:** Die Gold- und Silberfarne unserer Gewächshäuser. — pag. 1. 12. 32. c. figg.
Koch: Syn. Fl. Germ. ed. II. **3**. 1845.	**W. D. J. Koch:** Synopsis Florae germanicae et helveticae . . . Leipzig. 8°.
König: Naturforsch. Halle **21**. 1785.	**D. König:** Auszug eines Schreibens des Herrn D. König aus Trankebar vom 28. März 1783 an den Herrn Conferenzrath O. F. Müller. — Der Naturforscher Halle. 21 Stück. — pag. 107 t. 3.
Komarov: Fl. Mansh. 1901.	**V. L. Komarov:** Flora Manshuriae. In Acta hort. Petrop. **20**. 1901.
Krasser: Ann. Hofmus. Wien **15**. 1900.	**F. Krasser:** Filices, Lycopodiaceae et Selaginellaceae ap. Zahlbrückner: Plantae Pentherianae. — pag. 3 t. 1, 2 f. 1-5.
Krug: Engl. Jahrb. **24**. 1897.	**L. Krug:** Pteridophyta Herbarii Krug et Urban, e determinationibus cl. J. G. Baker et cl. H. Christ et beati M. Kuhn excerpsit atque composuit Leopoldus Krug. — pag. 97. — (in Ig. Urban: Additamenta ad cognitionem florae Indiae occidentalis, particula IV).
Kuhn: Fil. Deck. 1867.	**Max. Kuhn:** 1. Filices Deckenianae. Dissertatio. Lipsiae. 8°. (etiam in n. 2. Fil. Afr.).
— App. Ind. Sem. Berol. 1867.	vide A. Braun nr. 8.
— Fil. Afr. 1868.	— 2. Filices Africanae, Revisio critica omnium hucusque cognitorum cormophytorum Africae indigenorum additamentis Braunianis Novisque Africanis speciebus ex Reliquiis Mettenianis adaucta. Accedunt Filices Deckenianae et Petersianae. Lipsiae. 8°.
— Bot. Zeit. **1868**.	— 3. Filices quaedam novae et indescriptae. — pag. 40.
— JoB. **1868**.	— 4. Adnotationes de filicibus nonnullis Chinae indigenis. — pag. 268.
— Linn. **35**. 1868. **36**. 1869.	— 5. Reliquiae Mettenianae. **35**. pag. 385. **36**. pag. 40

Kuhn: Abh. Nat. Ges. Halle **11**. 1869.	**Max. Kuhn:** 6. Beiträge zur mexicanischen Farnflora. — pag. 23.
— Ann. Lugd. Bat. **4**. 1869.	— 7. Filices. — pag. 276 (vide Miquel et Mett. n. 11).
— Verh. zool. bot. Ges. **19**. 1869.	— 8. Filices Novarum Hebridarum. — pag. 569.
— Bot. Zeit. **1869**. **1870**.	— 9. Analecta pteridographica. **1869**. pag. 130, 144, 162, 391, 437, 458. **1870**. pag. 89, 135, (525).
— ——— **1871**.	Recensio pag. 503. vide Mac Ken.
— *Ind. Sem. Ht. Berol. 1875.	—10. Descriptio filicis arboreae novae: Cyathea Hildebrandtii. — pag. 20.
— Verh. zool. bot. Ges. **25**. 1876.	—11. Bemerkungen über einige Farne von der Insel Celebes. — pag. 593.
— JoB. **1877**.	—12. Cormophyta ap. Polakowsky: Bryophytas et Cormophytas Costaricenses anno 1875 lectas, enumeravit. — pag. 230.
— v. Deck. Reis. 3^3. Bot. 1879.	—13. Cryptogamae vasculares in »Botanik von Ostafrika« in Cl. v. d. Decken's Reisen in Ostafrika. 3 Bd. 3 Abth. Berlin. 8^0.
— Jahrb. bot. Gart. Berlin **1**. 1881.	—14. Uebersicht über die Arten der Gattung Adiantum. — pag. 337.
— Chaetopt. 1882.	—15. Die Gruppe der Chaetopterides unter den Polypodiaceen. Berlin. 8^0. 2 Tab. — (In Festschrift zu 50 jähr. Jubil. d. kgl. Realschule zu Berlin 1882).
— Ber. deutsch. bot. Ges. **1**. 1883.	—16. Ueber Farne und Charen von der Insel Socotra. — pag. 238.
— Fl. bras. 1^2. 1884.	—17. Isoëtaceae, Marsiliaceae, Salviniaceae in Martius: Flora brasiliensis. **1**. pars 2. — pag. 645 t. 81-82.
— Forschreis. Gazelle **4**. Farne 1889.	—18. Farne (Filicinae) und bärlappartige Gewächse (Lycopodinae) in »Die Forschungsreise S. M S. »Gazelle« in den Jahren 1874 bis 1876. **4**. Botanik.« Berlin. 4^0. 3 Tab.
— in *Schum. Fl. Kais. Wilh. Land 1889.	—19. Filices [novae] in K. Schumann und M. Hollrung: Die Flora von Kaiser Wilhelmsland. Beiheft zu den Nachrichten über Kaiser Wilhelmsland und den Bismarcks Archipel.
— Verh. bot. Ver. Brand. **31**. 1889.	—20. Filices ap. R. Büttner: Neue Arten von Guinea, dem Kongo und dem Quango. — pag. 64.
— in Engl. Hochgbfl. trop. Afr. 1892.	—21. Filices in Engler: Über die Hochgebirgsflora des tropischen Afrika. Berlin. 4^0. (Aus d. Abhdl. d. k. preuss. Akad. d. Wiss. v. Jahre 1891).
— Hedwigia **35**. 1896.	vide Schenck.
— Engl. Jahrb. **24**. 1897.	vide Krug.
Kunth: Syn. pl. **1**. 1822.	**K. S. Kunth:** Synopsis plantarum quas in itinere ad plagam aequinoctialem orbis novis collegerunt A. de Humboldt et A. Bonpland. Paris. 8.
O. Ktze. Flora 1880.	**O. Kuntze:** 1. Miscellen über Hybriden aus der Leipziger Flora. — pag. 291.

O. Ktze. Rev. Gen. Pl. **2.** 1891. $\mathbf{3}^{2}$. 1898.	**O. Kuntze:** 2. Revisio generum plantarum . . .
Kze. Flora $\mathbf{1833}^{2}$.	**Gustav Kunze:** 1. Hypodematium, eine neue Farrngattung. — pag. 689.
— Linn. **9.** 1834.	— 2. Synopsis plantarum cryptogamicarum ab Eduardo Poeppig in Cuba insula et in America meridionali collectarum. — pag. 1.
— —— —	— 3. Thyrsopteris, eine neue Farrngattung. — pag. 506.
— — **10.** 1836.	— 4. Acotyledonearum Africae australioris recensio nova. I. Filices. — pag. 481.
— Anal. 1837.	— 5. Analecta pteridographica, sive descriptio et illustratio Filicum aut novarum aut minus cognitarum. Lipsiae 1837. Folio. 30 Tab.
— Flora $\mathbf{1837}^{1}$.	— 6. Zur Erläuterung der in Arrabida's Flora Fluminensis dargestellten Kryptogamen. — pag. 321.
— — $\mathbf{1839}^{1}$. Beibl.	— 7. Filices in Martii Herbarium Florae Brasiliensis.
— Linn. **13** 1839.	— 8. Additamentum enumerationis filicum mexicanarum, partim a B. Schiedeo, partim a cl. Car. Ehrenbergio aliisque collectarum — pag. 129.
— Farrnkr. 1840-1851.	— 9. Die Farrnkräuter in kolorirten Abbildungen naturgetreu erläutert und beschrieben von — Schkuhr's Farrnkräuter, Supplement. 2 vol. Leipzig. 4^{0}. 140 Tab. col.
	vol. **1.** Lief. 1-2. pag. 1- 62. t. 1- 30: 1840.
	» 3-5. pag. 63-108. t. 31- 50: 1841-1842.
	» 6. pag. 109-138. t. 51- 60: 1843.
	» 7. pag. 139-166. t. 61- 70: 1844.
	» 8. pag. 167-190. t. 71- 80: 1845.
	» 9. pag. 191-222. t. 81- 90: 1846.
	» 10. pag. 223-262. t. 91-100: 1847.
	vol. **2.** » 11. pag. 1- 26. t. 101-110: 1848.
	» 12. pag. 27- 50. t. 111-120: 1849.
	» 13. pag. 51- 74. t. 121-130: 1850 (?).
	» 14. pag. 75- 98. t. 131-140: 1851.
— Linn. **16.** 1842. **17.** 1843.	—10. Pugillus primus plantarum adhuc ineditarum, seu minus cognitarum . . . . . coluit Hortus bot. Univ. Litt. Lipsiensis. **16.** pag. 309. Pugillus secundus etc. **17.** pag. 566 (vide n. 19).
— Bot. Zeit. **1843.**	—11. Nothochlaena pulchella, ein neues Farrnkraut der Gärten. — pag. 633.
— —— **1844.**	—12. Generis Polypodiacearum Jamesonia Hook. skiagraphia. — pag. 737.
— —— **1844. 1845. 1847. 1850.**	—13. Hooker: Species filicum. (Recensio).
— Linn. **18.** 1844.	—14. Filicum in Promontorio Bonae Spei et Portum Natalensem a Gueinzio nuperius collectarum. — pag. 113.

Kze. Linn. 18. 1844.	**Gustav Kunze:** 15. Filices a Leiboldio in Mexico lectae. — pag. 303.
— Bot. Zeit. **1845.**	—16. Filices a cl. Moritz in Caracas lectae. — pag. 281.
— ——— **1846. 1848.**	—17. In filices Javae Zollingerianas aliasque ex herbario Moricandiano observationes. **1846.** pag. 417 etc. **1848.** pag. 97 etc.
— in Lehm. Pl. Preiss. **2.** 1847.	—18. Filices in Lehmann: Plantae Preissianae in Australia collectae. **2.** pag. 109.
— Linn. **20.** 1847.	—19. Pugillus tertius plantarum etc. (vide n. 10). — pag. 1.
— Bot. Zeit. **1848.**	—20. Pteridographia Japonica. Uebersicht der Farrn der Flora von Japan, sowie der Inseln Loo Choo, Peel und Bonin Sima. — pag. 489 etc.
— ——— —	—21. Ueber ein noch unbeschriebenes und ein neuerlich beschriebenes Antrophyum. — pag. 702.
— Flora **1848.**	—22. Ueber drei bisher mehrfältig verwechselte deutsche Farrn: Aspidium lobatum Sm., aculeatum Sm. und Braunii Spenn. — pag. 353.
— Linn. **21.** 1848.	—23. Filices in Plantae Kegelianae Surinamenses. — pag. 201, 284.
— Amer. Journ. Sci. II. **6.** 1848.	—24. Notes on some Ferns of the United States. — pag. 80. — (descript. spec. nov. etiam in Bot. Zeit. **1850.** 481).
— Bot. Zeit. **1849. 1850.**	—25. Pteridologische Studien. **1849.** pag. 865, 881. **1850.** pag. 57.
— Linn. **22.** 1849.	—26. Filices et Lycopodiaceae in »Plantae Regnellianae«. — pag. 575.
— — **23.** 1850.	—27. Index Filicum (sensu latissimo) adhuc, quantum innotuit, in hortis Europaeis cultarum. — pag. 209. Supplementum pag. 408.
— Bot. Zeit. **1851.**	—28. Oleandra Cav. species in herbario suo servatas sciagraphice disposuit. — pag. 345.
— Linn. **24.** 1851.	—29. Filices Nilagiricae. — pag. 239.
— — **25.** 1853.	—30. Cryptogamae vasculares ap. Schlechtendal: Plantae Wagenerianae Columbicae. — pag. 745.
Kurz: Journ. As. Soc. Bengal 39^2. 1870.	**S. Kurz:** On some new or imperfectly known Indian Plants. — pag. 61. t. 5. — conf. J. Amann.
Lab. Nov. Holl. pl. spec. **2.** 1806.	**de Labillardière:** 1. Novae Hollandiae plantarum specimen. Parisiis. 4^0. Tabb.
— Sert. Austr. Cal. 1824.	— 2. Sertum Austro-Caledonicum, pars prior. Parisiis. 4^0. 13 Tab.
Lag. Garcia et Clem. *Anal. Cienc. **5.** 1802.	**M. Lagasca, D. Donato Garcia, y Don Simon de Roxas Clemente:** Introducion à la Criptogamia de España. — pag. 135. (Anales de Ciencias naturales. Madrid). — (t. A. Gepp).
Lag. Gen. et sp. 1816	**M. Lagasca:** Genera et species plantarum, quae aut novae sunt aut nondum recte cognoscuntur. Matriti. 4^0.

Laguna: Anal. Hist. Nat. 7. 1878.	**M. Laguna:** Cien Helechos de Filipinas. (Anales de la Soc. españa de Historia Natural. Madrid).
Lam. Fl. franç. **1**. 1778.	**de Lamarck:** 1. Flore françoise . . . Paris. 8^0.
— Enc. **1-4**. 1783-1797.	— 2. Encyclopédie méthodique. Botanique. Paris. 4^0. **1**. 1783. **2**. 1786. **3**. 1789. **4**. 1789. (vol. seq. **5-8** et Suppl. **1-5** Poiret elaboravit: vide Poiret). — Gen. Lemma (**3**. 720) et Meniscium (**4**. 93) Desrousseaux, Gen. Osmunda (**4**. 649) Savigny elaboravit.
— **et DC.** Fl. franç. **2**. 1805. **5**. 1815.	— **et De Candolle:** Flore française . . . 3me édition. Paris. 8^0.
Langsd. et Fisch. Ic. Fil. 1810-(1818).	**G. H. Langsdorff et F. Fischer:** Plantes recueillies pendant le Voyage des Russes autour du Monde. Expédition dirigée par M. de Krusenstern. 1me partie. Icones Filicum. Tubingae. Fol. 30 Tab.
La Pylaie: *Mém. Soc. Linn. Paris **4**. 1826.	**B. de La Pylaie:** Notice sur l'île de Terre Neuve et quelques îles voisines. — pag. 417.
Lasch: Bot. Zeit. 1856.	**W. Lasch:** Ueber Bastarde unter der wildwachsenden Farrn. — pag. 433.
Ledeb. Fl. ross. **4**. 1853.	**C. F. a Ledebour:** Flora rossica . . . Stuttgartiae. 8^0.
Le Grand: Bull. Soc. Fr. **30**. 1883.	**M. A. Legrand:** Notice sur quelques plantes critiques ou peu communes. — pag. 68.
Le Prieur: Ann. sc nat. **19**. 1830.	**Le Prieur:** Note sur le Pteris cornuta de Palisot-Beauvois, espèce du genre Ceratopteris. — pag. 99 t. 4 A.
Leybold: Flora **1855**.	**F. Leybold:** Asplenium Seelosii, ein neuer Farrn aus Südtirol. — pag. 81, 348 t. 15.
L'Hérit. Sert. angl. 1788.	**L'Héritier de Brutelle:** Sertum anglicum . . . . Paris. Fol. Tab.
Liebm. Vid. Selsk. Skr. V. **1**. 1849.	**F. Liebmann:** Mexicos Bregner, en systematisk. critisk, plantegeografisk Undersøgelse. — pag. 151. (etiam seorsim impressum).
Liljebl. Vet. Ak. Nya Handl. **1793**.	**S. Liljeblad:** Acrostichum Hyperboreum, en tilförene okänd Svensk Växt. — pag. 201 t. 8.
Lindm. Ark. för Bot. **1**. 1903.	**C. A. M. Lindman:** 1. Remarks on some American Species of Trichomanes Sm. sect. Didymoglossum Desv. — pag. 7. with 31 illustr.
— ———	— 2. Beiträge zur Kenntniss der tropisch-amerikanischen Farnflora. — pag. 189 t. 7-14.
— ——— 3^6. 1904.	— 3. Regnellidium, novum genus Marsiliacearum. Mit 10 Textfiguren. — pag. 1.
— Hedwigia **43**. 1904.	— 4. Neue Speziesnamen einiger südamerikanischen Farne. — pag. 308. Fig.
Link: Enum. alt. **2**. 1822.	**H. F. Link:** 1. Enumeratio plantarum horti regii botanici Berolinensis altera. Berolini. 8^0.
— Hort. Berol. **2**. 1833.	— 2. Hortus regius botanicus Berolinensis descriptus. Berolini. 8^0.
— Fil. sp. 1841.	— 3. Filicum species in Horto regio botanico Berolinensis cultae. Berolini. 8^0.

L sp. **2**. 1753; ed. II. **2**. 1763.	**C. v. Linné (Linnaeus):** 1. Species plantarum . . . Holmiae. 8⁰.
— Syst. nat. ed. X. **2**. 1759.	— 2. Systema Naturae . . . Holmiae 1759.
— Mant. 1767. 1771.	— 3. Mantissa plantarum . . . . Holmiae. 8⁰. — pag. 1-142. 1767. pag. 143-584. 1771.
— Syst. pl. ed. nov. **4**. 1780.	— 4. Systema plantarum . . . editio novissima curante J. J. Reichard. Francofurti a/M. 8⁰.
L.fil. Suppl. 1781.	**C. v. Linné fil.:** Supplementum plantarum Systematis vegetabilium editionis XIII. . . . Brunsvigae. 8⁰.
Lois. *Notice 1810.	**Loiseleur Deslongchamps:** Notice sur les plantes à ajouter à la Flore de France . . . Paris. 8⁰.
Lour. Fl. Cochinch. **2**. 1790.	**J. de Loureiro:** Flora cochinchinensis ... *Ulyssipone. 4⁰.
ed. in Germ. **2**. 1793.	— Denuo in Germania edita cum notis citata K. L. Willdenow. Berolini 1793. 8⁰. (Hanc editionem in Indice citavi).
Lowe: Ferns **1-8**. 1856-1860.	**E. J. Lowe:** 1. Ferns: British and exotic. London. 8⁰. Tab. col. — sec. edit. 1872. **1**. 1856. 50 Tab. **2**. 1858. 56 Tab. **3**. 1857. 50 Tab. **4**. 1859. 64 Tab. **5**. 1858. 56 Tab. **6**. 1857. 50 Tab. **7**. 1859. 66 Tab. **8**. ? (1859 ?).
— *Hist. new Ferns 1860-61 (62 ?).	— 2. A Natural History of new and rare Ferns. London. 8⁰. 72 Tab. col. — (t. A. Gepp).
Lowe: Tr. Cambr. Phil. Soc. **4**. 1830.	**R. T. Lowe:** 1. Primitiae Faunae et Florae Maderae et Portus Sancti. — pag. 1.
— JoB. **1**. 1834.	— 2. Notes and observations on the above »List of Plants« observed in the Island of Madeira. — pag. 22.
— Tr. Cambr. Phil. Soc. **6**. 1838.	— 3. Novitiae Florae Maderensis: or, Notes and Gleanings of Maderan Botany. — pag. 523.
Ludwig: Inst. ed. II. 1757.	**C. G. Ludwig:** Institutiones Historico Physicae Regni Vegetabilis. Lipsiae. 8⁰.
Luerss. Fil. Graeff. 1871.	**C. Luerssen:** 1. Filices Graeffeanae. Beitrag zur Kenntniss der Farnflora der Viti-, Samoa-, Tonga- und Ellice's Inseln; in Schenk und Luerssen: Mittheilungen aus der Botanik. **1**. pag. 57. t. 11-19.
— Farne d. Samoa 1874.	— 2. Die Farne der Samoa Inseln, ein Verzeichniss der bis jetzt von den Schiffer-Inseln bekannten Gefässkryptogamen, nebst allgemeinen Bemerkungen über die Systematik dieser Pflanzengruppe. — l. c. pag. 345.
— Journ. Mus. Godeffr. **3**. 1873. 1875.	— 3. Zur Flora von Queensland, Verzeichniss der von Frau Amalie Dietrich in den Jahren 1863 bis 1873 an den Nordostküste von Neuholland gesammelten Pflanzen. — pag. 1 ff. 1873. pag. 233 ff. t. 12-18. 1875.
— ——— —	— 4. Ein Beitrag zur Farnflora der Palaos- oder Pelew-Inseln. — pag 52.

Luerss. Journ. Mus. Godeffr. **3.** 1873.	**C. Luerssen:** 5. Ueber die Farnflora der Cooks- oder Hervey-Inseln. — pag. 59.
— Flora **1875.**	— 6. Gefässkryptogamen in H. Wawra: Beiträge zur Flora der Hawai'schen Inseln. — pag. 417, 433.
— — **1876.**	— 7. Verzeichniss der Gefässkryptogamen, welche Dr. H. Wawra auf seiner Erdumsegelung ... sammelte. — pag. 225, 285, 289.
— Abh. Nat. Ver. Bremen **7.** 1882.	— 8. Filices in »Reliquiae Rutenbergianae«. — pag. 1. t. 1.
— ——— —	— 9. Filices ap. Copeland: Ein Besuch auf der Insel Trinidad im südatlantischen Ocean. — pag. 277.
— Bot. Centralbl. **9.** 1882.	—10. Ueber einige Hymenophyllaceen Neuhollands und Polynesiens. — pag. 438.
— ——— —	—11. Eine neue Cheilanthes des tropischen Australiens. — pag. 442.
— ——— **11.** 1882.	—12. Zur Farnflora Hinterindiens und West-Sumatra's. — pag. 26, 76.
— Engl. Jahrb. **4.** 1883.	—13. Archegoniatae apud Engler: Beiträge zur Flora des südlichen Japans und der Liu-Kiu Inseln. — pag. 354.
— Rabenh. Kr. Fl. ed. II. **3.** 1884-89.	—14. Die Farnpflanzen oder Gefässbündelkryptogamen (Pteridophyta) in L. Rabenhorst's Kryptogamen Flora von Deutschland, Oessterreich und der Schweiz. Zweite Auflage. Leipzig. 8⁰. 223 Abbild.
Lyon: Bot. Gaz. **40.** 1905.	**Harold L. Lyon:** A new genus of Ophioglossaceae. — pag. 455. fig.
Mac Ken: *Ferns Natal 1869.	**M. Mac Ken:** The Ferns of Natal. Pietermaritzburg. 8⁰. (vide Kuhn: Bot. Zeit. **1871.** 503).
— **et Gerr.** *Syn. Fil. cap. 1870.	— **and W. J. Gerrard:** Synopsis Filicum capensium. Pietermaritzburg. 8⁰.
Mak. *Bot. Mag. Tokyo **6.** 1892.	**T. Makino:** 1. Notes on Japanese Plants. XV. — pag. 45. — (t. A. Gepp).
— *——— **8.** 1894.	— 2. Three Japanese Plagiogyriae. — pag. 332.
— *——— —	— 3. Gymnogramme Makinoi Maxim. — pag. 481. c. tab.
— *——— **8.** 1894. **9.** 1895.	— 4. Mr. H. Kuroiwa's collection of Liu-Kiu plants. **8.** pag. 370, 411. **9.** pag. 6.
— *——— **10.** 1896.	— 5. Review of some species of Japanese Ferns. — pag. 148, 177. — (t. A. Gepp).
— *——— **11.** 1897.	— 6. A new Aspidium from Liu-Kiu Island. — pag. 18.
— *——— **11.** 1897. **12.** 1898. **13.** 1899.	— 7. Plantae japonenses novae vel minus cognitae. **11.** pag 64, 82; **12.** pag. 26, 36, 120; **13.** pag. 12, 25, 44, 56, 61, 79, 111, 128.
— *——— **12.** 1898.	— 8. Contributions to the study of the Flora of Japan. — pag. 11, 86, 161, 192, 298, 372.
— *Phan. et Pter. Jap. icon. 1899 —.	— 9. Phanerogamae et Pteridophyta Japonicae iconibus illustratae, or, figures with brief de-

	scriptions and remarks of the flowering plants and ferns of Japan. Tokyo. **1**. t. 1-35: 1899; t. 36-45: 1900; t. 45-60: 1901.
Mak. Bot. Mag. Tokyo ***15**. 1901. **17**. 1903. **18**. 1904. **19**. 1905.	**T. Makino:** 10. Observations on the Flora of Japan. **15***. pag. 102; **17**. pag. 70; **18**. pag. 129; **19**. pag. 131.
Mann: Proc. Amer. Ac. **7**. 1868.	**H. Mann:** Enumeration of Hawaian Plants. Filices pag. 211. (Species novae ab Eaton descriptae).
Martens: Bull. Acad. Brux. **4**. 1837.	**M. Martens:** Notice sur un cas d'hybridité dans les fougères. — pag. 47.
Mart. et Gal. Mém. Ac. Brux. **15**. 1842.	**M. Martens et H. Galeotti:** Mémoire sur les Fougères du Mexique, et Considérations sur la Géographie botanique de cette Contrée. 23 Tab.
Marthe: *Cat. jard. méd. Paris 1801.	**F. Marthe:** Catalogue des plantes du jardin médical de Paris . . . Paris. 8⁰.
Mart. *Denkschr. Bot. Ges. Regensb. **2**. 1822.	**C. de Martius:** 1. De plantis nonnullis antediluvianis ope specierum intertropicos viventium illustrandas. — pag. 121.
— Ic. Cr. Bras. 1834.	— 2. Icones plantarum cryptogamicarum quas in itinere annis MDCCCXVII-MDCCCXX per Brasiliam . . . collegit et descripsit . . . Monachii. Fol. 76 tab. col. (1826-1834 descriptiones nov. spec. certe non antea 1834 editae sunt).
Maury: Journ. de Bot. **3**. 1889.	**M. P. Maury:** Énumération des plantes du Haut-Orénoque, recoltées par MM. J. Chaffanjon et A. Gaillard. — pag. 129. Fig.
Maxim. Mém. Ac. St. Pétersb. **9**. 1859.	**C. J. Maximowicz:** 1. Primitiae Florae Amurensis. Versuch einer Flora des Amurlandes. — pag. 1.
— Bull. Ac. St. Pétersb. **15**. 1871.	— 2. Diagnoses breves plantarum novarum Japoniae et Mandshuriae. Decas septima. — pag. 226.
— Mél. Biol. **11**. 1883. **12**. 1886.	— 3. Diagnoses plantarum novarum asiaticarum; in Mélanges biologiques tirés du Bull. Acad. Imp. Pétersb. — **11**. pag. 867. **12**. pag. 571.
Maxon: Bull. Torr. Cl. **27**. 1900.	**W. R. Maxon:** 1. A new Asplenium, hitherto referred to A. Trichomanes var. incisum Moore. — pag. 197.
— — —	— 2. A list of the Pteridophyta collected in Alaska in 1900 by Mr. J. B. Flett, with description of a new Dryopteris. — pag. 637.
— *Proc. Biol. Soc. Wash. **13**. 1900.	— 3. Polypodium hesperium, a new fern from Western North America. — pag. 199.
— Proc. U. S. Nat. Mus. **23**. 1901.	— 4. A List of the Ferns and Fern Allies of North America North of Mexico, with principal Synonyms and Distribution. — pag. 619.
— Fern Bull. **10**. 1902.	— 5. An interesting Japanese Polypody. — pag. 42.
— Contr. U. S. Nat. Herb. **8**. 1903.	— 6. A Study of Certain Mexican and Guatemalan Species of Polypodium. — pag. 271. t. 61-62.

Maxon: Proc. U. S. Nat. Mus. **27**. 1904. Bull. Dep. Agr. Jam. **2**. 1904.	**W. R. Maxon:** 7. Two New Ferns of the Genus Polypodium from Jamaica. Proc. pag. 741. Bull. pag. 176.
— Proc. U. S. Nat. Mus. **27**. 1904.	— 8. A new Fern. Goniophlebium pringlei, from Mexico. — pag. 953 t. 48.
— Bull. Torr. Cl. **31**. 1904.	— 9. A new Asplenium from Mexico. — pag. 657. c. fig.
— Fern Bull. **12**. 1904.	—10. Notes on American ferns VII. — pag. 101.
— Bot. Gaz. **39**. 1905.	—11. Adenoderris, a valid genus of ferns. — pag. 366. Fig.
— Bull. Torr. Cl. **32**. 1905.	—12. On the names of three Jamaican species of Polypodium. — pag. 73.
— ——————— —	—13. A new Botrychium from Jamaica. — pag. 219 t. 6.
— Smiths. Misc. Coll. **47**. 1905.	—14. A new species of fern of Genus Polypodium from Jamaica. — pag. 410 t. 57.
— Proc. Biol. Soc. Wash. **18**. 1905.	—15. A new fern from Porto Rico. — pag. 215.
— ——————— —	—16. A new name for a Middle American fern. — pag. 224.
— ——————— —	—17. A new cloak-fern from Mexico. — pag. 205.
— ——————— —	—18. A new name for Kaulfussia Blume, a genus of Marattiaceous Ferns. — pag. 239.
Mérat: *Fl. Paris. ed. II. 1821. ed. IV. **1**. 1836.	**F. V. Mérat:** Nouvelle Flore des environs de Paris ... Paris. 8⁰.
Merrill: Bureau Govt. Laborat. Manila n. **27**. 1905.	**Elmer D. Merrill:** A review of the identifications of the species described in Blanco's Flora de Filipinas. — Bureau of Government Laboratories, Manila nr. 27. April 1905. 8⁰.
Mett. Linn. **20**. 1847.	**G. Mettenius:** 1. Ueber Azolla. — pag. 259. t. 23.
— Fil. Lips. 1856.	— 2. Filices horti botanici Lipsiensis. Die Farne des botanischen Gartens zu Leipzig. — Leipzig. Fol. 30 Tab.
— Fil. Lechl. **1**. 1856. **2**. 1859.	— 3. Filices Lechlerianae, chilenses ac peruanae. Leipzig. 8⁰. 3 Tab.
	— Ueber einige Farngattungen (abgedruckt aus d. Abhandl. der Senckenb. Naturf. Ges. zu Frankfurt a. M. **2**. (I-IV). **3**. (V-VI)):
— Pol. 1857.	— 4. I. Polypodium. 3 Tab.
— Plag. 1858.	— 5. II. Plagiogyria. 1 Tab.
— Üb. Pteris 1858.	— 6. III. Über die mit einem Schleier versehenen Arten von Pteris. 1 Tab.
— Pheg. u. Asp. 1858.	— 7. IV. Phegopteris und Aspidium. 2 Tab.
— Cheil. 1859.	— 8 V. Cheilanthes } 3 Tab.
— Aspl. 1859.	— 9. VI. Asplenium } 3 Tab.
— Ann. sc. nat. IV. **15**. 1861.	—10. Filices Novae Caledoniae a cl. Vieillard collectae. — pag. 55. t. 3.
— Ann. Lugd. Bat. **1**. 1863-64. **2**. 1866. **4**. 1868-69.	—11. Filices praesertim indicae et japonicae in Miquel: Annales Musei Lugduno Batavi. (vide Miquel).

	1. pag. 46. t. 3. 1863 (Gleicheniaceae et Cyatheaceae). — pag. 222. t. 6. 1864 (Hypolepis, Phegopteris, Aspidium, Oleandra, Cystopteris).
	2. pag. 219. 1866 (Polypodium, Micropodium, Asplenium).
	4. pag. 170 t. 7. 1868-69 (Platycerium, Antrophyum, Taenitis, Vaginularia).
	vide Kuhn nr. 7 et Miquel nr. 2.
Mett. Abh. Sächs. Ges. Wiss. 7. 1864.	G. **Mettenius:** 12. Ueber die Hymenophyllaceae. 5 Tab.
— Ann. sc. nat. V. 2. 1864. 3. 1865.	—13. Filices in Triana et Planchon: Prodromus Florae Novo-Granatensis. 2. pag. 193. 3. pag. 310 (Rhizocarpeae).
— Mém. soc. sc. nat. Cherb. 10. 1864.	—14. ap. Le Jolis: Fougères de la Nouvelle Calédonie. — pag. 317. (nomina nuda).
— in Pl. Tinn. 1867.	—15. Rhizocarpeae in Kotschy & Peyritsch: Plantae Tinneanae ... Vindobonae. Fol. t. 25.
— Novara Reis. Bot. 1. 1870.	—16. Cryptogamae vasculares in »Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857-1859. Botanischer Theil. Erster Band. Sporenpflanzen«. — pag. 199. Wien. 4°.
Meyen: Reise 1. 1834.	F. J. F. **Meyen:** 1. Reise um die Erde. Berlin. 4°.
— Nova Acta 18. 1836.	— 2. Beiträge zur Kenntniss der Azollen. — pag. 505. t. 38.
C. A. **Mey.** Verz. Pfl. Cauc. 1831.	C. A. **Meyer:** Verzeichniss der Pflanzen ... im Caucasus etc. St. Petersburg. 4°.
Michx. Fl. bor. Am. 2. 1803.	A. **Michaux:** Flora boreali-americana ... Parisiis et Argentorati. 8°.
Milde: Jahresber. schles. Ges. 1855.	J. **Milde:** 1. Ueber einige neue, in Schlesien beobachtete Farne. — pag. 92.
— Bot. Zeit. 1857.	— 2. Europa's Gefässkryptogamen, zusammengestellt. — pag. 474.
— ——— —	— 3. Ueber Botrychium boreale Milde. — pag. 880.
— Nova Acta 26. 1858.	— 4. Die Gefäss-Cryptogamen in Schlesien, preussischen und österreichischen Antheils. — pag. 369. 25 Tab.
— Oest. Bot. Zeit. 8. 1858.	— 5. Ueber Botrychium matricariaefolium, simplex und anthemoides. — pag. 94.
— ——— —	— 6. Ueber Aspidium lobatum, aculeatum und Braunii. — pag. 184.
— Bot. Zeit. 1864.	— 7. Ueber Botrychien, deren Eintheilung und Unterscheidung. — pag. 101.
— Verh. zool. bot. Ges. 14. 1864.	— 8. Asplenium dolosum Milde (A. Adianto nigro × Trichomanes). — pag. 165. t. 4.
— ——— —	— 9. Scolopendrium hybridum Milde, Proles hybrida orta ex Scolopendrio vulgari Sym. et Ceterach officinarum Willd. — pag. 235. t. 18.

Milde: Sporenpfl. 1865.	**J. Milde:** 10. Die höheren Sporenpflanzen Deutschland's und der Schweiz.-Leipzig. 8⁰.
— Bot. Zeit. **1866**.	—11. Die höheren Sporenpflanzen Europa's und der Atlantis. — pag. 137.
— ——— —	—12. Ueber Hymenocystis caucasica C. A. Meyer. Ueber Pteris radiata Mett. — pag. 179.
— ——— — **1867**. **1868**.	—13. Filices criticae. **1866**. pag. 309, 384, 392. **1867**. pag. 25, 40, 57, 89, 148. **1868**. pag. 201, 375, 449, 614, 882.
— ——— **1866**.	—14. Das Genus Athyrium. — pag. 373.
— ——— **1866**. **1867**.	—15. Materialen zur Beurtheilung der Darwinschen Theorie. **1866**. pag. 397. **1867**. pag. 152.
— Fil. Eur. 1867.	—16. Filices Europae et Atlantidis, Asiae minoris et Sibiriae. Lipsiae. 8⁰.
— Bot. Zeit. **1867**.	—17. Ueber eine neue Eigenthümlichkeit bei Botrychium. — pag. 238.
— ——— —	—18. Zur Farn-Flora Klein-Asiens. — pag. 292.
— ——— —	—19. Das Wesen der Farn-Flora der Atlantis. — pag. 417.
— Verh. zool. bot. Ges. **17**. 1867.	—20. Ueber Asplenium fissum Kit. und A. lepidum Pr. — pag. 817.
— ——— —	—21. Bemerkungen über einige Sporenpflanzen der deutschen Flora. — pag. 825.
— Mon. Osm. 1868.	—22. Monographia generis Osmundae. Vindobonae. 8⁰. 8 Tab.
— Bot. Zeit. **1868**.	—23. Index Osmundarum. — pag. 49.
— ——— —	—24. Ueber Aspidium nevadensis Boiss. — pag. 359.
— Verh. zool. bot. Ges. **18**. 1868.	—25. Index Botrychiorum. — pag. 506.
— ——— **19**. 1869.	—26. Botrychiorum Monographia. — pag. 55. t. 7-9.
— Bot. Zeit. **1870**.	—27. Ueber Athyrium, Asplenium und Verwandte. — pag. 329, 345, 370.
— ——— et in Jahresb. Schles. Ges. **48**. 1870.	—28. Ueber Todea und Leptopteris. — pag. 470. — Schl. Ges. pag. 95.
Millsp. Field Columb. Mus. Bot. Ser. **3**. 1903.	**Ch F. Millspaugh:** Plantae Yucatanae (regionis Antillanae) etc. — pag. 1.
Miq. Verhdl. Nederl. Inst. III. **4**. 1851.	**F. A. V. Miquel:** 1. Analecta Botanica Indica; seu Commentationes de variis stirpibus Asiae Australioris. — pag. 13. t. 5-7.
— Ann. Lugd. Bat. **3**. 1867.	— 2. Prolusio Florae Japonicae. Filices. — pag. 169; in Annales Musei Lugduno-Batavi. Fol.
— ——— **4**. 1868-69.	— 3. Filices (Ophioglosseae, Pteris, Allosurus, pag. 91-98; Aspidium, Phegopteris, Asplenium, Micropodium, pag. 155-169).
— Cat. Mus. Lugd. Bat.	— 4. Catalogus Musei Botanici Lugduno-Batavi. Pars prima. Flora japonica. Hagae. 8⁰.
Mirbel: Bull. Soc. Philom. **2**. 1801.	**C. Brisseau-Mirbel:** 1. Mémoire sur le Ramondia, nouveau genre de fougère. — pag. 179. t. 12. f. 3.

Mirbel: *Hist. nat. Vég. 5. 1803.	**C. Brisseau-Mirbel:** 2. Histoire naturelle des Végétaux; in Buffon: Cours complet d'Histoire naturelle, ed. Déterville. Paris. — (t. A. Gepp).
— Hist. nat. pl. 4. 1803.	— 3. Histoire naturelle générale et particulière des plantes; ouvrage faisant suite aux Oeuvres de Buffon: Histoire naturelle générale et particulière, édit. nouv. rédigé par Sonnini. Paris. 8°.
Moench: Meth. 1794. Suppl. 1802.	**K. Moench:** Methodus plantas horti botanici et agri Marburgensis a staminum situ describendi. Marburg Cattorum. 8°.
Moore: JoB. 5. 1853.	**T. Moore:** 1. List of Mr. Plant's Natal Ferns. — pag. 225.
— Proc. Linn. Soc. 2. 1853.	— 2. On Venation as a generic character in Ferns; with Observations on the genera Hewardia J. Smith, and Cionidium Moore. — pag. 210.
— ——— 1854.	— 3. Notes on some Ferns in the Wallichian Herbarium. — pag. 285.
— Ind. 1857-1862.	— 4. Index Filicum, being a Synopsis of the Genera of Ferns with Characters and Illustrations and an Enumeration of the Species with geographical Distribution, Synonyms, References to Figures, Descriptions &c, &c. London. 8°. I. Synopsis of the Genera. — pag. I-CLXII. 84 Tab. 1857. II. Enumeration of the Species. — pag. 1-396. (incompletum; nomen ultimum: Goniophlebium semipinnatifidum). pag. 1- 72: 1857. pag. 73-108: 1858. pag. 109-156: 1859. pag. 157-216: 1860. pag. 217-336: 1861. pag. 337-396: 1862.
— Journ. Linn. Soc. 2. 1858.	— 5. Notes on some Suprasoriferous Ferns. — pag. 129.
— ——— 4. 1860.	— 6. Notice of the Discovery of Lastrea remota in England. — pag. 192.
— Gard. Chr. 1856-1887.	— 7. [Diagnoses specierum novarum in hortis cultivarum interdum sub titulo: New Garden Ferns].
— ——— III. 1. 1887.	— 8. A Decade of New Adiantums. — pag. 41, 110, 447, 547.
Moore et Houlst. *Gard. Mag. Bot. 3. 1851.	— **et Houlston:** Descriptions and Woodcuts of Cultivated Ferns; in Gardener's Magazine of Botany. 1851. — (t. A. Gepp).
Moritz: *Syst. Verz. 1845-1846.	**A. Moritz(i):** Systematisches Verzeichniss der von H. Zollinger in den Jahren 1842-44 auf Java gesammelten Pflanzen. Solothurn. 8°.
F. Muell. Fragm. 5-11. 1864-1881.	**F. v. Mueller:** 1. Fragmenta Phytographiae Australiae. Melbourne. 8°. — Filices: 4. pag. 165: 1864 — 5. pag. 52, 74: 1865; pag. 111, 187: 1866 — 6. pag. 124, 199: 1868

	— **7**. pag. 83, 104, 119: 1870; pag. 155: 1871 — **8**. pag. 32: 1873; pag. 157, 175, 224: 1874 — **10**. pag. 104: 1877 — **11**. pag. 131: 1881.
F. Muell. Veg. Chatham Isl. 1864.	**F. v. Mueller:** 2. The Vegetation of the Chatham Islands. Melbourne. 8^0.
— Descr. Not. Pap. Pl. **1-6**. 1875-1885.	— 3. Descriptive Notes on Papuan Plants. Melbourne. 8^0. **1**. 1875. **2**. **3**. **4**. 1876. **6**. 1885.
— Cens. **1**. 1882.	— 4. Systematic Census of Australian Plants with Chronologic, Literary and Geographic Annotations. Part. I. Vasculares. Melbourne. 4^0.
— Sec. Cens. **1**. 1889.	— Second Systematic Census etc. Part. I. Vasculares. Melbourne. 4^0 obl.
— JoB. **1884**.	— 5. On some Plants of Norfolk Island, with description of a new Asplenium. — pag. 289.
— Tr. Roy. Soc. Victoria **1**. 1889.	— 6. Records of Observations on Sir William Mac Gregor's Highland Plants from New Guinea. — pag. 1. — (vide et Bak. n. 73).
— **et Bak.** JoB. **1887**.	— **et Baker:** Note on a collection of Ferns from Queensland. — pag. 162.
K. Müll. Bot. Zeit. **1854**.	**K. Müller:** 1. Ueber einige bisher verwechselte Arten der Farrngruppe der Hymenophyllaceen. — pag. 713, 729, 745.
— ——— —	— 2. Einige Worte über die Bedeutung des Zellenbaus für die Klassifikation, nebst Beschreibung sechs neuen Arten der Farrngattung Vittaria. pag. 537 t. 13.
— ——— **1864**.	— 3. Adiantum Jordani, ein neues Farrnkraut Californiens. — pag. 25 t. 1.
Müll. Fl. Fridrichsd. 1767.	**O. F. Müller:** Flora Fridrichsdaliana . . . Argentorati. 8^0. Tabb.
Murb. Lunds Univ. Årsskr. 27^5. 1892.	**S. Murbeck:** 1. Beiträge zur Flora von Südbosnien und der Herzegovina.
— ——— 27^6. 1892.	— 2. Tvenne Asplenier, deras affiniteter och genesis. 2 tab.
Nadeaud: *Enum. 1873.	**Nadeaud:** 1. Énumération des plantes indigènes de l'île de Tahiti. Paris.
— Journ. de Bot. **11**. 1897.	— 2. Note sur quelques plantes rares ou peu connues de Tahiti. — pag. 103.
— ——— **13**. 1899.	— 3. Plantes nouvelles des îles de la Societé. — pag. 1.
Neck. *Act. pal. **3**. phys. 1775.	**N. J. de Necker:** 1. ?
— Element. **3**. 1790.	— 2. Elementa botanica . . . Neowedae ad Rhenum. 8^0.
Nees et Bl. Nova Acta **11**. 1823.	**Nees ab Esenbeck et C. F. Blume:** Pugillus plantarum javanicarum e Cryptogamicarum variis ordinibus selectus. — pag. 117 t. 12-16.
Nees: Linn. **19**. 1847.	**Nees ab Esenbeck et S. Schauer:** Enumeratio et descriptiones generum novorum specierumque plantarum in terris mexicanis crescentium, quas

	Ruhlandus, Coloniensis, De Berghes et Dr. Ascherborn collegerunt. — pag. 681.
Nelson: *Proc. Biol. Soc. Wash. **17**. 1904.	**A. Nelson:** Plantae Andrewseae. — pag. 173.
Newm. Mag. Nat. Hist. II. **2**. 1838.	**E. Newman:** 1. A few Notes on the British species of the genus Polypodium of Linnaeus. — pag. 141.
— *Hist. br. Ferns 1840; ed. II. 1844; ed. III. 1854.	— 2. A History of british ferns and allied plants. London. 8⁰.
— *Phytol. **2**. 1846.	— 3. On the identity of Nephrodium foenisecii var. alatum of Madeira with Lastrea recurva of Britain. — pag. 509.
— * —— **4**. 1851.	— 4. Attempt to characterize another apparently undescribed species of Lastrea. — pag. 256.
— * —— —	— 5. Proposed addition of three new species and three new genera to our list of British Ferns. — pag. 368.
— * —— **5**. 1854.	— 6. Contributions towards a history of an Irish Asplenium considered as a distinct species and as entitled to a place in the British Flora. — pag. 36.
— * —— —	— 7. Contributions towards a history of a British Botrychium, considered as a distinct species, and entitled to a place in the British Flora. — pag. 134.
— * —— —	— 8. Note on Pseudathyrium flexile. — pag. 239.
Nutt. Gen. **2**. 1818.	**T. Nuttall:** The genera of North American plants. Philadelphia. 8⁰.
Nyland. Spic. pl. fenn. **2**. 1844.	**F. Nylander:** Spicilegium plantarum fennicarum. Centuria altera. Helsingforsiae. 8⁰.
Opiz: *Böh. phan. kr. Gew. 1823.	**P. M. Opiz:** Böheims phanerogamische und kryptogamische Gewächse . . . Prag. 8⁰.
Pal.-Beauv. Fl. d'Ow. **1**. 1804. **2**. 1807.	**Palisot-Beauvois:** Flore d'Oware et de Benin en Afrique. Paris. Fol. Tabb. (vol. 2 usque ad annum 1818 prodiit).
Pappe et Raws. Syn. Fil. Afr. austr. 1858.	**L. Pappe et R. W. Rawson:** Synopsis Filicum Africae australis . . . Cape Town. 8⁰.
Parm. Ann. sc. nat. VIII. **9**. 1899.	**M. P. Parmentier:** 1. Recherches sur la structure de la feuille des Fougères et sur leur classification. — pag. 289.
— *Bull. Ac. Géogr. Bot. Mans **9**. 1900.	— 2. Une nouvelle fougère hybride. — pag. 40 c. tab.
Payot: Cat. Foug. Mt. Blanc 1860.	**M. V. Payot:** Catalogue des Fougères, Prêles et Lycopodiacées des environs du Mont Blanc. Paris. 8⁰.
Pérard: Bull. Soc. Fr. **16**. 1869.	**A. Pérard:** Quelques observations sur les Cryptogames de l'arrondissement de Montluçon et du département de l'Allier. — pag. 255.
Phil. Bot. Zeit. **1856**. (*Anal. Univ. Chil. **1856**).	**R. A. Philippi:** 1. Bemerkungen über die Flora der Insel Juan Fernandez. — pag. 625, 641. (etiam in Anales de Universidad de Chile).

Phil. Ann. sc. nat. IV. 7. 1857.	**R. A. Philippi:** 2. Remarques sur la Flore de l'île de Juan Fernandez. — pag. 87.
— Linn. **29**. 1857. **30**. 1859-60.	— 3. Plantarum novarum chilensium. Filices: **29**. pag. 104; **30**. pag. 207.
— — **33**. 1864.	— 4. Plantarum novarum chilensium centuriae inclusis quibusdam Mendocinis et Patagonicis. — pag. 1.
— *Anal. Univ. Chile. **43**. 1873.	— 5. Descripcion de las plantas nuevas incorporados últimamente en el herbario chileno. — pag. 479.
— *——— **94**. 1896.	— 6. Plantas nuevas Chilenas. — pag. 335.
Pirotta: Ann. Ist. bot. Roma **8**. 1903.	**R. Pirotta:** Flora della Colonia Eritrea. — pag. 1. t. 12.
Poirault: Ann. sc. nat. VII. **18**. 1893.	**M. G. Poirault:** Recherches anatomiques sur la Cryptogames vasculaires. — pag. 113.
Poir. Enc. **5-8**. 1804-1808. Enc. Suppl. **1-5**. 1811-1817.	**J. L. M. Poiret** in Lamarck: Encyclopédie méthodique. Botanique. (vide Lam.). **5**. 1804. (Polypodium pag. 508; Pteris pag. 708). **8**. 1808. (Trichomanes pag. 62). Suppl. **1**. 1810. **2**. 1811. **3**. 1813. **4**. 1816. **5**. 1817.
Potts: Tr. N. Zeal. Inst. **9**. 1877.	**F. H. Potts:** 1. Notes on a Lomaria collected in the Malvern district. — pag. 491.
— ——— **10**. 1878.	— 2. Notes on Ferns. — pag. 358.
Prain: Rec. Bot. Surv. Ind. 2^4. 1903.	**Prain:** Flora of the Sundribuns; in Records of the Botanical Survey of India **2** nr. 4.
Prantl: Hym. 1875.	**K. Prantl:** 1. Untersuchungen zur Morphologie der Gefässkryptogamen. I. Heft. Die Hymenophyllaceen, die niedrigste Entwicklungsreihe der Farne. Leipzig. 4^0. 6 Tab.
— Schiz. 1881.	— 2. Untersuchungen etc. II. Heft. Die Schizaeaceen, morphologisch und systematisch bearbeitet. Leipzig. 4^0. 8 Tab.
— Verh. zool. bot. Ges. **31**. 1882.	— 3. Verzeichniss der von v. Fridau auf Schmarda's Reise 1853 in Ceylon gesammelten Farne. — pag. 117.
— Engl. Jahrb. **3**. 1882.	— 4. Die Farngattungen Cryptogramme und Pellaea. — pag. 403.
— Ber. deutsch. bot. Ges. **1**. 1883.	— 5. Systematische Uebersicht der Ophioglosseen. — pag. 348.
— ——— —	— 6. Helminthostachys zeylanica und ihre Beziehungen zu Ophioglossum und Botrychium. — pag. 155.
— Gartenfl. **32**. 1883.	— 7. Adiantopsis alata. — pag. 99 t. 1115.
— Jahrb. bot. Gart. Berlin **3**. 1884.	— 8. Beiträge zur Systematik der Ophioglosseen. — pag. 297 t. 78.
— Arb. bot. Gart. Breslau **1**. 1892.	— 9. Das System der Farne. — pag. 1.
Prentice: JoB. **1869**.	**C. Prentice:** 1. On a new Species of Hypoderris. — pag. 240.

Prentice: JoB. 1873. — **C. Prentice:** 2. On Lindsaea linearis Sw., with descriptions of two new Species. — pag. 295.

— *Queensl. Dept. Agric. Bull. 7. 1891. — 3. vide Bailey n. 9.

Pr. Del. Prag. **1.** 1822. — **C. B. Presl:** 1a. Plantarum novarum Brasiliae, praesertim Filicum Linnei diagnoses et descriptiones. — pag. 158. In J. C. et C. B. Presl: Deliciae Pragenses, Historiam naturalem spectantes. Pragae. 8⁰.

1b. Descriptiones duarum plantarum cryptogamarum Čechiae indigenarum. — pag. 232.

— Rel. Haenk. **1.** 1825. — 2. Pteridophyta in Reliquiae Haenkeanae etc. **1.** fasc. 1. 1825. Pragae. Fol. 12 Tab.

— Verh. Ges. vaterl. Mus. **1836.** — 3. Beschreibung zweier neuen böhmischen Arten der Gattung Asplenium. — pag. 59 t. 3. (Verh. d. Gesellsch. d. vaterländ. Museums in Böhmen in den 14 allg. Versammlung am 6. April 1836).

— Tent. 1836. — 4. Tentamen Pteridographiae, seu Genera Filicacearum praesertim juxta venarum decursum et distributionem exposita. Pragae. 12 Tab.

— Hym. 1843. — 5. Hymenophyllaceae. Eine botanische Abhandlung. Prag. 4⁰. 12 Tab. (Aus d. Abhandl. d. k. böhm. Ges. d. Wiss. V. (Folge) **3** besonders abgedruckt. — pag. 93).

— Abh. böhm. Ges. V. **3.** 1845. — 6. Botanische Bemerkungen. — pag. 431. (Marsilea pag. 580).

— Suppl. 1845. — 7. Supplementum Tentaminis Pteridographiae. Pragae. 4⁰. — (Aus d. Abh. böhm. Ges. d. Wiss. V. **4**).

— Abh. böhm. Ges. V. **5.** 1848. — 8. Die Gefässbündel im Stipes der Farrn. — pag. 307 t. 1-7.

— Epim. 1849. — 9. Epimeliae botanicae. Pragae. 4⁰. 15 Tab. *Obs.* t. W. Hooker (in JoB. **4.** 286. 1852) hoc opus initio anni 1852 prodiit.

Pritch. *List St. Helena Pl. 1836. — **S. F. Pritchard:** An alphabetical list of indigenous and exotic plants growing on the island of St. Helena. Capetown. 8⁰.

Pursh: Fl. Am. sept. **2.** 1814. — **F. T. Pursh** (Pursch): Flora Americae septentrionalis . . . London. 8⁰.
*ed. I. 1814; ed. II. 1816.

Racib. Pterid. Buit. 1898. — **M. Raciborski:** 1. Die Pteridophyten der Flora von Buitenzorg. — Flore de Buitenzorg publiée par le Jardin Botanique de l'État. 1me partie. Leiden. 8⁰.

— Nat. Tijds. Ned. Ind. **59.** 1900. — 2. Die Farne von Tegal. — pag. 234 mit 2 Tafeln.

— Bull. int. Acad. Cracovie **1902.** — 3. O kilku nieznanych paprociach archipelagu malajskiego. (Ueber einige unbekannte Farne des malayischen Archipels). Bull. intern. de

	l'Acad. des sciences de Cracovie. Class. math. et nat. — pag. 54.
Racib. Bull. int. Acad. Cracovie **1905**.	**M. Raciborski:** 4. O rodzaju paproci Allandodia Wall. (Ueber die Farngattung Allantodia Wall.). — pag. 346.
Raddi: Opusc. sci. Bol. **3**. 1819.	**J. Raddi:** 1. Synopsis Filicum Brasiliensium. — pag. 279 t. 11-12. — (vide Kunze, Flora **1824**. 305).
— Pl. Bras. **1**. 1825.	— 2. Plantarum Brasiliensium Nova Genera et Species novae, vel minus cognitae. Pars I. Filices. Florentinae. Fol. 84 Tab.
Raoul: Ann. sc. nat. III. **2**. 1844.	**M. Raoul:** 1. Choix de Plantes de la Nouvelle-Zélande. — pag. 113. (vide seq.).
— Choix de pl. 1846.	— 2. Choix de Plantes de la Nouvelle-Zélande. Paris. 4^{0}. t. 1-2.
Regel: Ind. Sem. ht. Petr. 1855; Linn. **28**. 1856; Ann. sc. nat. IV. **12**. 1859.	**E. Regel:** 1. Index Seminum horti botanici Imperialis Petropolitani. — Diagnoses etiam in Linn. **28**. pag. 366 et in Ann. sc. nat. IV. **12**. pag. 372.
— Gartenfl. **9**. 1860.	— 2. Diplazium Katzeri Rgl. — pag. 35 t. 282.
— ——— —	— 3. Alsophila guianensis. — pag. 119 t. 288.
— Mém. Ac. Petersb. VII. 4^{4}. 1861.	— 4. Tentamen Florae Ussuriensis.
— Gartenfl. **15**. 1866.	— 5. Grammatosorus Blumeanus. — pag. 335. c. figg.
— Acta Hort. Petr. **7**. 1881.	— 6. Descriptiones plantarum novarum et minus cognitarum. — pag. 541.
Reich. Linn. Syst. pl.	vide Linné nr. 4.
Reich. Verh. zool. bot. Ges. **9**. 1859.	**H. W. Reichardt:** 1. Asplenium Heufleri, eine Hybride zwischen Asplenium germanicum Weis und A. Trichomanes. — pag. 93 t. 4.
— Denkschr. Ak. Wien 17^{2}. 1859.	— 2. Ueber die Gefässbundel-Vertheilung im Stamme und Stipes der Farne. — pag. 21.
Reichb. fil. Bot. Zeit. 1872.	**H. G. Reichenbach fil.:** Botanische Notizen. — pag. 487.
Reinw. Syll. pl. **2**. 1824.	**C. G. C. Reinwardt:** Nova plantarum indicarum genera; in Sylloge plantarum novarum itemque minus cognitarum a praestantissimis botanicis adhuc viventibus collecta et a societate regia botanica Ratisbonensi edita. — Ratisbonae. 8^{0}.
Remy: Ann. sc. nat. III. **6**. 1846.	**J. Remy:** Analecta Boliviana. — pag. 345.
Retz. Fl. Scand. Prod. 1799.	**A. J. Retzius:** 1. Florae Scandinaviae Prodromus . . . Holmiae. 8^{0}.
— Obs. bot. **1**. 1779. **2**. 1781. **6**. 1791.	— 2. Observationes botanicae sex fasciculis comprehensae. Lipsiae. Fol. Tabb.
A. Rich. Fl. N. Zél. 1832.	**A. Richard:** 1. Essai d'une Flore de la Nouvelle-Zélande; in Voyage de découvertes de l'Astrolabe. Botanique par A. Lesson et A. Richard. Paris 8^{0}. Atlas in Fol. t. 10-16.
— Sert. Astrol. 1834.	— 2. Sertum Astrolabianum . . . in Voyage de déc. de l'Astrolabe. (vide praeced.).

Rich. Act. Soc. hist. nat. Paris **1**. 1792.	**L. Richard:** Catalogus plantarum, ad societatem ineunte anno 1792 e Cayenna missarum a Domino Le Blond. — pag. 105.
Richards. in Frankl. Narr. of a journ. 1823.	**J. Richardson:** Botanical Appendix to John Franklin: Narrative of a journey to the shores of the Polar Sea in the years 1819-22. London. 4⁰. — pag. 754. — (n. sp. R. Brown descripsit).
Robins. et Fernald: Proc. Amer. Acad. **30**. 1895.	**B. L. Robinson and M. L. Fernald:** New Plants collected by Messrs. C. V. Hartman and C. E. Lloyd upon an Archaeological Expedition to Northwestern Mexico under the Direction of Dr. Carl Lumholtz. — pag. 114.
Rosenst. Mém. l'Herb. Boiss. nr. 9. 1900.	**E. Rosenstock:** 1. Aspidium libanoticum. n. sp.
— Hedwigia **43**. 1904.	— 2. Beiträge zur Pteridophytenflora Südbrasiliens. — pag. 210.
— Festschr. Alb. v. Bamberg 1905.	— 3. Einige neue Farne aus Südbrasilien; in Festschrift Albert von Bamberg. Gotha 1905. — pag. 56. 2 Tab.
Roth: Tent. Fl. Germ. **1**. 1788. **3**. 1800. (Röm. Arch. **2**¹. 1799).	**A. W. Roth:** 1. Tentamen Florae Germanicae. Lipsiae. 8⁰. (— Auszug eines Briefes von Hrn. Prof. Mertens in Bremen an den Herausgeber. — pag. 103. — In hac epistola nomina nova Rothii primum publicata sunt).
— Ust. neu. Ann. **2**¹⁰. 1794. **4**¹². 1796.	— 2. Observationes Botanicae. **2**¹⁰. pag. 34. **4**¹² pag. 1.
— Catal. bot. **1**. 1797.	— 3. Catalecta botanica . . . Lipsiae. 8⁰.
— Nov. pl. sp. 1821.	— 4. Novae plantarum species praesertim Indiae orientalis . . . Halberstadii. 8⁰.
Rouy: *Ill. pl. Eur. rar. 1896.	**Rouy:** 1. Illustrationes plantarum Europae rariorum. Paris.
— Bull. Soc. Fr. **44**. 1897.	— 2. Notices botaniques. — pag. 437.
Rovirosa: Naturaleza II. **2**. 1892.	**J. N. Rovirosa:** Observaciones sobre algunos helechos mexicanos de la tribu de las Asplenieas. — pag. 179 t. 10-11.
Roxb. in *Beats. St. Helen. 1816.	**W. Roxburgh:** 1. An alphabetical list of plants, seen by Dr. Roxburgh growing on the Island of St. Helena in 1813-14. — Appendix to A. Beatson: Tracts relative to the Island of St. Helena . . . London. 4⁰.
— Calc. Journ. **4**. 1844.	— 2. (W. Griffith): The Cryptogamous Plants of Dr. Roxburgh, forming the fourth and last part of the Flora Indica. — pag. 463 t. 2[illegible]-34.
Rudge: Pl. Guian. **1**. 1805.	**E. Rudge:** Plantarum Guianae rariorum icones et descriptiones hactenus ineditae. Londini. Fol. t. 35, 36.
Ruiz: *Mem. leg. Calaguala 1805.	**H. Ruiz Lopez:** Memoria sobre la legitima Calaguala y otras dos raices que con el mismo nombre nos vienen de la America meridional. Madrid. 4⁰. 1 Tab. (Polypodium Calaguala).

Rupr. Dist. cr. vasc. Ross. 1845.	**F. J. Ruprecht:** Distributio cryptogamicarum vascularium in Imperio Rossico; in Beiträge zur Pflanzenkunde des Russ. Reiches. Dritte Lieferung. St. Petersburg. 8^0.
Sadeb. Jahrb. Hamb. Wiss. Anst. **14**. Beih. 3. 1897.	**R. Sadebeck:** 1. Filices Camerunianae Dinklageanae. 1 Taf.
— Nat. Pfl. 1^4. 1899-1900.	— 2. Hymenophyllaceae 1899. Hydropterides 1900, in Engler und Prantl: Die natürl. Pflanzenfamilien. 1. Teil. Abt. 4.
Sadl. *Descr. pl. epiph. Hung. 1820.	**J. Sadler:** 1. Descriptio plantarum epiphyllospermarum Hungariae . . . Pestini. 8^0. — (Diagnoses spec. nov. etiam in Flora **1820**. 715).
— *Fil. Hung. 1830.	— 2. De Filicibus veris Hungariae, Transsylvaniae, Croatiae et Litoralis hungarici. Buda. 8^0.
Sagorski: Oest. Bot. Zeit. **53**. 1903.	**E. Sagorski:** Ueber Aspidium rigidum Sw. und Aspidium pallidum Bory, (sub Nephrodio). — pag. 76.
St.-Hil. Voy. Dist. Diam. 1833.	**A. de Saint-Hilaire:** 1. Voyage dans le district des diamans et sur le littoral du Brésil. Paris. 8^0. 2 voll.
— Ann. sc. nat. II. **6**. 1836.	— 2. Rapport sur un Mémoire relatif à la structure et au développement des organes générateurs d'une espèce de Marsilea trouvé par M. Esprit Fabre dans les environs d'Agde. — pag. 375.
Salisb. *Prod. 1796.	**R. A. Salisbury:** Prodromus stirpium in horto ad Chapel Allerton vigentium. Londini. 8^0. — (t. A. Gepp).
Sal. Nom. 1883.	**C. Salomon:** Nomenclator der Gefässkryptogamen oder alphabetische Aufzählung der Gattungen und Arten der bekannten Gefässkryptogamen mit ihrer Synonymen und ihrer geographischen Verbreitung. Leipzig. 8^0.
Sanio: Verh. Bot. Ver. Brand. **25**. 1884.	**Sanio:** Nachtrag zu Die Gefässkryptogamen und Characeen der Flora von Lyck in Preussen. — pag. 60.
Schenck: Hedwigia **35**. 1896.	**Schenck:** Brasilianische Pteridophyten. — pag. 141. (Determ. a Kuhn et Christ).
Schkuhr: Kr. Gew. **1**. 1804-1809.	**C. Schkuhr:** Vier und zwanzigste Klasse des Linnéischen Pflanzensystems oder Kryptogamische Gewächse. 1. Wittenberg. 4^0. 1809. 219 Tab col. (Fasciculus primus (pag. 1-40, tab. 1-38?) 1804 prodiit; in indice annum 1806 ex errore citavi).
Schlecht. Adumbr. 1825-1827 (1832).	**D. F. L. de Schlechtendal:** 1. Adumbratio Filicum in promontorio Bonae Spei provenientium. Berlin. 4^0. 24 tab. t. 1 18: 1825. t. 19-24: 1827. (t. 25-30: 1832 ineditae).
— Linn. **26**. 1853.	— 2. De plantis variis Mexicanis. — pag. 364.
— Bot. Zeit. **1856**.	— 3. Ueber Polypodium horridum L. Eine kritische Betrachtung. — pag. 449, 465.

Schlecht. et Cham. Linn. 5. 1830.	**D. F. L. de Schlechtendal et A. de Chamisso:** Plantarum mexicanarum a cel. viris Schiede et Deppe collectarum Recensio brevis. — Fil. pag. 605.
Schmid in Hoppe: Bot. Taschenb. **1803.**	**R. Schmid:** Ueber eine in Baiern entdeckte Osmunda. — pag. 1.
Scholtz: Enum. Fil. Silesiae 1836.	**H. Scholtz:** Enumeratio Filicum in Silesia sponte crescentium . . . Vratislaviae. 8⁰.
Schott: Gen. Fil. 1834.	**H. Schott:** Genera Filicum. Vindobonae. 4⁰ obl. 20 Tab.
Schrad. Gött. gel. Anz. **1818.**	**H. A. Schrader:** 1. Analecta ad Floram Capensem. Sect. I. Filices. — pag. 913.
— ——— **1824.**	— 2. Illustratio Filicum a Serenissimo Principe Neowidensi in Brasilia observatarum, praemissis animadversionibus de hujus familiae structura et oeconomia. — pag. 857.
Schrank: Bai. Fl. **2.** 1789.	**F. Schrank:** Baiersche Flora. München. 8⁰.
Schreb. Spic. Fl. Lips. 1771.	**J. Schreber:** 1. Spicilegium Florae Lipsicae. Lipsiae. 8⁰.
— Gen. pl. ed. VIII. **2.** 1791.	— 2. Linné: Genera plantarum. Francofurti a/M. 8⁰.
Schultz: Flora **1844. 1845.**	**F. W. Schultz:** Einige neue Pflanzenarten. **1844.** pag. 806. **1845.** pag. 738.
Schum. Enum. pl. Saell. **2.** 1803.	**C. F. Schumacher:** 1. Enumeratio plantarum in partibus Saellandiae septentrionalis et orientalis. Havniae. 8⁰.
— Vid. Selsk. Afh. **4.** 1827 (1829).	— 2. Beskrivelse af Guineeiske Planter, som ere fundne af Danske Botanikere, især af Etatsraad Thonning. — Vidensk. Selsk. natur.- og matemat. Afhandl. Kjøbenhavn. — pag. 1. 1829. — Seorsim impr., Kjøbenhavn 1827. 4⁰.
Schum. et Laut Fl. deut. Südsee 1901.	**K. Schumann und K. Lauterbach:** Die Flora der deutschen Schutzgebiete in der Südsee. Leipzig. 8⁰. Tab. (Filices Diels et Christ elab.).
— Nachtr. Fl. deut. Südsee 1905.	— Nachträge. Leipzig. 8⁰. Tab. (Filices Christ elab.).
Schur: *Verhandl. Siebenbürg. Verein **2.** 1851.	**F. Schur:** 1. Beiträge zur Kenntniss der Flora Siebenbürgens. — pag. 167.
— Oest. Bot. Zeit. **8.** 1858.	— 2. Die siebenbürgischen Farne. — pag. 133, 192, 322.
— Enum. Pl. Transs. 1866.	— 3. Enumeratio Plantarum Transsilvanicae . . . Vindobonae. 8⁰.
Schweinf. Bot. Zeit. **1871.**	**G. Schweinfurth:** Bericht über die botanischen Ergebnisse der ersten Niam-Niam Reise Januar-Juli 1870. — pag. 361. c. fig. — vide A. Braun nr. 7.
Scop. Fl. Carn. 1760.	**J. A. Scopoli:** Flora carniolica . . . Viennae. 8⁰.
Scott: Tr. Linn. Soc. **30.** 1874.	**J. Scott:** Notes on the Tree Ferns of British Sikkim, with Descriptions of three New Species, and a few supplemental remarks on their relations to Palms and Cycads. — pag. 1. t. 1-18.

Seaton: *Contr. U. S. Nat. Herb. **1**. 1890. — **H. E. Seaton:** Pteridophyta ap. J. M. Coulter: Upon a Collection of Plants made by Mr. G. C. Nealley in the region of the Rio Grande in Texas, from Brazos Santiago to El Paso County. — pag. 65.

Sessé et Mociño: Fl. Mex. 1896. — **M. Sessé et J. M. Mociño:** Flora mecicana (1790-1800?) edit. R. Ramirez in La Naturaleza II. **2**. apendice. Filices in nr. 11.

Seub. Fl. Azor. 1844. — **M. Seubert:** Flora Azorica . . . Bonnae. 4⁰.

Shimek: Bull. Lab. Nat. Hist. Univ. Iowa **4**. 1897. — **B. Shimek:** The Ferns of Nicaragua. An account of the Ferns collected by the Nicaragua Botanical Expedition of the State University of Iowa 1893. — pag. 116. 20 Tab.

Sim: *Handb. Ferns Kaffraria 1891. — **T. R. Sim:** 1. Handbook of the Ferns of Kaffraria. Capetown 1891. — (t. A. Gepp).

— *Ferns S. Afr. 1892. — — 2. The Ferns of South Africa, containing descriptions and figures with localities, cultural notes . . . Capetown. 8⁰. 159 Tab. — (t. A. Gepp).

Slosson: *Rhodora **6**. 1904. — **M. Slosson:** A new hybrid fern from Vermont. — pag. 75.

Donn. Smith: Bot. Gaz. **12-33**. 1887-1902. — **J. Donnell Smith:** Undescribed plants from Guatemala and other Central American Republics. **12**. pag. 131 t. 11: 1887 — **13**. pag. 77: 1888 — **15**. pag. 27 t. 4: 1890 — **19**. pag. 254 t. 26: 1894 — **20**. pag. 538: 1895 (nov. spec. Christ descripsit) — **23**. pag. 251: 1897 — **33**. pag. 262: 1902.

Sm. Pl. Ic. ined. **2**. 1790. **3**. 1791. — **J. E. Smith:** 1. Plantarum icones hactenus ineditae plerumque ad plantas in herbario Linneano conservatas delineata. London. Fol.

— Mém. Ac. Turin **5**. 1793 (1791?). — — 2. Tentamen botanicum de Filicum generibus dorsiferarum. — pag. 401 t. 9.

— Fl. brit. **3**. *1804. 1805. — — 3. Flora britannica . . . *Londini. 8⁰. — Edit. curavit J. J. Römer. Turici 1805.

— in *Rees: Cycl. 1802-1820. — — 4. in Rees: The Cyclopaedia, or, Universal Dictionary of Arts, Sciences and Literature. London. 4⁰. 39 vol.
Conf. Jackson in JoB. **1877**. p. 107.

— *Engl. Fl. **4**. 1828. — — 5. The English Flora. London. 8⁰.

J. Sm. JoB. **3**. 1840. — **J. Smith:** 1. Remarks on the Genus Sphaerostephanos among Ferns. — pag. 17.

— — 1841. — — 2. Enumeratio Filicum Philippinarum, or, a Systematic Arrangement of the Ferns collected by H. Cuming Esq. — pag. 392.

— — — — 3. On Hewardia, a new Genus of Ferns. — pag. 431 t. 16-17.

— — **4**. 1841. Lond. JoB. **1**. 1842. **2**. 1843. — — 4. An arrangement and definition of the Genera of Ferns with observations on the affinities of each Genus. **4**. pag. 38, 147 t. 18-19. — **1**. pag. 419, 659. — **2**. pag. 378.

— — **1**. 1842. — — 5. Filices in Bentham: Contributions towards a Flora of South America. Enumeration of

	Plants collected by Mr. Schomburgk in British Guiana. — pag. 193.
J. Sm. Lond. JoB. **4**. 1845.	**J. Smith:** 6. Observations on a New Genus of Ferns. — pag. 166 t. 7-8.
— Bot. Mag. **72**. Comp. 1846.	— 7. An Enumeration of Ferns cultivated in the Royal Gardens at Kew, in December 1845, with characters and observations on some of the Genera and Species. — Compendium to the Botanical Magazine vol. 72 (III. **2**). — pag. 7.
— Bot. Voy. Herald 1854-57.	— 8. Filices in Seemann: Botany of the Voyage of H. M. S. Herald. London. 4⁰. 4 Tab. Fil. (Filices mexicanae et panamenses. — pag. 226, 337. 1854; Filices in Flora of the Island of Hongkong. — pag. 425. 1857).
— *Cat. Kew Ferns 1856.	— 9. Catalogue of the Ferns in the Royal Gardens of Kew. —.
— Cat. cult. Ferns 1857.	—10. Cultivated Ferns, or, a Catalogue of exotic and indigenous Ferns cultivated in British Gardens. London. 8⁰.
— Bonplandia **9**. 1861.	—11. Filices apud Seemann: Plantae Vitienses. — pag. 253.
— Ferns br. a. for. 1866. ed. II. 1877.	—12. Ferns, British and Foreign. The History, Organography, Classification and Enumeration of the Species of Garden Ferns . . . London. 8⁰. Figg. — ed. II. New and enlarged edition 1877, reprint. 1896.
— Hist. Fil. 1875.	—13. Historia Filicum: an exposition of the Nature, Number, and Organography of Ferns, and review of the principles upon which genera are founded, and the systems of classification of the principal authors, with a new general arrangement; characters of the genera; remarks on their relationship to one another; their species; reference to authors; geographical distributions; etc. etc. with 30 lithografic plates illustrating the characters of the Tribes. London. 8⁰.
Sod. *Rec. 1883.	**A. Sodiro:** 1. Recensio Cryptogamarum vascularium Quitensium. Quiti. 8⁰. — (vide Bot. Centralbl. **26**. pag. 37. 1886).
— *Anal. Univ. Quit. **4**. 1890.	— 2. Acrostichum Yatesii Sod. — pag. 174.
— Cr. vasc. quit. 1893.	— 3. Cryptogamae vasculares quitenses, adjectis speciebus in aliis provinciis ditionis Ecuadorensis hactenus detectis. Quiti. 8⁰. 7 Tab.
— Sert. Fl. Ecuad. 1905.	— 4. Sertula Florae Ecuadorensis. I. Acrosticha Ecuadorensia nova. Quiti 1905. (Extracto de los »Anales« de la Universidad de Quito, Tomo XIX. nr. 135).

Sommerf. Vet. Ak. Handl. **1834**. 1835.	**S. C. Sommerfelt:** Aspidium crenatum, en ny Art af Ormbunke. — pag. 102.
Spenn. Fl. Friburg. **1**. 1825.	**F. C. L. Spenner:** Flora Friburgensis et regionum proxime adjacentium. Friburgi Brisgoviae. 8⁰.
Splitgb. Tijds. Nat. Gesch. 7. 1840.	**F. L. Splitgerber:** Enumeratio Filicum et Lycopodiacearum quas in Surinamo legit. — pag. 391. — (Hoeven en de Vriese: Tijdschrift voor Natuurlijke Geschiedenis en Physiologie. Amsterdam).
Spr. Schrad. Journ. **1799**².	**C. Sprengel:** 1. Bemerkungen über einige kryptogamische Pflanzen. — pag. 265.
— Anleit. **3**. 1804. ed. II. **2**. 1818.	— 2. Anleitung zur Kenntniss der Gewächse, in Briefen. Dritte Sammlung. Einleitung in das Studium der kryptogamischen Gewächse. Halle. 8⁰.
— Nov. prov. hort. 1819.	— 3. Novi proventus hortorum academicorum Halensis et Berolinensis. . . . Halae. 8⁰.
— Nova Acta **10**. 1821.	— 4. Filicum Novarum Manipulus. — pag. 223. t. 16-17.
— Neue Entd. **3**. 1822.	— 5. Neue Entdeckungen im ganzen Umfang der Pflanzenkunde. Leipzig. 8⁰.
— Syst. **4**. 1827.	— 6. Carolini Linnaei Systema vegetabilium ed. 16 curante Curtio Sprengel. — vol. **4**. Göttingae. 8⁰.
— Tent. Suppl. Syst. 1828.	— 7. Tentamen supplementi ad Systematis vegetabilium Linnaei editionem decimam sextam. 8⁰.
Steud. Nom. **2**. 1824.	**E. G. Steudel:** Nomenclator Botanicus . . . **2**. Cryptogamae. Stuttgardtiae et Tubingae. 8⁰.
Stokes: *Bot. Mat. Med. **4**. 1812.	**J. Stokes:** A botanical Materia medica . . London. 8⁰.
Strempel: Fil. Ber. Syn. 1822.	**C. F. Strempel:** Filicum Berolinensium Synopsis. Dissertatio inaug. Berolini. 8⁰.
Sturm: Flora **1853**.	**J. W. Sturm:** 1. Beschreibung zweier neuen Farne aus Valdivia. — pag. 361.
— *En. pl. vasc. cr. chil. 1858.	— 2. Enumeratio plantarum vascularium cryptogamicarum chilensium. Ein Beitrag zu Farn-Flora Chile's. Nürnberg. (Aus d. zweiten Hefte d. Abh. d. naturh. Ges. Nürnberg). vol. I. pag. 151). — (t. A. Gepp).
— Fl. bras. **1**². 1859.	— 3. Ophioglosseae, Marattiaceae, Osmundaceae, Schizaeaceae, Gleicheniaceae et Hymenophylleae in Martius: Flora brasiliensis vol. 1 pars 2. — pag. 137-362. t. 9-19.
— Bot. Zeit. **1859**.	— 4. Ueber einige neue Hymenophyllum-Arten aus der Verwandtschaft des Hymenophyllum sericeum Sw. — pag. 297.
Sw. Prod. 1788.	**O. Swartz:** 1. Nova Genera et Species Plantarum Prodromus descriptionum vegetabilium, maximam partem incognitorum quae sub itinere in Indiam occidentalem Annis 1783-87 digessit. Holmiae 1788. 8⁰.

Sw. Obs. bot. 1791.	**O. Swartz:** 2. Observationes botanicae etc. Erlangae. 8^0.
— Ges. Nat. Fr. Berl. Neu. Schr. **2**. 1799.	— 3. Vittaria, eine neue Farrenkraut-Gattung. — pag. 129 t. 7.
— Schrad. Journ. **1800^2**. 1801.	— 4. Genera et Species Filicum ordine systematico redactarum. — pag. 1.
— ——— **1801^2**. 1803.	— 5. Observationes botanicae genera et species Filicum illustrantes. — pag. 273 t. 1-2.
— Vet. Akad. Nya Handl. 1804.	— 6. Anmärkningar och Tillägg vid Ormbunke-slägtet Mertensia. — pag. 171 t. 5 f. 1-5.
— Fl. Ind. occ. **3**. 1806.	— 7. Flora Indiae occidentalis. Erlangae. 8^0.
— Syn. 1806.	— 8. Synopsis Filicum, earum Genera et Species systematice complectens. Kiliae. 8^0. 5 Tab.
— Berl. Mag. **4**. 1810.	— 9. Blechnum unilaterale. Eine neue peruanische Art. — pag. 79 t. 3.
— Vet. Akad. Handl. **1817**.	—10. Nya Arter af Ormbunkar (Filices) från Brasilien. — pag. 53 t. 3-4.
— Adnot. 1829.	—11. Adnotationes botanicae, quas reliquit Olavus Swartz. Edid. J. E. Wikström. Holmiae. 8^0.
Symons: *Syn. pl. 1798.	**J. Symons:** Synopsis plantarum insulis britannicis indigenarum . . . Londini. 8^0.
Szysz. *in Beck: It. Princ. Cob. **2**. 1888.	**Szyszylowicz:** Pteridophyta in G. Ritter von Beck: Itinera Principum Coburgi. Die botanische Ausbeute von der Reisen Ihrer Hoheiten der Prinzen von Sachsen Coburg-Gotha . . . . Wien. 4^0. — (t. A. Gepp).
Taschner: Dissert. 1843.	**C. F. Taschner:** Dissertatio inauguralis medico-botanica de duabus novis Trichomanarum speciebus de earum nec non aliarum huius generis plantarum structura. Jenae. 4^0. 2 Tab.
Thouars: *Mél. bot. et voy. 1811.	**Aubert du Petit-Thouars:** 1. Mélanges de botanique et voyages. Paris 1811. continens:
— *Gen. nov. Madag. (1809?) (Roem. Collect. 1809).	— 2. Genera nova Madagascarica; etiam impressa in Roemer: Collectanea ad omnem rem botanicam spectantia. Turici. 4^0.
— *Fl. Trist. d'Ac. (1804?)	— 3. Esquisse de la flore de Tristan d'Acugna c. tab. (t. Kuhn anno 1804 prodiit: certe tamen postea).
Taubert: Engl. Jahrb. **21**. 1896.	**P. Taubert:** Beiträge zur Kenntniss der Flora des centralbrasilianischen Staates Goyaz — pag. 402.
Tausch: Flora **1834**. **1836**. **1839**.	**J. F. Tausch:** Botanische Beobachtungen. **1834**. pag. 513. **1836**. pag. 417. **1839**. pag. 176. (Notizen).
Ten. Woods: Proc. Linn. Soc. N. S. Wales II. **4**. 1890.	**J. E. Tenison-Woods:** A List of the Ferns of the Malay Peninsula. — pag. 75.
Ten. *Fl. Nap. 1811-1838.	**M. Tenore:** 1. Flora Napolitana. Napoli. Fol. 5 voll.
— *Atti Ist. Incorr. Napoli **5**. 1830.	— 2. Sulla Felce maschia e sopra altra sette specie affini. — pag. 125. (Atti del Real Istituto d'Incorraggiamento alle Scienze Naturali di Napoli. 4^0).
— *Mem. nuova felc. 1832.	— 3. Memoria su di una nuova felce e su varie altre specie. Napoli. 4^0. 5 Tab.

Terracino: Rend. Ac. Napoli **25**. 1886.	**A. Terracino:** Felci australiane. — pag. 44. (Rendiconti Acad. fis. e mat. Napoli).
Thomson: *Ferns N. Zeal. 1882.	**George M. Thomson:** The Ferns and Fern-Allies of New Zealand, with Instructions for their Collection and Hints on their Cultivation. Dunedin. 5 tab. — (t. A. Gepp).
Thbg. Fl. Jap. 1784.	**C. P. Thunberg:** 1. Flora Japonica . . . Lipsiae. 8⁰. tabb.
— Tr. Linn. Soc. **2**. 1794.	— 2. Botanical Observations on the Flora Japonica. — pag. 326.
— Nova Acta Ac. Petr. **9**. 1795.	— 3. Descriptio Caenopteridis. — pag. 157 t. D-G.
— Prod. Pl. Cap. 1800.	— 4. Prodromus Plantarum Capensium, quas in Promontorio Bonae Spei Africes Annis 1772-1775 collegit. — Pars posterior. Upsaliae. 8⁰.
Thwait. En. Pl. Zeyl. 1864.	**G. H. K. Thwaites:** Enumeratio Plantarum Zeylaniae: An Enumeration of Ceylon Plants, with Descriptions of the new and little known Genera and Species. London. 8⁰. — pag. 321-483: 1864.
Todaro: Syn. pl. acot. vasc. Sic. 1866.	**A. Todaro:** Synopsis plantarum acotyledonearum vascularium sponte provenientium in Sicilia insulisque adjacentibus. Panormi. 4⁰.
Torr. Cat. Pl. N. York 1818.	**J. Torrey:** A Catalogue of Plants growing spontaneously within thirty Miles of the City of New York. Albany. 8⁰.
Trevir. Berl. Mag. **7**. 1816.	**C. L. Treviranus:** Observationes circa plantas orientis, cum descriptionibus novarum aliquot specierum. — pag. 145 t. 3 f. 18-19.
Trev. Atti Ist. Veneto II. **2**. 1851.	**V. Trevisan:** 1. Sopra alcuni nuovi generi, e trentadue nuove specie di Felci. — pag. 161. — (sine descript).
— ——— III. **14**. 1869.	— 2. Sopra el felci denominate Struthiopteris e delle loro piu'strette affini. — pag. 553.
— *Atti Soc. It. sc. nat. **17**. 1874.	— 3. Sylloge sporophytarum Italiae. I. Prothallogamae. Milano.
— Hedwigia **13**. 1874.	— 4. Zwei neue Arten des Gattung Aspidium. — pag. 145.
— Nu. Giorn. bot. It. **7**. 1895.	— 5. Nuova specie de felce. — pag. 155.
— Rend. Ist. Lombardo II. **9**. 1876.	— 6. Mildella, nuovo genere, tipo di nuova tribu di Felci Polipodiacee. — pag. 807.
— Atti Ist. Veneto V. **3**. 1877.	— 7. Cheilosoria, nuovo genere di Polipodiacee Platilomee. — pag. 575.
Trimen: JoB. 1885.	**H. Trimen:** Notes on the Flora of Ceylon. — pag. 266.
Turcz. Bull. Soc. Nat. Moscou **5**. 1832.	**N. Turczaninow:** 1. Decades tres plantarum novarum Chinae boreali et Mongoliae chinensi incolarum. — pag. 180.
— ————— '1838.	— 2. Catalogus Plantarum in regionibus Baicalensibus et in Dahuria sponte crescentium. — pag. 85.
Tutcher: Journ. Linn. Soc. **37**. 1905.	**W. J. Tutcher:** Descriptions of some New Species, and Notes on their Chinese Plants. — pag. 58.

Und. *Our Nat. Ferns ed. IV. 1893, ed. VI. 1900.	**L. Underwood:** 1. Our Native Ferns and their Allies, with synoptical descriptions of the American Pteridophyta North of Mexico. New York.
— Bot. Gaz. **22**. 1896.	— 2. The Habitats of the rarer Ferns of Alabama. — pag. 407 t. 21.
— Minn. bot. St. **1**. 1897.	vide Heller.
— Bull. Torr. Cl. **25**. 1898.	— 3. American Ferns. I. The ternate Species of Botrychium. — pag. 521.
— ——— **26**. 1899.	— 4. American Ferns. II. The Genus Phanerophlebia. — pag. 204 t. 359-360.
— *Mem. Torr. Cl. **6**. 1899.	— 5. A review of the genera of ferns proposed prior to 1832. — pag. 247.
— Fern Bull. **8**. 1900.	— 6. A new Botrychium from Jamaica. — pag. 59.
— Bull. Torr. Cl. **28**. 1901.	— 7. A new Adiantum from New Mexico. — pag. 46.
— ——— **29**. 1902.	— 8. American Ferns. III. Our Genera of Aspidieae. — pag. 121. — IV. The Genus Gymnogramme of the Synopsis Filicum. — pag. 617. — V. A Review of the Genus Danaea. — pag. 669.
— ——— **30**. 1903.	— 9. An Index to the described Species of Botrychium. — pag. 42. Fig.
— ——— — —	—10. A Summary of our Present Knowledge of the Ferns of the Philippines. — pag. 665.
— Bull. Dep. Agr. Jam. **1**. 1903.	—11. Four recently described Ferns from Jamaica. — pag. 136.
— *Torreya **3**. 1903.	—12. Notes on southern Ferns. — pag. 17.
— ——— **5**. 1905.	—13. A much-named Fern. — pag. 87.
— ——— — —	—14. Botrychium silaifolium Presl. — pag. 106.
— Bull. Torr. Cl. **32**. 1905.	—15. The Genus Alcicornium of Gaudichaud. — pag. 587.
Und. et Cook: ——— **14**. 1887.	— **and O. F. Cook:** Notes on the American Species of Marsilia. — pag. 89.
Und. et Maxon: ——— **29**. 1902.	— **and W. R. Maxon:** Notes on a Collection of Cuban Pteridophyta, with Descriptions of four new Species. — pag. 577. Fig.
Urban: Symb. Ant. **4**. 1903.	**J. Urban:** Symbolae Antillanae, seu Fundamenta Florae Indiae occidentalis. — Flora portoricensis.
Vahl: Symb. **1-3**. 1790-1794.	**M. Vahl:** 1. Symbolae Botanicae sive plantarum tam earum quas in itinere, imprimis orientali collegit Petrus Forskål . . . Hauniae. Fol. tabb. **1**. 1790. **2**. 1791. **3**. 1794.
— Ecl. Amer. **3**. 1807.	— 2. Eclogae americanae, seu descriptiones plantarum praesertim America meridionalis nondum cognitarum. fasc. 3. Hauniae. Fol.
Vell. Fl. flum. **11**. 1827.	**Vellozo de Miranda:** Florae Fluminensis Icones . . . edidit Antonius da Arrabida. Parisiis. Fol. Tabb. (sine descr.).
Arch. Mus Nac. Rio Jan. **5**. 1881.	— Florae Fluminensis seu descriptionum plantarum praefectura fluminensi sponte nascentium liber primus . . . per manis A. de Vasconcellos & Souza sistit Fr. J. M. a Conceptione Vellozo

	1790. in Archivos do Museu Nacional do Rio Janeiro vol. 5. 4^0.
Vill. Fl. Delph. 1785.	**D. Villars:** 1. Flora Delphinalis, in Gilibert: Caroli Linnaei Systema plantarum Europae . . . **1.**
— Hist. pl. Dauph. **1.** 1786. **3.** 1789.	— 2. Histoire des plantes de Dauphiné. Grenoble. 8^0.
Vis. Fl. dalm. **1.** 1842. Suppl. 1872.	**R. de Visiani:** 1. Flora dalmatica . . . Lipsiae. 4^0.
— Atti Ist. Veneto III. **12.** 1867.	— 2. Illustrazione della Cheilantes Szovitsii F. et M. — pag. 656. t. 10-11.
Viv. Ann. bot. $\mathbf{1}^{2}$. 1804.	**D. Viviani:** 1. Florae italicae fragmenta; in Viviani: Annales botanici. **1.** pars 2. — pag. 135.
— *Fl. ital. fragm. 1808.	— 2. Florae italicae fragmenta ... fasc. 1. Genuae. 4^0.
— *Fl. Cors. 1824.	— 3. Florae Corsicae specierum novarum . . . . Genuae. 4^0 . . . Appendix 1825. Appendix altera 1830.
— Fl. Lib. 1824.	— 4. Florae Libycae Specimen . . . Genuae. Fol.
Vogler: *Dissert. 1781.	**J. A. Vogler:** Dissertatio inauguralis, sistens Polypodii speciem nuperis auctoribus ignotam, Polypodium montanum vocatam. Gissae. 4^0.
de Vriese: Ned. kr. Arch. **1.** 1846.	**W. H. de Vriese:** 1. Bijdragen tot de Kennis der Flora von Sumatra. — pag. 1.
— ——— — —	— 2. Goniophlebium Reinwardtii de Vriese, Polypodiacea nova horti Academici Lugduno-Batavi. — pag. 257.
— Tijdschr. Wis. en Nat. Wet. Amsterd. **1.** 1848.	— 3. Lastraea microchlamys, nova Filicis species Javanica, in Horto Acad. Bot. Lugd. Bat. culta. — pag. 155. (Tijdschrift voor de Wis-en Natuurkundige Wetenschappen, Letterkunde . . . Amsterdam. 8^0).
— Ned. kr. Arch. $\mathbf{2}^{1}$. 1850. $\mathbf{2}^{2}$. 1851.	— 4. Epimetrum ad indicem seminum horti Academici Lugduno-Batavi. Anno 1849: $\mathbf{2}^{1}$. p. 228. — Anno 1850: $\mathbf{2}^{2}$. p. 284. — Anno 1851:
— ——— **3.** 1852.	— 5. (Marattiaceae Indiae Bataviae orientalis, imprimis Junghuhnianae). — pag. 183.
— JoB. **4.** 1852.	— 6. Angiopteris longifolia Grev. et Hook. of Sir William Jackson Hooker's Herbarium and its synonyms. — pag. 323.
— Mon. 1853.	— 7. (et P. Hartig): Monographie des Marattiacées. Leide. Fol. 8 Tab.
Waisb. Öst. Bot. Zeit. **49.** 1899.	**A. Waisbecker:** 1. Beiträge zur Flora des Eisenburger Comitats. — pag. 60.
— Mag. bot. lapok **2.** 1903.	— 2. Új odatok Vas vármegye flórájához. (Neue Beiträge zur Flora des Eisenburger Comitats). Magyar botanikai lapok. — pag. 63.
Wall. Tidsskr. f. Naturvid. **1.** 1822. Flora 1823^{1}.	**N. Wallich:** 1. Brev til Professor J. W. Hornemann. dat. Katmandu i Nepal d. 18. Juli 1821. — Tidsskrift for Naturvidenskaberne. Kjøbenhavn. — pag. 257. Flora pag. 65.
— List 1828-1829.	— 2. A numerical List of dried specimens of plants in the East India Company's Museum . . .

London. Fol. — (nomina nuda). n. 1-2160: 1828; n. 2161 ...: 1829.

Wall. Pl. as. rar. 1. 1829-30. — **N. Wallich:** 3. Plantae asiaticae rariores ... London. Fol. t. 16. 48: 1829. t. 52. 1830.

Wallr. Fl. Crypt. Germ. 1. 1831. — **F. G. Wallroth:** Flora Cryptogamica Germaniae: in A. J. Bluff et C. A. Fingerhuth: Compendium Florae Germanicae. Sectio II. — Norimbergae. 12⁰.

Walt. Fl. carol. 1788. — **Th. Walter:** Flora Caroliniana . . . Londini. 8⁰.

Watelet: Bull. Soc. Fr. 5. 1858. — **M. A. Watelet:** Sur quelques fougères observées dans le département de l'Aisne. — pag. 15.

Watson: *Phytol. 2. 1846. — **H. C. Watson:** Notes on the Lastrea foenesecii as a species including both forms of Nephrodium foenesecii, Lowe, and Aspidium dilatatum var. recurvum Bree. — pag. 568.

Watson: Proc. Amer. Ac. 26. 1891. — **Sereno Watson:** Notes upon a collection of Plants from the Island of Ascension. — pag. 161.

Watt: JoB. 1874. — **D. A. Watt:** On North American Species of Cheilanthes. — pag. 47.

Wawra: Oest. Bot. Zeit. 13. 1863. — **H. Wawra:** 1. Neue Pflanzenarten gesammelt auf der transatlantischen Expedition S. K. Hoheit des durchl. Herrn Erzherzogs Ferdinand Maximilian. — pag. 142.

— Maxim. Reise 1866. — — 2. Botanische Ergebnisse der Reise seiner Majestät des Kaisers von Mexico Maximilian I nach Brasilien (1859-60). Wien. Fol. t. 96-103.

Webb: in Hk. Niger Fl. 1849. — **P. B. Webb:** Specilegia Gorgonea; in W. J. Hooker: Niger Flora . . . London. 8⁰.

Webb et Berth: Hist. nat. Can. Phyt. 3. 1847. — — **et S. Berthelot:** Histoire naturelle des Iles Canaries. Tome III. Deuxième partie: Phytographia canariensis. Sectio III. Paris. 4⁰.

Weis: Pl. Cr. Gott. 1770. — **F. G. Weis:** Plantae Cryptogamicae Florae Gottingensis. Gottingae. 8⁰.

Whitwell: JoB. 1898. — **W. Whitwell:** Botrychium matricariaefolium A.Br. and B. lanceolatum Angstr. in Britain. — pag. 291. t. 388.

Wikstr. Vet. Ak. Handl. 1825. 1826. — **J. E. Wikström:** Nya eller mindre kända arter af Ormbunkar (Filices). — pag. 434.

Willd. Fl. Berol. Prod. 1787. — **K. L. Willdenow:** 1. Florae Berolinensis Prodromus. Berolini. 8⁰.

— Phyt. 1794. — — 2. Phytographia, seu descriptio rariorum minus cognitarum plantarum; fasc. prim. Erlangae. Fol.

— Akad. Wiss. Erfurt 1. 1802. — — 3. Bemerkungen über einige seltene Farrenkräuter. 3 Tab. (seorsim impr. in Willdenow u. Bernhardi: Zwey botanische Abhandlungen. Erfurt. 8⁰).

— Vet. Ak. Nya Handl. 1804. — — 4. Mertensia, ett nytt slägte af Ormbunkarne — pag. 163. t. 4-5.

— Berl. Mag. 3. 1809. — — 5. Einige Bemerkungen über die Gattung Onoclea. — pag. 160.

Willd. Enum. 1809; Suppl. 1813.	K. L **Willdenow:** 6. Enumeratio plantarum horti regii botanici Berolinensis . . . Berolini. 8⁰. — Suppl. ed. Schlechtendal. 1813.
— sp. 5. 1810.	— 7. Caroli a Linné Species plantarum . . . Editio quarta. Berolini. 8⁰.
H.B.Willd.	Humboldt et Bonpland in Willd. sp.
Willem. Usteri Ann. 18. 1796.	P. **Willemet:** Herbarium mauritianum. — pag. 1. (etiam seorsim impressum).
Wither. Bot. Arr. ed. III. 1796.	W. **Withering:** A botanical arrangement of all the vegetables naturally growing in Great Britain . . . *ed. I. 1776. *ed. II. 1787-1793. ed. III. 1796. London. 8⁰.
Wollaston: *Phytol. n. s. 1. 1855-56.	G. B. **Wollaston:** Lastrea filix mas. — pag. 171.
Woolson: Fern Bull. 12. 1904.	G. A. **Woolson:** Nephrodium pittsfordensis. — pag. 106.
Wright: Tr. Linn. Soc. II. Bot. 6. 1901.	C. H. **Wright:** Pteridophyta in Report on two Botanical Collections made by Messr. F. v. Mc Connell and J. J. Quelch at Mount Roraima in British Guiana. — pag. 97.
Yabe: Bot. Mag. Tokyo 16. 1902.	Y. **Yabe:** 1. A note of Ferns from the Island of Kōtō (Botol-Tobago). — pag. 45. pars japonica.
——— 17. 1903.	— 2. Florula Tsusimensis. — pag. 93.
— ——— — —	— 3. Filices Koreae Uchiyamanae; An Enumeration of Ferns collected by Mr. T. Uchiyama from Korea. — pag. 63.
— ——— 19. 1905.	— 4. Trichomanes Formosense et Loochooense. — pag. 31.
Yapp: Ann. Bot. 16. 1902.	R. H. **Yapp:** Two Malayan »Myrmecophilous« Ferns, Polypodium (Lecanopteris) carnosum (Blume), and Polypodium sinuosum Wall. — pag. 185 t. 10-12.
Yatabe: *Bot. Mag. Tokyo 5. 1891.	**Yatabe:** 1. A new Japanese Polypodium. — pag. 35. t. 21.
— *——— — —	— 2. A new Japanese Acrostichum: A Yoshinagae — pag. 109. t. 23.
— *——— — —	— 3. Acrostichum Tosaense. — pag. 149. t. 24.
Zeiller: Bull. Soc. Fr. 32. 1885.	**Zeiller:** Fougères recueillies dans la Péninsule malaise par M. de Morgan. — pag. 70.
Zenker: Plant. Ind. 1835.	J. C. **Zenker:** Plantae indicae quas in montibus Coimbaturicis coeruleis Nilagiri sive Neilgherries dictis collegit Rev. Bernh. Schmid. Jena. Fol. — (vide et Ann. sc. nat. II. 4. 379).
Zoll. Nat. Gen. Arch. Neer. Ind. 1. 1844. 2. 1845.	H. **Zollinger:** 1. Observationes Phytographicae praecipue genera et species nova nonnulla respicientes. 1. pag. 396. 2. pag. 200. — vide Hassk. nr. 2.
— Syst. Verz. 1854.	— 2. Systematisches Verzeichniss der im indischen Archipel in den Jahren 1842-1848 gesammelten sowie der aus Japan empfangenen Pflanzen. Zürich. 8⁰.

II.

ENUMERATIO SYSTEMATICA

Numeri post nomina auctorum abbreviata numeros operum in enumeratione alphabetica et numeri crassiusculi opera principalia indicant.

ITINERA CIRCA TERRAM.

Bory 5. 7 — Brack. — A. Brongn. 3 — Gaud. 3. 4 — Goldm — Klf. 3 — Kuhn 18 — Luerss. 7 — Mett. 16 — Pr. 2 — Szysz.

EUROPA. (Opera majora).

Milde 4. 10. **16**.

Europa centralis: Aschers et Graebn. — Christ 25 — Luerss. **14**.

Hispania et Portugal: Colmeiro.

AFRICA.

Kuhn 1. **2**. **13** — Bak. 85.

Insulae Atlantidis: Desf. — Milde 11. **16**. 19 — *Ins. Canar.* Bory 1 — L. v. Buch 1. 2 — Webb et Berth. **1** — Bolle **3** — *Madeira:* Lowe 1. 2. 3 — Johns. — *Ins. Azor.* Seub. — *Ins. Caboverd.* Bolle 1. 2.

St. Helena: Hemsl. **2** — Pritch. — Roxb. 1.

Ascension: Hemsl. **2** — Watson.

Africa occidentalis tropica (Guinea-Angola): Bak. 56 (S. Thomé) — Carr. 3 (Angola) — Fritsch (Angola) — Pal. Beauv. — Sadeb. 1 — Schum. 2 — Webb.

Africa centralis et orientalis tropica: Bak. 24. 42. 44. 64 — A.Br. 2. 7 — Christ 13. 46. 60 — Hier. **1**. 4 — Kuhn **20**. **21** — Schweinf.

Africa australis: Bak. 7. 11. 39 — Klf. 6 — Krasser — Kze. 4. 14 — Mac Ken — Mac Ken et Gerr. — Moore 1 — Pappe et Raws. **1** — Schrad. 1 — Schlecht. **1** — Sim **1**. **2** — Thbg. 4.

Madagascar: Bak. 22. 27. 34. 37. 43. 45. 50. 61. 71. 77. 85 — Christ 32. 42 — Kuhn **18** — Luerss. 8.

Ins. Comorenses: Bak. 47 — Kuhn **18**.

Réunion: Bory 2 — Cord. 1. **2.** 3 — Kuhn **13**.
Mauritius: Bak. **23**. 87 — Bojer — Bory 2 — Kuhn **13** — Willem.
Rodriguez: J. B. Balfour.
Ins. Sechellarum: Bak. **18**. 23 — Kuhn **13**.
Socotra: Kuhn 16.

ASIA.

Asia minor: Milde 16. 18.
Arabia: Dcne. 2 — Forsk.
Sibiria: Ledeb. — Milde **16** — Rupr.
Amur. China bor. Korea: Eat. 1. 3 — Komarov — Maxim. 1. 2. 3 — Turcz. 1. 2 — Yabe 3. (Korea).
Japonia: Bak. 26. 35 — Christ 9. 23. 34. 41. 52 — Eat. 1. 3 — Franch. et Sav. **1.** — Kze. 20 — Luerss. 13 — Mak. 1. 2 3. 5. 7. 8. 9. 10 — Maxim. 2 — Mett. 11 — Miq. 2. 4 — Thbg. 1. **2** — Yabe **2** — Yatabe 1. **2.** 3.
Liu-Kiu-Formosa: Bak. 48 — Christ 24. 51 — Hance 11 — Hk. et Arn. — Harr. — Luerss. 11 — Mak 4. 6 — Yabe 1. 4.
China centralis et meridionalis. Tibet: Bak. 15. 16. 32. 35. 60. 66. 69. 80. 81 — Benth. 1 — Christ 14. 17. 24. 36. **38**. 44. 49. **57** — Diels 2 — Franch. 1. **2** — Hance 1-13 — Hk. 15. 19. 24 — Kuhn 4.
India et Ceylon: Bak. 6 — Bedd. **1. 2. 3. 4.** 7 8. **9.** 11 — Blanf. 1. 2 — Bory 7 — Burm. — Christ 47 — Clarke 1. **2**. 3. 4 — Clarke et Bak. — Dcne. 3 — Don 1 — Gammie — Griff. 3. 4 — Hope 1. 2. **4**. — Kze. 29 — Prain — Prantl 3 — Roxb. 2 — Scott — Thwait. — Wall. 1. 2. 3. — Zenker.
Birma: Bedd. **2. 3. 4.** 5. 7. 8. 9.
Annam: Christ 59.
Tonkin: Bak. 74 — Christ 1. 4. 20 — Lour.
Siam: Christ 29.
Perak, Penang, Singapore . . .: Bedd. **2. 3. 4.** 5. 6. **9.** 10 — Luerss. 12 — Ten. Woods. — Zeiller.
Malesia: Bak. 37 (72. Christmas Ins.) — Blume 1. 2 — Bory 7 — Burck 2 — Christ 8. 24 — Kuhn **7** — Mett. **11** — Miq. 1. **3** — Racib. 3 — Zoll. 2.
— **Sumatra:** Bak. 33. 83 — Luerss. 12 — de Vriese 1.
— **Java:** Blume **1**. **2** — v.d.B. 1. **4** — Christ 6 — Hassk. 1-5 — Horsf. — Jungh. — Kze. **17** — Moritz — Nees et Bl. — Racib. **1**. 2 — Zoll. 1.
— **Borneo:** Bak. 29. 55. 62. 67. 78. 84 — Becc. 2 — Burck **1** — Ces. **1** — Christ **54**.
— **Celebes:** Christ 5. **12**. 48 — Kuhn 11.
— **Insulae minores:** Amann (Bangka) — Becc. 3 (Sumba; Timor) — Dcne. 1 Timor) — Hemsl. **2** (Aru) — G. Karst. (Molucc.) — Kuhn 18.
Insulae Philippinarum: Blanco — Brack. — Christ 15 — Copeland 1. **2** — Harr. — Laguna — J. Sm. 2 — Und. 10.

AUSTRALIA.

Bail. 5 — Benth. **2** — R.Br. 1 — French — F. Müll. 1. **4** — Kze. 18 — Lab. 1 — Luerss. 10 — Terracino.
Queensland: Bail. **1**-11 — F. Müll. et Bak. — Luerss. 3.
Tasmania: Hk.fil. **3**.

POLYNESIA.

Bak. 21 — Bory 5 — Brack. — Carr. 1 — Drake — Forst. 1. 2 — Hk. 4 — Hk. et Arn. — Luerss. 7 — A. Rich. 2.

Melanesia: Kuhn 18. 19 — Schum. et Laut. **1** — *N. Guinea:* Bail. 12 — Bak. 73. 86 — Becc. 2 — Ces. 2 — Christ 31. 61 — Forbes — F. Müll. 3. 6 — *N. Pommern:* Hier. 3 — *Salomon Ins.:* Bak. 9. 41 — *Sulu:* Bak. 30.
Novae Hebridae: Kuhn **8**.
Nova Caledonia: v.d.B. 6 — Fourn. 3. **9**. 12 — Lab. 2 — Mett. 10 — Terracino.
Fiji: Bak. 31. 53 — Brack. — Carr. **1** — Luerss. **1** — J. Sm. 11.
Samoa: Bak. 19. 20 — Brack. — Christ 10 — Luerss. 1. **2**.
Tonga: Luerss. 1 — Hemsl. **3**.
Ellice Ins.: Luerss. 1.
Ins. Societatis: Brack. — Drake **1** — Guill. — Nadeaud 1-3.
Cook Ins.: Cheesem. **3**.
Pomotou: Drake **1**.
Marquesas, Gambier et Wallis: Drake **1**.
Pelew et Cook Ins.: Luerss. 4. 5.
Hawaii: Brack. — Eat. 15. 16 — Heller — Hill. **1** — Klf. 3 — Luerss. 6 — Mann.
Lord Howe: Bak. 5. 8. 12 — Benth. 2 — F. Müll. 1.
Norfolk: Endl. 1 — Hew. 2 — F. Müll. 1. 5.
N. Zealand: Armstr. 1. **2**. 3 — Cheesem. 1. 2 — Col. 1-19 — A. Cunn. — Emys — Field **1**. 2 — Gilb. 5 — Hk.fil. **2**. **5** — Kirk 1-10 — Potts 1. 2 — Raoul 1. 2 — A. Rich 1 — Thomson.

INSULAE ANTARCTICAE.

Hemsl. **2** — Hombr. et Jacq. — Hk.fil. **1**.
Tristan d'Acunha: Carm. — Kuhn **2** — Thouars 3.
Amsterdam et St. Paul: Bak. 17 — Fourn. 11.

AMERICA.

America borealis: Dav. 1-6 — Eat. 11. **14** — Maxon 4. 2 (Alaska) — Und. **1**.
America centralis: Hemsl. **1** — Donn. Smith — Fourn. 2. 7.
— **Mexico:** Christ 56 — Dav. 7. 8 — Eat. 10. 17. 19. 21 — Fée 8. 9 — Fourn. **6** — Hk. 8 — Kuhn 6 — Kze. 8. 16 — Liebm. — Mart. et Gal. — Maxon 6 — Millsp. — Nees — Rovirosa — Schlecht. 2 — Schlecht. et Cham. — J. Sm. 8.
— **Guatemala:** Bertol. 3 — Hier. 5 — Donn. Smith — Maxon 6.
— **Nicaragua:** Fourn. 8. 10 — Shimek **1**.
— **Costa Rica:** Bak. 46. 58 — Bomm. et Christ 1. **2** — Christ **37**. **53** — Hemsl. 1 — Hier. 5 — Kuhn 12.
— **Panama:** J. Sm. 8.
India occidentalis: Fée **11** — Gris. 1. **2** — Husnot — Jenm. **12** — Krug — Sw. 1. 2. 7 — Wikstr.
— **Cuba:** Eat. 5. **6**. 9 — Gris. 3 — Kze. 2 — Und. et Maxon.
— **Haiti (S. Domingo):** Bak. 65.
— **Jamaica:** Bak. 59 — Clute 3 — Gilb. 3. 6 — Hew. 1 — Jenm. 1-7. **9**. 10 — Maxon 7. 12-14 — Sw. 1. 2 — Und. 11.
— **Puerto Rico:** Urb. 1.
— **Martinique:** Klf. 2.
— **St. Vincent:** Bak. **75**.
— **Grenada:** Bak. **79**.
— **Trinidad:** Bak. 68 — Eat. 13 — Jenm. 8.
— **Bermuda:** Gilb. 4 — Hemsl. **2**.

Guiana: Aublet — Bak. 63 — Jenm. 11. **12** — Kze. 23 — Rich. — Rudge — J. Sm. 5 — Splitgb. — Wright.
Venezuela: Eat. 6 — Kl. 2 — Maury.
Columbia: Bak. 36 — Fée 5-8 — Hier. **5** — Karst. 1. 2 — Kl. 2 — Kze. 30 Mett. **13**.
Ecuador: Bak. 25 — Hier. 5 — Kl. 2 — Sod. 1. 2. **3**.
Galapagos: Anders. — Hk.fil. 4.
Peru: Christ 58 — Harr. — Hk. 3 — Hk. et Grev. 2 — Kze. 2 — Mett. 3.
Chile: Colla — Gay — Mett. 3 — Phil. 3-6 — Sturm 1. **2**.
Juan Fernandez: Colla — Hemsl. **2** — Johow **1** — Kze. 5 — Phil. 1. 2.
Tierra del Fuego: Dusén — Hk.fil. **1**.
Ins. falklandicae: Gaud. **2** — Hk.fil. **1**.
Argentina: Gris. 4. 5 — Hier. **2**.
Uruguay: Hier. 2.
Paraguay: Bak. 28 — Christ 19. 22 — Hier. 2 — Lindm. 2.
Bolivia: Britton 1. 2 — Gilb. 2 — Hier. 2.
Brasilia: Bak. **4**. 10. 40. 49. 82 — Brack. — Christ 18. 22. 27. 30. 39. 58 — C. Chr. 1 — Fée **12** — Gardn. 3 — Kuhn 17 — Kze. 6. 7. 26 — Lindm. 1-4 — Mart. 2 — Pr. 1 a — Raddi 1. **2** — Rosenst. 2. 3 — St. Hil. 1 — Schenck — Schrad. 2 — Sturm **3** — Sw. 10 — Taubert — Vell. — Wawra 1. 2.
Trinidad australis: Hemsl. **2** — Luerss. 9.

OPERA SYSTEMATICA.

Bernh. 1. 2. 5 — Bomm. 1 — Christ **11** — Desv. **5** — Diels **1** — Endl. 2 — Fée 1. **5** — J. F. Gmel. — Grev. et Hk. 1 — Hk. 6. **11** — Hk. et Bak. **1** — Houtt. 1. 2 — Juss. — Keys. 1 — Klf. 4 — Kze. 13 — L. 1 — Mett. **2** — Moore **4** — Pr. **4** — Prantl 9 — Schott — Sm. 2 — J. Sm. **4**. **13** — Spr. 6 — Sw. 4. **8** — Trev. 6 — Willd. **7**.

MISCELLANEA (Icones etc.).

Bak. **3**. 76. 83 — Desv. 1. 2 — Ettingsh. 1. **2** — Fée 6. 7. 8. 10 — Hk. 1. **5**. **20**. 21. 22. **23** — Hk. et Grev. **1** — Jacq. 5 — Kuhn 5 — Kze. **5**. **9** — Langsd. et Fisch. — Pr. 9 — Schkuhr **1**.

OPERA MONOGRAPHICA

Acrophorus: Hk. 18 — Moore 3.
Acrostichum: Fée **2** — Pr. 9.
Actiniopteris: Milde 12.
Adenoderris: Maxon 11.
Adiantopsis: Prantl **7**.
Adiantum: Keys. 2 — Kuhn **14** — K. Müll. 3 — J. Sm. 3.
Aneimia: Gardn. 1.
Anogramma: Kuhn 15.
Antrophyum: Fée **4** — Kze. 21 — Luerss. 1 — Trev. 7.
Aspidium: Mett. **7**.
Asplenium: Berg. 1 (Caenopteris) — Bernh. 3 — Christ 45 — Cord. 3 — Heufl. — Hier. 8 — Luerss. 1. 2 — Mett. **9** — Milde 27 — Pr. 9 — Thbg. 3.

Ophioglossum: Bower — Britton 3 — Luerss. **3** — Prantl 8.
Paesia: Mett. **6** — Kuhn 15.
Pellaea: Fourn. **4** — Prantl 4 — Trev. 6.
Phanerophlebia: Und. **4**.
Photinopteris: Pr. 9.
Plagiogyria: Mett. **5**.
Platycerium: Fée **2** — Pr. 9 — Und. **15**.
Polybotrya: Fée **2**.
Polypodium: Desv. 4 — Hier. 6 — Mett. **4** — Pr. 9 — Yapp.
Polystichum: Christ 2. 55 — Hance 12 — Mett. **7** — Pr. 9.
Pteridium: Clute 1.
Pteris: Ag. **1**.
Salvinia: Griff. 2.
Schizaeceae: Diels **1** — Prantl **2** — Pr. **7**.
Stenochlaena: Fée **2** — Pr. 9.
Syngramma: Kuhn **15** — J. Sm. 6 — Trev. 7.
Trichomanes: Bak. 2 — Lindm. **1** — Prantl **1** — Pr. 9 — Taschner.
Vittaria: Fée **3** — Luerss. **1** — K. Müll. 2 — Sw. 3.
Woodsia: R.Br. 3 — Eat. 7 — Milde 12 — Trev 5.
Woodwardia: Hance 5 — Pr. 9.

OPERA DE FILICIBUS CULTIS.

Ait. — Bak. 87 — Fisch. et Mey 1 — K. Koch — Kze. 10. 19. **27** — Link 1. 2. **3** — E. J. Lowe **1**. **2** — Mett. **2** — Moore 7-8 — Moore et Houlst. — J. Sm. 7. 9. 10. **12**.

Adde nomina seq. omissa:

pag. 164. [Caenopteris] rutaefolia Berg. Acta Petr. **1782**[2]. 249 t. 7 f. 2. 1786 = A. achilleifolium.

— 655. [Vittaria] pumila Goebel ap. Krug, Engl. Jahrb. **24**. 136. 1897 = Hecistopteris.

— 506. dele **Lomagramme** [subgen. sub Polypodio] et lege **Loxogramme.**